# UNCLASSIFIED

# AD NUMBER

## AD849603

## LIMITATION CHANGES

## TO:

Approved for public release; distribution is unlimited.

# FROM:

Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; JAN 1969. Other requests shall be referred to Air Force Rocket Propulsion Laboratory, Edward, CA 93523. This document contains export-controlled technical data.

# AUTHORITY

AFRPL ltr, 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD849603

## INVESTIGATION AND COMPILATION OF THE THERMODYNAMIC PROPERTIES OF HIGH TEMPERATURE CHEMICAL SPECIES

## QUARTERLY TECHNICAL REPORT AFRPL-TR-69-70

(1 October 1968 - 31 December 1968)

January 1969

## AIR FORCE ROCKET PROPULSION LABORATORY RESEARCH AND TECHNOLOGY DIVISION AIR FORCE SYSTEMS COMMAND EDWARDS AIR FORCE BASE, CALIFORNIA 93523

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of AFRPL (RPPR-STINFO), Edwards, California 93523

> (Prepared under Contract Nr. F04611-67-C-0009 The Dow Chemical Company Midland, Michigan 48640)

> > 39

11

MAR 28 1969

#### NOTICE

The information in this report is presented in good faith, but no warranty is made, nor is freedom from any patent to be inferred.

#### LEGAL NOTICE

When U. S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise, or in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may be in any way related thereto.

#### DISTRIBUTION

Distribution has been made in accordance with the automatic distribution list, plus Categories B-1 through B-4 of the CPIA Publication No. 147, Chemical Propulsion List, May, 1967, with subsequent changes.



#### QUARTERLY TECHNICAL REPORT NR. 8

.

(1 October - 31 December 1968)

#### INVESTIGATION AND COMPILATION OF THE THERMODYNAMIC PROPERTIES OF HIGH TEMPERATURE CHEMICAL SPECIES

January 1969

Dow Report Nr. T0009-4Q-68

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of AFRPL (RPPR-STINFO), Edwards, California 93523.

> AIR FORCE SYSTEMS COMMAND RESEARCH AND TECHNOLOGY DIVISION ROCKET PROPULSION LABORATORY EDWARDS, CALIFORNIA CONTRACT NR. F04611-67-C-0009

> > THERMAL RESEARCH LABORATORY THE DOW CHEMICAL COMPANY MIDLAND, MICHIGAN 48640

#### FOREWORD

٠

.

•

This report was prepared by The Dow Chemical Company, Midland, Michigan, under USAF Contract Nr. F04611-67-C-0009. The Contract was initiated under Air Force Project Nr. 3148, 'Investigation and Compilation of the Thermodynamic Froperties of High Temperature Chemical Species.' The work was administered under the direction of the Rocket Propulsion Laboratory, Edwards Air Force Base, with Mr. Curtis C. Selph acting as Air Force Project Officer.

This is the eighth quarterly report, covering the work performed during 1 October - 31 December 1968. The Dow Report number is T0009-4Q-68.

The work was performed by I. H. Carr, J. Chao, A. T. Hu, G. C. Karris, H. Prophet, A. N. Syverud and D. U. Webb under the technical supervision of D. R. Stull.

Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

> W. H. Ebelke, Colonel, USAF Chief, Propellant Division

and indefension in a second side s also

the state state of the state of the state of

#### ABSTRACT

Supplement No. 30 to the JANAF Thermochemical Tables was completed and sent to the printers on December 27, 1968. This Supplement contains the new element calcium and its fluoride, revisions of several ionic species due to a determination of the electron affinity of BO<sub>2</sub>, and revisions of several fluorides due to a new table for HF.

A complete description of the measurements used in the simultaneous solution analysis of the interrelated HF heat of formation data is given in the Appendix.

----

#### JANAF THERMOCHEMICAL TABLES

Supplement No. 30 to the JANAF Thermochemical Tables was completed, for issue to the panel, and sent to the printers on December 27, 1968. This issue contains 58 tables representing a two-man level of effort. There are 25 new tables including the element calcium and its fluorides, and 33 revised tables. The revised tables include several ionic species, changed because of a reported value for the electron affinity of BO<sub>2</sub>, and several fluorides affected by the new HF table.

This supplement brings the total number of species covered to 1072 and the number of tables issued during the present contract to 280. The table of contents for this supplement is shown in Figure 1.

Twenty three thermochemical reactions from the literature are analyzed to obtain new selected values of  $\Box Hf_{298}^{\bullet}$  for HF(g), HF(50 H<sub>2</sub>0), NaF(c), BF<sub>3</sub>(g), NF<sub>3</sub>(g), CF<sub>4</sub>(g), and C<sub>2</sub>F<sub>4</sub>(polymer). A new method, called simultaneous adjustment, is used in place of the traditional method of sequential adjustment. The selected values are self-consistent and "best" in the least squares sense.

### Figure 1

### JANAF THERMOCHEMICAL DATA

Supplement No. 30, Issued December 31, 1968

Title Page	*CaF(g)	FNa(c)
AlF <sub>6</sub> Na <sub>3</sub> (c)	*CaF <sub>2</sub> (c)	FNa(8)
*ALF, Na, ( & )	*CaF,(&)	FNa(g)
AlHO,(g)	*CaF,(g)	*F2K (g)
A10, (g)	CITI(g)	*F,Li (g)
A10, (g)	Cl <sub>3</sub> Ti(g)	F,Li,(g)
BC1, (g)	*C1_Mo(c)	*F,Na (g)
BF <sub>2</sub> <sup>=</sup> (g)	*C1_Mo(&)	F,Na,(g)
B0, - (g)	$*Cl_{\mu}Mo(g)$	F,Si(g)
cci,(g)	*Cl <sub>2</sub> Mo(c)	F,Ti(g)
CC1 (g)	*C1_Mo(8)	F <sub>2</sub> Li <sub>2</sub> (g)
CHC1 <sub>2</sub> (g)	*Cl <sub>s</sub> Mo(g)	ITi(g)
CH <sub>2</sub> Cl <sub>2</sub> (g)	*Cl <sub>6</sub> Mo(c)	I,Ti(c)
*C,C1,(g)	*Cl <sub>6</sub> Mo(g)	I,Ti(g)
*C,HCI(g)	*Cs0(g)	I <sub>3</sub> Ti(c)
*Ca (ref)	*Cs,0(g)	I <sub>3</sub> Ti(c)
*Ca (α)	FH(g)	I <sub>u</sub> Ti(c)
*Ca (β)	FLi(c)	I <sub>L</sub> Ti(&)
*Ca (\$)	FLi(%)	I <sub>u</sub> Ti(g)
*Ca (g)	FLi(g)	т

\*New Table

ir.

The states

### APPENDIX

•

•

.

٠

•

•

### Enthalpies of Formation of Hydrogen Fluoride

### and Five Closely Related Fluorides

Ъу

A. N. Syverud

Thermal Research Laboratory The Dow Chemical Company Midland, Michigan 48640

#### Introduction

Recent research has provided much data pertinent to the values of  $\Delta Hf^{\circ}$  for HF(g) and HF(n H<sub>2</sub>O). More accurate values for these quantities are desirable in themselves and because they are prime values on which many other fluorides are based. These quantities have remained relatively uncertain despite many calorimetric studies in the period from 1880 to 1950. Subsequent work emphasized important systematic errors, namely, various side reactions involving impurities, corrosion of containers and self-polymerization of HF(g).

Date and Reference		<sup></sup> Hf <sup>•</sup> 298.15	
	HF(g)	HF (50 H,0)	HF (∞ H <sub>2</sub> O)
1936 ( <u>1</u> )	-64.0* -75	.56*	-78.2*
	-0.2	-0.10	-0.46
1952 ( <u>2</u> )	-64.2	-75.66	-78.66
	-0.6	-0.66	-0.46
1965 ( <u>3</u> )	-64.8±0.3	-76.316	-79.50
	-0.34	-0.46	-0.32
1968 (This selection)	-65.14±0.2	-76.78±0.1	-79.82±0.2

The history of values selected from critical analyses  $(\underline{1}, \underline{2}, \underline{3})$  of the data may be summarized as follows:

\* Values correspond to 18°C rather than 25°C.

It is apparent from the differences tabulated between the lines that there has been a continual progression to more negative values. The present selections are no exception. The systematic trend in the values appears to be a reflection of increasing knowledge of the side reactions and increasing emphasis on minimizing them or correcting for them. Therefore, thorough analysis of the chemical reaction is essential for calorimetry involving hydrogen fluoride, fluorine, or, for that matter, almost any fluoride. This criterion is satisfied by several recent and pertinent calorimetric studies which make possible the selection of more reliable values for hydrogen fluoride.

-4-

This paper documents the newly selected values of  $\Delta Hf_{298.15}^{\bullet}$ for HF(g)\*, HF(50 H<sub>2</sub>0), HF( $\omega$  H<sub>2</sub>0)\* and five closely related fluorides. First, a new approach to the problem, namely simultaneous adjustment, will be formulated, which will be used instead of the normal sequential adjustment (<u>1</u>, <u>2</u>, <u>3</u>). Next will be presented the observations selected for consideration and their resulting simultaneous adjustments. Finally, the results will be discussed and some suggestions offered for future experiments.

#### Simultaneous Adjustment

The mathematical problem involves the determination of values for several unknowns from an over-determined set of experimental observations of linear combinations of the unknowns. The problem may be formulated as follows: Experimental data are available for a series of reactions ( $\underline{i} = 1, 2...\underline{m}$ ) involving a number of species ( $\underline{j} = 1, 2...\underline{n}$ ). It is assumed that the data may be reduced to standard heats of reaction,  $R_{\underline{i}} = \Delta Hr_{298}^{\bullet}$ , which deviate from the true values by the errors  $E_{\underline{i}}$ . The heats of formation,  $F_{\underline{i}} = \Delta Hr_{298}^{\bullet}$ , are defined by  $\underline{m}$  heat balance equations of the form:  $\sum_{\underline{j}} c_{\underline{i}\underline{j}} F_{\underline{j}} = R_{\underline{i}} + E_{\underline{i}}$  [1]

where  $c_{\underline{ij}}$ , the stoichiometric coefficient for species  $\underline{j}$  in reaction  $\underline{i}$  is taken to be negative for reactants and positive for products. Values for  $c_{\underline{ij}}$  are presumed to be known exactly and a limiting uncertainty interval,  $\pm U_{\underline{i}}$ , is estimated for each observation  $R_{\underline{i}}$ . We wish to obtain values of  $F_{\underline{i}}$  which are "best" approximations for  $F_{\underline{i}}$ . Of course, neither  $F_{\underline{i}}$  nor  $E_{\underline{i}}$  will ever be known exactly; however, the deviations from the approximate solution are given exactly by:

$$E_{\underline{i}}' = (\Sigma c_{\underline{i}\underline{j}} F_{\underline{j}}') - R_{\underline{i}}.$$
 [2]

<sup>&</sup>quot;The description "(g)" denotes the ideal gas standard state, while "(cH\_0)" refers to the ionized, aqueous standard state of unit molality, which may be attained by appropriate extrapolation to infinite dilution.

The established procedure for adjustment of Equation [1] assumes that  $E_{i}$  = 0 for selected heat of reaction (or for a weighted average of selected values), and the heats of formation  $F_{i}$  are obtained one at a time by use of Equation [2] in some sequential fashion. Inconsistencies arising from multiple determinations or alternate paths are resolved in the selection process by attributing all of the deviations to those observations which are believed to be less accurate. This sequential selection of the variables will be called a sequential adjustment in order to distinguish it from the following alternative.

Simultaneous adjustments of Equation [1] are readily obtained (when  $\underline{m} > \underline{n}$ ) by minimization of a suitable function of the weighted deviation  $E_{\underline{i}}'/U_{\underline{i}}$ . Two different minimization criteria appear to be useful: least squares, which minimizes the sum of squares of  $E_{\underline{i}}'/U_{\underline{i}}$ , and least sum, which minimizes the sum of absolute magnitudes of  $E_{\underline{i}}'/U_{\underline{i}}$ . Least squares and least sum correspond, respectively, to the  $L_2$ -norm and the  $L_1$ -norm of approximation theory ( $\underline{4}$ ). The so-called  $L_{\underline{u}}$ -norm, which corresponds to minimization of the maximum value of  $E_{\underline{i}}'/U_{\underline{i}}$ , usually does not give acceptable adjustments ( $\underline{5}$ ) to Equation [1].

Least squares adjustments tend to distribute the deviations among all of the observations, while least sum adjustments assume at least as many zero deviations as there are variables ( $\underline{5}$ ). In the limit, as the simultaneous adjustment is reduced to the stepwise selection of  $F_{\underline{j}}'$ , least squares reduces to selection of the weighted average and least sum reduces to selection of the least uncertain observation.

Simultaneous adjustment may be extended (5) to include the Gibbs energy of formation and the entropy as additional variables. Thermodynamics then requires for each chemical species the linear constraint

$$\Delta Hf_{298}^{\bullet} = \Delta Gf_{298}^{\bullet} + 298.15 \Delta Sf_{298}^{\bullet}, \qquad [3]$$

where  $\Delta Sf_{298}^{\bullet}$  is the entropy of formation of that species from the

-6-

elements in their standard reference states. Linear constraints may be incorporated ( $\underline{6}$ ) without changing the adjustment problem. This problem, i.e. the solution of linear systems of algebraic equations, is the subject of many algorithms ( $\underline{7}$ ,  $\underline{8}$ ,  $\underline{9}$ ) which have been programmed for digital computers. Least squares adjustments are obtained with these algorithms, but least sum adjustments are usually obtained by the methods of linear programming ( $\underline{10}$ ,  $\underline{11}$ ).

#### Selected Observations

The thermodynamic data considered for possible inclusion in the analysis were those available at the end of 1968. The search included not only HF but also those fluorides having reliable, multiple links to HF. Recent data were concentrated mainly in the latter area. Several reviews  $(\underline{12} - \underline{16})$  were particularly helpful in screening the related fluorides for reliable and consistent links to HF. Data considered in the previous critical evaluations  $(\underline{1}, \underline{2}, \underline{3}, \underline{17})$  were newly screened, based on current knowledge. This process led to the selection of five fluorides which warrant consideration in the selection process. These fluorides include NaF(c),  $BF_3(g)$ ,  $NF_3(g)$ ,  $CF_4(g)$  and  $C_2F_4(poly$ mer).

The remainder of this section discusses the selected data for HF and these closely related fluorides. Since the selections are presumably neither definitive nor unique, suggestions of changes for use in future adjustments are welcome. There are several links, for example, which interrelate  $C_2F_6(g)$ ,  $C_2F_4(g)$ ,  $CHF_3(g)$ ,  $CClF_3(g)$ ,  $CBrF_3(g)$  and  $CF_3I(g)$  with the selected species. The inconsistencies for  $C_2F_4(g)$  and  $CHF_3(g)$  appear to be large enough so that these links will contribute little, if any, to a knowledge of HF. Thus, all of these species are subject to a secondary adjustment.

The experimental data for each species are reduced to a chosen standard state for convenience in the subsequent analysis. One exception is  $HF(n H_0)$  which is reduced to  $HF(50 H_0)$ . Other

-7-

aqueous species in the ionized standard state of unit molality will hereafter be denoted by the description '(aq)". Data for aqueous species are reduced by means of the selected curves of Reference (<u>17</u>) whenever possible. Thermodynamic functions for gaseous and condensed phases are taken from the JANAF Tables (<u>18</u>). Uncertainty intervals for calorimetric data are calculated from  $2\overline{s'}_Q$ , as recommended by Rossini (<u>19</u>), unless otherwise indicated. In some cases, the interval is increased by an additional contribution for the estimated systematic error. When the overall  $\Delta$ Hr is the sum of two or more independent values, then the overall uncertainty interval is calculated by means of Equation 17 of Rossini (<u>19</u>). The twenty three selected reactions are discussed under the headings Rl through R23.

R1. 0.5  $H_2(g)$  + 0.5  $F_2(g)$  + HF(50  $H_20$ )  $\Delta Hr_{298}^{\circ} = -76.68\pm0.05$  kcal/gfw King and Armstrong (2) measured the heats of reaction of fluorine, oxygen and  $0F_2$  with hydrogen by flame calorimetry at 30°C. The oxygen data agree with the accepted  $\Delta Hf^{\circ}$  for  $H_20(l)$  and the fluorine data reduce to the value shown above. This is a very thorough study which serves well to illustrate the importance of side reactions such as corrosion. The latter of the two papers (20) increases the uncertainty to 0.09 kcal/gfw based on estimates of the systematic uncertainty. The smaller value of 0.05 kcal/gfw, exclusive of the systematic uncertainty, has been assigned to weigh this observation more heavily.

R2.  $HF(g) \rightarrow H(g) + F(g)$   $\exists Hr_{298}^{\bullet} = -136.051\pm0.3 \text{ kcal/gfw}$ Johns and Barrow (21) obtained  $D_{3}' = 49310 \pm 100 \text{ cm}^{-1}$  from rotational predissociation of the vibrational levels in the UV spectra of HF. From this the value  $D_{0}' = -135.067 \pm 0.3 \text{ kcal/gfw}$  and the corresponding value at 298.15°K is derived.

-8-

and the second state of th

R3.  $HF(g) + NaOH(aq) + NaF(c) + H_2O(2) = Hr_{298}^{\circ} = -28.24\pm0.10 \text{ kcal/gfw}$ Rodenburg and Vanderzee (22) obtained values for the reaction  $HF(g) + NaOH(aq) + NaF(aq) + H_2O(2)$  from calorimetric studies with HF(real gas) at pressures of about 350 and 150 mm Hg. The results are  $\Delta Hr^{\circ} = -27.80 \pm 0.16$  and  $-28.01 \pm 0.10$  kcal/gfw, respectively, with preference being given to the latter value on experimental grounds (23).  $\Delta Hr^{\circ} = -28.01 \pm 0.10$  kcal/gfw includes a non-ideality correction of approximately  $0.20 \pm 0.025$ kcal/gfw. Combining this  $\Delta Hr^{\circ}$  with  $-\Delta H_{c}^{\circ} soln = -0.23 \pm 0.01$ kcal/gfw (24) for NaF(c), the selected observation given above is obtained.

R4.  $HF(50 H_20) + NaCl(c) + NaF(c) + HCl(aq)$   $\Delta Hr_{298}^{\circ} = -2.41\pm0.07 \text{ kcal/gfw}$ Coughlin (25) obtained  $\Delta Hr^{\circ} = -1.53 \pm 0.06 \text{ kcal/gfw}$  at 30°C for NaCl(c) + HF(5.716 H<sub>2</sub>0) + NaF(c) + HCl(12.731 H<sub>2</sub>0) from combination of five heat of solution measurements involving an acid solution of aluminum. Values (<u>17</u>, <u>18</u>) of Cp°,  $\phi_c$  and  $\phi_L$  yield a reduction to 25°C of 0.132 kcal/gfw, a reduction to HF(50 H<sub>2</sub>0) of 0.128 ± 0.03 kcal/gfw, and a reduction to HCl(aq) of -1.14 kcal/mol.

-9-

to equilibrium. Other aspects of the study appear to be thorough and well documented. It would be desirable to test for similar drifts in comparable studies (26) of BaCl\_-BaF, and NiCl\_-NiF,, but this was not done because of uncertainties in the free energy functions. Instead, it is noted that the third law  $\Delta Hr^{\bullet}$  is consistent with R3 within about 0.08 kcal/gfw, while the second law value is clearly inconsistent. Such behavior is observed in systems where the equilibrium constants are essentially correct at one extreme in temperature but show minor and increasing error toward the other extreme in temperature. Corrosion is suspected in this case, and the results at the lowest temperature are the most reliable. The individual third law values of AHr<sup>®</sup>, ranging from 3.93 kcal/gfw at 796°K to 3.56 kcal/gfw at 942°K, yield a mean of 3.81 kcal/gfw, which is influenced by the preponderance of points at lower temperatures. This mean is adopted and the uncertainty interval is estimated as ±0.25 kcal/gfw, which exceeds the range of the values and corresponds approximately to the maximum effect arising from uncertainty in the third law  $\Delta Sr^{\bullet}$ .

R6. 0.5  $F_2(g)$  + NaCl(c) + NaF(c) + 0.5 Cl<sub>2</sub>(g)

a ... which are the second and the second

ΔHr<sup>•</sup><sub>298</sub> =-39.30±0.4 kcal/gfw

Calorimetric data for this reaction at  $20^{\circ}$ C are reported by von Wartenberg and Fitzner (27). The reduction to  $25^{\circ}$ C is insignificant. Although the study is old and incompletely documented by current standards, the results are consistent with recent work. An uncertainty of 0.4 kcal/gfw based on 0.28 for the calorimetric experiments is assumed, and an estimate of 0.12 for other contributions.

R7.  $HF(50 H_20) + NaOH(aq) + NaF(c) + H_2O(l)$   $\Delta Hr_{298}^{\bullet} = -16.57\pm0.05 \text{ kcal/gfw}$ Kolesov and Skuratov (28) report  $\Delta Hn = -16.45 \pm 0.03 \text{ kcal/gfw}$  at 21.5°C for  $HF(16 H_20) + \text{LiOH}(3800 H_20) + \text{LiF}(3800 H_20) + H_2O(l)$ . It appears that this is the only modern determination of  $\Delta Hn$  for

and the second se

HF with any alkali hydroxide. With auxiliary data from  $(\underline{17}, \underline{18})$ , a reduction to 25°C of 0.074 ± 0.02 kcal/gfw is obtained, a reduction to HF(50 H<sub>2</sub>0) or 0.036 ± 0.01 kcal/gfw, and a combined reduction to LiOH(aq) and LiF(aq) of 0.001 kcal/gfw. The last mentioned standard states may be considered as OH<sup>-</sup>(aq) and F<sup>-</sup>(aq), or as NaOH(aq) and NaF(aq), without affecting the  $\Delta$ Hn = -16.34 ± 0.05 kcal/gfw. Combination of this value with  $-\Delta H_{c^+}^{\circ} = -0.23 \pm 0.01$ kcal/gfw (24) for NaF(c) gives the selected  $\Delta$ Hr<sup>\*</sup>.

R8.  $H^{+}(aq) + NaF(c) + HF(50 H_{2}0) + Na^{+}(aq)$ 

ΔHr<sup>•</sup><sub>298</sub> = 3.414±0.3 kcal/gfw This 'observation' is a combination of  $\phi_{1}$  = 3.184 ± 0.3 kcal/gfw (17) for HF(50 H<sub>2</sub>0) and  $\Delta H_c^{\circ}$  soln = 0.23 ± 0.01 kcal/gfw (24) for NaF(c). Itrepresents the composite of equilibrium and enthalpy data from which the dilution curve (17) was derived. Aqueous HF is a weak electrolyte which ionizes in dilute solutions according to the equilibria (a)  $HF = H^+ + F^-$  and (b)  $HF + F^- = HF_2^-$ . The two ionization reactions make an overall contribution of about 3 kcal/gfw to  $\phi_1$ . Parker (<u>17</u>) used selected values of  $\Box$ Ha = -3.0 and  $\Delta Hb = +0.662$  kcal/gfw, along with concentrations based on selected equilibrium data, to calculate  $\phi_T$  in dilute solutions. Recent data suggest that minor changes in these calculations may be desirable. In particular, the calculated dilution curve shows increasing deviations at lower molalities when compared with the new measurements of Cox and Harrop (29). Agreement is satisfactory near HF(1100  $H_00$ ), but near HF(6000  $H_00$ ) the calculated heats of dilution are roughly 240 cal/gfw (or 32%) larger than the observed values. Alternative selections for Ka, Kb,  $\pi$ Ha and  $\pi$ Hb might reduce this discrepancy considerably. The equilibrium data, both new (30) and old (31), are sufficiently divergent to allow changes in  $\pi$ Ha and  $\pi$ Hb of perhaps -0.2 and +0.5 kcal/gfw, respectively. Based on this review of the data, the uncertainty of  $\phi_{1}$  (50H<sub>2</sub>0) is estimated to be 0.3 kcal/gfw.

R9. C(graphite) +  $2F_2(g) \rightarrow CF_4(g) \qquad \Delta Hr_{298}^{\circ} = -223.05\pm0.18 \text{ kcal/gfw}$ Greenberg and Hubbard (32) report this result from a thorough study of the combustion in fluorine of natural and synthetic graphite.

R10.  $C_2F_4$  (polymer) +  $O_2(g)$  +  $2H_2O(2)$  +  $4HF(50 H_2O)$  +  $2CO_2(g)$  $\Delta Hr_{298}^{\circ} = -160.62\pm0.9$  kcal/gfw See the following discussion.

Rll.  $C_2F_4(polymer) + O_2(g) + CF_4(g) + CO_2(g)$   $\Delta Hr_{298}^{\circ} = -118.8\pm0.5 \text{ kcal/gfw}$ Good et al. (33) report the calorimetric combustion of various Teflon-oil mixtures. Reactions RlO and Rll are derived by extrapolation from the observed product ratios of HF(10 H<sub>2</sub>O)/CF<sub>4</sub> to the limits of only HF(10 H<sub>2</sub>O) and only CF<sub>4</sub>(g). RlO includes a reduction to HF(50 H<sub>2</sub>O) of -0.32\pm0.08 kcal/gfw.

R12.  $C_2F_4$ (polymer) +  $2F_2(g) \rightarrow 2CF_4(g)$ 

 $\Box Hr_{298}^{\circ} = -247.85\pm0.3 \text{ kcal/gfw}$ This  $\Delta Hr^{\circ}$  appears to be a minor adjustment (12) of the earlier data of Domalski and Armstrong (13) who report a value of  $-247.92\pm0.07$ kcal/gfw. The earlier value is labeled as Reaction Rl2a. Wood, Lagow and Margrave (34) reported -246.84 kcal/gfw for the same reaction. This is Rl2b and is reassigned an uncertainty of 0.2 kcal/gfw, since the authors' uncertainties and terminology are inconsistent with (19). No reason is apparent for the discrepancy of 1.0 kcal/gfw between Rl2 and Rl2b. Effects due to phase transitions of  $C_2F_4$  (polymer) near 20 and 30°C might be involved, but the overall  $\Delta Ht$  for these transitions is only 0.2 kcal/gfw (35).

R13. C(graphite) +  $2F_2(g)$  +  $CF_4(g)$   $\exists Hr_{298}^{\bullet} = -222.87\pm0.38$  kcal/gfw Domalski and Armstrong (<u>13</u>) report this result from their combustions in fluorine of graphite-Teflon mixtures.

こうちょうちょう しましょう あいていない ないないない い しんしていないない あいんできない あいかいない しょうない いろうちょう ないない しょうちょう

R14.  $CF_{\mu}(g) + 2H_{2}O(l) \rightarrow 4HF(50 H_{2}O) + CO_{2}(g)$  $\Delta Hr_{298}^{\bullet} = -41.506 \pm 0.3 \text{ kcal/gfw}$ Cox, Gundry and Head (36) report calorimetric combustions of mixtures of docosafluorobicyclohexyl  $(C_{1,2}F_{2,2})$  with benzoic acid. This study, which is analogous to that of Good et al. (33) with Teflon-oil mixtures, yields two reactions having quite different product ratios of HF(20  $H_2$ O)/CF<sub>4</sub>. By taking the difference of these two reactions, dividing by 3.3, and adding a reduction to HF(50 H<sub>2</sub>0) of  $-0.12 \pm 0.02$  kcal, the above observation is obtained. R15.  $8NF_3(g) + 3C_2N_2(g) + 6 CF_4(g) + 7N_2(g)$  $\Delta Hr_{298}^{\bullet} = -1308.8 \pm 1.3 \text{ kcal/gfw}$ Walker (37) reported a calorimetric value at 298.15°K of AHr° = -218.13 ± 0.15 kcal/gfw for 1/6 of the stoichiometry given above. The uncertainty is reassigned as 0.21 kcal/mol by reference to the original data, since the manuscript (37) is inconsistent in value and terminology with (19). Finally, the observation is multiplied by 6 in order to eliminate stoichiometric coefficients with infinite decimals.  $NF_3(g) + 1.5 H_2(g) + 3HF(50 H_20) + 0.5 N_2(g)$ R16.  $\Delta Hr_{298}^{\bullet} = -199.40 \pm 0.22 \text{ kcal/gfw}$ Sinke (38) reported a calorimetric value at 298.15°K of  $\Delta Hr^*$  = -199.49 $\pm$ 0.22 kcal/gfw for a final state of HF(123 H<sub>2</sub>0). The reduction to  $HF(50 H_20)$  is +0.09 ± 0.01 kcal/gfw.  $2NF_3(g) + S(c, rh) + SF_6(g) + N_2(g)$ R17. ∆Hr<sup>•</sup><sub>298</sub> = -228.26±0.25 kcal/gfw Walker (39) reported a calorimetric value at 298.15°K of AHr° = -228.26 ± 0.2 kcal/gfw. The uncertainty is reassigned as 0.25 kcal/gfw by reference to the original data, since the paper (39) is inconsistent in value and terminology with (19).

R18.  $NF_3(g) + 0.5 N_2(g) + 1.5 F_2(g) \Delta Hr_{298}^{\circ} = 31.44\pm0.3 \text{ kcal/gfw}$ Sinke (40) reported calorimetric data for the explosion of mixtures of hydrogen with 10% excess of  $NF_3$  and with 100% excess of  $NF_3$ . The complex final states were not corrected to standard states; instead, the experiments were designed to minimize the corrections. The difference between the two sets of data yields the result given above.

R19.  $NF_3(g) + B(c, \beta) + BF_3(g) + 0.5 N_2(g)$ 

 $\Delta Hr_{298}^{\bullet} = -239.46\pm1.2 \text{ kcal/gfw}$ Ludwig and Cooper (<u>41</u>) reported a calorimetric value of  $\Delta E_{c}^{\bullet}/M =$ -22177 cal/g after correction for 0.44% impurities in the sample of boron. Using 10.811 for the atomic weight of boron,  $\Delta E_{c}^{\bullet} =$ -239.76 ± 1.2 and  $\Delta Hr^{\bullet} = -239.46 \pm 1.2 \text{ kcal/gfw}$ . The amount of unburned boron remaining after combustion averaged 5% as determined by chemical analysis. This analysis was complicated by the retention on the bomb surfaces of some of the BF<sub>3</sub>(g), apparently in the form of a metal fluoride - BF<sub>3</sub> adduct which was not identified. It is likely that the metal fluoride in the adduct was formed, at least in part, by corrosion. No correction was made for adduct formation, so the calorimetric result may be biased. Bias may also arise from the impurity correction (cf. R20); however, the existing bias appears to be much smaller than the assigned uncertainty.

R20. 1.5  $F_2(g) + B(c, \beta) + BF_3(g)$   $\Delta Hr_{298}^* = -271.6\pm0.9$  kcal/gfw Wise et al. (<u>43</u>) reported the calorimetric result  $\Delta Hr^* = -270.10 \pm 0.24$  kcal/(10.82g B), based on correction for 0.62% impurities in the sample of boron. Reanalysis (<u>42</u>) of the sample with more reliable techniques showed the presence of 1.32% impurities. This made  $\Delta Hr^*$  more negative by 1.5 kcal/gfw and increased the uncertainty by more than a factor of three. Here is a pointed example of bias caused by side reactions due to impurities. Calorimetric combustions in fluorine have also been reported by

-14-

Gross et al.  $(\underline{44})$ , but their boron (the zone-refined sample) was not analyzed for C, N, O and N. The calorimetric result,  $\Delta Hr_{298}^{\bullet} =$ -270.58 + 0.3 kcal/(10.811g B), becomes -270.9 ± ? depending on whether the correction for CHON impurities is estimated from (<u>42</u>) or (<u>45</u>). Since such estimates may give only lower limits to the correction, the data of Gross et al. (<u>44</u>) are not considered.

R21. 1.5  $F_2(g) + B(C, \beta) + BF_3(g) \Delta Hr_{298}^{\circ} = -271.65\pm0.22$  kcal/gfw Johnson et al. (42) remeasured the heat of combustion in fluorine using two different experimental techniques with a zone-refined sample of boron. The sample contained 0.13% impurities (0, C, N and H), for which the corrections amounted to about 0.12% of  $\Delta E_c^{\circ}$ . Experimental determination of the isotopic ratio in the sample gave an atomic weight of 10.805 ± 0.004 which was used in calculating  $\Delta Hr^{\circ}$ . This value remains essentially unchanged in terms of the accepted atomic weight of 10.811 for natural boron.

R22. 1.5  $F_2(g) + B(c, \beta) + BF_3(g)$   $\Delta Hr_{298}^{\bullet} = -271.03\pm0.51$  kcal/gfw Domalski and Armstrong (<u>45</u>) report the combustion in fluorine of pelleted mixtures of boron and Teflon. The boron contributed roughly 35% of the energy. Corrections for impurities in the boron (0.12% metallic impurities and 0.20% C, 0 and N) amounted to about 0.24% of  $\Delta E_c^{\bullet}$  (boron). Experimental determination of the isotopic ratio gave 10.812  $\pm$  0.005; thus, the accepted atomic weight of 10.811 was used in calculating  $\Delta Hr^{\bullet}$ . The authors' estimate of the overall experimental uncertainty is accepted instead of a value based only on random error (<u>19</u>).

R23. 3HF(50 H<sub>2</sub>0) + B(c, β) + 0.75 0<sub>2</sub>(g) + BF<sub>3</sub>(g) + 1.5 H<sub>2</sub>0(*t*)  $\Delta Hr_{298}^{\bullet} = -142.77\pm0.5 \text{ kcal/gfw}$ Gunn (<u>46</u>) reported  $\Delta Hr_{298}^{\bullet} = -28.29 \pm 0.07 \text{ kcal/gfw}$  for the reaction BF<sub>3</sub>(g) + 15.67 HF(3.747 H<sub>2</sub>0) + [solution], while Good and Mansson (<u>47</u>) reported  $\Delta Hr_{298}^{\bullet} = -173.406 \pm 0.2 \text{ kcal/gfw}$  for B(c, β) + 0.75 0<sub>2</sub>(g) + 18.57 HF(3.065 H<sub>2</sub>0) + [solution] +

1.5  $H_2O($)$ , where the resulting solutions have the same composition. A reduction to  $HF(3.747 H_2O)$  of  $\pm 1.624$  kcal/gfw with an estimated uncertainty of 10 to 25%, or roughly 0.2 to 0.4 kcal/gfw, is applied to the latter reaction. The difference between the two reactions and a reduction to  $HF(50 H_2O)$  of  $\pm 0.717 \pm 0.1$  kcal/gfw are used to obtain the result given above. The overall uncertainty is estimated as 0.3 to 0.5 kcal/gfw, depending on the choice of uncertainty for reduction to  $HF(3.747 H_2O)$ .

#### Input Data

The computer program (<u>48</u>) is a generalized version which allows  $\Delta Hf_{298}^{\bullet}$ ,  $\Delta Gf_{298}^{\bullet}$  and  $S_{298}^{\bullet}$  as variables; however, only  $\Delta Hf_{298}^{\bullet}$ is involved in this study. The program performs two functions: it edits the observations and then obtains the least squares adjustment. Editing consists of transforming the observations from the form given previously into the system of linear equations ready for solution.

The input to the program consists of (i) species with variables, (ii) species with fixed values, (iii) observations, and (iv) constraints, if any. There are seven variables in this study, namely, the values of  $\Delta Hf_{298}^{*}$  for HF(g), HF(50 H<sub>2</sub>0), NaF(c), BF<sub>3</sub>(g), NF<sub>3</sub>(g), CF<sub>4</sub>(g) and C<sub>2</sub>F<sub>4</sub>(polymer). Species with fixed values of  $\Delta Hf_{298}^{*}$  are listed in Table I. These include nine elements or ions in their standard reference states and eleven compounds with assumed fixed enthalpies of formation. The twenty three observations, each consisting of the reaction, enthalpy of reaction, and uncertainty, are summarized\* in Table II. The last six columns in the table indicate variations in the observations which were used in six preliminary adjustments. No constraints are involved.

The edited observations, shown in Table III, summarize the relationships among the variables for the final adjustment (HF-7). All 'fixed' species have been removed by substitution of their heats of formation and uncertainties. All uncertainties are assumed to be independent and are combined by use of Equation 17 of Rossini (<u>19</u>). The resulting values of  $R_{\underline{i}}+U_{\underline{i}}$ , involving only variables, provide the most direct comparison among the observations. Table III is also the most concise representation of the input data. It is the starting point for both sequential and simultaneous adjustments.

<sup>\*</sup>Table II is an example of a catalog of observed thermochemical processes. Armstrong (12) has suggested the value of such catalogs. They summarize the input data used for adjustment, either sequential or simultaneous.

#### Results of the Simultaneous Adjustments

Results of the final adjustment (HF-7) and six preliminary adjustments (HF-1 through HF-6) are given in Table IV. Also shown is the least sum adjustment of HF-7. Below each least squares value of  $\Delta Hf_{298}^{\bullet}$  is the corresponding standard error (54) based on the overall consistency rather than the internal consistency of each observation. This point will be emphasized in the discussion.

The preliminary adjustments represent various stages in the review of the input data as noted in Table II. All preliminary adjustments used our tentative estimate of 0.05 kcal/gfw for the uncertainty of R3 rather than the proper value (23) of 0.10 kcal/ gfw. The effect of this change is insignificant (cf. HF-7 and HF-6 in Table IV). Adjustments prior to HF-6 do not include R7, the heat of neutralization from Kolesov and Skuratov (28). Inclusion of this observation (cf. HF-7 and HF-4) changes the values of both HF(g) and NaF(c) by about +0.07 kcal/gfw and reduces the standard error of the latter from 0.11 to 0.07. Other changes are insignificant. It is noted in Rl2 that the Teflon combustions of Wood et al. (34) and Armstrong (12, 13) differ by 1 kcal/gfw or twice the sum of their uncertainties. When the value of Wood is substituted for that of Armstrong (cf. HF-5 and HF-4),  $C_{2}F_{\mu}$ (polymer) is changed by +0.7 kcal/gfw and  $CF_{\mu}(g)$  by +0.08 kcal/ gfw. Adjustments prior to HF-4 use 0.3 rather than 0.5 kcal/gfw as the estimated uncertainty for R23. The only significant effect of this is to shift BF<sub>3</sub>(g) from -271.42 to -271.29 kcal/gfw. Adjustments HF-3 and HF-2 indicate the effects of different combinations of R12, R12a and R12b. Omission of R6 and R19 has negligible effect (cf. HF-2 and HF-1), since these observations are relatively consistent but uncertain.

Not listed in Table IV are several other adjustments in which  $\phi_L$  for HF(50 H<sub>2</sub>0) was changed by ±0.2 kcal/gfw and the uncertainty was reduced from 0.3 to 0.2 kcal/gfw. These results are not significantly different from those already tabulated.

-18-

#### Discussion

There is an excellent precedent for use of the simultaneous method in adjustments. The fundamental constants of physics and chemistry, formerly treated via sequential adjustment by Birge (55), have been obtained for over two decades by simultaneous adjustment with the criterion of least squares (56, 57, 58). Experience derived from these studies of the physical constants will serve as the basis of the following discussion.

The species with fixed values in Table I correspond roughly to the "auxiliary constants" of physics (58). In this classification are included standard reference states, which are zero by definition, and experimental values of relatively high precision. Since some of these experimental values have uncertainties comparable with the data to be analyzed, both the fixed values and their uncertainties are substituted into the observations.

Error assignments must be expressed on as comparable a basis as possible, since they are to be used in weighting the observations. Assignments for the HF system are based on the random uncertainty supplemented in some cases by estimates of the systematic uncertainty. Reliable estimates of the latter appear to be particularly important in the present system due to the common occurrence of side reactions. It must be emphasized that the observations are usually a combination of at least three measurements including an energy calibration, an energy determination, and an analysis of the amount of chemical reaction.

Cohen and DuMond  $(\underline{58})$  stress the importance of overdetermination in testing the observations. They suggest that consistency of the data as a whole is the ultimate test and that this test becomes more searching with each new path of measurement. It is this sort of testing which revealed the bias in early data for HF and prompted the new data analyzed herein. Over-determination in this system remains extensive even after elimination of those experiments which are probably biased. The selected observations include only two variables which are

-19-

involved in as few as three different experiments. The remainder are involved in from five to eight experiments.

Birge's ratio test may be used as a measure of the overall consistency of the data. Cohen and DuMond (58) note that this ratio consists of the quotient of the external or <u>a posteriori</u> error with the internal or <u>a priori</u> error. The <u>a priori</u> error is derived from the uncertainties of the individual observations, while the <u>a posteriori</u> error is based on their deviations from the adjustment. When the ratio is close to one, there is no strong evidence of the presence of systematic error. The ratio is evaluated from:

$$(Birge's ratio)^2 = (Chi)^2/(m-n)$$
 [4]

and

A PROPERTY OF THE CONST. CONST. IN STR. 19 6

3 # 2 # 4 B" ( A 1 3 # 5 1 1 1 1 1 1

$$(Chi)^{2} = \sum_{i=1}^{m} (E_{i}'/U_{i})^{2},$$
 [5]

where  $E_i'/U_i$  is the weighted deviation or normalized residual.

Results of the ratio test are summarized in Table V. It may be concluded that there is no strong evidence of systematic error in HF-7, HF-6 and HF-4. Larger ratio values in the other adjustments suggest the possibility of systematic errors or underestimates of the <u>a priori</u> uncertainties. Sharp increases in the ratio occur with the inclusion of Rl2a and Rl2b due to their mutual inconsistency. In such cases, Cohen and DuMond (<u>58</u>) recommend the rejection of one or both observations upon location of a physical cause of unreliability in the experiment. Thus Rl2b is rejected without a probable physical cause, since our estimate of the uncertainty is based on inadequate information. Likewise, Rl2a is replaced with Rl2, which involves the later (and presumably more reliable) estimate of the uncertainty.

Individual deviations of the observations from the final adjustments by least squares and least sum are shown in the last four columns of Table III. Note that least sum assigns zero deviations to seven observations: R3, R7, R9, R12, R14, R18 and R20. These same observations in least squares have small but non-zero deviations, the largest magnitude being 0.18 kcal/gfw for R20.

The largest weighted deviations are -2.21 (R16), +2.04 (R1), +1.87 (R17) and + 1.55 (R23). Most of the inconsistency is concentrated in these four observations. R1 is the direct measurement of HF(50  $H_2$ 0). R16, R17 and R23 are quite sensitive to HF(50  $H_2$ 0) and they tend toward a more negative value than the direct measurement. The strain would be partially relieved by increasing the uncertainty of R1 to 0.09 kcal/gfw, which includes a reasonable estimate for systematic uncertainty (20). Lack of comparable estimates for R16, R17 and R23 makes this undesirable.

Sequential adjustments of the observations are not attempted in this paper. Interested readers may use Table III to make their own adjustments. No two individuals will get the same answers, since the number of possible paths to each variable is guite large.

As stated by DuMond and Cohen  $(\underline{58})$ , simultaneous adjustment is the logical approach in situations where there is considerable over-determination. However, least squares is not the only logical criterion in a particular situation. Least sum considers all of the data, yet provides a logical means of selecting a preferred subset just sufficient to determine the constants. DuMond and Cohen ( $\underline{58}$ ) imply, to the contrary, that such a selection must ignore all of the other data. Least sum is also less sensitive to outlying observations than is least squares. It is true that a least sum adjustment is not necessarily unique. There may be multiple solutions in a particular case, but this has not appeared in our systems. It is not likely in cases with considerable overdetermination. We conclude that least sum is complementary with least squares and that the two should be used together whenever practical.

There is little reason to prefer one criterion over the other in the HF system. The only significant difference is that least sum agrees slightly better with the direct determinations of  $BF_2(g)$ . Preference for least sum could be based on the

-21-

likelihood of outlying observations in this system. The difference is so small, however, that the least squares adjustment is preferred.

The selected values are summarized in Table VI, along with overall uncertainties estimated from twice the standard errors of the least squares adjustment. The user should be warned that the simple formula for propagation of errors (<u>19</u>) is only the first approximation when two or more of the selected values are combined in calculation of a third quantity. Correlation of errors is likely among some of the selected values; giving rise to additional terms, which may be either positive or negative, in the general formula for propagation of errors (<u>58</u>, <u>59</u>). Interpretation of the derived uncertainties is further complicated by the input data being weighted in some cases by inclusion of reasonable bounds for the systematic error.

Values of  $\Box Hf_{298}^{\bullet}$  for HF(n H<sub>2</sub>0) should be obtained from that for HF(50 H<sub>2</sub>0) using differences taken from the dilution curve of Parker (<u>17</u>). Additional error should be small, except for n > 1000. The values selected for HF(50 H<sub>2</sub>0) and NaF(c) imply that  $\phi_{L}$  for HF(50 H<sub>2</sub>0) is about 0.15 ± 0.2 kcal/gfw less positive than tabulated by Parker (<u>17</u>). This difference arises partly from the heat of neutralization (<u>28</u>) and partly from other data.

The analysis reveals some paths which should receive further attention. Since NaF is so closely linked with HF(aq), it is most surprising that good, modern data are not available for the heat of neutralization of NaOH with HF or for the heat of dilution of NaF. Parker's estimate (<u>17</u>) for the latter is reasonable, but it should be verified by experimental data. New measurements of the heat of dilution of HF are also desirable in dilute solutions, and would be an accurate way of testing the calculated  $\phi_L$  curve (cf. earlier discussion for R8). It has already been stressed that links such as R16 are very sensitive for testing the enthalpy of formation of HF(50 H<sub>2</sub>O). Other links would add to the test. Finally, a thorough analysis of the data for HF(real gas) could

-22-

and a second second

remove the non-ideality corrections as a source of uncertainty for HF(g).

#### Acknowledgment

The National Standard Reference Data System, National Bureau of Standards, and the U. S. Department of Commerce supported the development of simultaneous adjustment, upon which this paper is based. The least squares computer program was prepared under this support.

#### References

State and the second state and

Stanks .

· · · · · · · ·

 F. R. Bichowsky and F. D. Rossini, "The Thermochemistry of Chemical Substances," Reinhold Publishing Corp., New York, 1936.

- F. D. Rossini et al., "Selected Values of Chemical Thermodynamic Properties," National Bureau of Standards Circular 500, U. S. Government Printing Office, Washington, D. C., 1950.
- 3. D. D. Wagman et al., "Selected Values of Chemical Thermodynamic Properties," NBS Technical Note 270-1, 1965, and NBS Technical Note 270-3, 1968.
- 4. J. R. Rice, "The Approximation of Functions," Vol. I, Chap. 1, Addison-Wesley Publishing Co. Reading, Mass., 1964.
- 5. A. Syverud and M. Klein, Progress Reports I, II, III, IV under National Bureau of Standards Contract CST-1165, The Dow Chemical Co., Midland, Mich., 1964-1965.
- 6. J. G. Hust and R. D. McCarty, Cryog., 7, 200 (1967).
- G. Forsythe and C. B. Moler, "Computer Solution of Linear Algebraic Systems," Prentice-Hall, Inc., Englewood Cliffs, N. J., 1967.
- A. Ralston and H. S. Wilf, "Mathematical Methods for Digital Computers," John Wiley & Sons, Inc., New York, 1964.
- A. Ralston and H. S. Wilf, "Mathematical Methods for Digital Computers," Vol. II, John Wiley & Sons, Inc., New York, 1967.
- 10. Ref. (8), Chapter 25.
- N. P. Loomba, "Linear Programming," McGraw-Hill Book Company, New York, 1964.
- 12. G. T. Armstrong, U. S. Nat. Bur. Stand., Report 9803, 1-18 (1968).
- E. S. Domalski and G. T. Armstrong, J. Res. NBS, <u>71A</u>, 105-118 (1967).
- J. R. Lacher and H. A. Skinner, J. Chem. Soc., <u>A1968</u>, 1034-8.

and the second second

- E. Rudzitis, private communication, parts IV and V of "Thermodynamic Properties of Hydrogen Fluoride," Argonne National Laboratory, 1963 and 1965.
- G. T. Armstrong and L. A. Krieger, U. S. Nat. Bur. Stand. Report 7192, 92-203 (1961).
- 17. V. B. Parker, "Thermal Properties of Aqueous Uni-Univalent Electrolytes," NSRDS-NBS 2, 1965.
- JANAF Thermochemical Tables, The Dow Chemical Co., Midland, Mich., December 1968.
- F. D. Rossini, Chap. 14, "Experimental Thermochemistry," Interscience Publishers, Inc., New York, 1956.
- 20. R. C. King and G. T. Armstrong, J. Res. NBS, <u>72A</u>, 113-131 (1968) and NBS Report 9500, 140-226 (1967).
- J. W. C. Johns and R. F. Barrow, Proc. Roy. Soc. (London), <u>A251</u>, 504-518 (1959).
- 22. W. W. Rodenburg and C. E. Vanderzee, paper presented at the 22nd Annual Calorimetry Conference, Thousand Oaks, Calif., June 1967.
- C. E. Vanderzee, private communication, Univ. of Nebraska, Jan. 23, 1969.
- 24. L. G. Hepler, W. L. Jolly and W. M. Latimer, J. Amer. Chem. Soc., <u>75</u>, 2809 (1953); W. M. Latimer and W. L. Jolly, ibid., 1548 (1953); D. H. Davies and G. C. Benson, Can. J. Chem., <u>43</u>, 3100 (1965).
- J. P. Coughlin, J. Amer. Chem. Soc., <u>80</u>, 1802 (1958); <u>79</u>, 2397 (1957).
- 26. G. C. Hood and M. M. Woyski, J. Amer. Chem. Soc., <u>73</u>, 2738-41 (1951).
- H. von Wartenberg and O. Fitzner, Z. Anorg. Allgem. Chem., <u>151</u>, 313-25 (1926).
- V. P. Kolesov and S. M. Skuratov, Russ. J. Inorg. Chem. (English Transl.), <u>6</u>, 889-91 (1961).
- 29. J. D. Cox and D. Harrop, Trans. Faraday Soc., 61, 1328 (1965).

30. A. J. Ellis, J. Chem. Soc. <u>1963</u>, 4300-4; H. N. Farrer and F. J. C. Rossotti, J. Inorg. Nucl. Chem., <u>26</u>, 1959 (1964);
L. G. Sillen and A. E. Martell, "Stability Constants,"
Special Pub. No. 17, pp. 256-7, The Chemical Society, London, 1964; T. Erdey-Gruz, L. Majthenye and E. Kugler, Acta Chim. Acad. Sci. Hung., <u>37(4)</u>, <u>393-403</u> (1963).

a construction of the second second

-1.48Ta

"Y THE REAL PROPERTY OF THE PARTY OF

- 31. L. G. Hepler, W. L. Jolly and W. M. Latimer, J. Amer. Chem. Soc., <u>75</u>, 2809 (1953); K. L. Elmore, J. D. Hatfield,
  C. M. Mason and A. D. Jones, ibid., <u>71</u>, 2710 (1949);
  H. H. Broene and T. DeVries, ibid., <u>69</u>, 1644 (1957);
  C. B. Wooster, ibid., <u>60</u>, 1609 (1938); W. A. Roth, Ann., 542, 35 (1939).
- 32. E. Greenberg and W. N. Hubbard, J. Phys. Chem., <u>72</u>, 222-7 (1968).
- W. D. Good, D. W. Scott and G. Waddington, J. Phys. Chem.,
   <u>60</u>, 1080 (1956); D. W. Scott, W. D. Good and G. Waddington,
   J. Amer. Chem. Soc., 77, 245 (1955).
- 34. J. L. Wood, R. J. Lagow and J. L. Margrave, J. Chem. Eng. Data, <u>12</u>, 255 (1967).
- 35. G. M. Martin and R. K. Eby, J. Res. NBS, 72A, 467-70 (1968).
- 36. J. D. Cox, H. A. Gundry and A. J. Head, Trans. Faraday Soc., 61, 1594-1600 (1965).
- 37. L. C. Walker, paper presented at the 22nd Annual Calorimetry Conference, Thousand Oaks, Calif., June 1967.
- 38. G. C. Sinke, J. Chem. Eng. Data, 10, 295-6 (1965).
- 39. L. C. Walker, J. Phys. Chem., 71, 361-3 (1967).
- 40. G. C. Sinke, J. Phys. Chem., 71, 359-60 (1967).
- J. R. Ludwig and W. J. Cooper, J. Chem. Eng. Data, <u>8</u>, 76 (1963).
- 42. G. K. Johnson, H. M. Feder and W. N. Hubbard, J. Phys. Chem., 70, 1-6 (1966).
- S. S. Wise, J. L. Margrave, H. M. Feder and W. N. Hubbard, J. Phys. Chem., 65, 2157 (1961).

- 44. P. Gross, C. Hayman, D. L. Levi and M. C. Stuart, Fulmer Res. Inst. Report R.146/4/23, November 1960.
- E. S. Domalski and G. T. Armstrong, J. Res. NBS, <u>71A</u>, 195-202 (1967).
- 46. S. R. Gunn, J. Phys. Chem., <u>69</u>, 1010-5 (1965).
- 47. W. D. Good and M. Mansson, J. Phys. Chem., 70, 97-104 (1966).
- 48. The computer program was prepared under Contract No. CST-1165, National Bureau of Standards, U. S. Dept. of Commerce.
- 49. Value assigned in this study.
- S. R. Gunn and L. G. Green, J. Chem. Eng. Data, <u>8</u>, 180 (1963);
   S. R. Gunn, UCRL-7992, Livermore, Calif., July 29, 1964.
- 51. C. E. Vanderzee and J. D. Nutter, J. Phys. Chem., <u>67</u>, 2521 (1963).
- 52. S. R. Gunn, J. Phys. Chem., 71, 1386 (1967).
- 53. P. A. G. O'Hare, J. L. Settle and W. N. Hubbard, Trans. Faraday Soc., 62, 558 (1966).
- 54. Ref. 8, Chapter 17.
- 55. R. T. Birge, Rev. Mod. Phys., 1, 1 (1929).
- 56. F. G. Dunnington, Rev. Mod. Phys., 11, 68 (1939).
- 57. E. R. Cohen, K. M. Crowe and J. W. M. DuMond, "The Fundamental Constants of Physics," Interscience Publishers, Inc., New York, 1957.
- 58. E. R. Cohen and J. W. M. DuMond, Rev. Mod. Phys., <u>37</u>, 537 (1965).
- 59. Ref. 57, Chapter 7.

		TABI	LEI		
Species	with	Fixed	Values	of	LHF <sup>•</sup> 298

MILSON AS IN SURGE

Species	<sup>4</sup> Hf <sup>2</sup> 98, k	cal/gfw	Reference
B(c, β)	0.0	± 0.0	Defined
C(graphite)	0.0	± 0.0	Defined
Cl <sub>2</sub> (g)	0.0	± 0.0	Defined
C0, (g)	-94.051	± 0.030	( <u>3</u> )
$C_2 \tilde{N}_2(g)$	73.84	± 0.43	( <u>3</u> )
F(g)	18.86	± 0.3	( <u>18, 49</u> )
F,(g)	0.0	± 0.0	Defined
H(g)	52.095	± 0.002	( <u>3</u> )
HCl(g)	-22.062	± 0.02	( <u>3</u> )
HCl(aq)	-39.932	± 0.02	( <u>49, 50, 51</u> )
H <sup>+</sup> (aq)	0.0	± 0.0	Defined
H <sub>2</sub> (g)	0.0	± 0.0	Defined
H_0(1)	-68.315	± 0.01	( <u>3</u> )
N_(g)	0.0	± 0.0	Defined
NaCl(c)	-98.32	± 0.06	( <u>49, 52</u> )
NaOH(aq)	-112.44	± 0.04	( <u>49, 52</u> )
Na <sup>+</sup> (aq)	-57.47	± 0.04	( <u>49, 52</u> )
0,(g)	0.0	± 0.0	Defined
S(c, rh)	0.0	± 0.0	Defined
SF <sub>6</sub> (g)	-291.77	± 0.24	( <u>53</u> )

| 1                                |   |  |   |  |   
   
  |   |  |  |  |   |                     |   |  |  
  |   |   |  
   |   |   |   |  |   |   |   |  
  |
|----------------------------------|---|--|---|--
--
--|---|--|--
--|---|---------------------|---|--
--
---|---|---|--|---
---|---|--|---|---|---|---|
| HF-1                             |   |  | ±0.05   |  |   
   
  | omit  | omit   |  |  |   |                     |   | 12a-12b  |  
  |   | than .  |  
   |   |   | omit  |  |   |   |   | ±0.3   
  |
| HF-2                             |   |  | ±0.05   |  |   
   
  |   | omit   |  |  |   |                     |   | 12 <b>4.1</b> 2b                                       | •  
  | tion 12b  | .4 rather   |  
   |   |   |   |  |   |   |   | £0.3   
  |
| HF-3                             |   |  | ±0.05   |  |   
   
  |   | omit   |  |  |   |                     |   | 12.12b*  | •  
  | for Read  | gned as   | -  
   |   |   |   |  |   |   |   | ±G.3   
  |
| HF-4                             |   |  | ±0.05   |  |   
   
  |   | omit   |  |  |   |                     |   |  |  
  | rtaintv   | 15 3551   | cal/ gru   
   |   |   |   |  |   |   |   |  
  |
| HF-5                             |   |  | ±0.05   |  |   
   
  |   | omit   |  |  |   |                     |   | 12b  |  
  | *Unce   | HF-3  |  
   |   |   |   |  |   |   |   |  
  |
| HF-6                             |   |  | ±0.05   |  |   
   
  |   |  |  |  |   |                     |   |  |  
  |   |   |  
   |   |   |   |  |   |   |   |  
  |
| JHf<br>kca <b>198</b> fw<br>HF-7 | -76.68±0.05   | 136.051±0.3  | -28.24±0.10   | -2.41±0.07   | 3.81±0.25   
   
  | -39.30±0.4  | -16.57±0.05  | 3.414±0.3  |  |   | -160.6220.9         | -118.8±0.5  | -247.85±0.3  |  
  | -41.506±0.3   | -1308.8±1.3   | -199.4±0.22  
   | -228.26±0.25  | 31.44±0.3   | -239.4621.2   | -271.6±0.9   | -271.65±0.22  | -271.03±0.51  |   | -142.77±0.5  
  |
| ło. Reaction Adjustment No. →    | 1. 0.5 H <sub>2</sub> (g)+0.5 F <sub>2</sub> (g) + HF(50 H <sub>2</sub> 0)              | 2. HF(g) + H(g) + F(g)   | 3. $HF(g)+NaOH(aq) + NaF(c) + H_{2}O(f)$  | <pre>4. HF(50 H<sub>2</sub>0)+NaCl(c) + NaF(c)+HCl(aq)</pre>   | 5. $HF(g)+NaCl(c) + NaF(c) + HCl(g)$  
   
  | <pre>6. 0.5 F<sub>2</sub>(g)+NaCl(c) + NaF(c)+0.5 Cl<sub>2</sub>(g)</pre>   | 7. HF(50 H <sub>2</sub> 0)+NaOH(aq) + NaF(c)+H <sub>2</sub> O(1) | <b>B</b> : $H^{+}(aq) + NaF(c) + HF(50 H_{2}0) + Na^{+}(aq)$ | 9. C(graph)+2F <sub>9</sub> (g) + CF <sub>h</sub> (g) -223.05±0.18 | 10. C <sub>2</sub> F <sub>k</sub> (polymer)+0,(g)+2H,0(f) + | HHF(50 H,0)+2CO,(g) | $(1. C_2F_k(polymer)+0_2(g) + CF_k(g)+C0_2(g)$  | 12. $C_2F_4(\text{polymer})+2F_2(g) + 2CF_4(g)$        | 13. C(graph)+2F <sub>2</sub> (g) + CF <sub>k</sub> (g) -222.87±0.38  
  | U4. CF <sub>4</sub> (g)+2H <sub>2</sub> O(g) + WHF(50 H <sub>2</sub> 0)+CO <sub>2</sub> (g)   | $(5. 8NF_3(g)+3C_3N_2(g) + 6CF_4(g)+7N_3(g)$  | L6. NF3(g)+1.5 H2(g) + 2HF(50 H20)+0.5N2(g)  
   | $(7. 2NF_{3}(g)+S(c, rh) + SF_{6}(g)+N_{2}(g)$  | $(6. NF_{3}(g) + 0.5 N_{2}(g) + 1.5 F_{2}(g)$   | (9. NF <sub>3</sub> (g)+B(c, p) + BF <sub>3</sub> (g)+0.5N <sub>2</sub> (g)   | :0. 1.5 F <sub>2</sub> (g)+B(c, β) + BF <sub>3</sub> (g)   | $1. 1.5 F_2(g) + B(c, p) + BF_3(g)$   | 2. 1.5 F <sub>2</sub> (g)+B(c, p) + BF <sub>3</sub> (g)   | 3. 3HF(50 H,0)+B(c, B)+0.75 0,(g) +   | BF <sub>3</sub> (g)+1.5 H <sub>2</sub> O(£)  
  |
|                                  | JHf 2985<br>kca1985<br>No. Reaction Adjustment No. → HF-7 HF-6 HF-5 HF-4 HF-3 HF-2 HF-1 | Mo. Reaction Adjustment No. + HE-7 HF-6 HF-5 HF-4 HF-3 HF-2 HF-1<br>1. 0.5 H <sub>2</sub> (g)+0.5 F <sub>2</sub> (g) + HF(50 H <sub>2</sub> 0) -76.68±0.05 | Mo.         Reaction         Adjustment No. +         HF-7         HF-6         HF-5         HF-3         HF-2         HF-1           1.         0.5         H_2(g)+0.5         F_2(g)         HF(50         H_20)         -76.68±0.05           2.         HF(g)         H(g)         F(g)         136.051±0.3         136.051±0.3 | Mo.     Reaction     Adjustment No. +     JHF-9       1.     0.5     H_2(g)+0.5     F_2(g) + HF(50     H_20)       2.     HF(g) + H(g) + F(g)     136.051±0.3       3.     HF(g)+NaOH(aq) + NaF(c) + H_20(g)     -28.24±0.10     ±0.05     ±0.05     ±0.05 | MC.       Reaction       Adjustment No. +       HF-7       HF-6       HF-5       HF-4       HF-3       HF-2       HF-1         1. 0.5 H <sub>2</sub> (g)+0.5 F <sub>2</sub> (g) + HF(50 H <sub>2</sub> 0)       -76.68±0.05       -76.68±0.05       136.051±0.3       3         2. HF(g) + H(g) + F(g)       136.051±0.3       136.051±0.3       3       40.05       ±0.05 </td <td>Mo.ReactionAdjustment No. +<math>JF_7^{29}</math><br/>kcal?9<br/>kcal?9<br/>HF-7HF-6HF-5HF-4HF-3HF-2HF-11. 0.5 H_2(g)+0.5 F_2(g) + HF(50 H_20)-76.68±0.05-76.68±0.05136.051±0.3136.051±0.32. HF(g) + H(g) + F(g)136.051±0.3136.051±0.3136.051±0.340.05<math>±0.05</math><math>±0.05</math><math>±0.05</math><math>±0.05</math><math>±0.05</math>3. HF(g)+NaCH(aq) + MaF(c) + H_2O(8)-28.24±0.10<math>±0.05</math><math>±0.05</math><math>±0.05</math><math>±0.05</math><math>±0.05</math><math>±0.05</math><math>±0.05</math>4. HF(50 H_2O)+NaCl(c) + NaF(c) + HCl(aq)-2.41±0.073.61±0.25<math>±0.16c</math><math>±0.16c</math><math>±0.05c</math><math>±0.05</math><math>±0.05</math><math>±0.05</math><math>±0.05</math></td> <td></td> <td></td> <td></td> <td></td> <td>JHF 29<br/></td> <td>O.         Reaction         Adjustment No. +         <math>\frac{JHF^2}{Kcall9}</math>         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + 0.5 F_2(g) + HF(50 H_2O)         -76.68±0.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         136.051±0.3         136.051±0.3         136.051±0.3         HF-2         HF-1           3.         HF(g) + H(g) + F(g)         136.051±0.3         136.051±0.3         136.051±0.3         100.5         ±0.05</td> <td><math display="block"> \begin{array}{c ccccccccccccccccccccccccccccccccccc</math></td> <td>Oc.         Reaction         Adjustment No.         <math>HF-5</math> <math>HF-5</math> <math>HF-3</math> <math>HF-2</math> <math>HF-1</math>           1.         0.5         <math>H_c(g) + F(g)</math> <math>HF(50 H_2O)</math> <math>-76.6820.05</math> <math>HF-5</math> <math>HF-3</math> <math>HF-2</math> <math>HF-1</math>           2.         <math>HF(g) + H(g) + F(g)</math> <math>136.05140.3</math> <math>136.05140.3</math> <math>136.05140.3</math> <math>136.05140.3</math>           3.         <math>HF(g) + H(g) + F(g)</math> <math>136.05140.3</math> <math>28.24\pm0.10 \pm 20.05</math> <math>20.05 \pm 20.05</math> <math>20.05 \pm 20.05</math> <math>20.05</math>           4.         <math>HF(50 H_2O) + WaCl(c) + WaF(c) + HCl(aq)</math> <math>-2.44120.07</math> <math>-2.44120.07</math> <math>2.44120.07</math> <math>2.4120.07</math>           5.         <math>HF(50 H_2O) + WaCl(c) + WaF(c) + HCl(aq)</math> <math>-2.44120.07</math> <math>-2.44120.07</math> <math>00005 \pm 20.05 \pm 20.05</math> <math>20.05</math> <math>20.05</math><td>Oc.         Reaction         Adjustment No.         <math>\frac{JHF-3}{kca13}</math>         HF-6         HF-5         HF-3         HF-2         HF-1           1         0.5         <math>H_{\gamma}(g)^{+0.5}</math> <math>F(g)</math> <math>HF(50 H_{2}^{0})</math> <math>-76.6820.05</math> <math>HF-5</math> <math>HF-3</math> <math>HF-2</math> <math>HF-1</math> <math>HF-1</math></td><td>O.         Reaction         Milustment No. +         <math>\frac{M_{12}}{K_{ca}}</math>         HF-6         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_{2}(u) + H(g) + F(g)         HE(50 H_{2}(0))         -76.6820.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         Mark(c) + H_{2}(0(1))         -76.6820.05         20.05</td><td>No.         Reaction         Adjustment No.         <math>\frac{117}{12}</math>         HF-6         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + H(g)         F_2(g) + H(50 H_2O)         -76.6820.05         136.05120.3         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.05120.3         156.6820.05         156.05120.3         156.6820.05         100.05         20.05</td><td>MFG         MFG         MF-5         MF-4         MF-3         MF-2         MF-1           1         0.5 <math>H_{\zeta}(g) + 0.5 \Gamma_{\zeta}(g) + HT(50 H_{\zeta}0)</math>         -76.6820.05         MF-5         MF-4         MF-3         MF-2         MF-1           1         0.5 <math>H_{\zeta}(g) + M(g) + F(g)</math>         136.05120.3         136.05120.3         136.05120.3         20.05</td><td>No.         Reaction         Adjustment No.         <math>\frac{ME_1}{N-3}</math>         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + HE(50 H_2())         -76.6820.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         136.05120.3         136.05120.3         136.05120.3         20.05</td><td>O.         Reaction         Adjustment No.         <math>\frac{ME}{MT-3}</math>         HF-6         HF-5         HF-3         HF-2         HF-1           1         0.5 H<sub>2</sub>(g) + H(g) + F(g)         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         100.05         20.05</td><td>Model         Match         <math>\frac{116-5}{18-7}</math>         HF-5         HF-4         HF-3         HF-2         HF-1           1         0.5 H<sub>2</sub>(g) + K(g)         H2 (g) + K(g)         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.051         136.051         136.051         136.051         136.05         100.5         20.05         2</td><td>Mit of the section         Mature No.         <math>\frac{117}{10}</math>         Hr-6         Hr-5         Hr-3         Hr-2         Hr-1           1         0.5 H<sub>2</sub>(g<sup>1</sup>)0.5 F<sub>2</sub>(g) + Hr(50 H<sub>2</sub>(0)         -78.680.05         136.05140.3         136.05140.3         136.05140.3           2         Hr(g) + H(g)         Hr(g)         Hr(g)         136.05140.3         136.05140.3           3         Hr(g) + H(g)         Hr(g)         28.15(g)         136.05140.3         20.05</td><td>Million         Million         &lt;</td><td>Million         Million         &lt;</td><td>Mit Spin         Mit Spin</td></td> | Mo.ReactionAdjustment No. + $JF_7^{29}$<br>kcal?9<br>kcal?9<br>HF-7HF-6HF-5HF-4HF-3HF-2HF-11. 0.5 H_2(g)+0.5 F_2(g) + HF(50 H_20)-76.68±0.05-76.68±0.05136.051±0.3136.051±0.32. HF(g) + H(g) + F(g)136.051±0.3136.051±0.3136.051±0.340.05 $±0.05$ $±0.05$ $±0.05$ $±0.05$ $±0.05$ 3. HF(g)+NaCH(aq) + MaF(c) + H_2O(8)-28.24±0.10 $±0.05$ $±0.05$ $±0.05$ $±0.05$ $±0.05$ $±0.05$ $±0.05$ 4. HF(50 H_2O)+NaCl(c) + NaF(c) + HCl(aq)-2.41±0.073.61±0.25 $±0.16c$ $±0.16c$ $±0.05c$ $±0.05$ $±0.05$ $±0.05$ $±0.05$ |  |  |  |   | JHF 29<br>          | O.         Reaction         Adjustment No. + $\frac{JHF^2}{Kcall9}$ HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + 0.5 F_2(g) + HF(50 H_2O)         -76.68±0.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         136.051±0.3         136.051±0.3         136.051±0.3         HF-2         HF-1           3.         HF(g) + H(g) + F(g)         136.051±0.3         136.051±0.3         136.051±0.3         100.5         ±0.05 | $ \begin{array}{c ccccccccccccccccccccccccccccccccccc$ | Oc.         Reaction         Adjustment No. $HF-5$ $HF-5$ $HF-3$ $HF-2$ $HF-1$ 1.         0.5 $H_c(g) + F(g)$ $HF(50 H_2O)$ $-76.6820.05$ $HF-5$ $HF-3$ $HF-2$ $HF-1$ 2. $HF(g) + H(g) + F(g)$ $136.05140.3$ $136.05140.3$ $136.05140.3$ $136.05140.3$ 3. $HF(g) + H(g) + F(g)$ $136.05140.3$ $28.24\pm0.10 \pm 20.05$ $20.05 \pm 20.05$ $20.05 \pm 20.05$ $20.05$ 4. $HF(50 H_2O) + WaCl(c) + WaF(c) + HCl(aq)$ $-2.44120.07$ $-2.44120.07$ $2.44120.07$ $2.4120.07$ 5. $HF(50 H_2O) + WaCl(c) + WaF(c) + HCl(aq)$ $-2.44120.07$ $-2.44120.07$ $00005 \pm 20.05 \pm 20.05$ $20.05$ <td>Oc.         Reaction         Adjustment No.         <math>\frac{JHF-3}{kca13}</math>         HF-6         HF-5         HF-3         HF-2         HF-1           1         0.5         <math>H_{\gamma}(g)^{+0.5}</math> <math>F(g)</math> <math>HF(50 H_{2}^{0})</math> <math>-76.6820.05</math> <math>HF-5</math> <math>HF-3</math> <math>HF-2</math> <math>HF-1</math> <math>HF-1</math></td> <td>O.         Reaction         Milustment No. +         <math>\frac{M_{12}}{K_{ca}}</math>         HF-6         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_{2}(u) + H(g) + F(g)         HE(50 H_{2}(0))         -76.6820.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         Mark(c) + H_{2}(0(1))         -76.6820.05         20.05</td> <td>No.         Reaction         Adjustment No.         <math>\frac{117}{12}</math>         HF-6         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + H(g)         F_2(g) + H(50 H_2O)         -76.6820.05         136.05120.3         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.05120.3         156.6820.05         156.05120.3         156.6820.05         100.05         20.05</td> <td>MFG         MFG         MF-5         MF-4         MF-3         MF-2         MF-1           1         0.5 <math>H_{\zeta}(g) + 0.5 \Gamma_{\zeta}(g) + HT(50 H_{\zeta}0)</math>         -76.6820.05         MF-5         MF-4         MF-3         MF-2         MF-1           1         0.5 <math>H_{\zeta}(g) + M(g) + F(g)</math>         136.05120.3         136.05120.3         136.05120.3         20.05</td> <td>No.         Reaction         Adjustment No.         <math>\frac{ME_1}{N-3}</math>         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + HE(50 H_2())         -76.6820.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         136.05120.3         136.05120.3         136.05120.3         20.05</td> <td>O.         Reaction         Adjustment No.         <math>\frac{ME}{MT-3}</math>         HF-6         HF-5         HF-3         HF-2         HF-1           1         0.5 H<sub>2</sub>(g) + H(g) + F(g)         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         100.05         20.05</td> <td>Model         Match         <math>\frac{116-5}{18-7}</math>         HF-5         HF-4         HF-3         HF-2         HF-1           1         0.5 H<sub>2</sub>(g) + K(g)         H2 (g) + K(g)         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.051         136.051         136.051         136.051         136.05         100.5         20.05         2</td> <td>Mit of the section         Mature No.         <math>\frac{117}{10}</math>         Hr-6         Hr-5         Hr-3         Hr-2         Hr-1           1         0.5 H<sub>2</sub>(g<sup>1</sup>)0.5 F<sub>2</sub>(g) + Hr(50 H<sub>2</sub>(0)         -78.680.05         136.05140.3         136.05140.3         136.05140.3           2         Hr(g) + H(g)         Hr(g)         Hr(g)         136.05140.3         136.05140.3           3         Hr(g) + H(g)         Hr(g)         28.15(g)         136.05140.3         20.05</td> <td>Million         Million         &lt;</td> <td>Million         Million         &lt;</td> <td>Mit Spin         Mit Spin</td> | Oc.         Reaction         Adjustment No. $\frac{JHF-3}{kca13}$ HF-6         HF-5         HF-3         HF-2         HF-1           1         0.5 $H_{\gamma}(g)^{+0.5}$ $F(g)$ $HF(50 H_{2}^{0})$ $-76.6820.05$ $HF-5$ $HF-3$ $HF-2$ $HF-1$ | O.         Reaction         Milustment No. + $\frac{M_{12}}{K_{ca}}$ HF-6         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_{2}(u) + H(g) + F(g)         HE(50 H_{2}(0))         -76.6820.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         Mark(c) + H_{2}(0(1))         -76.6820.05         20.05 | No.         Reaction         Adjustment No. $\frac{117}{12}$ HF-6         HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + H(g)         F_2(g) + H(50 H_2O)         -76.6820.05         136.05120.3         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.6820.05         156.05120.3         156.6820.05         156.05120.3         156.6820.05         100.05         20.05 | MFG         MFG         MF-5         MF-4         MF-3         MF-2         MF-1           1         0.5 $H_{\zeta}(g) + 0.5 \Gamma_{\zeta}(g) + HT(50 H_{\zeta}0)$ -76.6820.05         MF-5         MF-4         MF-3         MF-2         MF-1           1         0.5 $H_{\zeta}(g) + M(g) + F(g)$ 136.05120.3         136.05120.3         136.05120.3         20.05 | No.         Reaction         Adjustment No. $\frac{ME_1}{N-3}$ HF-5         HF-4         HF-3         HF-2         HF-1           1.         0.5 H_2(g) + HE(50 H_2())         -76.6820.05         HF-6         HF-3         HF-2         HF-1           2.         HF(g) + H(g) + F(g)         136.05120.3         136.05120.3         136.05120.3         20.05 | O.         Reaction         Adjustment No. $\frac{ME}{MT-3}$ HF-6         HF-5         HF-3         HF-2         HF-1           1         0.5 H <sub>2</sub> (g) + H(g) + F(g)         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         100.05         20.05 | Model         Match $\frac{116-5}{18-7}$ HF-5         HF-4         HF-3         HF-2         HF-1           1         0.5 H <sub>2</sub> (g) + K(g)         H2 (g) + K(g)         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.05120.3         136.051         136.051         136.051         136.051         136.05         100.5         20.05         2 | Mit of the section         Mature No. $\frac{117}{10}$ Hr-6         Hr-5         Hr-3         Hr-2         Hr-1           1         0.5 H <sub>2</sub> (g <sup>1</sup> )0.5 F <sub>2</sub> (g) + Hr(50 H <sub>2</sub> (0)         -78.680.05         136.05140.3         136.05140.3         136.05140.3           2         Hr(g) + H(g)         Hr(g)         Hr(g)         136.05140.3         136.05140.3           3         Hr(g) + H(g)         Hr(g)         28.15(g)         136.05140.3         20.05 | Million         < | Million         < | Mit Spin         Mit Spin |

A. B. Sugar

287

ALLY MADE DO THE

TABLE II

AFRPL-TR-69-70

10.30

TAB	LE III. <u>Relationships among</u>	the Variables	as Obtained from	the Input	for the	Final Adi	ustment(HF-7)
			R; + U;	Dev.	Dev./U;	Dev.	Dev./U;
	Relationships among the ${}_{2}\mathrm{Hf}_{2}^{\bullet}$	98 Variables	-kcal/gfu	Least	Squares	Least	Sum
1.	HF(50 H <sub>2</sub> 0)		-76.680±0.050	0.102	2.035	0.104	2.075
2.	-HF(g)		65.096±0.424	-0.040	-0.094	-0.018	-0.042
ë.	Naf(c) - Hf(g)		-72.365±0.108	0.015	0.135	0.0	0.0
æ	Naf(c) - Hf(50 H <sub>2</sub> 0)		-60.798±0.094	-0-064	-0.680	-0.103	-1.096
s.	NaF(c) - $HF(g)$		-72.448±0.258	-0.068	-0.265	-0.083	-0.322
<b>e</b> .	NaF(c)		-137.620±0.404	-0.104	-0.258	-0.141	-0.350
٦.	<b>Naf</b> (c) - HF(50 H <sub>2</sub> 0)		-60.695±0.065	0.039	0.600	0.0	0.0
<b></b>	-[NaF(c) - HF(50 H <sub>2</sub> 0)]		60.884±0.303	0.150	0.496	0.189	0.624
	CF4 (g)		-223.050±0.180	-0.015	-0.083	0.0	0.0
ю.	4 × HF(50 H20) - C2F4 (polym	er)	-109.148±0.902	-0.239	-0.265	-0.263	-0.292
н.	CF4(g) - C2F4(polymer)		-24.749±0.501	0.068	0.136	0, 051	0.102
12.	2 x CF <sub>4</sub> (g) - C <sub>2</sub> F <sub>4</sub> (polymer)		-247.850±0.300	0.002	0.007	0.0	0.0
13.	CF4 (g)		-222.870±0.380	0.165	0.434	0.180	0.474
14.	4 × HF(50 H <sub>2</sub> 0) - CF <sub>4</sub> (g)		-84.085±0.302	0.007	0.023	0.0	0.0
15.	6 x CF <sub>4</sub> (g) - 8 x NF <sub>3</sub> (g)		-1087.280±1.831	-0.516	-0.282	-0.500	-0.273
16.	3 × HF(50 H <sub>2</sub> 0) - NF <sub>3</sub> (g)		-199.400±0.220	-0.486	-2.207	-0.489	-2.222
17.	-2 x NF <sub>3</sub> (g)		63.510±0.347	0.648	1.071	0.630	1.016
19.	-NF <sub>3</sub> (g)		31.440±0.300	0.009	0.030	0.0	0.0
19.	BF <sub>3</sub> (g) - NF <sub>3</sub> (g)		-239.460±1.200	0.529	144.0	0.700	0.583
20.	BF <sub>3</sub> (g)		-271.600±0.900	-0.180	-0.200	0.0	0.0
21.	BF <sub>3</sub> (g)		-271.650±0.220	-0.230	-1.045	-0.050	-0.227
22.	BF <sub>3</sub> (g)		-271.030±0.510	0.390	0.765	0.570	1.118
23.	<b>BF<sub>3</sub>(g)</b> - 3 × HF(50 H <sub>2</sub> 0)		-40.297±0.500	0.777	1.554	0.952	1.904

.1

Sum of squares = 18.436 \*Dev. =  $-E_{\frac{1}{2}}$  = observed value minus calculated value in kcal/gfw.

All and THE AREA STATEMENT AND A STATEMENT OF THE STATEMENT OF TH A ALASA 12-4126-1-1-1

AFRPL-TR-69-70

1.904 13.517 20.013

Sum of absolute values = 13.906

-30-

TABLE IV

Values of LHf<sup>•</sup><sub>298</sub> from the Simultaneous Adjustments

0.204 0.067 0.173 0.211 0.234 0.307 -223.043 0.478 -65.223 -76.796 -137.596 -271.293 -31.438 -198.283 HF-1 0.219 0.188 -76.796 0.062 -271.286 0.197 -223.044 -198.284 -137.597 0.158 0.285 -31.442 -65.224 0.447 HF-2 0.046 0.145 0.116 0.210 0.138 -223.020 0.164 0.388 -65.224 -76.795 -137.596 -271.285 -31.437 -198.476 HF-3 0.040 0.100 0.196 0.125 -65.2100.119 -137.583 -271.420 -31.430 -223.034 0.141 -198.216 0.353 -76.781 HF-4 Least Squares\* kcal/gfw 0.132 0.045 111.0 0.218 0.138 -222.949 0.155 -198.900 -65.209 -76.779 -137.581 -271.419 -31.414 0.354 HF-5 0.040 0.140 0.094 -271.420 0.195 0.124 -137.515 -223.035 -198.218 -65.144 -76.782 0.068 -31.431 0.351 HF-6 0.122 0.040 -271.420 0.195 0.124 0.140 -198.218 -137.516 0.068 -65.136 -76.782 -31.431 -223.035 0.351 HF-7 -31.440 -223.050 -198.250 -65.114 -76.784 -137.479 -271.600 kcal/gfw HF-7 Least Sum C<sub>2</sub>F<sub>4</sub>(polymer) HF(50 H<sub>2</sub>0) Species BF<sub>3</sub>(g) CF4 (g) NaF(c) NF<sub>3</sub>(g) HF(g)

\*The number immediately below  $\Delta Hf_{298}^{\bullet}$  is the standard error (54) as based on the overall consistency of the least squares adjustment.

AFRPL-TR-69-70

-31-

Adjustment	Unknowns	Observations*	Degrees of <u>Freedom(m-n)</u>	$(Chi)^2$	Birge's Ratio**
Hz <sup>.</sup> -7	7	23	16	18.44	1.073
(Least S	Sum) 7	23	16	(20.01)	(1.118)
HF-6	7	23	16	18.45	1.074
HF-5	7	22	15	21.46	1.196
HF-4	7	22	15	17.46	1.079
HF-3	7	23	16	25.37	1.259
HF-2	7	23	16	46.77	<b>1.7</b> 10
HF-1	7	21	14	46.66	1.826

 TABLE V

 Statistical Summary of the Simultaneous Adjustments

Standard - States All States

\* Changes in the observations are shown in Table II. \*\*See Equations 4 and 5.

- 20

## TABLE VI Selected Values of $\Delta Hf_{298}^{\bullet}$

Species	<sup>4</sup> Hf <sup>•</sup> <sub>298</sub> , kcal/gfw
HF(g)	-65.14±0.2
HF(50 H,0)	-76.78±0.1
HF(~ H_0)	-79.82±0.2
NaF(c)	-137.52±0.15
BF <sub>3</sub> (g)	-271.42:0.4
NF <sub>3</sub> (g)	-31.43±0.3
CF <sub>u</sub> (g)	-223.04±0.3
C,F <sub>u</sub> (polymer)	-198.2 ±0.7

Scoulty Classification	uu. =		
DOCUMENT C	ONTROL DATA - RE	0	
1 SAISHATH & ACTIVITY (Company auton)			
The Dow Chemical Company		Une	classified
Midland, Michigan 48640		N.	Α.
i merent title Investigation and Compilati High Temperature Chemical S	on of the The pecies	rnody	namic Properties of
Quarterly Technical Report	No. 8 - 1 Oct	. 195	- 31 Dec. 1968
Stull, D. R.; and Prophet,	н.		
A REPORT DATE	70 TOTAL NO 07 P		
A CONTRACT OF GRANT NO	1 84 - ORIGINATOR'S 81		
F04611-67-C-0009	T0009-4Q-	68	
•	AFRPL-TR-	69-70	effer punkert ther any be costigned
to AVAILABLETY LIMITATION MOVIES This docus trols and each transmittal to fe tionals may be made only with po Edwards, California 93523.	ment is subje oreign govern rior approval	ct to ments of Al	special export con- or foreign na- FRPL(RPPR/STINFO)
II SUPPLEMENTARY NOTES	AFRPL, RT Edwards	D AFB, C	viv California 93523
Supplement No. 30 to the JANAF T and sent to the printers on Decen contains the new element calcium several ionic species due to a du affinity of BO <sub>2</sub> , and revisions of table for HF.	hermochemical mber 27, 1968 and its fluo atermination f several flu	Table . Thi ride, of the orides	s was completed is Supplement revisions of electron due to a new
A complete description of the measure ous solution analysis of the interdata is given in the Appendix. (	asurements us srrelated HF	ed in heat c	the simultane- of formation
DD		UNICL	SSIFTED

UNCLASSIFIED

Security Classification		,				
		A See	LIN	K B	LIN	K C
KET BONDS	ROLE		ROLE		ROLE	
JANAF Thermochemical Table Exhaust Species						
Propellant Ingredients						
Hydrogen Fluoride						
INST I. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of De- innee activity or other organization (corporate suthor) issuing the report.	UCTIONS 10. AVAILABIL itations on furth imposed by secu such as:	IT Y/LIM	TATION N Instion of the	IOTICES the report	Enter and t, other th andard star	ny lim en the
a. REPORT SECURITY CLASSIFICATION: Enter the over- il security classification of the report. Indicate whether 'Restricted Date'' is included. Marking is to be in accord- ince with appropriate security regulations.	(1) "Qualif report f (2) "Foreig	rom DDC.	coment and	dissem	opies of th	this
15. GROUP: Automatic downgrading is specified in DoD Di- ective 5200. 10 and Armed Forces Industrial Manual. Enter he group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as author- ised.	(3) "U.S. this repusers s	Governme ort direct hall requi	not author ni agencie ly from DD pat through	nay of	n qualified	DDC
. REPORT TITLE: Enter the complete report title in all apital letters. Titles in all cases should be unclassified.	(4) "U. S. report d	military a	om DDC.	by obtain Other qu	copies of	this me

capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(5): Enter the name(s) of suthor(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of sarvice. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7s. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

76. NUMBER OF REFERENCES Enter the total number of references cited in the report.

Sa. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

Sb, &c, & Sd. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

96. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unque to this report.

9b. OTHER REPORT NUMBER(S): If the report has been essigned any other report numbers (either by the originator or by the sponsor), also enter this number(s). If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicete this fect and enter the price, if known.

(5) "All distribution of this report is controlled. Qualified DDC users shall request through

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports he unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (T5), (5), (C), or (U)

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

ever, the suggested length is from 150 to 225 words. 14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no necurity classification is required. Idenfiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

> UNCLASSIFIED Security Classification

-40-