

UNCLASSIFIED

AD NUMBER
AD869934
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAY 1970. Other requests shall be referred to Department of Army, Fort Detrick, Attn: Technical Release Branch/TID, Frederick, MD 21701.
AUTHORITY
BDRL, D/A ltr, 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

20
CB

AD 869934

TECHNICAL MANUSCRIPT 613

LOSS OF PICLORAM AND 2,4,5-T
FROM THE ROOTS
OF ASH AND MAPLE SEEDLINGS

AD No. —
DDC FILE COPY

William A. Wells
Woodland Hurtt
Charles P. P. Reid

MAY 1970

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch, TID, Frederick, Maryland 21701

DEPARTMENT OF THE ARMY

Fort Detrick
Frederick, Maryland

DDC
RECEIVED
JUN 10 1970
C

49834

DD FORM 1300		
DDC		
EXAMINED		
CLASSIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
DDC	AVAIL. and/or SPECIAL	
2		

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 613

LOSS OF PICLORAM AND 2,4,5-T FROM THE ROOTS
OF ASH AND MAPLE SEEDLINGS

William A. Wells

Woodland Hurtt

Charles P.P. Reid

Plant Physiology Division
PLANT SCIENCES LABORATORIES

Project 1B562602AD04

May 1970

ABSTRACT

Sublethal concentrations of C^{14} -picloram (4-amino-3,5,6-trichloropicolinic acid) and C^{14} -2,4,5-T (2,4,5-trichlorophenoxyacetic acid) were foliarly applied to red maple and green ash seedlings grown in nutrient solution. C^{14} -picloram was also applied to the leaves of white ash.

Radioactivity was detected in nutrient solution samples from all treated plants within 24 hours. C^{14} -picloram loss through the roots increased over a 3-week period, with the exception of a slight decrease in the amount lost on the 6th and 16th day from the green ash. The total radioactivity in the white ash nutrient solution increased over the 9-day experiment.

The nutrient solution samples of the red maple treated with C^{14} -2,4,5-T yielded radioactivity within 24 hours; this amount increased over the 3-week sampling period except for a decrease on the 16th day. The green ash nutrient solution showed C^{14} -2,4,5-T activity at 48 hours and increased throughout the 3 weeks. Greater loss of both labeled herbicides occurred from red maple than from green ash.

The fact that C^{14} activity in the exudates was associated with unaltered herbicide molecules was substantiated by bioassays.

I. INTRODUCTION*

Ash (Fraxinus spp.) has been reported to be generally resistant to picloram, although red maple (Acer rubrum) is susceptible. Both ash and maple are also reported to be relatively tolerant to 2,4,5-T.¹⁻³ These differences in the responses of ash and maple to picloram and 2,4,5-T have not been completely explained, although attempts to explain the differences have been made through studies of phloem and xylem transport systems and foliar uptake mechanisms.^{1,3,4}

Root excretion as a means of detoxifying foliarly applied herbicides has been suggested,⁵ and numerous reports state that both exogenous and endogenous substances are root-excreted by a wide range of plants.⁶⁻⁹ In view of this and the fact that picloram has been reported to be excreted from the roots of bean,^{10,11} it was considered worthwhile to study the possibility that ash and maple tolerance to picloram or 2,4,5-T is related to root excretion. Therefore, the goal of these experiments was twofold, to determine (i) if picloram and 2,4,5-T are root-excreted by ash and maple, and (ii) if so, are they in sufficient quantities to explain differential tolerance to the two herbicides.

II. MATERIALS AND METHODS

Uniform seedlings of red maple (Acer rubrum L.), green ash (Fraxinus pennsylvanica Marsh.), and white ash (Fraxinus americana L.) were maintained in containers of 0.5X Hoagland's solution under controlled environmental conditions in a growth chamber.

After one day in solution culture, sublethal dosages of C¹⁴-labeled picloram (4.25 µc/mg) or C¹⁴-labeled 2,4,5-T (4.25 µc/mg) in 95% ethanol were applied to the two leaves at the fourth whorl above the root collar. Each plant received ten 5-µl droplets of the labeled herbicide, which were applied within lanolin rings on the upper surface of the leaf with a calibrated microliter syringe.

The red maple and green ash seedlings were treated with both labeled herbicides; the white ash seedlings received only labeled picloram. The containers were brought up to a predetermined volume before 5-ml samples were taken daily for 7 days and then every 2 to 3 days until the termination of the red maple and green ash studies at 22 days. The white ash experiment covered a 9-day period; samples were taken each day.

* This report should not be used as a literature citation in material to be published in the open literature. Readers interested in referencing the information contained herein should contact the senior author to ascertain when and where it may appear in citable form.

The samples of the nutrient solution were evaporated to dryness under vacuum and counted in a liquid scintillation counter to determine the amount of C^{14} present. The background count was determined from nutrient solution samples of untreated plants.

Bioassays were conducted to determine if the areas of biological activity and radioactivity in the nutrient solution were in agreement. In order to do this, the nutrient solution was filtered and evaporated to a 435-fold decrease in volume. The solution was then spotted on Whatman No. 20 chromatograph paper and co-chromatographed with labeled picloram and 2,4,5-T standards. Both the biological activity and radioactivity of the developed chromatographs were determined, the former by a lettuce seed bioassay¹² and the latter by a strip-scanner.

III. RESULTS

The radioactivity of the mean cumulative total of C^{14} for replicates at each sampling period for the red maple and green ash is shown in Figure 1.

Loss of picloram was observed within 24 hours in both red maple and green ash. The amount of picloram lost increased over the 22-day period, except for slight decreases on the 6th and 16th days of sampling.

The maple nutrient solution showed the presence of labeled 2,4,5-T within 24 hours, and the amount increased over the entire sampling period except for a decrease on the 16th day. Loss of 2,4,5-T was evident in the ash nutrient solution within 48 hours, and increased over the 3 weeks of the experiment.

A greater loss of both herbicides was noted in the nutrient solution samples of the red maple seedlings compared with those from the green ash. The decrease in amount of radioactivity in samples taken on the 16th day has not been satisfactorily explained.

The cumulative total C^{14} activity of the nutrient solution samples of each replicate of the white ash seedlings is shown in Figure 2. The amount of C^{14} lost by the plant increased over the 9-day treatment period. There was considerable variability in the quantity of C^{14} lost by the replicates, but the maxima and minima generally are in agreement for all of the replicates on a given sampling. The 19,200 counts per minute (cpm) on day 9 represents 4.3% of the total picloram applied.

The lettuce seed bioassays of the nutrient solution showed that biological activity indicating the presence of picloram closely coincided with the areas of radioactivity when the nutrient solution was co-chromatographed with picloram and 2,4,5-T standards.




FIGURE 1. Cumulative ^{14}C Loss from Roots of Red Maple and Green Ash after Foliar Treatment with 0.1 μg (24 μg) of ^{16}C -Picloram and ^{14}C -2,4,5-T. Each plotted line represents the mean of six replicates.




FIGURE 2. Cumulative C^{14} Loss from Roots of White Ash after Foliar Treatment with 0.1 μ Ci (24 μ g) of C^{14} -Picloram. Each plotted line represents one replicate consisting of two trees in 500 ml of nutrient solution.

IV. DISCUSSION

Data from these tests show that significant quantities of picloram and 2,4,5-T are lost from the roots of red maple and of green and white ash seedlings. The amounts of herbicides lost from the roots appears to have no consistent relationship with species tolerance or resistance. This would suggest that root exudation, as a detoxification mechanism, does not determine species response to the particular herbicide.

The amounts of picloram and 2,4,5-T lost by the roots of these woody species may, however, be of considerable ecological importance. The effects of root-excreted substances on other organisms (allelopathy) has been recognized by others.^{9,13-15} Although Mitchell and Linder⁹ believed that root exudation of exogenous compounds is of no practical significance, they did acknowledge that the growth of subsequent crops or nearby plants might be affected by exuded substances that persist in the soil. The reported persistence of picloram in soils^{16,17} may make its exudation by woody species a factor of considerable ecological importance.

Mitchell and Linder⁹ state in their review article that regulating substances exuded from roots were confined to the α -methoxyphenylacetic acids and chlorinated benzoic acids. Since the time of their article, the exudation of 2,4-D^{5,18} and picloram^{10,11} have been reported. Neither of these compounds or 2,4,5-T falls within the two-family classification reported by Mitchell and Linder.

It should be realized that many factors found in the environment of soil-grown plants may differ from those observed in nutrient solution culture. These would include variation in temperature, humidity, light conditions, and the presence of microbes that could cause bio-degradation, as well as a host of unknown variables. The importance of these factors is illustrated by the findings of Meikle et al,¹⁹ who have found that the combination of living plant and soil was more active in decomposition of picloram than either the plant or soil alone.

Thus, although a difference in species response can be seen at the physiological level, i.e., greater loss of both herbicides by the red maple, it is difficult to correlate this with previously reported differences in species tolerance. However, the fact that detectable amounts of exogenous compounds may be lost to the rhizosphere from the roots of treated plants should be considered in order to obtain a better understanding of the physiological and possible ecological aspects of the use of growth-regulating compounds for woody plant control.

LITERATURE CITED

1. Perry, P.W.; Upchurch, R.P. 1968. Growth analysis of red maple and white ash seedlings treated with eight herbicides. *Weed Sci.* 16:32-37.
2. Nation, H.A.; Lichy, C.T. 1964. Tordon herbicide for brush control in Southern United States. *Proc. Southern Weed Control Conference.* 17:287-294.
3. Pallas, J.E., Jr. 1963. Absorption and translocation of the triethylamine salt of 2,4-D and 2,4,5-T in four woody species. *Forest Sci.* 9:485-491.
4. Leonard, O.A.; Bayer, D.E.; Glenn, R.K. 1966. Translocation of herbicides and assimilates in red maple and white ash. *Bot. Gaz.* 127:193-201.
5. Fites, R.C.; Slife, F.W.; Hanson, J.B. 1964. Translocation and metabolism of radioactive 2,4-D in jimsonweed. *Weeds* 12:180-183.
6. Garb, S. 1961. Differential growth-inhibitors produced by plants. *Bot. Rev.* 27:422-443.
7. Borner, H. 1960. Liberation of organic substances from higher plants and their role in the soil sickness problem. *Bot. Rev.* 26:393-424.
8. Moreland, E.E.; Egley, G.H.; Worsham, A.D.; Monaco, T.J. 1966. Regulation of plant growth by constituents from higher plants, p. 112-141. *In* *Natural Pest Control Agents*, (Advances in Chemistry Series 53). American Chemical Society, Washington, D.C.
9. Mitchell, J.W.; Linder, P.J. 1963. Absorption, translocation, exudation, and metabolism of plant growth-regulating substances in relation to residues, p. 51-76. *In* Francis A. Gunther, (ed.) *Residue reviews*, Volume 2. Springer-Verlag New York Inc., New York, N.Y.
10. Hurtt, W.; Foy, C.L. 1965. Excretion of foliarly applied dicamba and picloram from roots of Black Valentine beans grown in soil, sand, and culture solution. *Proc. Northeastern Weed Control Conference* 19:602.
11. Hurtt, W.; Foy, C.L. 1965. Some factors influencing the excretion of foliarly applied dicamba and picloram from roots of Black Valentine beans. *Plant Physiol.* 40(Suppl.):xlvi. (Abstr.)
12. Reid, C.P.P.; Hurtt, W. 1969. A rapid bioassay for simultaneous identification and quantitation of picloram in aqueous solution. *Weed Res.* 9:136-141.

13. Mitchell, J.W.; Smale, B.C.; Preston, W.H., Jr. 1959. New plant regulators that exude from roots. *J. Agr. Food Chem.* 7:841-843.
14. Woods, F.W. 1960. Biological antagonisms due to phytotoxic root exudates. *Bot. Rev.* 26:546-569.
15. van Overbeek, J. 1966. Plant hormones and regulators. *Science* 152:721-731.
16. Hamaker, J.W.; Johnston, H.; Martin, R.T.; Redemann, C.T. 1963. A picolinic acid derivative: A plant growth regulator. *Science* 141:363.
17. Grover, R. 1967. Studies on the degradation of 4-amino-3,5,6-trichloropicolinic acid in soil. *Weed Res.* 7:61-67.
18. Crafts, A.S.; Yamaguchi, S. 1964. The autoradiography of plant materials. *Calif. Agr. Expt. Sta. Manual* 35, 143 p.
19. Meikle, R.W.; Williams, E.A.; Redemann, C.T. 1966. Metabolism of Tordon herbicide (4-amino-3,5,6-trichloropicolinic acid) in cotton and decomposition in soil. *J. Agr. Food Chem.* 14:384-387.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R & D						
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)						
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION				
Department of the Army Fort Detrick, Frederick, Maryland, 21701		Unclassified				
2b. GROUP						
3. REPORT TITLE						
LOSS OF PICLORAM AND 2,4,5-T FROM THE ROOTS OF ASH AND MAPLE SEEDLINGS						
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)						
5. AUTHOR(S) (First name, middle initial, last name)						
William A. Wells Charles P.P. Reid Woodland (NMI) Hurtt						
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS				
May 1970	13	19				
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S)				
a. PROJECT NO. 1B562602ADO4		Technical Manuscript 613				
c. Task-Work Unit 02-001		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)				
4. DD 1498 Agency Access. DA OL 0003		CMs 6609, AMXFD-AE-T 49834				
10. DISTRIBUTION STATEMENT						
Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized.						
Release or announcement to the public is not authorized.						
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY				
Plant Physiology Division		Department of the Army Fort Detrick, Frederick, Md., 21701				
13. ABSTRACT						
Sublethal concentrations of C¹⁴-picloram (4-amino-3,5,6-trichloro-picolinic acid) and C¹⁴-2,4,5-T (2,4,5-trichlorophenoxyacetic acid) were foliarly applied to red maple and green ash seedlings grown in nutrient solution. C¹⁴-picloram was also applied to the leaves of white ash. Radioactivity was detected in nutrient solution samples from all treated plants within 24 hours. C¹⁴-picloram loss through the roots increased over a 3-week period, with the exception of a slight decrease in the amount lost on the 6th and 16th day from the green ash. The total radioactivity in the white ash nutrient solution increased over the 9-day experiment. The nutrient solution samples of the red maple treated with C¹⁴-2,4,5-T yielded radioactivity within 24 hours; this amount increased over the 3-week sampling period except for a decrease on the 16th day. The green ash nutrient solution showed C¹⁴-2,4,5-T activity at 48 hours and increased throughout the 3 weeks. Greater loss of both labeled herbicides occurred from red maple than from green ash. The fact that C¹⁴ activity in the exudates was associated with unaltered herbicide molecules was substantiated by bioassays. ()						
14. Key Words						
<table border="0"> <tr> <td>Picloram</td> <td>Tree roots</td> </tr> <tr> <td>2,4,5-T</td> <td>Excretion</td> </tr> </table>			Picloram	Tree roots	2,4,5-T	Excretion
Picloram	Tree roots					
2,4,5-T	Excretion					

DD FORM 1473
1 NOV 66

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified