


**REPORT DOCUMENTATION PAGE**

Form Approved  
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.  
**PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

<b>1. REPORT DATE (DD-MM-YYYY)</b> 01/01/1997		<b>2. REPORT TYPE</b> Technical Report - Other		<b>3. DATES COVERED (From - To)</b>	
<b>4. TITLE AND SUBTITLE</b> On Two Problems in the Multivariate Approximation				<b>5a. CONTRACT NUMBER</b>	
				<b>5b. GRANT NUMBER</b>	
				<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b>				<b>5d. PROJECT NUMBER</b>	
				<b>5e. TASK NUMBER</b>	
				<b>5f. WORK UNIT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> University of South Carolina Columbia SC United States				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> Office of Naval Research Arlington VA United States				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
				<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>	
<b>12. DISTRIBUTION/AVAILABILITY STATEMENT</b> A = Approved For Public Release 12/2/2015 No					
<b>13. SUPPLEMENTARY NOTES</b>					
<b>14. ABSTRACT</b> The paper contains two theorems on approximation of functions with bounded mixed derivative. These theorems give some progress in two old open problems. The first one gives, in particular, an upper estimate in the Bernstein L1-inequality for trigonometric polynomials on two variables with harmonics in hyperbolic crosses. The second one gives the order of the entropy numbers and Kolmogorov's widths in the $L_\infty$ -norm of the class $MW/r_\infty$					
<b>15. SUBJECT TERMS</b>					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>	<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b>
<b>a. REPORT</b> U	<b>b. ABSTRACT</b> U	<b>c. THIS PAGE</b> U			<b>19b. TELEPHONE NUMBER (Include area code)</b>



INDUSTRIAL  
MATHEMATICS  
INSTITUTE

1998:22

On two problems in the  
multivariate approximation

V.N. Temlyakov

IMI  
Preprint Series

Department of Mathematics  
University of South Carolina

# ON TWO PROBLEMS IN THE MULTIVARIATE APPROXIMATION<sup>1</sup>

V.N. TEMLYAKOV

Department of Mathematics, University of South Carolina, Columbia, SC 29208

ABSTRACT. The paper contains two theorems on approximation of functions with bounded mixed derivative. These theorems give some progress in two old open problems. The first one gives, in particular, an upper estimate in the Bernstein  $L_1$ -inequality for trigonometric polynomials on two variables with harmonics in hyperbolic crosses. The second one gives the order of the entropy numbers and Kolmogorov's widths in the  $L_\infty$ -norm of the class  $MW_{\infty, \alpha}^r$  of functions of two variables.

## 1. INTRODUCTION

In this paper we report some progress in two old problems from approximation of multivariate functions with bounded mixed derivative. These two problems originate from the very first publications (see [B1], [B2], [M], [Te]) on the subject. We introduce first some notations. We give these notations in the general  $d$ -dimensional (functions on  $d$  variables) case and point out that the results in this paper concern only the 2-dimensional case. Let  $s = (s_1, \dots, s_d)$  be a vector whose coordinates are nonnegative integers and

$$\rho(s) := \{k \in Z^d : [2^{s_j-1}] \leq |k_j| < 2^{s_j}, j = 1, \dots, d\},$$

where  $[a]$  means the integer part of a number  $a$ . Denote

$$Q_m := \bigcup_{\|s\|_1 \leq m} \rho(s)$$

and denote by  $\mathcal{T}(Q_m)$  the subspace of trigonometric polynomials with harmonics in  $Q_m$ .

**Problem 1. (The Bernstein inequality in  $L_1$ ).** *Find the order in  $m$  of the sequence*

$$b(m, r, L_1) := \sup_{f \in \mathcal{T}(Q_m)} \|f^{(r, \dots, r)}\|_1 / \|f\|_1, \quad m = 1, 2, \dots$$

This problem is open for all  $r > 0$  and  $d \geq 2$ . The corresponding problem for the  $L_p$ -norm,  $1 < p \leq \infty$ , was solved in early 60-th. K.I. Babenko [B2] proved the upper bound

$$b(m, r, L_\infty) \ll m^{d-1} 2^{rm}$$

---

<sup>1</sup>This research was supported by the National Science Foundation Grant DMS 9622925 and by ONR Grant N0014-96-1-1003

and S.A. Telyakovskii [Te] proved the lower bound

$$b(m, r, L_\infty) \gg m^{d-1} 2^{rm}.$$

B.S. Mityagin [M] proved for  $1 < p < \infty$

$$b(m, r, L_p) \asymp 2^{rm}.$$

For these results see also [T1], Ch.1, s.1. The known bounds for  $b(m, r, L_1)$  are simple

$$2^{rm} \ll b(m, r, L_1) \ll m^{d-1} 2^{rm}.$$

We prove here a new upper bound in the 2-dimensional case:

$$b(m, r, L_1) \ll m^{1/2} 2^{rm}.$$

The second problem we are dealing with here is about the Kolmogorov widths of classes  $MW_{\infty, \alpha}^r$  in the  $L_\infty$ -norm. Let for  $r > 0$  and  $\alpha \in \mathbb{R}$

$$F_r(t, \alpha) = 1 + 2 \sum_{k=1}^{\infty} k^{-r} \cos(kt - \alpha\pi/2), \quad t \in [0, 2\pi],$$

and for  $x = (x_1, \dots, x_d)$  and  $\alpha = (\alpha_1, \dots, \alpha_d)$

$$F_r(x, \alpha) = \prod_{j=1}^d F_r(x_j, \alpha_j).$$

Define

$$MW_{q, \alpha}^r = \{f : f = F_r(\cdot, \alpha) * \phi(\cdot), \quad \|\phi\|_q \leq 1\},$$

where  $*$  means the convolution. We recall the definition of the Kolmogorov width of a centrally symmetric set  $A$  in a Banach space  $X$

$$d_n(A, X) := \inf_{g_j \in X, j=1, \dots, n} \sup_{f \in A} \inf_{c_j, j=1, \dots, n} \|f - \sum_{j=1}^n c_j g_j\|_X.$$

**Problem 2.** Find the order of the sequence  $\{d_n(MW_{\infty, \alpha}^r, L_\infty)\}_{n=1}^{\infty}$ .

The first upper bounds in this problem were obtained by K.I. Babenko [B2]

$$d_n(MW_{\infty, \alpha}^r, L_\infty) \ll n^{-r} (\log n)^{(d-1)(r+1)}.$$

We prove in this paper the lower bound

$$d_n(MW_{\infty, \alpha}^r, L_\infty) \gg n^{-r} (\log n)^{r+1/2}, \quad r > 1/2,$$

in the 2-dimensional case. This lower estimate combined with the corresponding known upper estimate (see [Be]) gives the solution to Problem 2 in the case  $d = 2$  and  $r > 1/2$  :

$$d_n(MW_{\infty, \alpha}^r, L_\infty) \asymp n^{-r} (\log n)^{r+1/2}.$$

Problem 2 is open in the case  $d > 2$ .

## 2. THE BERNSTEIN INEQUALITIES

We introduce some more notations. Let for two integers  $a \geq 1$  and  $0 \leq b < a$   $AP(a, b)$  denote an arithmetic progression of the form  $al + b$ ,  $l = 0, 1, \dots$  and

$$H_n(a, b) := \{s : s \in \mathbb{Z}_+^2, \quad \|s\|_1 = n, \quad s_1, s_2 \geq a, \quad s_1 \in AP(a, b)\}.$$

It will be convenient for us to consider subspaces  $\mathcal{T}(\rho'(s))$  of trigonometric polynomials with harmonics in

$$\rho'(s) := \{k \in \mathbb{Z}^2 : [2^{s_j-2}] \leq |k_j| < 2^{s_j}, \quad j = 1, 2\}.$$

For a subspace  $Y$  in  $L_2(\mathbb{T}^d)$  we denote by  $Y^\perp$  its orthogonal complement.

**Lemma 2.1.** *Take any trigonometric polynomials  $t_s \in \mathcal{T}(\rho'(s))$  and form the function*

$$\Phi(x) := \prod_{s \in H_n(a, b)} (1 + t_s(x)).$$

*Then for any  $a \geq 6$  and any  $0 \leq b < a$  this function admits the representation*

$$\Phi(x) = 1 + \sum_{s \in H_n(a, b)} t_s(x) + R(x)$$

*with  $R \in \mathcal{T}(Q_{n+a-6})^\perp$ .*

The proof of this lemma is similar to the proof of Lemma 2.1 from [T2] (for the idea of the proof see also [T1], p. 57(60)).

**Remark 2.1.** *For any real numbers  $|y_l| \leq 1$ ,  $l = 1, \dots, N$ , we have ( $i^2 = -1$ )*

$$\left| \prod_{l=1}^N \left(1 + \frac{iy_l}{\sqrt{N}}\right) \right| \leq C.$$

**Lemma 2.2.** *For any function  $f$  of the form*

$$f = \sum_{s \in H_n(a, b)} t_s$$

*with  $a \geq 6$ ,  $0 \leq b < a$ , and  $t_s$ ,  $s \in H_n(a, b)$ , is a real trigonometric polynomial in  $\mathcal{T}(\rho'(s))$  such that  $\|t_s\|_\infty \leq 1$  we have*

$$E_{Q_{n+a-6}}^\perp(f)_\infty := \inf_{g \in \mathcal{T}(Q_{n+a-6})^\perp} \|f - g\|_\infty \ll (1 + n/a)^{1/2}.$$

*Proof.* Let us form the function

$$RP(f) := \text{Im} \prod_{s \in H_n(a, b)} (1 + it_s(1 + n/a)^{-1/2}),$$

which is an analog of the Riesz product. Then by Remark 2.1 we have

$$(2.1) \quad \|RP(f)\|_\infty \leq C.$$

Lemma 2.1 provides the representation

$$(2.2) \quad RP(f) = (1 + n/a)^{-1/2} \sum_{s \in H_n(a, b)} t_s + g, \quad g \in \mathcal{T}(Q_{n+a-6})^\perp.$$

Combining (2.1) and (2.2) we get the statement of Lemma 2.2.

**Remark 2.2.** *It is clear that in Lemma 2.2 we can drop the assumption that the  $t_s$  are real polynomials.*

**Lemma 2.3.** *For any function  $f$  of the form*

$$f = \sum_{\|s\|_1=n} t_s, \quad t_s \in \mathcal{T}(\rho'(s)), \quad \|t_s\|_\infty \leq 1,$$

*we have for any  $a \geq 6$*

$$E_{Q_{n+a-6}}^\perp(f)_\infty \leq Ca(1 + n/a)^{1/2}.$$

*Proof.* Let us introduce some more notations. Denote

$$\theta_n := \{s : \|s\|_1 = n\}; \quad \theta_{n,a} := \{s \in \theta_n, \quad s_1 < a \quad \text{or} \quad s_2 < a\}.$$

Then

$$f = \sum_{s \in \theta_n} t_s = \sum_{s \in \theta_{n,a}} t_s + \sum_{b=0}^{a-1} \sum_{s \in H_n(a,b)} t_s$$

and

$$E_{Q_{n+a-6}}^\perp(f)_\infty \leq \sum_{s \in \theta_{n,a}} \|t_s\|_\infty + \sum_{b=0}^{a-1} E_{Q_{n+a-6}}^\perp\left(\sum_{s \in H_n(a,b)} t_s\right)_\infty.$$

Using the assumption  $\|t_s\|_\infty \leq 1$ , Lemma 2.2 and Remark 2.2 we get from here the required estimate.

Lemma 2.3 is proved.

It proved to be useful in studying approximation of functions with bounded mixed derivative to consider along with the  $L_p$ -norms the Besov type norms. Let  $V_n(t)$  be the de la Vallée-Poussin polynomials,  $t \in [0, 2\pi]$ . We define

$$\mathcal{A}_0(t) := 1, \quad \mathcal{A}_1(t) := V_1(t) - 1, \quad \mathcal{A}_n(t) := V_{2^{n-1}}(t) - V_{2^{n-2}}(t), \quad n \geq 2,$$

and for  $x = (x_1, x_2)$ ,  $s = (s_1, s_2)$

$$\mathcal{A}_s(x) := \mathcal{A}_{s_1}(x_1)\mathcal{A}_{s_2}(x_2).$$

Consider the convolution operator  $A_s$  with the kernel  $\mathcal{A}_s(x)$ ,

$$A_s(f) := f * \mathcal{A}_s,$$

and define the  $B_{1,1}^r$ -norm as follows

$$\|f\|_{B_{1,1}^r} := \sum_s 2^{r\|s\|_1} \|A_s(f)\|_1.$$

**Theorem 2.1.** *Let  $r > 0$  be given. For any  $f \in \mathcal{T}(Q_m)$  we have*

$$\|f\|_{B_{1,1}^r} \leq C(r)m^{1/2}2^{rm}\|f\|_1.$$

*Proof.* Take any  $n \leq m + 2$  and consider the sum

$$S_n := \sum_{\|s\|_1=n} \|A_s(f)\|_1.$$

Define the polynomials  $t_s := A_s(\text{sign } A_s(f))$  and

$$g_n := \sum_{\|s\|_1=n} \bar{t}_s.$$

It is clear from the definition of  $A_s(\cdot)$  that  $\bar{t}_s \in \mathcal{T}(\rho'(s))$ . Next, on one hand we have

$$(2.3) \quad \langle f, g_n \rangle = \sum_{\|s\|_1=n} \|A_s(f)\|_1$$

and on the other hand we have

$$(2.4) \quad \langle f, g_n \rangle \leq \|f\|_1 E_{Q_m}^\perp(g_n)_\infty.$$

By Lemma 2.3 with  $a = \max(m - n + 6, 6)$  we get

$$(2.5) \quad E_{Q_m}^\perp(g_n)_\infty \ll (m - n + 6)(1 + n/(m - n + 6))^{1/2}.$$

The relations (2.3)–(2.5) imply the inequality

$$\sum_s 2^{r\|s\|_1} \|A_s(f)\|_1 = \sum_{n \leq m+2} 2^{rn} S_n \leq C(r)m^{1/2}2^{rm}\|f\|_1,$$

that is what was required.

We consider a more general derivative than standard mixed derivative. For a polynomial  $f \in \mathcal{T}(Q_m)$  we define its  $(r, \alpha)$ -derivative as the convolution with the kernel

$$U_m^r(x, \alpha) := 4 \sum_{k \in Q_m, k > 0} (k_1 k_2)^r \cos(k_1 x_1 + \alpha_1 \pi/2) \cos(k_2 x_2 + \alpha_2 \pi/2).$$

Thus,

$$f^{(r)}(x, \alpha) := D^{r, \alpha} f := f(\cdot) * U_m^r(\cdot, \alpha).$$

**Corollary 2.1.** *For any  $r > 0$ ,  $\alpha \in \mathbb{R}^2$ , we have for any function  $f \in \mathcal{T}(Q_m)$  on two variables the inequality*

$$\|f^{(r)}(x, \alpha)\|_1 \leq C(r)m^{1/2}2^{rm}\|f\|_1.$$

*Proof.* By Bernstein inequality for trigonometric polynomials with harmonics in rectangles we get

$$\|f^{(r)}(x, \alpha)\|_1 \leq \sum_s \|D^{r,\alpha} A_s(f)\|_1 \leq C(r) \sum_s 2^{r\|s\|_1} \|A_s(f)\|_1 \leq$$

and using Theorem 2.1 we continue

$$\leq C(r)m^{1/2}2^{rm}\|f\|_1.$$

**Remark 2.3.** *The inequality in Theorem 2.1 is sharp. The multiplier  $m^{1/2}2^{rm}$  can not be replaced by smaller function on  $m$  even if we write  $\|f\|_p$ ,  $p < \infty$ , instead of  $\|f\|_1$  in Theorem 2.1.*

*Proof.* Indeed, take

$$f = \sum_{\|s\|_1=m} \cos 2^{s_1} x_1 \cos 2^{s_2} x_2.$$

Then

$$\sum_s 2^{r\|s\|_1} \|A_s(f)\|_1 \asymp m2^{rm}$$

and by Littlewood-Paley Theorem

$$\|f\|_p \ll m^{1/2}, \quad p < \infty.$$

The next theorem shows that we can improve the inequality in Theorem 2.1 if replace  $\|f\|_1$  by  $\|f\|_\infty$ .

**Theorem 2.2.** *Let  $r > 0$  be given. For any  $f \in \mathcal{T}(Q_m)$  we have*

$$\|f\|_{B_{1,1}^r} \leq C(r)2^{rm}\|f\|_\infty.$$

*Proof.* Using functions  $\Phi(x)$  defined in Lemma 2.1 and their property  $\|\Phi\|_1 = 1$  instead of functions  $RP(f)$  defined in the proof of Lemma 2.2 and their property  $\|RP(f)\|_\infty \leq C$  we get the following analog of Lemma 2.3 (see also Lemma 2.1 in [T2]).

**Lemma 2.4.** *For any function  $f$  of the form*

$$f = \sum_{\|s\|_1=n} t_s, \quad t_s \in \mathcal{T}(\rho'(s)), \quad \|t_s\|_\infty \leq 1,$$

*we have for any  $a \geq 6$*

$$E_{Q_{n+a-6}}^\perp(f)_1 \leq Ca.$$

We repeat now the arguments from the proof of Theorem 2.1 with (2.4) replaced by

$$\langle f, g_n \rangle \leq \|f\|_\infty E_{Q_m}^\perp(g_n)_1$$

and Lemma 2.3 replaced by Lemma 2.4.



### 3. THE KOLMOGOROV WIDTHS

We prove in this section estimates for the entropy numbers and Kolmogorov's widths of the class  $MW_{\infty, \alpha}^r$ . We use the following definition of the entropy numbers

$$\epsilon_m(F, X) = \inf\{\epsilon : \exists f_1, \dots, f_{2^m} \in X : F \subset \cup_{j=1}^{2^m} (f_j + \epsilon B(X))\},$$

where  $B(X)$  is the unit ball of the Banach space  $X$ .

**Theorem 3.1.** *Let  $r > 1/2$ . We have for the class  $MW_{\infty, \alpha}^r$  of functions on two variables the asymptotic relations*

$$\epsilon_m(MW_{\infty, \alpha}^r, L_\infty) \asymp m^{-r} (\log m)^{r+1/2},$$

$$d_m(MW_{\infty, \alpha}^r, L_\infty) \asymp m^{-r} (\log m)^{r+1/2}.$$

*Proof.* The upper estimates are known (see [Be]). We prove here the corresponding lower estimates. We consider first the entropy numbers and use the interpolation property of these numbers (see [P], s. 12.1.12)

$$(3.1) \quad \epsilon_{2m}(MW_{4, \alpha}^r, L_\infty) \leq 2\epsilon_m(MW_{2, \alpha}^r, L_\infty)^{1/2} \epsilon_m(MW_{\infty, \alpha}^r, L_\infty)^{1/2}.$$

Next, from [T2] we get for  $r > 1/4$

$$\epsilon_{2m}(MW_{4, \alpha}^r, L_\infty) \geq C(r) m^{-r} (\log m)^{r+1/2}$$

and from [Be] we have for  $r > 1/2$

$$\epsilon_m(MW_{2, \alpha}^r, L_\infty) \leq C(r) m^{-r} (\log m)^{r+1/2}.$$

Substituting these two estimates into (3.1) we get the required lower estimate for the entropy numbers.

It is known (see [L]) that the entropy numbers give in a certain sense the lower bounds for the Kolmogorov widths. We formulate this in a way convenient for us (see [KT] and also [T3]).

**Lemma 3.1.** *Let  $A$  be a compact set in a separable Banach space  $X$ . Assume that for two real numbers  $r > 0$  and  $b$  we have*

$$\epsilon_m(A, X) \asymp m^{-r} (\log m)^b.$$

*Then for Kolmogorov's widths of this set we have*

$$d_m(A, X) \gg m^{-r} (\log m)^b.$$

Application of this lemma completes the proof of Theorem 3.1.

## REFERENCES

- [B1] K.I. Babenko, *On the approximation of periodic functions of several variables by trigonometric polynomials*, Dokl. Akad. Nauk USSR **132** (1960), 247–250; English transl. in Soviet Math. Dokl. **1** (1960).
- [B2] K.I. Babenko, *On the approximation of a certain class of periodic functions of several variables by trigonometric polynomials*, Dokl. Akad. Nauk USSR **132** (1960), 982–985; English transl. in Soviet Math. Dokl. **1** (1960).
- [Be] E.S. Belinskii, *The asymptotic characteristics of classes of functions with restrictions on the mixed derivative (mixed difference)*, Investigations in the theory of functions on several real variables, Yaroslavskii Univ. (1990), 22–37. (Russian)
- [KT] B.S. Kashin and V.N. Temlyakov, *On estimating approximative characteristics of classes of functions with bounded mixed derivative*, Math. Zametki **58** (1995), 922–925; English transl. in Math. Notes **58**.
- [L] G.G. Lorentz, *Metric entropy and approximation*, Bull. Amer. Math. Soc. **72** (1966), 903–937.
- [M] B.S. Mityagin, *Approximation of functions in the spaces  $L^p$  and  $C$  on the torus*, Matem. Sbornik **58** (1962), 397–414. (Russian)
- [P] A. Pietsch, *Operator Ideals*, VEB, Berlin, 1978.
- [Te] S.A. Telyakovskii, *On estimates of the derivatives of trigonometric polynomials in several variables*, Siberian Math. Zh. **4** (1963), 1404–1411. (Russian)
- [T1] V. N. Temlyakov, *Approximation of functions with bounded mixed derivative*, Trudy MIAN **178** (1986); English transl. in Proc. Steklov Inst. Math. (1989), no. 1.
- [T2] V.N. Temlyakov, *An inequality for trigonometric polynomials and its application for estimating the entropy numbers*, J. Complexity **11** (1995), 293–307.
- [T3] V.N. Temlyakov, *An inequality for trigonometric polynomials and its application for estimating the Kolmogorov widths*, East J. on Approximations **2** (1996), 253–262.