



## BIOMEDICAL HYPOTHESIS

# Injury, inflammation and the emergence of human-specific genes

Andrew Baird, PhD; Todd Costantini, MD; Raul Coimbra, MD, PhD; Brian P. Eliceiri, PhD

Division of Trauma, Surgical Critical Care, Burns and Acute Care Surgery, Department of Surgery, University of California San Diego School of Medicine, La Jolla, San Diego, California

**Reprint requests:**

Andrew Baird, PhD, Division of Trauma, Surgical Care, Burns, and Acute Care Surgery, Department of Surgery, UC San Diego Health Sciences, 212 Dickinson Street, MC 8236, San Diego, CA 92103, USA.

Tel: 619-471-9027;

Fax: 619-543-2325;

Email: [anbaird@ucsd.edu](mailto:anbaird@ucsd.edu)

Manuscript received: November 24, 2015

Accepted in final form: February 8, 2016

DOI:10.1111/wrr.12422

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

The fact that new genes emerge during speciation is not unexpected, although their significance to the injury response remains largely enigmatic. Yet, for over 60 years, a tenet of modern molecular biology has held that essential genes, like those needed for tissue repair, have significant protein and/or DNA sequence homology across multiple species.<sup>1</sup> As a consequence, the contributions of human-specific genes to the onset and progression of human disease have been largely understudied.<sup>2</sup> In contrast, the genetic mechanisms to explain the emergence of these taxonomically restricted genes are well described. They include complete and/or partial gene duplication, gene rearrangement, silencing, mutagenesis and, the endogenization of foreign DNA.<sup>3</sup>

If an increased incidence of injury can be associated with or be the result of newly acquired behaviors, then it is reasonable to presume that plasticity in the inflammatory response can contribute to the emergence of new genes. This raises the possibility that selection pressures conferred by injury, and specifically the inflammatory responses that are tied to injury, might drive the adaptation of newly emerged genes to new functions in inflammatory cells. To this end, it is noteworthy that the ancestors of homo sapiens acquired three new injury-prone behaviors that are hallmarks of the species. First, when early hominid moved between trees and the savannah, they became erect and bipedal. Second, they learned to use tools and weapons so as to modify

**ABSTRACT**

In light of the central role of inflammation in normal wound repair and regeneration, we hypothesize that the preponderance of human-specific genes expressed in human inflammatory cells is commensurate with the genetic versatility of inflammatory response and the emergence of injuries associated with uniquely hominid behaviors, like a bipedal posture and the use of tools, weapons and fire. The hypothesis underscores the need to study human-specific signaling pathways in experimental models of injury and infers that a selection of human-specific genes, driven in part by the response to injury, may have facilitated the emergence of multifunctional genes expressed in other tissues.

their environment for hunting and compete for habitation. Finally, when hominid harnessed fire and turned to cooking food, their gut microbiome was fundamentally altered. Presumably, the acquisition of these behaviors came with significant risk for trauma and burn injury: bipedal motion and tool/weapon-making skills led to the appearance of new kinds of pressure and penetrating injuries while the use of fire, another uniquely human activity, was likely associated with accidental burn of skin and hair and smoke inhalation. Like no other species, some early human societies are even known for using injury response (scar formation) to confer reproductive advantage.

In modern humans, there are approximately 300 human-specific genes that distinguish the human genome from that of great apes.<sup>4</sup> There are perhaps another 2,000 that differentiate primates from other species.<sup>3</sup> For the most part, however, the reasons for the retention of these genes in the human genome remain to be investigated. Neuroscientists posit that human-specific genes were selected to regulate traits that are uniquely human: size of the human brain, cognition, memory, and consciousness. To others, their very existence has been enigmatic, their functions unknown and the reasons for their over-representation in complex human disease,<sup>5</sup> an interesting, albeit cursory problem in translational research. We hypothesize here that the opposite may be true, and that the very existence


**Table 1.** Human-specific genes and inflammatory cells

Gene name and/or gene family*	Gene ID (Ensembl)	Known, putative and/or inferred function(s)*	Immune and injury cell expression	Reference*
CHRFAM7A	ENSG00000166664	Receptor antagonist	BM, WB, L, LN, T	J Leukoc Biol. 2015; 97:247
TBC1D3	ENSG00000274611	Endocytosis/pinocytosis	BM, WB, L, LN, T	Genomics. 2006; 88:731
ARHGAP11B	ENSG00000187951	Rho GTPase activating	WB, L, LN	Science. 2015; 347:1465
CCL18	ENSG00000275385	Immunoregulation	BM, WB, LN, T	Genomics. 1999; 55:353
CCL23 (MIP-3)	ENSG00000274736	Leukocyte chemotaxis	BM, WB, LN, T	J Exp Med. 1997; 185:1163
NLGN4X (Neuroigin4X)	ENSG00000146938	Tissue remodeling	BM, WB, LN, T	Nat Genet. 2003; 34:27
Interleukin 26	ENSG00000111536	T cell responsiveness	BM, WB, LN, T	J Virol. 2000; 74: 3881
IFNL1	ENSG00000182393	Antiviral host defense	WB, LN	J Biol Chem. 2004; 279:32269
ANGPTL5 (Angiotensin-like)	ENSG00000187151	Tissue regeneration	WB LN	J Hum Genet. 2003; 48:159
Alpha Defensin (N = 5; DEFA1B, A3, A4, A5, A6)	ENSG00000240247	Antimicrobial peptides	BM, WB, L, LN, T	Physiol. Gen 2004; 20: 1
Beta Defensin (N = 5; DEF4B, B104A, B104B, B114, B132)	ENSG00000239839	Antimicrobial peptides	BM, WB, LN, T	Genome Biol. 2003; 4:R31
NPIP genes (N = 2; NPIP-A1 and B11)	ENSG00000164821	Nuclear pore proteins	BM, WB, LN, T	Nature. 2001; 413:514
CSAG (N = 2; CSAG1, CSAG2)	ENSG00000164816	Cell growth and tumor antigen	WB, LN	Gene. 1999; 229:75
c20orf203 (Alugen)	ENSG00000177257	Neurodegeneration	WB	PLoS Comput Biol. 2010; 6:e1000734
SPANX genes (N = 9; SPANXB1, D, A2, C, A1, N3, N2, N5, N1)	ENSG0000016782	Nuclear proteins, spermatogenesis and cancer antigens	BM, WB, LN, T	Proc Natl Acad Sci USA. 2004; 101:3077
	ENSG00000227234			
	ENSG00000196406			
	ENSG00000203926			
	ENSG00000196573			
	ENSG00000198021			
	ENSG00000189252			
	ENSG00000266988			
	ENSG00000204363			
	ENSG00000203923			

**Table 1.** Continued.

Gene name and/or gene family*	Gene ID (Ensembl)	Known, putative and/or inferred function(s)*	Immune and injury cell expression	Reference*
VCX genes (N = 3; VCX2, VCX3A, VCX3B)	ENSG00000182583 ENSG00000177504 ENSG00000169059	mRNA stability reproduction	WB	Cancer Res. 2014; 74:4694
VCY	ENSG00000205642 ENSG00000129864	Nuclear protein	BM, WB, LN, T	Mol Reprod Dev. 2008; 75:219
OPN genes OPN1MW	ENSG00000268221 ENTREZ: 100534624	Locus control for opsin trichromatic vision	BM, WB, LN, T	Trends Ecol Evol 2003; 18:198
VN1R genes (VN1R3)	ENSG00000180663 (mouse, no rat)	Olfactory chemosensory	BM, WB, LN, T	Genome Res. 2010;20:10
STRA6	ENSG00000137868	light adaptation	BM, WB, LN, T	PLoS One. 2014; 9:e108388
NBPF	ENSG00000162825	Development	BM, WB, LN, T	Mol Biol Evol. 2005; 22:2265
CT45 genes (N = 4; CT45A1, A2, A3, A4)	ENSG00000268940 ENSG00000271449 ENSG00000269096 ENSG00000271449	B cell lymphoma	BL	Proc Natl Acad Sci USA. 2005;102:7940

\*See www.genecards.org for details on genes, gene ontogeny, orthologs, paralogs, inferred and known functions and relative tissue expression. BM, bone marrow; BL, B cell lymphoma; WB, whole blood; LN, lymph node; T, thymus; LL, leukocytes and lymphocytes as determined by microarray.



**Figure 1.** Injury, inflammation, and adaptation. New hominid behaviors, like the emergence of bipedal movement (trauma), harnessing of fire (burn injury) and the use of tools and weapons (injury/infection) is a positive selection for new genes. Positive adaptive selection enables new inflammatory activities (A) in response to new types of injuries. In this model, genes like *CHRFAM7A*, *TBC1D3*, and *ARHGAP11B* are initially selected in leukocytes and the in the human inflammatory response for their capacity to regulate ligand binding, internalization, and signaling, respectively. When these same genes are expressed in other tissues however (B), they can elicit new activities that also contribute to positive selection for example regulating neurotransmitter action (*CHRFAM7A*), cell–cell communication (*TBC1D3*) and progenitor cell growth (*ARHGAP11B*).

of human-specific genes may help explain intrinsic problems in the representation of human inflammatory disease in animal models.

In an *in silico* survey of candidate taxonomically restricted genes, we were struck by the number of human-specific genes that are expressed in blood-derived cells, most often leukocytes and lymphocytes (Table 1). We also noted that the expression of these genes is often associated with the biology of injury, namely infection, inflammation, and tissue repair and regeneration. These genes include well-known anti-infection and human-specific defensin genes (Table 1), some of which specifically localize to human neutrophils. They also include more recently uncovered genes, like *CHRFAM7A*, *ARHGAP11B* and *TBC1D3* that are either linked injury responses like the cholinergic regulation of inflammation by the  $\alpha 7$ -nicotinic acetyl choline receptor ( $\alpha 7$ -nAChR), to growth factor-mediated signal transduction through *rho*-mediated signal transduction or to the regulation of microvescle exocytosis and endocytosis, respectively (Table 1).

Numerous investigators have noted the significant variability that exists in the resilience of different species to different pathogens, but species differences in injury and scarring is largely underinvestigated or often minimized. Yet, trauma and burn surgeons attest to the intrinsic variability of the inflammatory response after burn and traumatic injury in humans that are otherwise indistinguishable.<sup>6</sup> Interestingly, just as we noted the expression of human-specific genes in human immune cells

(Table 1), Long and colleagues noted the wide and variable representation of primate- and human-specific protein-coding sequences in brain development.<sup>3</sup> We, therefore, considered the possibility that genes, originally selected to modulate human resilience to injury and infection, could have multifunctional biological effects, for example, when expressed in other tissues (Figure 1). In one such example, *CHRFAM7A*<sup>7</sup> is a proinflammatory human-specific gene that is a dominant negative inhibitor of the anti-inflammatory  $\alpha 7$ -nicotinic acetylcholine receptor ( $\alpha 7$ AChR) in peripheral tissues like macrophages. When expressed in the central nervous system, it presumably alters  $\alpha 7$ AChR activities on human cognition and memory. In other examples, the human antimicrobial defensins are highly multifunctional in different tissues<sup>8</sup> and the human-specific *TBC1D3* gene regulates human macropinocytosis and the formation of human microvesicles.<sup>9</sup>

So what are the consequences of an anthropogenic injury hypothesis to better understanding the mechanistic basis to human disease, and particularly wound repair and regeneration? First, like most genes, the expression of human-specific genes in different tissues is highly contextual. Their multifunctional activities are dependent on where and when they are expressed (Figure 1). In this proposed paradigm, a human-specific gene in monocytes may have arisen to alter responsiveness after injury but may be retained in the human genome for unrelated activities, for example, in the central nervous system. Second, current experimental models of human disease do not test the contribution(s) of human-specific genes to disease. To this end, the mouse still remains the most robust and experimentally tractable vertebrate animal model of human disease and it is possible to ask whether imparting mice with a human-specific gene alters their injury response. Plasmids encoding human-specific genes-like *ARHGAP11B* have been injected into mice to generate a human phenotype.<sup>10</sup> Alternatively, human genes (e.g., *CHRFAM7A*) that encode antagonists (e.g., for *CHRNA7*) can be studied in mice to determine if over expression mimics the proinflammatory and behavioral phenotype of target gene knockout (e.g., *CHRNA7*). Finally, endogenous human-specific genes in circulating and resident human immune cells can be studied in mice after the transplantation and engraftment of human hemato-lymphoid immune systems.<sup>11</sup> With heterologous or autologous human skin grafts, it is even possible to create precision models of human diseases, including for wound repair and regeneration.

On a final note, the possibility that injury, inflammation, and infection responses contribute to the selection of species-specific genes and regulatory pathways implies that they contribute factors to injury and wound healing responses in modern humans. If so, their study should be integrated into animal models of injury, wound healing and inflammation research. This is particularly relevant to “characteristically human” diseases like keloids and certain inflammatory bowel diseases that are difficult to model in animals and whose origins might trace to human behaviors like scarring and changes in the gut microbiome. Along this line, it is interesting to note that genomic expression data,<sup>12</sup> albeit controversial,<sup>13–15</sup> already points to the existence of characteristically human gene expression patterns in response to infection, trauma, and burn injury. We propose that while a better understanding of the contributions of species-specific genes to the injury and inflammation response adds another level of complexity to biomedical research, it could lead to the identification of

completely new drugs, new drug targets and the development of new therapeutics.

## ACKNOWLEDGMENTS

*Conflict of Interest:* The authors declare that they have no conflicts of interest.

*Source of Funding:* The biomedical hypothesis presented in this manuscript initially arose during Supplementary Research funding from NIGMS (NIGM78421) to AB and BPE through the American Recovery and Reinvestment Act (ARRA) and was further developed with support of the National Cancer Institute (CA170140) to BPE, the Hammond Fund (RC), the Division of Trauma Reinvestment Fund (AB, TC, RC and BPE), the American Surgical Association Foundation Research Fellowship (TC), and a Department of Defense grant (W81XWH-10-1-0527) to RC that is administered by the American Burn Association.

## REFERENCES

- Acharya D, Mukherjee D, Podder S, Ghosh TC. Investigating different duplication pattern of essential genes in mouse and human. *PLoS One* 2015; 10: e0120784.
- O'Bleness M, Searles VB, Varki A, Gagneux P, Sikela JM. Evolution of genetic and genomic features unique to the human lineage. *Nat Rev Genet* 2012; 13: 853–66.
- Zhang YE, Landback P, Vibranovski M, Long M. New genes expressed in human brains: implications for annotating evolving genomes. *Bioessays* 2012; 34: 982–91.
- Long M, VanKuren NW, Chen S, Vibranovski MD. New gene evolution: little did we know. *Ann Rev Genet* 2013; 47: 307–33.
- Cooper DN, Kehrer-Sawatzki H. Exploring the potential relevance of human-specific genes to complex disease. *Hum Genomics* 2011; 5: 99–107.
- Harrison C. Sepsis: calming the cytokine storm. *Nat Rev Drug Discov* 2010; 9: 360–1.
- Costantini TW, Dang X, Coimbra R, Eliceiri BP, Baird A. CHRFAM7A, a human-specific and partially duplicated alpha7-nicotinic acetylcholine receptor gene with the potential to specify a human-specific inflammatory response to injury. *J Leukoc Biol* 2015; 97: 247–57.
- Cheng DQ, Li Y, Huang JF. Molecular evolution of the primate alpha-/theta-defensin multigene family. *PLoS One* 2014; 9: e97425.
- Frittoli E, Palamidessi A, Pizzigoni A, Lanzetti L, Garre M, Troglio F, et al. The primate-specific protein TBC1D3 is required for optimal macropinocytosis in a novel ARF6-dependent pathway. *Mol Biol Cell* 2008; 19: 1304–16.
- Florio M, Albert M, Taverna E, Namba T, Brandl H, Lewitus E, et al. Human-specific gene ARHGAP11B promotes basal progenitor amplification and neocortex expansion. *Science* 2015; 347: 1465–70.
- Rongvaux A, Willinger T, Martinek J, Strowig T, Gearty SV, Teichmann LL, et al. Development and function of human innate immune cells in a humanized mouse model. *Nat Biotechnol* 2014; 32: 364–72.
- Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker HV, Xu W, et al. Genomic responses in mouse models poorly mimic human inflammatory diseases. *Proc Natl Acad Sci USA* 2013; 110: 3507–12.
- Takao K, Miyakawa T. Genomic responses in mouse models greatly mimic human inflammatory diseases. *Proc Natl Acad Sci USA* 2015; 112: 1167–72.
- Warren HS, Tompkins RG, Moldawer LL, Seok J, Xu W, Mindrinos MN, et al. Mice are not men. *Proc Natl Acad Sci USA* 2015; 112: E345.
- Osuchowski MF, Remick DG, Lederer JA, Lang CH, Aasen AO, Aibiki M, et al. Abandon the mouse research ship? Not just yet! *Shock* 2014; 41: 463–75.