



UNIFORMED SERVICES UNIVERSITY, SCHOOL OF MEDICINE GRADUATE PROGRAMS
Graduate Education Office (A 1045), 4301 Jones Bridge Road, Bethesda, MD 20814




DISSERTATION APPROVAL FOR THE DOCTORAL DISSERTATION IN THE NEUROSCIENCE
GRADUATE PROGRAM

Title of Dissertation: "GABA_A receptor-mediated activity in a model of cortical dysplasia"


Name of Candidate: Joseph Abbah
Doctor of Philosophy Degree
June 29, 2012

DISSERTATION AND ABSTRACT APPROVED:



Dr. Maria F. Braga
DEPARTMENT OF ANATOMY, PHYSIOLOGY, AND GENETICS
Committee Chairperson

DATE:


6/29/12


Dr. Sharon L. Juliano
DEPARTMENT OF ANATOMY, PHYSIOLOGY, AND GENETICS
Dissertation Advisor

6/29/12


Dr. Martin Doughty
DEPARTMENT OF ANATOMY, PHYSIOLOGY, AND GENETICS
Committee Member

6/29/12


Dr. Xin Xiang
DEPARTMENT OF BIOCHEMISTRY
Committee Member

6/29/12


Dr. Joshua Corbin
CHILDREN'S NATIONAL MEDICAL CENTER AND GEORGE WASHINGTON UNIVERSITY

6/29/12



UNIFORMED SERVICES UNIVERSITY, SCHOOL OF MEDICINE GRADUATE PROGRAMS
Graduate Education Office (A 1045), 4301 Jones Bridge Road, Bethesda, MD 20814



FINAL EXAMINATION/PRIVATE DEFENSE FOR THE DEGREE OF DOCTOR OF PHILOSOPHY IN
IN THE NEUROSCIENCE GRADUATE PROGRAM






Name of Student: Joseph Abbah

Date of Examination: June 29, 2012

Time:

Place:

DECISION OF EXAMINATION COMMITTEE MEMBERS:

	PASS	FAIL
 Dr. Maria R. Braga DEPARTMENT OF ANATOMY, PHYSIOLOGY, AND GENETICS Committee Chairperson	<u>X</u>	—
 Dr. Sharon L. Juliano DEPARTMENT OF ANATOMY, PHYSIOLOGY, AND GENETICS Dissertation Advisor	<u>X</u>	—
 Dr. Martin Doughty DEPARTMENT OF ANATOMY, PHYSIOLOGY, AND GENETICS Committee Member	<u>X</u>	—
 Dr. Xin Xiang DEPARTMENT OF BIOCHEMISTRY Committee Member	<u>X</u>	—
 Dr. Joshua Corbin CHILDREN'S NATIONAL MEDICAL CENTER AND GEORGE WASHINGTON UNIVERSITY	<u>X</u>	—

The author hereby certifies that the use of any copyrighted material in the thesis manuscript entitled:

'GABA_A receptor-mediated activity in a model of cortical dysplasia'

is appropriately acknowledged and, beyond brief excerpts, is with the permission of the copyright owner.



Joseph Abbah

Program in Neuroscience

Uniformed Services University

07/09/2012

COPYRIGHT AND DISCLAIMER

The author certifies that the use of any copyrighted material in this dissertation manuscript entitled:

“GABA_A receptor-mediated activity in a model of cortical dysplasia”

and aside from brief passages, it is with the permission of the copyright holder. The author holds harmless the Uniformed Services University from damage which may arise from any copyright infringement.

The views expressed in this work are those of the author and do not reflect those of the United States Department of Defense, or the United States Governmental Institution.

Joseph Abbah

Graduate Program in Neuroscience

Uniformed Services University of the Health Sciences

Title of Dissertation: GABA_A RECEPTOR-MEDIATED ACTIVITY IN A
MODEL OF CORTICAL DYSPLASIA

By

Joseph Abbah, Doctor of Philosophy, 2012

Thesis directed by: Sharon L. Juliano, Ph.D.
Professor, Department of Anatomy, Physiology and
Genetics, Neuroscience Program Director

ABSTRACT

Cortical dysplasia is a developmental abnormality characterized by changes in the structure and function of the neocortex. This disorder is implicated in many neuropsychiatric disorders such as epilepsy, autism and schizophrenia, and results from failure of immature neurons to appropriately migrate to their cortical targets. We developed a model of cortical dysplasia by administering methylazoxymethanol (MAM), an anti-mitotic, to pregnant ferrets on gestational day 33 leading to abnormal cortical layering, increased expression of GABA_A receptors (GABA_{AR}) and altered distribution of neurons derived from the ganglionic eminence (GE).

We evaluated the impact of MAM treatment on the kinetic behavior of migrating GE cells and examined the relationship between changes in the ambient level of GABA_{AR}-mediated activity and alterations in neuronal migration

in an organotypic cultures of postnatal day 0 to 1 (P0 - P1). In addition, we investigated the functional implication of altered neuronal migration in the cortex of dysplastic animals using whole-cell patch-clamp recording and western blot assay. The normal dynamic pattern of migration of GE-derived cells alters after MAM treatment. Cells originating in the GE of normal ferrets migrate significantly faster and greater proportion of these cells demonstrates 'exploratory behavior' compared with cells in model animals. Treatment with MAM also increased the expression of GABA_{Aα2}, GABA_{Aα3} receptor subunits and the neuron-specific potassium chloride co-transporter in the cortex of P0 – P1 animals suggesting that GABA_{AR}-mediated activity increases in our model animals. In support, the deficit in migration in MAM-treated GE cells improves when GABA_{AR} activity is blocked.

Functional analysis showed that treatment with MAM significantly increased the amplitude and frequency of GABA_{AR}-mediated spontaneous inhibitory post-synaptic currents (sIPSCs) in GE cells. In older MAM-treated animals, the amplitude of sIPSCs in layer 2/3 pyramidal cells increased with no significant effect on frequency. The amplitude and frequency of miniature IPSCs, however, were significantly increased in the MAM-treated pyramidal cells. The expression of GABA_{Aα3} receptor subunit was increased following MAM treatment. These data suggest that treatment with MAM increases GABA signaling within the neocortex, which impairs migration and distribution of GE-derived cells to alter the microcircuitry and function of the resultant cortex.

**GABA_A RECEPTOR-MEDIATED ACTIVITY IN A MODEL OF CORTICAL
DYSPLASIA**

By

Joseph Abbah

Doctoral Dissertation submitted to the Faculty of the Neuroscience Graduate
Program at the Uniformed Services University of the Health Sciences in partial
fulfillment of the requirements for the degree of
Doctor of Philosophy, 2012.

PREFACE

This thesis is based upon work performed in the laboratory of Dr. Sharon Juliano at the Uniformed Services University of the Health Sciences between September 3, 2007 and June 22, 2012.

DEDICATION

To my beloved wife, Faith Abbah, our beautiful son, Joseph (jnr) Owoicho Adah Joseph, and the living memory of our lovely daughter, Emily Enayi Joseph (who left us too soon to be with the Lord) - for their love and inspiration; and to the beautiful memories of my parents- Enayi Abah and Audu Abah- for making me recognize the value of education very early in life.

ACKNOWLEDGEMENTS

This work is the product of the efforts of many individuals who made tremendous contributions in building my capacity as a professional and scientist; the tribute I pay here is an inadequate reflection of my personal sense of gratitude to them and what they have meant to me.

I am deeply indebted to Dr Sharon L. Juliano for her faith in me, for her support and mentorship, which extend beyond the confines of academia. Throughout my time in the program, she was a steady influence and guidance to me and my family. Although my journey into graduate school did not begin with my chanced meeting with her in South Africa, nonetheless, she was a major catalyst in my quest for admission; I cannot imagine under what scenario I would have ended up at USUHS had I not met her. My personal experience with her is both a reflection and a tiny representation of her dedication towards expanding the culture of neuroscience to technologically and scientifically-challenged regions of the world, sometimes at great risk to her security and life.

I want to specially thank members of my thesis committee-Dr Maria Braga (Chair), Dr Martin Doughty, Dr Joshua Corbin, Dr Xing Xiang, for their invaluable inputs, criticisms, suggestions and useful comments, which greatly improved the quality of this work.

Graduate school is a challenging task made more daunting when undertaken outside a socio-cultural comfort zone. As an international student from Nigeria, a lot of what I had to deal with in the beginning were completely

new to me and a good amount of effort was expended on adjusting rapidly to my new environment and responsibilities. This effort was aided immensely by the warmth with which I was received, the guidance and support that I enjoyed from my colleagues and faculties, and the generous outpouring of encouragement from members of the USUHS community. In particular, I like to thank my classmates- Mike Bentley, Anton Dmitriev, Nicole Flint, Nora Hibits and especially Megan Rose Hershfield for creating a very cordial class room environment into which I was seamlessly integrated. Thank you guys!

My colleagues-current and former- in Dr Juliano's lab have an unbelievable capacity for fun that eases the stress of running lab experiments; they are an incredible bunch of fellas to work with. I appreciate the scientific assistance Dr Marcin Gierdalski and Dr Sylvie Poluch freely rendered; their advice and criticisms were very useful. Dr Tom McFate and Dr John Trentini's empathy and support towards securing a visa for my wife will forever be remembered. Latoya Hyson's patient support and extra efforts in practically 'baby sitting' the ferrets was exemplary and outstanding. Other members of the lab- Dr Deb McLaughlin, Dr Alisa Schaeffer, Dr Nate Craymer, Dr Susan Schwerin, Dr Kapinga Ngalula, Dr Kwame Affram, Dr Olu Akinola, Mara Schindell, Mitali Charterjee, Neha Datta, were quite supportive throughout. Dr Volodymyr Pidoplichko was very helpful with some aspects of the project.

I owe a great depth of gratitude to Dr Gamaniel Karniyus and Dr Yakubu Ngwai who recruited me to my first job and gave me the initial opportunity to deeply appreciate science. Dr Gamaniel's support and mentorship has

consistently guided me in every step of this journey, which he, with a touch of clairvoyance, predicted. I will always remember the tutelage I enjoyed under Dr Habiba Usman, Dr Amos Samson, Dr Chindo Ben, and the rest of my colleagues in Pharmacology and Toxicology Dept of NIPRD.

My gratitude goes to my family members who, in various measures, have been a constant positive force in my life, which continues to propell me towards greater heights. My parents, Audu Abah and Enayi Abah, who exposed me to the values of education and applied their meager incomes towards that goal, and who inculcated in me the ideals of hard work, commitment, dedication, and a belief that I could achieve whatever goal I set my sights on, they are the architects of this success. Although they are no longer here with us to bear witness to this achievement, am certain that as they look down from heaven, they would be extremely proud of this moment, which I want to dedicate to them. I want to specially thank Fred Abah for his selfless investment and constant encouragement to all of us to reach for our greatest potential. I cannot thank Salamatu E. Abah enough for her crucial financial support from the time I was in high school through college and beyond. She has been a defacto Mum to us. I acknowledge with gratitude the generosity of Adah Abah whose 'widow's might' assistance meant so much more to me than she could ever imagine. Alice N. Odeh has persistently supported my efforts and maintained a sustained interest in my progress. I want to thank my brothers Ibrahim Abba, Suleiman Abba, Mohammed F. Abah, Alfred Abah, Ongaji Abah, Mohammed Onoja, Sunday Oluma, Adakole Awuja, Godwin Ugana, Audu Ugana, Moses A. Odeh, Adah

Adugba, Emmanuel Oche, Elijah Abah for your love, prayers and the rich memories we shared; my sisters, Maminetu Abah, Alice Abba, Agnes Edoh, Onyema Onum, Cecilia Abah, Margaret Abah, Fatima Okodeje, Onyoibo Abah; my aunt, Aladi Abah; my in-laws, Agnes Abah, Helen Abah, Husseina Abba, Stephen Edeh, Victoria A. Edeh, Cynthia Edeh and those who, as a result of space constraints, are not mentioned here, for all you have meant to me.

I deeply appreciate the role played by John Oduh and his wife Aisha Oduh, in laying the foundation for my professional career. I would have literally starved and dropped out of school had it not been for their generosity when I was in college. They have maintained the same level of commitment to me up to this moment. I benefited immensely from Mike Onum's humility and selfless support at crucial periods during the program. The world needs more John Oduhs and Mike Onums.

My friends Bernard Akpa, Tony Eti-Ukwu, Nicodemus Nanven, Tags Zacharias, Jiru Bako, Julius Ogunleye, have been an unbelievable source of inspiration and support to me.

The Idoma community in the DMV area is quite remarkable for the friendship and sense of solidarity they have built over the years. I benefited from my association and friendship with all of you. In particular, I want to thank Matthew Abah and Deborah Abah for their fantastic support. I also want to express my appreciation to Patrick Akatu, George Akoji, Grace Akoji, Francis Obeya, Dr Ene Obeya and Lawrence Ogbe for their care and concern throughout

this program. Pharm. Steve Onyilokwu and Beatrice O. Onyilokwu were magnificent in their support for the family. Pharm Prince A. Adekoya's professional assistance will always be appreciated.

To my beautiful wife, Faith Ene Abbah, whose love and care for the family allowed me time and space to selfishly focus on my career, I will eternally remain grateful. Life took a different meaning, expanding in scope and richness when you came into my life. Your unwavering support and belief in this project was very reassuring at very tough periods and helped strengthened my resolve and faith. I love you. Our son Joseph is a bundle of joy that sustained my spirit in the latter stage of study. Joseph, I know even in your immature thoughts (your brain cells are still migrating and forming networks) you have always wondered why am always away from home, why am unavailable to play soccer with you, this is the reason why. It is my hope that you will one day come to appreciate this and explain to your own children. My daughter, Emily Enayi, gave me so much hope but left us rather too soon and took a lot away from us. The cloud of gloom and despair that enveloped us when you departed was profoundly felt, but our hearts are lifted by the belief that you are resting with the Lord. We will always love you.

Finally, I want to thank God for making everything possible, for his faithfulness and blessings. The early phase of my life never offered any clue that I will be a high school graduate let alone attend graduate school in the US. But he made all things possible. Thank you Lord.

TABLE OF CONTENTS

APPROVAL SHEET.....	i
COPYRIGHT STATEMENT.....	ii
ABSTRACT.....	iii
TITLE PAGE.....	v
PREFACE.....	vi
DEDICATION.....	vii
ACKNOWLEDGEMENTS.....	viii
TABLE OF CONTENTS.....	xiii
LIST OF ABBREVIATIONS.....	xvi
CHAPTER 1- INTRODUCTION.....	1
CHAPTER 2- ALTERED KINETIC BEHAVIOR UNDERLIES REDISTRIBUTION OF INTERNEURONS IN A MODEL OF CORTICAL DYSPLASIA: THE INFLUENCE OF ELEVATED GABA _A RECEPTOR ACTIVITY.	

Title Page.....	28
Abstract.....	29
Introduction.....	30

Materials and Methods.....	33
Results.....	37
Discussion.....	45
Acknowledgements.....	52
References.....	53
Figures.....	64

Chapter 3- SELECTIVE DISRUPTION OF LAYER 4 FORMATION ENHANCES GABA_A RECEPTOR-MEDIATED ACTIVITY IN LAYER 2/3 OF FERRET SOMATOSENSORY CORTEX.

Title Page.....	75
Abstract.....	76
Introduction.....	78
Materials and Methods.....	82
Results.....	87
Discussion.....	93
Acknowledgements.....	102
References.....	103
Figures.....	119

CHAPTER 4- DISCUSSION.....	125
REFERENCES.....	140

LIST OF ABBREVIATIONS

aCSF	Artificial cerebrospinal fluid
AMPA	α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
AP	Action potential
BMI	Bicuculline methiodide
BSA	Bovine serum albumin
CGE	Caudal ganglionic eminence
CNQX	6-Cyano-7-nitroquinoxaline-2,3-dione
CVZ	Cortical ventricular zone
Dil	1,1'-Diocetadecyl-3,3,3',3'-Tetramethylindocarbocyanine perchlorate
E33	Embryonic day 33 (E1= day vaginal plug is seen)
GABA	Gamma (γ)-aminobutyric acid
GE	Ganglionic eminence
KCC2	Potassium- Chloride Co-transporter
LGE	Lateral ganglionic eminence
MAM	Methylazoxymethanol acetate
MGE	Medial ganglionic eminence
MK801	(5R, 10S)-(-)-5-Methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5,10-imine maleate
MZ	Marginal zone
NB media	Neurobasal media (containing B27, N2 and G1.2. supplements)
NKCC1	Sodium Potassium Chloride Co-transporter
NMDA	N-methyl-D-aspartate

PBS	Phosphate buffered saline
RA	Retinoic acid
RG	Radial glia cells
SCH50911	(2S)-(+)-5,5-Dimethyl-2-morpholine acetic acid
SVZ	Subventricular zone
TTX	Octahydro-12-(hydroxymethyl)-2-imino-5,9:7,10a-dimethano-10a <i>H</i> - [1,3]dioxocino[6,5- <i>d</i>]pyrimidine-4,7,10,11,12-pentol citrate
VZ	Ventricular zone

CHAPTER 1

INTRODUCTION

Cortical dysplasia (CD) is a developmental abnormality characterized by changes in the structure and function of the cerebral cortex. This disorder is implicated in many neuropsychiatric disorders such as intractable epilepsy, schizophrenia, and autism, and results from failure of immature neurons to appropriately migrate and reach their cortical targets (Taylor et al., 1971; Choi and Mathias, 1987; Tassi et al., 2002; Calcagnotto and Baraban, 2003; Moroni et al., 2008). By some estimates, CD is responsible for 8-12% of all clinical cases of epilepsy that fail to respond to standard drug treatment, and 14-26% of pediatric cases of epilepsy in which surgical intervention is indicated (Li et al., 1995; Polky, 1996; Fauser et al., 2012). During development, the cerebral cortex, which consists of six distinct cortical layers, is constructed through a sequence of time-dependent processes that includes generation, migration and subsequent differentiation of two principal neuronal cell types: projection neurons, primarily pyramidal cells, and interneurons. These two major classes of neurons derive from distinct neurogenic regions of the developing cortex and use different modes of migration to reach their cortical targets (Parnavelas et al., 1991; Mione et al., 1994; Parnavelas, 2000; Nadarajah and Parnavelas, 2002; Marin and Rubenstein, 2003; Kriegstein and Noctor, 2004; Ayala et al., 2007). Over the years, studies conducted in invertebrates and rodents revealed some of the molecular mechanisms underlying the complex processes of neurogenesis and neuronal migration; these investigations demonstrate that the process of neurogenesis and neuronal migration are similar across species (Garcia-Bellido,

1979; Villares and Cabrera, 1987; Ghysen and Dambly-Chaudiere, 1988; Gonzalez et al., 1989; Jarman et al., 1993; Bertrand et al., 2002). These findings form the basis for extrapolation of developmental events observed in lower animals to higher organisms including humans. However, it is apparent from recent studies that important differences exist in the process of cortical development between lissencephalic animals such as rodents and gyrencephalic organisms, such as ferrets and primates. For instance, gyrencephalic animals such as ferrets and primates have protracted neurogenesis leading to a significant increase in population of neurons generated, as well as the size and features of the resulting cortex (Poluch et al., 2008; Fietz et al., 2010; Martinez-Cerdeno et al., 2012). In addition, the site of origin and migratory characteristics of interneurons in higher organisms differ from lower animals (Poluch et al., 2008; Letinic et al., 2002; Fietz et al., 2010). These differences underscore the need to study the process of corticogenesis under normal and diseased conditions in animals with a convoluted cortex.

Origin of Projection neurons. Projection neurons are usually characterized by a pyramidal morphology and use glutamate as their neurotransmitter. It is widely accepted that despite the unique pyramidal morphology of these cells, cortical projection neurons in different cortical layers have different molecular signatures, and are generated sequentially from stem and progenitor cells in neocortical germinal zones in the dorsal telencephalon including the ventricular zone (VZ) and the subventricular zone (SVZ) (Rakic, 1988; Walsh and Cepko, 1988, 1993). Both symmetric and asymmetric division

of stem cells in the germinal zones and their progeny generates the repertoire of projection neurons in the cortex (Noctor et al., 2001a, 2004). Neurogenesis initiates when stem cells in the neuroepithelium undergo symmetric division to generate daughter stem cells (with same molecular properties as the parent stem cells) followed by asymmetric division to produce radial glia (RG) cells (Haubensak et al., 2004; Noctor et al., 2004; Gotz and Huttner, 2005; Kriegstein and Alvarez-Buylla, 2009; Hansen et al., 2011). Within the ventricular zone (VZ), RG cells, which have stem cell properties, divide symmetrically or asymmetrically to generate daughter cells that are also RG cells, while simultaneously generating other daughter cells that represent committed neuronal progenitors. Subsequently, the progenitor cells in the subventricular zone (SZ) undergo further cycles of symmetric cell division to produce committed neuroblasts that exit the cell cycle and migrate to their final cortical target (Gray et al., 1988; Luskin et al., 1988; Price and Thurlow, 1988; Grove et al., 1993; Chen and McConnell, 1995; Kornack and Rakic, 1995; McConnell, 1995; Rakic, 1995; Mione et al., 1997; Reid et al., 1997; Qian et al., 1998 and 2000; Malatesta et al., 2000; Miyata et al., 2001; Noctor et al., 2001a; Shen et al., 2002; Gotz et al., 2002; Fishell and Kriegstein, 2003; Kriegstein and Gotz, 2003; Noctor et al., 2004; Gotz and Huttner, 2005; Gal et al., 2006; Kriegstein and Alvarez-Buylla, 2009).

Mammals have six cortical layers populated by pyramidal cells and interneurons; the position of projection neurons is determined by the time of birth, corresponding to exit of the cell cycle (Caviness and Sidman, 1973; Caviness,

1982; Holt et al., 1988; Takahashi et al., 1999; Ohnuma and Harris, 2003). The initial group of cells migrate from the VZ; the cortical hem is also likely to provide a cohort of Cajal Retzius cells to form the preplate (Yoshida et al., 2006). This zone is split by a surge of cortical cells to create the cortical plate, now sandwiched between the more superficial MZ that includes Cajal Retzius cells and the deeper subplate layer (Marin-Padilla, 1998). Subsequent waves of cortical cells situate themselves between these two layers (in the cortical plate) in such a way that cells occupying the deep layers of 6 and 5 are born first followed by cells of the more superficial layers, described as an inside-out pattern of corticogenesis (Angevine and Sidman, 1961; Rakic, 1974; Bayer and Altman, 1991; Shen et al., 2006).

The specification and progressive generation of pyramidal cells is regulated by a molecular program involving several transcriptional factors with distinct spatial expressions. Whereas all pyramidal neurons, irrespective of layer identity, are specified by *Neurogenin 1* and *Neurogenin 2* (*Neurog1* and *Neurog2*), basic-helix-loop-helix transcriptional factors expressed in the VZ of the dorsal telencephalon (Fode et al., 2000; Bertrand et al., 2002; Ross et al., 2003; Schummanns et al., 2004; Guillemot, 2007), the molecular identity of pyramidal cells is controlled by downstream effector genes that regulate both the layer identity and axonal projections and targets of the different types of these cells (Fishell and Hanashima, 2008). For instance, projection neurons destined to the deep layers of 5 and 6 express *Sox5*, *Fezf2*, *Ctip2* and *Tbr1* and project to subcortical regions of the brain (DeFelipe and Farinas, 1992; Koster and O'leary,

1993; Kristin et al., 2002; Arlotta et al., 2005; Chen et al., 2005a, 2005b; Molyneaux et al., 2005, 2007; Chen et al., 2008). On the other hand, pyramidal cells in upper cortical layers of 2 to 4 express *Satb2* and project to intracortical regions (Britanova et al., 2008).

Migration of projection neurons. Regardless of laminar position, projection neurons born in the neocortical VZ or SVZ migrate radially, using two migratory approaches to reach their final cortical position (Marin and Rubenstein, 2003). A subset of migrating projection neurons use radial glial support and adopt 'glial guided' locomotion (Rakic, 1971, 1972; O'Rourke et al., 1992; Anton et al., 1996; Nadarajah et al., 2001). Projection neurons that use this migratory strategy attach to radial glial fibres that stretch across the entire cortical wall, as a supporting framework. Other projection neurons reach the cortical plate via a somal translocation mechanism in which cells migrate independent of radial glia support (Morest, 1970; Rakic, 1972, O'Rourke et al., 1992; Nadarajah and Parnavelas, 2002; Nadarajah et al., 2001). Migration of pyramidal cells proceeds through distinct phases that include the following steps: (1) migrating cells become physically attached to radial glia fibres (those that adopt glia-guided locomotion), (2) development and extension of a leading process, (3) nucleokinesis that involves translocation of the nucleus in the direction of the leading process (4) retraction of the trailing process and (5) detachment of cell from radial glia fiber (for glial guided migrating cells) and adoption of a cortical position (de Rouvroit and Goffinet, 2001; Honda et al., 2003; Marin and Rubenstein, 2003).

Regulation of radial migration. Each stage of the migratory process is regulated by both environmental cues and intrinsic signaling mechanisms that act in concert to modulate the cytoskeletal network of cells. Among the identified factors implicated in influencing pyramidal cell migration are motogens such as neurotrophins (NT4 and BDNF), epidermal growth factors, neurotransmitters such as GABA acting on GABA_A and GABA_B receptors and glutamate acting on both NMDA and AMPA receptors (Behar et al., 1996, 1998, 1999, 2000, 2001; Komuro and Rakic, 1993; Kornblum et al., 1997; Nakagawa et al., 1998; Marin and Rubenstein, 2001, 2003). Genes (e.g. *Lis1*, *Doublecortin*, *Filamin*), which regulate the function of the neuronal cytoskeleton to control locomotion, and cues that signal migrating cells to cease locomotion and detach from radial glia cells are also crucial in regulating the migratory process (Cavinness, 1982; Reiner et al., 1993; Hattori et al., 1994; D'Arcangelo et al., 1995; Hirotsune et al., 1995; Ogawa et al., 1995; Eksioglu et al., 1996; Frotscher 1997; Sheppard and Pearlman, 1997; Dulabon et al., 2000; Feng and Walsh, 2001).

Genes including *Neurogenin 1* and *Neurogenin 2* (*Neuog1* and *Neuog2*) play crucial roles in the migration of these cells (Hand et al., 2005; Ge et al., 2006; Nguyen et al., 2006). *Neuog1/2* promotes neuronal migration by inducing expression of *Dcx* and *p35*, while simultaneously repressing the activity of RhoA, a GTPase (Ge et al., 2006; Nobrega-Pereira et al., 2009). *Neuog1/2* also influence the migration of pyramidal cells by regulating (increasing) the expression of *Rnd2*, another GTPase (Nakamura et al., 2006; Heng et al., 2008).

On reaching the cortical plate, migrating cortical projection neurons assume laminar positions within the developing cortical wall and terminate migration. The ability of pyramidal cells to find an appropriate cortical target is regulated by *Brn1* and *Brn2*, POU-domain transcription factors that regulate the reelin/DAB1 pathway, and cyclin-dependent kinase (CDK5) signaling; this pathway plays critical roles in normal cortical layer formation (McEvilly et al., 2002; Sugitani et al., 2002; Nobrega-Pereira et al., 2009). Understandably, loss of function of any of the key players above due to mutation leads to abnormal cortical layering with inversion of cortex (D'Arcangelo et al., 1995; Hirotsune et al., 1995; Ogawa et al., 1995; Ohshima et al., 1996; Chae et al., 1997; Kwon and Tsai, 1998; Howell et al., 1999).

Interneurons. Cortical interneurons are a heterogeneous population of cells that originate from germinal zones under the regulatory control of a molecular program (Markram et al., 2008; Ascoli et al., 2008; Corbin and Butt, 2011). In rodents, these cells arise from three distinct germinal zones within the ventral telencephalon: the VZs of the lateral ganglionic eminence (LGE), the medial ganglionic eminence (MGE) and the caudal ganglionic eminence (CGE) (Van Eden et al., 1989; DeDiego et al., 1994; De Carlos et al., 1996; Anderson et al., 1997; Tamamaki et al., 1997; Lavdas et al., 1999; Sussel et al., 1999; Wichterle et al., 1999; Corbin et al., 2001; Letinic et al., 2002; Nery et al., 2002; Yuste, 2005; Wonders and Anderson, 2006; Batista-Brito and Fishell, 2009). An overwhelming majority of cortical interneurons are generated in the GE (Corbin et al., 2001; Wonders and Anderson, 2006). These cells express *GAD65/67* and

make up roughly 20% of the total neuronal cell population in the neocortex and use GABA as their primary neurotransmitter (Hendry et al., 1987; Meinecke and Peters, 1987; Hatten, 2002; Tamamaki et al., 2003). Similar to pyramidal cells, the concerted action of proneural genes of the bHLH motif such as *Mash1*, homeodomain transcription factors such as *Dlx1/2*, *Dlx 5/6*, *Nkx2.1*, *Nkx6.2*, *Lhx6* and *Gli1*, and other signaling molecules such as Sonic hedgehog in the GE specify and generate the different types of interneurons destined for the neocortex and non-cortical structures (Chiang et al., 1996; Anderson et al., 1997; Casarosa et al., 1999; Sussel et al., 1999; Xu et al., 2004; Butt et al., 2008; Corbin et al., 2008; Welagen and Anderson, 2011). The ability of the GE to generate diverse interneuron subtypes is due to the expression of distinct transcription factors in different regions of the GE (Butt et al., 2007; Corbin and Butt, 2011). For instance, *Gli1* and *Nkx6.1* show restricted expression in the dorsal part of the MGE (Corbin et al., 2008; Welagen and Anderson, 2011), this fact has been used to explain why the dorsal MGE is the source of somatostatin-expressing GE cells while the ventral MGE generates parvalbumin-expressing GE cells (Fogarty et al., 2007; Wonders et al., 2008), a view not supported by others (Yu et al., 2009). Similarly, *COUPTI* and *COUPTII* are expressed within the caudal ganglionic eminence and regulate the identity of cells generated within this region (Kanatani et al., 2008; Lodato et al., 2009).

Biochemically, interneurons are classified on the basis of their expression of different calcium binding proteins such as calretinin, calbindin, and parvalbumin, neuropeptide Y and somatostatin (Hendry et al., 1987; De Felipe,

1993; Fairen et al., 1984; Houser et al., 1983; Gupta et al., 2000; Hendry et al., 1989; Jones, 1975; Kawaguchi and Kubota, 1997; Lund and Lewis, 1993; Ascoli et al., 2008). Physiological classifications are based on intrinsic firing properties of the cells (Butt et al., 2005) while morphological classification relies on the distinctive appearance of the cells.

In rodents, the MGE gives rise to two broad types of interneurons: (a) somatostatin (SST) positive cells with a characteristic burst spiking firing property and a bitufted morphology, exemplified by the Martinotti cell, and (b) parvalbumin (PV) positive cells with fast spiking intrinsic properties, such as basket cells and chandelier cells (reviewed in Corbin and Butt, 2011). The cells generated in the MGE provide the bulk of cortical interneurons and also make significant contributions to interneurons in the hippocampus (Lavdas et al., 1999; Sussel et al., 1999; Witcherle et al., 1999; Pleasure et al., 2000; Anderson et al., 2001). Interneurons generated in the LGE migrate within the rostral migratory stream to reach the olfactory bulb; other LGE-derived interneurons eventually settle in the hippocampus, the striatum and amygdala (Corbin et al., 2001; Marin et al., 2000; Witcherle et al., 2001; Yun et al., 2001; Kohwi et al., 2007; Young et al., 2007; Cocas et al., 2011). The caudal ganglionic eminence (CGE) gives rise to a diverse sub-population of interneurons that express reelin, calretinin (CR), and neuropeptide Y (NPY) (Miyoshi and Fishell, 2011). These cells include the late spiking neurogliaform cells, the bipolar and tripolar cells capable of adaptation in the frequency of their firing, and the fast spiking double bouquet cells (Butt et al., 2005; Corbin and Butt, 2011).

The contribution of the different GE regions to the overall population of cortical interneurons varies across animal species. In rodents, the majority of interneurons are generated in the MGE even though a good number of these cells also originate from the lateral and caudal ganglionic eminences (Marin and Rubenstein, 2001; Lopez-Bendito et al., 2004). In ferrets, however, which demonstrates an early fusion of the LGE and MGE after midcortical development, the majority of interneurons originate from the lateral ganglionic eminence (Figure 1B; Poluch et al., 2008). In humans, many cortical interneurons are produced within the CVZ (Letinic et al., 2002; Yu and Zeevic, 2011).

Similar to pyramidal cells, the laminar position of interneurons appears dependent on the time of birth and their phenotype and reflects their source within the GE (Miller 1985; Fairen et al., 1986; Peduzzi, 1988; Leone et al., 2008). For these reasons, MGE-derived cells, which are born early during development, occupy both deep and upper cortical layers, whereas cells from the CGE, generated much later, occupy the upper cortical layers (Cavanagh and Parnavelas, 1988 and 1989; Miyoshi et al., 2007; Rymar and Sadikot, 2007; Miyoshi and Fishell, 2011). Thus the inside-out pattern of neurogenesis demonstrated by MGE-derived cells is absent in the CGE. Interestingly, interneurons and projection neurons born at a similar time period occupy the same cortical layer (Hevner et al., 2004).

Migration of cortical interneurons. The sequence of events involved in the migration of interneurons is similar to the process of migration of pyramidal cells described earlier. In general, migrating interneurons display either a bipolar

morphology with leading and trailing processes or branched leading processes (Anderson et al., 1997; Bellion et al., 2005; Friocourt et al., 2007; Kappeler et al., 2006; Lavdas et al., 1999; Marin and Rubenstein, 2001; Polleux et al., 2002; Martini et al., 2009). Migration is achieved with a step-wise development and extension of a leading process followed by nucleokinesis and retraction of a trailing process (Marin and Rubenstein, 2001, 2003). Interneurons travelling to the neocortex migrate through multiple routes and complex pathways. Interneurons first migrate within the subpallium towards the corticostriatal border; the path taken by GE-derived cells within this region is influenced by the time of their birth. In general, interneurons fated to lower cortical layers (early born) migrate within the IZ of the GE whereas upper cortical layers (late-born) interneurons migrate through the VZ/SVZ of the striatum before entering the neocortex (Anderson et al., 2001; Ang et al., 2003). As they cross the corticostriatal boundary, migrating GE-derived cells, which are oriented tangentially, use two major routes to reach the neocortex: a subset of interneurons, particularly those originating from the MGE, migrate through the IZ and SVZ; other MGE-derived cells travel through the MZ and the preplate (Anton et al., 1996; Kriegstein and Noctor, 2004; Poluch and Juliano, 2008; Lavdas et al., 1999; Anderson et al., 2001; Wichterle et al., 2001; Jimenez et al., 2002; Ang et al., 2003; Tanaka et al., 2003). LGE-derived cells are also believed to travel tangentially within the SVZ (Anderson et al., 2001).

On reaching the neocortex, migrating interneurons (in the IZ/SVZ and MZ) switch their orientation from tangential to radial mode to reach the cortical plate

and take up a laminar position (Anderson et al., 2001; Nadarajah et al., 2001; Anderson et al., 2002; Nadarajah and Parnavelas, 2002; Wichterle et al., 2001; 2003; Ang et al., 2003; Tanaka et al., 2003, 2006; Poluch and Juliano 2007, 2008). Some interneurons that migrate tangentially within the IZ/SVZ switch to radial orientation and migrate past the cortical plate to reach the MZ before reversing direction to migrate downwards to enter the cortical plate (Tanaka et al., 2003). It has also been observed that populations of interneurons exhibit a 'ventricle-directed' migration. These cells, after reaching the neocortex, reverse direction and migrate to the ventricle before proceeding to migrate back to the cortical plate (Nadarajah et al., 2002). Various reports suggest that the radial phase of interneuron migration use radial glia or axonal support (O'Rourke et al., 1995; Denaxa et al., 2001; Letinic et al., 2002; Polleux et al., 2002; Nadarajah et al., 2002; Ang et al., 2003; Poluch and Juliano, 2007; Yokota et al., 2007).

Functionally, interneurons play important roles in local cortical circuits and in modulating the excitatory output of pyramidal neurons (Cobb et al., 1995; Thomson et al., 1996; Anderson et al., 1997; Wichterle et al., 1999; Parnavelas, 2000; Tamas et al., 2000; Marin and Rubenstein, 2001).

Regulation of interneuron migration. As mentioned earlier, in addition to cortical interneurons, the GE also provides interneurons to other telencephalic structures (such as the hippocampus, striatum and olfactory bulb), which raises the question as to what mechanisms ensure appropriate migration of these cells to their ultimate target destinations. Although the full picture has yet to emerge, available evidence suggests that the combination of intrinsic mechanisms and

environmental cues ensure guided migration of GE-derived cells to their final targets (Figure 1A, Mason et al., 2001; Marin et al., 2003; Wichterle et al., 2003; Corbin and Butt, 2011). The factors responsible for guiding interneurons from the GE to their various targets can be roughly classified into: (a) motogenic factors (b) repulsive signals (c) attractants (d) neurotransmitters (e) genetic factors and (f) stop signals (Marin and Rubenstein, 2001; 2003; Ayala et al., 2007; Corbin and Butt, 2011). Motogenic factors trigger kinetic activity in GE-derived cells leading to random movement in all directions. For instance, hepatocyte growth factor (HGF), a motogen, acts on μ -PAR receptors to selectively promote the migration of parvalbumin-positive interneurons to the neocortex (Powell et al., 2001). Thus, in loss-of-function experiments such as in μ -PAR^{-/-} mutants, interneuron migration is impaired leading to significant reduction in GE-derived cells in the neocortex. Neurotrophins such as NT4 and BDNF activate TrkB neurotrophin receptors to stimulate migration of GE cells, which underscores the migration deficit seen when neurotrophin signaling is impaired (Brunstrom et al., 1997; Polleux et al., 2002).

Repulsive signals regulate the movement of GE-derived cells to cross the cortico-striatal boundary enroute to the neocortex. The ability of cortical interneurons to sort out from GE-derived cells destined to the striatum is in part due to the repellant action of semaphorins mediated by semaphorin-neuropillin signaling. Cortical interneurons express two transmembrane receptors, neuropillin 1 and neuropilin 2 (*Nrp1/2*), enabling them to respond to repulsive signals of semaphorin 3A and semaphorin 3AF expressed in the striatum to thus

avoid settling in the striatum (Marin et al., 2001b; Tamamaki et al., 2003). Other studies suggest a role for Slit proteins (*Slit1* and *Slit2*), which are expressed in the subpallium during development, and *Netrin* as repulsive signals to migrating interneurons, but this claim is undermined by observations in *Slit1* and *Slit2* double mutants or in *Slit1* and *Slit2* and *Netrin1* deficient animals, in which the number of cortical interneurons were not significantly altered (Zhu et al., 1999; Hamasaki et al., 2001; Marin et al., 2003; Wichterle et al., 2003). Recently, it has been suggested that repulsive signals expressed in the cortical hem regulate the switch in orientation of migrating GE cells from tangential to radial as cells migrate to the cortical plate (Caronina-Brown and Grove, 2011).

In various *in-vitro* migration assays, migrating interneurons are attracted to cortical cells as well as ectopically placed cortical explants, suggesting that neocortical diffusible attractive factors modulate the directional movement of interneurons (Marin et al., 2003; Poluch et al., 2008). This suggestion is supported by observations that genetic disruption of embryonic cortex in *Emx1* and *Emx2* double mutants impairs the migration of interneurons, indicating that factors intrinsic to the cortex influence interneuron migration (Shinozaki et al., 2002). Furthermore, when normal and dysplastic GE explants are placed together in coculture experiments, improved orientation of interneurons migrating toward the normal cortex indicate that the neocortical environment plays an important role in the proper migration of interneurons (Poluch et al., 2008). Taken together, these observations provide strong evidence that the neocortical environment is a critical determinant of proper migration of GE-derived neuronal

cells destined for the cortex. However, the identities of candidate neocortical cues that modulate interneuron migration are yet to be conclusively established. Nonetheless, there have been suggestions that GDNF and Neuregulin (Flames et al., 2004; Pozas and Ibanez, 2005; Ghashghaei et al., 2006), and chemokines via CXCR7 and CXCR4 receptors (Stumm et al., 2003; Lopez-Bendito et al., 2008; Sanchiz-Alcaniz et al., 2011; Wang et al., 2011) are chemoattractant to migrating GE cells.

Neurotransmitters also regulate migration of GE-derived neurons. Migrating interneurons express AMPA, NMDA, GABA_A and GABA_B and dopamine receptors and may therefore respond to activity mediated by these molecules (Lopez-Bendito et al., 2003; Represa and Ben-Ari, 2005; Manent et al., 2006; Crandall et al., 2007). Indeed, the role of these agents as modulators of interneuron migration has been demonstrated in a number of migratory models in which the actions of these agents operated via a Ca²⁺-dependent mechanism (Behar et al., 1996, 1998, 1999, 2001; Komuro and Rakic, 1992, 1993; Metin et al., 2000; Poluch and Konig, 2002; Poluch et al., 2008; Soria and Valdemolillos, 2002).

Proneural genes also play important roles in the specification of interneurons and regulate migration of these cells. For instance, the homeodomain transcription factor, *Nkx2.1*, expressed in the MGE, negatively regulates the expression of *Nrp* receptors in migrating GE-derived cells that control the sorting of cortical interneurons (Nobrega-Pereira et al., 2008). Thus, in the presence of *Nkx2.1*, *Nrp* is downregulated, resulting in the failure of MGE

cells to respond to semaphorin signaling leading to the reduction of interneurons entering the neocortex (Nobrega-Pereira et al., 2008, Nobrega-Pereira and Marin, 2009). A similar role has been suggested for the homeobox transcription factor, *Dlx1/2* in regulating interneuron migration, presumably by controlling the expression of *Nrp* receptors and *Arx* (Le et al., 2007; Cobos et al., 2005a, 2005b, 2007). The aristaless related homeobox gene, *Arx*, has been implicated in modulating migration of interneurons both in rodents and humans (Kitamura et al., 2002; Alifragis et al., 2004; Liodis et al., 2007; Friocourt et al., 2008).

Retinoic acid (RA) also influence the migration of cells leaving the GE (Crandall et al., 2011). RA inhibits GE cell migration probably through regulation of dopamine D2 receptor expression. Activation of dopamine D2 receptors inhibits GE cell migration (Crandall et al., 2007). Consequently, blocking D2-receptors abolishes the effect of RA leading to enhancement of GE cell migration. Recently, it was demonstrated that a subset of interneurons that are generated in the CGE depend on activity for migration and development of axonal arbors for network integration (Natalia et al., 2011).

On reaching their target destination in the cortex, migrating GE cells terminate migration and take up a cortical position. How this is achieved still remains unclear. Local signals from projection neurons within the same cortical layer and/or from the meninges may instruct migrating GE cells to take up cortical position (Tiveron et al., 2006; Li et al., 2008; Li and Pleasure, 2011; Lodato et al., 2011; Caronia-Brown and Grove, 2011; Zarbalis et al., 2012). The neuron-specific potassium- chloride co-transporter, KCC2, has also been found

to act as a stop signal (Bortone and Polleux, 2010) (for detailed discussion on this topic, see below).

Role of GABA in cortical development and function. Within the cerebral cortex of adult mammals, inhibitory neurotransmission is mediated primarily by GABA, through two receptor subtypes: the fast conducting ionotropic GABA_A receptors and the slow conducting metabotropic GABA_B receptor (Tija et al., 2008). Thus GABA signaling modulates excitatory outputs of glutamatergic projection neurons to establish a balance between excitation and inhibition to ensure normal cortical function (Gilbert and Wiesel 1985; Jones 1986; Cobb et al., 1995; Thomson et al., 1996; Tamas et al., 2000; Freund, 2003; Hensch, 2005; Somogyi and Klausberger 2005; Caronia-Brown and Grove, 2011). Consequently, alterations in GABAergic signaling have been implicated in many neuropsychiatric disorders such as epilepsy (Luhmann et al., 1995; Loup et al., 2000; Kobayashi and Buckmaster, 2003; Fristch et al., 2009; Gant et al., 2009; Kang and Barnes; 2012; Ma and Prince et al., 2012), Alzheimers Disease (Young, 1987; Fonseca et al., 1993; Lanctot et al., 2004; Limon et al., 2012), schizophrenia (Keverne, 1999; Hashimoto et al., 2003; Lewis et al., 2004, 2005; Bernstein et al., 2007; Chattopaddhyaya and Di Cristo, 2012; Moyer et al., 2012) autism (Fatemi et al., 2006 , 2009; Chao et al., 2010; Oblak et al., 2010; Chattopaddhyaya and Di Cristo, 2012; Kang and Barnes; 2012), and X-linked lissencephaly with abnormal genitalia (Colombo et al., 2007).

Interestingly, the role of GABA as an inhibitory neurotransmitter is not innate but evolves. During development of the neocortex, migrating neurons first

express GABA receptors and receive signals from striatally released GABA, followed by subsequent activation of excitatory glutamate receptors (Owens et al., 1999; Tyzio et al., 1999; Hennou et al., 2002). There is good evidence that GABA plays a multifaceted role during development, in regulating DNA synthesis, cell proliferation, migration, differentiation, development of AMPA receptors, excitatory synapse formation and possibly acting as a stop signal for migrating GE cells (Barbin et al., 1993; Heck et al., 2007; LoTurco et al., 1995; Wang and Kriegstein, 2008, 2009; Bortone and Polleux, 2010; Inada et al., 2011). It has been known for several years that GABA acts as an excitatory neurotransmitter early in development to facilitate cell migration, but switches to mediate inhibition as cells mature (Plotkin et al., 1997; Clayton et al., 1998; Rivera et al., 1999). The dual role of GABA is attributed to the temporal expression of two cation-chloride transporters, the sodium-potassium-chloride co-transporter (NKCC1) and the potassium-chloride co-transporter (KCC2). Migrating neurons initially express NKCC1, which imports chloride ions (Cl^-) into the cell to maintain a higher intracellular Cl^- concentration (Clayton et al., 1998). In response to GABA_A activation, the outward movement of Cl^- results in depolarization of the cells and triggers Ca^{2+} -mediated events that influence the cytoskeleton and stimulate migration (Behar et al., 2001; Soria & Valdeolmillos, 2002). As cells mature, NKCC1 downregulates and coincides with increased expression of KCC2 (Plotkin et al., 1997; Delpire, 2000; Li et al., 2002; Owens and Kriegstein, 2002). KCC2 is a Cl^- exporter that maintains a higher extracellular gradient of Cl^- . Consequently, GABA_{AR} activation when KCC2 is

increased results in hyperpolarization and cell inhibition (Payne, 1997; Rivera et al., 1999). In addition, GABA activity is believed to be an important factor in promoting the expression of KCC2 during development (Ganguly et al., 1999).

The role of GABA in neuronal migration has been evaluated in various *in vitro* and *in vivo* migratory models, with researchers finding both enhancement and inhibition of neuronal migration (Behar et al., 1996, 1998, 1999, 2001, Heck et al., 2007). These discrepancies may arise from differences in the composition of cell type and the design of the various studies. In addition, many experiments did not distinguish between effects elicited on cells arising from the GE vs CVZ. The dose of GABAergic agonists and antagonists appears to have differential effects, as low concentrations of GABA cause directed migration, whereas high concentrations appear to cause random movement (Behar et al., 1996; 1998). In a study by Heck et al. (2007) both GABA_{AR} agonists and antagonists delivered by elvax result in ectopic migration in the neocortex in neonatal animals. Bolteus and Bordey (2004) demonstrated that enhancement of GABAergic activity in acute brain slices slows the rate of cellular migration; although this study evaluated chain migration in the olfactory system, which may use other mechanisms than interneurons migrating into the neocortex. In addition to differences in cell types studied and the type of assay used, the seemingly complex and contradictory role of GABA in modulating migration of interneurons could also be due in part to the dynamic and temporal expression of the different subsets of GABA receptors in migrating GE cells, which determines the nature of response of these cells to GABA (Laurie et al., 1992; Poulter et al., 1992;

Fritschy et al., 1994; Cuzon et al., 2006; Wang and Kriegstein, 2009; Carlson and Yeh, 2010). Taken together, these studies indicate that GABA has an effect on neuronal migration, but the exact process needs continued clarification.

In addition to affecting migration, GABA, as a result of KCC2 expression, can also act as a stop signal for migrating GE cells (Bortone and Polleux, 2010). The effect of GABA on neuronal migration is also likely to be dependent on Ca^{2+} . In the neocortex, stimulation of GABA_A receptors activates changes in intracellular calcium levels in tangentially migrating cells and this effect may influence growth cone dynamics and thus migratory behavior (Behar et al., 2001; Soria & Valdeolmillos, 2002).

The excitatory action of GABA during early neocortical development (Ben-Ari et al., 1989; Cherubini et al., 1991) is also implicated in the establishment of AMPA receptors and formation of synapses- an effect mediated via NMDA signaling (Wang and Kriegstein, 2008). Recently, our lab demonstrated that GABA_{AR} mechanisms are important for both the migration and orientation of tangentially migrating interneurons (Poluch et al., 2008). Taken together, these observations underline an important role of GABA signaling during migration and cortical development.

Cortical dysplasia. Perturbation in the process of cell generation, migration or terminal differentiation as a result of genetic or environmental factors can profoundly disrupt corticogenesis leading to cortical dysplasia. These aberrations, which are often associated with abnormal GABA signaling (Rosen et al., 1998; Roper et al., 1999; Talos et al., 2012; Cepeda et al., 2012), underlie a

vast number of neurological/neuropsychiatric disorders including drug-resistant epilepsies requiring surgery, depression and schizophrenia (Chugani et al., 1990; Palmini et al., 1991; Gleeson and Walsh, 2000; Zhu and Roper, 2000; Ross and Walsh, 2001; Calcagnotto et al., 2002; Calcagnotto and Baraban, 2003; Colombo et al., 2003; Lerner et al., 2009). The etiological factors underlying the development of cortical dysplasia may either be genetic or environmental in nature. Among the identified genetic factors are mutations in key genes that play a pivotal role in neuronal migration such as *Lis1* (which results in lissencephaly), and *Doublecortin* (that leads to double cortex in females and heterotopias in males) (des Portes et al., 1998; Feng and Walsh, 2001; Gleeson et al., 1998; 1999). Prenatal alcohol exposure, carbon monoxide intoxication from cigarette smoke, alcohol intoxication are known environmental factors mediating cortical dysplasia (Ginsburg and Myers, 1974a and b; Choi and Matthias, 1987; Barinaga, 1993; Komatsu et al., 2001; Gorman et al., 2001; Autti-Ramo et al., 2002).

MAM Model of Cortical dysplasia: In order to gain critical insight into the underlying pathology in cortical dysplasia, many animal models of the syndrome have been developed to mirror specific deficits of the disorder observed in humans. Genetic models are generated through manipulation of key genes involved in cortical development such as *Lis1*, *Doublecortin*, *Reelin*, *Otx1*, *TSC1* *cKO*; this generates phenotypes such as smooth brain, double cortex, reeler mouse, flathead, Tish, and Eker rats (Caviness & Yorke, 1976; Acampora et al., 1996; Lee et al., 1997; Yeung et al., 1997; Hirotsune et al., 1995, 1998; Roberts

et al., 2000; Uhlmann et al., 2002; Moroni et al., 2008). Other models leading to CD not involving genetic manipulation, such as freeze lesions and cortical undercut, rely on physical assault to the developing brain to induce injury and subsequent aberrant migration (Rosen et al., 1992 and 1998; Jacobs et al., 1996). In addition, some models use toxins such as carmustine, MAM, and radiation to disrupt cortical development in-utero in rodents (Ferrer, 1993; Ferrer et al., 1993; Roper, 1998; Colacitti et al., 1999; Benardete and Kriegstein, 2002; Moroni et al., 2008).

MAM is an antimitotic that transiently interferes with the cell cycle to inhibit neurogenesis. MAM methylates the 7' position of the guanine residue, which results in inhibition of DNA polymerase activity, and thus reduces nucleic acid synthesis (Matsumoto and Higa, 1966). Our laboratory developed a model of cortical dysplasia, in which we administer MAM intraperitoneally to pregnant ferrets at specific gestational periods. Ferrets are the smallest mammal with a gyrencephalic cortex and thus make excellent models for studying development of the neocortex. The presence of sulci and gyri is evolutionarily significant since these features, which expand the cortical surface area, enhance the capacity for higher cognitive function (Kriegstein et al., 2006). These features are present in humans and underlie the need to develop and study this disorder in more advanced animal models.

MAM transiently disrupts cell production resulting in different types of cortical dysplasia depending on the time of administration (Noctor et al., 1999; Gierdalski and Juliano., 2003; Hasling et al., 2003; Gierdalski and Juliano, 2005;

Poluch and Juliano, 2007, 2010). Administration of MAM during the birth of layer 4 (E33) results in diminished or near total loss of this layer and widespread terminations of thalamic afferents in the neocortex (Figure 3A; Noctor et al., 1999, 2001b; Palmer et al., 2001). Of significance, ferrets treated with MAM on E33 demonstrate aberrant distribution of GABAergic interneurons (Poluch et al., 2008), increased expression of GABA_{AR} in the cortex (Figure 3B; Jablonska et al., 2005), alteration in the balance of excitation and inhibition, and overall changes in the pattern of cortical information transfer (Jablonska et al., 2004; McLaughlin and Juliano, 2005; Noctor et al., 2001b). Although the morphology and overall number of GABAergic interneurons in the neocortex is not altered, the laminar position of certain categories of interneurons is changed in the E33 MAM model, suggesting that these neurons do not migrate into their proper target sites (Poluch et al., 2008). An important question that arises from this finding of aberrant interneuron redistribution is whether they occur as a result of changes in the kinetic behavior of migrating cells due to intrinsic mechanisms, or as a result of changes in the neocortical environment, or both. Ferrets treated with MAM display an altered distribution of subsets of interneurons and functional deficits similar to those seen in other models of cortical dysplasia (Rosen et al., 1998; Palmer et al., 2001; Cacagnotto et al., 2002; McLaughlin et al., 2005; Poluch et al., 2008). Although we do not observe clinical symptoms of epilepsy in our model as has been reported following similar treatments in rats (Baraban and Schwartzkroin, 1996; Chevassus-au-Louis et al., 1998; de Feo et al., 1995; Germano et al., 1996; Kimberle et al., 1999), the aberrant redistribution and

position of interneurons seen in our model raises the likelihood that the inhibitory function of these cells might be compromised, which may impair overall cortical function. This study investigates the impact of MAM treatment on the dynamic pattern of migration of GE cells as it relates to the basal level of GABA_A activity and function of the resulting cortex.

A

B




Figure 1: (A). During development, projection neurons and interneurons are generated from distinct neurogenic regions. Projection neurons form in the cortical ventricular zone and migrate radially to the cortical plate. Cortical interneurons are generated in the ganglionic eminences (GE) but migrate tangentially and radially to settle in the cortex. (B) Several extrinsic cues, including GABA, act in concert with intrinsic signaling mechanisms to regulate the process of migration of interneurons. The anatomic layout of the GE varies across species. Unlike rodents with three distinct subtypes of GE, the GE in ferrets fuses at midcortical development. Broken lines signify the corticostriatal boundary.




Figure 2: Pyramidal cells are generated from cortical ventricular zone and migrate radially to different cortical layers. In this picture, cells within the CVZ were electroporated with a plasmid producing red fluorescent protein. (A) Labeled radial glia cells that span the cortex can be seen aposed with migrating projection neurons (B) High power view of a section of migrating projection neurons on radial glial cells.




Figure 3: Administration of MAM on E33 induces several developmental malformations on the cortex. (A) As a result of transient inhibition of neurogenesis, layer 4 cells, which are born during this time period, are not generated leading to reduction in the thickness of cortical layer 4. (B) After the reduction in thickness of layer 4, GABA_A receptors, which are normally expressed in high density in this region, expand to upper cortical layers.

CHAPTER 2

Altered kinetic behavior underlies redistribution of interneurons in a model of cortical dysplasia: the influence of elevated GABA_A receptor activity.

J. Abbah¹ and S. L. Juliano^{1,2}

¹Program in Neuroscience and ²Anatomy, Physiology and Genetics, Uniformed Services University of the Health Sciences, Bethesda, Maryland, 20814-4799

Corresponding Author: Sharon L Juliano: Anatomy, Physiology and Genetics, Uniformed Services University of the Health Sciences, 4301 Jones Bridge Rd., Bethesda, Maryland, 20814-4799

Phone: 301-295-3673

Email: sjuliano@usuhs.edu

Running Title: GABA activity is critical for migration of GE cells

Abstract

Appropriate function of the neocortex depends on timely generation and migration of cells produced in the germinal zones of the neocortex and ganglionic eminence (GE). Failure to accurately complete migration results in cortical dysplasia, a developmental syndrome implicated in many neurologic disorders. We developed a model of cortical dysplasia in ferrets involving administration of methylazoxymethanol acetate (MAM), an antimitotic, to pregnant ferrets on gestational day 33, leading to dramatic reduction of layer 4 in the neocortex. Here, using time-lapse video imaging, we investigate kinetic behavior of migrating GE cells in ferrets and the role of GABA_A activity. Treatment with MAM significantly reduced migration speed and the relative proportion of GE cells demonstrating exploratory behavior. Pharmacologic inhibition of GABA_A receptors (GABA_{AR}) improved the speed of migration and exploratory ability of migrating MAM-treated GE cells. Additionally, the expression of $\alpha 2$ and $\alpha 3$ subunits of GABA_{AR} and the potassium chloride co-transporter (KCC2) increased in the neocortex of MAM-treated animals. After MAM treatment, increases in endogenous KCC2 and GABA_{AR} combine to alter the kinetic properties and exploratory behavior of migrating interneurons in ferrets. We show a direct correlation between increased GABA_A and KCC2 expression with impaired migration and ability to explore the environment.

Keywords: development, ferret, KCC2, MAM, neuronal migration

Introduction

The laminar organization of the neocortex and its ability to function depends on timely generation and proper migration of cells originating from the ventricular zones of the neocortex (CVZ) and GE. Events that adversely impact the migratory process can lead to cortical dysplasia, a developmental abnormality characterized by aberrant cell clustering, altered gyral patterns, and changes in electrophysiological profile (Taylor et al. 1971; Choi and Mathias 1987; Tassi et al. 2002; Calcagnotto and Baraban 2003; Moroni et al. 2008). The aberrations underlie a vast number of neurological/neuropsychiatric disorders including drug-resistant epilepsy, depression, and schizophrenia (Palmini et al. 1991; Gleeson and Walsh 2000; Zhu and Roper 2000; Ross and Walsh 2001; Calcagnotto et al. 2002). Understanding the process of neuronal migration into the neocortex is critical to comprehending the many disorders that result from disrupted migration.

We developed a model of cortical dysplasia by administering a short acting anti-mitotic, methylazoxymethanol (MAM), to pregnant ferrets on gestational day (E33). Ferrets are the smallest animal with a convoluted cortex, which make them an important model for neocortical development. The tangential expansion of the neocortex, which assists in the formation of the sulci and gyri, relies on proliferation of outer subventricular progenitors, a cell population found in ferrets and humans, but absent in rodents (Fietz et al. 2010; Lui et al. 2011; also see Martinez et al 2012 for a different point of view). Our lab also found distinctions in the migratory patterns of interneurons in rodents and

ferrets, indicating the importance of studying a more developed model of neocortex (Poluch et al. 2008).

Treatment with MAM on E33 coincides with the generation of layer 4 and results in its dramatic reduction and in widespread re-distribution of GABA_{AαR} (Noctor et al. 1999; Palmer et al. 2001; Jablonska et al. 2004). Although the morphology and overall number of interneurons in the neocortex is not altered, the laminar position and orientation of interneurons is changed in the MAM model, suggesting that these neurons do not migrate into their proper target sites (Poluch et al. 2008). The aberrant redistribution of interneurons and GABA_{AαR} within the neocortex of MAM-treated animals raises important questions. 1) Do changes in the distribution of cells arising from the GE occur as a result of altered kinetic behavior of migrating GE cells, and, if so 2) are these changes influenced by the ambient activity of GABA?

Using real-time video imaging and an *in-vitro* migration assay of organotypic cultures of neonatal ferrets, we observed the dynamic movement patterns of neurons leaving the GE and CVZ. Treatment with MAM impairs the speed and exploratory potential of tangentially migrating interneurons leaving the GE without any significant effects on neurons arising from the CVZ. GE cells in MAM-treated animals exposed to GABA antagonists showed a significant improvement in the speed of migration and the proportion of cells that display exploratory activity. This reinforces the idea that E33 MAM treatment alters the migration kinetics of GE cells. The changes in the dynamic movement of GE cells are related to abnormal levels of GABA_A-mediated activity, which diminishes the

capacity of these cells to explore the environment and effectively migrate to their target within the neocortex.

Materials and Methods

Timed pregnant ferrets were obtained from Marshal Farms (New Rose, NY) and maintained in the animal facilities of the Uniformed Services University of the Health Sciences (USUHS). Pregnant ferrets were injected with 14mg/kg of MAM (Midwest Research Institute, Kansas City, MO) IP on E33 under isoflurane anesthesia (1-2%). After recovery from anesthesia, ferrets were maintained in the animal facility until their kits were delivered. Handling of animals complied with the USUHS Institutional Animal Care and Use Committee policy on the humane use and treatment of animals.

Preparation of Organotypic slices: Preparation of organotypic cultures was accomplished as previously described (Palmer et al. 2001). Postnatal day 0 to day 1 (P0-P1) ferret kits were anesthetized with sodium pentobarbital (50mg/kg), their brains removed and placed in ice-cold artificial cerebrospinal fluid (aCSF) composed of (in mM): 124 NaCl, 3.2 KCl, 2.4 CaCl₂, 1.2 MgCl₂, 26 NaHCO₃, 1.2 NaH₂PO₄ and 10 glucose bubbled with 95% O₂ and 5% CO₂ under a laminar flow hood. Coronal slices (350-500 µm thick) were prepared from each hemisphere using a tissue chopper (Stoelting Co., Wood Dale, Illinois). Brain slices were transferred into 0.4 µm culture plate inserts (Millicell-CM, Bedford, MA) placed in six-well plates containing Neurobasal media with B27, N2, and G1.2 (containing gentamycin and glutamine) supplements. Slices were incubated at 37°C under 95% O₂ and 5% CO₂. In some experiments, bicuculline methiodide (BMI; Tocris Bioscience, Park Ellisville, MO) at a final concentration of 10 µM was added to the media.

Cell labeling: Two approaches were used to label migrating neurons. In one approach, crystals of 1,l'-dioctadecyl-3,3,3,3'-tetramethylindocarbocyanine percholate (Dil, Invitrogen, Carlsbad, CA) were placed in the GE or CVZ using pulled borosilicate glass pipettes. In another approach, we used electroporation to focally transfect cells within the ventricular zone (VZ) of the GE using a modification of the method described by Flames et al, 2004. Transfection was accomplished using a plasmid that codes for red fluorescent protein (RFP), which was cloned into pCAGGS expression vector (a gift from Dr Tarik Haydar). Between 1-3 μ l of plasmid DNA (2.7-3.6 μ g/ μ l) was injected into the VZ of the GE of organotypic slices of ferrets. The cathode of a gene paddle electrode (Harvard Apparatus, Inc., Holliston, MA) was placed within the lateral ventricle close to the VZ of the GE while the anode was placed closed to the pial surface in an appropriate position. A pulse of 60V was applied four times, each lasting for 50 ms at intervals of 950 ms using a BTX ECM830 pulse generator (Gal et al 2006; Harvard Apparatus, Inc., Holliston, MA). Slices were incubated as above for at least 24 hrs prior to video imaging.

Video imaging: After incubating organotypic slices for 24 h (CVZ-derived) or 48 h (GE-derived) for analysis, migrating neurons were continuously visualized using an Axiovert 200 inverted microscope fitted with an apotome and Axiovision software (Carl Zeiss AG, Oberkochen, Germany). The microscope was fitted with an incubation chamber and a holder for the slices, which were maintained with humidification at 37°C with 95% O₂ and 5% CO₂. Serial stacks of images taken through the thickness of the slice were collected using a 10X objective every 30

min for 24 hrs. The image stacks were collapsed into a single frame prior to analysis of migration. Migrating GE cells were analyzed after crossing the corticostriatal boundary into the neocortex.

Analysis of migration: To measure speed, migrating cells captured in real time were tracked using ImageJ software (<http://rsb.info.nih.gov/ij>) and the tracked distances measured and expressed over time (to obtain the speed in real time) using a program written in R statistical language (R Development, 2009). Also, the orientation of migrating GE cells (either tangential or radial) and exploratory activities were quantified for each migrating cell. The distribution of migrating GE cells within the neocortex in both normal and MAM-treated animals was determined using Axiovision software. To do this, the neocortex was divided into three roughly concentric regions (0-350 μm , 350-700 μm and >700 μm) from the border of the GE VZ and labeled neurons found within the various bins were counted and expressed as a percentage of the total number of cells.

Western Blot analysis: The neocortex was dissected from organotypic slices of normal and MAM-treated P0-P1 ferret brains (prepared as described above), frozen on dry ice and preserved at -82°C prior to use. Tissues were homogenized using RIPA lysis buffer (Santa Cruz Biotech, Santa Cruz, CA) followed by centrifugation at 14,000 g at 4°C . Protein concentration was estimated using a colorimetric assay. Proteins were separated by SDS-PAGE using 10% Bis-Tris gel and electrophoretically transferred to a PVDF membrane (Invitrogen, Carlsbad, CA). A loading volume of 10 μl containing 1-2 μg of protein was used for each analysis. Membranes were incubated with Casein blocking

buffer (PBS (0.5M NaCl) + 3% Casein + 0.5% Tween-20) overnight, followed by affinity purified rabbit polyclonal antibodies directed against GABA_{Aα2} (1:200; ProSci Inc., Poway, CA), GABA_{Aα3} (1:2000; Sigma, St. Louis, MO), KCC2 (1:500, Millipore, Billerica, MA), and monoclonal anti-actin (1:3000, NeoMarkers, Fremont, CA) for 24 hrs. Following several washes with PBS, protein bands were detected using HRP-conjugated anti-rabbit secondary antibodies (1:1000, Jackson Lab., West Grove, PA) and HRP-conjugated anti mouse (1:3000, Thermo Scientific, Rockford, IL) and visualized using enhanced chemiluminescence detection. Signal intensities were quantified using Image j software (<http://rsb.info.nih.gov/ij>).

Statistics: The chi-square distribution test was used to analyze distribution of GE cells within the neocortex. For all other analysis, a Student's t-test was applied and differences evaluated at $P < 0.05$.

Results

General properties of normal and MAM- treated migrating cells leaving the GE

We previously reported that cells leaving the GE in MAM-treated animals exhibit abnormal characteristics, specifically showing orientations that differed from control cells (Poluch et al., 2008). To further characterize the migratory behavior of cells leaving the GE or the CVZ, we applied an *in-vitro* live video-imaging assay to continuously view migrating cells. Our earlier work demonstrates that at P0-P1 in the ferret, the medial and lateral GE are fused, thus we will refer to our label as directed to the fused GE (Poluch et al. 2008). Migrating cells exiting the GE or CVZ of organotypic slices were labeled using electroporation with plasmids that code for RFP or injections of Dil. An example of a slice labeled by electroporation can be seen in Figure 1A. Higher power views of cells en route to the cortical plate are shown in Figures 1B and C. These cells have varied orientations as well as morphology; the dynamic features change when observed in real time. After allowing the cells to move away from the GE or CVZ, the dynamic pattern of migration was captured in real time. GE cells moving into the neocortex adopt different orientations and take different routes. A number of migrating neurons course through the VZ or SVZ (subventricular zone), many orient in a tangential direction (Figure 2A, B red arrows). A different group of more superficially labeled GE cells display both tangential (yellow arrows) and radial (green arrows) orientations. Many of the radially oriented cells point in the direction of the pial surface. However, some of

the radially-oriented cells have their leading edges directed toward the ventricular zone (Figure 2A, B, blue arrows). These observations correspond to our previous report in the ferret that similar to rodents, tangentially migrating cells in the GE follow 3 paths including one close to the VZ, one in the IZ, and one superficially near the marginal zone (not shown in this case) (Poluch et al. 2008). We also observed that migrating GE cells alter their morphology as they transit from point to point, generating distinct morphological features. In static images the labeled cells are either unipolar (the cell possesses a single leading unbranched process (Figure 2C-a), display a branched process (Figure 2C-b), are bipolar (both leading and trailing processes Figure 2C-c), or multipolar (multiple processes emanate from the soma of the cell (Figure 2C-d).

Although the general trend of movement for cells leaving the GE is towards the neocortex, individual cells move in all directions and often alter their course of direction (Movie 1). Cells may emit one or more processes and move in the direction of the new branch (e.g. Movie 2), or emit branches and continue in the same direction or not move at all, but only explore the environment for a time (Movie 3). We also frequently observed nucleokinesis in which the nucleus moves within the cytoplasm of a cell in the direction of the leading process (as described by others such as Nadarajah et al. 2001, Martini et al. 2009, Movies 2 and 3). This process usually precedes the point-to-point translocation of the cell during migration.

Kinetic behavior of cells exiting the GE in normal and MAM-treated slices

Our earlier work indicates that after treatment with MAM, in addition to changes in orientation while migrating into the cortical plate, several subtypes of interneurons are redistributed in the neocortex of juvenile ferrets (Poluch et al. 2008). To assess the factors contributing to the altered distribution, we evaluated specific dynamic parameters of cells leaving the ganglionic eminence in normal and MAM-treated animals, including the speed, orientation of movement, and exploratory behavior (defined as extension of new processes and changes in the direction of movement).

Speed and direction. The migratory speed of GE and CVZ cells was measured as described in the Methods. Cells in organotypic cultures of normal ferrets migrated significantly faster ($37.435 \mu\text{m} \pm 0.87/\text{hr}$) compared with similar cells in animals treated with MAM ($32.98499 \mu\text{m} \pm 0.73/\text{hr}$) (Figure 3A). Because migrating interneurons from the GE have different orientations (e.g. Figure 2A and also Ang et al. 2003; Tanaka et al. 2003 and 2006), we examined the relationship of speed to orientation during movement and assessed distinctions between normal and MAM treatment for radial versus tangential travel. Normal cells traveling in both radial and tangential directions migrated significantly faster than MAM-treated cells (Figure 3B).

To investigate if changes in the speed of migration affect the final distribution of migrating neurons, we assessed their positions in organotypic slices of normal and MAM-treated animals after 48 hrs of incubation. We divided the neocortex into three roughly concentric regions (0-350 μm , 350-700 μm and greater than 700 μm) from the point of Dil injection in the GE using Axiovision

software (Figure 4A). The total number of migrating GE cells in each sector was computed; the amount of cells in each sector of the normal and MAM-treated organotypic slices were compared and presented as percentages of the total. Although slightly more cells in normal organotypic slices migrated further away from the GE as revealed by the higher percent in the region 700 μ m from the GE, the overall distribution across the three sectors was not significantly different (Figure 4B). In other words, the cells leaving the MAM-treated GE moved slower, but eventually situate at similar distances away from the GE.

Exploratory behavior of GE-derived neurons in normal and MAM-treated slices

Migrating GE cells are influenced by a variety of signaling molecules that induce stereotypic migratory activity, which we refer to as exploratory behavior. Two parameters defined exploratory behavior in our study: (i) process extension or the ability of migrating cells to extend more than one leading process and (ii) turns or changes in direction of movement, in which a deviation from the initial course of movement occurred. In Figure 5A, the green arrow indicates a cell changing direction, while the red arrow shows a cell branching and also changing its direction of movement. To determine if treatment with MAM alters the ability of migrating GE cells to explore the neocortical environment, migrating cells were observed during live video imaging for 24 hrs. For several examples, the path of multiple migrating cells was traced using ImageJ. The movement was tracked for a period of at least 8 hours and the position of each cell recorded. Figure 6A and

B illustrate that the general pattern of movement in the MAM-treated tracks of cells (Figure 6B) show fewer changes in their direction when compared with the normal patterns of movement (Figure 6A). The quantification of movement patterns is seen in Figure 5B, which demonstrates that the tracks of normal cells completed more changes of direction in their movement patterns (i.e., turns) compared with the MAM-treated cells. The percent of cells displaying process extension or turning in each slice was calculated for normal and MAM-treated organotypic cultures. Cells were scored negative if they did not display any of these behaviors during the observed period of time. In MAM-treated slices, fewer cells leaving the GE extended more than one leading process in comparison with cells in normal animals. In addition, the percent of cells that performed turns was fewer after MAM treatment (Figure 5B).

Interference with GABA_A-mediated activity affects migratory behavior of GE cells in MAM-treated animals

The migratory behavior of interneurons entering the neocortex is regulated by a variety of signaling molecules including GABA (Cuzon et al. 2006; Heck et al. 2007; Lopez-Bendito et al. 2003 and 2008). We previously demonstrated that treatment with MAM leads to a widespread redistribution of GABA_{Aα} receptors; we also found that increasing GABA_{AR}-mediated activity in normal organotypic cultures interferes with the orientation of interneurons as they migrate into the cortical plate similar to the effect of MAM treatment (Jablonska et al. 2004; Poluch et al. 2008). These findings suggest that sensitivity to GABA is unusually

high after MAM treatment and inhibiting GABA signaling may alter the positioning of neocortical interneurons. We predicted that treatment with GABA_A antagonists would shift the kinetic behavior of neurons originating from MAM-treated slices toward normal. The speed and exploratory behavior of migrating cells in normal and MAM-treated slices was evaluated as described previously after adding the GABA_A antagonist BMI to the media. This treatment significantly increased the speed of migration in the MAM-treated group to be closer to normal (Figure 7A).

When organotypic slices of E33 MAM-treated animals were exposed to BMI, the percent of migrating cells extending more than one leading process increased and was nearly identical to those of normal migrating GE cells. This matches our prediction that decreasing GABA_A activity in MAM-treated brains results in more normal behavior (Figure 7B). The number of turns, or alterations in the direction of movement, did not change however, after adding BMI to the media.

Treatment with MAM does not significantly alter the migration kinetics of neurons originating from the cortical ventricular zone

Our previous observations indicated that tangentially migrating GABAergic neurons were specifically disoriented in MAM-treated ferrets (Poluch et al. 2008). To test if the changes in the kinetic behavior of migrating interneurons were limited to the neurons originating from the GE, we examined the migration of neurons from the CVZ to the neocortex using live-video imaging. Organotypic slices were injected with Dil in the neocortical ventricular zone and only slices in

which the Dil label was confined to the VZ were used for analysis. In this regard we limited labeling cells in the vicinity of the CVZ, but arising from the GE. We hoped to insure that the cells migrating away originated from the CVZ. An example of an injection site and cells migrating away from the CVZ can be seen in Figure 8. We measured the speed and exploratory behavior of cells leaving the CVZ in normal and MAM-treated slices. Although we found no significant differences in the parameters analyzed between the two groups, there was a noticeable, but statistically insignificant, decrease in the speed of migration for the MAM-treated cells (Figure 9A-B).

Treatment with MAM increases the expression of GABA_{Aα2}, GABA_{Aα3} receptor subunits and KCC2

Our previous studies strongly suggest that GABA_A receptors and GABA activity are increased in the neocortex of MAM-treated animals (Jablonska et al. 2004; Poluch et al. 2008). Here we also show that blockade of GABA_A-mediated activity ameliorates the deficits in dynamic activity after MAM treatment. To further assess the role of GABA in mediating the kinetics of migrating GE cells following treatment with MAM, we determined the level of expression of GABA_{Aα2} and GABA_{Aα3} receptor subunits using Western blots. GABA_{Aα2} and GABA_{Aα3} show high expression in both the neocortex and GE early in development, and may play important roles in migration and other developmental events in GE cells (Laurie et al. 1992; Poulter et al. 1992; Fritschy et al. 1994; Wang and Kriegstein 2009; Carlson et al. 2010). In our study, both α2 and α3 subunits are increased in

MAM- treated cortex compared to normal, furthering support for increased GABA_A-mediated activity in our model animals (Figure 10A and B).

To further evaluate the potential of altered environmental GABA to influence migration, we established the amount of KCC2 in normal and MAM-treated cortex. During development, KCC2 maintains a high extracellular Cl⁻ gradient and induces hyperpolarization of cells in response to GABA, playing a role in the developmental switch from the GABA-induced depolarization to hyperpolarization, which may also terminate cell migration (Rivera et al 1999; Bortone and Polleux 2009). After MAM treatment, both monomeric and oligomeric forms of KCC2 were increased in the neocortex as demonstrated using Western blots (Figure 10C, right upper panel). Furthermore, ratiometric analysis reveals a decrease in KCC2 monomer: oligomer ratio (Figure 10C, right lower panel).

Discussion

The kinetic behavior of migrating interneurons, influenced by both intrinsic and environmental factors, plays an important role in the proper positioning of these cells in the neocortex. We developed a model of cortical dysplasia in ferrets by interrupting the development of layer 4 cells resulting in abnormal placement of specific populations of interneurons (Noctor et al. 1999; Poluch et al. 2008). In this study we show that transient and layer-specific interference with corticogenesis leads to: a) alterations in the basal activity of GABA_A and in KCC2 expression b) this change directly influences the kinetic behavior of migrating GE cells. These findings directly and uniquely implicate increased GABA_{AR} and KCC2 activity in impairing dynamic aspects of migration and in the ability of GE cells to actively explore the environment during migration to the cerebral cortex. We demonstrate that details of migration not shown before involving active exploration are strongly influence by ambient GABA_{AR} and KCC2 activity. This may underlie the modified migratory behavior of later born interneurons.

Ferrets signify a valuable developmental model. As the smallest mammal with a convoluted cortex, they represent an important link between studying lissencephalic rodents and non-human primates. Several groups have identified important developmental features in ferrets, including our previous findings that many neurons populating the neocortex are generated in lateral portions of the GE (Poluch et al. 2008). Several recent articles describe a distinct outer subventricular zone (oSVZ) in ferrets, a region found in primates, and suggest that this region may be important for generating cells that populate and expand

the gyrencephalic neocortex (Fietz et al. 2010; Reillo and Borell 2011; Reillo et al. 2011; Liu et al. 2011). The oSVZ contains a population of cells called outer radial glia (oRG; also called tRG), which appear to be the cell type responsible for generating additional cortical cells leading to the expansion of the developing cortex. This idea is contested recently by a group suggesting that the outer SVZ in ferrets does not differ substantially from that region in the rat (Martinez-Cerdeno et al. 2012). Nevertheless, these examples demonstrate the importance of studying developmental processes using several animal models.

MAM treatment alters the kinetic behavior of migrating GE cells

The ability of migrating GE cells to reach appropriate positions within the neocortex depends on the concerted action of extrinsic factors within the neocortex and intrinsic signaling mechanisms that inhibit, facilitate, and guide migrating neurons (Anderso et al. 1997; Powell et al. 2001; Lopez-Bendito et al. 2003 and 2008; Polleux et al. 2003; Stumm et al. 2003; Alifragis et al. 2004; Flames et al. 2004; Cuzon et al. 2006; Cobos et al. 2007; Liodis et al. 2007; Friocourt et al. 2008). Events leading to changes in neocortical architecture may cause variability in the environment that alters the pattern of migration of neuronal cells and results in abnormal cell positioning and cortical dysplasia. In our model of cortical dysplasia, layer 4 of ferret neocortex is diminished resulting in widespread termination of thalamocortical afferents (Palmer et al. 2004), redistribution of GABA_A receptors and parvalbumin and calbindin positive interneurons (Jablonska et al. 2004, Poluch et al. 2008) and changes in the

sequence of information transfer in the somatosensory cortex (McLaughlin et al. 2005). To better understand the mechanisms contributing to the redistribution of interneurons, we evaluated several parameters of kinetic behavior in migrating GE cells. Cells leaving the GE in normal animals migrated significantly faster compared to similar cells in MAM-treated animals. In addition, GE generated neurons in MAM-treated animals exhibited less exploration of the neocortical environment. Exploratory behavior, including extension of multiple leading processes and changes in direction of movement, results from coordinated signaling from various sites, which communicate cues that determine the final location of these cells. The change in the speed of migration and exploratory activity of interneurons following treatment with MAM therefore has important implications on the overall positioning of cells within the neocortex.

Inhibiting GABA_A-mediated activity improves migration of interneurons in MAM-treated animals

GABA is an important influence on migrating cells, although its precise role in the migration of telencephalic cells is complex. GABA receptors are important during different phases of migration and mediate aspects of the initiation, continuation, and termination of cellular movement (Behar et al. 1996, 2001; Bolteus and Bordey 2004; Heck et al. 2007; Bortone and Polleux 2009). Cuzon and colleagues (2006) report that GABA_A activity is present throughout the path of migration of mouse GE cells into the neocortex, but increases in activity as neurons approach and enter their cortical target. Our studies suggest

that MAM treatment results in increased ambient GABA_A-mediated activity, which may interfere with migration of GE cells as they move en route to the neocortex before the final phase identified by Cuzon et al. 2006. This is supported by our original observation of expanded GABA_A immunoreactivity in MAM-treated cortex (Jablonska et al. 2004) and the current finding of increased GABA_{A,2} and GABA_{A,3} receptor subunits as demonstrated by Western blot. Both $\alpha 2$ and $\alpha 3$ subunits of the GABA_A receptor are expressed early during development and may mediate the initial roles of GABA during development (Laurie et al. 1992; Poulter et al. 1992; Fritschy et al. 1994; Carlson et al. 2010). This assists in explaining our finding that elevated levels of GABA interfere with migration of GE cells to slow the pace but not prevent their movement.

To investigate the involvement of GABA_A-mediated transmission in the migration of interneurons, we manipulated the level of GABA_A activity in MAM-treated organotypic cultures using BMI to reduce the ambient GABA_A activity. We predicted that limiting the increased GABA_A activity would change the cellular actions in MAM-treated cortex to be more like normal. In fact, blocking GABA_A activity in MAM-treated animals resulted in significant enhancement of the migration speed of cells leaving the GE as well as increasing the percentage of MAM-treated GE cells extending more than one leading process. This suggests that reducing the effectiveness of ambient GABA in the MAM-treated ferret encourages migratory activity to become more normal.

Increased GABA signaling interferes with migration

Both GABA_{Aα2} and GABA_{Aα3} expression are increased in MAM-treated animals compared to controls. Does increased GABA_{Aα} expression in MAM-treated animals alter the kinetic behavior of GE cell migration? It has been known for several years that GABA acts as an excitatory neurotransmitter early in development to facilitate cell migration, but switches to mediate inhibition as cells mature (Plotkin et al. 1997; Clayton et al. 1998; Rivera et al. 1999). The dual role of GABA is attributed to the temporal expression of two cation-chloride transporters, sodium-potassium-chloride (NKCC1) and KCC2. Migrating neurons initially express NKCC1, which imports chloride ions (Cl⁻) into the cell to maintain a higher intracellular Cl⁻ concentration (Clayton et al. 1998). In response to GABA_A activation, the outward movement of Cl⁻ results in depolarization of the cells and triggers Ca²⁺-mediated events that influence the cytoskeleton and stimulate migration (Behar et al. 2001; Soria & Valdeolmillos 2002). As cells mature, NKCC1 downregulates and coincides with increased expression of KCC2 (Plotkin et al. 1997). KCC2 is a Cl⁻ exporter that maintains a higher extracellular gradient of Cl⁻. Consequently, GABA_A activation when KCC2 is increased results in hyperpolarization and cell inhibition (Payne, 1997; Rivera et al. 1999). Interestingly, in dissociated hippocampal cell cultures, activation of GABA_{AR} induces the expression of NKCC1 and KCC2 (Ganguly et al. 2001). In addition, pharmacologic blockade of GABA_A delayed the expression of KCC2, while increased activity of GABA_{AR} induces precocious expression of the transporter. Premature expression of KCC2 as a result of a heightened state of GABA_A-mediated activity, as we see here, may be responsible for the migration

defects observed in MAM-treated ferrets. KCC2 is implicated in several important processes, including decreased motility as a neuron migrates, which leads to cell slowing, stopping, and differentiating. As a consequence, a premature switch of the GABA_A response may occur and produce early arrest of migrating interneurons. Bortone and Polleux 2009, demonstrated that depolarization through GABA_A receptors stimulates motility of GABAergic migrating neurons while hyperpolarization of GABA_A receptors, induced by KCC2 expression, provides a stop signal for migrating cells. We observed increased expression of both KCC2 monomer and oligomers in the neocortex of young MAM-treated animals. The ratio of KCC2 monomer to oligomer was also reduced in MAM-treated animals, which indicates greater activity of this transporter in our model animals. Oligomerization of KCC2 is an important step in the activation and function of KCC2 as a Cl⁻ exporter (Blaesse et al. 2006). Although others have demonstrated that GABA_{AR} and KCC2 activity are involved in neuronal migration, we demonstrate here that high endogenous levels of these substances directly interfere with migration. In addition they appear specifically important in dynamic and exploratory behavior in cells leaving the ganglionic eminence.

MAM treatment has no effect on the kinetic profile of migrating cortical VZ cells

To determine whether the effect of MAM treatment on the kinetics of migration is restricted to interneurons alone, we also investigated the kinetics of migrating neurons arising from the cortical VZ. Although the migration speed of

CVZ cells was reduced in MAM-treated animals, the deficit was not significant. We also did not observe any significant differences in exploratory behavior between cells in normal animals and those that received MAM treatment, indicating the impact might be restricted to tangentially migrating interneurons. The differential impact of MAM-treatment on the kinetic behavior of GE and CVZ-derived cells relative to normal cells is not surprising since these two cell populations have different neurogenic origin and migrate under the influence of similar but not identical signaling mechanisms (Marin and Rubenstein, 2003).

A working model. A plethora of signaling molecules influence migrating interneurons. GABA initially acts as a stimulating signal to facilitate migration and then acts as a stop signal to arrest the migratory process, an effect that correlates with the sequential expression of NKCC1 and KCC2. An appropriate level of GABA is necessary for the proper migration and positioning of interneurons. Our data suggests that following treatment with MAM, the basal level of GABA_A receptor (GABA_{AR}) activity increases, and the elevated level of GABA_{AR} activity impairs the migration of interneurons, through premature expression of KCC2. Thus, treatments that decrease the level of GABA_{AR} activity enhance the migration of interneurons in part by slowing the development of KCC2.

Acknowledgements: We thank Drs Sylive Poluch, Tarik Haydar, and Marcin Gierdalski for their helpful discussions and scientific assistance. We also thank Latoya Hyson for excellent animal care and technical assistance. This work was supported by funding from the National Institute of Neurological Disorders and Stroke, grant number RO1 24014.

References

Alifragis P, Liapi A, Parnavelas JG. 2004. Lhx6 regulates the migration of cortical interneurons from the ventral telencephalon but does not specify their GABA phenotype. *J Neurosci.* 24:5643-5648.

Anderson AA, Eisenstat SL, Rubenstein JL. 1997. Interneuron migration from basal forebrain to neocortex: dependence on Dlx genes. *Science.* 278:474-476.

Ang ESBC, Haydar TF, Gluncic V, Rakic P. 2003. Four-dimensional migratory coordinates of GABAergic interneurons in the developing mouse cortex. *J Neurosci.* 23(13):5805-5815.

Behar TN, Li YX, Tran HT, Ma W, Dunlap V, Scott C, Barker JL. 1996. GABA stimulates chemotaxis and chemokinesis of embryonic cortical neurons via calcium-dependent mechanisms. *J Neurosci.* 16:1808-1818.

Behar TN, Smith SV, Kennedy RT, McKenzie JM, Maric I, Barker JL. 2001. GABA_B receptors mediate motility signals for migrating embryonic cortical cells. *Cereb Cortex.* 11(8):744-753.

Blaesse P, Guillemain I, Schindler J, Schweizer M, Delpire E, Khiroug L, Friauf E, Nothwang HG. 2006. Oligomerization of KCC2 correlates with development of inhibitory neurotransmission. *J Neurosci.* 26(41):10407-10419.

Bolteus AJ, Bordey A. 2004. GABA release and uptake regulate neuronal precursor migration in the postnatal subventricular zone. *J Neurosci.* 24:7623-7631.

Bortone D, Polleux F. 2009. KCC2 expression promotes the termination of cortical interneuron migration in a voltage-sensitive calcium-dependent manner. *Neuron.* 62:53-71.

Calcagno ME, Paredes MF, Scott C, Baraban SC. 2002. Heterotopic neurons with altered inhibitory synaptic function in an animal model of malformation associated epilepsy. *J Neurosci.* 22(17):7596-7605.

Calcagno ME, Baraban SC. 2003. An examination of calcium current function on heterotopic neurons in hippocampal slices from rats exposed to methylazoxymethanol. *Epilepsia.* 44:315-321.

Carlson VCC, Yeh HH. 2010. GABA_A receptor subunit profiles of tangentially migrating neurons derived from the medial ganglionic eminence. *Cereb Cortex.* 21:1792-1802.

Choi BH, Mathias SC. 1987. Cortical dysplasia associated with massive ectopia of neurons and glial cells within the subarachnoid space. *Acta Neuropathol (Berl)*. 73:105-109.

Clayton GH, Owens GC, Wolff JS, Smith RL. 1998. Ontogeny of cation-Cl⁻ cotransporter expression in rat neocortex. *Brain Res Dev Brain Res*. 109:281-292.

Cobos I, Borello U, Rubenstein JL. 2007. Dlx transcription factors promote migration through repression of axon and dendrite growth. *Neuron*. 54:873-888.

Carlson VCC, Yeh HH. 2010. GABAA receptor subunit profiles of tangentially migrating neurons derived from the medial ganglionic eminence. *Cereb Cortex*. 21:1792-1802.

Cuzon VC, Yeh PWL, Cheng Q, Yeh HH. 2006. Ambient GABA promotes cortical entry of tangentially migrating cells derived from the medial ganglionic eminence. *Cereb Cortex*. 16(10):1577-1588.

Flames N, Long JE, Garratt AN, Fischer TM, Gassmann M, Birchmeier C, Lai C, Rubenstein JLR, Marín O. 2004. Short- and long-range attraction of cortical GABAergic interneurons by neuregulin-1. *Neuron*. 44:251-261.

Fietz SA, Kelava I, Vogt J, Wilsch-Brauninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R, Huttner WB. 2010. OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat Neurosci.* 13:690-699.

Friocourt G, Kanatani S, Tabata H, Yozu M, Takahashi T, Antypa M, Raguene O, Chelly J, Ferec C, Nakajima K, Parnavelas JG. 2008. Cell autonomous roles of ARX in cell proliferation and neuronal migration during corticogenesis. *J Neurosci.* 28:5794-5805.

Fristchy J-M, Paysan J, Enna A, Mohlerl H. 1994. Switch in the expression of rat GABAA-receptor subtypes during postnatal development: an immunohistochemical study. *J Neurosci.* 14(9):5302-5324.

Gal JS, Morozov YM, Ayoub AE, Chatterjee M, Rakic P, Haydar TF. 2006. Molecular and morphological heterogeneity of neural precursors in the mouse neocortical proliferative zones. *J Neurosci.* 26(3):1045-1056.

Ganguly K, Schinder AF, Wong ST, Poo M. 2001. GABA itself promotes the Developmental-switch of neuronal GABAergic responses from excitation to inhibition. *Cell.* 105:521-532.

Gleeson GJ, Walsh CA. 2000. Neuronal migration disorders: from genetic diseases to developmental mechanisms. *Trends Neurosci.* 23:352–359.

Heck N, Kilb W, Reiprich P, Kubota H, Furukawa T, Fukuda A, Luhmann HJ. 2007. GABA_A receptors regulate neocortical neuronal migration in vitro and in vivo. *Cereb. Cortex.* 17:138-148.

Jablonska B, Smith AL, Palmer SL, Noctor SC, Juliano SL. 2004. GABA_A receptors reorganize when layer 4 in ferret somatosensory cortex is disrupted by methylazoxymethanol (MAM). *Cereb. Cortex.* 14(4):432-40.

Laurie DJ, Wisden W, Seeburg PH. 1992. The distribution of thirteen GABA_A receptor subunit mRNAs in the rat brain. III. embryonic and postnatal development. *J Neurosci.* 12(11):4151-4172.

Liodis P, Denaxa M, Grigoriou M, Akufo-Addo C, Yanagawa Y, Pachnis V. 2007. Lhx6 activity is required for the normal migration and specification of cortical interneuron subtypes. *J Neurosci.* 27:3078-3089.

López-Bendito G, Luján R, Shigemoto R, Ganter P, Paulsen O, Molnar Z. 2003. Blockade of GABA_B receptors alters the tangential migration of cortical neurons. *Cereb Cortex.* 13:932-942.

Lopez-Bendito G, Sanchez-Alcaniz JA, Pla R, Borrell V, Pico E, Valdeolmillos M, Marin O. 2008. Chemokine signaling controls intracortical migration and final distribution of GABAergic interneurons. *J Neurosci*. 28:1613-1624.

Lui JH, Hansen DV, Kriegstein AR. 2011. Development and evolution of the human neocortex. *Cell*. 146:18-36.

Marín O, Rubenstein JLR. 2003. Cell migration in the forebrain. *Ann Rev Neurosci*. 26:441-483.

Martinez-Cerdeno V, Cunningham CL, Camacho J, Antczak JL, Prakash AN, Cziep ME, Walker AI, Noctor C. 2012. Comparative analysis of the subventricular zone in rat, ferret and macaque: evidence for an outer subventricular zone in rodents. *Plos one* 7(1):1-37.

Martini FJ, Valiente M, Lopez-Bendito G, Szabo G, Moya F, Valdeolmillos M, Marin O. 2009. Biased selection of leading process branches mediates chemotaxis during tangential neuronal migration. *Development*. 136:41-50.

McLaughlin DF, Juliano S. 2005. Disruption of layer 4 development alters laminar processing in ferret somatosensory cortex. *Cereb Cortex*. 15:1791-1803.

Moroni RF, Inverardi F, Regondi MC, Panzica F, Spreafico R, Frassoni C. 2008. Altered spatial distribution of PV-cortical cells and dysmorphic neurons in the somatosensory cortex of BCNU-treated rat model of cortical dysplasia. *Epilepsia*. 49(5):872-887.

Nadarajah B, Brunstrom JE, Grutzendler J, Wong RO, Pearlman AL. 2001. Two modes of radial migration in early development of the cerebral cortex. *Nat Neurosci*. 4:143-150.

Noctor SC, Palmer SL, McLaughlin DF, Juliano SL. 2001. Disruption of layers 3 and 4 during development results in altered thalamocortical projections in ferret somatosensory cortex. *J Neurosci*. 21:3184-3195.

Palmer SL, Noctor SC, Jablonska B, Juliano SL. 2001. Laminar specific alterations of thalamocortical projections in organotypic cultures following layer 4 disruption in ferret somatosensory cortex. *Eur J Neurosci*. 13:1559-1571.

Palmini A, Andermann F, Olivier A, Tampieri D, Robitaille Y, Andermann E, Wright G. 1991. Focal neuronal migration disorders and intractable partial epilepsy: a study of 30 patients. *Ann Neurol*. 30:741–749.

Payne JA. 1997. Functional characterization of the neuronal-specific K-Cl cotransporter: implications for [K⁺]_o regulation. *Am J Physiol.* 273: C1516-C1525.

Plotkin MD, Snyder EY, Hebert SC, and Delpire E. 1997. Expression of the Na-K-2Cl cotransporter is developmentally regulated in postnatal rat brains: a possible mechanism underlying GABA's excitatory role in immature brain. *J Neurobiol.* 33: 781-795.

Polleux F, Whitford KL, Dijkhuizen PA, Vitalis T, Ghosh A. 2003. Control of cortical interneuron migration by neurotrophins and PI3-kinase signaling. *Development.* 129:3147-3160.

Poluch S, Jablonska B, Juliano SL. 2008. Alteration of interneuron migration in a ferret model of cortical dysplasia. *Cereb Cortex.* 18(1):78-92.

Poulter MO, Barker JL, O'Carroll AM, Lolait SJ, Mahan LC. 1992. Differential and transient expression of GABAA receptor α -subunit mRNAs in the developing rat CNS. *J Neurosci.* 12(8):2888-2900.

Powell EM, Mars WM, Levitt P. 2001. Hepatocyte growth factor/scatter factor is a motogen for interneurons migrating from the ventral to dorsal telencephalon. *Neuron.* 30:79-89.

R Development Core Team. 2009. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.

Reillo I, Borrell V. 2011. Germinal Zones in the Developing Cerebral Cortex of Ferret: Ontogeny, Cell Cycle Kinetics, and Diversity of Progenitors. *Cereb Cortex*.

Reillo I, de Juan Romero C, Garcia-Cabezas MA, Borrell V. 2011. A role for intermediate radial glia in the tangential expansion of the mammalian cerebral cortex. *Cereb Cortex* 21: 1674-1694.

Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtenin H, Lamsa K, Pirvola U, Saarma M, Kaila K. 1999. The K⁺/Cl⁻ co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature*. 397:251-255.

Ross EM, Walsh CA. 2001. Human brain malformations and their lessons for neuronal migration. *Annu Rev Neurosci*. 24:1041-70.

Soria JM, Valdeolmillos M. 2002. Receptor-activated calcium signals in tangentially migrating cortical cells. *Cereb Cortex*. 12(8):831-839.

Stumm RK, Zhou C, Ara T, Lazarini F, Dubois-Dalcq M, Nagasawa T, Holt V, Schulz S. 2003. CXCR4 regulates interneuron migration in the developing neocortex. *J Neurosci.* 23:5123-5130.

Tassi L, Colombo N, Garbelli R, Francione S, Lo Russo G, Mai R, Cossu M, Ferrario A, Galli C, Bramerio M, Citterio A, Spreafico R. 2002. Focal cortical dysplasia: neuropathological subtypes, EEG, neuroimaging and surgical outcome. *Brain.* 125:1719-1732.

Tanaka D, Nakaya Y, Yanagawa Y, Obata K, Murakami F. 2003. Multimodal tangential migration of neocortical GABAergic neurons independent of GPI-anchored proteins. *Development.* 130:5803-5813.

Tanaka D, Maekawa K, Yanagawa Y, Obata K, Murakami F. 2006. Multidirectional and multizonal tangential migration of GABAergic interneurons in the developing cerebral cortex. *Development.* 133:2167-2176.

Taylor DC, Falconer MA, Bruton CJ, Corsellis JAN. 1971. Focal dysplasia of the cerebral cortex in epilepsy. *J Neurol Neurosurg Psychiatry.* 34:369-387.

Wang DD, Kriegstein AR. 2009. Defining the role of GABA in cortical development. *J Neurophys.* 587(9):1873-1879.

Zhu WJ, Roper, SN. 2000. Reduced Inhibition in an Animal Model of Cortical Dysplasia. *J Neurosci.* 20(23):8925-8931.




Figure 1. Images of migrating neurons leaving the ganglionic eminence (GE) of a normal ferret. (A) Example of an organotypic culture obtained from a P0 ferret labeled by electroporation using a plasmid that codes for red fluorescent protein (RFP). (B-C) Higher power views of cells en route to the cortex revealing varied orientations and morphologies of migrating GE cells. B is taken from the boxed in region of A and C is from a different normal slice culture. The arrows indicate the path of the migrating cells. GE: ganglionic eminence; CVZ: Cortical ventricular zone




Figure 2. Migrating GE-derived neurons labeled with Dil. (A) Neurons migrating from the GE in an organotypic culture of a MAM-treated animal display different orientations and positions within the neocortex: tangentially migrating GE cells within the VZ/SVZ region (red arrows), tangentially migrating GE cells within the intermediate zone (yellow arrows), radially-oriented GE cells in the intermediate zone (green arrows) and radially-oriented GE cell directed towards the ventricle (blue arrow). (B) A different organotypic culture of a MAM-treated animal labeled with Dil, showing similar orientations of migrating cells. Arrows mark the different orientations and positions of cells within the neocortex as described above. (C) Higher power images of cells obtained from several labeled organotypic cultures from both normal and MAM-treated animals (the images on the left and right were taken from normal slices); various morphologies can be seen including those with a single leading unbranched process (a), those with single leading but branched processes (b), bipolar cells (c), and cells with multiple leading processes (d). GE: ganglionic eminence; LV: Lateral ventricle




Figure 3. Effect of MAM treatment on the speed and orientation of migrating interneurons leaving the ganglionic eminence and traveling to the neocortex. (A) Neurons leaving the GE travel faster in normal slices compared with those in MAM-treated slices, $n = 390$ cells (normal) and 467 cells (MAM-treated). (B) Cells traveling in either the radial or tangential direction travel slower in MAM treated slices. Tangentially migrating cells: $n = 129$ (normal), 105 (MAM-treated); radially migrating cells: $n = 223$ (normal), 348 (MAM-treated). Significance determined using a Student's t-test; * $p < 0.05$. Error bars = Standard error of mean.




Figure 4. Distribution of cells leaving the GE and traveling to the neocortex. (A) To determine if cells migrated different distances after treatment with MAM roughly concentric regions were created in order to count the neurons that traveled up to 350 μm from the border of the GE VZ, from 350 to 700 μm from that border, or greater than 700 μm from the center of Dil injection. (B) After 48 hrs in culture there were no significant differences in the overall distribution of cells leaving the GE in normal slices vs those in MAM- treated slices, A Chi-Square distribution showed no significant differences. $n=12$ slices (MAM-treated), 10 slices (normal). Error bars = Standard error of mean. GE: ganglionic eminence.




Figure 5. Effect of MAM treatment on the ability of migrating GE cells to extend multiple leading processes and initiate a change in direction of movement. (A) Sequence of images of Dil-labeled GE cells during time-lapse imaging for 8.5 h. Two cells are indicated with either a red or green arrow and followed over time. The cell labeled with the red arrow moves diagonally from the upper left to the approximate center of the image. At 6.5 h it branches and moves toward the right. The cell labeled with a green arrow remains relatively still from 0 - 4h. At this point it begins to move upward and curves to the left, moving out of view at 8.5h. (B) The percent of migrating cells extending new processes was evaluated in normal and MAM-treated slices. After MAM treatment, migrating cells extended fewer new processes and displayed fewer turns/changes in the direction of movement. $n=23$ slices/366 cells (MAM-treated), 21 slices/239 cells (normal); $*P \leq 0.05$, Student's t-Test. Error bars = Standard error of mean.




Figure 6. Tracks of cells in normal and MAM treated slices. (A) The positions of migrating cells were tracked every 30 minutes over a period of 24 h. Each cell is indicated with a different color and a number. This is an example taken from a normal cortex. (B) The tracks of cells leaving the GE of MAM-treated cortex are indicated here. The position of each cell is illustrated with a different color and number. The cells leaving the normal GE show more variability in their route, compared with those leaving the GE of MAM-treated brains. The numbers on the x and y axis indicate the size of the area of the neocortex analyzed while the length of each track indicates a rough estimate of the distance traveled in micrometers. The outline of ferret cortex in A and B indicates the approximate path of cell movement (red arrow).




Figure 7. Administration of bicuculline (BMI) to MAM-treated slices results in an increase in the speed and exploratory behavior of migrating cells. MAM-treated slices were placed in culture with BMI added to the medium. (A) The addition of BMI increased the speed of migration in the treated slices. (B) The addition of BMI also improved the ability of MAM-treated cells to extend processes. The number of turns in the direction of movement did not increase. Significant differences are only shown between the 2 conditions of MAM treated slices, i.e. MAM-Tx alone (red bars) or MAM-Tx with BMI added (gray bars). The normal condition is shown for reference (black bars). BMI treatment also eliminated significant differences between normal and MAM-Tx for speed and extension of processes. Speed: $n = 467$ cells (MAM-treated), 270 cells (MAM-treated + BMI); exploratory behavior: $n = 23$ slices/366 cells (MAM-treated), 16 slices/107 cells (MAM-treated + BMI), * $P \leq 0.05$, Student's t-Test. Error bars = Standard error of mean.




Figure 8. Example of an injection of Dil into the neocortical ventricular zone of an organotypic culture obtained from a MAM-treated animal. The ventricular zone is outlined with a white line. Labeled neurons can be seen migrating toward the neocortex. The inset shows labeled migrating cells from the same image moving into the cortex. The image in the inset was taken from the region indicated with an arrow. LV, lateral ventricle.




Figure 9. Effect of MAM treatment on the speed of migration and exploratory behavior of cells leaving the cortical ventricular zone (CVZ) to the neocortex. (A) There were no significant differences in the speed of migration of cells leaving the CVZ from either normal or MAM-treated slices $n= 191$ cells (normal) and 185 cells (E33 MAM), or (B) in the ability of cells to extend processes or change the direction of movement or turn. Exploratory behavior: $n= 13$ slices/85 cells (normal) and $n= 12$ slices/75 cells (MAM-treated).




Figure 10. Expression of GABA_A receptor subunits and KCC2 following treatment with MAM. (A) Treatment with MAM results in an increase in the expression of GABA_A $\alpha 2$, and (B) GABA_A $\alpha 3$ receptor subunits in the neocortex as demonstrated by Western blots. (C) Both monomeric and oligomeric forms of KCC2 are increased (top bar graph) with significant elevation in the ratio of KCC2 oligomer:monomer (bottom graph). Protein levels were determined by densitometric analysis using ImageJ software. For both normal and MAM-Tx animals, $n = 4$. Significance determined using the Student's t-test, * $P < 0.05$. Error bars = Standard error of mean.

Movie 1. View of cells leaving the ganglionic eminence en route to the cerebral cortex. This movie shows cells leaving the ganglionic eminence that move towards the cerebral cortex, which is located in the upper left of the images. It took place over a period of 16 hours, images were obtained every 30 minutes. Although the overall flow of movement was toward the neocortex, many cells make turns on their way or travel in opposite directions, toward the ventricular zone or in more lateral directions. Cells were labeled with Dil. This is taken from a MAM-treated cortical slice at P0.

Movie 2. Cells exiting the ganglionic eminence migrating to the cerebral cortex. There are fewer cells in this image, which allows easier viewing of details. Individual cells stop and explore the environment without obvious movement. Other cells change direction of movement or extend processes. Nucleokinesis occurs in many cells. This takes place over a period of 16 hours, with images obtained every 30 minutes. This is taken from a MAM-treated slice at P0.

Movie 3. Single cell branching and exploring. This movie shows a single cell that exited the ganglionic eminence and moves toward the cerebral cortex. The cell initially shifts ventrally and then changes direction toward the cortex. It also appears to explore the environment by extending processes. This movie takes place over a period of 9 hours, with sampling every 30 minutes. It is taken from a normal slice at P0.

CHAPTER 3

Targeted disruption of layer 4 formation increases GABA_A receptor-mediated activity in upper cortical layers.

J. Abbah¹ M. F. Braga^{1, 2} and S. L. Juliano^{1, 2}

¹Program in Neuroscience and ²Anatomy, Physiology and Genetics, Uniformed Services University of the Health Sciences, Bethesda, Maryland, 20814-4799

Corresponding Author: Sharon L Juliano: Anatomy, Physiology and Genetics, Uniformed Services University of the Health Sciences, 4301 Jones Bridge Rd., Bethesda, Maryland, 20814-4799

Phone: 301-295-3673

Email: sjuliano@usuhs.edu

Abstract

Cortical dysplasia is frequently associated with various clinical pathologies including epilepsy and mental retardation. This disorder results from impaired migration of immature neurons to their cortical targets, leading to anomalous clustering of neural cells and changes in cortical properties. We developed a model of cortical dysplasia by administering methylazoxymethanol (MAM), an anti-mitotic, to pregnant ferrets on embryonic day 33 (E33) to disrupt the birth of layer 4 cells leading to a significant reduction in the thickness of this layer and a redistribution and increased expression of GABA_A receptors. We evaluated the impact of MAM-treatment on GABA_A receptor-mediated synaptic transmission in both migrating neurons derived from the ganglionic eminence (GE) in organotypic cultures and on pyramidal cells in layer 2/3 of acute slices of postnatal day 28-38 (P28-P38), using whole-cell patch-clamp recording. E33 MAM treatment results in a significant increase in the amplitude and frequency of spontaneous GABA_A receptor-mediated inhibitory post-synaptic currents (IPSCs) in migrating GE cells. In older MAM-treated animals, the amplitude of GABA_A receptor-mediated spontaneous IPSCs in layer 2/3 pyramidal cells increased with no changes in frequency. The amplitude and frequency of miniature IPSCs, however, were significantly increased as a result of MAM-treatment in the older pyramidal cells. The kinetics of GABA_A receptor opening also altered following treatment with MAM. Western blot analysis shows that the expression of GABA_{Aα3} receptor subunit increased in our model animals. Taken together, these observations reinforce the idea that synaptic neurotransmission through GABA_{AR}

is enhanced following treatment with MAM and coincides with our previous finding of increased expression of GABA_{Aα} receptor expression within the upper cortical layers. Overall, we demonstrate that small amounts of toxins delivered during corticogenesis can result in long lasting changes in the neural architecture that manifest as subtle disorders of information processing.

Key words: cortical dysplasia, GABA, patch-clamp, western blotting.

Introduction

During development of the mammalian neocortex, immature neurons generated in the germinal zones of the neocortex and ganglionic eminences (GE) migrate to create the six-layered, inside-out model of the neocortex (Angevine and Sidman 1961; Rakic 1974). This laminar organization is crucial for establishing a balance between excitation and inhibition leading to the overall function of the neocortex. Events with adverse impact on neuronal migration lead to cortical dysplasia (CD), a developmental abnormality characterized by aberrant distribution and clustering of neocortical cells resulting in a plethora of neuropathological disorders such as epilepsy, schizophrenia, and mental retardation (Taylor et al. 1971; Choi and Mathias 1987; Palmini et al. 1991; Gleeson and Walsh 2000; Zhu and Roper 2000; Ross and Walsh 2001; Tassi et al. 2002; Calcagnotto et al. 2002; Calcagnotto and Baraban 2003; Colombo et al. 2003; Guerrini et al. 2003; Moroni et al. 2005, 2008). A common pathological finding in clinical cases and animal models of CD is anomalous distribution and clustering of neurons within the neocortex, which alters the microcircuitry and cortical architecture (Ferrer 1993; Roper 1998; Colacitti et al. 1999; Jacobs et al. 1999; Benardete and Kriegstein, 2002; Moroni et al. 2008). Importantly, abnormal GABA signaling frequently associates with cortical dysplasia leading to consequent alteration in the balance of excitation and inhibition and disruption of normal cortical function (Roper et al. 1999; Zhu and Roper 2000; Benardete and Kriegstein 2002; Xiang et al. 2006; Moroni et al. 2008; Zhou and Roper 2009; Brill and Huguenard 2010; Zhou and Roper 2010, 2011).

We developed a model of cortical dysplasia in ferrets using methylazoxymethanol (MAM). We took advantage of the pharmacokinetic property of MAM as a short-acting anti-mitotic, and the long developmental time course of ferret corticogenesis to selectively disrupt the birth of layer 4 by delivery on embryonic day 33 (E33) (Noctor et al. 1997; Noctor et al. 1999; Palmer et al. 2001). The ferret, as the smallest mammal with a gyrencephalic cortex, is crucial for the study of neocortical development. Because ferrets possess an expanded neocortex with sulci, gyri, and large amounts of white matter it is distinctive as a research subject. The protracted period of corticogenesis and long period of cell proliferation in the large outer subventricular zone may contribute to the expanded cerebral cortex in the ferret compared with the rodent (Fietz et al. 2010). Distinctive features in the ferret related to neocortical development make it essential to study developmental processes in this animal, since aspects of neurodevelopment differ in mammals with a convoluted versus lissencephalic cortex (Kriegstein et al. 2006; Poluch et al. 2008).

As a likely result of the loss of layer 4 in our model, thalamic afferents that normally synapse directly in this layer become redistributed to upper and lower cortical layers; the capacity for entrainment and information transfer is also lost within the somatosensory cortex (Noctor et al. 2001; Palmer et al. 2001; McLaughlin and Juliano, 2005). Ferrets treated with MAM on E33 additionally show an increased expression of GABA_A receptors (GABA_{AR}), which expands to upper cortical layers, and interneurons are disorganized in their laminar positions

(Jablonska et al. 2004; Poluch et al. 2008). Changes in the migratory behavior of GE-derived cells also occur (Abbah and Juliano 2012, under revision). These changes are specific to the antimitotic MAM delivery on E33, as administration of this drug on different embryonic dates leads to dramatically different results (Noctor et al. 1999; Gierdalski and Juliano 2003, 2005; Poluch and Juliano 2007, 2010). An important question emerging from these observations is whether the loss of layer 4 and subsequent redistribution of GABA_{AR} and interneurons alters GABA signaling within upper cortical layers and/or on the global neocortical environment that influences the migration of late-born neurons.

Because our earlier studies in MAM-treated animals showed increased expression of GABA_{AR} within the neocortex as well as an abnormal pattern of migration and distribution specifically for inhibitory cells leaving the GE, we wanted to characterize the electrophysiological properties of both the GABAergic cells migrating toward the neocortex as well as the functional impact of the changed neocortical environment on responses. To investigate if changes in cortical laminar formation alter the overall inhibitory tone within the neocortex and also affect neuronal migration, we applied whole-cell patch-clamp recording to measure spontaneous GABA_{AR}-mediated inhibitory postsynaptic currents (sIPSCs) in two types of cells. We studied cells migrating away from the GE relatively early during ferret cortical development (P0 – P1) and pyramidal cells of layer 2/3 of juvenile ferrets (P28 - P38). Choosing these two types of cells for functional analysis allows us to determine the effect of early environmental changes on migrating GABAergic cells and on inhibitory neurotransmission in the

somatosensory region of the resulting neocortex. Our results indicate that after E33 MAM treatment, GABAergic synaptic neurotransmission is increased significantly in cells migrating into the cortex from the GE and on cells populating the cortex in the upper layers.

Materials and Methods

Handling of animals complied with the Animal Care and Use Committee of the Uniformed Services University of the Health Sciences (USUHS). Pregnant ferrets were obtained from Marshal Farms (New Rose, NY) and maintained in the animal facilities of USUHS. On E33, ferrets were anesthetized with isoflurane (1-2%) and injected IP with 14mg/kg of MAM (Midwest Research Institute, Kansas City, MO). Ferrets were closely monitored and allowed to recover from anesthesia before returning to the animal facility to await delivery of their kits. Ferret kits were used for experiments at P0 - P1 or P28 - P38.

Preparation of Organotypic slices: Preparation of organotypic cultures followed the method described by Palmer et al. (2001). Briefly, P0 ferret kits were anesthetized with sodium pentobarbital (50mg/kg) and after observing insensitivity to pain, their brains were removed and maintained in ice-cold artificial cerebrospinal fluid (aCSF) bubbled with 95% O₂ and 5% CO₂ under aseptic conditions in a laminar flow hood. The aCSF is composed of (in mM): 124 NaCl, 3.2 KCl, 2.4 CaCl₂, 1.2 MgCl₂, 26 NaHCO₃, 1.2 NaH₂PO₄ and 10 glucose. Coronal slices of the neocortex (350-500 µm thick) were generated using a tissue chopper (Stoelting Co., Wood Dale, Illinois). Each slice was plated into a 0.4 µm culture plate insert (Millicell-CM, Bedford, MA) and placed into six-well plates containing Neurobasal media with B27, N2, and G1.2 (containing gentamycin and glutamine) supplements. Slices were incubated at 37°C with 95% O₂ and 5% CO₂.

Cell label: Migrating GE cells were labeled through electroporation using a modification of the method described by Flames et al. (2004). Transfection was achieved using a plasmid that codes for red fluorescent protein (RFP), which was cloned into pCAGGS expression vector (a gift from Dr Tarik Haydar). We applied approximately 1 μ l of the plasmid DNA (5.3 μ g/ μ l) to the ventricular region of the ganglionic eminence (GE) of organotypic slices of ferrets. The cathode of a gene paddle electrode (Harvard Apparatus, Inc., Holliston, MA) was placed within the lateral ventricle close to the GE and the anode situated close to the pia in an appropriate location; plasmid DNA was driven across the organotypic tissue wall by electric pulse generated by a BTX ECM830 pulse generator (Harvard Apparatus, Inc., Holliston, MA). A pulse of 60V current was delivered four times, each lasting for 50 ms at intervals of 950 ms (Gal et al. 2006).

Preparation of acute slices: P28-P38 ferrets were anesthetized with sodium pentobarbital (50 mg/kg i.p.) and then decapitated. Brains were quickly removed and placed in ice-cold aCSF bubbled with 95% O₂ and 5% CO₂. The parietal cortex was blocked and serial coronal sections (400 μ m thick) were cut on a vibratome (LEICA VT1000S, Buffalo Grove, IL). Slices were rapidly transferred into a holding chamber with aerated aCSF maintained at room temperature for 30 min to 1 h after dissection and before electrophysiological recordings. The aCSF was composed of (in mM): 125 NaCl, 2.5 KCl, 1.0 CaCl₂, 2.0 MgCl₂, 25 NaHCO₃, 1.25 NaH₂PO₄ and 25 glucose.

Recordings: Whole-cell patch-clamp recordings were performed on (a) single neocortical pyramidal cells within layer 2/3 of the parietal cortex in acute

slices and (b) GE-derived cells labeled with *RFP* in organotypic slices after 14-20 days in culture. Pyramidal cells were identified by morphology; GE-derived cells were identified under fluorescent microscopy using an upright microscope with 40X objective (ZEISS Axioscope 2FS PLUS). Differential interference optics (DIC) was used for visualizing pyramidal cells. Similarly, cells migrating away from the GE were initially identified using fluorescent microscopy but subsequently visualized under DIC. Only GE cells that have crossed the corticostriatal boundary into the neocortex and display characteristic elongated leading process were selected for recording. All recordings were performed in a submersion type recording chamber where slices were continuously perfused with aerated aCSF composed of (in mM): 125 NaCl, 2.5 KCl, 2.0 CaCl₂, 1.0 MgCl₂, 25 NaHCO₃, 1.25 NaH₂PO₄ and 25 glucose maintained at 28°C for pyramidal cells and 37 °C for cells migrating from the GE. Slices were perfused at a rate of 3-4ml/min. Recordings were performed using a pulled borosilicate pipette filled with an internal solution composed of (in mM): 135 Cs-gluconate, 10 MgCl₂, 0.1 CaCl₂, 1 EGTA, 10 HEPES, 2 Na-ATP, 0.2 Na₃ GTP, pH 7.3 (280–290 mOsm). In some experiments, neurobiotin at a final concentration of 0.2% was included in the internal solution for post-hoc identification. Whole-cell recordings from the soma of neuronal cells occurred in voltage-clamp mode using an Axopatch 200B amplifier (Axon Instruments, Foster City, CA) after a tight seal (1 GΩ) was formed between the recording electrode (with resistance 5-7 mΩ for pyramidal cells and 7-11 mΩ for GE-derived cells) and the cell body. For all recordings, a threshold of 2 kHz was applied to filter signals, which were

recorded and digitized (Digidata1322A, Axon Instruments), stored using pClamp software (Axon Instruments), and analyzed off-line with the Mini Analysis Program (Synaptosoft, Inc., Leonia, NJ). The access resistance was constantly monitored throughout the recording and cells that showed fluctuations beyond 15% were excluded from analysis.

Morphology: After recording, acute slices were immediately fixed in 4% paraformaldehyde overnight at 4°C. After several washes with phosphate buffered saline (PBS), slices were incubated with a solution containing 10% methanol and 0.6% hydrogen peroxide in PBS for 30 min at room temperature. After further washes with PBS, slices were incubated in a solution containing 2% bovine serum albumin (BSA) and 0.75% triton X100 in PBS for 1 h, and then in 2% BSA and 0.1 % triton X100 for 15 min. Slices were then incubated in a dilute solution of ABC elite kit overnight at 4°C. After multiple washes, cells were visualized with peroxidase staining using DAB (Immpact DAB, Vector Laboratories, Inc., Burlingame, CA.), and mounted in moviol.

Western blotting: Pieces of the neocortex cut from blocks of the parietal cortex prepared as described above were frozen on dry ice and preserved at -82°C prior to use. Protein samples were prepared first by homogenizing the tissues using RIPA lysis buffer (Santa Cruz Biotech, Santa Cruz, CA) followed by centrifugation at 14,000 g at 4°C. An estimate of the concentration of protein was obtained through colorimetric assay. Separation of proteins in the samples was accomplished by SDS-PAGE using 10% Bis-Tris gel and separated proteins were transferred to a PVDF membrane (Invitrogen, Carlsbad, CA) using

electrophoresis. A loading volume of 10 to 15 μ l containing 1 μ g of protein was used for each analysis. To limit non-specific binding of antibodies, membranes were initially incubated with Casein blocking buffer (PBS [0.5M NaCl] + 3% Casein + 0.5% Tween-20) for at least 2 h, followed by affinity purified rabbit polyclonal antibodies directed against GABA_{A α 2} (1:200; ProSci Inc., Poway, CA), GABA_{A α 3} (1:2000; Sigma, St. Louis, MO), GABA_{A γ 2} (1:400, Sigma-Aldrich, St Louis, MO.), and monoclonal anti-GAPDH (1:6000, Abcam, Cambridge, MA) for 24 h. After several washes with PBS, protein bands were detected using HRP-conjugated anti-rabbit secondary antibodies (1:1000, Jackson Lab., West Grove, PA) and HRP-conjugated anti mouse (1:6000, Thermo Scientific, Rockford, IL) and visualized using enhanced chemiluminescence detection. Signal intensities were quantified using Image j software (<http://rsb.info.nih.gov/ij>).

Statistics: Student's t-test was applied for all statistical analysis and differences evaluated at $P < 0.05$.

Drugs: The following drugs were added to the recording buffer to isolate sIPSCs and/or mIPSCs: 50 μ M D-(-)-2-Amino-5-phosphonopentanoic acid (D-APV), 20 μ M 5-methyl-10,11-dihydro-5H-dibenzo[a, d] cyclohepten- 5,10-imine [(+)-MK-801], both NMDA receptor antagonists, 20 μ M (–)-(2S)-5,5-dimethyl-2-morpholineacetic acid (SCH50911), GABA_B antagonist; 10 μ M bicuculline methiodide (BMI), a GABA_A receptor antagonist; 10 μ M 6-Cyano-7-nitroquinoxaline-2,3-dione (CNQX), an AMPA/Kainate receptor antagonist, all from Tocris Cookson, Ballwin, MO, 1 μ M Tetrodotoxin (TTX), a sodium channel blocker; (Sigma, St. Louis, MO).

Results

In utero exposure to MAM increases the inhibitory tone mediated by GABA_{AR} in migrating cells originating from the GE.

Migrating cells leaving the GE are influenced by a variety of signaling molecules including GABA. We previously demonstrated that the normal pattern of migration and orientation of GE-derived neurons is altered in MAM-treated animals, in part due to altered GABA_{AR}-mediated activity (Poluch et al. 2008). To determine whether the changes in neuronal migration result from increased GABA_{AR} activity in the neocortex, we evaluated GABA_{AR} receptor-mediated synaptic neurotransmission in cells leaving the GE by recording the spontaneous inhibitory post synaptic currents (sIPSCs). Migrating cells were labeled with a plasmid that codes for red fluorescent protein and identified under fluorescent microscopy. Whole-cell patch-clamp recording was performed to record sIPSCs on GE cells after allowing them to cross the corticostriatal boundary into the neocortex (Figure 1E). sIPSCs were isolated using CNQX (10 μ M), SCH50911 (20 μ M) and AP-5 (50 μ M) and recorded at a holding potential (V_h) of -70mV. The amplitude and frequency of sIPSCs are increased in migrating GE cells relative to the control (Figure 1 A-D), indicating a heightened state of GABAergic neurotransmission in this population of cells.

Treatment with MAM increases the amplitude of spontaneous IPSCs in the pyramidal cells in layer 2/3 of parietal cortex.

Following treatment with MAM on E33, the thickness of layer 4 in ferret somatosensory cortex diminishes leading to reduction in the overall cortical

thickness (Noctor et al. 1999). Additionally, the distribution of GABA_{Aα} receptor subunits is expanded and increased in the upper cortical layers; several subtypes of interneurons are also abnormally positioned (Jablonska et al. 2004; Poluch et al. 2008). To examine the functional implication of the increased expression of GABA_{Aα} receptor subunits on the inhibitory tone within the upper cortical layers, we performed whole-cell patch-clamp recording on pyramidal cells to record sIPSCs. sIPSCs were isolated using CNQX (10 μM), SCH50911 (20 μM) and AP-5 (50 μM) and recorded at a holding potential (V_h) of -70mV. We observed that in P28-38 MAM-treated animals, the amplitude of sIPSCs was significantly increased (40.08 ± 5.61 pA, $P \leq 0.05$) relative to control animals (29.45 ± 2.31 pA, $P \leq 0.05$) (Figure 2 A, B, and C). Our data also indicate that the frequency of sIPSCs was not significantly altered following treatment with MAM (Figure 2D). The increased amplitude of sIPSCs (Figure 2C) in MAM-treated animals with no change in frequency (Figure 2D) suggests that the change in GABA_{Aα}-mediated activity results from enhanced GABA_{Aα} expression in the postsynaptic membrane of layer 2/3 in the parietal cortex.

Several cells were injected with neurobiotin and reacted to visualize their morphology. An example of the types of cells from which we recorded can be seen in Figure 2E and F. All of the cells reported here were similar in morphology and position within the cortex.

In MAM-treated animals, both amplitude and frequency of miniature IPSCs were increased relative to control.

The increase in amplitude of sIPSCs indicates that GABA_{AR}-mediated activity is increased following treatment with MAM. The increase in sIPSC amplitude may be due to changes in postsynaptic response. To determine the underlying mechanism for increased GABA_{AR}-mediated activity, we recorded the action potential independent IPSCs. Miniature IPSCs (mIPSCs) were recorded in the presence of CNQX (10 μ M), SCH50911 (20 μ M), AP-5 (50 μ M) and TTX (1 μ M). Our results show that the amplitude of mIPSCs increased in MAM-treated animals (17.31 ± 1.30 , $n = 24$, $P \leq 0.05$) relative to control (13.08 ± 0.71 ; $n = 20$, $P \leq 0.05$) (Figure 3A, B, and C). Quantitative analysis of the frequency revealed a leftward shift in the inter-event interval indicating increased frequency of mIPSCs generation (Figure 3D). The addition of bicuculline eliminated this response. This result provides further indication of increased postsynaptic GABA_{AR} synaptic transmission in MAM-treated animals.

The kinetics of GABA_A receptors are altered in MAM-treated animals

The change in frequency of mIPSCs in MAM-treated animals suggests that alterations in presynaptic mechanisms may occur, such as alteration in number and the configuration of interneurons, whereas changes in the amplitude of response suggests changes in the postsynaptic membrane, probably due to the observed increase in expression of GABA_A receptors. To further evaluate the factors responsible for these alterations, we evaluated the kinetics of GABA_A receptors. First, we analyzed the 10-90 rise time and compared control with MAM-treated values. The 10-90% rise time provides spatial information about

key synaptic elements and potential alterations due to changes in presynaptic terminal number and positioning (Xie et al. 1997; Wierenga and Wadman 1999; Kobayashi and Buckmaster 2003; Alberto and Hirasawa 2010). In our study, the 10-90% rise time of the sIPSCs and mIPSCs was not significantly different between normal and MAM-treated animals (Figure 4A-I and B-I). This parameter was significantly reduced in cells migrating from the GE in our model animals during the recordings of these cells in organotypic cultures (Figure 4C-I). We next evaluated the decay time constant. In MAM-treated animals, the decay time constant of sIPSCs and mIPSCs in the pyramidal cells was significantly prolonged (Figure 4A-II and B-II). This is also true for the migrating cells originating from the GE (Figure 4C-II). Analysis of the charge transfer (measured as area) also reveals a significant increase in this parameter for the sIPSCs of the pyramidal cells in layer 2-3 (Figure 4A-III) and the cells originating in the GE (Figure 4C-III). This was also true for the recorded mIPSCs (Figure 4B-III).

Passive properties of layer 2/3 pyramidal cells and migrating GE cells are unaffected as a result of treatment with MAM.

We measured the passive properties of both pyramidal cells of layer 2/3 and migrating neurons to evaluate whether the changes in IPSC reported is related to intrinsic properties of the cell. Specifically, we recorded the capacitance, membrane potential across the cell and internal and membrane resistances. These properties provide important developmental clues relating to the level of maturity of the cells. For this analysis, we compared data obtained

from normal cells leaving the GE in organotypic cultures with cells leaving the GE after MAM treatment. We also compared pyramidal cells in normal slices with pyramidal cells in MAM-treated slices. We found no significant differences in the intrinsic properties between normal and MAM-treated cells for either the GE derived cells or pyramidal cells, suggesting that the intrinsic properties are unaffected following MAM treatment (Figure 5 A and B). In each case, there was a reduction in the membrane resistance that was notable, but not significant.

Treatment with MAM increases the expression of GABA_{Aα3} receptor within the parietal cortex of ferrets.

The electrophysiological results presented so far support the idea that GABA_{AR}-mediated activity is increased following treatment with MAM. To further validate the aberrant elevation of GABA_{AR} activity in our model animals, we performed western blot analysis to measure the level of expression of different subtypes of the GABA_{AR} within the somatosensory cortex. We tested several receptor subtypes based on the likelihood of their expression in neocortex at this stage of development in ferrets. Although we expected that GABA_{Aα1} receptors would be present at this age in ferret cortex, we were not able to visualize them. GABA_{Aα2} receptors were present equally in control and MAM-treated cortex (Figure 6 top). The GABA_{Aα3} receptor subunit, however, was increased in the parietal cortex after MAM treatment (Figure 6 middle). We also tested the presence of the $\gamma 2$ subunit and found this to be slightly increased in E33 MAM-treated cortex (Figure 6 bottom). These results indicate that GABA_{AR} expression

is increased, underlying the enhanced GABA_{AR} -mediated synaptic transmission in MAM-treated animals.

Discussion

The current study uses a ferret model of cortical dysplasia to investigate the alterations of GABAergic synaptic transmission recorded in layer 2-3 pyramidal cells of the somatosensory cortex and on migrating GE-derived cells. We show that the basal activity mediated by GABA_{AR} is enhanced within layer 2/3 and in cells leaving the GE in our model animals. An increased expression of GABA_{AR} after MAM treatment is likely to underlie the changes in synaptic transmission. In our model, layer 4 formation is selectively disrupted by *in utero* administration of MAM on E33, leading to its profound reduction in size, a redistribution of interneurons, thalamic afferents, and GABA_{AR} (Noctor et al. 1999; Palmer et al. 2001; Jablonska et al. 2005; Poluch et al. 2008). We also observed that cells leaving the ganglionic eminence and traveling to the neocortex after MAM treatment display abnormal dynamic properties while *en route* (Abbah and Juliano, under revision). Our earlier studies using MAM treatment during mid corticogenesis show a specific disruption of GABA-related features, leading to an apparent overall increase in GABA signaling. This study evaluated the functional implications of these findings in 2 populations of cells, GABAergic neurons leaving the GE to become interneurons and pyramidal cells in layer 2-3 of neocortex, presumably influenced by inhibitory synapses.

Treatment with MAM increases the amplitude of sIPSCs in layer 2/3 pyramidal cells.

An appropriate balance between excitation and inhibition is crucial for normal cortical function. In cortical dysplasia, this equilibrium is disrupted resulting in abnormal cortical activity and function. In pyramidal cells of layer 2/3 we observed an increased amplitude of sIPSCs without any significant change in frequency. This indicates that synaptic transmission through GABA_{AR} is upregulated in MAM-treated animals, which is likely due to the increased GABA_A receptor density on the post synaptic membrane previously observed in MAM-treated animals (Jablonska et al. 2004). To further investigate the mechanism underlying the increase in sIPSC amplitude, we evaluated the mIPSCs in the same layer 2/3 pyramidal cells. mIPSCs are caused by the quantal release of GABA from presynaptic terminals of GABAergic interneurons in the absence of action potentials (Edwards et al. 1990). The number of presynaptic terminals of interneurons and/or the postsynaptic GABA_A receptor density on pyramidal cells determines the frequency of mIPSCs (Shao and Dudek, 2005). In our study, both amplitude and frequency of mIPSC were significantly increased in MAM-treated animals relative to control. The increase in mIPSCs frequency and amplitude signifies that both pre and postsynaptic mechanisms are involved in altering GABAergic neurotransmission in our model animals. Presynaptic factors such as alterations in the number of GABAergic interneurons and the quantal size of release determine the frequency of response. It is possible that the GABAergic neurons in our model are altered in some way to induce a greater number of synapses or greater quantal release. We also know that after MAM treatment, subpopulations of interneurons show altered distributions (Poluch et al. 2008).

Such redistributions may result in ectopic projections or altered synapses with upper cortical layer pyramidal cells. Such region-specific alterations in local circuit connections occur in other models of cortical dysplasia (Brill and Huguenard 2010). Postsynaptic mechanisms such as the density and internalization of postsynaptic protein modulate the size of GABAergic neurotransmission contributing to the change in mIPSC amplitude (DeFazio and Hablitz, 1999; Kittler et al. 2000; Loup et al. 2006; Goodkin et al. 2007; Michels and Moss, 2007; Hartmann et al. 2008). The frequency of sIPSCs, which is the summation of both action potential dependent and independent events, was not altered even though the frequency of mIPSC was increased in MAM-treated animals. A number of factors may be responsible for this, notably the nature of the network of interneurons within layer 2/3 (Tamas et al. 1998; Bacci et al. 2003).

Although reduction in amplitude and frequency of IPSCs are frequently associated with epilepsy and cortical dysplasia (McDonald et al. 1991; Henry et al. 1993; Buckmaster and Dudek 1997; Hablitz and Defazio, 1998; Jacobs et al. 1999; Rocha et al. 2007), the converse occurs in other models of these diseases. Our finding of increased GABAergic neurotransmission in layer 2/3 pyramidal cells corresponds with a recent finding of enhanced GABAergic activity in pediatric cortical dysplasia (Cepeda et al. 2012). Other studies demonstrate a similar increase in GABAergic activity. A freeze lesion model of cortical dysplasia reports increased IPSCs (Prince et al. 1997; DeFazio and Hablitz 1999; Brill and Huguenard 2010). In other models of cortical dysplasia, however, a reduction in

inhibitory tone occurs, such as *in utero* exposure to carmustine, MAM, or radiation in rats (Xiang et al. 2006; Zhou et al. 2009; Zhou and Roper 2010; Karlsson et al. 2011). In these models, the development of seizures is believed to be a consequence of reduced inhibitory drive similar to epileptogenic foci reported in animal models of epilepsy (Rice et al. 1996; Hirsch et al. 1999; Cossart et al. 2001; Kobayashi and Buckmaster 2003; Shao and Dudek 2005; Sun et al. 2007). This analysis is complicated further by evidence of elevated GABAergic signaling in other seizure models (Gibbs et al. 1997; Nusser et al. 1998; Cossart et al. 2001; Shao and Dudek 2005; Zhan and Nadler 2009).

We did not observe any significant variation in the passive properties between MAM-treated and sham animals indicating that the cellular intrinsic properties remain relatively unaffected by MAM treatment. For both the cells migrating away from the GE and the layer 2-3 pyramidal cells, however, the membrane resistance was lower after MAM treatment, compared with control, although the difference was not significant. It is not clear what might cause this change, but it is possible that the membrane characteristics of MAM-treated pyramidal cells have properties that would cause a slight difference in membrane resistance. One possibility is that the MAM-treated brains may mature at a different rate than the normal brains, leading to small distinctions in membrane characteristics (Goldberg et al. 2011).

GABA_A receptor-mediated activity is enhanced in migrating GE-derived neurons after MAM treatment.

The orientation and distribution of interneurons is altered in E33 MAM-treated animals, in part due to elevated GABA_{AR}-mediated activity (Poluch et al., 2008). Recently, we found that elevated GABA_{AR}-mediated activity plays a role in altering the kinetic behavior of migrating cells leaving the GE (Abbah and Juliano, under revision). To determine whether the cells leaving the GE receive abnormally increased GABA_{AR} signaling after MAM treatment, we measured the sIPSCs. Both amplitude and frequency of sIPSCs were increased in MAM-treated animals suggesting that GABA_{AR} synaptic transmission enhances in MAM-treated animals. GABA signaling plays an important role in promoting migration of GE cells and alterations in the basal level of GABA_{AR} activity is likely to be deleterious in the adequate migration of GE-derived cells. Specifically, elevated GABA_{AR}-mediated activity, while initially promoting migration of cells derived from the GE, triggers early maturation of these cells by inducing precocious expression of the potassium chloride co-transporter (KCC2) (Ganguly et al., 1999; Abbah and Juliano, in revision). During development, KCC2 maintains a high extracellular Cl⁻ gradient and induces hyperpolarization of cells in response to GABA, playing a role in the developmental switch from the GABA-induced depolarization to hyperpolarization, which may also terminate cell migration (Rivera et al, 1999; Bortone and Polleux, 2009). Thus the elevated functional responses to GABA_{AR} synaptic transmission in cells migrating away from the GE supports the idea that an aberrant GABAergic environment contributes to both abnormal migration, and disrupted processing of information.

Kinetics of GABA_A signaling is altered in MAM-treated animals.

The measure of 10-90% rise time gives relative spatial information with respect to postsynaptic sites; changes in rise time deduce the locus of change in presynaptic terminal number. It is widely accepted that receptors located in the soma of postsynaptic membrane give a faster rise time compared to those located on the dendrites (Shao and Dudek 2005). In our study there is no significant change in the rise time in either the sIPSCs or the mIPSCs of pyramidal cells in layer 2/3 indicating that there is no significant positional change in the active zones and receptor clusters between normal and MAM-treated animals (Xi et al. 1997; Alberto and Hirasawa 2011). In the cells migrating from the GE, however, we observed a significant reduction in rise time after MAM treatment compared to normal. Although, the reason for this change is unclear, one factor that may be responsible is the relative position of the migrating cells in the developing neocortex, which may affect the nature of synaptic contact they receive. The decay time constant is influenced by the density of postsynaptic receptors and receptor subtype composition (Xi et al. 1997; Shao and Dudek 2005); prolongation of decay time constant that we observed in both the sIPSCs and mIPSCs in MAM-treated animals is consistent with increased expression of GABA_{AR}. The amplitude of the IPSCs and decay time constant determines the charge transfer. In our study, the charge transfer was significantly increased for both sIPSC and mIPSC in cells leaving the GE and layer 2/3 pyramidal cells following treatment with MAM. This is consistent with the increased amplitude and decay time that we reported.

Expression of GABA_A receptor subunits are altered in MAM-treated animals.

Our results show that GABAergic neurotransmission is enhanced in the neocortex of MAM-treated animals, probably as a result of increased expression of GABA_{AR}. We also demonstrated that GABA_{AR} are increased after E33 MAM treatment, primarily in the upper cortical layers using receptor binding and immunohistochemistry (Jablonska et al. 2004). Here we also find an increase in the amplitude of sIPSCs and mIPSCs in pyramidal cells of layer 2/3 and a prolonged decay time. To further confirm that increased GABA_{AR} receptor expression is responsible for the enhanced GABAergic neurotransmission, we applied western blot techniques to measure the level of various subunits of GABA_A receptors. Our data shows that GABA_{Aα3} expression is increased in dysplastic cortex compared to control. The mammalian GABA_{AR} is made of five subunits generated from 19 possible subunits (α 1-6, β 1-3, γ 1-3, δ , ϵ , θ , ρ , and ρ 1-3) - (Olsen and Sieghart 2008). Compelling evidence indicates that these subunits have differences in temporal expression and regional distribution during development (Garrett et al. 1990; McKernan et al. 1991; Araki et al. 1992; Laurie et al. 1992; Fritschy et al. 1994; Pirker et al. 2000; Yu et al. 2006; Hashimoto et al. 2009; Carlson et al. 2010). GABA_{Aα3} subunit is expressed early in development and mediates a number of the developmental activities of GABA (Laurie et al. 1992; Poulter et al. 1992; Fritschy et al. 1994; Carlson et al. 2010). The increased expression of GABA_{Aα3} appears to underlie the enhanced

GABAergic neurotransmission reported here. We also observed increased expression of GABA α 2 and GABA α 3 subunits in the neocortex of MAM-treated animals at P0 (Abbah and Juliano, under review). A possible explanation for increased GABA AR expression in layer 2/3 of parietal cortex in MAM-treated animals is the redistribution of thalamic afferents, which associate closely with GABA AR during development (Noctor et al. 2001; Palmer et al. 2001; Jablonksa et al. 2004). In normal animals, thalamic afferents make synaptic contacts primarily in layer 4 (Jones and Burton 1976). This ensures that afferent information to the neocortex is first received in layer 4 before being transferred to the upper layers and lower layers 5 and 6 (White 1989; Senft and Woolsey 1991; Johnson and Alloway 1996). Thus, in the presence of diminished layer 4, the thalamic afferents and associated GABA AR become redistributed to upper cortical layers; this thalamic redistribution may account for the increase in GABAergic signaling in this region. Others report a close correlation between the expression of GABA AR and projection of thalamic afferents in the somatosensory and visual cortices of rodents (Schlaggar et al. 1994, Broide et al. 1996). Thalamic afferents also play a significant role in the expression and regional distribution of GABA AR subunits within the visual and somatosensory cortical areas of rodents (Paysan et al. 1997). In rats, the expression of the α 1 subunit, the predominant subunit of GABA AR in adult animals, is highest within the primary somatosensory (S1) and visual (V1) cortical areas; these regions also receive substantial terminations of thalamic afferents (Fritschy et al. 1994; Paysan et al. 1994). In S1, the distribution of α 1 receptors mirrors the arrangement of the barrels, the cortical

representation of whiskers. On the other hand, $\alpha 2$, $\alpha 3$ and $\alpha 5$ subunits have low expression in S1 but high expression in the association areas of the somatosensory cortex. Interestingly, perturbation of the thalamic afferents early during development differentially impacts the density and regional distribution of various α subunits within the parietal cortex (Paysan et al. 1997).

Acknowledgements: We thank Drs Volodymyr Pidoplichko and Sylive Poluch for their helpful discussions and scientific assistance. We also thank Latoya Hyson for excellent animal care and technical assistance. This work was supported by funding from the National Institute of Neurological Disorders and Stroke, grant number RO1 24014.

References

Alberto CO, Hirasawa M (2010) AMPA receptor-mediated miniature EPSCs have heterogeneous time courses in orexin neurons. *Biochem Biophys Res Comm* 400:707-712.

Angevine JB, Sidman RL (1961) Autoradiographic study of cell migration during histogenesis of cerebral cortex in the mouse. *Nature* 192:766-768.

Araki T, Kiyama H, Tohyama M (1992) GABA_A receptor subunit messenger RNAs show differential expression during cortical development in the rat brain. *Neuroscience* 51(3):583-591.

Bacci A, Rudolph U, Huguenard JR, and Prince DA (2003) Major differences in inhibitory synaptic transmission onto two neocortical interneuron subclasses. *J Neurosci* 23: 9664-9674.

Benardete EA, Kriegstein AR (2002) Increased excitability and decreased sensitivity to GABA in an animal model of dysplastic cortex. *Epilepsia* 43:970-982.

Bortone D, Polleux F (2009) KCC2 expression promotes the termination of cortical interneuron migration in a voltage-sensitive calcium-dependent manner. *Neuron* 62:53-71.

Brill J, Huguenard JR (2010) Enhanced infragranular and supragranular input onto layer 5 pyramidal neurons in a rat model of cortical dysplasia. *Cereb. Cortex* 20:2926-2938.

Broide RS, Robertson RT, Leslie FM (1996) Regulation of $\alpha 7$ Nicotinic acetylcholine receptors in the developing rat somatosensory cortex by thalamocortical afferents. *J Neurosci* 16(9):2956-2971

Buckmaster PS, Dudek FE (1997) Neuron loss, granule cell axon reorganization, and functional changes in the dentate gyrus of epileptic kainate-treated rats. *J Comp Neurol* 385:385-404.

Calcagnotto ME, Baraban SC (2003) An examination of calcium current function on heterotopic neurons in hippocampal slices from rats exposed to methylazoxymethanol. *Epilepsia* 44:315-321.

Calcagnotto ME, Paredes MF, Scott C, Baraban SC (2002) Heterotopic neurons with altered inhibitory synaptic function in an animal model of malformation associated epilepsy. *J Neurosci* 22(17):7596-7605.

Carlson VCC, Yeh HH (2010) GABA_A receptor subunit profiles of tangentially migrating neurons derived from the medial ganglionic eminence. *Cereb Cortex*. 21:1792-1802.

Cepeda C, Andre VM, Hauptman JS, Yamazaki I, Huynh MyN, Chang JW, Chen JY, Fisher RS, Vinters HV, Levine MS, Mathern GW (2012) Enhanced GABAergic network and receptor function in pediatric cortical dysplasia TyplIB compared with tuberous sclerosis complex. *Neurobiol Dis* 45:310-321.

Choi BH, Matthias SC (1987) Cortical dysplasia associated with massive ectopia of neurons and glial cells within the subarachnoid space. *Acta Neuropathol (Berl)* 73:105-109.

Colacitti C, Sancini G, DeBiasi S, Franceschetti S, Caputi A, Frassoni C, Cattabeni F, Avanzini G, Spreafico R, Di Luca M, Battaglia G (1999) Prenatal methylazoxymethanol treatment in rats produces brain abnormalities with morphological similarities to human developmental brain dysgeneses. *J Neuropathol Exp Neurol* 58:92-106.

Colombo N, Tassi L, Galli C, Citterio A, Lo Russo G, Scialfa G, Spreafico R (2003) Focal cortical dysplasias: MR imaging, histopathologic and clinical correlations in surgically treated patients with epilepsy. *Am J Neuroradiol* 24:724-733.

Cossart R, Dinocourt C, Hirsch JC, Merchan-Perez A, De Felipe J, Ben-Ari Y, Esclapez M, Bernard C (2001) Dendritic but not somatic GABAergic inhibition is decreased in experimental epilepsy. *Nat Neurosci* 4:52-62.

DeFazio RA, Hablitz JJ (1999) Reduction of zolpidem sensitivity in a freeze lesion model of neocortical dysgenesis. *J Neurophysiol* 81:404-407.

Edwards FA, Konnerth A, Sakmann B (1990) Quantal analysis of inhibitory synaptic transmission in the dentate gyrus of rat hippocampal slices: a patch-clamp study. *J Physiol* 430:213-249.

Ferrer I (1993) Experimentally induced cortical malformations in rats. *Childs Nerv Syst* 9:403-407.

Fietz SA, Kelava I, Vogt J, Wilsch-Brauninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R, Huttner WB (2010) oSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat Neurosci* 13:690-699.

Flames N, Long JE, Garrat AN, Fischer TM, Gassmann M, Birchmeier C, Lai C, Rubenstein JLR, Marin O (2004) Short- and long-range attraction of cortical GABAergic interneurons by neuregulin-1. *Neuron* 44:251-261.

Fristchy J-M, Paysan J, Enna A, Mohlerl H (1994) Switch in the expression of rat GABAA-receptor subtypes during postnatal development: an immunohistochemical study. *J Neurosci* 14(9):5302-5324.

Gal JS, Morozov YM, Ayoub AE, Chatterjee M, Rakic P, Haydar TF (2006) Molecular and morphological heterogeneity of neural precursors in the mouse neocortical proliferative zones. *J Neurosci* 26(3):1045-1056.

Garrett KM, Saito N, Duman RS, Abel MS, Ashton RA, Fujimori S, Beer B, Tallmann JF, Vitek MP, Blume AJ (1990) Differential expression of γ -aminobutyric acid_A receptor subunits. *Mol Pharmacol* 37:652-657.

Ganguly K, Schinder AF, Wong ST, Poo M (2001) GABA itself promotes the Developmental-switch of neuronal GABAergic responses from excitation to inhibition. *Cell* 105:521-532.

Gibbs JW, III, Shumate MD, and Coulter DA (1997) Differential epilepsy associated alterations in postsynaptic GABA(A) receptor function in dentate granule and CA1 neurons. *J Neurophysiol* 77:1924-1938.

Gierdalski M, Juliano SL (2003) Factors affecting the morphology of radial glia. *Cereb Cortex* 13(6):572-579.

Gierdalski M, Juliano SL (2005) Endogenous neuregulin restores radial glia in a (ferret) model of cortical dysplasia. *J Neurosci* 25(37):8498-84504.

Gleeson GJ, Walsh CA (2000) Neuronal migration disorders: from genetic diseases to developmental mechanisms. *Trends Neurosci* 23:352-359.

Goldberg EM, Jeong H-Y, Kruglikov I, Tremblay R, Lazarenko RM, Rudy B (2011) Rapid developmental maturation of neocortical FS cell intrinsic excitability. *Cereb Cortex* 21:666-682.

Goodkin HP, Sun CS, Yeh JL, Mangan PS, Kapur J. 2007. GABA(A) receptor internalization during seizures. *Epilepsia*. 48:109-113.

Guerrini R, Sicca F, Parmeggiani L (2003) Epilepsy and malformations of the cerebral cortex. *Epileptic Disord* 5:S9-S26.

Hablitz JJ, Defazio T (1998). Excitability changes in freeze-induced neo- cortical microgyria. *Epilepsy Res*. 32: 75-82.

Hartmann K, Bruehl C, Golovko T, Draguhn A (2008) Fast homeostatic plasticity of inhibition via activity-dependent vesicular filling. *PLoS ONE*. 3:e2979.

Hashimoto T, Nguyen QL, Rotaru D, Keenan T, Arion D, Beneyto M, Gonzalez-Burgos G, Lewis DA (2009) Protracted developmental trajectories of GABA_A

Henry TR, Frey KA, Sackellares JC, Gilman S, Koeppe RA, Brunberg JA, Ross DA, Berent S, Young AB, Kuhl DE (1993) In vivo cerebral metabolism and central benzodiazepine receptor binding in temporal lobe epilepsy. *Neurology* 43:1998-2006.

Hirsch JC, Agassandian C, Merchán-Pérez A, Ben-Ari Y, DeFelipe J, Esclapez M, Bernard C. 1999. Deficit of quantal release of GABA in experimental models of temporal lobe epilepsy. *Nat Neurosci.* 2:499-500.

Jacobs KM, Kharazia VN, Prince DA (1999) Mechanisms underlying epileptogenesis in cortical malformations. *Epilepsy Res* 36:165-188.

Jablonska B, Smith AL, Palmer SL, Noctor SC, Juliano SL (2004) GABA_A receptors reorganize when layer 4 in ferret somatosensory cortex is disrupted by methylazoxymethanol (MAM). *Cereb Cortex* 14(4):432-440.

Jones EG, Burton H (1976) Areal differences in the laminar distribution of thalamic afferents in cortical fields of the insular, parietal and temporal regions of primates. *J Comp Neurol* 168:197-247.

Johnson MJ, Alloway KD (1996) Cross-correlation analysis reveals laminar differences in thalamocortical interactions in the somatosensory system. *J Neurophysiol* 75:1444-1457.

Karlsson A, Lindquist C, Malmgren K, Asztely F (2011) Altered spontaneous synaptic inhibition in an animal model of cerebral heterotopias. *Brain Res* 1383:54-61.

Kittler JT, Delmas P, Jovanovic JN, Brown DA, Smart TG, Moss SJ (2000) Constitutive endocytosis of GABA(A) receptors by an association with the adaptin AP2 complex modulates inhibitory synaptic currents in hippocampal neurons. *J Neurosci* 20:7972-7977.

Kobayashi M, Buckmaster PS (2003) Reduced inhibition of dentate granule cells in a model of temporal lobe epilepsy. *J Neurosci* 23:2440-2452.

Kriegstein A, Noctor S, Martinez-Cerdeno (2006) Patterns of neural stem and progenitor cell division may underlie evolutionary cortical expansion. *Nat Rev Neurosci* 7:883-890.

Laurie DJ, Wisden W, Seeburg PH (1992) The distribution of thirteen GABA_A receptor subunit mRNAs in the rat brain. III. embryonic and postnatal development. *J Neurosci* 12(11):4151-4172.

Loup F, Picard F, Andre VM, Kehrli P, Yonekawa Y, Wieser HG, Fritschy JM (2006) Altered expression of alpha3-containing GABA(A) receptors in the neocortex of patients with focal epilepsy. *Brain* 129:3277-3289.

McDonald JW, Garofalo EA, Hood T, Sackellares JC, Gilman S, McKeever PE, Troncoso JC, Johnston MV (1991) Altered excitatory and inhibitory amino acid receptor binding in hippocampus of patients with temporal lobe epilepsy. *Ann Neurol* 29:529-4.

McKernan RM, Quirk K, Prince R, Cox PA, Gillard NP, Ragan CL, Whiting P (1991) GABA_A receptor subtypes immunopurified from rat brain with a subunit-specific antibodies have unique pharmacological properties. *Neuron* 7:667-76.

McLaughlin DF, Juliano S (2005) Disruption of layer 4 development alters laminar processing in ferret somatosensory cortex. *Cereb Cortex* 15:1791-1803.

Michels G, Moss SJ (2007) GABA(A) receptors: properties and trafficking. *Crit Rev Biochem Mol Biol*. 42:3-14.

Moroni R, Regondi MC, Inverardi F, Garbelli R, Spreafico R, Frassoni C (2005) BCNU-treated rat as a model of cortical dysplasia. In National Congress of the Italian Society for Neuroscience, October 1-4, 2005, Ischia, Italy. p. 647.

Moroni RF, Inverardi F, Regondi MC, Panzica F, Spreafico R, Frassoni C. (2008) Altered spatial distribution of PV-cortical cells and dysmorphic neurons in the somatosensory cortex of BCNU-treated rat model of cortical dysplasia. *Epilepsia* 49(5):872-887.

Noctor SC, Palmer SL, McLaughlin DF, Juliano SL (2001) Disruption of layers 3 and 4 during development results in altered thalamocortical projections in ferret somatosensory cortex. *J Neurosci* 21:3184-3195.

Noctor SC, Scholnicoff NJ, Juliano SL (1997) Histogenesis of ferret somatosensory cortex. *J Comp Neurol* 387:179-193.

Noctor SC, Palmer SL, Hasling T, Juliano SL (1999) Interference with the development of early generated neocortex results in disruption of radial glia and abnormal formation of neocortical layers. *Cereb Cortex* 9:121-136.

Nusser Z, Hajos N, Somogyi P, Mody I (1998) Increased number of synaptic GABA(A) receptors underlies potentiation at hippocampal inhibitory synapses. *Nature* 395:172-77.

Olsen RW, Sieghart W (2008) International Union of Pharmacology. LXX. Subtypes of gamma-aminobutyric acid(A) receptors: classification on the basis of

subunit composition, pharmacology, and function. Update. Pharm Rev 60(3):243-260.

Palmer SL, Noctor SC, Jablonska B, Juliano SL (2001) Laminar specific alterations of thalamocortical projections in organotypic cultures following layer 4 disruption in ferret somatosensory cortex. Eur J Neurosci 13:1559-1571.

Palmini A, Andermann F, Olivier A, Tampieri D, Robitaille Y, Andermann E, Wright G (1991) Focal neuronal migration disorders and intractable partial epilepsy: a study of 30 patients. Ann Neurol 30:741-749.

Paysan J, Bolz J, Mohler H, Fritschy J-M (1994) GABA_A receptor α 1 subunit, an early marker for area specification in developing rat cerebral cortex. J Comp Neurol 350:133-149.

Paysan J, Kossel A, Bolz J, Fritschy J-M (1997) Area specific regulation of γ -aminobutyric acid type A receptor subtypes by thalamic afferents in developing rat cortex. Proc Natl Acad Sci USA 94:6995-7000.

Poulter MO, Barker JL, O'Carroll AM, Lolait SJ, Mahan LC (1992) Differential and transient expression of GABA_A receptor α -subunit mRNAs in the developing rat CNS. J Neurosci 12(8):2888-2900.

Poluch S, Juliano SL (2007) A normal radial glial scaffold is necessary for migration of interneurons during neocortical development. *Glia* 55(8):822-830.

Poluch S, Jablonska B, Juliano SL (2008) Alteration of interneuron migration in a ferret model of cortical dysplasia. *Cereb Cortex* 18(1):78-92.

Poluch and Juliano (2010) Populations of radial glial cells respond differently to reelin and neuregulin1 in a ferret model of cortical dysplasia. *PLoS One* 5(10):1-17

Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G (2000) GABA_A receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 101(4):815-850.

Prince DA, Jacobs KM, Salin PA, Hoffman S, Parada I (1997) Chronic focal neocortical epileptogenesis: does disinhibition play a role? *Can J Physiol Pharmacol*. 75: 500-507.

Rakic P (1974) Neurons in rhesus monkey visual cortex: systematic relation between time of origin and eventual disposition. *Science* 183: 425-427.

Rice A, Rafiq A, Shapiro SM, Jakoi ER, Coulter DA, DeLorenzo RJ (1996) Long-lasting reduction of inhibitory function and gamma-aminobutyric acid type A

receptor subunit mRNA expression in a model of temporal lobe epilepsy. *Proc Natl Acad Sci USA*. 93:9665-69.

Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtenin H, Lamsa K, Pirvola U, Saarma M, Kaila K (1999) The K⁺/Cl⁻ co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* 397:251-255.

Rocha L, Suchomelová L, Mares P, Kubová H (2007) Effects of LiCl/pilocarpine-induced status epilepticus on brain mu and benzodiazepine receptor binding: regional and ontogenetic studies. *Brain Res* 1181:104-17.

Roper SN (1998) In-utero irradiation of rats as a model of human cerebrocortical dysgenesis: a review. *Epilepsy Res* 32:63-74.

Roper SN, Eisenschenk S, King MA (1999) Reduced density of parvalbumin- and calbindin D28-immunoreactive neurons in experimental cortical dysplasia. *Epilepsy Res* 37:63-71.

Ross EM, Walsh CA (2001) Human brain malformations and their lessons for neuronal migration. *Annu Rev Neurosci* 24:1041-70.

Schlaggar BL, O'Leary DDM (1994) Early development of the somatotopic map and barrel patterning in rat somatosensory cortex. *J Comp Neurol* 346:80-96.

Shao LR, Dudek FE (2005) Changes in mIPSCs and sIPSCs after kainate treatment: evidence for loss of inhibitory input to dentate granule cells and possible compensatory responses. *J Neurophysiol* 94:952-960.

Senft SL, Woolsey TA (1991) Growth of thalamic afferents into mouse barrel cortex. *Cereb Cortex* 1:308-335.

Sun C, Mtchedlishvili Z, Bertram EH, Erisir A, Kapur J (2007) Selective loss of dentate hilar interneurons contributes to reduced synaptic inhibition of granule cells in an electrical stimulation-based animal model of temporal lobe epilepsy. *J Comp Neurol* 500:876-93.

Tamas G, Somogyi P, and Buhl EH (1998) Differentially interconnected networks of GABAergic interneurons in the visual cortex of the cat. *J Neurosci* 18:4255-4270.

Tassi L, Colombo N, Garbelli R, Francione S, Lo Russo G, Mai R, Cossu M, Ferrario A, Galli C, Brammerio M, Citterio A, Spreafico R (2002) Focal cortical dysplasia: neuropathological subtypes, EEG, neuroimaging and surgical outcome. *Brain* 125:1719-1732.

Taylor DC, Falconer MA, Bruton CJ, Corsellis JAN (1971) Focal dysplasia of the cerebral cortex in epilepsy. *J Neurol Neurosurg Psychiatry* 34:369-387.

Wierenga CJ and Wadman WJ (1999) Miniature inhibitory postsynaptic currents in CA1 pyramidal neurons after kindling epileptogenesis. *J Neurophysiol* 82:1352-1362.

White EL (1989) *Cortical circuits: synaptic organization of the cerebral cortex-structure, function and theory*. Boston, MA: Birkhauser.

Xiang H, Chen HX, Yu XX, King MA, Roper SN (2006) Reduced excitatory drive in interneurons in an animal model of cortical dysplasia. *J Neurophysiol.* 96:569-578.

Xie X, Liaw JS, Baudry M, Berger TW (1997) Novel expression mechanism for synaptic potentiation: Alignment of presynaptic release site and postsynaptic receptor. *Proc. Natl. Acad. Sci. USA* 94 (1997) 6983-6988.

Yu Z-Y, Wang W, Fritschy J-M, Witte OW, Redecker C (2006) Changes in neocortical and hippocampal GABAA receptor subunit distribution during brain maturation and aging. *Brain Res.* 1099:73-81.

Zhan R-Z, Nadler V (2009) Enhanced tonic GABA current in normotopic and hilar ectopic dentate granule cells after pilocarpine-induced status epilepticus. *J Neurophysiol* 102:670-681.

Zhou FW, Chen HX, Roper SN (2009) Balance of inhibitory and excitatory synaptic activity is altered in fast-spiking interneurons in experimental cortical dysplasia. *J Neurophysiol* 102:2514-2525.

Zhou FW, Roper SN (2010) Densities of glutamatergic and GABAergic pre-synaptic terminals are altered in experimental cortical dysplasia. *Epilepsia* 51:1468-1476.

Zhou FW, Roper SN (2011) Altered Firing rates and patterns in interneurons in experimental cortical dysplasia. *Cereb Cortex* 21:1645-1658.

Zhu WJ, Roper, SN (2000) Reduced Inhibition in an Animal Model of Cortical Dysplasia. *J Neurosci* 20(23):8925-8931.




Figure 1. In MAM-treated animals, both amplitude and frequency of sIPSC is increased in migrating GE cells within the developing cortex. sIPSCs were recorded in a voltage clamp mode at a holding potential of -70mV and isolated using CNQX (10 μ m), SCH50911 (20 μ m), AP-V (50 μ m) and TTX (1 μ m). Representative traces of sIPSCs recorded in a cell leaving the GE in normal (A) or MAM-treated cortex. (B). sIPSCs in both normal and MAM-treated GE cells were blocked by bath application of 10 μ m bicuculline. (C) The amplitude of the sIPSCs is significantly increased in MAM-treated animals relative to control. (D) The cumulative probability plots of sIPSC frequency show a left-ward shift following treatment with MAM, indicating a reduction in inter-event interval and increase in frequency. (E) A schematic diagram of an organotypic slice illustrating the positioning of the *RFP* construct in a pipette and the migratory path of the cells exiting the GE. (F) Image of a cell migrating away from the GE during whole-cell patch-clamp recording. An arrow points to the recording electrode. For group analysis of amplitude and frequency of sIPSC, $n=$ 16 cells (normal) and 21 cells (MAM-treated), $*P < 0.05$, Student T-test. Error bars = Standard error of mean.




Figure 2: Amplitude but not frequency of spontaneous IPSCs (sIPSCs) are increased in pyramidal cells of layer 2/3 of ferret parietal cortex. sIPSC were recorded in voltage clamp mode at a holding potential of -70mV in the presence of CNQX (10 μ m), SCH50911 (20 μ m), AP-V (50 μ m). (A) Representative traces of recorded sIPSCs in pyramidal cells of layer 2/3 of normal cortex. (B) Representative traces of recorded sIPSC in pyramidal cells of layer 2/3 of a MAM-treated cortex. sIPSCs in both normal and MAM-treated pyramidal cells were blocked by bath application of 10 μ m of bicuculline. (C) The amplitude of sIPSC is significantly increased in MAM-treated animals relative to control. (D) The cumulative probability curve of the frequency of sIPSCs indicates that the inter-event interval was not altered following treatment with MAM. For group analysis of amplitude and frequency of sIPSC, $n = 40$ cells (normal) and 24 cells (MAM-treated), * $P < 0.05$, Student T-test. Error bars = Standard error of mean. (E) Example of a pyramidal cell in parietal cortex during whole-cell patch-clamp recording. (F) Post-hoc identification of pyramidal cell in layer 2/3 of the parietal cortex after whole-cell patch-clamp recording.




Figure 3. In MAM-treated animals, both amplitude and frequency of miniature IPSC (mIPSC) is increased in pyramidal cells of layer 2/3 of ferret parietal cortex. mIPSCs were recorded in voltage clamp mode at a holding potential of -70mV and isolated using CNQX (10 μ m), SCH50911 (20 μ m), AP-5 (50 μ m) and TTX (1 μ m). (A) Representative traces of mIPSCs recorded in pyramidal cells of layer 2/3 of a normal cortex. (B) Representative traces of mIPSCs recorded in pyramidal cells of layer 2/3 of a MAM-treated cortex. mIPSCs in both normal and MAM-treated pyramidal cells were block by application of 10 μ m bicuculline. (C) The amplitude of the mIPSCs is significantly increased in MAM-treated animals relative to control. (D) Cumulative probability plots of frequency of mIPSC show a leftward shift following treatment with MAM indicating a reduction in inter-event interval and increase in frequency. For group analysis of the amplitude and frequency of mIPSCs, $n= 20$ cells (normal) and 24 cells (MAM-treated), * $P < 0.05$, Student T-test. Error bars = Standard error of mean.




Figure 4. Kinetics of GABA_A receptors are altered following treatment with MAM. (A) In pyramidal cells, the 10-90 rise time of the sIPSCs was not significantly altered following treatment with MAM (I). However, the decay time constant was significantly prolonged (II); the charge transfer was also significantly increased in MAM-treated animals (III). $n = 40$ cells (normal) and 24 cells (MAM-treated), $*P < 0.05$, Student T-test. (B) Analysis of GABA_R kinetics of the mIPSCs indicates that the 10-90% rise time was unchanged in MAM-treated animals (I). The decay time constant, however, was significantly prolonged (II) and the charge transfer also significantly decreased (III). $n = 20$ cells (normal) and 24 cells (MAM-treated), $*P < 0.05$, Student T-test. (C) Kinetics of GABA_R in migrating GE cells. (I) The 10-90% rise time of recorded sIPSC in migrating GE cells was shortened in MAM-treated animals relative to control. The decay time constant (II) and charge transfer (III) however, increased significantly. $n = 16$ cells (normal) and 21 cells (MAM-treated), $*P < 0.05$, Student T-test. Error bars = Standard error of mean




Figure 5. Treatment with MAM did not significantly alter the passive properties of both pyramidal cells of layer 2/3 and migrating GE-derived cells. (A) Group data from MAM-treated pyramidal cells ($n= 14$) normalized relative to control animals. (B) Group data from MAM-treated migrating GE cells ($n= 13$) normalized relative to control animals. $P < 0.05$, Student T-test. Error bars = Standard error of mean.




Figure 6. Expression of GABA_A receptor subunits following treatment with MAM. Treatment with MAM results in a significant increase in the expression of GABA_{Aα3} receptor subunits and slight elevation of GABA_{Aα2} receptor subunits, but those of GABA_{Aα2} subunits were unchanged, in the neocortex as demonstrated by Western blots. Protein bands were detected through enhanced chemiluminescence and quantitative analysis of the bands performed by densitometric analysis using image J software for both normal

CHAPTER 4

DISCUSSION

Cortical dysplasia (CD) is a developmental abnormality characterized by changes in structure and function of the cerebral cortex. CD often results from the failure of immature neurons to migrate appropriately to their cortical targets. This failure compromises the establishment of the six-layered structure of the cortex and impairs its ability to process information and perform its role to integrate and command. The lack of a satisfactory procedure for managing disorders due to CD underlies the need for development of alternative approaches; this requires critical insight into the mechanisms underlying neuronal migration to comprehend the many disorders that result from disrupted migration. Although recent studies improve our understanding of neuronal migration, significant gaps in knowledge still remain. Moreover, most of these studies were performed in lower animals and rodents, which have important differences compared with higher gyrencephalic mammals like humans and thus necessitate the study of neuronal migration in more advanced animals.

We developed a model of cortical dysplasia in ferrets by interrupting the development of layer 4 cells, which results in abnormal placement of specific populations of interneurons and the redistribution of thalamic afferents and GABA_A receptors (Noctor et al., 1999; Jablonska et al, 2005; Poluch et al., 2008). We demonstrate that the dynamic movement of migrating GE-derived cells is impaired in a ferret model of cortical dysplasia in part due to enhanced GABA_A

receptor (GABA_{AR}) transmission. We also demonstrated that changes in cortical lamination in our model animals impact the microcircuit in upper cortical layers to alter the balance between excitation and inhibition.

Treatment with MAM alters the kinetic behavior of GE-derived cells

In our first set of experiments, we investigated the impact of MAM-treatment on the dynamic pattern of movement of both GE cells and CVZ cells. During development, a well regulated process of neuronal migration involving both intrinsic factors and environmental cues ensures that projection neurons and interneurons that make up each layer of the cortex reach their appropriate targets to create the normal laminar pattern of the cortex (Anderson et al, 1997; Powell et al, 2001; Lopez-Bendito et al., 2003 and 2008; Polleux et al, 2003; Stumm et al., 2003; Alifragis et al., 2004; Flames et al., 2004; Cuzon et al, 2006; Cobos et al., 2007; Liodis et al., 2007; Friocourt et al., 2008). Changes in the proper function of any of the numerous factors that guide migration can lead to abnormal cell positioning and CD. In our model of CD, the neocortical environment is altered as a result of redistribution of thalamic afferents, GABA_A receptors, and a sub-population of interneurons, following the loss of layer 4 (Noctor et al., 1999; Jablonska et al., 2005; Poluch et al., 2008). To better understand the mechanisms contributing to the redistribution of interneurons, we evaluated several parameters of kinetic behavior in migrating GE cells including speed and exploratory tendencies. Cells leaving the GE in normal animals migrated significantly faster compared to similar cells in MAM-treated animals. In addition,

GE generated neurons in MAM-treated animals exhibited less exploration of the neocortical environment. Exploratory behavior, including extension of multiple leading processes and changes in direction of movement, results from coordinated signaling from various sites, which communicate cues that determine the final location of these cells. The change in the speed of migration and exploratory activity of interneurons following treatment with MAM therefore has important implications on the overall positioning of cells within the neocortex.

Is the impact of MAM treatment restricted to cells generated in the GE?

To determine if the effect of E33 MAM treatment on the kinetics of migration is restricted to interneurons alone, we also investigated the dynamic pattern of migration of neurons arising from the cortical VZ. Although the migration speed of CVZ cells was reduced in MAM-treated animals, the deficit was not significant. We also did not observe any significant differences in exploratory behavior between cells arising from the CVZ in normal animals and those receiving E33 MAM treatment, indicating that the impact might be restricted to tangentially migrating GABAergic interneurons. The differential impact of MAM-treatment on the kinetic behavior of GE and CVZ-derived cells relative to normal cells is not surprising since these two cell populations have different neurogenic origin and migrate under the influence of similar but not identical signaling mechanisms (Marin and Rubenstein, 2003). For instance, CVZ cells, which are derived from RG cells, rely predominantly on the structural support of RG for their radial migration (Noctor et al., 2001, 2004). Migration of GE-derived cells is largely

independent of radial glia support. In addition, different transcription programs regulate migration of CVZ and GE-derived cells. Furthermore, different signaling mechanisms provide positional guidance to these two population of cells within the neocortex (for details, see above).

Does treatment with MAM alter the functional properties of migrating GE cells?

Next, we investigated the impact of MAM treatment on the GABA_A receptor activity of migrating GE cells. The orientation and distribution of interneurons is altered in MAM-treated animals in part due to elevated GABA_A-mediated activity (Poluch et al, 2008). To determine whether migrating GE cells receive abnormally increased GABA_{AR} signaling, we recorded sIPSCs using whole-cell patch-clamp recording. Both the amplitude and frequency of sIPSCs were increased in MAM-treated animals suggesting that GABA_{AR}-mediated synaptic transmission is enhanced in our model animals. sIPSC amplitude is influenced by presynaptic factors such as number of inhibitory interneurons in synaptic contact with migrating GE cells as well as the density of GABA_{AR} expressed on the membrane of GE cells migrating to the neocortex (Braga et al., 2003; Shao and Dudek 2005). The frequency is determined by number of presynaptic GABAergic neurons. The GE cells that we recorded from were still in transit enroute to their target destination in the neocortex. Consequently, synaptic contacts with these cells are transient and different regions of the neocortical environment or route of transit influence the level of GABA signaling. In our model, loss of layer 4 leads to redistribution of thalamic afferents and associated GABA_{AR}; this alters the

neocortical environment precisely with respect to GABA signaling to influence migration and positioning of migrating GE cells. The increase in both amplitude and frequency of sIPSCs suggests that both pre-synaptic mechanisms and postsynaptic GABA_{AR} expression contribute to the elevated GABA_{AR}-mediated increase in inhibitory synaptic neurotransmission.

Abnormal elevation in GABA_{AR}-mediated activity may underlie the altered migration pattern in GE-derived cells.

The increase in amplitude and frequency of sIPSCs in cells migrating away from the GE suggests that inhibitory neurotransmission received by GE-derived cells is elevated in E33 MAM-treated animals. In the second set of experiments, we examined the relationship between the abnormal increase in GABA_{AR}-mediated activity and the alteration in GE cell migration. First, we evaluated the basal level of GABA_{AR} by measuring the expression of GABA_A α 2 and GABA_A α 3 subunits in the neocortex. The mammalian GABA_{AR} is made of five subunits generated from 19 possible subunits (α 1-6, β 1-3, γ 1-3, δ , ϵ , θ , ρ , and ρ 1-3) - a pentameric hetero-oligomer (Olsen and Sieghart, 2008). There is compelling evidence based on work from various labs indicating that these subunits have different temporal expression and regional distribution during development (Garrett et al., 1990; Fritschy et al., 1994; Yu et al., 2006; Araki et al., 1992; Laurie et al., 1992; Hashimoto et al., 2009; McKernan et al., 1991; Pirker et al., 2000). Both α 2 and α 3 subunits of the GABA_A receptor are expressed early during development and may mediate the initial roles of GABA during development that includes

regulation of DNA synthesis, cell proliferation, migration, differentiation, development of AMPA receptors, excitatory synapse formation and possibly acting as a stop signal for migrating GE cells (Poulter et al., 1992; Barbin et al., 1993; Heck et al., 2007; LoTurco et al., 1995; Wang and Kriegstein, 2008, 2009; Bortone and Polleux, 2010; Inada et al., 2011). In our study, GABA_{Aα2} and GABA_{Aα3} expression are increased in the neocortex of young (P0 – P1) MAM-treated animals compared to controls. We did not observe any noticeable expression of the α1 subunit in the neocortex of both normal and MAM-treated animals at this age. This is not surprising since this subunit typically show delayed expression during development (Araki et al., 1992; Laurie et al., 1992). The expression of the α1 subunit is accompanied by decrease in the expression of the α2 and α3 subunits; this transition occurs as animals mature (Fristchy et al., 1992).

GABA initially acts as an excitatory neurotransmitter during development to facilitate cell migration, but switches to mediate inhibition as cells mature (Plotkin et al., 1997; Clayton et al., 1998; Rivera et al., 1999). The dual role of GABA is due to the sequential expression of NKCC1 and KCC2. Migrating neurons initially express NKCC1, which imports chloride ions (Cl⁻) into the cell to maintain a higher intracellular Cl⁻ concentration (Clayton et al., 1998). In response to GABA_{AR} activation, the outward movement of Cl⁻ results in depolarization of the cells and triggers Ca²⁺-mediated events that influence the cytoskeleton and stimulate migration (Behar et al., 2001; Soria & Valdeolmillos, 2002). As cells mature, NKCC1 downregulates and coincides with increased

expression of KCC2 (Plotkin et al., 1997). KCC2 is a Cl^- exporter that maintains a higher extracellular gradient of Cl^- . Consequently, GABA_{AR} activation when KCC2 is increased results in hyperpolarization and cell inhibition (Payne, 1997; Rivera et al., 1999). Interestingly, in dissociated hippocampal cell cultures, activation of GABA_{AR} induces the expression of NKCC1 and KCC2 (Ganguly et al., 2001). In addition, pharmacologic blockade of GABA_{A} delayed the expression of KCC2, while increased activity of GABA_{AR} induces precocious expression of the transporter. Premature expression of KCC2 as a result of a heightened state of GABA_{AR} -mediated activity, as we see here, may be responsible for the migration defects observed in MAM-treated ferrets. KCC2 is implicated in several important processes, including decreased motility as a neuron migrates, which leads to cell slowing, stopping, and differentiating. As a consequence, a premature switch of the GABA_{A} response may occur and produce early arrest of migrating interneurons. Bortone and Polleux (2009) demonstrated that depolarization through GABA_{A} receptors stimulates motility of GABAergic migrating neurons while hyperpolarization via GABA_{AR} , induced by KCC2 expression, provides a stop signal for migrating cells. We observed an increased expression of both KCC2 monomer and oligomers in the neocortex of young MAM-treated animals. The ratio of KCC2 monomer to oligomer was also reduced in MAM-treated animals, which indicates greater activity of this transporter in our model animals. Oligomerization of KCC2 is an important step in the activation and function of KCC2 as a Cl^- exporter (Blaesse et al., 2006). These overall changes in KCC2 expression, as a consequence of increased GABA_{AR} actions, may result in a

slowing of interneuron migration, causing early differentiation and maturation of GE-derived cells.

Does blocking GABA_{AR} activity ameliorate the deficit in interneuron migration in MAM-treated animals?

GABA is an important influence on migrating cells, although its precise role in the migration of telencephalic cells is complex. GABA receptors are important during different phases of migration and mediate aspects of the initiation, continuation, and termination of cellular movement (Behar et al., 1996, 2001; Bolteus and Bordey, 2004; Heck et al., 2007; Bortone and Polleux, 2009). Cuzon and colleagues (2006) report that ambient GABA_A activity is present throughout the path of migration of mouse GE cells into the neocortex, but increases in activity as neurons approach and enter their cortical target. Our studies suggest that MAM treatment results in increased ambient GABA_{AR}-mediated activity, which may interfere with migration of GE cells as they move *en route* to the neocortex. The region between the cortical striatal junction and developing cortical plate is identified by Cuzon et al (2006) as a site of normally high levels of GABA, which act as an important factor in the proper movement of GE derived cells. This is supported by our original observation of expanded GABA_{A_α} immunoreactivity in MAM-treated cortex (Jablonska et al., 2004) and the current finding of increased GABA_{A_α3} receptor subunit as demonstrated by Western blot. Both α2 and α3 subunits of the GABA_A receptor are expressed early during development and may mediate the initial roles of GABA during development (Laurie et al., 1992;

Poulter et al., 1992; Fritschy et al., 1994; Carlson et al., 2010). This assists in explaining our finding that elevated levels of GABA interfere with migration of GE cells to slow the pace but not prevent their movement.

To investigate the involvement of GABA_{AR}-mediated transmission in the migration of interneurons, we manipulated the level of GABA_{AR} activity in MAM-treated organotypic cultures using BMI to reduce the ambient GABA_{AR} activity. We predicted that limiting the abnormal increased GABA_{AR} activity would change the cellular actions in MAM-treated cortex to be more like normal. In fact, blocking GABA_{AR} activity in MAM-treated animals resulted in significant enhancement of the migration speed of cells leaving the GE as well as the percentage of MAM-treated GE cells extending more than one leading process. This suggests that reducing GABA_{AR}-mediated signalling in the MAM-treated ferret encourages migratory activity to become more normal.

What is the functional implication of impaired migration of GE-derived neurons?

The data we present here indicates that elevated GABA_{AR}-mediated activity in MAM-treated animals plays a role in the impaired migration of GE-derived cells. An important question that arises from this finding is what impact these changes have on GABAergic neurotransmission in the resulting cortex, which then influences function. We addressed this question by applying whole-cell patch clamp recording to measure IPSCs in pyramidal cells in layer 2/3. We observed increased amplitude of sIPSCs in pyramidal cells of layer 2/3 without any

significant change in frequency. This indicates that GABA_{AR}-mediated activity is higher in MAM-treated animals. To investigate the mechanism underlying the increase in sIPSC amplitude, we evaluated the mIPSC in layer 2/3 pyramidal cells. mIPSCs are caused by quantal release of GABA from presynaptic GABAergic interneurons in the absence of action potentials and thus reflect release probability of GABA (Edwards et al., 1990). The number of presynaptic terminals of interneurons and/or postsynaptic GABA_{AR} density on pyramidal cells determines the amplitude and frequency of mIPSC (Shao and Dudek, 2005). In our study, both amplitude and frequency of mIPSC were significantly increased in MAM-treated animals relative to control. The increase in mIPSC frequency and amplitude that we observed here signifies that both pre and postsynaptic mechanisms may be involved.

We also observed changes in the kinetics of opening of GABA_{AR} in our model animals. The 10-90% rise time gives relative spatial information with respect to postsynaptic sites and changes in rise time could be used to deduce the locus of change in presynaptic terminal number. It is widely accepted that receptors located in the soma of postsynaptic membrane give a faster rise time compared to those located on the dendrites (Shao and Dudek, 2005). In our study there is no significant change in the rise time in the sIPSC and mIPSC of pyramidal cells of layer 2/3 indicating no significant positional change in the active zones and receptor clusters between normal and MAM-treated animals (Xie et al., 1997; Alberto and Hirasawa, 2011). In the migrating GE cells, however, we observed a significant reduction in rise time. Although, the reason

for this change is unclear, one factor that may be responsible is the relative position of the migrating GE cells in the emerging neocortex that may affect the nature of synaptic contact they receive. The decay time constant is influenced by the density of postsynaptic receptors (Xi et al., 1997, Shao and Dudek, 2005); prolongation of decay time constant that we observed in both sIPSC and mIPSC in MAM-treated animals is consistent with increased expression of GABA_A receptors. The amplitude of IPSC and decay time constant determines the charge transfer. In our study, the charge transfer was significantly increased in the sIPSC and mIPSC recorded in both migrating GE cells and layer 2/3 pyramidal cells following treatment with MAM suggests. This is consistent with the increased amplitude and decay time that we reported.

Expression of GABA_A receptor subunits is altered in dysplastic cortex

The result we present above suggests that GABAergic neurotransmission is enhanced in the neocortex of MAM-treated animals, probably as a result of the increased expression of GABA_A receptors. This is evidenced by the increased amplitude of sIPSCs and mIPSCs in pyramidal cells of layer 2/3 and the prolonged decay time in animals P28-38. To further confirm that increased GABA_{AR} expression is responsible for the enhanced GABAergic neurotransmission, we applied western blot to measure the level of various subunits of GABA_{AR} in the older cortex. Our data shows that GABA_{Aα3} expression was significantly increased in our model animals, as it was in the P0 animals. We also observed a slight increase in the expression of GABA_{Aγ2} and no change in

the GABA_{A α 2} subunits in dysplastic cortex compared to control. The increase in GABA_{A α 3} receptor subunit may underscore the elevated GABA transmission in pyramidal cells of MAM-treated juvenile ferrets. The GABA_{A γ 2} receptor subunit is presumed to also be important for synaptic transmission and for the postsynaptic clustering of GABA_{AR} during the development of synapses (Schweitzer et al., 2003). In this regard, it could clearly play a role in the abnormal synaptic transmission observed here. It is not clear why we did not observe an increase or change in the α 1 receptors in either the P0 – P1 or P28 – P38 animals, as we would expect these to increase as the animal matures. It may be that the ferret neocortex is still relatively immature, even at P28 – P38, and does not express high levels of this receptor until a later date. In any case, the obvious increase and probable change in the composition of at least 2 GABA_{AR} subtypes can clearly underlie the change in kinetics that we observed in our recordings.

Conclusion and future directions.

In this set of experiments, we demonstrated that *in utero* administration of MAM on E33 leading to development of CD, selectively interferes with migration of GE-derived cells by altering their kinetic properties. We also showed that the aberrant migration of MAM-treated GE cells results from anomalous increase in GABA_{AR} synaptic transmission within the neocortical environment. This increase appears to enhance the rate of maturation of GE derived cells by inducing the precocious expression of KCC2 (Figure 4). As a result, GE-derived neurons migrate slowly and terminate at aberrant sites within the neocortex; resulting in a

change in cortical microcircuitry that impairs information processing within the cortex. This finding provides evidence that abnormal elevation in GABA signaling during neocortical maturation may underlie the development of CD.

Several neuropsychiatric disorders such as epilepsy, schizophrenia, mental retardation and autism result from abnormal cortical development. Most of these disorders present with impaired GABA function due to loss of inhibitory interneurons resulting in unrestrained excitatory drive in the neocortex. This alters cortical information processing leading to disease phenotypes. At present, the factors responsible for loss of inhibitory interneurons are unknown, although it has been suggested that selective vulnerability of these populations of cells may be responsible. In cortical dysplasia due to defective migration of specific population of cells, aberrant positioning, rather than loss of inhibitory interneurons, impairs cortical function. Therefore, ameliorating the deficit in migration is an appealing approach to management of CD. We postulate that an optimal level of GABA activity is central to proper migration of GE cells and a reduction, as well as increase, in GABA signaling compromises neuronal migration to contribute to disease phenotype. GABA also plays a multifaceted role during development, including regulation of DNA synthesis and intracellular Ca^{2+} levels; these processes may influence aspects of neuronal migration and their functional integration into circuits. A fascinating, but yet unexplored role of GABA during development, is its involvement in regulating gene transcription to influence proliferation and specification of interneurons. This area of thought deserves greater investigation and exploration. In light of strong evidence linking

alterations of GABA signaling to several neuropsychiatric disorders, an intriguing question/possibility is whether early intervention involving restoring/maintaining an optimum level of GABA activity will reduce the severity and improve treatment outcome.

In the future, it will be interesting to specifically evaluate the migratory properties of the GE cells derived from different regions of the LGE or CGE in our model. In addition, functional analysis of other regions of the neocortex of juvenile ferrets will provide further evidence as to whether similar alterations in IPSCs occur in other regions of the MAM-treated neocortex.




Figure 4: Schematic illustration of the cellular mechanisms contributing to defective migration and increase GABAergic neurotransmission in the MAM model of cortical dysplasia. During development, GABA initially acts as an excitatory neurotransmitter due to outward movement of chloride ion that leads to cell depolarization and cell migration in response to GABA_{AR} activation. As cells matured, increased expression of KCC2 reverses the direction of chloride ion movement to cause depolarization. MAM treatment triggers increased GABA_{AR} activity to promote premature expression of KCC2 to cause deficit in GE cell migration.

Reference:

Acampora D, Mazan S, Avantaggiato V, Barone P, Tuorto F, Lallemand Y, Brulet P, Simeone A (1996) Epilepsy and brain abnormalities in mice lacking the Otx1 gene. *Nat Genet* 14:218-222.

Alberto CO, Hirasawa M (2010) AMPA receptor-mediated miniature EPSCs have heterogenous time courses in orexin neurons. *Biochem Biophys Res Comm* 400:707-712.

Alifragis P, Liapi A, Parnavelas JG (2004) Lhx6 regulates the migration of cortical interneurons from the ventral telencephalon but does not specify their GABA phenotype. *J Neurosci* 24:5643-5648.

Anderson SA, Eisenstat DD, Shi L, Rubenstein JL (1997) Interneuron migration from basal forebrain to neocortex: dependence on Dlx genes. *Science* 278:474-476.

Anderson SA, Marin O, Horn C, Jennings K, Rubenstein JLR (2001) Distinct cortical migrations from the medial and lateral ganglionic eminences. *Development* 128:353-363.

Anderson SA, Kaznowski EC, Horn C, Rubenstein JLR, McConnell SK (2002) Distinct origins of neocortical projection neurons and interneurons in vivo. *Cereb Cortex* 12:702-709.

Angevine JB, Sidman RL (1961) Autoradiographic study of cell migration during histogenesis of cerebral cortex in the mouse. *Nature* 192:766-768.

Ang ESBC, Hadar TF, Gluncic V, Rakic P (2003) Four-dimensional migratory coordinates of GABAergic interneurons in the developing mouse cortex. *J Neurosci* 23(13):5805-5815.

Anton ES, Cameron RS, Rakic P (1996) Role of neuron-glial junctional domain proteins in the maintenance and termination of neuronal migration across the embryonic cerebral wall. *J Neurosci* 16:2283-2293.

Arlotta P, Molyneaux BJ, Chen J, Inoue J, Kominami R, Macklis JD (2005) Neuronal subtype-specific genes that control corticospinal motor neuron development in vivo. *Neuron* 45: 207-221.

Araki T, Kiyama H, Tohyama M (1992) GABA_A receptor subunit messenger RNAs show differential expression during cortical development in the rat brain. *Neuroscience* 51(3):583-591.

Ascoli GA, Alonso-Nanclares L, Anderson SA, Barrionuevo G, Benavides-Piccione R, Burkhalter A, Buzsa'ki G, Cauli B, Defelipe J, Fairen A, Feldmeyer D, Fishell G, Frenac Y, Freund TF, Gardner D, Gardner EP, Goldberg JH, Helmstaedter M, Hestrin S, Karube F, Kisvarday ZF, Lambolez B, Lewis DA, Marin O, Markram H, Munoz A, Packer A, Petersen CC, Rockland KS, Rossier J, Rudy B, Somogyi P, Staiger JF, Tamas G, Thomson AM, Toledo-Rodriguez M, Wang Y, West DC, Yuste R (2008) Petilla terminology: nomenclature of features of GABAergic interneurons of the cerebral cortex. *Nat Rev Neurosci* 9:557-568.

Autti-Ramo I, Autti T, Kettunen S, Salonen O, Valanne L (2002) MRI findings in children with school problems who had been exposed prenatally to alcohol. *Dev Med and Child Neurol* 44:98-106.

Ayala R, Shu T, Tsai L-H (2007) Trekking across the brain: the journey of neuronal migration. *Cell* 128:29-43.

Baraban SC, Schwartzkroin PA (1996) Flurothyl seizure susceptibility in rats following prenatal methylazoxymethanol treatment. *Epilepsy Res* 23:189-194.

Barbin G, Pollard H, Gaiarsa JL, Ben-Ari Y (1993) Involvement of GABA_A receptors in the outgrowth of cultured hippocampal neurons. *Neurosci Lett* 152:150-154.

Barinaga M (1993) Carbon monoxide: killer to brain messenger in one step. Science 259: 309.

Batista-Brito R, Fishell G (2009) The developmental integration of cortical interneurons into a functional network. Curr Top Dev Biol 87:81-118.

Bayer SA, Altman J (1991) Neocortical Development. (Raven Press, New York).

Behar TN, Li YX, Tran HT, Ma W, Dunlap V, Scott C, Barke JL (1996) GABA stimulates chemotaxis and chemokinesis of embryonic cortical neurons via calcium-dependent mechanisms. J Neurosci 16:1808-1818.

Behar TN, Schaffner AE, Scott CA, O'Connell C, Barker JL (1998) Differential response of cortical plate and ventricular zone cells to GABA as a migration stimulus. J Neurosci 18:6378-6387.

Behar TN, Scott CA, Greene CL, Wen X, Smith SV, Maric D, Liu Q-Y, Colton CA, Barker JL (1999) Glutamate acting at NMDA receptors stimulates embryonic cortical neuronal migration. J Neurosci 19(11):4449-4461.

Behar TN, Schaffner AE, Scott CA, Greene CL, Barker JL (2000) GABA receptor antagonists modulate postmitotic cell migration in slice cultures of embryonic rat cortex. Cereb Cortex 10:899-909.

Behar TN, Smith SV, Kennedy RT, McKenzie JM, Maric I, Barker JL (2001) GABA_B receptors mediate motility signals for migrating embryonic cortical cells. *Cereb Cortex* 11(8):744-753.

Bellion A, Baudoin JP, Alvarez C, Bornens M, Métin C (2005) Nucleokinesis in tangentially migrating neurons comprises two alternating phases: forward migration of the Golgi/centrosome associated with centrosome splitting and myosin contraction at the rear. *J Neurosci* 25:5691-5699.

Benardete EA, Kriegstein AR (2002) Increased excitability and decreased sensitivity to GABA in an animal model of dysplastic cortex. *Epilepsia* 43:970-982.

Ben-Ari Y, Cherubini E, Corradetti R, Gaiarsa JL (1989) Giant synaptic potentials in immature rat CA3 hippocampal neurons. *J Physiol* 416:303-325.

Bernstein HG, Krause S, Krell D, Dobrowolny H, Wolter M, Stauch R, Ranft K, Danos P, Jirikowski GF, Bogerts (2007) Strongly reduced number of arvalbumin-immunoreactive projection neurons in the mammillary bodies in schizophrenia: Further evidence for limbic neuropathology. *Ann N Y Acad Sci* 1096:120-127.

Bertrand N, Castro DS, Guillemot F (2002) Proneural genes and the specification of neural cell types. *Nature Rev Neurosci* 3:517-530.

Blaesse P, Guillemain I, Schindler J, Schweizer M, Delpire E, Khiroug L, Friauf E, Nothwang HG (2006) Oligomerization of KCC2 correlates with development of inhibitory neurotransmission. *J Neurosci* 26(41):10407-10419.

Bolteus AJ, Bordey A (2004) GABA release and uptake regulate neuronal precursor migration in the postnatal subventricular zone. *J Neurosci* 24:7623-7631.

Bortone D, Polleux F (2009) KCC2 expression promotes the termination of cortical interneuron migration in a voltage-sensitive calcium-dependent manner. *Neuron* 62:53-71.

Braga MFM, Aroniadou-Anderjaska V, Xie J, Li H (2003) Bidirectional modulation of GABA release by presynaptic glutamate receptor 5 Kainate receptors in the basolateral amygdala. *J Neurosci* 23(2):442-452.

Brill J, Huguenard JR (2010) Enhanced infragranular and supragranular synaptic input onto layer 5 pyramidal neurons in a rat model of cortical dysplasia. *Cereb Cortex* 20:2926-2938.

Britanova O, de Juan Romero C, Cheung A, Kwan KY, Schwark M, Gyorgy A, Vogel T, Akopov S, Mitkovski M, Agoston D, Sestan N, Molnar Z, Tarabykin V (2008) Satb2 is a postmitotic determinant for upper-layer neuron specification in the neocortex. *Neuron* 57:378-392.

Brunstrom JE, Gray-Swain MR, Osborne PA, Pearlman AL (1997) Neuronal heterotopias in the developing cerebral cortex produced by neurotrophin-4. *Neuron* 18:505-517.

Butt SJ, Fuccillo M, Nery S, Noctor S, Kriegstein A, Corbin JG, Fishell G (2005) The temporal and spatial origins of cortical interneurons predict their physiological subtype. *Neuron* 48:591-604.

Butt SJ, Cobos I, Golden J, Kessar N, Pachnis V, Anderson S (2007) Transcriptional regulation of cortical interneuron development. *J Neurosci* 27(44):11847-11850.

Butt SJ, Sousa VH, Fuccillo MV, Hjerling-Leffler J, Miyoshi G, Kimura S, Fishell G (2008) The requirement of Nkx2-1 in the temporal specification of cortical interneuron subtypes. *Neuron* 59:722-732.

Calcagnotto ME, Paredes MF, Scott C, Baraban SC (2002) Heterotopic neurons with altered inhibitory synaptic function in an animal model of malformation associated epilepsy. *J Neurosci* 22(17):7596-7605.

Calcagnotto ME, Baraban SC (2003) An examination of calcium current function on heterotopic neurons in hippocampal slices from rats exposed to methylazoxymethanol. *Epilepsia* 44:315-321.

Carlson VCC, Yeh HH (2010) GABA_A receptor subunit profiles of tangentially migrating neurons derived from the medial ganglionic eminence. *Cereb Cortex* 21:1792-1802.

Caronia-Brown G, Grove EA (2011) Timing of cortical interneuron migration is influenced by the cortical hem. *Cereb Cortex* 21(4):748-755.

Casarosa S, Fode C, Guillemot F (1999) Mash1 regulates neurogenesis in the ventral telencephalon. *Development* 126:525-534.

Cavanagh ME, Parnavelas JG (1988) Development of somatostatin immunoreactive neurons in the rat occipital cortex: a combined immunocytochemical-autoradiographic study. *J Comp Neurol* 268:1-12.

Cavanagh ME, Parnavelas JG (1989) Development of vasoactive-Intestinal polypeptide-immunoreactive neurons in the rat occipital cortex: a combined immunohistochemical-autoradiographic study. *J Comp Neurol* 284:637-645.

Caviness VS, Sidman RL (1973) Time of origin of corresponding cell classes in the cerebral cortex of normal and reeler mutant mice: an autoradiographic analysis. *J Comp Neurol* 148:141-151.

Caviness, VS (1982) Neocortical histogenesis in normal and reeler mice: a developmental study based upon [3H]thymidine autoradiography. *Brain Res* 256:293-302.

Caviness VS, Yorke CH (1976) Interhemispheric neocortical connections of the corpus callosum in the reeler mutant mouse: a study based on anterograde and retrograde methods. *J Comp Neurol* 170:449-459.

Cepeda C, Andre VM, Hauptman JS, Yamazaki I, Huynh MyN, Chang JW, Chen JY, Fisher RS, Vinters HV, Levine MS, Mathern GW (2012) Enhanced GABAergic network and receptor function in pediatric cortical dysplasia TpyIIB compared with tuberous sclerosis complex. *Neurobiol Dis* 45: 310-321.

Chae T, Kwon YT, Bronson R, Dikkes P, Li E, Tsai LH (1997) Mice lacking p35, a neuronal specific activator of Cdk5, display cortical lamination defects, seizures, and adult lethality. *Neuron* 18:29-42.

Chao HT, Chen H, Samaco RC, Xue M, Chahrour M, Yoo J, Neul JL, Gong S, Lu H-C, Heintz N, Ekker M, Rubenstein JLR, Noebels JL, Rosenmund C, Zoghbi HY (2010) Dysfunction in GABA signaling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature* 468:263-269.

Chattopaddhyaya B, Di Cristo G (2012) GABAergic circuit dysfunctions in neurodevelopmental disorders. *Front Psych* 3(51):1-9.

Chen B, Schaevitz LR, McConnell SK (2005a) Fezl regulates the differentiation and axon targeting of layer 5 subcortical projection neurons in cerebral cortex. *Proc Natl Acad Sci USA* 102:17184-17189.

Chen JG, Rasin MR, Kwan KY, Sestan N (2005b) Zfp312 is required for subcortical axonal projections and dendritic morphology of deep-layer pyramidal neurons of the cerebral cortex. *Proc. Natl Acad Sci USA* 102: 17792-17797.

Chen B, Wang SS, Hattox AM, Rayburn H, Nelson SB, McConnell SK (2008) The Fezf2-Ctip2 genetic pathway regulates the fate choice of subcortical

projection neurons in the developing cerebral cortex. *Proc Natl Acad Sci USA* 105:11382-11387.

Chenn A, McConnell SK (1995) Cleavage orientation and the asymmetric inheritance of Notch1 immunoreactivity in mammalian neurogenesis. *Cell* 82:631-641.

Cherubini E, Gaiarsa JL, Ben-Ari Y (1991). GABA: an excitatory transmitter in early postnatal life. *Trends Neurosci* 14(12):515-519.

Chevassus-au-Louis N, Ben-Ari Y, Vergnes M (1998) Decreased seizure threshold and more rapid rate of kindling in rats with cortical malformation induced by prenatal treatment with methylazoxymethanol. *Brain Res* 812:252-255.

Chiang C, Litingtung Y, Lee E, Young KE, Corden JL, Westphal H, Beachy PA (1996) Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* 383:407-413.

Choi BH, Matthias SC (1987) Cortical dysplasia associated with massive ectopia of neurons and glial cells within the subarachnoid space. *Acta Neuropathol* (Berl) 73:105-109.

Chugani HT, Shields WD, Shewmon DA, Olson DM, Phelps ME, Peacock WJ (1990) Infantile spasms: I. PET identifies focal cortical dysgenesis in cryptogenic cases for surgical treatment. *Ann Neurol* 27:406-413.

Clayton GH, Owens GC, Wolff JS, Smith RL (1998) Ontogeny of cation-Cl⁻ cotransporter expression in rat neocortex. *Brain Res Dev Brain Res* 109:281-292.

Cobb SR, Buhl EH, Halasy K, Paulsen O, Somogyi P (1995) Synchronization of neuronal activity in hippocampus by individual GABAergic interneurons. *Nature* 378:75-78.

Cobos I, Broccoli V, Rubenstein JL (2005a) The vertebrate ortholog of *Aristaless* is regulated by *Dlx* genes in the developing forebrain. *J Comp Neurol* 483:292-303.

Cobos I, Calcagnotto ME, Vilaythong AJ, Thwin MT, Noebels JL, Baraban SC, Rubenstein JL (2005b) Mice lacking *Dlx1* show subtypespecific loss of interneurons, reduced inhibition and epilepsy. *Nat Neurosci* 8:1059-1068.

Cobos I, Borello U, Rubenstein JL (2007) *Dlx* transcription factors promote migration through repression of axon and dendrite growth. *Neuron* 54:873-888.

Cocas LA, Georgala PA, Mangin JM, Clegg JM, Kessarar N, Haydar TF, Gallo V, Price DJ, Corbin JG (2011) Pax6 is required at the telencephalic pallial-subpallial boundary for the generation of neuronal diversity in the postnatal limbic system. *J Neurosci* 6:5313-5324.

Colacitti C, Sancini G, DeBiasi S, Franceschetti S, Caputi A, Frassoni C, Cattabeni F, Avanzini G, Spreafico R, Di Luca M, Battaglia G. (1999) Prenatal methylazoxymethanol treatment in rats produces brain abnormalities with morphological similarities to human developmental brain dysgeneses. *J Neuropathol Exp Neurol* 58:92-106.

Colombo N, Tassi L, Galli C, Citterio A, Lo Russo G, Scialfa G, Spreafico R (2003) Focal cortical dysplasias: MR imaging, histopathologic and clinical correlations in surgically treated patients with epilepsy. *Am J Neuroradiol* 24:724-733.

Colombo E, Collombat P, Colasante G, Bianchi M, Long J, Mansouri A, Rubenstein JLR, Broccoli V (2007). Inactivation of Arx, the murine ortholog of the X-linked lissencephaly with ambiguous genitalia gene, leads to severe disorganization of the ventral telencephalon with impaired neuronal migration and differentiation. *J Neurosci* 27:4786-4798.

Corbin JG, Nery S, Fishell G (2001) Telencephalic cells take a tangent: non-radial migration in the mammalian forebrain. *Nat Neurosci* 4:1177-1182.

Corbin JG, Gaiano N, Juliano SL, Poluch S, Stancik E, Haydar TF (2008) Regulation of neural progenitor cell development in the nervous system. *J Neurochem* 106:2272-2287.

Corbin JG, Butt SJB (2011) Developmental mechanisms for the generation of telencephalic interneurons. *Dev Neurol* 71(8):710-732.

Crandall JE, McCarthy DM, Araki KY, Sims JR, Ren JQ, Bhide PG (2007) Dopamine receptor activation modulates GABA neuron migration from the basal forebrain to the cerebral cortex. *J Neurosci* 27:3813-3822.

Crandall JE, Goodman T, McCarthy MD, Gregg D, Bhide PG, Grager UC, McCaffery P (2011) Retinoic acid influences neuronal migration from the ganglionic eminence to the cerebral cortex. *J Neurochem* 119:723-735

Cuzon VC, Yeh PWL, Cheng Q, Yeh HH (2006) Ambient GABA promotes cortical entry of tangentially migrating cells derived from the medial ganglionic eminence. *Cereb Cortex* 16(10):1577-1588.

D'Arcangelo G, Miao GG, Chen SC, Soares HD, Morgan JI, Curran T (1995) A protein related to extracellular matrix proteins deleted in the mouse mutant reeler. *Nature* 374:719-723.

De Carlos JA, L'opez-Mascaraque L, Valverde F (1996) Dynamics of cell migration from the lateral ganglionic eminence in the rat. *J Neurosci* 16:6146-6156.

DeDiego I, Smith-Fernandez A, Fairen A (1994) Cortical cells that migrate beyond area boundaries: characterization of an early neuronal population in the lower intermediate zone of prenatal rats. *Eur J Neurosci* 6:983-997.

DeFelipe J, Farinas I (1992) The pyramidal neuron of the cerebral cortex: morphological and chemical characteristics of the synaptic inputs. *Prog Neurobio* 39:563-607.

De Felipe J (1993) Neocortical neuronal diversity: chemical heterogeneity revealed by colocalization studies of classic neurotransmitters, neuropeptides, calcium-binding proteins, and cell surface molecules. *Cereb.Cortex* 3:273-289.

de Feo MR, Mecarelli O, Ricci GF (1995) Seizure susceptibility in immature rats with microencephaly induced by prenatal exposure to methylazoxymethanol acetate. *Pharmacol. Res.* 31:109-114.

Delpire E (2000) Cation-chloride cotransporters in neuronal communication. *News Physiol Sci* 15:309-312.

Denaxa M, Chan CH, Schachner M, Parnavelas JG, Karagogeos D (2001) The adhesion molecule TAG-1 mediates the migration of cortical interneurons from the ganglionic eminence along the corticofugal fiber system. *Development* 128:4635-4644.

de Rouvroit CL, Goffinet AM (2001) Neuronal migration. *Mech Dev* 105:47-56.

des Portes V, Francis F, Pinard JM, Desguerre I, Moutard ML, Snoeck I, Meiners LC, Capron F, Cusmai R, Ricci S, Motte J, Echenne B, Ponsot G, Dulac O, Chelly J, Beldjord C (1998) Doublecortin is the major gene causing X-linked subcortical laminar heterotopia (SCLH). *Hum. Mol Genet* 7:1063-1070.

Dulabon L, Olson EC, Taglienti MG, Eisenhuth S, McGrath B, Walsh CA, Kreidberg JA, Anton ES (2000) Reelin binds $\alpha 3 \beta 1$ integrin and inhibits neuronal migration. *Neuron* 27:33-44.

Edwards FA, Konnerth A, Sakmann B (1990) Quantal analysis of inhibitory synaptic transmission in the dentate gyrus of rat hippocampal slices: a patch-clamp study. *J Physiol* 430:213-249.

Eksioglu YZ, Scheffer IE, Cardenas P, Knoll J, DiMario F, Ramsby G, Berg M, Kamuro K, Berkovic SF, Duyk GM, Parisi J, Huttenlocher PR, Walsh CA (1996) Periventricular heterotopia: an X-linked dominant epilepsy locus causing aberrant cerebral cortical development. *Neuron* 16:77-87.

Fairen A, DeFelipe J, Regidor J (1984) Nonpyramidal neurons: general account. In *Cereb. Cortex.* (1) (ed. A. Peters and E. G. Jones), pp. 201-253. New York: Plenum Press.

Fairen A, Cobas A, Fonseca M (1986) Times of generation of glutamic acid decarboxylase immunoreactive neurons in mouse somatosensory cortex. *J Comp Neurol* 251:67-83.

Fatemi SH, Reutiman TJ, Folsom TD, Rooney RJ, Patel DH, Thuras PD (2010) mRNA and protein levels for GABA α 4, α 5, β 1 and GABABR1 receptors are altered in brains from subjects with autism. *J Autism Dev Disord* 40:743-750.

Fatemi SH, Reutiman TJ, Folsom TD, Thuras PD (2009) GABA α receptor down regulation in brains of subjects with autism. *J Autism Dev Disord* 39:223-230.

Fauser S, Zentner J (2012) Management of cortical dysplasia in epilepsy. *Adv Tech Stand Neurosurg* 38:137-163.

Feng Y, Walsh CA (2001) Protein-protein interactions, cytoskeletal regulation and neuronal migration. *Nat Rev Neurosci* 2:408-416.

Ferrer I (1993) Experimentally induced cortical malformations in rats. *Childs Nerv Syst* 9:403-407.

Ferrer I, Alcantara S, Z'ujar MJ, Cinos C (1993) Structure and pathogenesis of cortical nodules induced by prenatal X-irradiation in the rat. *Acta Neuropathol* 85:205-212.

Fietz SA, Kelava I, Vogt J, Wilsch-Brauninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R, Huttner WB (2010) oSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat Neurosci* 13:690-699.

Fishell G, Kriegstein AR (2003) Neurons from radial glia: the consequences of asymmetric inheritance. *Curr Opin Neurobiol* 13: 34-41.

Fishell G, Hanashima C (2008) Pyramidal neurons grow up and change their mind. *Neuron* 57:333-338.

Flames N, Long JE, Garratt AN, Fischer TM, Gassmann M, Birchmeier C, Lai C, Rubenstein JLR, Marín O (2004) Short- and long-range attraction of cortical GABAergic interneurons by neuregulin-1. *Neuron* 44:251-261.

Fode C, Ma Q, Casarosa S, Ang S-L, Anderson DJ, Guillemot F (2000) A role for neural determination genes in specifying the dorsoventral identity of telencephalic neurons. *Genes Dev* 14:67-80.

Fogarty M, Grist M, Gelman D, Marín O, Pachnis V, Kessaris N (2007) Spatial genetic patterning of the embryonic neuroepithelium generates GABAergic interneuron diversity in the adult cortex. *J Neurosci* 27:10935-10946.

Fonseca M, Soriano E, Ferrer I, Martinez A, Tunon T (1993) Chandelier cell axons identified by parvalbumin-immunoreactivity in the normal human temporal cortex and in Alzheimer's disease. *Neuroscience* 55:1107-1116.

Freund TF (2003) Interneuron diversity series: rhythm and mood in perisomatic inhibition. *Trends Neurosci* 26:489-495.

Friocourt G, Liu JS, Antypa M, Rakic S, Walsh CA, Parnavelas JG (2007) Both doublecortin and doublecortin-like kinase play a role in cortical interneuron migration. *J Neurosci* 27:3875-3883.

Friocourt G, Kanatani S, Tabata H, Yozu M, Takahashi T, Antypa M, Raguene O, Chelly J, Ferec C, Nakajima K, Parnavelas JG (2008) Cell autonomous roles of *ARX* in cell proliferation and neuronal migration during corticogenesis. *J Neurosci* 28:5794-5805.

Fristch B, Qashu F, Figueiredo TH, Aroniadou-Anderjaska V, Rogawski MA, Braga MFM (2009). Pathological alterations in GABAergic interneurons and reduced tonic inhibition in the basolateral amygdala during epileptogenesis. *Neuroscience* 163(1):415-429.

Fristchy J-M, Paysan J, Enna A, Mohlerl H (1994) Switch in the expression of rat GABA-receptor subtypes during postnatal development: an immunohistochemical study. *J Neurosci* 14(9):5302-5324.

Frotscher M (1997) Dual role of Cajal-Retzius cells and reelin in cortical development. *Cell Tissue Res* 290:315-322.

Gal JS, Morozov YM, Ayoub AE, Chatterjee M, Rakic P, Haydar TF (2006) Molecular and morphological heterogeneity of neural precursors in the mouse neocortical proliferative zones. *J Neurosci* 26(3):1045-1056.

Ganguly K, Schinder AF, Wong ST, Poo M (2001) GABA itself promotes the Developmental-switch of neuronal GABAergic responses from excitation to inhibition. *Cell* 105:521-532.

Gant JC, Thibault O, Blalock EM, Yang J, Bachstetter A, Kotick J, Schauwecker PE, Hauser KF, Smith GM, Mervis R, Li Y, Barnes GN (2009) Decreased number of interneurons and increased seizures in neuropilin 2 deficient mice: implications for autism and epilepsy. *Epilepsia* 50:629-645.

Garrett KM, Saito N, Duman RS, Abel MS, Ashton RA, Fujimori S, Beer B, Tallmann JF, Vitek MP, Blume AJ (1990) Differential expression of γ - aminobutyric acid_A receptor subunits. *Mol Pharmacol* 37:652-657.

Garcia-Bellido A (1979) Genetic analysis of the achaete-scute system of *Drosophila melanogaster*. *Genetics* 91:491-520.

Ge W, He F, Kim KJ, Blanch B, Coskun V, Nguyen L, Wu X, Zhao J, Heng JI-T, Martinowich K, Tao J, Wu H, Castro D, Sobeih M, Corfas G, Gleeson JG, Greenberg ME, Guillemot F, Sun YE (2006) Coupling of cell migration with neurogenesis by proneural bHLH factors. *Proc Natl Acad Sci USA*.103:1319-1324.

Germano IM, Zhang YF, Sperber EF, Moshe SL (1996) Neuronal migration disorders increase susceptibility to hyperthermia-induced seizures in developing rats. *Epilepsia* 37:902-910.

Ghashghaei HT, Weber J, Pevny L, Schmid R, Schwab MH, Lloyd KC, Eisenstat DD, Lai C, Anton ES. 2006. The role of neuregulin-ErbB4 interactions on the proliferation and organization of cells in the subventricular zone. *Proc Natl Acad Sci U S A*. 103:1930-1935.

Ghysen A, Dambly-Chaudiere C (1988) From DNA to form: the achaete-scute complex. *Genes Dev* 2:495-501.

Gierdalski M, Juliano SL (2003) Factors affecting the morphology of radial glia. *Cereb Cortex* 13(6):572-579.

Gierdalski M, Juliano SL (2005) Endogenous neuregulin restores radial glia in a (ferret) model of cortical dysplasia. *J Neurosci* 25(37):8498-84504.

Gilbert CD, Wiesel TN (1985) Intrinsic connectivity and receptive field properties in visual cortex. *Vision Res* 25:365-374.

Ginsberg MD, Myers RE (1974) Experimental carbon monoxide encephalopathy in the primate. I. Physiologic and metabolic aspects. *Arch Neurol* 30:202-208.

Ginsberg MD, Myers RE, McDonagh BF (1974) Experimental carbon monoxide encephalopathy in the primate. II. Clinical aspects, neuropathology, and physiologic correlation. Arch Neurol 30:209-216.

Gleeson JG, Allen KM, Fox JW, Lamperti ED, Berkovic S, Scheffer I, Cooper EC, Dobyns WB, Minnerath SR, Ross MH, Walsh CA (1998) doublecortin, a brain-specific gene mutated in human X-linked lissencephaly and double cortex syndrome, encodes a putative signaling protein. Cell 92: 63-72.

Gleeson JG, Minnerath, SR, Fox JW, Allen KM, Luo RF, Hong SE, Berg MJ, Kuzniecky R, Reitnauer PJ, Borgatti R, Mira AP, Guerrini R, Holmes GL, Rooney CM, Berkovic S, Scheffer I, Cooper EC, Ricci S, Cusmai R, Crawford TO, Leroy R, Andermann E, Wheless JW, Dobyns WB, Walsh CA (1999) Characterization of mutations in the gene doublecortin in patients with double cortex syndrome. Ann Neurol 45:146-153.

Gleeson GJ, Walsh CA (2000) Neuronal migration disorders: from genetic diseases to developmental mechanisms. Trends Neurosci 23:352-359.

Gorman DF, Huang YL, Williams C (2001) Prolonged exposure to one percent carbon monoxide causes a leucoencephalopathy in un-anaesthetised sheep. Toxicology 165: 97-107.

Gonzalez F, Romani S, Modolell J, Campuzano S (1989) Molecular analysis of the *asense* gene, a member of the *achaete-scute* complex of *Drosophila melanogaster*, and its novel role in optic lobe development. EMBO J 8:3553-3562.

Gotz M, Huttner WB (2005) The cell biology of neurogenesis. Nat Rev Neurosci 6:777-788.

Götz M, Hartfuss E, Malatesta P (2002) Radial glial cells as neuronal precursors: a new perspective on the correlation of morphology and lineage restriction in the developing cerebral cortex of mice. Brain Res Bull 57:777-788.

Gray GE, Glover JC, Majors J, Sanes JR (1988) Radial arrangement of clonally related cells in the chicken optic tectum: lineage analysis with a recombinant retrovirus. Proc Natl Acad Sci USA 85:7356-7360.

Grove EA, Williams BP, Li D-Q, Hajihosseini M, Friedrich A, Price J (1993) Multiple restricted lineages in the embryonic rat cerebral cortex. Development 117:553-561.

Guillemot F (2007) Spatial and temporal specification of neural fates by transcription factor codes. Development 134:3771-3780.

Gupta A, Wang Y, Markram H (2000) Organizing principles for a diversity of GABAergic interneurons and synapses in the neocortex. *Science* 287:273-278.

Hamasaki T, Goto S, Nishikawa S, Ushio Y (2001) A role of netrin-1 in the formation of the subcortical structure striatum: repulsive action on the migration of late-born striatal neurons. *J. Neurosci* 21:4271-4280.

Hand R, Bortone D, Mattar P, Nguyen L, Heng JI, Guerrier S, Boutt E, Peters E, Barnes AP, Parras C, Schuurmans C, Guillemot F, Polleux F (2005) Neurogenin2 specifies the migration properties and the dendritic morphology of pyramidal neurons in the neocortex. *Neuron* 48:45-62.

Hansen DV, Rubenstein JLR, Kriegstein, AR (2011) Deriving excitatory neurons of the neocortex from pluripotent stem cells. *Neuron* 70:645-660.

Hashimoto T, Volk DW, Eggan SM, Mirnics K, Pierri JN, Sun Z, Sampson AR, Lewis DA (2003) Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *J Neurosci* 23:6315-6326.

Hashimoto T, Nguyen QL, Rotaru D, Keenan T, Arion D, Beneyto M, Gonzalez-Burgos G, Lewis DA (2009) Protracted developmental trajectories of GABA_A

receptor $\alpha 1$ and $\alpha 2$ subunit expression in primate prefrontal cortex. *Biol Psychiatry* 65:1015-1023.

Hasling TA, Gierdalski M, Jablonska B, Juliano SL (2003) A radialization factor in normal cortical plate restores disorganized radial glia and disrupted migration in a model of cortical dysplasia. *Eur J Neurosci* 17:467-480.

Hatten ME (2002) New directions in neuronal migration. *Science* 297:1660-1663.

Hattori M, Adachi H, Tsujimoto M, Arai H, Inoue K (1994) Miller-Dieker lissencephaly gene encodes a subunit of brain platelet activating factor acetylhydrolase. *Nature* 370:216-218.

Haubensak W, Attardo A, Denk W, Huttner WB (2004) Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proc Natl Acad Sci USA* 101: 3196-3201.

Heck N, Kilb W, Reiprich P, Kubota H, Furukawa T, Fukuda A, Luhmann HJ (2007) GABA_A receptors regulate neocortical neuronal migration in vitro and in vivo. *Cereb Cortex* 17:138-148.

Hendry SH, Schwark HD, Jones EG, Yan J (1987) Numbers and proportions of GABA-immunoreactive neurons in different areas of monkey cerebral cortex. *J Neurosci.* 7:1503-1519.

Hendry SHC, Jones EG, Emsin PC, Lawson DEM, Heizmann CW, Streit P (1989) Two classes of cortical GABA neurons defined by differential calcium binding protein immunoreactivities. *Exp Brain Res* 76:467-472.

Heng JI, Nguyen L, Castro DS, Zimmer C, Wildner H, Armant O, Skowronska-Krawczyk D, Bedogni F, Matter JM, Hevner R, Guillemot F (2008) Neurogenin 2 controls cortical neuron migration through regulation of *Rnd2*. *Nature* 455:114-118.

Hennou, S., Khalilov, I., Diabira, D., Ben-Ari, Y., Gozlan, H. (2002) Early sequential formation of functional GABA_A and glutamatergic synapses on CA1 interneurons of the rat fetal hippocampus. *Eur J Neurosci* 16:197-208.

Hensch TK (2005) Critical period plasticity in local cortical circuits. *Nat Rev Neurosci* 6:877-888.

Hevner RF, Daza RA, Rubenstein JL, Stunnenberg H, Olavarria JF, Englund C (2003) Beyond laminar fate: toward a molecular classification of cortical projection/pyramidal neurons. *Dev Neurosci* 25:139-151.

Hevner RF, Daza RA, Englund C, Kohtz J, Fink A (2004) Postnatal shifts of interneuron position in the neocortex of normal and reeler mice: evidence for inward radial migration. *Neuroscience* 124:605-618.

Hirotsune S, Takahara T, Sasaki N, Hirose K, Yoshiki A, Ohashi T, Kusakabe M, Murakami Y, Muramatsu M, Watanabe S, Nakao K, Katsuki M, Hayashizaki Y (1995) The reeler gene encodes a protein with an EGF-like motif expressed by pioneer neurons. *Nat Genet* 10:77-83.

Hirotsune S, Fleck MW, Gambello MJ, Bix GJ, Chen A, Clark GD, Ledbetter DH, McBain CJ, Wynshaw-Boris A (1998) Graded reduction of Pafah1b1 (Lis1) activity results in neuronal migration defects and early embryonic lethality. *Nat Genet* 19:333-339.

Holt CE, Bertsch TW, Ellis HM, Harris WA (1988). Cellular determination in the *Xenopus* retina is independent of lineage and birth date. *Neuron* 1:15-26.

Honda T, Tabata H, Nakajima K (2003) Cellular and molecular mechanisms of neuronal migration in neocortical development. *Cell and Dev Biol* 14:169-174.

Houser CR, Hendry SH, Jones EG, Vaughn JE (1983) Morphological diversity of immunocytochemically identified GABA neurons in the monkey sensory-motor cortex. *J. Neurocytol.* 12:617-638.

Howell BW, Herrick TM, Cooper JA (1999) Neuronal position in the developing brain is regulated by mouse disabled-1. *Nature* 389:733-737.

Inada H, Watanabe M, Uchida T, Ishibashi H, Wake H, Nemoto T, Yanagawa Y, Fukuda A, Nabekura J (2011) GABA regulates the multidirectional tangential migration of GABAergic interneurons in living neonatal mice. *PLoS One* 6(12):1-11.

Jablonska B, Smith AL, Palmer SL, Noctor SC, Juliano SL (2004) GABA_A receptors reorganize when layer 4 in ferret somatosensory cortex is disrupted by methylazoxymethanol (MAM). *Cereb Cortex* 14(4):432-440.

Jacobs KM, Gutnick MJ, Price DA (1996) Hyperexcitability in a model of cortical maldevelopment. *Cereb Cortex* 6:514-523.

Jarman AP, Grau Y, Jan LY, Jan YN (1993) *atonal* is a proneural gene that directs chordotonal organ formation in the *Drosophila* peripheral nervous system. *Cell* 73:1307-1321.

Jimenez DL, Opez-Mascaraque L, Valverde F, De Carlos JA (2002) Tangential migration in neocortical development. *Developmental Biology* 244:155-169.

Jones EG (1975) Varieties and distribution of non-pyramidal cells in the somatic sensory cortex of the squirrel monkey. *J Comp Neuro* 160:205-267.

Jones EG (1986) Neurotransmitters in the cerebral cortex. *J Neurosurg* 65:135-153.

Kang J-Q, Barnes G (2012) A Common Susceptibility Factor of Both Autism and Epilepsy: Functional deficiency of GABAA Receptors. *J Autism Dev Disord*.

Kanatani S, Yozu M, Tabata H, Nakajima K (2008) COUPTFII is preferentially expressed in the caudal ganglionic eminence and is involved in the caudal migratory stream. *J Neurosci* 28:13582-13591.

Kappeler C, Saillour Y, Baudoin JP, Tuy FP, Alvarez C, Houbroun C, Gaspar P, Hamard G, Chelly J, Métin C, Francis F (2006) Branching and nucleokinesis defects in migrating interneurons derived from doublecortin knockout mice. *Hum Mol Genet* 15:1387-1400.

Kawaguchi Y, Kubota Y (1997) GABAergic cell subtypes and their synaptic connections in rat frontal cortex. *Cereb Cortex* 7:476-486.

Keverne EB (1999) GABA-ergic neurons and the neurobiology of schizophrenia and other psychoses. *Brain Res Bull* 48:467-473.

Kimberle MJ, Bryan JH, David AP (1999). Focal epileptogenesis in a rat model of polymicrogyria. *J Neurophysiol* 81:159-173.

Kitamura K, Yanazawa M, Sugiyama N, Miura H, Iizuka-Kogo A, Kusaka M, Omichi K, Suzuki R, Kato-Fukui Y, Kamiirisa K, Matsuo M, Kamiyo S, Kasahara M, Yoshioka H, Ogata T, Fukuda T, Kondo I, Kato M, Dobyns WB, Yokoyama M, Morohashi K (2002) Mutation of ARX causes abnormal development of forebrain and testes in mice and X-linked lissencephaly with abnormal genitalia in humans. *Nat Genet* 32:359-369.

Kobayashi M, Buckmaster PS (2003) Reduced inhibition of dentate granule cells in a model of temporal lobe epilepsy. *J Neurosci* 23:2440-2452.

Kohwi M, Petryniak MA, Long JE, Ekker M, Obata K, Yanagawa Y, Rubenstein JL, Alvarez-Buylla A (2007) A subpopulation of olfactory bulb GABAergic interneurons is derived from Emx1- and Dlx5/6-expressing progenitors. *J Neurosci* 27:6878-6891.

Komatsu S, Sakata-Haga H, Sawada K., Hisano, S., Fukui, Y. (2001) Prenatal exposure to ethanol induces leptomeningeal heterotopia in the cerebral cortex of the rat fetus. *Acta Neuropathol* 101:22-26.

Komuro H, Rakic P (1992) Selective Role of N-Type calcium channels in neuronal migration. *Science* 257: 806-809.

Komuro H, Rakic P (1993) Modulation of neuronal migration by NMDA receptors. *Science* 260:95-97.

Komuro H, Rakic P (1995) Dynamics of granule cell migration: a confocal microscopic study in acute cerebellar slice preparations. *J Neurosci.* 15(2):1110-1120.

Kornblum, H. I., Hussain, R. J., Bronstein, J. M., Gall, C. M., Lee, D. C., Seroogy, K. B. (1997). Prenatal ontogeny of the epidermal growth factor receptor and its ligand, transforming growth factor alpha, in the rat brain. *J Comp Neurol* 380:243-246.

Kornack DR, Rakic P (1995) Radial and horizontal deployment of clonally related cells in the primate neocortex: relationship to distinct mitotic lineages. *Neuron* 15, 311-321.

Koster SE, O'Leary DDM (1993) Axons of early generated neurons in cingulate cortex pioneer the corpus callosum. *J Neurosci* 14(11):6608-6620.

Kriegstein AR, Götz M (2003) Radial glia diversity: a matter of cell fate. *Glia* 43:37-43.

Kriegstein AR, Noctor SC (2004) Patterns of neuronal migration in the embryonic cortex. *Trends Neurosci* 27(7):4392-399.

Kriegstein A, Noctor S, Martinez-Cerdeno (2006) Patterns of neural stem and progenitor cell division may underlie evolutionary cortical expansion. *Nat Rev Neurosci* 7:883-890.

Kriegstein A, Alvarez-Buylla A (2009) The glial nature of embryonic and adult neural stem cells. *Annu Rev Neurosci* 32:149-184.

Kristin LW, Paul D, Anirvan G (2002) A molecular control of cortical dendrite development. *Annu Rev Neurosci* 25:127-49.

Kwon YT, Tsai LH (1998) A novel disruption of cortical development in p35 (-/-) mice distinct from reeler. *J Comp Neurol* 395:510-522.

Lancot KL, Herrmann N, Mazzotta P, Khan LR, Ingber N (2004) GABAergic function in Alzheimer's disease: Evidence for dysfunction and potential as a therapeutic target for the treatment of behavioural and psychological symptoms of dementia. *Can J Psychiatry* 49:439-453.

Laurie DJ, Wisden W, Seeburg PH (1992) The distribution of thirteen GABA_A receptor subunit mRNAs in the rat brain. III. embryonic and postnatal development. *J Neurosci* 12(11):4151-4172.

Lavdas AA, Grigoriou M, Pachnis V, Parnavelas JG (1999) The medial ganglionic eminence gives rise to a population of early neurons in the developing cerebral cortex. *J Neurosci* 19:7881-7888.

Le TN, Du G, Fonseca M, Zhou QP, Wigle JT, Eisenstat DD (2007) Dlx homeobox genes promote cortical interneuron migration from the basal forebrain by direct repression of the semaphorin receptor neuropilin-2. *J Biol Chem* 282:19071-19081.

Lee KS, Schottler F, Collins JL, Lanzino G, Couture D, Rao A, Hiramatsu KI, Goto Y, Hong SC, Caner H, Yamamoto H, Chen ZF, Bertram E, Berr S, Omary R, Scrable H, Jackson T, Goble J, Eisenman L (1997) A genetic animal model of human neocortical heterotopia associated with seizures. *J Neurosci* 17:6236-6242.

Leone DP, Srinivasan K, Chen B, Alcamo E, McConnell SK (2008) The determination of projection neuron identity in the developing cerebral cortex. *Curr Opin Neurobiol* 18:28-35.

Lerner JT, Salamon N, Hauptman JS, Velasco TR, Hemb M, Wu JY, Sankar R, Donald Shields W, Engel J Jr, Fried I, Cepeda C, Andre VM, Levine MS, Miyata H, Yong WH, Vinters HV, Mathern GW (2009) Assessment and surgical outcomes for mild type I and severe type II cortical dysplasia: a critical review and the UCLA experience. *Epilepsia* 50:1310-1335.

Letinic K, Zoncu R, Rakic P (2002) Origin of GABAergic neurons in the human neocortex. *Nature* 417:645-649.

Lewis DA, Cruz D, Eggan S, Erickson S (2004) Postnatal development of prefrontal inhibitory circuits and the pathophysiology of cognitive dysfunction in schizophrenia. *Ann N Y Acad Sci* 1021:64-76.

Lewis DA, Hashimoto T, Volk DW (2005) Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 6:312-324.

Li LM, Fish DR, Sisodiya SM, Shorvon SD, Alsanjari N, Stevens JM (1995) High resolution magnetic resonance imaging in adults with partial or secondary

generalized epilepsy attending a tertiary referral unit. *J Neurol Neurosurg Psychiatry* 59:384-387.

Li H, Tornberg J, Kaila K, Airaksinen MS, Rivera C (2002) Patterns of cation-chloride cotransporter expression during embryonic rodent CNS development. *Eur J Neurosci* 16:2358-2370.

Li G, Adesnik H, Li J, Long J, Nicoll RA, Rubenstein JL, Pleasure SJ (2008) Regional distribution of cortical interneurons and development of inhibitory tone are regulated by Cxcl12/Cxcr4 signaling. *J Neurosci* 28:1085-1098.

Li G, Pleasure SJ (2011) Exciting information for inhibitory neurons. *Neuron* 69(4): 585-587.

Limon A, Reyes-Ruiz JM, Miledi R (2012) Loss of functional GABAA receptors in the Alzheimer diseased brain. *Proc Natl Acad Sci USA* 109(25):10071-10076.

Liodis P, Denaxa M, Grigoriou M, Akufo-Addo C, Yanagawa Y, Pachnis V (2007) Lhx6 activity is required for the normal migration and specification of cortical interneuron subtypes. *J Neurosci* 27:3078-3089.

Lodato S, Tomassy GS, De Leonibus E, Uzcategui YG, Andolfi G, Armentano M, Touzot A, Gatelu, JM, Arlotta P, Menendez de la Prida L, Studer M (2011) Loss

of COUP-TFI alters the balance between caudal ganglionic eminence- and medial ganglionic eminence-derived cortical interneurons and results in resistance to epilepsy. *J Neurosci* 31:4650-4662.

López-Bendito G, Luján R, Shigemoto R, Ganter P, Paulsen O, Molnar Z (2003) Blockade of GABA_B receptors alters the tangential migration of cortical neurons. *Cereb Cortex* 13:932-942.

López-Bendito G, Sturgess K, Erdelyi F, Szabo G, Molnar Z, Paulsen O (2004) Preferential origin and layer destination of GAD65-GFP cortical interneurons. *Cereb Cortex* 14:1122-1133.

López-Bendito G, Sanchez-Alcaniz JA, Pla R, Borrell V, Pico E, Valdeolmillos M, Marin O (2008) Chemokine signaling controls intracortical migration and final distribution of GABAergic interneurons. *J Neurosci* 28:1613-1624.

LoTurco JJ, Owens DF, Heath MJ, Davis MB, Kriegstein AR (1995) GABA and glutamate depolarize cortical progenitor cells and inhibit DNA synthesis. *Neuron* 15:1287-1298.

Loup F, Wieser HG, Yonekawa Y, Aguzzi A, Fritschy JM (2000) Selective alterations in GABAA receptor subtypes in human temporal lobe epilepsy. *J Neurosci* 20:5401-5419.

Luhmann HJ, Mittmann T, van Luijtelaar G, Heinemann U (1995) Impairment of intracortical GABAergic inhibition in a rat model of absence epilepsy. *Epilepsy Res* 22:43-51.

Lund JS, Lewis DA (1993) Local circuit neurons of developing and mature macaque prefrontal cortex: Golgi and immunocytochemical characteristics. *J Comp Neurol* 328:282-312.

Luskin MB, Pearlman AL, Sanes JR (1988) Cell lineage in the cerebral cortex of the mouse studied *in-vivo* and *in-vitro* with a recombinant retrovirus. *Neuron* 1: 635-647.

Ma Y, Prince DA (2012) Functional alterations in GABAergic fast-spiking interneurons in chronically injured epileptogenic neocortex. *Neurobiol Dis* 47:102-113.

Malatesta P, Hartfuss E, Gotz M (2000) Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development* 127:5253-63.

Manent JB, Jorquera I, Ben-Ari Y, Aniksztejn L, Represa A (2006) Glutamate acting on AMPA but not NMDA receptors modulates the migration of hippocampal interneurons. *J Neurosci* 26:5901-5909.

Marín O, Anderson SA, Rubenstein JL (2000) Origin and molecular specification of striatal interneurons. *J Neurosci* 20:6063-6076.

Marín O, Rubenstein JLR (2001) A long, remarkable journey: tangential migration in the telencephalon. *Nat Rev Neurosci* 2:780-790.

Marín O, Yaron A, Bagri A, Tessier-Lavigne M, Rubenstein JLR (2001) Sorting of striatal and cortical interneurons regulated by semaphorin/neuropilin interactions. *Science* 293:872-75.

Marín O, Andrew S, Plump AS, Flames N, Sánchez-Camacho C, Tessier-Lavigne M, John LRR (2003) Directional guidance of interneuron migration to the cerebral cortex relies on subcortical Slit1/2-independent repulsion and cortical attraction. *Development* 130:1889-1901.

Marín O, Rubenstein JLR (2003) Cell migration in the forebrain. *Ann Rev Neurosci* 26: 441-483.

Marin-Padilla, M (1998) Cajal-Retzius cells and the development of the neocortex. *Trends Neurosci* 21:64-71.

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C (2008) Interneurons of the neocortical inhibitory system. *Nat Rev Neurosci* 5(10):793-807.

Martínez-Cerdeño V, Cunningham CL, Camacho J, Antczak JL, Prakash AN, Cziep ME, Walker AI, Noctor SC (2012) Comparative Analysis of the Subventricular Zone in Rat, Ferret and Macaque: Evidence for an Outer Subventricular Zone in Rodents. *PLoS One* 7(1):1-36.

Martini FJ, Valiente M, López-Bendito G, Szabó G, Moya F, Valdeolmillos M, Marín O (2009) Biased selection of leading process branches mediates chemotaxis during tangential neuronal migration. *Development* 136: 41-50.

Mason HA, Ito S, Corfas G (2001) Extracellular signals that regulate the tangential migration of olfactory bulb neuronal precursors: inducers, inhibitors, and repellants. *J Neurosci* 21:7654-7663.

Matsumoto H, Higa HH (1966) Studies on methylazoxymethanol, the aglycone of cycasin: methylation of nucleic acids in vitro. *Biochem J* 98:20-22C.

McConnell SK (1995) Constructing the cerebral cortex: neurogenesis and fate determination. *Neuron* 15:761-768.

McEvelly RJ, de Diaz MO, Schonemann MD, Hooshmand F, Rosenfeld MG (2002) Transcriptional regulation of cortical neuron migration by POU domain factors. *Science* 295:1528-1532.

McKernan RM, Quirk K, Prince R, Cox PA, Gillard NP, Ragan CL, Whiting P (1991) GABA_A receptor subtypes immunopurified from rat brain with a subunit-specific antibodies have unique pharmacological properties. *Neuron* 7:667-76

McLaughlin DF, Juliano S (2005) Disruption of layer 4 development alters laminar processing in ferret somatosensory cortex. *Cereb Cortex* 15:1791-1803.

Meinecke DL, Peters A (1987) GABA immunoreactive neurons in rat visual cortex. *J Comp Neuro* 261:388-404.

Me'tin C, Denizot J-P, Ropert N (2000) Intermediate zone cells express calcium-permeable AMPA receptors and establish close contact with growing axons. *J Neurosci* 20(2):696-708.

Miller MW (1985) Cogeneration of retrogradely labeled corticocortical projection and GABA-immunoreactive local circuit neurons in cerebral cortex. *Brain Res* 355:187-192.

Mione MC, Cavanagh JF, Harris B, Parnavelas JG (1997) Cell fate specification and symmetrical/asymmetrical divisions in the developing cerebral cortex. *J Neurosci* 17:2018-2029.

Miyata T, Kawaguchi A, Okano H, Ogawa M (2001) Asymmetric inheritance of radial glial fibers by cortical neurons. *Neuron* 31:727-741.

Miyoshi G, Butt SJ, Takebayashi H, Fishell G (2007) Physiologically distinct temporal cohorts of cortical interneurons arise from telencephalic Olig2-expressing precursors. *J Neurosci* 27:7786-7798.

Miyoshi G, Fishell G (2011) GABAergic Interneuron lineages selectively sort into specific cortical layers during early postnatal development. *Cereb Cortex* 21:845-852.

Molyneaux BJ, Arlotta P, Hirata T, Hibi M, Macklis JD (2005) Fezl is required for the birth and specification of corticospinal motor neurons. *Neuron* 47:817-831.

Molyneaux BJ, Arlotta P, Menezes JR, Macklis JD (2007). Neuronal subtypes specification in the cerebral cortex. *Nat Rev Neurosci* 8:427-437.

Morest DK (1970) A study of neurogenesis in the forebrain of opossum pouch young. *Z Anat Entwicklungsgesch* 130:265-305.

Moroni RF, Inverardi F, Regondi MC, Panzica F, Spreafico R, Frassoni C. (2008) Altered spatial distribution of PV-cortical cells and dysmorphic neurons in the somatosensory cortex of BCNU-treated rat model of cortical dysplasia. *Epilepsia* 49(5):872-887.

Moyer CE, Delevich KM, Fish KN, Asafu-Adjei JK, Simpson AR, Dorph-Petersen K-A, Lewis DA, Sweet RA (2012) Reduced glutamate decarboxylase 65 protein within the primary auditory cortex inhibitory boutons in schizophrenia. *Biol Psychiatry*

Nadarajah B, Brunstrom JE, Grutzendler J, Wong RO, Pearlman AL (2001) Two modes of radial migration in early development of the cerebral cortex. *Nat Neurosci* 4:143-150.

Nadarajah B, Parnavelas JG (2002) Modes of neuronal migration in the developing cerebral cortex. *Nat Rev Neurosci* 3:423-32.

Nakagawa T, Sasahara M, Hayase Y, Haneda M, Yasuda H, Kikkawa R, Higashiyama S, Hazama F (1998) Neuronal and glial expression of heparin-binding EGF-like growth factor in central nervous system of prenatal and early-postnatal rat. *Brain Res Dev Brain Res* 108:263-272.

Nakamura K, Yamashita Y, Tamamaki N, Katoh H, Kaneko T, Negishi M (2006) In vivo function of Rnd2 in the development of neocortical pyramidal neurons. *Neurosci Res* 54:149-153.

Natalia V, Garcia De Marco, Karayannis T, Fishell G (2011) Neuronal activity is required for the development of specific cortical interneuron subtypes. *Nature* 472:351-356.

Nery S, Fishell G, Corbin JG (2002) The caudal ganglionic eminence is a source of distinct cortical and subcortical cell populations. *Nat Neurosci* 5:1279-1287.

Nguyen L, Besson A, Heng JI, Schuurmans C, Teboul L, Parras C, Philpott A, Roberts JM, Guillemot F (2006) p27^{kip1} independently promotes neuronal differentiation and migration in the cerebral cortex. *Genes Dev* 20:1511-1524.

Nóbrega-Pereira S, Kessar N, Du T, Kimura S, Anderson SA, Marín O (2008) Postmitotic Nkx2-1 controls the migration of telencephalic interneurons by direct repression of guidance receptors. *Neuron* 59:733-45.

Nóbrega-Pereira S, Marín O (2009). Transcriptional control of interneuron migration in the developing mouse brain. *Cereb Cortex* 19:107-113.

Noctor SC, Palmer SL, Hasling T, Juliano SL (1999) Interference with the development of early generated neocortex results in disruption of radial glia and abnormal formation of neocortical layers. *Cereb Cortex* 9:121-136.

Noctor SC, Flint AC, Weissman TA, Dammerman RS, Kriegstein AR (2001a) Neurons derived from radial glial cells establish radial units in neocortex. *Nature* 409:714-720.

Noctor SC, Palmer SL, McLaughlin DF, Juliano SL (2001b) Disruption of layers 3 and 4 during development results in altered thalamocortical projections in ferret somatosensory cortex. *J Neurosci* 21:3184-3195.

Noctor SC, Martinez-Cerdeno V, Ivic L, Kriegstein AR (2004) Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nat Neurosci* 7:136-144.

Oblak L, Gibbs TT, Blatt GJ (2010) Decreased GABA- B receptors in the cingulate cortex and fusiform gyrus in Autism. *J Neurochem* 114:1414-1423.

Ogawa M, Miyata T, Nakajima K, Yagyu K, Seike M, Ikenaka K, Yamamoto H, Mikoshiba, K (1995) The reeler gene-associated antigen on Cajal-Retzius neurons is a crucial molecule for laminar organization of cortical neurons. *Neuron* 14:899-912.

Ohnuma S, Harris WA (2003) Neurogenesis and the cell cycle. *Neuron* 40:199-208.

Ohshima T, Ward JM, Huh CG, Longenecker G, Veeranna, Pant HC, Brady RO, Martin LJ, Kulkarni AB (1996) Targeted disruption of the cyclin-dependent kinase 5 gene results in abnormal corticogenesis, neuronal pathology and perinatal death. *Proc Natl Acad Sci USA* 93(11):173-178.

O'Leary DDM, Koester SE (1993) Development of projection neuron types, axon pathways, and patterned connections of the mammalian cortex. *Neuron* 10:991-1006.

Olsen RW, Sieghart W (2008) International Union of Pharmacology. LXX. Subtypes of gamma-aminobutyric acid(A) receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharm Rev* 60(3):243-260.

O'Rourke NA, Dailey ME, Smith SJ, McConnell SK (1992) Diverse migratory pathways in the developing cerebral cortex. *Science* 258:299-302.

O'Rourke NA, Sullivan DP, Kaznowski CE, Jacobs AA, McConnell SK (1995) Tangential migration of neurons in the developing cerebral cortex. *Development* 121:2165-2176.

Owens DF, Liu X, Kriegstein AR (1999) Changing properties of GABA(A) receptor-mediated signaling during early neocortical development. *J Neurophysiol* 82:570-583.

Owens DF, Kriegstein AR (2002) Is there more to GABA than synaptic inhibition? *Nat Rev Neurosci* 3:715-727.

Palmer SL, Noctor SC, Jablonska B, Juliano SL (2001) Laminar specific alterations of thalamocortical projections in organotypic cultures following layer 4 disruption in ferret somatosensory cortex. *Eur J Neurosci* 13:1559-1571.

Palmini A, Andermann F, Olivier A, Tampieri D, Robitaille Y, Andermann E, Wright G (1991) Focal neuronal migration disorders and intractable partial epilepsy: a study of 30 patients. *Ann Neurol* 30:741-749.

Parnavelas JG, Barfield JA, Franke E, Luskin MB (1991) Separate progenitor cells give rise to pyramidal and nonpyramidal neurons in the rat telencephalon. *Cereb Cortex* 1:463-468.

Parnavelas JG (2000) The origin and migration of cortical neurones: new vistas. *Trends Neurosci* 23:126-131.

Payne JA (1997) Functional characterization of the neuronal-specific K-Cl cotransporter: implications for $[K^+]_o$ regulation. *Am J Physiol* 273: C1516-C1525.

Peduzzi JD (1988) Genesis of GABA-immunoreactive neurons in the ferret visual cortex. *J Neurosci* 8:920-931.

Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G (2000). GABA_A receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 101(4):815-850.

Pleasure SJ, Anderson S, Hevner R, Bagri A, Marin O, Lowenstein DL, Rubenstein JLR (2000) Cell migration from the ganglionic eminences is required for the development of hippocampal GABAergic interneurons. *Neuron* 28:727-740.

Plotkin MD, Snyder EY, Hebert SC, and Delpire E (1997) Expression of the Na-K-2Cl cotransporter is developmentally regulated in postnatal rat brains: a possible mechanism underlying GABA's excitatory role in immature brain. *J Neurobiol* 33: 781-795.

Polkey CE (1996) Cortical dysplasia: resective surgery in children. In: Guerrini, R., Andermann, F., Canapicchi, R., Roger, J. Zifkin, B. G., Pfanner, P., editors. *Dysplasias of cerebral cortex and epilepsy*. Philadelphia: Lippincott-Raven, 435-439.

Polleux F, Whitford KL, Dijkhuizen PA, Vitalis T, Ghosh A (2002) Control of cortical interneuron migration by neurotrophins and PI3-kinase signaling. *Development* 129:3147-3160.

Poluch S, Konig N (2002) AMPA receptor activation induces GABA release from neurons migrating tangentially in the intermediate zone of embryonic rat neocortex. *Eur J Neurosci* 16:350-354.

Poluch S, Juliano SL (2007) A normal radial glial scaffold is necessary for migration of interneurons during neocortical development. *Glia* 55(8):822-830.

Poluch S, Jablonska B, Juliano SL (2008) Alteration of interneuron migration in a ferret model of cortical dysplasia. *Cereb Cortex* 18(1):78-92.

Poluch and Juliano (2010) Populations of radial glial cells respond differently to reelin and neuregulin1 in a ferret model of cortical dysplasia. PLoS One 5(10):1-17

Poulter MO, Barker JL, O'Carroll AM, Lolait SJ, Mahan LC (1992) Differential and transient expression of GABA_A receptor α -subunit mRNAs in the developing rat CNS. J Neurosci 12(8):2888-2900.

Powell EM, Mars WM, Levitt P (2001) Hepatocyte growth factor/scatter factor is a motogen for interneurons migrating from the ventral to dorsal telencephalon. Neuron 30:79-89.

Pozas E, Ibanez CF (2005) GDNF and GFR α 1 promote differentiation and tangential migration of cortical GABAergic neurons. Neuron 45:701-713.

Price J, Thurlow L (1988) Cell lineage in the rat cerebral cortex: a study using retroviral-mediated gene transfer. Development 104: 473-482.

Qian X, Goderie SK, Shen Q, Stern JH, Temple S (1998) Intrinsic programs of patterned cell lineages in isolated vertebrate CNS ventricular zone cells. Development 125:3143-3152.

Qian X, Shen Q, Goderie SK, He W, Capela A, Davis AA, Temple S (2000) Timing of CNS cell generation: a programmed sequence of neuron and glial cell production from isolated murine cortical stem cells. *Neuron* 28:69-80.

Rakic P (1971) Guidance of neurons migrating to the fetal monkey neocortex. *Brain Res* 33:471-476.

Rakic P (1972) Mode of cell migration to the superficial layers of fetal monkey neocortex. *J Comp Neurol* 145(1):61-83.

Rakic P (1974) Neurons in rhesus monkey visual cortex: systematic relation between time of origin and eventual disposition. *Science* 183: 425-427.

Rakic P (1988) Specification of cerebral cortical areas. *Science* 241:170-176.

Rakic P (1995) A small step for the cell, a giant leap for mankind: a hypothesis of neocortical expansion during evolution. *Trends Neurosci* 18:383-388.

Reid CB, Tavazoie SF, Walsh CA (1997) Clonal dispersion and evidence for asymmetric cell division in ferret cortex. *Development* 124:2441-2450.

Reiner O, Carrozzo R, Shen Y, Wehnert M, Faustinella F, Dobyns WB, Caskey TC, Ledbetter DH (1993) Isolation of a Miller-Dieker lissencephaly gene containing G protein beta-subunit-like repeats. *Nature* 364:717-721.

Represa A, Ben-Ari Y (2005) Trophic actions of GABA on neuronal development. *Trends Neurosci* 28:278-283.

Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtinen H, Lamsa K, Pirvola U, Saarma M, Kaila K (1999) The K⁺/Cl⁻ co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* 397:251-255.

Roberts MR, Bittman K, Li WW, French R, Mitchell B, LoTurco JJ, D'Mello SR (2000) The flathead mutation causes CNS-specific developmental abnormalities and apoptosis. *J Neurosci* 20:2295-2306.

Roper SN (1998) In-utero irradiation of rats as a model of human cerebrocortical dysgenesis: a review. *Epilepsy Res* 32:63-74.

Roper SN, Eisenschenk S, King MA (1999). Reduced density of parvalbumin- and calbindin D28-immunoreactive neurons in experimental cortical dysplasia. *Epilepsy Res* 37:6371.

Rosen GD, Press DM, Sherman GF, Galaburda AM (1992) The development of induced cerebrocortical microgyria in the rat. *J Neuropathol Exp Neurol* 51:601-611.

Rosen GD, Jacobs KM, Prince DA (1998) Effects of neonatal freeze lesions on expression of parvalbumin in rat neocortex. *Cereb Cortex* 8:753-761.

Ross EM, Walsh CA (2001) Human brain malformations and their lessons for neuronal migration. *Annu Rev Neurosci* 24:1041-70.

Ross SE, Greenberg ME, Stiles CD (2003) Basic helix-loop-helix factors in cortical development. *Neuron* 39:13-25.

Rymar VV, Sadikot AF (2007) Laminar fate of cortical GABAergic interneurons is dependent on both birthdate and phenotype. *J Comp Neurol* 501:369-380.

Sanchez-Alcaniz JA, Haege S, Mueller W, Pla R, Mackay F, Schulz S, Lopez-Bendito G, Stumm R, Marin O (2011) *Cxcr7* controls neuronal migration by regulating chemokine responsiveness. *Neuron* 69:77-90

Schuermans C, Armant O, Nieto M, Stenman J, Britz O, Klenin N, Brown C, Langevin L-M, Seibt J, Tang H, Cunningham JM, Dyck R, Walsh C, Campbell K, Polleux F, Guillemot F (2004) Sequential phases of neocortical fate specification

involve neurogenin-dependent and -independent pathways. EMBO J 23:2892-2902.

Shao LR, Dudek FE (2005) Changes in mIPSCs and sIPSCs after kainate treatment: evidence for loss of inhibitory input to dentate granule cells and possible compensatory responses. J Neurophysiol 94:952-960.

Shen Q, Zhong W, Jan YN, Temple S (2002) Asymmetric Numb distribution is critical for asymmetric cell division of mouse cerebral cortical stem cells and neuroblasts. Development 129:4843-4853.

Shen Q, Wang Y, Dimos JT, Fasano CA, Phoenix TN, Lemischka IR, Ivanova NB, Stifani S, Morrissey EE, Temple S (2006) The timing of cortical neurogenesis is encoded within lineages of individual progenitor cells. Nat Neurosci 9:743-751.

Sheppard AM, Pearlman AL (1997) Abnormal reorganization of preplate neurons and their associated extracellular matrix: an early manifestation of altered neocortical development in the reeler mutant mouse. J Comp Neurol 378:173-179.

Shinozaki K, Miyagi T, Yoshida M, Miyata T, Ogawa M, Aizawa S, Suda Y (2002) Absence of Cajal-Retzius cells and subplate neurons associated with defects of

tangential cell migration from ganglionic eminence in *Emx1/2* double mutant cerebral cortex. *Development* 129:3479-3492

Somogyi P, Klausberger T (2005) Defined types of cortical interneuron structure space and spike timing in the hippocampus. *J Physiol* 562:9-26.

Soria JM, Valdeolmillos M (2002) Receptor-activated calcium signals in tangentially migrating cortical cells. *Cereb Cortex* 12:831-839.

Stumm RK, Zhou C, Ara T, Lazarini F, Dubois-Dalcq M, Nagasawa T, Holtt, V, Schulz S (2003) CXCR4 regulates interneuron migration in the developing neocortex. *J Neurosci* 23:5123-5130.

Sugitani Y, Nakai S, Minowa O, Nishi M, Jishage K, Kawano H, Mori K, Ogawa M, Noda T (2002) Brn-1 and Brn-2 share crucial roles in the production and positioning of mouse neocortical neurons. *Genes Dev* 16:1760-1765.

Sussel L, Marín O, Kimura S, Rubenstein JL (1999) Loss of *Nkx2.1* homeobox gene function results in a ventral to dorsal molecular re-specification within the basal telencephalon: evidence for a transformation of the pallidum into the striatum. *Development* 126:3359-3370.

Takahashi T, Goto T, Miyama S, Nowakowski RS, Caviness VSJr (1999) Sequence of neuron origin and neocortical laminar fate: relation to cell cycle of origin in the developing murine cerebral wall. *J Neurosci* 19:10357-10371.

Talos DM, Sun H, Kosaras B, Joseph A, Folkerth RD, Poduri A, Madsen J, Black PM, Jensen FE (2012) Altered Inhibition in Tuberous Sclerosis and Type IIb Cortical Dysplasia. *Ann Neurol* 71:539-551.

Tamamaki N, Fujimori KE, Takauji R (1997) Origin and route of tangentially migrating neurons in the developing neocortical intermediate zone. *J Neurosci* 17:8313-8323.

Tamamaki N, Yanagawa Y, Tomioka R, Miyazaki J, Obata K, Kaneko T (2003) Green fluorescent protein expression and colocalization with calretinin, parvalbumin, and somatostatin in the GAD67-GFP knock-in mouse. *J Comp Neurol* 467:60-79.

Tamas G, Buhl EH, Lorincz A, Somogyi P (2000) Proximally targeted GABAergic synapses and gap junctions synchronize cortical interneurons. *Nat Neurosci* 3:366-371.

Tanaka D, Nakaya Y, Yanagawa Y, Obata K, Murakami F (2003) Multimodal tangential migration of neocortical GABAergic neurons independent of GPI-anchored proteins. *Development* 130:5803-5813.

Tanaka D, Maekawa K, Yanagawa Y, Obata K, Murakami F (2006) multidirectional and multizonal tangential migration of GABAergic interneurons in the developing cerebral cortex. *Development* 133:2167-2176.

Tassi L, Colombo N, Garbelli R, Francione S, Lo Russo G, Mai R, Cossu M, Ferrario A, Galli C, Brammerio M, Citterio A, Spreafico R (2002) Focal cortical dysplasia: neuropathological subtypes, EEG, neuroimaging and surgical outcome. *Brain* 125:1719-1732.

Taylor DC, Falconer MA, Bruton CJ, Corsellis JAN (1971) Focal dysplasia of the cerebral cortex in epilepsy. *J Neurol Neurosurg Psychiatry* 34:369-387.

Thomson AM, West DC, Hahn J, Deuchars J (1996) Single axon IPSPs elicited in pyramidal cells by three classes of interneurons in slices of rat neocortex. *J Physiol (Lond)* 496:81-102

Tija CJ, Stephen JM, Rachel J (2008) GABA_A receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat Rev Neuroscience* 9:331-343.

Tiveron MC, Rossel M, Moepps B, Zhang YL, Seidenfaden R, Favor J, Konig N, Cremer H (2006) Molecular interaction between projection neuron precursors and invading interneurons via stromal-derived factor 1 (CXCL12)/CXCR4 signaling in the cortical subventricular zone/intermediate zone. J Neurosci 26:13273-13278.

Tyzio R, Represa A, Jorquera I, Ben-Ari Y, Gozlan H, Aniksztejn L (1999) The establishment of GABAergic and glutamatergic synapses on CA1 pyramidal neurons is sequential and correlates with the development of the apical dendrite. J Neurosci 19:10372-10382.

Uhlmann EJ, Wong M, Baldwin RL, Bajenaru ML, Onda H, Kwiatkowski DJ, Yamada K, Gutmann DH (2002) Astrocyte-specific *TSC1* conditional knockout mice exhibit abnormal neuronal organization and seizures. Ann Neurol 52:285-296.

Van Eden CG, Mrzljak L, Voorn P, Uylings HB (1989) Prenatal development of GABAergic neurons in the neocortex of the rat. J Comp Neurol 289:213-227.

Villares R, Cabrera CV (1987) The *achaete-scute* gene complex of *D. melanogaster*: conserved domains in a subset of genes required for neurogenesis and their homology to *myc*. Cell 50:415-424.

Walsh C, Cepko CL (1988) Clonally related cortical cells show several migration patterns. *Science* 241:1342-1345.

Walsh C, Cepko CL (1993) Clonal dispersion in proliferative layers of the developing cerebral cortex. *Nature* 362: 632-635.

Wang DD, Kriegstein AR (2008) GABA Regulates Excitatory Synapse formation in the Neocortex via NMDA Receptor Activation. *J Neurosci* 28(21):5547-5558.

Wang DD, Kriegstein AR (2009) Defining the role of GABA in cortical development. *J Neurophys* 587(9):1873-1879.

Wang Y, Li G, Stanco A, Long JE, Crawford D, Potter GB, Pleasure SJ, Behrens T, Rubenstein JLR (2011) CXCR4 and CXCR7 have distinct functions in regulating interneuron migration. *Neuron* 69:61-76.

Welagen J, Anderson S (2011) Origins of neocortical interneurons in mice. *Dev Neurobiol* 1:10-17.

Wichterle H, Garcia-Verdugo JM, Herrera DG, Alvarez-Buylla A (1999) Young neurons from medial ganglionic eminence disperse in adult and embryonic brain. *Nat Neurosci* 2:461-466.

Wichterle H, Turnbull DH, Nery S, Fishell G, Alvarez-Buylla A (2001) In utero fate mapping reveals distinct migratory pathways and fates of neurons born in the mammalian basal forebrain. *Development* 128:3759-3771.

Wichterle H, Alvarez-Dolado M, Erskine L, Alvarez-Buylla A (2003) Permissive corridor and diffusible gradients direct medial ganglionic eminence cell migration to the neocortex. *Proc Natl Acad Sci USA* 100:727-732.

Wonders CP, Anderson SA (2006) The origin and specification of cortical interneurons. *Nat Rev Neurosci* 7:687-696.

Wonders CP, Taylor L, Welagen J, Mbata IC, Xiang JZ, Anderson SA (2008) A spatial bias for the origins of interneuron subgroups within the medial ganglionic eminence. *Dev Biol* 314:127-136.

Xie X, Liaw JS, Baudry M, Berger TW (1997) Novel expression mechanism for synaptic potentiation: alignment of presynaptic release site and postsynaptic receptor. *Proc Natl Acad Sci USA* 94:6983-6988.

Xu Q, Cobos I, De La Cruz E, Rubenstein JL, Anderson SA (2004) Origins of cortical interneuron subtypes. *J Neurosci* 24:2612-2622.

Yeung RS, Katsetos CD, Klein-Szanto A (1997) Subependymal astrocytic hamartomas in the Eker rat model of tubero sclerosis. *Am J Pathol* 151:1477.

Yokota Y, Gashghaei HT, Han C, Watson H, Campbell KJ, Anton ES (2007) Radial glial dependent and independent dynamics of interneuronal migration in the developing cerebral cortex. *PLoS One* 2(8):1-10.

Yoshida M, Assimacopoulos S, Jones KR, Grove EA (2006) Massive loss of Cajal-Retzius cells does not disrupt neocortical layer order. *Development* 133(3):537-545.

Young AB (1987) Cortical amino acidergic pathways in Alzheimer's disease. *J Neural Transm Suppl* 24:147-152.

Young KM, Fogarty M, Kessar N, Richardson WD (2007) Subventricular zone stem cells are heterogeneous with respect to their embryonic origins and neurogenic fates in the adult olfactory bulb. *J Neurosci* 27:8286-8296.

Yu X, Jakovcevski I, Mo Z, Zeevic N (2009) Cortical origin of calretinin-expressing neurons in the human fetal forebrain. *Soc Neurosci Abstract* 39:807.12/A12.

Yu X, Zeevic N (2011). Dorsal radial glial cells have the potential to generate cortical interneurons in human but not in mouse brain. *J Neurosci* 31:2413-2420.

Yu Z-Y, Wang W, Fritschy J-M, Witte OW, Redecker C (2006) Changes in neocortical and hippocampal GABAA receptor subunit distribution during brain maturation and aging. *Brain Res* 1099:73-81.

Yun K, Potter S, Rubenstein JL (2001) *Gsh2* and *Pax6* play complementary roles in dorsoventral patterning of the mammalian telencephalon. *Development* 128:193-205.

Yuste R (2005) Origin and classification of cortical interneurons. *Neuron* 48(4):524-527.

Zarbalis K, Choe Y, Siegenthaler JA, Orosco LA, Pleasure SJ (2012) Meningeal defects alter the tangential migration of cortical interneurons in *Foxc1^{hith/hith}* mice. *Neural Dev* 7:2.

Zhu WJ, Roper, SN (2000) Reduced Inhibition in an Animal Model of Cortical Dysplasia. *J Neurosci* 20(23):8925-8931.

Zhu Y, Li H, Zhou L, Wu JY, Rao Y (1999) Cellular and molecular guidance of GABAergic neuronal migration from an extracortical origin to the neocortex. *Neuron* 23:473-485.