

AWARD NUMBER: W81XWH-15-1-0420

TITLE: Cellular Energy Pathways as Novel Targets for the Therapy of Autosomal Dominant Polycystic Kidney Disease

PRINCIPAL INVESTIGATOR: Kenneth R. Hallows, MD, PhD, FASN

CONTRACTING ORGANIZATION: University of Southern California
Los Angeles, CA 90089-0701

REPORT DATE: September 2016

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE September 2016		2. REPORT TYPE Annual		3. DATES COVERED 1 Sep 2015 - 31 Aug 2016	
4. TITLE AND SUBTITLE Cellular Energy Pathways as Novel Targets for the Therapy of Autosomal Dominant Polycystic Kidney Disease				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-15-1-0420	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Kenneth R. Hallows, MD, PhD Michael J. Caplan, PhD, MD E-Mail: hallows@usc.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Southern California Los Angeles, CA 90089-0701 Yale University New Haven, CT 06520				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Autosomal dominant polycystic kidney disease (ADPKD) is a common inherited disorder where patients, over the course of decades, develop large fluid filled cysts that damage the normal kidney tissue and can lead ultimately to kidney failure that necessitates transplantation or dialysis. There are currently no FDA-approved medications for this condition. Recent research reveals that the formation of cysts is due in part both to inappropriate cell growth, fluid secretion, and dysregulation of cellular energy metabolism. The enzyme AMPK regulates a number of cellular pathways, including these disease-causing features. Drugs that activate AMPK, therefore, may constitute an effective therapeutic option for slowing or preventing cyst growth in ADPKD. This research project is aimed at examining the potential of approved, widely used, inexpensive and low-toxicity drugs that can activate AMPK (metformin, simvastatin, and salicylates) and or promote oxidative metabolism (dichloroacetic acid) as potential therapies for the treatment of ADPKD. During this past research period, we optimized the measurement and analysis of various metabolites and metabolic enzymes (metabolomic biomarkers) in samples derived from cell lysates and urine and collected preliminary data in ADPKD mouse in vitro cell culture models and urine specimens derived from a cross-section of patients with ADPKD.					
15. SUBJECT TERMS Autosomal Dominant Polycystic Kidney Disease, Polycystins, Metformin, Statins, Salicylates, Dichloroacetic acid, AMP-activated Protein Kinase, Biomarkers, Metabolism, Mass Spectrometry					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area code)
Unclassified	Unclassified	Unclassified	Unclassified	20	

Table of Contents

	<u>Page</u>
1. Introduction.....	4
2. Keywords.....	5
3. Accomplishments.....	6
4. Impact.....	14
5. Changes/Problems.....	15
6. Products.....	16
7. Participants & Other Collaborating Organizations.....	17
8. Special Reporting Requirements.....	20
9. Appendices.....	N/A

1. INTRODUCTION

Autosomal Dominant Polycystic Kidney Disease (ADPKD) is the most prevalent genetic disease, affecting at least 600,000 Americans. It is characterized by massive cystic growth and enlargement of the kidneys, progressing ultimately to renal failure in ~50% of affected individuals. There are currently no FDA-approved pharmaceutical therapies for ADPKD. The genes associated with ADPKD, *Pkd1* and *Pkd2*, encode polycystin-1 (PC1) and polycystin-2 (PC2), respectively. PC1 and PC2 associate with one another to form an ion channel complex in the cilium and the endoplasmic reticulum whose function and regulation are still poorly understood. Cells that are homozygous for ADPKD-causing mutations are hyper-proliferative and manifest substantial perturbations in energy production. These cells exhibit very high levels of aerobic glycolysis and low levels of oxidative metabolism, reminiscent of the Warburg effect that is seen in tumor cells. The Warburg effect has been recognized as a promising target in efforts to develop new drugs for cancer and other hyper-proliferative diseases. Compounds that force cells to derive their energy primarily from oxidative phosphorylation rather than glycolysis may slow the growth of tumor cells and diminish their capacity to thrive in hypoxic environments such as those that characterize solid tumors and renal cysts. In fact, recent work has shown that administration of 2-deoxyglucose perturbs the dependence of ADPKD cells on aerobic glycolysis and substantially slows cyst growth in mouse models of ADPKD.

The mechanism responsible for this metabolic perturbation has not been elaborated. Very recently, we have identified an entirely novel mechanism that we believe is responsible for this “Warburg-like” excessive glycolytic activity that is observed in ADPKD cells. We have discovered that the PC1/2 ion channel complex interacts with and is regulated by the cellular O₂ sensing machinery. We find that cellular O₂ levels modulate the channel activity of the polycystin proteins, which in turn modulates the level of mitochondrial oxidative metabolism. Pyruvate dehydrogenase (PDH) is a key regulator of aerobic metabolism, transforming pyruvate into acetyl-CoA, which can enter the tricarboxylic acid cycle. Our findings suggest that the activity of PDH constitutes a potential drug target in ADPKD. Dichloroacetic acid (DCA) is a small molecule inhibitor of PDH kinase and thus its administration increases PDH activity. DCA has been used in human clinical trials that tested its utility in treating hereditary lactic acidosis and various forms of cancer. The perturbed energy metabolism in ADPKD results in reduced activity of the AMP-activated protein kinase (AMPK). We have previously shown that activating AMPK with metformin slows cyst growth in in vitro and in vivo models of ADPKD. Salicylates and statins have been shown to synergize with metformin in activating AMPK. In the studies described in Aim 1 we have proposed to assess the potential of treatment with DCA, metformin, salicylates and statins, alone or in combination, to correct the perturbed energy metabolism in ADPKD and to slow cyst growth in vitro and in vivo.

One of the current limitations in the development and assessment of new therapies for ADPKD is the lack of clinically useful markers that report on disease severity and progression. The metabolic perturbations that characterize ADPKD-affected cells result in alterations in the inventory of small metabolites that these cells produce. Assessing the profile of metabolites produced by ADPKD cells may lead to the identification of new biomarkers that report on disease progression and on the efficacy of therapeutic interventions. The studies proposed in Aim 2 will seek to identify such metabolism-related markers in vitro and in vivo, and to determine the extent to which they accurately reflect disease severity in samples obtained from human ADPKD patients. These studies have the potential to identify new drug-targetable pathways that can be

exploited in the development of novel therapies for ADPKD and to identify new biomarkers that may prove to be extremely valuable in future clinical trials. The proposed work also has the potential to be translated rapidly into human clinical studies, as agents that will be examined in the proposed studies have been extensively tested in human clinical trials.

2. KEYWORDS

Autosomal Dominant Polycystic Kidney Disease (ADPKD)
Polycystin-1 (PC1)
Polycystin-2 (PC2)
Metformin
Statins
Salicylates
Dichloroacetic acid (DCA)
AMP-activated Protein Kinase (AMPK)
Biomarkers
Metabolism
Mass Spectrometry

3. ACCOMPLISHMENTS

During the first year of grant support, the Caplan laboratory has refined conditions for *in vitro* two- and three-dimensional culture of ADPKD cells and for testing the effects of metformin, statins, salicylates and DCA on their growth and metabolic properties. Extensive optimization procedures to analyze the various metabolomic biomarkers obtained from both cell lysates and from urine specimens have been performed in the Hallows laboratory during the first year of support for the grant.

Task 1. Assess whether PDH kinase inhibition or AMPK activation, alone or in combination, corrects the perturbed energy metabolism in ADPKD cells and slows cyst growth *in vitro* and *in vivo*. The studies encompassed in this task group are designed to determine whether and how reducing PDH kinase activity and/or increasing AMPK activity impacts upon the cellular pathways that are involved in cyst development and expansion. These experiments will make use of cultured ADPKD cells and a mouse model of ADPKD to analyze the potential therapeutic utility of reducing PDH kinase activity and/or increasing AMPK activity.


Task 1a: Determine whether treatment with DCA, metformin, salicylate or simvastatin, alone or in combination, correct the perturbed energy metabolism and slow cyst growth in culture models of ADPKD renal cyst formation.

Effects of energy pathway-targeted treatments on aerobic glycolysis in ADPKD cells: We will measure glucose utilization and lactate production using enzymatic assays and mass spectrometry to determine the quantities of these compounds that are present in media that had been bathing cells for various incubation intervals. Dose ranging studies will be performed to identify concentrations of DCA, metformin, salicylate or simvastatin that effectively correct the energy perturbations observed in ADPKD cells. Cell growth and media collection will be performed in the Caplan laboratory; biochemical analysis of the media will be performed by the Hallows laboratory in association with the studies outlined in **2a** and **2b (months 1-8)**.

Research Accomplishments: Initial studies have been performed in *Pkd1* ^{-/-} (ADPKD) vs. *Pkd1* flox/⁻ (control) cells grown in monolayers on Transwell filters (polarized) and in monolayers in multi-well plates (non-polarized) in the Caplan laboratory. These cells were treated with metformin (1 mM x 24 h) vs. vehicle control. Additional dose response and time course studies of metformin are being carried out in the Caplan laboratory on these cells *in vitro* along with growth studies in 3-D Matrigel cultures. Following treatment, lysates from multiple experiments were sent to the Hallows laboratory for initial targeted metabolomic biomarker analysis (pACC, pP70S6K, pAMPK, pERK, PKM2, PDK1, LDHA, cAMP, lactate, and pyruvate) and to the Pennathur laboratory for preliminary analyses of metabolic intermediates by mass spectrometry (e.g., hexose-6-phosphate, 2,3-diphosphoglycerate, phosphoenolpyruvate, citrate, succinate, and malate).

Of note, our preliminary data suggest that LDHA levels are higher in ADPKD (*Pkd1* null) cells than control cells and that metformin treatment reduces LDHA expression in the ADPKD cells (**Fig. 1A**). Similarly, PDK1 expression levels tended to be higher in the ADPKD cells than in control cells, and metformin treatment reduced PDK1 expression (**Fig. 1B**). Although PKM2

Figure 1: Effects of cell *Pkd1* KO (null) vs. control (het) and metformin treatment (1 mM x 24 h) in cells grown in vitro on key metabolomic biomarkers. A. Relative LDHA levels as detected by immunoblotting of cell lysates. B. Relative PDK1 levels as detected by immunoblotting of cell lysates. C. Relative PKM2 levels as detected by immunoblotting of cell lysates. D. Relative changes in cAMP levels as detected by enzymatic activity assays (*, $P < 0.05$, unpaired t-tests relative to null control group).



adenylyl cyclase (Miller, RA, et al., *Nature* 494: 256, 2013). Results of the other biomarkers tested appear inconclusive so far (not shown).

Effects of PDH kinase inhibition and/or AMPK activation on growth of cultured ADPKD cells: PC1 null (*Pkd1*^{-/-}) and heterozygous (*Pkd1*^{flox/-}) cells will be incorporated into three-dimensional Matrigel matrices, and the size and number of cysts will be determined by quantitative microscopy. Cell proliferation and apoptosis rates will be assessed by immunofluorescence microscopy. The activities of PDH, AMPK and of the mTOR pathway will be determined by Western blotting using phosphorylation-specific antibodies. Media from these cells will be subjected to the metabolomics analyses outlined in **2a** and **2b** to determine whether the doses of DCA, metformin, simvastatin or salicylate that modulate patterns of growth also correct metabolic disturbances. Analysis of cell growth will be performed in the Caplan laboratory, and the Hallows laboratory will analyze culture media in association with the studies outlined in **2a** and **2b** (months 6-12).

Research Accomplishments: The Caplan laboratory has tested the effects of metformin, DCA, simvastatin and salicylate on the growth of *Pkd1*^{-/-} cells in three-dimensional Matrigel culture. When grown in a three-dimensional collagen matrix, mouse *Pkd1*^{-/-} cells form large fluid-filled spherical cysts, whereas cells that express PC1 at native levels frequently grow in roughly

levels were higher in ADPKD cells, metformin did not appear to have any significant effect on these levels in our preliminary experiments (Fig. 1C). Finally, cAMP levels were significantly higher in the ADPKD cells than in control cells, and metformin treatment significantly reduced cAMP levels in the ADPKD cells (Fig. 1D). These results are consistent with the findings of Birnbaum and colleagues, who showed that metformin reduces cAMP production via inhibition of

cylindrical “tubule-like” structures. The extent of these morphological differences can be quantitated by using an ImageJ plug to measure the circularity index, which constitutes a ratio in which the magnitude of a structure’s narrowest dimension is divided by the magnitude of the structure’s widest dimension. For a sphere the circularity index will have a value of 1, whereas tubules will have values of less than 1. The Caplan laboratory has previously found that the mean circularity index for the structures formed by *Pkd1*^{-/-} cells is ~0.7, whereas the mean value for cells expressing native levels of PC1 is ~0.5. Importantly, re-expression of the full length PC1 protein in *Pkd1*^{-/-} cells results in the formation of “tubule-like” rather than cystic structures that have a mean circularity index of ~0.5. These observations indicate that the *in vitro* culture system recapitulates quantifiable aspects of cyst formation and that expression of full-length PC1 protein is sufficient to exert anti-cystogenic activity.

To date the Caplan laboratory has tested the effects on three-dimensional growth of two concentrations each of DCA, metformin, simvastatin and salicylate. Both circularity index and cyst cross-sectional area were measured. The results of these experiments are depicted in the images presented in **Fig. 2** and quantitated in **Fig. 3**. We find that all of the treatments reduce cyst cross-sectional area in a dose-dependent manner. Furthermore, at the highest concentrations applied DCA, simvastatin and salicylate reduce cyst circularity index, whereas metformin does not.

Planned studies for Task 1a: During the next reporting period, dose response and time course experiments will be completed for metformin, simvastatin, salicylate, and DCA. Morphometry, cell proliferation and apoptosis rates will be assayed under these conditions, along with metabolomic biomarker analyses. Media samples obtained from cell culture experiments will also be analyzed for biomarker analyses.

Task 1b: Determine whether DCA, metformin, aspirin or simvastatin slow cyst growth in a mouse model of ADPKD. (A total of at least 320 mice will be used in these proposed studies, as outlined below)




Figure 2: Images of the structures formed by *Pkd1*^{-/-} cells grown in three-dimensional culture in the absence (Control) and in the presence of the various indicated concentrations of DCA, metformin, simvastatin and salicylate.

Defining dose regimens: Prior to assessing the potential utility of DCA, metformin, aspirin or simvastatin administration in *Pkd1*^{flox/-};*Pax8* rtTA;Tet-O cre mice, we will establish dose and delivery regimens that are sufficient to inhibit the activity of PDH kinase and/or activate AMPK in the renal tissue of treated mice. Mice (8 per treatment group) will be treated for 7 days, after which they will be sacrificed and their renal tissue will be analyzed by immunofluorescence microscopy or by western blotting. PDH kinase activity will be determined by performing quantitative immunofluorescence microscopy to measure the pPDH/PDH ratio. AMPK activity will be determined by western blot analysis to assess the ratios of pAMPK/AMPK and pACC/ACC. **We anticipate that 200 mice will be required for these experiments.** Urine and tissue from a subset of these mice will be subjected to the metabolomics analyses outlined in **2a** and **2b**. Dose ranging studies will be performed in the Caplan laboratory in association with the Somlo laboratory, while metabolomics studies will be performed in the Hallows laboratory (**months 12-24**).

Assess the efficacy of DCA, metformin, aspirin or simvastatin *in vivo*: *Pkd1*^{flox/-};*Pax8* rtTA;Tet-O cre mice will be treated or not (8 mice/treatment group) according to the regimens selected in the studies performed in the previous section. Treatment will begin prior to doxycycline induction and will be carried out for 14 weeks. At the end of the treatment period the mice will be sacrificed and the severity of cyst disease will be determined as outlined in the proposal. We expect to repeat this experiment at least three times. Thus, **at least 120 mice will be required to complete these studies.** Urine and tissue from a subset of these mice will be subjected to the metabolomics analyses outlined in **2a** and **2b**. Treatment and assessment of treatment efficacy will be performed by the Caplan laboratory in association with the Somlo laboratory, while metabolomics studies will be




Figure 3: Measurements of cyst cross-sectional area (top) and circularity index (bottom). Each of the treatments reduces cyst area, while DCA, simvastatin and salicylate reduce circularity index whereas metformin does not.

performed in the Hallows laboratory (**months 24-36**).

Research Accomplishments: None of these studies have yet been performed. They were proposed to occur between months 12-36 of the project.

Planned studies for **Task 1b**: During the next reporting period the Caplan laboratory in collaboration with S. Somlo (Co-I) plans to initiate and complete experiments to define the relevant in vivo dosing regimens sufficient to inhibit the activity of PDH kinase and/or activate AMPK in the renal tissue of treated mice. Metabolomics studies pertaining to this work will also be performed by the Hallows laboratory in parallel.

Task 2. Measure candidate metabolomic biomarkers and screen for additional biomarkers that correlate with disease severity and progression in ADPKD mice and cells and that are modulated by reduction of PDH kinase activity or AMPK activation. Cross-correlate biomarker results obtained from mice with banked human urine samples from a cross-section of ADPKD patients (months 0-36).

Task 2a. Investigation of candidate metabolomic biomarkers and profile comparisons as a function of ADPKD severity and treatments.

***In vitro* studies**: Media and cell lysates from experiments outlined in **1a** will be enzymatically assayed for 4 metabolites (lactate, pyruvate, succinate, and cAMP), for 3 glycolytic enzymes (LDHA, PDK1, and PKM2), and for markers of PDH and AMPK activity (total and phosphorylated forms of PDH, AMPK, ACC, and P70S6K). Assays will be performed on triplicate cell lysates from each treatment time point and for each treatment regimen. In addition, media collected over 24-h prior to each time point will also be analyzed. Analysis of metabolites will be performed in the Hallows laboratory, while cell growth and collection will be performed in the Caplan laboratory (**months 1-12**).

Research Accomplishments: See above results outlined under **Task 1a**.

***In vivo* studies**: Urine and renal tissue obtained from animals used in the studies performed in **1b** will be enzymatically assayed for 5 metabolites (lactate, pyruvate, succinate, cAMP, and creatinine), for 3 glycolytic enzymes (LDHA, PDK1, and PKM2), and for markers of PDH and AMPK activity (total and phosphorylated forms of PDH, AMPK, ACC, and P70S6K). Assays will be performed on triplicate tissue homogenates from each treatment time point and for each treatment regimen. In addition, urine collected over 24-h prior to each time point will also be analyzed. Analysis of metabolites will be performed in the Hallows laboratory, while animal treatments and sample collection (**tissue and urine from a total of at least 320 mice; these samples are derived from the same animals employed in Task 1b**) will be performed in the Caplan laboratory in association with the Somlo laboratory (**months 12-36**).

Research Accomplishments: None of these studies have yet been performed. They were proposed to occur between months 12-36 of the project.

Planned studies for **Task 2a**: During the next reporting period, metabolomic biomarker analyses

will be performed by the Hallows laboratory for both in vitro cell culture experiments and in vivo kidney tissue homogenates and urine specimens as these samples become available from the Caplan laboratory. In vivo samples will be compared as a function of disease stage in the animals before and after inducible knockout of the *Pkd1* gene by doxycycline treatment. Media samples obtained from cell culture experiments will also be analyzed for biomarker analyses.

Task 2b. Perform broader targeted metabolomic screens by mass spectrometry (MS) to discover novel metabolite biomarkers that may be informative for ADPKD disease severity and response to treatments.

MS analysis to identify new potential markers: The analytes measured in **2a** may be only a subset of the informative biomarkers correlate with ADPKD disease severity, progression, and/or response to therapies. A broad range of metabolic markers will be screened using cell lysate and culture media samples collected *in vitro* (**1a**) and urine and kidney tissue collected *in vivo* (**1b**). The presence of metabolites will be interrogated in targeted platforms by MS, including all glycolytic and TCA cycle intermediates, nucleotides, and amino acids. Metabolite analysis will be performed by the Hallows laboratory with assistance from Dr. S. Pennathur, Michigan Metabolomics and Obesity Center (MMOC) at U. Michigan and statistical consultation from Dr. K. Abebe at U. Pittsburgh. Cell culture and animal-derived samples (**from a total of at least 320 animals-these samples are derived from the same animals employed in Task 1b**) will be provided by the Caplan laboratory (**months 0-36**).


Research Accomplishments: These studies have been initiated with the in vitro cell lysate samples described above under **Task 1a** with Dr. S. Pennathur (Co-I: University of Michigan).

Planned studies for Task 2b: During the next reporting period, metabolite analyses by MS will be performed by the Pennathur laboratory for both in vitro cell culture experiments and in vivo kidney tissue homogenates and urine specimens as these samples become available from the Caplan laboratory. In vivo samples will be compared as a function of disease stage in the animals before and after inducible knockout of the *Pkd1* gene by doxycycline treatment. Media samples obtained from cell culture experiments will also be analyzed for MS analyses.

Task 2c. Correlate metabolomic biomarkers from banked ADPKD patient urine samples with disease severity and progression and compare these results with those obtained in mice.

A subset of the measurements outlined in **2a** and **2b** will be performed on aliquots of patient urine specimens. Metabolite measurements will be correlated with disease severity parameters. **95 banked, coded ADPKD patient urine specimens and clinical data** were provided by Dr. T. Watnick (U. Maryland) and used for this study. Analysis will be performed in the Hallows laboratory with assistance from Drs. S. Pennathur and K. Abebe (**months 20-36**).

Research Accomplishments: Preliminary analyses for all of the 95 banked, coded ADPKD urine samples have been performed by the Hallows laboratory. Although the data are scattered, there was a significant correlation observed with PKM2 levels in urine samples from patients that were stratified by estimated glomerular filtration rate (eGFR) as estimated by CKD-Epi (Tent, H, et al., *Clin J Am Soc Nephrol* 5: 1960, 2010). Specifically, PKM2 levels tended to be higher in patients with more advanced CKD (<60 ml/min; Stage 3 or higher) than in patients with milder CKD



(≥ 60 ml/min; Stage 1 or 2) (see **Fig. 4A**). However, urinary lactate or pyruvate concentrations did not appear to stratify as a function of CKD severity in this cross-section of ADPKD patients (**Figs. 4B** and **4C**), nor did many of the other biomarkers tested (not shown).

Planned studies for Task 2c: Once additional samples are generated and measurements obtained for the *in vivo* experiments described above in **Tasks 1b** and **2a**, we will perform correlations of the biomarkers obtained from the banked urine specimens derived from our cross-section of ADPKD patients with those obtained from ADPKD mouse tissue at different stages of disease. Dr. Pennathur will also check a subset of the metabolite measurements on the banked ADPKD patient samples by MS for further correlation once the most relevant ones are identified in the earlier *in vitro* and *in vivo* studies described above. Dr. Abebe (paid consultant) will oversee all of the statistical analyses at the end of the study.

Opportunities for Training and Professional Development Provided by the Project:

Nothing to Report.

Dissemination of Results to Communities of Interest:

An overview of preliminary findings obtained so far in this study were included as part of a 30-minute invited talk by Dr. Hallows at the 2016 Baltimore PKD Symposium that was held on September 19, 2016. His talk was titled “Effects of Metformin on the AMPK Pathway and

Metabolomic Biomarkers in ADPKD Kidney Epithelial Cells”. In addition, some of these preliminary results will be shared in a poster presentation at the annual ASN Kidney Week meeting to be held in Chicago, IL in November, 2016. Dr. Caplan has presented aspects of this work in a talk delivered at the University of Oklahoma Medical Center on April 8, 2016.

Research Plans During the Next Reporting Period to Accomplish the Goals:

Please see responses above under each individual task.

4. IMPACT

Impact on the development of the principal discipline(s) of the project? The studies performed to date in the Caplan and Hallows laboratories have permitted both groups to establish the experimental parameters that will be required to move the project forward into its next stages. The three-dimensional culture model has been validated, and the conditions that need to be used in the assays of metabolic activity have been established. Thus, the groundwork has been successfully laid for the subsequent phases of the project. These studies have provided tools that will prove extremely useful to the ADPKD research community.

Impact on other disciplines? Nothing to report

Impact on technology transfer? Nothing to report

Impact on society beyond science and technology? Nothing to report

5. CHANGES/PROBLEMS

Nothing to report

6. PRODUCTS

Nothing to report

7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

Name: Michael Caplan
Project Role: Initiating PI
Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 2.4
Contribution to Project: Dr. Caplan is responsible for co-supervision, with Dr. Hallows, of all of the studies described in the proposal. He is directly involved in every aspect of experimental design and data interpretation.
Funding Support: NA

Name: Valeria Padovano
Project Role: Postdoctoral Associate
Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 12
Contribution to Project: Dr. Padovano has conducted the in vitro cell growth studies and performed all of the relevant analyses
Funding Support: NA

Name: Vanathy Rajendran
Project Role: Research Associate II
Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 4.8
Contribution to Project: Ms. Vanathy Rajendran carries out the laboratory's tissue culture work required for the in vitro studies. She also participates in molecular biologic, biochemical and immunocytochemical procedures and maintains laboratory stock solutions.
Funding Support: NA

Name: Stefan Somlo
Project Role: Collaborator
Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 0.36
Contribution to Project: Dr. Somlo is supervising the breeding of the mouse stocks that will be used in years 2 and 3 of the project.
Funding Support: NA

Name: Seung Lee
Project Role: Postdoctoral Associate

Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 3
Contribution to Project: Dr. Lee is carrying out the breeding of the mouse stocks that will be used in years 2 and 3 of the project.
Funding Support: NA

Name: Kenneth Hallows
Project Role: Partnering PI
Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 1.8
Contribution to Project: Dr. Hallows is responsible for co-supervision, with Dr. Caplan, of all of the studies described in the proposal. He is directly involved in every aspect of experimental design and data interpretation.
Funding Support: NA

Name: Hui Li
Project Role: Senior Investigator
Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 4.2
Contribution to Project: Dr. Li has overseen the daily operation of the Hallows lab biomarker measurements and has personally performed all of the immunoblotting and some of the multi-well plate reader studies. He has also performed all of the relevant analyses.
Funding Support: NA

Name: Daniel Rivera
Project Role: Research Lab Technician
Researcher Identifier (e.g., ORCID ID): NA
Nearest person month worked: 6
Contribution to Project: Mr. Rivera has conducted most of the multi-well plate reader studies of metabolomic biomarkers on the urine and cell lysate samples along with the relevant analyses.

Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period? Nothing to report

What other organizations were involved as partners? The University of Michigan Metabolomics Core run by Dr. Subramaniam Pennathur (Associate Professor co-investigator on this project) and his team of personnel, including Drs. Jaeman Byun and Lixia Zeng, were

engaged during this past year of support to perform a broader array of analysis of metabolites by mass spectrometry on the experimental samples generated by the Caplan laboratory. They have received a sub-contract awarded through Dr. Hallows' grant award.

8. SPECIAL REPORTING REQUIREMENTS

Collaborative Awards: Reports will be submitted directly by both Dr. Caplan (Initiating PI) and by Dr. Hallows (Partnering PI). These reports indicate the site at which each of the studies described is conducted.