



AFRL-AFOSR-VA-TR-2017-0031

Reaching higher Gamma in ultracold neutral plasmas through
disorder-induced heating control

Scott Bergeson
BRIGHAM YOUNG UNIVERSITY
UNIVERSITY HILL A-261, ASB
PROVO, UT 84602 - 1231

02/07/2017
Final Report

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory
AF Office Of Scientific Research (AFOSR)/RTB1

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.					
1. REPORT DATE (DD-MM-YYYY) 27-06-2016		2. REPORT TYPE Final Report		3. DATES COVERED (From - To) 01 June 2012 - 31 May 2016	
4. TITLE AND SUBTITLE Reaching higher Gamma in ultracold neutral plasmas through disorder-induced heating control				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER FA9550-12-1-0308	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Scott D. Bergeson				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Brigham Young University Office of research and creative activities A-41 ASB Provo, UT 84602-1014				8. PERFORMING ORGANIZATION REPORT NUMBER R0202364	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) USAF, AFRL DUNS 143574726 Air Force Office of Scientific Research 875 N. Randolph St. Room 3112 Arlington VA 22203				10. SPONSOR/MONITOR'S ACRONYM(S) AFOSR	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) FA9550-12-1-0308	
12. DISTRIBUTION/AVAILABILITY STATEMENT DISTRIBUTION A: Distribution approved for public release.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT This is the final report on project FA9950-12-1-0308, "Reaching higher Gamma through disorder-induced heating control." Highlights from the grant include making the first ion temperature measurements in a multiply-ionized ultracold neutral plasma (Ca/Ca+), making the first measurements of the ion temperature in a plasma that evolves from a dense Rydberg gas, and starting work on the first dual-species MOT and plasma.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF: a. REPORT b. ABSTRACT c. THIS PAGE			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Marilyn Webb 19b. TELEPHONE NUMBER (Include area code) 801-422-3607

INSTRUCTIONS FOR COMPLETING SF 298

1. REPORT DATE. Full publication date, including day, month, if available. Must cite at least the year and be Year 2000 compliant, e.g. 30-06-1998; xx-06-1998; xx-xx-1998.

2. REPORT TYPE. State the type of report, such as final, technical, interim, memorandum, master's thesis, progress, quarterly, research, special, group study, etc.

3. DATES COVERED. Indicate the time during which the work was performed and the report was written, e.g., Jun 1997 - Jun 1998; 1-10 Jun 1996; May - Nov 1998; Nov 1998.

4. TITLE. Enter title and subtitle with volume number and part number, if applicable. On classified documents, enter the title classification in parentheses.

5a. CONTRACT NUMBER. Enter all contract numbers as they appear in the report, e.g. F33615-86-C-5169.

5b. GRANT NUMBER. Enter all grant numbers as they appear in the report, e.g. AFOSR-82-1234.

5c. PROGRAM ELEMENT NUMBER. Enter all program element numbers as they appear in the report, e.g. 61101A.

5d. PROJECT NUMBER. Enter all project numbers as they appear in the report, e.g. 1F665702D1257; ILIR.

5e. TASK NUMBER. Enter all task numbers as they appear in the report, e.g. 05; RF0330201; T4112.

5f. WORK UNIT NUMBER. Enter all work unit numbers as they appear in the report, e.g. 001; AFAPL30480105.

6. AUTHOR(S). Enter name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. The form of entry is the last name, first name, middle initial, and additional qualifiers separated by commas, e.g. Smith, Richard, J, Jr.

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES). Self-explanatory.

8. PERFORMING ORGANIZATION REPORT NUMBER. Enter all unique alphanumeric report numbers assigned by the performing organization, e.g. BRL-1234; AFWL-TR-85-4017-Vol-21-PT-2.

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES). Enter the name and address of the organization(s) financially responsible for and monitoring the work.

10. SPONSOR/MONITOR'S ACRONYM(S). Enter, if available, e.g. BRL, ARDEC, NADC.

11. SPONSOR/MONITOR'S REPORT NUMBER(S). Enter report number as assigned by the sponsoring/monitoring agency, if available, e.g. BRL-TR-829; -215.

12. DISTRIBUTION/AVAILABILITY STATEMENT. Use agency-mandated availability statements to indicate the public availability or distribution limitations of the report. If additional limitations/ restrictions or special markings are indicated, follow agency authorization procedures, e.g. RD/FRD, PROPIN, ITAR, etc. Include copyright information.

13. SUPPLEMENTARY NOTES. Enter information not included elsewhere such as: prepared in cooperation with; translation of; report supersedes; old edition number, etc.

14. ABSTRACT. A brief (approximately 200 words) factual summary of the most significant information.

15. SUBJECT TERMS. Key words or phrases identifying major concepts in the report.

16. SECURITY CLASSIFICATION. Enter security classification in accordance with security classification regulations, e.g. U, C, S, etc. If this form contains classified information, stamp classification level on the top and bottom of this page.

17. LIMITATION OF ABSTRACT. This block must be completed to assign a distribution limitation to the abstract. Enter UU (Unclassified Unlimited) or SAR (Same as Report). An entry in this block is necessary if the abstract is to be limited.

Final report: Achieving higher Gamma in ultracold neutral plasmas through disorder-induced heating control

S. D. Bergeson

Department of Physics and Astronomy,
Brigham Young University, Provo, UT 84602, USA
scott.bergeson@byu.edu

June 28, 2016

Abstract

This is the final report on project FA9950-12-1-0308, “Reaching higher Gamma through disorder-induced heating control.” Highlights from the grant include making the first ion temperature measurements in a multiply-ionized ultracold neutral plasma (Ca/Ca⁺), making the first measurements of the ion temperature in a plasma that evolves from a dense Rydberg gas, and starting work on the first dual-species MOT and plasma.

This project included three major activities. The first was to explore the possibility of increasing the strong-coupling parameter in an ultracold neutral plasma by sequentially ionizing to higher ionization states. This work was motivated by a calculation indicating that it would be possible to quadruple the coupling parameter. The second major activity was to explore the possibility of using the Rydberg excitation blockade to pre-order the neutral gas before ionization. The third major activity was to build a new experiment to study energy relaxation in a dual-species ultracold neutral plasma.

This document summarizes achievements at Brigham Young University during the award FA9950-12-1-0308 from June 1, 2012 to May 31, 2016. The primary focus of this effort was to find ways to increase the value of the strong coupling parameter in ultracold neutral plasmas. This parameter is defined as the ratio of the nearest-neighbor Coulomb energy to the average ion kinetic energy in the plasma,

$$\Gamma = \frac{Z^2 e^2}{4\pi\epsilon_0 a_{\text{ws}}} \frac{1}{k_B T}, \quad (1)$$

where Z is the ion charge state, $a_{\text{ws}} = (3/4\pi n)^{1/3}$ is the Wigner-Seitz radius, n is the ion (number) density, and T is the ion temperature.

The value of the strong-coupling parameter is limited by disorder-induced heating in ultracold neutral plasmas. This heating occurs as the ions relax towards equilibrium. The plasma ions have essentially no kinetic energy when they are first ionized. However, due to the random spatial distribution in the atom trap, the ions have a comparatively large value of electrical potential energy. Energy relaxation converts the potential energy to kinetic energy and the plasma ions heat up dramatically. In our experiment, laser-cooled atoms ($T \sim 1$ mK) heat to temperatures near 2 K in a time frame of 100 to 500 ns.

Achieving higher values of the strong coupling parameter remains a major priority for this field. Molecular dynamics simulations show that thermodynamic and kinetic properties of these plasmas scale with the strong-coupling parameter. However, connecting these computationally-intensive simulations with extensions of kinetic theories remains a challenge. Carefully diagnosed and high accuracy experiments that study the characteristics of neutral plasmas over a range of strong-coupling parameter values can help in this effort. Because plasma properties scale with Γ , measurements in ultracold neutral plasmas can inform the diagnostics of high-energy-density plasmas. Better understanding of collision rates, equilibration, and energy transport in warm dense matter may lead to important technological and scientific advances in fusion class plasmas. Given the failure to achieve ignition at NIF, a better understanding of strongly-coupled neutral plasmas remains a national priority.

Table 1: Undergraduate students supported by this project. The titles of the senior theses written by these students is also give.

Student name	Senior thesis title	Post-BYU employment
Meredith Gold Dahl	Determining the accuracy of a partially-stabilized frequency comb	Grad. student at U. Rochester
Daniel Woodbury	Construction of a compact two-dimensional magneto-optical trap for a cold calcium beam	Grad. student at U. Maryland
Joshua Wilson	Ultracold neutral plasma physics at room temperature	Grad. student at Penn State
Alex Erikson	A permanent magnet Zeeman slower (in process)	Medical school, Fall 2017
Nicholas Harrison	Radius and intensity dependence of laser plasma expansion	Private sector
Daniel Crunkleton	Strong field ionization of neutral strongly-coupled plasmas in a gas jet	USAF
Stephen Rubber	Frequency locking circuits	Sandia National Lab (starting graduate school at BYU 2016)

1 Personnel

This grant supported one graduate student, Mary E. Lyon. Dr. Lyon received her Ph.D. in December 2014. She is the lead author on five of the six archival publications from the grant. After receiving her degree, she accepted a post-doctoral research position at the Joint Quantum Institute at the University of Maryland.

This grant also supported seven undergraduate students At Brigham Young University. They are listed in Table 1. This research project is continuing under NSF support. This fall two graduate students (Stephen Rupper and Tucker Sprenkle) and three undergraduate students (Heather Longstaff, Quin McNight, and Michael Peterson) will be working in my lab on this and related projects.

2 Publications

During the grant, six peer-reviewed journal articles were published. During this past year, four conference presentations were also made. These are enumerated in Tables 2 and 3.

Table 2: Peer-reviewed publications from this grant.

1.	M. Lyon, S. D. Bergeson, G. Hart, and M. S. Murillo, "Strongly-coupled plasmas formed from laser-heated solids," Scientific Reports 5, 15693 (2015), DOI:10.1038/srep15693
2.	S. D. Bergeson, M. Lyon, J. B. Peatross, N. Harrison, D. Crunkleton, J. Wilson, S. Rupper, A. Diaw and M. S. Murillo, "A neutral strongly coupled laser-produced plasma by strong-field ionization in a gas jet," AIP Conf. Proc. 1668, 040001 (2015), December 2014, Takamatsu, Japan, DOI:10.1063/1.4923114
3.	M. Lyon and S. D. Bergeson, "Towards stronger Coulomb coupling in an ultracold neutral plasma," Contrib. Plasma Phys. 55, 399-406 (2015), DOI:10.1002/ctpp.201400097
4.	M. Lyon, S. D. Bergeson, A. Diaw, and M. S. Murillo, "Using higher ionization states to increase Coulomb coupling in an ultracold neutral plasma," Phys. Rev. E 91, 033101 (2015)
5.	M. Lyon and S. D. Bergeson, "Precision spectroscopy using a partially stabilized frequency comb," Appl. Opt. 53, 5163-5168 (2014)
6.	M. Lyon, S. D. Bergeson and M. S. Murillo, "The limit of strong ion coupling due to electron shielding," Phys. Rev. E 87, 033101 (2013)

3 Higher Gamma using electron screening

The first publication in this grant explored the possibility of using electron screening to reduce the temperature of the ions and thereby increase the strong coupling parameter. In previous work in our lab we showed that the electron screening results in very low temperatures. Working with Michael Murillo, we showed that unfortunately the electron screening reduces the nearest-neighbor potential energy by the same amount. Therefore the physical value of the strong coupling parameter remains unchanged. It is important to understand this point. Although some groups report somewhat higher values of Γ , these can only be viewed as a scaled inverse ion temperature and not a physically meaningful ratio of potential-to-kinetic energy.

Publications:

- M. Lyon, S. D. Bergeson and M. S. Murillo, “The limit of strong ion coupling due to electron shielding,” Phys. Rev. E 87, 033101 (2013)

4 Sequential ionization of ultracold plasma ions

A simulation published in 2007 by Michael Murillo showed that the quasi-coherent kinetic energy oscillations of ions in ultracold neutral plasmas could be used to increase the strong coupling parameter by a factor of 4 (see Eq. 1 and Fig. 1).

We built an experiment to explore this possibility in neutral Calcium. We used five pulsed lasers to reach the second ionization stage. While the disorder-induced heating may not increase when the second ionization event occurs, the electron temperature increases significantly. This increases the plasma expansion rate, as shown in Fig. 2.

One of the limitations in this approach is that the strong laser fields required for ionization Stark-shift the intermediate energy levels out of resonance with the excitation lasers. The maximum achievable second ionization fraction was therefore 35%, consistent with a rate-equation model for the excitation. The limited second-ionization resulted in some additional ion heating, as verified by molecular dynamics simulations of the experiment (see Figs. 3 and 4).

Publications:

- M. Lyon and S. D. Bergeson, “Towards stronger Coulomb coupling in an ultracold neutral plasma,” Contrib. Plasma Phys. 55, 399-406 (2015)
- M. Lyon, S. D. Bergeson, A. Diaw, and M. S. Murillo, “Using higher ionization states to increase Coulomb coupling in an ultracold neutral plasma,” Phys. Rev. E 91, 033101 (2015)

5 Generating spatial order via the Rydberg excitation blockade

Several demonstrations of the Rydberg blockade have been published in ultracold atom experiments. The blockade occurs when the atom-atom interaction results in an energy level shift that is greater than linewidth of both the laser and the atomic level. At long range, the energy level shift can be described as a van der Waals interaction between excited atoms using a relatively simple atomic theory. Most of the research in

Table 3: Contributed conference presentations from this grant during the past year.

1. (Invited) S. D. Bergeson, “What 1-Kelvin plasmas can tell you about thermonuclear fusion,” APS 4-Corners Meeting, Tempe AZ, October 2015
 2. (Invited) S. Bergeson, “Some progress towards an ultracold Ca/Yb plasma,” AFOSR workshop on ultracold neutral plasmas, Santa Fe, NM, May 2016
 3. M. Lyon, S. Bergeson, G. Hart, M. Murillo, “Strongly-coupled plasmas formed from laser-heated solids,” DAMOP, Providence RI, May 2016
 4. S. Bergeson and M. Lyon, “Measurements of the ion velocity distribution in an ultracold neutral plasma derived from a cold, dense Rydberg gas,” DAMOP, Providence RI, May 2016
-




Figure 1: Temperature as a function of scaled time for ions in an ultracold neutral plasma. The blue line shows the temperature evolution due to disorder-induced heating. The red line shows the predicted temperature evolution if the ions are promoted to the second ionization state at the peak of the blue curve. In this case, no additional disorder-induced heating occurs because ionization occurs at the minimum in the ion potential energy evolution. The green line shows the predicted temperature increase when the ions are promoted to the second ionization state at later times. From M. S. Murillo, Phys. Plasmas 14, 055702 (2007).




Figure 2: RMS ion velocity in an ultracold neutral calcium plasma. The inset shows the excitation pathway. The velocity distribution was probed using laser-induced fluorescence at 397 nm on the residual Ca^+ ions. The second ionization event occurred at 100 ns for this data. Scattered light, optical pumping, and spontaneous emission prevented meaningful velocity measurements closer than ± 30 ns after the ionization event. The gray line is the rms velocity without the second ionization event. The black dots represent the rms velocity when the second ionization event occurs. The red lines are fits of the data to expansion models. From M. Lyon, S. D. Bergeson, A. Diaw, and M. S. Murillo, Phys. Rev. E 91, 033101 (2015).




Figure 3: Extracted increase in the ion temperature as a function of time delay between the first and second ionization event. The green line shows the predicted increase in temperature from a simulation of the experiment. From M. Lyon, S. D. Bergeson, A. Diaw, and M. S. Murillo, Phys. Rev. E 91, 033101 (2015).




Figure 4: Molecular dynamics results showing the increase in temperature and also the value of the strong-coupling parameter Γ as a function of ionization fraction. From M. Lyon, S. D. Bergeson, A. Diaw, and M. S. Murillo, Phys. Rev. E 91, 033101 (2015).

this area occurs in this low-energy regime. That research is generally motivated by applications in quantum information.

In certain configurations, the long-range repulsive interaction between Rydberg atoms is used to generate spatial order in neutral gases. While most demonstrations of this effect have been made using degenerate quantum gases or atoms in optical dipole traps, a few publications suggest that spatial ordering might be achievable in a magneto-optical trap like the one used in our lab. If the Rydberg system could be rapidly ionized, the spatial ordering from the blockade is predicted to be sufficient to increase the value of the strong-coupling parameter to $\Gamma \leq 30$.

We explored the possibility of using the Rydberg blockade to generate a spatially ordered atomic sample. We used an excitation scheme that is typical of Rydberg excitation experiments (see Fig. 5). Our initial work using cw laser excitation proved to be insufficient to achieve enough excitation before the calcium atoms moved significantly. When the atoms move during the excitation pulse, they collide with neighboring atoms and collisionally ionize, thereby compromising the blockade effect.

We further explored the possibility of using ns-duration pulsed lasers to achieve the Rydberg blockade. Typical excitation efficiency measurements are shown in Fig. 6 for excitation to the $n = 30$ state. We measured these kinds of efficiency curves for a number of different Rydberg levels. The results are shown in Fig. 7.

Following excitation, we used a weak cw laser tuned to the ion resonance transition at 397 nm to probe the velocity distribution of the Rydberg atoms. Results are shown in Fig. 8. The probe laser beam caused ionization of the Rydberg atoms due to core-interaction. Therefore we introduced a variable time delay, between 0 and 10 μ s in order to let the Rydberg system evolve unperturbed before probing the system. Any reduction in disorder-induced heating should be apparent in the first few μ s of the data. However, a reduction is not clearly evident and further work is required to understand the system in detail. In particular, excitation with the large Rabi frequencies necessary to achieve collective excitation in a few ns is probably inadequately described by a simple van der Waals analysis of the Rydberg-Rydberg atom interaction. Full molecular potential curves need to be calculated and analyzed for this project to move forward.

Publications:

- S. D. Bergeson and M. Lyon, “Measurements of the ion velocity distribution in an ultracold neutral plasma derived from a cold, dense Rydberg gas,” arXiv:1601.07439.




Figure 5: Partial energy level diagram for calcium. a) The two-photon Rydberg excitation pathway using laser pulses at 423 and 390 nm. b) The excitation and fluorescence scheme used to diagnose the Ca^+ ions formed after Rydberg excitation.




Figure 7: Rydberg atom excitation efficiency for a range of different principle quantum numbers and a range of different laser frequencies. These data are suggestive of blocked excitation. However more work needs to be done to understand the excitation dynamics. The densities and excitation energies are high enough that a simple van der Waals treatment is probably insufficient. The legend indicates the excitation laser intensity in units of 10^7 mW/cm^2 . From S. D. Bergeson and M. Lyon, arXiv:1601.07439.




Figure 6: Rydberg excitation efficiency as a function of Ω_2 and fixed $\Omega_1 = 350 \text{ MHz}$ for a principle quantum number of 30. As the laser intensity increases (Ω_2 increases), the excitation efficiency also increases. At high intensities the Rydberg excitation frequency is Stark-shifted by the pulsed laser.




Figure 8: RMS velocity distribution of the Rydberg atoms/plasma generated after excitation to $n = 50$ at high laser power shown in Figure 7. Because of the ionization resulting from core-excited Rydberg atoms, a time delay was introduced in between Rydberg excitation and the probe laser. Evidence of reduced disorder-induced heating should be apparent in the first 1 or 2 μs of the data. However, any reduction is not clear. From S. D. Bergeson and M. Lyon, arXiv:1601.07439.

6 Dual-species plasma work

During a one-year extension to this project, we built a dual-species MOT for Yb and Ca. These atoms differ from one another by a factor of 4 in mass. The plan is to ionize both of these species and to study energy relaxation and velocity diffusion. We have trapped Ca and Yb atoms simultaneously in our MOT. An initial study of the trap losses indicate that these atoms do not interact strongly. There is no Yb-induced loss in the Ca MOT and no Ca-induced loss in the Yb MOT. In other dual-species MOTs, simultaneous trapping of different atoms is often impractical because the losses are too great. However, there is no hyperfine splitting in Ca-40, and the hyperfine splitting in Yb is all in the excited state. This bodes well for our future work.

The atomic densities in the MOT are approximately 10^{16} cm^{-3} . As of this writing, work is underway to increase the number of atoms in the MOT and to optimize trapping parameters.

This experiment requires four frequency-doubled ti:sapphire lasers, all of which are in place in our lab. It also requires four pulsed lasers for ionization, three of which are in our lab. The final one should be in place before the end of the year.

This work will continue with funding from the joint NSF/DOE plasma physics program. This is a collaborative research proposal that includes Professor Michaela Kleinert at Willamette University in Oregon, Michael Murillo at the New Mexico Consortium, and Scott Bergeson at Brigham Young University.

1.

1. Report Type

Final Report

Primary Contact E-mail**Contact email if there is a problem with the report.**

scott.bergeson@byu.edu

Primary Contact Phone Number**Contact phone number if there is a problem with the report**

8014226161

Organization / Institution name

Brigham Young University

Grant/Contract Title**The full title of the funded effort.**

Reaching higher Gamma in ultracold neutral plasmas through disorder-induced heating control

Grant/Contract Number**AFOSR assigned control number. It must begin with "FA9550" or "F49620" or "FA2386".**

FA9550-12-1-0308

Principal Investigator Name**The full name of the principal investigator on the grant or contract.**

Scott Bergeson

Program Manager**The AFOSR Program Manager currently assigned to the award**

Tatjana Curcic

Reporting Period Start Date

06/01/2015

Reporting Period End Date

05/31/2016

Abstract

This is the final report on project FA9950-12-1-0308, "Reaching higher Gamma through disorder-induced heating control." Highlights from the grant include making the first ion temperature measurements in a multiply-ionized ultracold neutral plasma ($\text{Ca}/\text{Ca}^{\wedge+}$), making the first measurements of the ion temperature in a plasma that evolves from a dense Rydberg gas, and starting work on the first dual-species MOT and plasma. This project included three major activities. The first was to explore the possibility of increasing the strong-coupling parameter in an ultracold neutral plasma by sequentially ionizing to higher ionization states. This work was motivated by a calculation indicating that it would be possible to quadruple the coupling parameter. The second major activity was to explore the possibility of using the Rydberg excitation blockade to pre-order the neutral gas before ionization. The third major activity was to build a new experiment to study energy relaxation in a dual-species ultracold neutral plasma.

Distribution Statement**This is block 12 on the SF298 form.**

Distribution A - Approved for Public Release

Explanation for Distribution Statement**If this is not approved for public release, please provide a short explanation. E.g., contains proprietary information.**

DISTRIBUTION A: Distribution approved for public release.

SF298 Form

Please attach your [SF298](#) form. A blank SF298 can be found [here](#). Please do not password protect or secure the PDF. The maximum file size for an SF298 is 50MB.

[AFD-070820-035.pdf](#)

Upload the Report Document. File must be a PDF. Please do not password protect or secure the PDF. The maximum file size for the Report Document is 50MB.

[afosr2016FinalReport.pdf](#)

Upload a Report Document, if any. The maximum file size for the Report Document is 50MB.

Archival Publications (published) during reporting period:

1. M. Lyon, S. D. Bergeson, G. Hart, and M. S. Murillo, "Strongly-coupled plasmas formed from laser-heated solids," Scientific Reports 5, 15693 (2015), DOI:10.1038/srep15693
2. S. D. Bergeson, M. Lyon, J. B. Peatross, N. Harrison, D. Crunkleton, J. Wilson, S. Rupper, A. Diaw and M. S. Murillo, "A neutral strongly coupled laser-produced plasma by strong-field ionization in a gas jet," AIP Conf. Proc. 1668, 040001 (2015), December 2014, Takamatsu, Japan, DOI:10.1063/1.4923114
3. M. Lyon and S. D. Bergeson, "Towards stronger Coulomb coupling in an ultracold neutral plasma," Contrib. Plasma Phys. 55, 399-406 (2015), DOI:10.1002/ctpp.201400097
4. M. Lyon, S. D. Bergeson, A. Diaw, and M. S. Murillo, "Using higher ionization states to increase Coulomb coupling in an ultracold neutral plasma," Phys. Rev. E 91, 033101 (2015)
5. M. Lyon and S. D. Bergeson, "Precision spectroscopy using a partially stabilized frequency comb," Appl. Opt. 53, 5163-5168 (2014)
6. M. Lyon, S. D. Bergeson and M. S. Murillo, "The limit of strong ion coupling due to electron shielding," Phys. Rev. E 87, 033101 (2013)
7. S. D. Bergeson and M. Lyon, "Measurements of the ion velocity distribution in an ultracold neutral plasma derived from a cold, dense Rydberg gas," arXiv:1601.07439 (2016)

2. New discoveries, inventions, or patent disclosures:

Do you have any discoveries, inventions, or patent disclosures to report for this period?

No

Please describe and include any notable dates

Do you plan to pursue a claim for personal or organizational intellectual property?

Changes in research objectives (if any):

A one-year extension was granted that allowed us to build a dual-species Yb/Ca MOT. This is a new extension of the originally proposed work. It is major stepping stone towards realizing a dual-species ultracold neutral plasma.

Change in AFOSR Program Manager, if any:

None

Extensions granted or milestones slipped, if any:

A one-year extension was granted that allowed us to build a dual-species Yb/Ca MOT. This is a major stepping stone towards realizing a dual-species ultracold neutral plasma.

AFOSR LRIR Number

LRIR Title

Reporting Period

Laboratory Task Manager

Program Officer

Research Objectives

Technical Summary

Funding Summary by Cost Category (by FY, \$K)

	Starting FY	FY+1	FY+2
Salary			
Equipment/Facilities			
Supplies			
Total			

Report Document

Report Document - Text Analysis

Report Document - Text Analysis

Appendix Documents

2. Thank You

E-mail user

Jun 28, 2016 14:40:21 Success: Email Sent to: scott.bergeson@byu.edu