

Award Number: W81XWH-12-1-0585

TITLE: Intranasal Insulin: A Novel Treatment for Gulf War Multisymptom Illness

PRINCIPAL INVESTIGATOR: Dr. Julia Golier

CONTRACTING ORGANIZATION: Bronx Veterans Medical Research Foundation, Inc.
Bronx, NY 10468

REPORT DATE: October, 2017

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE October 2017		2. REPORT TYPE Annual		3. DATES COVERED 30Sep2016 - 29Sep2017	
4. TITLE AND SUBTITLE Intranasal Insulin: A Novel Treatment for Gulf War Multisymptom Illness			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER W81XWH-12-1-0585		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Julia Golier E-Mail: Julia.Golier@va.gov			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Bronx Veterans Medical Research Foundation, Inc. Bronx, NY 10468			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Following their deployment to the 1991 Gulf War, many veterans (GWV) reported a constellation of unexplained health symptoms; common among them were attention and memory difficulties, fatigue, joint pain, headaches, gastrointestinal complaints, and mood and sleep problems (Proctor et al., 1998; Sullivan et al., 2003). Despite the passage of time, the symptom complex persists for many veterans. Indeed, it is estimated that at least 25 percent of GWV (nearly 170,000 veterans) have a persistent form of chronic multisymptom illness (CMI) (Kang et al., 2009; Gulf War Research Advisory Committee (RAC), 2008; IOM, 2010). GW deployed veterans are also developing significantly more chronic diseases such as diabetes, hypertension, arthritis, and coronary heart disease than their non-deployed veteran peers (Toomey et al., 2009; Chao et al., 2010; Chao et al., 2011; Li et al., 2011) putting these individuals at risk for accelerated aging-related diseases of the peripheral and central nervous system (CNS). Over the years it has been found that cognitive complaints have been particularly troublesome to GWV. Recent studies have shown a slowing of response speed that affects mental flexibility across multiple cognitive domains (memory, attention, visuospatial functions) especially on tests that were timed and computerized and where small differences in cognitive reaction times could be measured (Anger et al., 1999; RAC, 2008; Krengel and Sullivan, 2008; Toomey et al., 2009; Chao et al., 2011). Recent studies also have suggested that the response inhibition deficits shown in GWV may reflect executive system dysfunction (Tillman et al., 2010) as reflected by slower motor responses across multiple cognitive domains (RAC, 2008). To date, there are no treatments that have been shown to substantially improve cognitive impairments or health symptoms of GWVs. Thus, it is of paramount importance to identify effective, safe, and tolerable treatments for Gulf War CMI.					
15. SUBJECT TERMS Nothing listed					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	4
Keywords.....	4
Overall Project Summary.....	4
Key Research Accomplishments.....	5
Reportable Outcomes.....	N/A
Conclusion.....	N/A
References.....	6

INTRODUCTION:

Background: Following their deployment to the 1991 Gulf War, many veterans (GWV) reported a constellation of unexplained health symptoms; common among them were attention and memory difficulties, fatigue, joint pain, headaches, gastrointestinal complaints, and mood and sleep problems (Proctor et al., 1998; Sullivan et al., 2003). Despite the passage of time, the symptom complex persists for many veterans. Indeed, it is estimated that at least 25 percent of GWV (nearly 170,000 veterans) have a persistent form of chronic multisymptom illness (CMI) (Kang et al., 2009; Gulf War Research Advisory Committee (RAC), 2008; IOM, 2010). GW deployed veterans are also developing significantly more chronic diseases such as diabetes, hypertension, arthritis, and coronary heart disease than their non-deployed veteran peers (Toomey et al., 2009; Chao et al., 2010; Chao et al., 2011; Li et al., 2011) putting these individuals at risk for accelerated aging-related diseases of the peripheral and central nervous system (CNS). Over the years it has been found that cognitive complaints have been particularly troublesome to GWV. Recent studies have shown a slowing of response speed that affects mental flexibility across multiple cognitive domains (memory, attention, visuospatial functions) especially on tests that were timed and computerized and where small differences in cognitive reaction times could be measured (Anger et al., 1999; RAC, 2008; Kregel and Sullivan, 2008; Toomey et al., 2009; Chao et al., 2011). Recent studies also have suggested that the response inhibition deficits shown in GWV may reflect executive system dysfunction (Tillman et al., 2010) as reflected by slower motor responses across multiple cognitive domains (RAC, 2008). To date, there are no treatments that have been shown to substantially improve cognitive impairments or health symptoms of GWVs. Thus, it is of paramount importance to identify effective, safe, and tolerable treatments for Gulf War CMI.

KEYWORDS: Insulin, clinical trial, novel therapeutics, inflammation, cortisol, chronic fatigue, multi-symptom illness, malaise, deployment.

OVERALL PROJECT SUMMARY:

Objective: To test whether insulin, administered intranasally, improves the health and functioning of GWV with CMI.

Specific Aims: (1) To assess the efficacy of two different doses (10 IU BID and 20 IU BID) of daily intranasal insulin for eight weeks on memory and attention functioning in GWV with CMI. (2) To assess the efficacy of two different doses of intranasal insulin on overall physical health and mood in GWV with CMI. (3) To characterize the effect of two different doses of intranasal insulin on other symptoms that are characteristic of or associated with CMI (e.g., fatigue, pain, sleep quality, subjective cognitive function). (4) To assess the safety of two different doses of self-administered intranasal insulin in GWV with CMI.

Study Design: 114 eligible GWVs with CMI will be randomly assigned in parallel groups to treatment with 20 IU (i.e., 10 IU BID (after breakfast and dinner)), 40 IU (i.e., 20 IU BID (after breakfast and dinner)), or placebo for eight weeks and assessed for clinical outcomes at treatment endpoint. The treatment groups will self-administer 10 IU insulin or 20 IU insulin through

a nasal infusion pump twice daily through the nose. The placebo group will administer saline through a nasal infusion pump twice daily as well. The primary outcome measure will be neuropsychological outcome (verbal memory and selective attention). As this will be the first trial of intranasal insulin in Gulf War veterans, a dose-finding clinical trial is proposed using two doses within the range that has been shown to be effective and safe in cognitively impaired older adults. Treatment duration of eight weeks was chosen in order to assess the effect of sustained intranasal treatment on cognition, mood, and overall health; a post-treatment follow-up assessment will be performed to characterize the sustainability of treatment effects.

KEY RESEARCH ACCOMPLISHMENTS:

Study recruitment has shown a sharp increase since last year due to increased advertising and outreach. Participants are currently enrolled in study procedures and being monitored throughout the process. The Bronx team plans to continue recruiting Gulf War Veteran's by visiting other VA hospitals and vet centers in the surrounding areas.

A protocol amendment was submitted and approved by the IRB to mainstream questionnaires to decrease participant burden. An additional blood draw was added to the protocol to monitor for health and safety at visit 6 and visit 7. The consent form was also amended to reflect the protocol changes. Both the protocol amendment and consent form amendment were approved by the IRB and currently in use. A two year no-cost extension has been approved by the DoD.

There is currently a new, updated version of the intranasal device that has improved upon the old design. We are currently speaking with the company to look into updating the devices for the study, as some participants have complained of issues with the current devices. We are awaiting approval on our no-cost extension before making any financial decisions. The team will continue to work through issues through established weekly conference calls and further training.

Total enrollment to-date: 38 subjects.

REPORTABLE OUTCOMES

(none). Trial is actively recruiting.

REFERENCES:

- Anger WK, Storzbach D, Binder LM, Campbell KA, Rohlman DS, McCauley L, Kovera CA, Davis KL. (1999) Neurobehavioral deficits in Persian Gulf veterans: evidence from a population-based study. Portland Environmental Hazards Research Center. *J Int Neuropsychol Soc*, 5(3):203-12.
- Chao LL, Abadjian L, Hlavin J, Meyerhoff DJ, Weiner MW. (2011, June 29). Effects of low-level sarin and cyclosarin exposure and Gulf War Illness on Brain Structure and Function: A study at 4T. *Neurotoxicology*.
- Chao LL, Rothlind JC, Cardenas VA, Meyerhoff DJ, Weiner MW. (2010). Effects of low-level exposure to sarin and cyclosarin during the 1991 Gulf War on brain function and brain structure in US veterans. *Neurotoxicology*, 31(5):493-501.
- Institute of Medicine (IOM). (2010). Gulf War and Health: Volume 8. Health Effects of Serving in the Gulf War. Washington, DC: *National Academies Press*.
- Kang H, Li B, Mahan C, Eisen S, Engel C. (2009). Health of US Veterans of 1991 Gulf War: A follow up Survey in 10 years. *JOEM*, 51(4): 1-10.
- Krengel M, Sullivan K. (2008). Neuropsychological Functioning in Gulf War veterans exposed to pesticides and pyridostigmine bromide. Fort Detrick, MD: *US Army Medical Research and Materiel Command*.
- Li B, Mahan CM, Kang HK, Eisen SA, Engel CC. (2011, July 27). Longitudinal Health Study of US 1991 Gulf War Veterans: Changes in Health Status at 10-Year Follow-up. *Am J Epidemiol*.
- Proctor SP, Heeren T, White RF, Wolfe J, Borgos MS, Davis JD, Pepper L, Clapp R, Sutker PB, Vasterling JJ, Ozonoff D. (1998). Health status of Persian Gulf War veterans: self-reported symptoms, environmental exposures and the effect of stress. *Int J Epidemiol*, 27(6):1000-10.
- Research Advisory Committee (RAC) on Gulf War Veterans' Illnesses. (2008). Gulf War illness and the Health of Gulf War Veterans: Scientific Findings and Recommendations. Washington, DC: US GPO.
- Sullivan K, Krengel M, Proctor SP, Devine S, Heeren T, White RF. (2003). Cognitive functioning in treatment-seeking Gulf War veterans: pyridostigmine bromide use and PTSD. *Journal of Psychopathology and Behavioral Assessment*. 25(2):95-103.
- Tillman GD, Green TA, Ferree TC, Calley CS, Maguire MJ, Briggs R, Hart J Jr, Haley RW, Kraut MA. (2010, Oct 15). Impaired response inhibition in ill Gulf War veterans. *J Neurol Sci*, 297(1-2):1-5.
- Toomey R, et al. (2009). Neuropsychological functioning of U.S. Gulf War veterans 10 years after the war. *Journal of the International Neuropsychological Society*, 15(5):717-29.