AWARD NUMBER: W81XWH-14-1-0403

TITLE: The Roles of the Bone Marrow Microenvironment in Controlling Tumor Dormancy

PRINCIPAL INVESTIGATOR: Yusuke Shiozawa, M.D., Ph.D.

CONTRACTING ORGANIZATION: Wake Forest University Health Sciences Winston Salem, NC 27157

REPORT DATE: October 2017

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE					Form Approved		
Public reporting burden for this	collection of information is estin	wing instructions, sear	. searching existing data sources, gathering and maintaining the				
data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the based of 0.000 (1990) and the bas							
4302. Respondents should be	aware that notwithstanding any	other provision of law, no perso	n shall be subject to any penalty	for failing to comply wit	h a collection of information if it does not display a currently		
1 REPORT DATE	EASE DO NOT RETURN YOU	R FORM TO THE ABOVE ADDE 2 REPORT TYPE	RESS.	3	DATES COVERED		
October 2017		Annual		8	Sep 2016 - 7 Sep 2017		
4. TITLE AND SUBTIT	LE			5a.	CONTRACT NUMBER		
The Poles of the B	one Marrow Microe	nvironment in Contr	olling Tumor Dorma				
			oning runor Donna	5b.	GRANT NUMBER		
				W	81XWH-14-1-0403		
				5c.	PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d.	PROJECT NUMBER		
Yusuke Shiozawa	MD. PhD			5e.	TASK NUMBER		
	, ,						
				5f.	WORK UNIT NUMBER		
E-Mail: vshiozaw@	wakehealth.edu						
7. PERFORMING ORC	SANIZATION NAME(S)	AND ADDRESS(ES)		8.1	PERFORMING ORGANIZATION REPORT		
				1	NUMBER		
Wake Forest Univ	ersity Health Sciend	ces					
Medical Center Blv	vd.						
Winston-Salem NO	C 27157						
9. SPONSORING / MC	NITORING AGENCY N	AME(S) AND ADDRES	S(ES)	10.	SPONSOR/MONITOR'S ACRONYM(S)		
U.S. Army Medica	I Research and Ma	teriel Command					
Fort Detrick, Maryland 21702-5012				11.	SPONSOR/MONITOR'S REPORT		
					NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT							
Approved for Public Release; Distribution Unlimited							
13. SUPPLEMENTARY NOTES							
14. ADSTRACT							
The nurnese of this study is to identify the mechanisms whereby the hone marrow microenvironment is involved in the							
regulation of tum	nio olday io lo lac or dormancy Aim1	will identify and e	volore how dissem	inated tumor	cells (DTCs) stay dormant for long		
neriods of time M	/e postulate that D	TCs drive the hone	marrow niche into	dormancy th	rough the GAS6 pathway Aim2 will		
determine how DT	Cs escape dorman	cv consequently re	ndering them more	suscentible to	the chemotherany		
		cy, consequently re			and chemotherapy.		
During this period	we found that a	fast arowing less d	ifferentiated osteob	last may con	tribute to growth of bone metastatic		
prostate cancer Ir	, we loand that a	loned a new and cr	ative in vitro cell cu	ilture system	that allows us to evaluate the effects		
of osteoblastic activation on the growth of disseminated prostate cancer cells							
or obcobladilo derivation on the growth of disseminated prostate cancer cens.							
15. SUBJECT TERMS	Dono mata-t!	Discominate - Lite	n coller Dar		ironmonti Tumor domocrani 0400		
Prostate Cancer;	Done metastasis;	Disseminated tumo	or cells; Bone mari	ow microenv	ironment, rumor dormancy; GAS6;		
BINIP2							
16. SECURITY CLASSIFICATION OF:			17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON		
			OF ABSTRACT	OF PAGES	USAMRMC		
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area		
			Unclassified	73	codej		
Unclassified	Unclassified	Unclassified					

Table of Contents

Page

1. Introduction	4
2. Keywords	4
3. Accomplishments	5
4. Impact	11
5. Changes/Problems	12
6. Products	13
7. Participants & Other Collaborating Organizations	16
8. Special Reporting Requirements	16
9. Appendices	17

1. INTRODUCTION:

Despite improvements in treatments for primary prostate cancer (PCa), bone metastasis remains a major cause of death in PCa patients. Several studies have shown that disseminated tumor cells (DTCs) shed from a primary tumor may lie dormant in distant tissues for long periods of time, retaining the potential for activation resulting in metastatic growth. Understanding the underlying mechanisms of metastasis is therefore crucial for effective treatment of this disease. Bone marrow has been well established as a regulatory site for hematopoietic function. In the marrow, hematopoietic stem cells (HSCs) are believed to localize to a specific microenvironment, the "niche", where they reside in a dormant state. Likewise, growing evidence has suggested that disseminated PCa also resides within the marrow niche. In fact, disseminated PCa uses similar mechanisms as HSCs in order to gain access to the marrow microenvironment, and DTCs target and displace HSCs, establishing metastatic foci within the hematopoietic niche. As a result, these cells parasitize the niche to become dormant, utilizing the mechanisms that keep HSCs in a dormant state. Although bone marrow is known as a fertile microenvironment ("soil") for metastatic tumor cells ("seed"), little is known about how dormancy is established or what leads to re-activation of the dormant cells. Therefore, we hypothesize that **once DTCs become dormant within the bone marrow niche, they stay dormant by stimulating the niche to remain dormant, and eventually escape from dormancy when the niche matures.**

To address our hypothesis the following aims are proposed:

Aim1: Determine the mechanisms whereby DTCs control the dormancy of the niche cells.

Sub hypothesis: DTCs drive the niche into dormancy via GAS6 signaling.

Aim2: Determine if the differentiation of the niche cells triggers the regrowth of DTCs.

Sub hypothesis: Dormant DTCs exit from dormancy when the niche is differentiated via BMP2 signaling.

The proposed studies will provide significant insight into the mechanisms whereby the bone marrow microenvironment is involved in regulation of tumor dormancy. Aim 1 allows us to identify and explore how DTCs stay dormant for long periods of time. We postulate that DTCs drive the bone marrow niche into dormancy through the GAS6 pathway. Aim2 will determine how DTCs escape dormancy, consequently rendering them more susceptible to the chemotherapy. Results from this work will lead to a greater understanding of niche aging effects on metastatic growth, and could result in valuable new treatment approaches.

2. KEYWORDS:

Prostate Cancer; Bone metastasis; Disseminated tumor cells; Bone marrow microenvironment; Tumor dormancy; GAS6; BMP2

3. ACCOMPLISHMENTS:

What were the major goals and objectives of the project?

The goal of this project is to understand the mechanisms of tumor dormancy and metastatic outgrowth of disseminated prostate cancer within the bone marrow microenvironment.

Task 1: Complete the grant transfer from University of Michigan to Wake Forest School of Medicine.

Months 1-3.

• Upon arrival at Wake Forest School of Medicine, the PI will seek to obtain the necessary approvals (IACUC, IRB, IBC) to complete the grant transfer, and then will initiate the proposed research as soon as possible (**Months 1-3**).

Task 2: Determine the mechanisms whereby DTCs control the dormancy of the niche cells.

Months 4-18.

• To determine the effects of GAS6 on the dormancy of niche cells *in vitro*, co-culture of bone marrow stromal cells (**BMSCs**) (pre-stained with DiD fluorescent dye) with either GAS6-downregulated PCa cells (**PCa**^{shGAS6}) or control PCa (**PCa**^{Control}) will be performed. At the termination of experiments, BMSCs will be harvested, and the retention of DiD dye will be measured with FACS (**Months 4-7**).

To further characterize the difference, gene and protein expression of proliferation markers and cell cycle status will be analyzed using those isolated BMSCs (**Months 7-9**).

• To determine the effects of GAS6 on the dormancy of niche cells *in vivo*, we will perform a vertebral body implant (vossicle) experiment. We will implant BrdU-incorporated vossicles directly injected with PCa^{shGAS6} or PCa^{Control} into immunocompromized mice, and then will determine the effects of GAS6 on the dormancy of the microenvironment by immunohistochemistry for BrdU (**Months 9-14**).

Additionally, using immunohistochemistry we will also visualize co-localization of PCa cells with the dormant microenvironment cells using these vossicles (Months 14-19).

Task 3: Determine if the differentiation of the niche cells triggers the regrowth of DTCs.

Months 19-36.

• To determine if the differentiation of the niche following exogenous BMP2 treatment stimulates the regrowth of DTCs *in vitro*, co-culture of BMSCs with G1-Red and SG2M-Cyan co-infected PCa cells will be performed. The differentiation of the niche, and the dormancy, proliferation, and cell

cycle status of PCa cells after treatment with recombinant mouse (**rm**) BMP2 will be analyzed (**Months 19-22**).

- To determine if the differentiation of the niche following the exogenous BMP2 treatment stimulates the regrowth of DTCs *in vivo*, we will implant vossicles directly injected with G1-Red and SG2M-Cyan co-infected PCa cells into immunocompromized mice. The differentiation of the niche, and the dormancy, proliferation, and cell cycle status of PCa cells after treatment with rm BMP2 will be analyzed (**Months 22-26**).
- To determine whether BMP2 expressed by DTCs is crucial for metastatic progression *in vitro*, coculture of BMSCs with BMP2-downregulated PCa (**PCa**^{shBMP2}), upregulated PCa (**PCa**^{BMP2OE}), or control PCa (**PCa**^{Control}) will be performed. Thereafter, the differentiation of the niche, and the dormancy, proliferation, and cell cycle status of PCa cells will be analyzed (**Months 27-30**)
- To determine whether BMP2 expressed by DTCs is crucial for metastatic progression *in vivo*, we will implant vossicles directly injected with PCa^{shBMP2}, PCa^{BMP2OE}, or PCa^{Control}. Thereafter, the differentiation of the niche, and the dormancy, proliferation, and cell cycle status of PCa cells will be analyzed (**Months 31-36**).

What was accomplished under these goals?

(~2016)

The Award transfer.

As of 03/01/15, thanks to receiving this Idea Development Award for Young Investigators, the PI, Dr. Yusuke Shiozawa started an independent faculty job as an Assistant Professor at Wake Forest School of Medicine. Upon his arrival at Wake Forest School of Medicine, the PI obtained the necessary institutional approvals (IACUC, IRB, IBC) and submitted the grant transfer request (06/11/2015) to gain approval from the Department of Defense for a transfer of the award from the University of Michigan to Wake Forest School of Medicine. As of 07/01/16, a transfer of the award from the University of Michigan to Wake Forest School of Medicine was completed.

The development of a mouse model to measure tumor growth and bone remodeling.

To evaluate the interaction between the bone marrow niche and disseminated PCa cells, we must be able to measure (i) the growth of bone metastatic PCa and (ii) bone remodeling within the same animal. To address this concern, we first attempted to establish an innovative and powerful mouse model. For this experiment, PCa cells (DU145) were inoculated intrafemorally into severe combined immunodeficient (SCID) mice to establish bone metastases. Thereafter, we measured changes in tumor growth [bioluminescent imaging (BLI), immunohistochemistry (IHC)] and bone remodeling (microCT, IHC). Using this strategy, we found (i) tumor growth by BLI, (ii) tumor burden in the marrow by histology, and decreased bone volume density and connective density in tumor-burdened mice on microCT.

The neuropeptide CGRP expressed by sensory neurons around bone influences PCa proliferation through CRLR/JNK pathway.

We recently reported that PCa cells parasitize the mechanisms whereby HSCs home to the marrow to gain

access to bone. Nerves are a major component of the microenvironment for HSCs, and are also involved in the metastatic process of PCa to bone. Yet whether the interactions between DTCs and sensory neurons in the bone play a crucial role in controlling later bone metastatic progression remains unclear. Interestingly, our preliminary data demonstrated that bone metastatic PCa increases neuronal hypertrophy of calcitonin-gene related peptide (CGRP)-expressing sensory nerves in the periosteum in a time-dependent manner. It has been demonstrated that levels of CGRP are increased in the serum of patients with advanced PCa compared to low-grade PCa, and that CGRP induces osteoblastic differentiation. Next, we wondered if CGRP affects PCa progression. We found that CGRP enhances proliferation of PCa cells *in vitro*. PCa patients with metastases express higher levels of CGRP receptor Calcitonin receptor like receptor (CRLR) (gene name CALCRL), compared to PCa patients without metastases. Additionally, CGRP activates JNK in PCa.

(2016-2017)

The fast growing less differentiated osteoblasts may contribute more to tumor growth, compared to mature osteoblasts *in vitro*.

To test the effects of differentiation status on osteoblast proliferation we used an MTT assay, as well as a live cell imaging device (Incucyte). We used the mouse calvarial preosteoblast cell line MC3T3-E1 transduced to constitutively express green fluorescent protein (eGFP) for these studies. We found that BMP2 stimulated a modest increase in osteoblast proliferation by (Fig. 1A) as well as expression of the differentiation osteoblast marker osteocalcin (Ocn) (Fig. 1C). We used bone mineralization media (BMM) consisting of 0.5mM L-Ascorbic acid and 2mM β-Glycerophosphate as a positive control for osteoblast differentiation and found that only this treatment increased osteoblast proliferation, when measured by Incucyte live cell imaging (Fig. 1B). We were unable to test this by MTT as ascorbic acid interferes with this assay. We also tested the effect of histamine dihydrochloride (HC) and found little to no effect on osteoblast proliferation and differentiation (Fig. 1). Strangely, we found that BMM had no effect on Ocn expression (Fig. 1C). Also, we were surprised to find that another osteoblast differentiation marker. osteopontin (**Opn**), actually decreased after treatment with BMM, BMP2 and HC (Fig. 1D).



Fig. 1: The effects of differentiation on osteoblast growth

MC3T3-eGFP were seeded into 96-well plates and serum starved for 24 hours before treatment with 25 ng/mL BMP2, 100nM histamine dihydrochloride (HC), or 0.1% DMSO (Vehicle). (A) Relative cell numbers measured at 48 and 72 hours with MTT (500µg/mL) by optical density (OD) at 560nm minus background at 650nm. (B) Confluency change measured for 72 hours using the Incucvte after addition of bone mineralization media (BMM) supplemented with 0.5mM L-Ascorbic acid and 2mM β-Glycerophosphate, or BMP2, HC, or Vehicle. (C) Osteocalcin expression levels were tested by qPCR after 14 day treatments of BMM, BMP2, HC, or Vehicle in complete media (10% FBS). (D) Osteopontin levels were also measured (data presented as fold change using the $\Delta\Delta$ CT method with Gapdh as reference gene). All data presented as mean \pm SD, *P<0.05, **P<0.005, ***P<0.0005. ****P<0.0005 (Two-way ANOVA Bonferroni with multiple comparisons)

Next we tested the effects of these treatments on the proliferation of the PCa cell lines PC3 and DU145 alone and in coculture with MC3T3-E1. We used a luciferase reporter to monitor tumor cell growth via bioluminescent imaging (BLI) using the IVIS system. Not surprisingly, we found that PC3 and DU145 grew slower in coculture with osteoblasts (**Fig. 2A&D**). Additionally, we found that BMM, BMP2 and HC had no effect on PC3 growth when cultured alone (**Fig. 2B**), whereas BMM slightly increased DU145 growth in the

same conditions (Fig. 2E). A different trend was revealed when the same treatments were tested on the cocultures. There was a significant reduction in PC3 and DU145 growth only when treated with BMM (Fig. 2C&E). We performed the test again using the DiD membrane dye method of tracking cellular divisions and tumor cell dormancy by flow cytometry. We found that coculture of PC3 and DU145 with MC3T3-E1 led to higher retention of DiD, indicating slower cancer cell growth (Fig. **2G&H**). Treatment with BMM resulted in even higher retention of DiD in PC3 and DU145 (Fig. 2G&H). Treatment of BMP2 caused a similar effect only in DU145 (Fig. In addition, we **2H**). found that the percentage of PC3 and DU145 cells in the cocultures decreased after treatments with BMM and BMP2 (Fig.



Fig. 2: The effects of osteoblast differentiation on prostate cancer growth DiD-labelled, luciferase expressing PC3 (PC3-luc) and DU145 (DU145-luc) cells were sparsely seeded on top of a monolayer of MC3T3-E1-eGFP (Coculture), or directly into 24well tissue culture plates (Alone). PCa cell growth was measured by BLI and media replaced with BMM, BMP2, HC, or Vehicle on Day 0, 3, and 5. Cultures harvested on Day 6. (**A&D**) Luciferase signals from the Vehicle treated PCa Alone and Coculture plates (One-way ANOVA). (**B&E**) Treatment effects from the Alone and (**C&F**) Coculture plates (Two-way ANOVA with Bonferroni multiple comparisons). (**G&H**) BD Accuri flow cytometry for mean fluorescence intensity of DiD in the eGFP negative cell population and (**I**) Percent DiD positive, eGFP negative PCa cells in Coculture (Two-way ANOVA with Bonferroni multiple comparisons). All data presented as mean \pm SD, *P<0.05, **P<0.005, ***P<0.0005

2I), although we cannot say whether this is because cancer cell growth was slowed, or osteoblast proliferation was increased. These findings made us reconsider the initial hypothesis as the differentiation status or maturity of osteoblasts may actually slow tumor growth, contributing to dormancy. However, the unexpected result of BMM having no significant effect on Ocn expression, as well as decreasing Opn expression made us question the validity of MC3T3-E1 as a reliable model of osteoblast differentiation.

We thought back to an interesting observation when we perform intracardiac injection of tumor cells as a model of metastasis. Often times these animals develop tumors in the mandible, and sometimes only in the

mandible, as observed by BLI (**Fig. 3A**). It is not uncommon that these mandible mettastases grow faster than those in the hind limbs of the same animal (**Fig. 3B&C**). It may be that in these cases there was higher seeding of tumor cells in the mandible than the hind limbs, or as we suspect, the mouse mandible microenvironment is less dormant and allows for faster tumor cell growth.

To test this we harvested adult osteoblasts from the mouse mandible and hind limbs using an explant model. Briefly, adult male C57BL/6 mice were sacrificed according to approved protocols and hind limbs and mandibles dissected and muscles completely removed. The epiphyses from the femurs and tibias were cut and the marrow was flushed from the diaphyses and discarded. The empty diaphyses were placed in a tissue culture dish and minced into smaller pieces using a scalpel. The molars, incisors and dental pulp of the mandibles were removed and the remaining pieces (ascending ramus containing the coronoid, condylar and angular processes) were placed in a dish and minced. A 10 minute trypsin digestion was performed on all the bone pieces and discarded. Complete media containing 10nM dexamethasone was added to select for osteoblast outgrowth. The bone cells were allowed to migrate out of the bone and mandible pieces and expand for two weeks before experimentation.



Fig. 3: Mandible metastasis after intracardiac injection of prostate cancer

SCID hairless outbread (SHO) mice were given intracardiac injections of 1 million PC3-luc cells and tumor growth was measured by BLI weekly. (A) Mouse at day 100 with single large met in the mandible by BLI. (B) Mouse showing both mandible and hind limb tumor growth at days 8, 15, 22, and 29; and (C) quantification of the BLI signals.

We first compared the proliferation of the mandible osteoblasts (MaOB) and the hind limb osteoblasts (HLOB) using an MTT assay. Indeed, we found the MaOB to be much more proliferative than the HLOB (**Fig. 4A**). We tested for osteoblast differentiation markers and found the basal expression of Ocn and Opn to be less in the MaOB than the HLOB, however after treatment with BMM we found significantly higher expression of Ocn and a trend toward increased Opn in the MaOB (**Fig. 4B&C**). These findings suggested to us that the MaOB are more proliferative and less mature, but with a higher potential for differentiation than the HLOB. Finally, we hypothesized that cancer cell proliferation would increase when cocultured with the MaOB, if the microenvironment of the mandible is truly involved in the phenomena observed in Figure 3. Indeed, when we performed this experiment we found that both PC3 and DU145 grew significantly faster in coculture with MaOB compared to HLOB (**Fig. 4D&E**). Similar to the MC3T3-1 studies in Fig. 2, we found that treatment with BMM decreased this proliferation effect in PC3 (**Fig. 4D**). Again, the results suggest that a more mature osteoblast may contribute more to tumor dormancy than a fast growing less differentiated osteoblast.

AWARD: PC130359

TITLE: The Role of the Bone Marrow Microenvironment in Controlling Tumor Dormancy PI: Yusuke Shiozawa, M.D., Ph.D.

What opportunities for training and professional development did the project provide?

(~2016)

Thanks to receiving an Idea Development Award for Young Investigators, the PI obtained independent status at Wake Forest School of Medicine with lab space, office space, and start-up costs provided.

Thanks to the Department of Defense, the PI attended IMPaCT Young Investigator Meeting, Baltimore, MD, USA, August 4-5, 2016.

The PI was chosen for the 2016-2018 cohort of the Wake Forest Clinical and Translational Science Institute's Translational Scholar Academy, which supports the scientific and career development of early-stage investigators.

(2016-2017)

The PI was invited to attend The U.S. Bone and Joint Initiative (USBJI) and Bone and Joint Canada (BJC) Young Investigator Initiative Program will be held at Toronto, Canada on 11/10/17-11/12-17.

How were the results disseminated to communities of interest?

There is nothing to report at this time.

What do you plan to do during the next reporting period to accomplish the goals and objectives?



(OD) at 560nm minus background at 650nm, normalized to Day 1. (B) Osteocalcin expression levels were tested by qPCR after 14 day treatments of BMM or Vehicle in complete media (10% FBS). (C) Osteopontin levels were also measured (data presented as fold change using the $\Delta\Delta$ CT method with Gapdh as reference gene). (D&E) PC3-luc and DU145-luc were sparsely seeded on top of a monolayer of MaOB or HLOB in a 96-well plate and PCa cell growth was measured by BLI and media replaced with BMM, BMP2, HC, or Vehicle on Day 1, 3, and 5. All data presented as mean \pm SD, *P<0.05, **P<0.005, ***P<0.0005, ****P<0.0005 (Two-way ANOVA with Bonferroni multiple comparisons).

In the next year of the award, we will characterize the differences between the mandible and hind limb osteoblasts in order to determine the detailed molecular mechanisms whereby active osteoblasts influence the progression of bone metastatic prostate cancer (Aim 2). We will also further elucidate the role of osteoblastic activation in the progression of prostate cancer bone metastasis *in vivo* (Aim 2). In addition, we will continue to pursue the effects of the disseminated tumor cells on the dormancy of the bone marrow niche (Aim 1).

4. IMPACT:

What was the impact on the development of the principal discipline(s) of the project?

(~2016)

We developed an innovative and powerful mouse models that enable us to measure <u>within the same animal</u>: (i) the growth of bone metastatic PCa and (ii) bone remodeling.

(2016-2017)

We developed a new and creative tool (mandible vs. hind limb osteoblast) to test the hypothesis that active osteoblasts enhance the re-growth of cancer in the bone.

Using this technique, we found that a fast growing less differentiated osteoblast may contribute to growth of bone metastatic prostate cancer.

What was the impact on other disciplines?

There is nothing to report at this time.

What was the impact on technology transfer?

There is nothing to report at this time.

What was the impact on society beyond science and technology?

There is nothing to report at this time.

5. CHANGES/PROBLEMS:

Changes in approach and reasons for change

Nothing to report.

Actual or anticipated problems or delays and actions or plans to resolve them

Nothing to report.

Changes that have a significant impact on expenditures

Nothing to report.

Significant changes in use or care of human subjects, vertebrate animals, biohazards, and/or select agents

Nothing to report.

6. PRODUCTS:

Publications, conference papers, and presentations

Journal Publications

Peer reviewed journal

(~2016)

- Seib FP, Berry JE, Shiozawa Y, Taichman RS, Kaplan DL. Tissue engineering a surrogate niche for metastatic cancer cells. Biomaterials. 2015;51:313-9. PMID: 25771021. PMCID: PMC4367489. Status of Publication: Published Acknowledgement of federal support: Yes
- Zalucha JL, Jung Y, Joseph J, Wang J, Berry JE, Shiozawa Y, Taichman RS. The Role of Osteoclasts in Early Dissemination of Prostate Cancer Tumor Cells. J Cancer Stem Cell Res. 2015;3:e1005. PMID: 26097863. PMCID: PMC4469294.
 Status of Publication: Published Acknowledgement of federal support: Yes
- Shiozawa Y, Berry JE, Eber MR, Jung Y, Yumoto K, Cackowski FC, Yoon HJ, Parsana P, Mehra R, Wang J, McGee S, Lee E, Nagrath S, Pienta KJ, Taichman RS. The Marrow Niche Controls The Cancer Stem Cell Phenotype Of Disseminated Prostate Cancer. Oncotarget. In Press. PMID: 27172799. PMCID: PMC5173053. Status of Publication: Published Acknowledgement of federal support: Yes
- Sharma S, Xing F, Liu Y, Wu K, Said N, Pochampally R, Shiozawa Y, Lin HK, Balaji KC, Watabe K. Secreted Protein Acidic and Rich in Cysteine (SPARC) Mediates Metastatic Dormancy of Prostate Cancer in the Bone. J Biol Chem. 2016;291:19351-63. PMID: 27422817. PMCID: PCM5016675. Status of Publication: Published Acknowledgement of federal support: No

(2016-2017)

 Cackowski F, Eber MR, Rhee J, Decker A, Yumoto K, Berry JE, Lee E, Shiozawa Y, Jung Y, Aguirre-Ghiso JA, Taichman RS. Mer Tyrosine Kinase Regulates Disseminated Prostate Cancer Cellular Dormancy. J Cell Biochem. 2017;118:891-902. PMID: 27753136. PMCID: PMC5296338. Status of Publication: Published Acknowledgement of federal support: Yes

Kast RE, Skuli N, Cos S, Karpel-Massler G, Shiozawa Y, Goshen R, Halatsch ME. The ABC7 regimen: a new approach to metastatic breast cancer using seven common drugs to inhibit epithelial-to-mesenchymal transition and augment capecitabine efficacy. Breast Cancer (Dove Med Press). 2017;9:495-514. PMID: 28744157. PMCID: PMC5513700 Status of Publication: Published Acknowledgement of federal support: Yes

Invited reviews

(~2016)

- Shiozawa Y^{*}, Eber MR, Berry JE, Taichman RS^{*}. Bone marrow as a metastatic niche for disseminated tumor cells from solid tumors. BoneKEy Rep. 2015;4:689. PMID: 26029360. PMCID: PMC4440229. (* Co-corresponding authors) Status of Publication: Published Acknowledgement of federal support: Yes
- Dai J, Hensel J, Wang N, Kruithof-de Julio M, Shiozawa Y. Mouse models for studying prostate cancer bone metastasis. BoneKEy Rep. 2016;5:777. PMID: 26916039. PMCID: PMC4757481. Status of Publication: Published Acknowledgement of federal support: Yes
- Tsuzuki S, Park SH, Eber MR, Peters CM, Shiozawa Y. Skeletal complications in cancer patients with bone metastases. Int J Urol. In Press. PMID: 27488133. PMCID: In Progress. Status of Publication: Accepted Acknowledgement of federal support: Yes
 - The Figure 1 is chosen as the Cover Figure of Int J Urol Vol. 23 No. 10

<u>Book</u>

(~2016)

 Miler SF, Thomas CY, Shiozawa Y. (2016) Molecular involvement of the bone marrow microenvironment in bone metastasis. In Ahmad A (Ed.), *Introduction to Cancer Metastasis*. In Press. Philadelphia, Elsevier. Status of Publication: Accepted Acknowledgement of federal support: Yes

Presentation

(~2016)

1. Tsuzuki S, Eber MR, Miler SF, Park SH, Widner DB, **Shiozawa Y**. The effects of neuropeptides on the prostate cancer progression. IMPaCT Young Investigator Meeting, Baltimore, MD, USA, August 4-5, 2016. Poster.

(2016-2017)

2. Mamo M, Park SH, Eber MR, Widner DB, Tsuzuki S, Warren CC, Strickland TB, Peters CM, **Shiozawa Y**. The Interaction between Disseminated Prostate Cancer Cells and Sensory Nerves. the 2016 Annual Biomedical Research Conference for Minority Students (ABRCMS), Tampa, FL, USA, November 9-12, 2016. Poster.

Website(s) or other Internet site(s)

Nothing to report.

Technologies or techniques

Nothing to report.

Inventions, patent applications, and/or licenses

Nothing to report.

Other products

Nothing to report.

7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

What individuals have worked on the project?

Name: Yusuke Shiozawa Project Role: PI Researcher Identifier (e.g. ORCID ID): orcid.org/0000-0001-9814-9230 Nearest person month worked: 2.4 Contribution to Project: Dr. Shiozawa provides oversight of the entire program and development and implementation of all policies, procedures, and processes. In this role, Dr. Shiozawa is responsible for the implementation of the specific aims, and for ensuring that systems are in place to guarantee institutional compliance with US laws, including biosafety and animal research, data and facilities. Dr. Shiozawa supervises other personnel on the project to ensure timely and effective studies. Funding Support: National Cancer Institution, Department of Defense

Has there been a change in the other active support of the PD/PI(s) or senior/key personnel since the last reporting period?

Nothing to report.

What other organizations have been involved as partners?

Nothing to report.

8. SPECIAL REPORTING REQUIREMENTS:

N/A.

9. APPENDICES:

The original copies of manuscript and a curriculum vitae are attached.



Mer Tyrosine Kinase Regulates Disseminated Prostate Cancer Cellular Dormancy

Frank C. Cackowski,^{1,2} Matthew R. Eber,^{1,3} James Rhee,¹ Ann M. Decker,¹ Kenji Yumoto,¹ Janice E. Berry,^{1§} Eunsohl Lee,¹ Yusuke Shiozawa,³ Younghun Jung,¹ Julio A. Aguirre-Ghiso,⁴ and Russell S. Taichman¹*

¹Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, Michigan ²Division of Hematology and Oncology, Department of Medicine, University of Michigan School of Medicine, Ann Arbor, Michigan

³Department of Cancer Biology and Comprehensive Cancer Center, Wake Forest University School of Medicine, Winston-Salem, North Carolina

⁴Division of Hematology and Oncology, Tisch Cancer Institute, Departments of Medicine, Otolaryngology, and Black Family Stem Cell Institute, Icahn School of Medicine at Mount Sinai, New York, New York

ABSTRACT

Many prostate cancer (PCa) recurrences are thought to be due to reactivation of disseminated tumor cells (DTCs). We previously found a role of the TAM family of receptor tyrosine kinases TYRO3, AXL, and MERTK in PCa dormancy regulation. However, the mechanism and contributions of the individual TAM receptors is largely unknown. Knockdown of MERTK, but not AXL or TYRO3 by shRNA in PCa cells induced a decreased ratio of P-Erk1/2 to P-p38, increased expression of p27, NR2F1, SOX2, and NANOG, induced higher levels of histone H3K9me3 and H3K27me3, and induced a G1/G0 arrest, all of which are associated with dormancy. Similar effects were also observed with siRNA. Most importantly, knockdown of MERTK in PCa cells increased metastasis free survival in an intra-cardiac injection mouse xenograft model. MERTK knockdown also failed to inhibit PCa growth in vitro and subcutaneous growth in vivo, which suggests that MERTK has specificity for dormancy regulation or requires a signal from the PCa microenvironment. The effects of MERTK on the cell cycle and histone methylation were reversed by p38 inhibitor SB203580, which indicates the importance of MAP kinases for MERTK dormancy regulation. Overall, this study shows that MERTK stimulates PCa dormancy escape through a MAP kinase dependent mechanism, also involving p27, pluripotency transcription factors, and histone methylation. J. Cell. Biochem. 118: 891–902, 2017. © 2016 Wiley Periodicals, Inc.

KEY WORDS: MERTK; AXL; TYRO3; PROSTATE CANCER; DORMANCY; DISSEMINATED TUMOR CELL

P rostate cancer patients often have long time periods between curative intent surgery or radiation therapy until the time of biochemical recurrence or metastatic disease visible with current imaging, which marks incurable disease with current treatment options. For example, in a large series of patients treated with radical prostatectomy, nearly 20% recurrences occurred at least 5 years after surgery [Amling et al., 2000]. Greater than half of prostate cancer patients with no evidence of disease, soon after radical prostatectomy were found to have disseminated prostate tumor cells (DTCs) in their bone marrow, which are thought to be a major source of distant

recurrences [Morgan et al., 2009]. This finding implies that many of these tumor cells die, never grow, or grow very slowly. Many investigators refer to this ability of cancer cells to remain viable but not have detectable growth as "cellular dormancy."

There is significant interest in regulators of cancer cellular dormancy. Several studies have identified a low ratio of phosphorylated MAPK3/MAPK1 (Erk 1/2) to phosphorylated MAPK14 (p38) as marking dormant tumor cells. TGFB2 (TGF- β 2) was proposed to be the major ligand responsible for dormant behavior of head and neck squamous cell carcinoma cells. The cell cycle inhibitor CDKN1B

[§]Deceased on February 4, 2016.

Grant sponsor: NIH/NCI; Grant number: P01-CA093900; Grant sponsor: NIH/NCI Tumor Microenvironment Network; Grant number: U54-CA163124; Grant sponsor: Department of Defense; Grant numbers: W81XWH-14-1-0403, W81XWH-15-1-0637, W81XWH-15-1-0413; Grant sponsor: NIH/NCI T32; Grant number: 5T32CA009357-32; Grant sponsor: Prostate Cancer Foundation Grant (PCF); Grant number: 2016CHAL1503. *Correspondence to: Russell S. Taichman, D.M.D., D.M.Sc., University of Michigan School of Dentistry, 1011 North University Avenue, Ann Arbor, MI 48109. E-mail: rtaich@umich.edu

Manuscript Received: 17 October 2016; Manuscript Accepted: 17 October 2016

Accepted manuscript online in Wiley Online Library (wileyonlinelibrary.com): 17 October 2016 DOI 10.1002/jcb.25768 • © 2016 Wiley Periodicals, Inc.

891

(p27) and transcription factor BHLHE41 (DEC2) were implicated as nuclear signals [Bragado et al., 2013]. More recently, pluripotency associated transcription factors NR2F1, SOX2, SOX9, NANOG, and RARB were identified as transcriptional regulators of dormancy in head and neck, prostate and breast cancers [Sosa et al., 2015]. Similarly, others have shown that TGF- β family member BMP7 maintains prostate cancer dormancy through autocrine SPARC [Kobayashi et al., 2011; Sharma et al., 2016]. A role of the epigenome in regulating cellular dormancy is also becoming apparent. Histone H3 tri-methylated lysine 9 and tri-methylated lysine 27 were shown to identify and to be important for dormant cells, primarily in head and neck cancer [Sosa et al., 2015].

Our group has established a role of the TYRO3, AXL, and MERTK (TAM) family of receptors and one of their ligands, growth arrestspecific 6 (GAS6), in regulation of prostate cancer cell dormancy in the bone marrow [Shiozawa et al., 2010; Jung et al., 2012; Taichman et al., 2013]. We also found that GAS6 and MERTK are important for cancer stem like cell formation [Jung et al., 2016; Shiozawa et al., 2016]. This receptor family has an established role in the regulation of the innate immune system, but more recently has been shown to be important for cancer growth and metastasis as well. For example, MERTK was recently identified in a screen of wild-type kinases as a mediator of prostate cancer metastasis [Faltermeier et al., 2015]. The TAM family of receptors have a high degree of homology, but have been shown to have different functions, which might relate to differences in ligand binding affinities and downstream pathways [Graham and DeRyckere, 2014]. There are at least four vitamin K dependent γ -carboxyalated protein ligands that bind to at least one of the TAM receptors including GAS6, Protein S (PROS1), Tubby (TUB), and Tubby like protein 1 (TULP1) [Caberoy, 2010]. We found that GAS6 decreased prostate cancer proliferation and protected the cells from chemotherapy induced apoptosis [Shiozawa et al., 2010; Lee et al., 2016]. We also found that prostate cancer bone metastases grew larger in the absence of GAS6 [Jung et al., 2012].

However, these studies did not identify which of the TAM receptors are responsible for the ability of GAS6 to slow prostate cancer growth, while also preventing apoptosis-findings which are consistent with cellular dormancy. However, the role of TYRO3 and MERTK remained unclear. To begin to answer this question, we previously studied the relative expression level TYRO3 and AXL in prostate cancer primary tumors, DTCs, and in gross metastases. However, MERTK was not included in these studies. We found that TYRO3 was expressed highly in the primary tumors, but that AXL was expressed highly in disseminated but dormant disease [Taichman et al., 2013]. Based on these results, we hypothesized that TYR03 might play a role when prostate cancer was actively growing but that AXL might play a role when it is dormant. However, these studies did not include experiments to test this hypothesis further than gene expression. Most recently, in work that is currently in press, we reported that AXL is required for TGF-B2 to induce prostate cancer dormancy [Yumoto et al., 2016]. However, the contributions of MERTK and TYRO3 remain unclear.

In the current study, we took an unbiased approach to discern which of the TAM receptors, including MERTK, are required for prostate cancer dormancy escape. With shRNA and siRNA technology, we knocked down the expression of each of the three receptors in three different prostate cancer cell lines. We found that loss of MERTK, but not the other receptors decreased the ratio of P-Erk to P-p38, increased the expression of p27 and pluripotency associated transcription factors, increased the levels of dormancy associated histone H3 and caused accumulation of cells in the G1 and G0 phases of the cell cycle, and decreased apoptosis, all of which characterize dormant cells. Importantly, the effect of MERTK on the cell cycle and histone H3 post-translational modifications was reversed by altering MAP kinase signaling with a p38 inhibitor. We also found that MERTK knockdown increased metastasis free survival in an intra-cardiac injection mouse prostate cancer xenograft model, but did not inhibit in vitro cell growth or subcutaneous tumor growth showing that it did not compromise global growth characteristics. Thus, our studies implicate MERTK in stimulation of prostate cancer dormancy escape by a mechanism particular to the metastatic microenvironment and involving MAP kinases.

MATERIALS AND METHODS

CELL CULTURE

Human PCa cell lines, PC3, Du145, and LNCaP C4-2B (C4-2B) were obtained from American Type Culture Collection (Rockville, MD [PC3 and Du145]) and UroCor (Oklahoma City, OK [C4-2B]). PCa cells were maintained in RPMI 1640 with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (P/S) in a humidified incubator with 5% CO₂. For in vitro assays, unless indicated otherwise, cells were seeded at a density 1×10^5 /ml, allowed to rest for 1 day in 10% serum and then changed to reduced serum concentrations as indicated. For the p38 inhibitor experiments, cells were first cultured for 14 days under routine conditions with 10% serum and 5 μ M SB203580 (EMD Millipore #A8254) dissolved in DMSO or 0.05% DMSO control.

TAM RECEPTOR STABLE shRNA KNOCKDOWNS

GFP and luciferase expressing PCa cell lines (PC3^{*GFP*}, Du145^{*GFP*}, and C4-2B ^{*GFP*} cells) were first established by lentiviral transduction. Stable knockdowns of the TAM receptors (TYR03, AXL, and MERTK) were then generated by lentiviral infection. Lentiviruses were constructed by the University of Michigan Vector Core using pGIPZ lentiviral vectors containing either a shRNA targeting one of the TAM receptors or a nonsilencing (shControl) shRNA (Open Biosystems). Stable lines were selected with puromycin. Knockdown of greater than 80% was verified by Western blotting and qRT-PCR. qRT-PCR gene expression data are presented as mean \pm SEM of independent cultures.

MerTK TRANSIENT siRNA KNOCKDOWNS

siRNAs targeting MerTK (# s20474, s20473, and s20472) and control siRNA (siControl) (# 4390843) were purchased from Thermo-Fisher Scientific. Transient transfection in C4-2B and PC3 cells was performed using 10 mM of each siRNA with Lipofectamine RNAiMAX reagent (Thermo-Fisher) using the reverse transfection protocol, followed by 3 days incubation. Knockdown was verified by real time qRT-PCR. Data are presented as mean \pm SEM of triplicate PCR reactions.

WESTERN BLOTTING

Cells were serum starved overnight unless indicated otherwise. Lysates were prepared in cOmplete lysis M (Roche #04 719 956 001) supplemented with proteinase inhibitor Mini cOmplete Tablets (Roche #04705378) and phosphatase inhibitor PhosSTOP EASYpack Tablets (Roche #04 906 837 001). Protein concentration was determined by the BCA method. Twenty micrograms of total protein was added per lane of 4-20% reducing SDS polyacrylamide Tris-Glycine gels after sample preparation in Laemmli sample buffer. The samples were transferred to PVDF membranes and blocked for 1 h in 5% dry milk in TBS with 0.1% Tween-20 (TBST). Antibodies for phosphorylated proteins were applied at 4°C overnight in 5% BSA TBST, washed, and visualized with a horseradish peroxidase conjugated anti-rabbit IgG secondary antibody (Cell Signalling #7074S) and SuperSignal West Dura Chemiluminescent Substrate (Thermo Scientific #34075). Images were acquired with a ChemiDoc Touch imager (BioRad). Membranes were stripped with Restore PLUS stripping buffer (Thermo Scientific #46430). They were blocked and re-probed for antibodies to total proteins, and again striped and reprobed for GAPDH or β -actin to normalize for protein loading. All primary antibodies were monoclonal rabbit from Cell Signaling Technology. Catalog numbers and dilutions were as follows; Phosphorylated-Erk 1/2 (P-Erk) Y204(#4377S, diluted 1:500), total Erk (#4695, 1:500), Phospho-p38 (P-p38) T190/Y182 (#4511, 1:500), total p38, (#9212, total Axl (#4939, 1:500), total Tyro3 (#5585, 1:500), total MerTK (#4319, 1:500), Sox2 (# 3579, 1:500), Caspase-9 (#9502, 1:1000), $\beta\text{-actin}$ (#4970, 1:2000), and GAPDH (#2118, 1:2000). Images representative of biological replicates are shown and cropped for presentation. For P-Erk, P-p38, and p27 quantification, images from five independent experiments were quantified relative to each vehicle treated scrambled shRNA control with BioRad ImageLab software and then normalized to housekeeping gene expression. The P-Erk to P-p38 ratio was obtained by dividing the normalized P-Erk and P-p38 values for each independent experiment. All data are shown as fold change from control.

REAL TIME REVERSE TRANSCRIPTASE PCR (qRT-PCR)

Cells were lysed and RNA was harvested using the Qiagen RNeasy kit followed by reverse transcription using Invitrogen SuperScript II Reverse Transcriptase. Real time qPCR was performed using TaqMan Universal PCR Master Mix and Gene Expression Assays on a Applied Biosystems ViiA 7 instrument. TaqMan MGB probes (Applied Biosystems) were as follows: MERTK (Hs00179024_m1), p27/CDKN1B (Hs00153277), SOX2 (Hs01053049_s1), and NANOG (Hs02387400). We designed primers and a probe to specifically detect NR2F1/TFCOUP1: forward; CAAAGCCATCGTGCTGTTCAC, reverse; CCTGCAGGCTCTCGATGT, and probe; TCA-GACGCCTGTGGCCTG. \beta-actin (Hs01060665_g1) was used as an internal control for the normalization of target gene expression.

FLOW CYTOMETRY FOR HISTONE POST-TRANSLATIONAL MODIFICATIONS AND Ki67

Cell pellets were fixed and permeabilized with dropwise addition of 1 ml of cold 70% ethanol and then incubated overnight. All steps were at 4° C or on ice. Samples were then washed, blocked, and

incubated for 1 h in flow buffer (PBS with 2% FBS and 2 mM EDTA) with each of the following antibodies; Alexa 647 conjugated rabbit anti histone H3 tri-methylated lysine 27 diluted 1:50 (Cell Signaling Technology #12158), unconjugated rabbit polyclonal anti histone H3 tri-methylated lysine 9 (Abcam #8898), or APC conjugated rabbit anti-human Ki-67 antibody (Biolegend #350513). Cells were washed twice with flow buffer. The unconjugated histone H3 tri-methylated lysine 9 antibody was detected with an Alexa 647 conjugated antirabbit IgG diluted 1:250 (Cell Signaling Technology #4414). Data were acquired with a three laser (405, 488, and 640 nm) Becton Dickinson FACS Aria IIu flow cytometer. Gating was forward scatter versus side scatter, single cells (linear on FSC-A vs. FSC-H), then the 670/30 filter (APC or Alexa 647) versus forward scatter or histogram. An isotype control antibody was used for setting the gates. Negative and dim cells were selected for methylated histones. Negative cells were selected for Ki67. Data are presented as representative plots or mean \pm SEM of triplicate wells from replicate experiments.

CELL CYCLE ANALYSIS

Cells were cultured for 3 days as indicated and pulsed with 10μ M bromodeoxyuridine (BrdU) for 30 min. The cells were collected with trypsin as necessary and then fixed and stained for total DNA with 7-AAD and BrdU incorporated into DNA using the Becton Dickinson APC BrdU flow kit (#552598). Data were acquired with a Becton Dickinson FACS Aria IIu flow cytometer. Gating was forward scatter versus side scatter, single cells (linear on FSC-A vs. FSC-H), then APC (BrdU) versus 7-AAD (DNA).

LEFT VENTRICLE INTRACARDIAC INJECTION XENOGRAFT MODEL OF PROSTATE CANCER METASTASIS

Stable shRNA infected PC3^{GFP} (1×10^6) or Du145^{GFP} (2×10^5) prostate cancer cells were suspended in 100 µl of PBS and injected into male CB.17. SCID mice (6-8 weeks of age: Charles River Labs) by left ventricle intracardiac injection. For analysis of metastasis free survival, bioluminescence images were acquired after injection of luciferin twice weekly using a PerkinElmer IVIS 2000 system. Animals that had a large portion of the signal in the lungs (indicative of a right ventricle injection) were removed from analysis a priori. After removing mice that had a right ventricular injections or did not survive the procedure, the following numbers of animals were analyzed; PC3 shControl; 6, PC3 shMER; 7, Du145 shControl; 20, and Du145 shMER; 18. Time to metastasis formation visible by bioluminescence (or death in rare cases) was then determined from the images. The data were analyzed by Kaplan-Meier analysis. For analysis of transit to the bone marrow, different mice, five mice per group, were sacrificed 24 h after tumor cell injection, and their pelvis, femora, and tibiae were harvested. The bones were crushed with a mortar and pestle and strained to remove debris. All steps used PBS buffer with 2% FBS unless otherwise noted. Cells were first depleted of mouse cells with a Mouse Cell Depletion Kit magnetic labeling system (Miltenyi Biotec # 130-104-694) and anti-Biotin MicroBeads and an AutoMACS machine (Miltenyi Biotec). The enriched cells were incubated with an APC-Cy7 conjugated anti-HLA-ABC antibody (BioLegend #311426) and a PerCP-Cy5.5 conjugated anti-mouse lineage cocktail (CD3e, CD11b, B220, Ter-119, Ly-6G, and Ly-6C) (BD Biosciences #561317), for an hour at 4°C, washed and resuspended in

PBS with 2% FBS, 2 mM EDTA and 0.5 μ g/ml DAPI. Thereafter, the percentage of disseminated prostate cancer cells (DTCs) was determined by gating on single, viable, lineage negative, HLA⁺ cells with a FACSAria IIu flow cytometer. Mice injected with PBS rather than PCa cells were used as a negative control for flow cytometry. Data represents three independent experiments.

PROSTATE CANCER SUBCUTANEOUS TUMOR MODEL

One million prostate cancer cells suspended in 50 μ l complete media were mixed with an equal volume of cold collagen solution and then slowly injected under the skin of the back of SCID mice; five mice per group. Bioluminescence images were acquired weekly. Animals were sacrificed before tumors grew to 1 cm³. All experimental procedures were approved by the University of Michigan Committee for the Use and Care of Animals.

PROSTATE CANCER CELL VIABILITY/MTS ASSAY

Prostate cancer cells were seeded at 2000 cells per well in 96 well plates and rested for 1 day in 10% FCS RPMI media. The media was subsequently changed to the indicated serum concentrations and the cells were cultured for an additional 3 days. The total viable cell number was then assayed with the Cell Titer Aqueous One Solution MTS Proliferation Assay System (Promega #G3580) by absorbance at 490 nm. Data represent means of three independent experiments.

STATISTICAL ANALYSES

The type I error rate (α) was set to 0.05 for all analyses. Two-sample, two-tailed Student's *t*-tests were used to compare means of two groups. One-way repeated measures analysis of variance (ANOVA) with Bonferroni post-hoc testing was used for data normalized to housekeeping genes (blots and PCR). Standard one-way ANOVA with Tukey's Honest Significant Difference post-hoc testing was used for multiple comparisons in other experiments. The Log-rank test was used for Kaplan–Meier survival analyses. Growth curves for subcutaneous tumors were analyzed with a mixed design (split plot) ANOVA with repeated measures. All analyses were conducted with SPSS software, except for *t*-tests, which were performed in Microsoft Excel.

RESULTS

MERTK KNOCKDOWN CAUSES DORMANCY ASSOCIATED CHANGES IN MAPK ACTIVITY AND p27 EXPRESSION

To study the importance of TAM signaling on dormancy, each TAM receptor was stably knocked down with shRNA in PC3, Du145, and





C4-2B prostate cancer cell lines. Protein expression of each receptor was decreased by at least 80% (Fig. S1). A decreased ratio of P-Erk 1/2 to P-p38 MAPK marks cellular dormancy in prostate and other cancers [Kobayashi et al., 2011; Bragado et al., 2013; Chery et al., 2014]. Therefore, we first examined P-Erk1/2 and P-p38 levels in PC3 cells with each of the TAM receptors knocked down. The ratio of P-Erk 1/2 to P-p38 was significantly decreased in serum starved MERTK knockdown cells, but not in TYRO3 or AXL knockdown cells (Fig. 1A and B). Similarly, the cell cycle inhibitor p27 was also previously found to be a dormancy marker [Kobayashi et al., 2011; Bragado et al., 2013]. In agreement with the Erk and p38 data, we found higher basal p27 protein expression in the MERTK knockdown cells (Fig. 1C and D). These data are consistent with a dormant phenotype in prostate cancer cells as a result of chronically reduced expression of MERTK.

MERTK KNOCKDOWN CAUSES EXPRESSION OF DORMANCY AND PLURIPOTENCY ASSOCIATED TRANSCRIPTION FACTORS

Transcription factors first studied in embryonic stem cells have been found to promote cancer dormancy [Sosa et al., 2015]. Therefore, we determined the basal expression level of three of these transcription factors in PC3 cells with each of the TAM receptors knocked down by shRNA. In parallel with the MAP kinase and p27 data, we saw marked upregulation of SOX2 message and protein in shMER but not shAXL or shTYRO3 cells (Fig. 2A and B). Similarly, we also observed increased expression of SOX2 in shMER C4-2B cells (Fig. 2A). We also observed increased NR2F1 and NANOG mRNA in shMER, but not shAXL or shTYRO3 cells (Fig. 2C and D). Because of the possibility of off target effects of shRNAs, we performed analogous studies with siRNA rather than shRNA and found that a siRNA targeting MERTK increased expression of SOX2 and NANOG in PC3 and C4-2B cells.







Fig. 3. Cell cycle analysis of TAM receptor knockdown PC3 cells by flow cytometry with BrdU and total DNA labeling. (A) Example flow plots for each cell type. Top: histograms of total DNA content labeled with7–AAD. Bottom: BrdU versus total DNA plots. The significance of each population is as follows: <2n DNA and BrdU negative; Sub–GO (apoptotic and necrotic), 2n DNA and BrdU negative; GO and G1 phases, 4n DNA and BrdU negative; G2 and M phases. BrdU positive; S–phase, >4n DNA; polyploid cells. (B) Quantification of the above cell cycle data. The table lists *P* values for each cell type compared to control with significant comparisons marked with an asterisk. Error bars are shown for the GOG1 and G2M populations and represent mean \pm SEM. *Represents *P* < 0.05 compared to shRNA control cells. (C) Western blots for total caspase–9 to verify the changes in Sub–G0 cells observed by flow cytometry.

MERTK KNOCKDOWN INDUCES CELL CYCLE CHANGES ASSOCIATED WITH CELLULAR DORMANCY

Cellular dormancy and decreased Erk 1/2 activity are also characterized by arrest in the G1 and G0 phases of the cell cycle [Aguirre-Ghiso et al., 2004]. Therefore, we compared cell cycle characteristics of TAM receptor knockdown PC3 cells cultured in 0.1% serum using flow cytometry to detect antibody labeled pulsed bromodeoxyuridine (BrdU) incorporated into DNA and 7-AAD to quantify total DNA content. This assay identifies BrdU positive cells as S-phase, less than 2n DNA as apoptotic or necrotic cells, 2n BrdU negative cells as G1 and G0 phases, 4n BrdU negative cells as G2 and M phases and cells with >4n DNA as polyploid (Fig. 3A). In agreement with our other data, MERTK knockdown cells showed a pattern consistent with dormancy, with a higher percentage of cells in G0/G1 and lower percentage in G2/M compared to control (Fig. 3B). Curiously, TYRO3 knockdown cells showed the opposite pattern with fewer cells in G0/G1 and more cells in the G2 and M phases. We did not convincingly see other analogous results for TYRO3 in our other experiments. We did not see differences between the different TAM knockdown cells in the percentage of cells in S-phase, but note that there are very few cells in S-phase in this study because of the low serum conditions. However, we did observe a reduction in the sub-G0 (apoptotic and necrotic) population in the shMER cells. Further, Western blots showed decreased levels of cleaved Caspase-9 in the shMER cells, thus suggesting that this reduced sub-G0 population represented reduced apoptosis. There is precedence in the literature for correlation of reduced apoptosis with a dormant phenotype [Aguirre-Ghiso et al., 2004]. Further, p38 stimulated cellular dormancy and reduced apoptosis have been proposed to be adaptive responses to allow DTCs to survive when conditions are not conducive to growth [Ranganathan et al., 2006].



Fig. 4. MERTK knockdown and metastasis free survival in a prostate cancer left ventricle injection xenograft model. (A) Experimental design. (B) Kaplan–Meier analysis of time to formation of metastases visible by bioluminescence imaging or death in mice injected with luciferase labeled control or shMER PC3 or Du145 cells. Left panels: metastases to any site. Right panels: cranial metastases only. *Indicates P < 0.05 versus control cells. (C) Strategy for quantification of the percentage of DTCs in mouse bone marrow by flow cytometry after first depleting the number of mouse cells with immunomagnetic beads. (D) Comparison of the percentage of DTCs in mouse bone marrow in control versus shMER PC3 cells 1 day after intracardiac injection. Error bars represent mean \pm SEM.

DEPLETION OF MERTK INCREASES METASTASIS FREE SURVIVAL IN VIVO

We next tested the importance for MERTK for dormancy escape in vivo. GFP and luciferase labeled control shRNA or MERTK shRNA PC3 or Du145 PCa cells were injected in the left ventricle of SCID mice and time to metastases visible by bioluminescence imaging, or death in rare cases, was evaluated with Kaplan–Meier analysis (Fig. 4A). Metastases are primarily to bone with both cell lines. Metastasis free survival in this model is established in the literature as a measure of dormancy [Kobayashi et al., 2011]. In agreement



Fig. 5. Effect of MERTK knockdown on growth of prostate cancer cells in culture and in subcutaneous tumors. (A) Subcutaneous tumor growth of control or shMER PC3 or Du145 cells as measured by bioluminescence imaging. *Indicates P < 0.05 versus shcontrol group. (B) Relative cell number quantified by XTT assay after 3 days of culture in 10%, 1%, or 0.1% serum for PC3, Du145, and C4–2B cells with each of the TAM receptors knocked down. Data are presented as mean \pm SEM. No pairwise comparisons were statistically significant.

with the in vitro data, metastasis free survival was prolonged in mice injected with shMER PC3 cells relative to control cells (Fig. 4B, top left). With Du145 cells, the increase in metastasis free survival with MERTK knockdown was small and not statistically significant (Fig. 4B, lower left). However, we noted that shMER Du145 cells appeared to develop metastases to the head more slowly, which approached statistical significance (Fig. 4B, lower right). Time to cranial metastases was again significantly different with PC3 cells. A defect in transit to the bone marrow could also explain delayed metastasis formation but would not involve dormancy escape. Therefore, in separate experiments, we used flow cytometry to quantify the percentage of control or shMER PC3 cells in marrow 1 day after injection and found no difference (Fig. 4C and D). This further supports the conclusion that MERTK is selectively important for dormancy escape rather than transit to the bone marrow in this model.

MERTK KNOCKDOWN DOES NOT DECREASE PROSTATE CANCER GROWTH IN VITRO OR AT A SUBCUTANEOUS SITE IN VIVO

We hypothesized that the role of MERTK was somewhat specific to cellular dormancy and therefore, it should not greatly affect overall cellular growth in vitro or in vivo at a site not dependent on the usual microenvironment, such as a subcutaneous site with an artificial extracellular matrix. Indeed, other groups who have derived dormant and tumorigenic cancer cell lines have observed similar growth in culture [Bragado et al., 2013; Sharma et al., 2016]. In agreement with these expectations, we saw no difference in subcutaneous growth in vivo between control and shMER PC3 cells and saw slightly higher growth of Du145 shMER cells compared to control (Fig. 5A). Similarly, we saw no significant differences in relative cell number, as measured by MTS assay, between any of the TAM shRNA knockdowns in PC3, Du145, and C4-2B cells cultured for 3 days in 0.1%, 1%, or 10% serum (Fig. 5B). The Du145 shMER



Fig. 6. Reversal of the shMER cell cycle phenotype with long term p38 inhibition. (A) Control or shMER PC3 cells cultured with 5 μ M SB203580 or 0.05% DMSO solvent control for 14 days in 10% serum and 3 days in 0.1% serum followed by cell cycle analysis with total DNA and BrdU labeling as described for Figure 3. Top: Total DNA flow cytometry histograms. Bottom: Quantified data. Error bars are shown for the GOG1 and G2M populations and represent mean \pm SEM. *Represents *P* < 0.05 for comparisons of interest. (B) Cell cycle analysis as above of Du145 cells grown for 17 days in 10% serum with or without SB203580.

cells trended toward higher relative cell number but did not reach statistical significance after multiple comparison testing.

CELL CYCLE CHANGES OF MERTK KNOCKDOWN ARE MAP KINASE DEPENDENT

Because of the well-established role of MAP kinases in regulation of cancer cellular dormancy, we investigated if p38 was required for the dormancy associated cell cycle changes induced by knockdown of MERTK. In appreciation of the known role of epigenetic changes in dormancy regulation and the time period required for these changes to occur, we cultured control and MERTK shRNA PC3 and Du145 cells with p38 inhibitor SB203580 or 0.05% DMSO solvent control for 2 weeks before performing experiments [Sosa et al., 2015]. The expected compensatory increase in P-p38 in response to p38 active site inhibition was observed by Western blot (Fig. S2A). We again observed an increased percentage of G0/G1 and decreased percentage

of G2/M cells with MERTK knockdown both in PC3 and Du145 cells (Fig. 6). This change induced by MERTK knockdown was completely reversed by p38 inhibition, thus showing involvement of MAP kinases. Similarly, we also observed a increased percentage of Ki67 negative cells (non-cycling) with MERTK knockdown in PC3 cells, which was also reversed by p38 inhibition (Fig. S2C). Curiously, we did not observe reversal of shMER induced p27 upregulation with p38 inhibition (Fig. S2B). This suggests that not all of the effects of MERTK knockdown are MAP kinase dependent.

MERTK KNOCKDOWN INDUCES DORMANCY ASSOCIATED CHANGES IN HISTONE H3 METHYLATION BY A MAP KINASE DEPENDENT MECHANISM

Lastly, we used flow cytometry with specific antibodies to determine the effect of shRNA knockdown in PC3 cells of each of the TAM receptors on histone H3 tri-methylated lysine 9 (H3 K9 me3) and





histone H3 tri-methylated lysine 27 (H3 K27 me3). Both of these histone marks are increased in dormant cells [Sosa et al., 2015]. Because the majority of cells were positive, we gated on the negative and dim populations rather than the positive population (Fig. 7A). We observed the expected dormancy associated change in H3 K9me3, and a trend toward significance for H3 K27me3 with MERTK knockdown but no significant differences for AXL or TYR03 knockdown (Fig. 7B). In cells treated with the p38 inhibitor SB203580 or solvent control, we saw significant dormancy associated changes with MERTK knockdown, which were partially reversed by p38 inhibition (Fig. 7C).

DISCUSSION

Overall, these studies implicate MERTK in prostate cancer dormancy escape through a MAP kinase dependent mechanism linked to transcriptional and epigenetic regulation. Knockdown of MERTK consistently induced the changes expected for dormant cells; a decreased ratio of P-Erk to P-p38, increased p27 expression, expression of dormancy, and pluripotency associated transcription factors, and G0/G1 arrest. Further, these findings translated to an increased metastasis free survival in vivo. This identifies MERTK as being important for the process whereby one or a few cancer cells progress to a small tumor (i.e., escape from cellular dormancy). As expected, MERTK knockdown did not inhibit growth in culture or growth of subcutaneous tumors implanted in an artificial matrix. This lack of a general growth inhibitory effect of MERTK knockdown suggests specificity for dormancy regulation in bone and a requirement for a signal from the microenvironment rather than dysregulation of multiple cellular processes.

Our data do not identify which signal(s) from the microenvironment interact with MERTK to regulate dormancy. The four other TAM receptor ligands, other than GAS6 (Tubby, Tubby like protein 1, Galectin 2, and Protein S), may also play a role. Although not a MERTK ligand, retinoic acid may be indirectly involved as well because it has been shown to stimulate cancer dormancy through NR2F1 and also interacts indirectly with MERTK in immune cells [Garabuczi, 2015; Sosa et al., 2015]. Because no recombinant protein or other treatment other than low serum conditions was required for



Fig. 8. Results summary for MERTK stimulates PCa dormancy escape through a MAP kinase dependent mechanism, also involving p27, pluripotency transcription factors, and histone methylation.

our observed in vitro effects, knockdown of MERTK may have caused epigenetic reprogramming of the cells as a result of chronic alterations in signaling from autocrine TAM receptor ligands. Prostate cancer cells are known to express MERTK ligands GAS6 and PROS1 [Jung et al., 2016; Ning et al., 2016]. The observed changes in histone H3 methylation also suggests that part of the effects of MERTK may be due to epigenetic reprogramming rather than through a more immediate response to ligands.

Previously, we examined the expression of TYRO3 and AXL, but not MERTK in disseminated tumor cells (DTCs) versus prostate cancer primary tumors and gross metastases [Taichman et al., 2013]. We reported higher expression of AXL in DTCs and higher expression of TYRO3 in primary tumors and gross metastases. Thus, we hypothesized that AXL might be important for maintenance of dormancy and TYRO3 might be important for escape from dormancy or growth of a gross tumor. In retrospect, it would have been useful to include MERTK in these studies. Our most recent work, currently in press, confirms the earlier hypothesis regarding AXL and shows that AXL is required for TGF- β 2 induced prostate cancer dormancy [Yumoto et al., 2016]. However, prior to the current work, the contribution of MERTK to prostate cancer dormancy remained completely unclear. Here, we took an unbiased approach and were consistently guided by the data to focus on MERTK. However, our data does not rule out contributions from AXL or TYRO3 in stimulation of dormancy escape. For example, the percent knockdown might have been insufficient to observe effects with shAXL or shTYRO3.

We recently reported a role for MERTK in formation of prostate cancer stem-like cells (CSCs) [Jung et al., 2016; Shiozawa et al., 2016]. However, the proliferative rate of CSCs was not examined in these studies and other literature reports that CSCs can be either slowly or rapidly cycling [Sharma et al., 2016; Takeishi and

Nakayama, 2016]. Additionally, others identified MERTK in an unbiased screen of over 100 kinases as a stimulator of prostate cancer metastasis. Therefore, our data showing a new role for MERTK in dormancy escape combined with the other studies implicating MERTK in CSC formation and metastasis, suggests that MERTK plays a role in prostate cancer progression by regulating several processes.

In summary, we are beginning to understand the roles of the TAM kinases in prostate cancer dormancy regulation. AXL plays a role in dormancy maintenance through TGF- β 2 and microenvironment dependent effects. The role of TYRO3 remains less clear. Conversely, this study provides the first evidence that MERTK stimulates prostate cancer dormancy escape. Our studies with a p38 inhibitor show MAP kinases are required for MERTK to stimulate dormancy escape. Furthermore, MERTK causes changes observed in histone H3 methylation, pluripotency associated transcription factors and cell cycle regulatory proteins observed in dormancy regulation of other cancers (Fig. 8). Thus, MERTK appears to reprogram DTCs to grow rather than remain dormant. This understanding of the role of MERTK should inform future studies of prostate cancer dormancy and may provide a therapeutic target for prevention of recurrence.

ACKNOWLEDGMENTS

The authors wish to thank Taocong Jin with assistance with primer design and qRT-PCR. Portions of Figure 8 are derived from the Servier powerpoint image bank.

REFERENCES

Aguirre-Ghiso JA, Ossowski L, Rosenbaum SK. 2004. Green fluorescent protein tagging of extracellular signal-regulated kinase and p38 pathways reveals novel dynamics of pathway activation during primary and metastatic growth. Cancer Res 64(20):7336–7345.

Amling CL, Blute ML, Bergstralh EJ, Seay TM, Slezak J, Zincke H. 2000. Longterm hazard of progression after radical prostatectomy for clinically localized prostate cancer: Continued risk of biochemical failure after 5 years. J Urol 164(1):101–105.

Bragado P, Estrada Y, Parikh F, Krause S, Capobianco C, Farina HG, Schewe DM, Aguirre-Ghiso JA. 2013. TGF-beta2 dictates disseminated tumour cell fate in target organs through TGF-beta-RIII and p38alpha/beta signalling. Nat Cell Biol 15(11):1351–1361.

Caberoy NB, Zhou Y, Li W. 2010. Tubby and tubby-like protein 1 are new MerTK ligands for phagocytosis. EMBO J 29(23):3898–3910.

Chery L, Lam HM, Coleman I, Lakely B, Coleman R, Larson S, Aguirre-Ghiso JA, Xia J, Gulati R, Nelson PS, Montgomery B, Lange P, Snyder LA, Vessella RL, Morrissey C. 2014. Characterization of single disseminated prostate cancer cells reveals tumor cell heterogeneity and identifies dormancy associated pathways. Oncotarget 5(20):9939–9951.

Faltermeier CM, Drake JM, Clark PM, Smith BA, Zong Y, Volpe C, Mathis C, Morrissey C, Castor B, Huang J, Witte ON. 2015. Functional screen identifies kinases driving prostate cancer visceral and bone metastasis. Proc Natl Acad Sci USA 113(2):E172–E181.

Garabuczi E, Sarang Z, Szondy Z. 2015. Glucocorticoids enhance prolonged clearance of apoptotic cells by upregulating liver X receptor, peroxisome proliferator-activated receptor-delta and UCP2. Biochim Biophys Acta 1853(3):573–582.

Graham DK, DeRyckere D, Davies KD, Earp HS. 2014. The TAM family: Phosphatidylserine sensing receptor tyrosine kinases gone awry in cancer. Nat Rev Cancer 14(12):769–785.

Jung Y, Decker AM, Wang J, Lee E, Kana LA, Yumoto K, Cackowski FC, Rhee J, Carmeliet P, Buttitta L, Morgan TM, Taichman RS. 2016. Endogenous GAS6 and Mer receptor signaling regulate prostate cancer stem cells in bone marrow. Oncotarget 7(18):25698–25711.

Jung Y, Shiozawa Y, Wang J, McGregor N, Dai J, Park SI, Berry JE, Havens AM, Joseph J, Kim JK, Patel L, Carmeliet P, Daignault S, Keller ET, McCauley LK, Pienta KJ, Taichman RS. 2012. Prevalence of prostate cancer metastases after intravenous inoculation provides clues into the molecular basis of dormancy in the bone marrow microenvironment. Neoplasia 14(5):429–439.

Kobayashi A, Okuda H, Xing F, Pandey PR, Watabe M, Hirota S, Pai SK, Liu W, Fukuda K, Chambers C, Wilber A, Watabe K. 2011. Bone morphogenetic protein 7 in dormancy and metastasis of prostate cancer stem-like cells in bone. J Exp Med 208(13):2641–2655.

Lee E, Decker AM, Cackowski FC, Kana LA, Yumoto K, Jung Y, Wang J, Buttitta L, Morgan TM, Taichman RS. 2016. Growth arrest-specific 6 (GAS6) promotes prostate cancer survival by G1 Arrest/S phase delay and inhibition of apoptotic pathway during chemotherapy in bone marrow. J Cell Biochem 117(12):2815–2824.

Morgan TM, Lange PH, Porter MP, Lin DW, Ellis WJ, Gallaher IS, Vessella RL. 2009. Disseminated tumor cells in prostate cancer patients after radical prostatectomy and without evidence of disease predicts biochemical recurrence. Clin Cancer Res 15(2):677–683.

Ning P, Zhong JG, Jiang F, Zhang Y, Zhao J, Tian F, Li W. 2016. Role of protein S in castration-resistant prostate cancer-like cells. Endocr Relat Cancer 23(8):595–607.

Ranganathan AC, Adam AP, Zhang L, Aguirre-Ghiso JA. 2006. Tumor cell dormancy induced by p38SAPK and ER-stress signaling: An adaptive advantage for metastatic cells? Cancer Biol Ther 5(7):729–735.

Sharma S, Xing F, Liu Y, Wu K, Said N, Pochampally R, Shiozawa Y, Lin HK, Balaji KC, Watabe K. 2016. Secreted protein acidic and rich in cysteine

(SPARC) mediates metastatic dormancy of prostate cancer in the bone. J Biol Chem 291(37):19351–19363.

Shiozawa Y, Berry JE, Eber MR, Jung Y, Yumoto K, Cackowski FC, Yoon HJ, Parsana P, Mehra R, Wang J, McGee S, Lee E, Nagrath S, Pienta KJ, Taichman RS. 2016. The marrow niche controls the cancer stem cell phenotype of disseminated prostate cancer. Oncotarget. doi: 10.18632/oncotarget.9251 [Epub ahead of print].

Shiozawa Y, Pedersen EA, Patel LR, Ziegler AM, Havens AM, Jung Y, Wang J, Zalucha S, Loberg RD, Pienta KJ, Taichman RS. 2010. GAS6/AXL axis regulates prostate cancer invasion, proliferation, and survival in the bone marrow niche. Neoplasia 12(2):116–127.

Sosa MS, Parikh F, Maia AG, Estrada Y, Bosch A, Bragado P, Ekpin E, George A, Zheng Y, Lam HM, Morrissey C, Chung CY, Farias EF, Bernstein E, Aguirre-Ghiso JA. 2015. NR2F1 controls tumour cell dormancy via SOX9- and RARbeta-driven quiescence programmes. Nat Commun 6:6170.

Taichman RS, Patel LR, Bedenis R, Wang J, Weidner S, Schumann T, Yumoto K, Berry JE, Shiozawa Y, Pienta KJ. 2013. GAS6 receptor status is associated with dormancy and bone metastatic tumor formation. PLoS ONE 8(4):e61873.

Takeishi S, Nakayama KI. 2016. To wake up cancer stem cells, or to let them sleep, that is the question. Cancer Sci 107(7):875–881.

Yumoto K, Eber M, Wang J, Cackowski F, Lee E, Nobre AR, Aguirre-Ghiso J, Jung Y, Taichman R. 2016. Axl is required for TGF- β 2-induced dormancy of prostate cancer cells in the bone marrow. Sci Rep 6:36520. doi: 10.1038./ srep36520

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher's web-site.

∂ Open Access Full Text Article

PERSPECTIVES

495

The ABC7 regimen: a new approach to metastatic breast cancer using seven common drugs to inhibit epithelial-to-mesenchymal transition and augment capecitabine efficacy

Richard E Kast¹ Nicolas Skuli² Samuel Cos³ Georg Karpel-Massler⁴ Yusuke Shiozawa⁵ Ran Goshen⁶ Marc-Eric Halatsch⁴

'IIAIGC Study Center, Burlington, VT, USA; ²INSERM, Centre de Recherches en Cancérologie de Toulouse - CRCT, UMR1037 Inserm/Université Toulouse III – Paul Sabatier, Toulouse, France; ³Department of Physiology and Pharmacology, School of Medicine, University of Cantabria and Valdecilla Research Institute (IDIVAL), Santander, Spain; ⁴Department of Neurosurgery, Ulm University Hospital, Ulm, Germany; ⁵Department of Cancer Biology, Comprehensive Cancer Center, Wake Forest School of Medicine, Winston-Salem, NC, USA; ⁶Eliaso Consulting Ltd., Tel Aviv-Yafo, Israel

Correspondence: Richard E Kast IIAIGC Study Center, 22 Church Street, Burlington, VT 05401, USA Tel +1 802 557 7278 Email richarderickast@gmail.com



Abstract: Breast cancer metastatic to bone has a poor prognosis despite recent advances in our understanding of the biology of both bone and breast cancer. This article presents a new approach, the ABC7 regimen (Adjuvant for Breast Cancer treatment using seven repurposed drugs), to metastatic breast cancer. ABC7 aims to defeat aspects of epithelial-to-mesenchymal transition (EMT) that lead to dissemination of breast cancer to bone. As add-on to current standard treatment with capecitabine, ABC7 uses ancillary attributes of seven already-marketed noncancer treatment drugs to stop both the natural EMT process inherent to breast cancer and the added EMT occurring as a response to current treatment modalities. Chemotherapy, radiation, and surgery provoke EMT in cancer generally and in breast cancer specifically. ABC7 uses standard doses of capecitabine as used in treating breast cancer today. In addition, ABC7 uses 1) an older psychiatric drug, quetiapine, to block RANK signaling; 2) pirfenidone, an anti-fibrosis drug to block TGF-beta signaling; 3) rifabutin, an antibiotic to block beta-catenin signaling; 4) metformin, a first-line antidiabetic drug to stimulate AMPK and inhibit mammalian target of rapamycin, (mTOR); 5) propranolol, a beta-blocker to block beta-adrenergic signaling; 6) agomelatine, a melatonergic antidepressant to stimulate M1 and M2 melatonergic receptors; and 7) ribavirin, an antiviral drug to prevent eIF4E phosphorylation. All these block the signaling pathways – RANK, TGF-beta, mTOR, beta-adrenergic receptors, and phosphorylated eIF4E – that have been shown to trigger EMT and enhance breast cancer growth and so are worthwhile targets to inhibit. Agonism at MT1 and MT2 melatonergic receptors has been shown to inhibit both breast cancer EMT and growth. This ensemble was designed to be safe and augment capecitabine efficacy. Given the expected outcome of metastatic breast cancer as it stands today, ABC7 warrants a cautious trial.

Keywords: ABC7, breast cancer, agomelatine, capecitabine, metformin, pirfenidone, propranolol, quetiapine, repurposing, ribavirin, rifabutin, TGF-beta

Plain language summary

This article presents the rationale and thinking behind the ABC7 regimen for metastatic breast cancer. Since there is currently no cure for breast cancer once it has spread to bone and other organs beyond the breast itself, the ABC7 regimen was designed to take advantage of ancillary attributes of seven common and readily available noncancer treatment drugs that, in theory, should make current traditional cytotoxic chemotherapy with capecitabine more effective. The ABC7 regimen has not been shown to be safe or effective yet. In the current article, we discuss an untested proposal for a new treatment approach to metastatic breast cancer.

with the set of the se

Introduction

Estrogen-positive metastatic breast cancer cannot be cured currently.¹ One major metastatic site of breast cancer is bone. Once breast cancer metastasizes to bone, the survival rate declines despite recent advances in local treatments of breast cancer. Current treatment strategies for bone metastasis, including bone-targeted agents (bisphosphonate and denosumab), provide only palliation. New and effective therapeutic strategies for this still incurable disease are therefore urgently needed.

This article reviews the attributes of seven currently marketed drugs that, as indicated by prior research data, will block or partially block the escape pathways from current traditional treatments. The seven drugs of ABC7 were chosen by first identifying the basic pathways by which EMT is initiated and maintained. We then reviewed 1000 of the most commonly used drugs² for which we have both usual plasma levels and published data showing potential inhibitory interaction with these pathways. The resulting list was reduced by semi-subjective evaluation of the strength of data on their EMT inhibition benefit versus the drugs' expected tolerability. The better the tolerability, the weaker the data had to be for inclusion.

This ABC7 regimen is designed to block several core breast cancer growth signals in a coordinated manner, thereby augmenting the cytotoxicity of a currently used cytotoxic chemotherapy drug, capecitabine. Figure 1 shows an overview schematic of the biochemistry that ABC7 is designed to influence. This is explained in detail in the respective drug discussions in the "Drugs to inhibit EMT" section. Table 1 gives an overview of the ABC7 drugs and their intended targets in treating breast cancer.

The ABC7 regimen follows the approach of previous cancer treatment regimens, for example, MTZ regimen,³ COMBAT regimen,⁴ MEMMAT regimen, a current trial of Peyrl et al's seven-drug cocktail (ClinicalTrials.gov Identifier: NCT01356290), and CUSP9 regimen.^{5,6} In all of the studies, extensive use is made of drugs not primarily marketed to treat cancer but that have ancillary attributes that research data indicate would enhance the anticancer effect of a cytotoxic, traditional cancer treatment drug. The ancillary drugs exert anticancer effects by blocking various growth-enhancing survival pathways used by the target cancer or as for agomelatine are agonists at growth-retarding receptors.

Similar to other cancers, breast cancer has heterogeneous regions within the same tumor – different areas that depend on or use different growth-signaling pathways. This is related to but distinct from the idea of clonal evolution driven by cytotoxic chemotherapy selection. Both forms of heterogeneity exist in a typical breast cancer, proteomic and genomic. ABC7 aims to inhibit breast cancer by pharmacological manipulation of what genes are expressed and what genes are not, as well as by targeting different clonal variants of the original breast cancer clone.

Because of these limitations, we do not expect testing for molecular markers to be predictive or useful. In addition, further intensifying the spatial and temporal diversity of the molecular status, particularly for EMT markers, are the diversity-driving effects of chemotherapy⁷ and discussed in greater detail in the following sections.

Cytotoxic chemotherapy also induces important receptor status changes in a large minority of breast cancer cases.^{8,9} Typical findings are as follows: 13% changed from HR+ to HR-, 5% changed from HR- to HR+, 6% changed from HER2+ to HER2, 3% changed from HER2 to HER2+, and 13% changed to triple negative.⁸

Multiple signaling systems have been identified that drive metastatic breast cancer.^{1,9–11} These growth-driving receptors can cross cover for each other.^{1,9–11} When one is pharmacologically blocked, several parallel growth-driving pathways can become active, taking the place of the blocked pathways. Growth factor signaling converges from a wide variety of outer membrane receptors to more restricted, fewer, intracellular pathways. This is, more elegantly stated, the spatial–temporal genomic and proteomic range, the "genetic collectives [that] dominate the landscape of advanced-stage (malignant) disease."^{11,12} We see this as mandating an integrated, coordinated polypharmacy to successfully address these malignancy attributes.

Capecitabine is intracellularly metabolized to 5-FU; the details are given in the "Capecitabine: 359 Da, half-life <1 hour" section. ABC7 drugs are designed to make 5-FU more effective.

The results of several recent ER+ metastatic breast cancer studies are listed in Table 2. These studies cannot be judged simply by overall survival in that entry requirements were different, with different kinds and number of prior treatments. These numbers in Table 2 are for general idea only.¹ One cannot conclude that one of these is better than another.

In general, post-progression survival durations in recent Phase III studies of combination therapy ranged from approximately 16 to 33 months.¹

EMT is a feature of cancers generally 13 and breast cancer specifically. 14,15

Table 3 lists several features and behaviors associated with the two (epithelial and mesenchymal) states. Interestingly, a transcription factor ZEB1, known to control EMT,



Figure I The basic biochemical, intracellular, and receptor pathways relating to interventions of the ABC7 regimen for advanced breast cancer. Notes: This is the basic breast cancer intracellular circuitry that the ABC7 regimen attempts to address. The two major controllers or stimuli for eIF4E activation are indicated by yellow arrows, the p-mTOR and the MNK paths.

Table I	The drugs	of ABC7, t	their targets	during treat	ment of breast	cancer, and	suggested	doses
		, -	· · · · · · · · · · · · · · · · · · ·			,		

Drug	Target in breast cancer treatment	Starting dose	Target dose
Capecitabine	DNA synthesis	600 mg/m ² twice daily.	1250 mg/m² twice daily. 7 days on, 7 days off or 14 days on,
		7 days on, 7 days off	7 off
Quetiapine	RANK/RANKL	50 mg once at bedtime	300–600 mg once at bedtime
Pirfenidone	TGF-beta	200 mg three times daily	600 mg three times daily
Rifabutin	BCL-6; beta-catenin	150 mg/day	300 mg/day
Metformin	AMPK*; mTOR; mitochondria oxphos	500 mg once daily	1000 mg twice daily
Propranolol	Beta-adrenergic receptors	10 mg twice daily	Uptitrate as tolerated
Agomelatine	Melatonergic receptors*	25 mg once at bedtime	50 mg once at bedtime
Ribavirin**	elF4E; MNK; IMPDH	600 mg/day	1200 mg/day

Notes: The drugs are listed in a suggested order of addition. Pace of drug addition is individualized per patient and physician estimations of risk/benefit. *Note that all entries denote inhibition of named target except for metformin that activates AMPK and agomelatine that stimulates melatonin receptors. All drugs, except capecitabine, are given continuously without interruption. Capecitabine is given on 7 days off, or 14 days on, 7 days off cycles. **Ribavirin is likely to give unpleasant side effects and depressed mood-but is potentially a beneficial enough drug to try.

Table	2 Representative	recent	trials	in	metastatic	HR+	breast
cancer							

Intervention	Months survival	Trial
Anastrozole + fulvestrant	48	SWOG
Anastrozole + fulvestrant	38	FACT
Anastrozole + fulvestrant	21	SoFEA
Letrozole + fulvestrant	52	LEA
Everolimus + exemestane	31	BOLERO 2

contributes to breast cancer osteolytic bone metastasis, but not brain or lung metastasis.^{16,17}

EMT is a phase transition, where flat, sessile, mutually adherent epithelioid cells take on a rounded, non-adherent, motile mesenchymal shape and behavior.^{18–20} The reverseless transient state and process, MET, also occurs and is also a feature of robust or aggressive cancer growth.²¹ Post-EMT cells tend to be invasive but proliferation restricted. Post-MET cells tend to be proliferative but have limited invasiveness.^{18–20} Breast cancers develop in proximity to adipose tissue. Adipocytes are a further trigger to EMT.²² The relationship between stem cell subpopulations within a cancer and EMT process is also unclear,²³ and the two populations probably largely overlap.

Perhaps our deepest insight into the EMT process in breast cancer came from a study by Bulfoni et al.²⁴ They showed that all patients with metastatic breast cancer had CTCs. These circulating cancer cells split into four groups: those with epithelial, those with mesenchymal, those with both, and those with neither marker. Patients with higher numbers of circulating cancer cells that expressed both markers had shorter overall survival.

Survival as a function of E-cadherin expression, representative of epithelial state, and fibronectin, representative of mesenchymal state, was examined by immunohistochemistry on 1495 breast cancer biopsies.²⁵ More E-cadherin and less fibronectin are associated with longer survival.²⁵ Breast cancer patients whose tissue expresses greater EMT-related protein have shorter survival.²⁶ EMT drives chemotherapy resistance and other poor prognosis features in breast cancer.^{27–29} In addition, a greater degree of metabolic changes characteristic of EMT in breast cancer predicts shorter overall survival.³⁰

Medical research discusses vimentin, fibronectin, and N-cadherin as markers of EMT process, but these proteins would be more accurately viewed as mediators of the attributes we designate EMT.

As indicated in Table 3, fibronectin is a characteristic marker of EMT. Higher breast cancer tissue expression of fibronectin correlates with shorter survival.^{31,32} A range of other characteristic behavioral and morphological attributes of EMT and MET states is also outlined in Table 3.

 Table 3 Characteristic protein markers and mediators of EMT

 in breast cancer

Marker	Epithelial state	Mesenchymal state
E-cadherin	Increased	Decreased
N-cadherin	Decreased	Increased
ZO-I	Increased	Decreased
Occludin	Increased	Decreased
Vimentin	Decreased	Increased
Fibronectin	Decreased	Increased
MMP-2	Decreased	Increased
MMP-9	Decreased	Increased
Phenotype	Epithelial state	Mesenchymal state
Motility	Sessile	Motile
Shape	Elongated	Rounded
Adherence	Adherent to neighbors	Non-adherent to neighbors
Invasion	Noninvasive	Invasive
Proliferation	Higher proliferation	Lower proliferation
Microtentacles	Absent	Present

Abbreviation: EMT, epithelial-to-mesenchymal transition.

Chemotherapy triggers EMT

Paclitaxel triggers EMT in breast cancer, increasing mesenchymal markers, vimentin and fibronectin, and decreasing epithelial marker, ZO-1.³³ Experimental inhibitors of TGFbeta signaling block paclitaxel-induced EMT and suppress paclitaxel-induced CSC properties.³³ Paclitaxel also increases EMT markers in mouse breast cancer cell line, MCF-7/PAX.³⁴ Doxorubicin exposure enhances gastric cancer's EMT marker expression.³⁵

These reports, combined with similar findings in other cancers, allow a general statement of a core principle of oncology: cytotoxic chemotherapies tend to provoke EMT. Such a conjecture is amply supported by the recent work of Yoshimasu et al³⁶ who reported that cisplatin, 5-FU, gemcitabine, paclitaxel, and vinorelbine show hormesis when tested individually.

Surgical trauma or fine-needle biopsy triggers EMT

Of concern in current medical practice, there is a tendency for cancers generally, and breast cancer specifically, to be triggered by any kind of tissue disruption – including fineneedle biopsy – to undergo EMT with consequent cancer cell shedding to circulation. Such hematogenous tumor cell dissemination could be the origin of later overt metastases.

Breast cancers in mice release a flood of CTCs after simple fine-needle biopsy.³⁷ Clinical needle biopsy of breast cancer triggers recruitment of inflammatory cells to the biopsy site and causes increased tumor cell mitoses in the biopsied area.³⁸ In a second murine breast cancer study, both fineneedle biopsy and surgical resection resulted in the release of a flood of CTCs, but noteworthy in this work is that biopsy resulted in greater and longer lasting appearance of circulating cancer cells than did surgical resection.³⁹ These murine data were replicated by Kaigorodova et al⁴⁰ who showed that simple fine-needle biopsy of human breast cancers releases a flood of breast cancer cells into the general circulation. These authors found that although some released CTCs had CSC markers and attributes and some did not, none of them had the particular EMT markers for which they were tested.

This worrisome situation in breast cancer is similar to data collected in other cancers. For example, treatment with radioactive needle insertion into prostate cancer results in significant hematological shedding of tumor cells post procedure.⁴¹ Standard transrectal ultrasound-guided prostate needle biopsy results in detectable prostate cancer cells in the circulation in half of patients.⁴² Oral squamous cell carcinoma biopsies result in 16% of patients having post-biopsy

CTCs.⁴³ Simple wide excision that does not disrupt the tumor tissue integrity did not result in postoperative CTCs, whereas incisional biopsy did.⁴⁴

A study is therefore required comparing the long-term outcome potential difference between those having fineneedle biopsy versus those having initial wide lesion excision. If initial excision that leaves the suspicious mass intact does result in fewer later disseminated metastases, it might be worth the iatrogenic morbidity incurred by the consequent excision of some benign masses.

Radiation triggers EMT

Above we reviewed some evidence that chemotherapy and mechanical tissue disruption give rise to CTC and EMT in surviving cells. Below we review data showing radiation causes CTC and EMT as well.^{45,46} For a few specific examples, as in other cancers,⁴⁷ breast cancers synthesize GM-CSF that then functions as a growth factor for them.⁴⁸ Clinically used, radiation treatment not only kills breast cancer cells and prolongs survival in breast cancer but also triggers exposed residual cells that are not killed to undergo EMT, to start migrating, and to synthesize increased amounts of autocrine growth factor, GM-CSF.⁴⁹ Radiation also increases IL-6, migration, and EMT markers in murine and human breast cancer cell lines.⁵⁰ The subject of radiation-induced EMT and radiation-induced increase in CTCs was recently reviewed by Lee et al.⁵¹

Clinically, finding greater post-EMT CTCs confers a worse prognosis with more aggressive disease course and greater metastatic proclivity in colon cancer⁵² and finding circulating clusters of vimentin-positive gastric cancer cells confers a worse prognosis⁵³ as did finding circulating cancer cell clusters and vimentin-positive CTC in colon cancer.⁵⁴ Surgery for epithelial ovarian cancer causes an increase in both EMT-positive and EMT-negative CTCs, but there is a disproportionate increase in EMT positive. The increase in EMT-positive CTCs was even stronger after platinum-based chemotherapy.⁷ As Kolbl et al⁵⁵ point out, EMT precedes the release of CTCs but after entering circulation CTC can revert to epithelial or partial epithelial phenotype.

Based on all these evidences, it seems that inhibiting EMT is a worthwhile goal during breast cancer treatment and that current common cancer treatments have elements of cancer growth stimulation inherent to them, or as Niccolo Machiavelli (born 1469–died 1527) said in 1513:

People should either be caressed or crushed. If you do them minor damage they will get their revenge; but if you cripple them there is nothing they can do. If you need to injure someone, do it in such a way that you do not have to fear their vengeance.

ABC7 regimen was crafted with that in mind.

Drugs to inhibit EMT Quetiapine: 384 Da, cyp3A4 to norquetiapine, 6-hour half-life

The RANK, its ligand (RANKL), and the soluble decoy receptor OPG (or bone protector) are central elements in breast cancer's establishment of metastases to bone.⁵⁶ Early indications are that quetiapine inhibits the RANK/RANKL signaling system.⁵⁷

Several forms of pro-RANKL are expressed on osteoblasts. After proteolytic release, RANKL binds to RANK leading to osteoclast syncytium formation then osteoclasts' resorption of bone. Osteopetrosis results when RANK/ RANKL system is nonfunctional. RANK/RANKL also functions in guiding normal breast gland ontogeny. There occurs an ebb and flow of RANK expression in mammary duct epithelial cells during the menstrual cycle, the increase occurring in late luteal phase. RANK/RANKL function is essential to the luminal epithelial proliferation seen particularly where ducts branch.⁵⁶ Higher levels of RANK/RANKL in human breast cancer biopsy tissue correlate with higher metastasis likelihood and shorter survival.⁵⁸

PR-negative cells are affected through RANKL-induced paracrine actions leading to proliferation of mammary epithelial PR-negative cells.⁵⁹

RANK/RANKL is a core physiologic signaling system allowing circulating breast cancer cells to metastasize to bone.^{60,61} RANK/RANKL is a principal part of the complex signaling giving rise to breast cancer's propensity to metastasize to bone.

Since breast cancer commonly metastasizes to bone with consequent bone pain, pathological fractures, vertebrae compressions, and hypercalcemia, this process is important to block. Breast cancer cells are continuously shed into circulation from the primary and metastatic sites. Why then do these CTCs choose to establish growth preferentially in bone? This is because these CTCs can establish growth-enhancing communication with bone cells, specifically osteoblasts, and they do so primarily via RANK/RANKL.⁶² Muscle, skin, liver, spleen, fat, and other sites of less common breast cancer metastasis cannot so reciprocally communicate.

Osteoblasts receiving RANKL signaling transform to syncytial osteoclasts that resorb bone and increase TGF-beta. TGF-beta is also stored in bone, then released with any bone

499

dissolution. This creates room for the CTCs to grow and free TGF-beta signaling prompting them to do so,^{62,63} making the TGF-beta blocking drug pirfenidone, which is discussed in the following sections, an ideal partner drug for quetiapine during metastatic breast cancer treatment.

An initial dose can be 50 mg once at bedtime, uptitrating to a target dose of quetiapine 300 mg or more as tolerated, given once at bedtime. Tiredness for a few hours on awakening is common upon starting quetiapine. It then abates after a week or so but reappears after each dose increase. Some weight increase due to increased appetite can be expected. Otherwise, side effects are not common.

Pirfenidone: 185 Da, cyp1A2, 3-hour half-life

Pirfenidone is a 185 Da drug approved and marketed to treat idiopathic pulmonary fibrosis.⁶⁴ Mild-to-moderate, reversible, nausea, dyspepsia, and rash are side effects in about one-third of treated patients, but these often resolve with continued use. Approximately 2403 mg/day divided into three equal doses is a common pirfenidone dose in treating its marketed indication, idiopathic pulmonary fibrosis.^{64,65}

Pirfenidone blocks TGF-beta signaling.^{66–70} TGF-beta is a 25 kDa signaling protein proteolytically clipped from a larger precursor protein. Carboplatin induces elevation of TGF-beta and triggers EMT in NSCLC, as given in the "Chemotherapy triggers EMT" section,⁷¹ both effects blocked by coadministration with pirfenidone.⁷¹

TGF-beta signaling is a major driver of EMT in cancer generally⁷²⁻⁷⁴ and in breast cancer EMT specifically.⁷⁵⁻⁷⁸ TGFbeta is a facilitating element of many cancers by promoting angiogenesis and differentiation, by immune suppression, by promoting loss of cell-to-cell contact, and particularly by promoting EMT. Pirfenidone inhibits TGF-beta-induced phosphorylation of SMAD3, p38, and AKT. TGF-beta provides a "get up and go" signal for breast cancer.⁷⁹ In a murine breast cancer model, TGF-beta exposure also enhances normal lung's ability to better support establishment of breast cancer metastases.⁸⁰ TGF-beta drives breast cancer's EMT and various biochemical, morphological, and behavioral changes characteristic of EMT.^{10,81-83} The manifold paths by which TGF-beta signaling leads to or enhances EMT specifically in breast cancer were outlined by Tan et al,¹⁸ Chen et al,⁸¹ Nooshinfar et al,⁸⁴ and Felipe Lima et al.⁷⁵

TGF-beta dependency for taking on typical mesenchymal morphology, increased motility, and increased vimentin expression after radiation exposure was shown in breast, colon, and lung adenocarcinoma cell lines.⁸⁵ Preclinical studies have shown activity in pirfenidone's enhancing cisplatin cytotoxicity to NSCLCs.⁸⁶ In addition, pirfenidone enhances radiation and sunitinib cytotoxicity in Lewis lung cancer cells⁸⁷ and reduces desmoplasia in pancreatic cancer.⁸⁸ Growth of human TNBC tissue (ER negative, PR negative, HER2 negative) xenografted to nude mice was inhibited more by pirfenidone and doxorubicin than by doxorubicin alone.⁸⁹ In another murine breast cancer model, pirfenidone reduced intratumoral collagen and hyaluronan by TGF-beta inhibition with consequent improvement of doxorubicin efficacy.⁹⁰

Pirfenidone disrupts Hh signaling in parallel with TGFbeta inhibition, a worthwhile added benefit during breast cancer treatment.⁹¹

The starting dose of pirfenidone is 267 mg three times a day. This is gradually increased at 14-day intervals as tolerated to 801 mg three times daily. Pirfenidone at 400 mg three times daily (1200 mg/day) used to treat potential progression of hepatitis C-related fibrosis reduced circulating TGF-beta and IL-6.⁹² Abdominal pain, rash, and nausea were seen in a half of treated patients, but these side effects tended to subside within a month or two and no patient dropped out due to them.

Ribavirin: 244 Da, 6-day half-life for a single oral dose, up to 12 days after continuous use

Since its introduction to clinical practice in the late 1970s, ribavirin had been used to treat various viral infections, later becoming central to a now-outmoded hepatitis C treatment. Ribavirin remains useful in treating human respiratory syncytial virus infections and selected other rarer virus infections such as those of the hantavirus group.⁹³ Ribavirin is currently being investigated in numerous clinical trials for its therapeutic activity in various cancers, particularly acute myeloid leukemia (NCT02109744, NCT02073838), head and neck cancer (NCT01268579), and notably for ABC7, metastatic breast cancer (NCT01056757).

Although ribavirin's mechanisms of antiviral and anticancer action are uncertain and probably will vary between viruses, several potential mechanisms of action have been identified. One proposes that ribavirin enters the cell via a nucleoside transport mechanism, intermingling itself within the viral RNA, thus inhibiting/altering viral RNA synthesis. However, ribavirin, particularly when paired with interferonalpha, activates anti-inflammatory responses in various other ways. Alternatively, due to the fact that ribavirin is structurally analogous to GTP, a purine nucleoside, ribavirin can be incorporated into the cell passively, thereafter competitively binding to, and inhibiting, RNA polymerase, and RNA synthesis as a whole; ribavirin often achieves this via blocking the IMPDH pathway, among other pathways such as the eIF4E pathway. Ultimately, five major mechanisms of action have been proposed:^{94–96}

- 1. Immunostimulation by upregulating cytokines to shift Th1/2 cell balance to Th1 dominance.
- 2. Inhibition of 24 kDa eIF4E function, thereby inhibiting mRNA capping and translation initiation.
- 3. Modulation of interferon-alpha-related gene expression.
- 4. Direct inhibition of IMPDH with consequent depletion of intracellular GTP.
- After triphosphorylation, ribavirin triphosphate is incorporated into replicating RNA viral RNA polymerases with consequent induction of viral mutagenesis.

How ribavirin acts vis-a-vis eIF4E is as follows:

eIF4E forms part of the multimeric cap-dependent mRNA translation initiation complex. Mammalian cap-dependent translation starts with that complex binding to an RNA methyl-nucleotide. eIF4E has many positive and negative control points, two of which are 1) posttranslational phosphorylation and 2) 4E-BPs.⁹⁷⁻⁹⁹ There are several variants of 4E-BP protein, hereafter designated simply as 4E-BP. 4E-BP is in turn controlled by its phosphorylation status.

eIF4E non-covalently bound to 4E-BP is inactive in translation initiation. Both currently recognized complexes of mTOR (mTORC1, loosely associated with growth and mTORC2, loosely associated with cell survival and apoptosis resistance) can phosphorylate 4E-BP.¹⁰⁰ Unphosphorylated 4E-BP has non-covalent affinity to and prevents transcription initiation activity of eIF4E. When phosphorylated, 4E-BP1 loses that affinity and separates from eIF4E, thereby allowing eIF4E to function in cap-dependent mRNA translation.^{97,99–101}

MAP kinase interacting kinases (hereafter referred to as MNK) can also phosphorylate 4E-BP, releasing it from eIF-4E.^{99,102–105} A wide variety of internal and extracellular events converge on mTOR and/or MNK to enhance or inhibit their activity. Development of resistance to mTOR inhibitors such as everolimus is often caused by eIF4E amplification or MNK upregulation.^{97,105}

eIF4E overexpression has been identified in 30% of human cancers generally,^{97,106–110} including in invasive breast cancer where the degree of eIF4E, both gene and protein overexpression, has been positively correlated with occurrence, recurrence, and metastasis.^{111–118} eIF4E protein expression was associated with shorter survival, higher tumor mitotic index, and higher-grade breast cancer.¹¹⁵ Increased phosphorylated 4E-BP confers a worse prognosis and faster disease progression in breast, ovary, and prostate cancers.¹⁰³

A crucially important oddity of eIF4E in breast cancer is the homogenous spatial uniformity of phosphorylated eIF4E protein overexpression in breast cancer tissues, both metastatic and primary.¹¹⁶ This is particularly notable given the spatial heterogeneity of ER, PR, HER2, mTOR, and other commonly overexpressed markers in breast cancer.

In addition, in 200 patients with Stage 4 breast cancer, immunohistochemistry analysis revealed that greater increase in eIF4E phosphorylation in response to chemotherapy with doxorubicin, cyclophosphamide, or FU was correlated with shorter median overall survival,¹¹⁴ 4.7 years in patients with a two- to fourfold increase in eIF4E phosphorylation versus 3.1 years in patients with a 9-11-fold increase. A second study a few years later found similar results.¹¹⁸ Among patients undergoing primary debulking for a node-positive breast cancer when nodes were positive, after 4-year followup, systemic recurrence occurred in 22% of women with low eIF4E protein expression, 27% of the intermediate group, and in 49% expressing large amounts of eIF4E.¹¹⁹ Even more serious was the presence of multiple distant metastases in 60% of women whose primary expressed large amounts of eIF4E but in 15% of women whose primary expressed low amounts of eIF4E, again after 4-year follow-up.119

In an unusually exciting and instructive study, Li et al¹²⁰ studied breast cancer biopsy tissue by immunohistochemistry both before and after chemotherapy. After cytotoxic chemotherapy with doxorubicin, or cyclophosphamide or 5-FU, the expression of phosphorylated eIF4E increased in the posttreatment biopsy material, as given in the "Chemotherapy triggers EMT" section, and chemotherapy-activated Wnt/beta-catenin, as given in the "Rifabutin: 847 Da, 2-day half-life" section, signaling in a phosphorylated eIF4E-dependent manner.¹²⁰

Although the significance of eIF4E phosphorylation or its range of functions is not fully understood, some aspects are predominantly the empirical data in the abovementioned paragraph. Regulation of eIF4E function is partly achieved through this phosphorylation process. Untreated GBMs show an excess of phosphorylated (unbound) 4E-BP.^{98,102} Inhibition of 4E-BP phosphorylation with consequent retention of its association with 4E-BP leads to inhibition of protein synthesis, inhibition of glioma cell proliferation in vitro, and tumor growth in vivo, in an orthotopic GBM mouse model.^{98,102} We know ribavirin gets good brain tissue levels based on the psychiatric morbidities associated with its use in treating hepatitis C. Volpin et al¹²¹ suggested using ribavirin to treat GBM based on these considerations.

That metformin inhibits 4E-BP1 phosphorylation via mTOR inhibition¹²² makes metformin a good coordinated partner drug to ribavirin. That ribavirin also inhibits MNK and since MNK phosphorylation of eIF4E is an alternate eIF4E activation pathway particularly used during the development of resistance to the mTOR inhibitor everolimus, 98,99,102,123,124 ribavirin might be combined with everolimus or metformin to advantage. This would be a good example of the phenomenon mentioned in the "Introduction" section that when one growth pathway is pharmacologically blocked other parallel growth-driving pathways can become active, taking the place of blocked paths. mTOR phosphorylates 4E-BP1, or if mTOR is inhibited then MNK can take over, phosphorylating 4E-BP1. This would also explain why and how mTOR inhibitors have not been successful in treating some tumors such as GBM even though they express an overabundance of mTOR. MNK simply takes over when mTOR is blocked.

TGF-beta promotion of EMT that occurs largely through phosphorylation of eIF4E by MNK (with multiple intermediates between the two)¹⁰³ makes pirfenidone a good partner drug for both metformin and ribavirin.

In addition, experimental MNK inhibitors decrease eIF4E phosphorylation levels in breast cancers,¹²⁰ and GBM,⁹⁸ where MNK inhibition enhanced temozolomide cytotoxicity. In parallel fashion, in 103 cases of astrocytomas, high expression of phosphorylated eIF4E was significantly correlated with shorter overall survival rates.¹⁰⁷

All treated breast cancers were found to overexpress phosphorylated (activated) eIF4E,116,120 a remarkable and unique finding in any cancer. Decreased eIF4E phosphorylation in breast cancer also resulted in increased E-cadherin and betacatenin protein levels¹²⁵ reflecting a shift from mesenchymal toward epithelial attributes. The abovementioned combined data suggest that ribavirin could be of potential benefit by inhibiting eIF4E in breast cancer. Kentsis et al^{126,127} have demonstrated that ribavirin inhibits m⁷G mRNA cap binding to eIF4E. Ribavirin directly bound to eIF4E with a micromolar affinity at the functional site used by m7G mRNA cap, reducing eIF4E/mRNA binding and disrupting the translation process. Of note, not all mRNA translation is eIF4E dependent, but important mRNAs in breast cancer are, for instance, the one coding cyclin D1/3, c-Myc, VEGF, FGF2/4, and MCL-1. Some preclinical studies in several murine models of breast cancer revealed that ribavirin inhibits breast cancer cell proliferation through eIF4E blockage.128,129 Moreover, in these studies, multiple-aspect characteristics of EMT were reversed or diminished by ribavirin.128,129

More recently, two studies demonstrated significant glioma cell killing by ribavirin, ^{121,130} confirming a 2014 study

showing that ribavirin induced G0/G1 arrest in seven glioma cell lines at a median 55 μ M IC₅₀ (range 28–664).¹³¹ This latter study positively correlated mRNA expression of PDGF receptor-alpha, a major driver of GBM growth, with better glioma cell sensitivity (lower IC₅₀) to ribavirin. That PDGF receptor is also a major driver of breast cancer^{132,133} and can cross cover for the ER^{134,135} forming one of the many escape paths from aromatase inhibitor suppression of breast cancer growth. This fact favors the possibility of this path contributing to ribavirin's inhibitory effect in breast cancer as well.

Similar to the abovementioned data on breast cancer, targeting eIF4E using ribavirin to block migration and EMT in NSCLC has been highlighted.¹³⁶ In this study, inhibition of eIF4E after ribavirin treatment led to decreased migration, differentiation, and expression of several EMT-related genes such as *ERa*, *SMAD5*, *NF-* κ *B*, *cyclin D1*, *c-MYC*, *or HIF-1a*.¹³⁶ As we expect to do but using ribavirin, an engineered short hairpin RNA interfering with eIF4E transcription inhibited breast cancer cell migration, primary tumor growth, and metastasis establishment.¹³⁷

TGF-beta-induced eIF4E phosphorylation enhanced metastases, invasion, and EMT in a mouse breast cancer model, all of which were inhibited when an un-phosphorylatable eIF4E was present.¹³⁸

IMPDH is a pivotal enzyme for biosynthesis of GTP and is frequently increased in tumor cells.¹³⁹ It has been shown that ribavirin via IMPDH inhibition was effective against chronic lymphocytic leukemia cells.¹⁴⁰ Recently, Isakovic et al¹³⁰ demonstrated in glioma cells that ribavirin inhibits IMPDH activity and induces autophagy inhibiting the activity of mTORC1 and the SRC/AKT pathway.

Of deep significance for understanding breast cancer growth and the ABC7 regimen to inhibit it, is the study by Decarlo et al,¹⁴¹ where they demonstrated a feed-forward amplification loop between TGF-beta and eIF4E (that we intend to block with pirfenidone and ribavirin, respectively). In addition, TGF-beta agonism drives eIF4E activation¹³⁸ confirming pirfenidone as a good partner drug for ribavirin.

Ribavirin has also been shown to inhibit mTOR/eIF4E signaling increasing paclitaxel and imatinib activity in squamous cell carcinoma¹⁴² and leukemic cells,¹²⁴ respectively.

Discussion here of ribavirin strikes at the heart of why pharmaceutical mTOR inhibitors such as everolimus have not been as clinically useful as the biochemistry of cancer indicates it should be. The data in this section paint a consistent picture of eIF4E as a central element in breast cancer malignancy degree and as such a worthwhile target to inhibit. Ribavirin can be expected to do this effectively but it will be the most difficult of the ABC7 drugs to tolerate. Ribavirin is

For personal use only.
the problematic drug of ABC7. When used over months to treat hepatitis C, 1000 mg/day would have been a common dose. Depressed mood, anemia, weight loss, and a severe but ill-defined malaise were common side effects and not rarely required dose reduction or even stopping ribavirin entirely.¹⁴³ Given ribavirin's propensity to give unpleasant side effects, it should be increased with caution from starting dose of 100 mg once daily with frequent mood and CBC evaluations.

Rifabutin: 847 Da, 2-day half-life

Rifabutin is an old antibiotic closely related to the even older drug rifampin (same as rifampicin). Rifabutin is active against *Mycobacterium tuberculosis*, atypical mycobacteria, staphylococci, group A streptococci, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Haemophilus influenzae*, *Haemophilus ducreyi*, *Campylobacter* spp., *Helicobacter pylori*, chlamydia, and *Toxoplama gondii*.^{144,145}

In 2016, rifabutin was reported to have blunted the growth of a patient's NSCLC, subsequently studied in vitro and found to be active in inhibiting lung cancer cell growth and suppressing Ki67 staining.¹⁴⁶ Rifabutin suppressed eIF4E phosphorylation with consequent decreased beta-catenin phosphorylation and increased beta-catenin destruction consequent to that.¹⁴⁶ Thus, rifabutin could coordinate to advantage with ribavirin to thoroughly block eIF4E.

Erlotinib inhibits epidermal growth factor receptors (HER1, EGFR) and is effective initially in stopping some lung cancers' growth. As resistance to erlotinib develops, EGFR mutations resulting in EGFR affinity to beta-catenin, thereby shifting growth drive to beta-catenin system.^{147,148} eIF4E–beta-catenin axis is inhibited by several of the ABC7 drugs.

BCL6 is a 95 kDa protein transcription factor of selected genes, inhibiting expression of some and triggering transcription of others' in cancers generally¹⁴⁹ and in breast cancer specifically.^{150–152} The result is a BCL-6-mediated antiapoptosis effect. Breast cancer cells' survival is enhanced by BCL6.¹⁵¹ Rifabutin binds to BCL6, preventing its function in translation inhibition.¹⁵³ This would be expected to be of benefit on multiple accounts during breast cancer treatment. Interestingly, miR-544 inhibition of BCL6 in TNBC cells inhibited proliferation, migration, and invasion in vitro.¹⁵⁴ BCL6 promoted invasion, migration, and EMT marker expression in breast cancer with indication that greater expression of BCL-6 correlates with shorter overall survival in breast cancer.¹⁵⁰

Not all malignant cells within a strongly ER+ breast cancer will express ERs. The minority population not expressing ERs is relatively chemotherapy resistant with some of that extra chemotherapy resistance mediated by upregulated BCL-6 specifically in that subpopulation.¹⁵⁵

Metformin: 129 Da, not metabolized, 6-hour half-life

Metformin is the most prescribed initial drug treatment for type 2 diabetes worldwide. Despite 60 years of use, the mechanism of action in lowering average glucose is not entirely clear.¹⁵⁶ Hepatic gluconeogenesis is decreased by metformin and insulin sensitivity is increased but how that occurs is uncertain. Metformin in vivo and in vitro increases AMPK, a major regulator of energy homeostasis, metabolism, and protein synthesis.¹⁵⁶ Thus, activated AMPK results in inhibition of mTOR. Breast cancer cell expression of beta-catenin was decreased by metformin concomitantly and proportionately to AMPK phosphorylation.¹⁵⁷

Decreased insulin/insulin-like growth factor-I signaling and inhibition of mitochondrial electron transport chain complex are other documented actions of metformin. Across many cancers, a large chart review has shown decreased mortality in patients treated with metformin.¹⁵⁸ Experimental data support the notions that increased lactate secretion, reduced oxygen consumption, and activated AMPK signaling are plausible mechanisms for metformin's anticancer effects.¹⁵⁹ Metformin also decreased breast cancer cells' intracellular adenosine triphosphate, viability, and anti-apoptotic protein BCL6 concomitant with increased intracellular ROS,¹⁶⁰ the conclusion being that metformin acts primarily on mitochondria, other effects being secondary to that. That work confirmed a related earlier breast cancer study where the mode of viability loss mediated by metformin was found to be by oxidative stress increase and BCL-2 decrease.¹⁶¹

Silvestri et al¹⁶² showed that metformin was indeed cytotoxic to breast cancer cells but 1) only in low glucose conditions – high glucose in vitro could subvert metformin's growth inhibition and 2) although AMPK activation was a requirement for cytotoxicity, mTOR was not. However, Wu et al¹⁶³ showed that metformin both increased lifespan of the nematode *Caenorhabditis elegans* and showed growth inhibition of pancreatic cancer and melanoma cells by an AMPK-independent interference with mitochondrial respiration mechanism. Likewise, Ben Sahra et al¹⁶⁴ demonstrated that metformin cytotoxicity to androgen-sensitive human prostate adenocarcinoma cells was AMPK independent but mTOR inactivation dependent. Furthering complicating delineation of metformin's mechanism of action in treating cancer, Gui et al¹⁶⁵ showed that metformin's anticancer effect

submit your manuscript | www.dovepress.con

Dovepress

was by inhibiting mitochondrial regeneration of oxidized NAD+ regeneration and lowering aspartate levels.

Just in 2016, five extensive reviews appeared recounting evidence favoring the use of metformin as treatment adjunct in cancer generally.^{166–170}

In a study particularly relevant to ABC7 regimen considering that capecitabine metabolizes into 5-FU within cancer cells, Qu et al¹⁷¹ showed that breast cancer cells that had become resistant to 5-FU regained cytotoxic sensitivity to 5-FU by simultaneous exposure with metformin. Metformin synergy with 5-FU could also be demonstrated to breast cancer cells in both the stem and non-stem subpopulations.¹⁷² Of central importance to the ABC7 regimen, IC₅₀ of 5-FU to esophageal cancer cells was lowered by metformin¹⁷³ and correlated with increased AMPK activation and decreased mTOR function and lactate production. Metformin plus 5-FU combination was also active in slowing esophageal cancer growth in a xenotransplant model more than either agent alone.¹⁷⁴

YAP is a small protein transcription factor promoting the growth of many cancers. When phosphorylated, it is retained in cytoplasm and therefore nonfunctional in promoting growth or inhibiting apoptosis. Metformin treatment of hepa-tocellular carcinoma patients increased YAP phosphorylation via AMPK phosphorylation and prolonged survival, half deceased at ~31 months without compared to ~44 months with metformin.¹⁷⁵ Adding metformin to exemestane also increased survival in ER+ breast cancers that overexpressed IGF1R.¹⁷⁶

An ongoing trial (ClinicalTrials.gov Identifier NCT01589367) is studying potential survival benefits of adding metformin 2000 mg/day to standard antiestrogen aromatase inhibitor, letrozole 2.5 mg/day, in nondiabetic postmenopausal women with ER+ breast cancer.

Preoperative treatment of breast cancer patients with metformin has given mixed results. Some studies showed reduced mitotic rate after metformin 2000 mg/day¹⁷⁷ and 1500 mg/ day,¹⁷⁸ while others showed no reduction using 1500 mg/ day.¹⁷⁹ A similarly designed study using 1700 mg/day found marginally lower Ki67 only in women with increased insulin resistance.¹⁸⁰

A pivotal study supporting metformin use during the treatment of breast cancer was reported back in 2011. In women undergoing primary resection for breast cancer, 1 g twice daily metformin was given 14 days prior to surgery.¹⁷⁷ By immunohistochemistry, the diagnostic biopsy was compared to resected tissue for p-AMPK, p-AKT, insulin receptor, cleaved caspase-3, and Ki67. In metformin pretreated,

increased p-AMPK and decreased p-AKT were seen compared to those not treated with metformin in the interval between biopsy and surgery. Ki67 and cleaved caspase-3 were diminished in metformin-treated women compared to those not so treated. These changes were not large but were statistically significant and large enough to expect some clinical benefit.¹⁷⁷

Although metformin decreases breast cancer cell survival in vitro,^{160–162,181} the clinical benefit would seem small given the equivocal human trials and evidence that the small benefit seen tended to be restricted to diabetic/prediabetic people. However, small benefit is not no benefit.

Metformin despite being hydrophilic achieves approximately equal plasma and brain tissue levels. In rats, after single-dose oral metformin administration, 28 µmol/L plasma and 14 nmol/g brain tissue (14 µM) were seen.¹⁸² Average metformin plasma levels typically seen in asymptomatic diabetes patients were 2.7 ± 7.3 mg/L, \sim 3 µM. The unusually wide drug range seen, ±7.3 mg/L (±57 µM), reflects metformin's safety.¹⁸³ Metformin's side effects are limited to diarrhea, nausea, and vomiting. Some cases of lactic acidosis could occur but at a low frequency and when metformin is implicated as the cause of lactic acidosis, metformin plasma levels greater than 5 µg/mL are generally found. Target dose of metformin is the standard dose used in past breast cancer studies of metformin – 1700–2000 mg/day.

Propranolol: 259 Da, cyp 1A2, 2D6, 9-hour half-life

Propranolol was the first beta-blocker introduced to clinical practice. Introduced in the 1960s, it is still in wide use to treat hypertension, migraine, angina, selected arrhythmias, essential tremor, resolution of infantile hematomas, and in reducing the cardiac effects (tachycardia) due to acute anxiety. Propranolol's general cancer process inhibiting attributes were recently reviewed.¹⁸⁴ Below are selected data supporting propranolol's use specifically as adjunct in breast cancer treatment.

A study of 404 breast cancer patients to compare the proliferation rates of breast cancers in women who had taken beta-blockers compared to those who had not found a clear reduction in Ki67 only in those with Stage 1 disease.¹⁸⁵ A single ER+, HER2– patient was treated with 25 days of propranolol 1.5 mg/kg per day after diagnostic biopsy but before resection. Resection of tumor tissue showed a 23% reduction in Ki67 staining compared to biopsy tissue 25 days earlier, before any propranolol.¹⁸⁵ Of important note, beta-1-selective beta-blockers did not work to reduce Ki67, only

nonselective beta-blockers did. However, a large European epidemiological study found no survival benefit from propranolol use after a breast cancer diagnosis.¹⁸⁶

Of particular interest to ABC7 regimen, Rico et al¹⁸⁷ examined the effects of metformin and propranolol singly and combined in several preclinical TNBC models, finding additive to synergistic growth-inhibiting effects.

In a cohort of 800 women with early TNBC, 9% used beta-blockers. The beta-blocker use and nonuse groups were well matched. At 5 years, 19% of the nonusers had died of breast cancer while 8% of beta-blocker users had died of breast cancer.¹⁸⁸

In examining a cohort of 1971 multiple myeloma patients, those who took any beta-blocker, had a 24% disease-specific mortality at 5 years. Those who took a beta-blocker plus other cardiac drugs had 32% while those on no cardiac or blood pressure medicines had 41% myeloma-specific mortality at 5 years.¹⁸⁹

An interesting study from Choy et al¹⁹⁰ showed that among 1000 breast cancer patients those on a beta-blocker had a lower recurrence rate, and specifically TNBC expressed particularly high levels of beta-adrenergic receptors. Their brain metastases expressed more beta-adrenergic receptors per cell than did the primary tumors.¹⁹⁰ This study also gave evidence of propranolol's inhibition of proliferation and migration in breast cancer cells expressing the beta-adrenergic receptor.

In reviewing seven epidemiological studies prior to 2015 on beta-blocker use in breast cancer, Childers et al¹⁹¹ concluded that, although results were mixed between these studies, slightly lower risk of death was associated with beta-blocker use. Beta-blocker use is associated with improved relapse-free survival (but not in overall survival) also in patients with TNBC.¹⁹²

Bone is richly supplied with sympathetic nerve endings. When specifically osteoblasts' beta-adrenergic receptor is stimulated by norepinephrine from these nerve endings, the osteoblasts secrete RANKL.¹⁹³ Thus, propranolol should harmonize with quetiapine (vide supra) in treating and preventing bone metastases in breast cancer. CA125 is a high molecular weight mucin commonly elevated in ovarian cancer. Patients given perioperative propranolol showed an 83% CA125 decrease on postoperative day 7 when those given placebo had a 72% decrease.¹⁹⁴

Although the data were mixed, a review of 10 studies completed by 2015 of beta-blocker use in breast cancer concluded that specifically propranolol use was indeed associated with slightly reduced breast cancer-specific mortality.¹⁹⁵ Propranolol-blocked beta-adrenergic agonist induced increased migration and decreased breast cancer cell-to-cell adhesion.¹⁹⁶ Propranolol inhibited breast cancer cell migration in vitro.¹⁹⁷ Breast cancer cells express beta-adrenergic receptors. Blocking these with propranolol lowers their glucose uptake.¹⁹⁸

Campbell et al¹⁹⁹ demonstrated that beta-adrenergic stimulation of bone increased osteoblasts' RANKL expression. That induced RANKL increased breast cancer establishment of metastases in bone.¹⁹⁹ Thus, the combination with quetiapine might be particularly beneficial.

Beta-adrenergic stimulation did not change the growth of an orthotopic murine breast cancer but did induce a remarkable 30-fold increase in metastases, an effect partially blocked by propranolol.²⁰⁰ Of clinical importance to ABC7, Shaashua et al²⁰¹ showed that combining propranolol with a COX-2 inhibitor in perioperative breast cancer decreased EMT, serum IL-6, and C-reactive protein levels.

There is risk of symptomatic iatrogenic hypotension with propranolol. The propranolol dose must therefore be slowly uptitrated as tolerated, monitoring blood pressure.

Capecitabine: 359 Da, half-life <1 hour

Capecitabine is a 359 Da pro-drug giving rise to intracellular release of 130 Da 5-FU.^{202,203} 5-FU inhibits thymidylate synthase, which mediates the synthesis of thymidine monophosphate, the active form of thymidine required DNA synthesis.

Despite ~20 years of clinical use in treating breast cancer, there remains some unclarity on the ideal dosing schedule for capecitabine.^{204–206} A comparison of cycles of 1000 mg/m² twice daily for 14 days, 7 days off with 1250 mg/m² twice daily for 14 days, and 7 days off indicated lower side effect burden with 1000 mg/m² twice daily.^{207,208}

Several reports indicate that dosing capecitabine at just high enough level to generate palmar–plantar erythrodysesthesia might be most effective dosing regimen.^{206,209} This would be analogous to erlotinib dosing where titrating to rash might be most effective.²¹⁰

Capecitabine is best given with Coke[™] or fresh squeezed lemon juice to assure low enough gastric pH for adequate and uniform absorption. This would be particularly important for those on proton pump inhibitors.

Principle toxicity is palmar–plantar erythrodysesthesia (synonyms hand-foot syndrome or chemotherapy induced acral erythema), diarrhea, and nausea, although cytopenias, fatigue, dyspnea, or cardiac abnormalities can be seen.²¹¹ The common dose for capecitabine in breast cancer is 1250 mg/m² orally twice daily for 14 days, none for 7 days, every 21 days.

Agomelatine: 243 Da, 2-hour half-life

Agomelatine is a 243 Da pharmaceutical melatonergic agonist at both melatonin's receptors, M1 and M2.²¹² It has many advantages over the use of melatonin itself^{213,214} In short, these advantages are: 1) agomelatine is Health Canada and EMA approved and marketed as an antidepressant. As such, it is a well-standardized product, as opposed to over-thecounter melatonin preparations which are exempt from the strict standards of approved medicines; 2) agomelatine has considerably tighter affinity to both M1 and M2 receptors than does the natural ligand (melatonin); 3) agomelatine has a much longer dwell time in the body than does melatonin, and; 4) absorption is more uniform and reliable than is absorption of melatonin.

Although agomelatine is available for import into the USA, it is not FDA approved. Ramelteon is an equally potent melatonergic agonist at M1 and M2 as is agomelatine. Ramelteon is FDA approved and marketed in the USA. It has similar actions and advantages over melatonin as does agomelatine^{213,215} and can be substituted for agomelatine in the ABC7 regimen.

Elevation of hepatic transaminases is of potential concern when using agomelatine. This requires regular monitoring. Elevation is dose dependent, occurring in $\sim 3\%$ of those receiving 50 mg once at bedtime.²¹⁶ It is usually reversible.

Work pointing to diminished breast cancer cell malignant behavior during exposure to melatonin dates back at least 3 decades.²¹⁷ There are numerous studies about oncostatic effects of melatonin on several tumors as well as recent reviews summarizing the different mechanisms of cancer inhibition by melatonin.^{84,218–221} These include regulation of estrogen pathway, melatonin as SERM and SEEM, modulation of the cell cycle, differentiation and the induction of apoptosis, inhibition of telomerase activity, inhibition of oxidative stress, inhibition of angiogenesis, regulation of circadian rhythms, avoidance of circadian disruption, inhibition of tumor metastasis, invasiveness and motility decline, and enhancement of immune system and epigenetic regulation.^{218,221}

Briefly and empirically, melatonin has readily demonstrable growth-inhibiting effects in both in vivo animal models, with chemically induced mammary tumors in rodents, and in vitro assays in estrogen-positive human breast cancer cells.^{221–224} Melatonin inhibits invasive and metastatic properties of human breast cancer cells in different xenograft models.^{225–228} Due to the broad spectrum of melatonin's actions, the mechanisms through which it interferes with metastases are varied. These include modulation of cell–cell and cell–matrix interaction, extracellular matrix remodeling by matrix metalloproteinases, cytoskeleton reorganization, EMT, and angiogenesis.²²⁹

Melatonin shifts human breast cancer cells to a lower invasive status by upregulating E-cadherin and B1-integrin expression and decreasing OCT4, N-cadherin, and vimentin.^{219,227,228,230} These findings suggest that melatonin modulates both cell-cell and cell-matrix interactions in breast cancer and reduces the metastatic potential of the tumor. Melatonin also has regulatory actions on matrix metalloproteinases in breast cancer. It has been described that melatonin inhibits the induction, catalytic activity, and expression of MMP-9 and MMP-2.^{229,231} In addition to modulating the metalloproteinase activity, melatonin reduces cancer cell migration through the downregulation of ROCK-1 and MCLCK, two kinases that control the cytoskeletal rearrangement associated with cell-cell and cell-matrix adhesion.229,232 The attenuation of HER2-Rsk2 signaling by melatonin plays a main role in the melatonin-mediated suppression of EMT and late-staged metastasis in breast cancer cells.226,229

In tumor angiogenesis, there is a crosstalk between cancer cells and surrounding endothelial cells. Melatonin interferes in the paracrine interactions between malignant epithelial cells and proximal endothelial cells through a downregulatory action on VEGF expression in human breast cancer cells, which decrease the levels of VEGF around endothelial cells.^{230,231} In addition, melatonin directly exerts antiangiogenic actions by reducing endothelial cell proliferation, invasion, migration, and tube formation, through a downregulation of VEGF expression.^{219,233–235} Melatonin also impedes the EMT process and cancer cell dissemination through downregulatory actions of the p38 pathway²²⁷ and interferences with NF- κ B signaling in tumor cells.^{217,229,236}

Recently, a review of the effects of melatonin and chemotherapeutic agents in combination in cancer treatments has been published.²³⁷ Although the information available is limited, the results obtained suggest that melatonin sensitizes tumor cells to the cytotoxic effects of chemotherapeutic agents.

In addition, in a rat ER+ breast cancer model, melatonin reduced tumor weight, prolonged survival, and increased E-cadherin without giving apparent side effects.²³⁸ In this model, doxorubicin cytotoxicity to the breast cancers was augmented by giving simultaneous melatonin.²³⁸ Melatonin reduced in vitro migration and in vivo growth, proliferation index, and metastases in a murine xenograft model.²³²

Of particular relevance to ABC7, earlier in year 2017, melatonin was shown to increase 5-FU inhibition of colon cancer cell proliferation, in vitro colony formation, migration,

506

and invasion, showing a corresponding in vivo synergy with 5-FU in colon cancer tumor growth inhibition in a xenograft model.²³⁹

Similarly, melatonin moderately enhanced cytotoxicity to cisplatin and doxorubicin, while slightly but significantly enhancing 5-FU cytotoxicity to HeLa cells.²⁴⁰ In an in vitro study, in rat pancreatic adenocarcinoma, melatonin augmented cytotoxicity of 5-FU, cisplatin, and doxorubicin.²⁴¹ Melatonin decreased pancreas cancers in hamsters given a carcinogen (*N*-nitrosobis (2-oxopropyl) amine), as did capecitabine. Giving both melatonin and capecitabine decreased this incidence further.²⁴² Melatonin augmented doxorubicin cytotoxicity to lymphocytic leukemia cells without having cytotoxicity to normal lymphocytes.²⁴³

Melatonin sensitized human breast cancer cells to radiation via 1) reduction in estrogen-synthesizing proteins, and 2) induction of a twofold change in p53 expression, and 3) downregulation of proteins involved in double-strand DNA break repair, such as RAD51 and DNA-PKcs.²⁴⁴ Melatonin enhanced cytotoxicity of 5-FU to esophageal squamous carcinoma cells both in vitro and in a xenograft model.245 These authors used 20 mg/ kg per day melatonin in the xenograft model, corresponding to a nominal 1400 mg/day for a 70 kg adult human. The common over-the-counter melatonin used is 3-20 mg once at bedtime. The tighter affinity to melatonin receptors and much longer half-life of agomelatine compared to melatonin would go some of the way toward generating a stronger agonist signal to M1 and M2 than today's commonly used melatonin doses. Another felicitous aspect of melatonergic agonism is a potential increase in NK cell numbers and function.246

A remarkable epidemiological study of cancer-free postmenopausal women showed that higher urinary melatonin levels were associated with a slightly reduced risk of later developing breast cancer,²⁴⁷ although these data are not uncontested. A review of all studies on urinary melatonin would indicate that this matter remains unsettled.²⁴⁸

The suggested dose of agomelatine is 50 mg once at bedtime, twice the EMA and Health Canada recommended dose for treating depression. If ramelteon is used instead 16 mg at bedtime, twice the FDA-approved dose is recommended.

Conclusion

Once breast cancer has metastasized to bone, liver, or lungs, the prognosis becomes poor. No current treatment has a reliable and robust disease control rate at that point.

Animal study of the complete ABC7 regimen would be advisable. Based on clinical experience with these drugs individually and in pairs in general medical practice, the predicted safety and tolerability of the ABC7 regimen should be safe. As a further safety measure, the ABC7 drugs should be added one at a time at weekly intervals, thereby catching any unwanted interactions early and the offending drug more easily identified.

In this article, we propose that seven common and already FDA-approved drugs, such as agomelatine (or ramelteon), metformin, pirfenidone, propranolol, quetiapine, ribavirin, and rifabutin, can have the ability to reduce EMT and breast cancer cell tumorigenesis. These ancillary drugs have demonstrated that attributes that we have reason to believe will inhibit EMT and enhance capecitabine's efficacy. The predicted safety and tolerability of the ABC7 regimen is good. A clinical trial is warranted given the fatal outcome of metastatic breast cancer as things now stand.

Abbreviations

ABC7, Adjuvant for Breast Cancer treatment using seven repurposed drugs; AMPK, AMP-activated protein kinase; BCL-6, B-cell lymphoma-6; CBC, complete blood count; COMBAT regimen, combined oral metronomic biodifferentiating antiangiogenic treatment regimen; CSCs, cancer stem cells; CTCs, circulating tumor cells; CUSP9 regimen, coordinated undermining of survival paths with nine regimens; EGFR, epidermal growth factor receptor; EMA, European Medicines Agency; EMT, epithelial-to-mesenchymal transition; ER, estrogen receptor; ER+, ER positive; 4E-BP, eIF4Ebinding protein; 4E-BP1, eIF4E-binding protein-1; FDA, Food and Drug Administration; 5-FU, 5-fluorouracil; GBM, glioblastoma; GM-CSF, granulocyte-monocyte colonystimulating factor; GTP, guanosine-5'-triphosphate; HER2, human epidermal growth factor receptor 2; Hh, hedgehog; HR, hormone receptor; HR+, hormone receptor positive; HR-, hormone receptor negative; IGF1R, insulin-like growth factor type 1 receptor; IL-6, interleukin-6; IMPDH, inosine monophosphate dehydrogenase; MCLCK, myosin lightchain kinase; MEMMAT regimen, metronomic and targeted anti-angiogenesis therapy regimen; MET, mesenchymal-toepithelial transition; MNK, MAP kinase-interacting kinase; mTOR, mammalian target of rapamycin; MTZ regimen, minocycline telmisartan and zoledronic acid regimen; m⁷G, 7-methylguanosine; NAD+, nicotinamide adenine dinucleotide; NCATS, National Center for Advancing Translational Sciences; NK, natural killer; NSCLC, non-small cell lung cancer; OPG, osteoprotegerin; PDGF, platelet-derived growth factor; PR, progesterone receptor; RANK, receptor activator of nuclear factor-κB; RANKL, RANK ligand; ROCK-1, rho-associated protein kinase; ROS, reactive oxygen species; SEEM, selective estrogen enzyme modulator; SERM, selective estrogen receptor modulator; TGF-beta, transforming growth factor-beta; Th, T helper; TNBC, triple-negative breast cancer; ZO-1, zonula occludens-1.

Acknowledgments

SC was supported by grants from the Spanish Economy and Competitiveness Ministry (SAF2016-77103-P) and from Instituto de Investigación Sanitaria Valdecilla (IDIVAL) (APG/12). YS was supported by the Department of Defense (W81XWH-14-1-0403), the Wake Forest School of Medicine Internal Pilot Funding, and the Translational Research Academy which is supported by the NCATS, National Institutes of Health, through Grant Award Number UL1TR001420.

Disclosure

The authors report no conflicts of interest in this work.

References

- Jerusalem G, Bachelot T, Barrios C, et al. A new era of improving progression-free survival with dual blockade in postmenopausal HR(+), HER2(-) advanced breast cancer. *Cancer Treat Rev.* 2015;41(2):94–104.
- 2. Schulz M, Iwersen-Bergmann S, Andresen H, Schmoldt A. Therapeutic and toxic blood concentrations of nearly 1,000 drugs and other xenobiotics. *Crit Care*. 2012;16(4):R136.
- Salacz ME, Kast RE, Saki N, Brüning A, Karpel-Massler G, Halatsch ME. Toward a noncytotoxic glioblastoma therapy: blocking MCP-1 with the MTZ regimen. *Onco Targets Ther.* 2016;9:2535–2545.
- Zapletalova D, Andre N, Deak L, et al. Metronomic chemotherapy with the COMBAT regimen in advanced pediatric malignancies: a multicenter experience. *Oncology*. 2012;82(5):249–260.
- Kast RE, Boockvar JA, Brüning A, et al. A conceptually new treatment approach for relapsed glioblastoma: coordinated undermining of survival paths with nine repurposed drugs (CUSP9) by the International Initiative for Accelerated Improvement of Glioblastoma Care. *Oncotarget*. 2013;4(4):502–530.
- Kast RE, Karpel-Massler G, Halatsch ME. CUSP9* treatment protocol for recurrent glioblastoma: aprepitant, artesunate, auranofin, captopril, celecoxib, disulfiram, itraconazole, ritonavir, sertraline augmenting continuous low dose temozolomide. *Oncotarget*. 2014;5(18):8052–8082.
- Chebouti I, Kasimir-Bauer S, Buderath P, et al. EMT-like circulating tumor cells in ovarian cancer patients are enriched by platinumbased chemotherapy. *Oncotarget*. Epub 2017 Mar 14. doi: 10.18632/ oncotarget.16179.
- Jin X, Jiang YZ, Chen S, Yu KD, Shao ZM, Di GH. Prognostic value of receptor conversion after neoadjuvant chemotherapy in breast cancer patients: a prospective observational study. *Oncotarget*. 2015;6(11):9600–9611.
- Toss A, Venturelli M, Peterle C, Piacentini F, Cascinu S, Cortesi L. Molecular biomarkers for prediction of targeted therapy response in metastatic breast cancer: trick or treat? *Int J Mol Sci.* 2017;18(1):E85. doi: 10.3390/ijms18010085.
- Sambi M, Haq S, Samuel V, et al. Alternative therapies for metastatic breast cancer: multimodal approach targeting tumor cell heterogeneity. *Breast Cancer (Dove Med Press)*. 2017;9:85–93.
- Jonsson VD, Blakely CM, Lin L, et al. Novel computational method for predicting polytherapy switching strategies to overcome tumor heterogeneity and evolution. *Sci Rep.* 2017;7:44206.

- Blakely CM, Watkins TBK, Wu W, et al. Evolution and clinical impact of genetic epistasis within EGFR-mutant lung cancers. *bioRxiv*. 2017. Available from: http://biorxiv.org/content/biorxiv/ early/2017/03/16/117291.full.pdf. Accessed March 16, 2017.
- 13. Yeung KT, Yang J. Epithelial-mesenchymal transition in tumor metastasis. *Mol Oncol.* 2017;11(1):28–39.
- Creighton CJ, Gibbons DL, Kurie JM. The role of epithelialmesenchymal transition programming in invasion and metastasis: a clinical perspective. *Cancer Manag Res.* 2013;5:187–195.
- 15. Wu Y, Sarkissyan M, Vadgama JV. Epithelial-mesenchymal transition and breast cancer. *J Clin Med.* 2016;5(2):E13.
- Mock K, Preca BT, Brummer T, Brabletz S, Stemmler MP, Brabletz T. The EMT-activator ZEB1 induces bone metastasis associated genes including BMP-inhibitors. *Oncotarget*. 2015;6(16):14399–14412.
- Preca BT, Bajdak K, Mock K, et al. A novel ZEB1/HAS2 positive feedback loop promotes EMT in breast cancer. *Oncotarget*. 2017;8(7):11530–11543.
- Tan EJ, Olsson AK, Moustakas A. Reprogramming during epithelial to mesenchymal transition under the control of TGFβ. *Cell Adh Migr*. 2015;9(3):233–246.
- 19. Sato R, Semba T, Saya H, Arima Y. Concise review: stem cells and epithelial-mesenchymal transition in cancer: biological implications and therapeutic targets. *Stem Cells*. 2016;34(8):1997–2007.
- 20. Chaffer CL, San Juan BP, Lim E, Weinberg RA. EMT, cell plasticity and metastasis. *Cancer Metastasis Rev.* 2016;35(4):645–654.
- Yeung SF, Tong JH, Law PP, et al. Profiling of oncogenic driver events in lung adenocarcinoma revealed MET mutation as independent prognostic factor. *J Thorac Oncol.* 2015;10(9):1292–1300.
- 22. Lee Y, Jung WH, Koo JS. Adipocytes can induce epithelialmesenchymal transition in breast cancer cells. *Breast Cancer Res Treat*. 2015;153(2):323–335.
- 23. Ingthorsson S, Briem E, Bergthorsson JT, Gudjonsson T. Epithelial plasticity during human breast morphogenesis and cancer progression. *J Mammary Gland Biol Neoplasia*. 2016;21(3–4):139–148.
- Bulfoni M, Gerratana L, Del Ben F, et al. In patients with metastatic breast cancer the identification of circulating tumor cells in epithelialto-mesenchymal transition is associated with a poor prognosis. *Breast Cancer Res.* 2016;18(1):30.
- 25. Bae YK, Choi JE, Kang SH, Lee SJ. Epithelial-mesenchymal transition phenotype is associated with clinicopathological factors that indicate aggressive biological behavior and poor clinical outcomes in invasive breast cancer. *J Breast Cancer*. 2015;18(3):256–263.
- 26. Jin MS, Hyun CL, Park IA, et al. SIRT1 induces tumor invasion by targeting epithelial mesenchymal transition-related pathway and is a prognostic marker in triple negative breast cancer. *Tumour Biol.* 2016;37(4):4743–4753.
- Kotiyal S, Bhattacharya S. Epithelial mesenchymal transition and vascular mimicry in breast cancer stem cells. *Crit Rev Eukaryot Gene Expr.* 2015;25(3):269–280.
- 28. Wu ZH, Tao ZH, Zhang J, et al. MiRNA-21 induces epithelial to mesenchymal transition and gemcitabine resistance via the PTEN/AKT pathway in breast cancer. *Tumour Biol.* 2016;37(6): 7245–7254.
- Scimeca M, Antonacci C, Colombo D, Bonfiglio R, Buonomo OC, Bonanno E. Emerging prognostic markers related to mesenchymal characteristics of poorly differentiated breast cancers. *Tumour Biol.* 2016;37(4):5427–5435.
- Bhowmik SK, Ramirez-Pena E, Arnold JM, et al. EMT-induced metabolite signature identifies poor clinical outcome. *Oncotarget*. 2015;6(40):42651–42660.
- Fernandez-Garcia B, Eiro N, Marin L, et al. Expression and prognostic significance of fibronectin and matrix metalloproteases in breast cancer metastasis. *Histopathology*. 2014;64:512–522.
- Zhou X, Zhang J, Yun H, et al. Alterations of biomarker profiles after neoadjuvant chemotherapy in breast cancer: tumor heterogeneity should be taken into consideration. *Oncotarget*. 2015;6(34): 36894–36902.

- Park SY, Kim MJ, Park SA, et al. Combinatorial TGF-beta attenuation with paclitaxel inhibits the epithelial-to-mesenchymal transition and breast cancer stem-like cells. *Oncotarget*. 2015;6(35):37526–37543.
- Choi HS, Cho SG, Kim MK, et al. SH003 enhances paclitaxel chemosensitivity in MCF-7/PAX breast cancer cells through inhibition of MDR1 activity. *Mol Cell Biochem*. 2017;426(1–2):1–8.
- Xu J, Liu D, Niu H, et al. Resveratrol reverses doxorubicin resistance by inhibiting epithelial-mesenchymal transition (EMT) through modulating PTEN/Akt signaling pathway in gastric cancer. *J Exp Clin Cancer Res.* 2017;36(1):19.
- Yoshimasu T, Ohashi T, Oura S, et al. A theoretical model for the hormetic dose-response curve for anticancer agents. *Anticancer Res.* 2015;35(11):5851–5855.
- Mathenge EG, Dean CA, Clements D, et al. Core needle biopsy of breast cancer tumors increases distant metastases in a mouse model. *Neoplasia*. 2014;16(11):950–960.
- Szalayova G, Ogrodnik A, Spencer B, et al. Human breast cancer biopsies induce eosinophil recruitment and enhance adjacent cancer cell proliferation. *Breast Cancer Res Treat*. 2016;157(3):461–474.
- Juratli MA, Siegel ER, Nedosekin DA, et al. In vivo long-term monitoring of circulating tumor cells fluctuation during medical interventions. *PLoS One.* 2015;10(9):e0137613.
- 40. Kaigorodova EV, Tarabanovskaya NA, Staheeva MN, et al. Effect of small and radical surgical injury on the level of different populations of circulating tumor cells in the blood of breast cancer patients. *Neoplasma*. 2017;64(3):437–443. doi: 10.4149/neo_2017_315.
- 41. Tsumura H, Satoh T, Ishiyama H, et al. Perioperative search for circulating tumor cells in patients undergoing prostate brachytherapy for clinically nonmetastatic prostate cancer. *Int J Mol Sci.* 2017;18(1):E128.
- Hara N, Kasahara T, Kawasaki T, et al. Frequency of PSA-mRNAbearing cells in the peripheral blood of patients after prostate biopsy. *Br J Cancer*. 2001;85(4):557–562.
- 43. Dyavanagoudar S, Kale A, Bhat K, Hallikerimath S. Reverse transcriptase polymerase chain reaction study to evaluate dissemination of cancer cells into circulation after incision biopsy in oral squamous cell carcinoma. *Indian J Dent Res.* 2008;19(4):315–319.
- 44. Kusukawa J, Suefuji Y, Ryu F, Noguchi R, Iwamoto O, Kameyama T. Dissemination of cancer cells into circulation occurs by incisional biopsy of oral squamous cell carcinoma. *J Oral Pathol Med.* 2000;29(7):303–307.
- 45. Moncharmont C, Levy A, Guy JB, et al. Radiation-enhanced cell migration/invasion process: a review. *Crit Rev Oncol Hematol*. 2014;92(2):133–142.
- 46. Lee HM, Hwang KA, Choi KC. Diverse pathways of epithelial mesenchymal transition related with cancer progression and metastasis and potential effects of endocrine disrupting chemicals on epithelial mesenchymal transition process. *Mol Cell Endocrinol*. Epub 2016 Dec 29.
- Kast RE, Hill QA, Wion D, et al. Glioblastoma-synthesized G-CSF and GM-CSF contribute to growth and immunosuppression: potential therapeutic benefit from dapsone, fenofibrate, and ribavirin. *Tumour Biol.* 2017;39(5):1010428317699797. doi: 10.1177/1010428317699797.
- 48. Leyland-Jones B, Bondarenko I, Nemsadze G, et al. A randomized, open-label, multicenter, phase III study of epoetin alfa versus best standard of care in anemic patients with metastatic breast cancer receiving standard chemotherapy. *J Clin Oncol.* 2016;34(11):1197–1207.
- Vilalta M, Rafat M, Giaccia AJ, Graves EE. Recruitment of circulating breast cancer cells is stimulated by radiotherapy. *Cell Rep.* 2014;8(2):402–409.
- Kim RK, Kaushik N, Suh Y, et al. Radiation driven epithelialmesenchymal transition is mediated by Notch signaling in breast cancer. *Oncotarget*. 2016;7(33):53430–53442.
- Lee SY, Jeong EK, Ju MK, et al. Induction of metastasis, cancer stem cell phenotype, and oncogenic metabolism in cancer cells by ionizing radiation. *Mol Cancer*. 2017;16(1):10.

- Zhao R, Cai Z, Li S, et al. Expression and clinical relevance of epithelial and mesenchymal markers in circulating tumor cells from colorectal cancer. *Oncotarget*. 2017;8(6):9293–9302.
- Zheng X, Fan L, Zhou P, et al. Detection of circulating tumor cells and circulating tumor microemboli in gastric cancer. *Transl Oncol.* 2017;10(3):431–441.
- 54. Zhang D, Zhao L, Zhou P, et al. Circulating tumor microemboli (CTM) and vimentin+ circulating tumor cells (CTCs) detected by a size-based platform predict worse prognosis in advanced colorectal cancer patients during chemotherapy. *Cancer Cell Int.* 2017;17:6.
- Kolbl AC, Jeschke U, Andergassen U. The significance of epithelialto-mesenchymal transition for circulating tumor cells. *Int J Mol Sci.* 2016;17(8):E1308.
- Kiesel L, Kohl A. Role of the RANK/RANKL pathway in breast cancer. Maturitas. 2016;86:10–16.
- 57. Wang H, Shen W, Hu X, et al. Quetiapine inhibits osteoclastogenesis and prevents human breast cancer-induced bone loss through suppression of the RANKL-mediated MAPK and NF-κB signaling pathways. *Breast Cancer Res Treat*. 2015;149(3):705–714.
- Palafox M, Ferrer I, Pellegrini P, et al. RANK induces epithelialmesenchymal transition and stemness in human mammary epithelial cells and promotes tumorigenesis and metastasis. *Cancer Res.* 2012;72(11):2879–2888.
- Beleut M, Rajaram RD, Caikovski M, et al. Two distinct mechanisms underlie progesterone-induced proliferation in the mammary gland. *Proc Natl Acad Sci U S A*. 2010;107(7):2989–2994.
- 60. Kan C, Vargas G, Pape FL, Clézardin P. Cancer cell colonisation in the bone microenvironment. *Int J Mol Sci.* 2016;17(10):E1674.
- Yoneda T, Tanaka S, Hata K. Role of RANKL/RANK in primary and secondary breast cancer. *World J Orthop*. 2013;4(4):178–185.
- 62. Futakuchi M, Fukamachi K, Suzui M. Heterogeneity of tumor cells in the bone microenvironment: mechanisms and therapeutic targets for bone metastasis of prostate or breast cancer. *Adv Drug Deliv Rev.* 2016;99(pt B):206–211.
- 63. Lamora A, Talbot J, Mullard M, Brounais-Le Royer B, Redini F, Verrecchia F. TGF- β signaling in bone remodeling and osteosarcoma progression. *J Clin Med.* 2016;5(11):E96.
- 64. King TE Jr, Bradford WZ, Castro-Bernardini S, et al; ASCEND Study Group. A phase 3 trial of pirfenidone in patients with idiopathic pulmonary fibrosis. *N Engl J Med.* 2014;370(22):2083–2092.
- 65. Nathan SD, Albera C, Bradford WZ, et al. Effect of continued treatment with pirfenidone following clinically meaningful declines in forced vital capacity: analysis of data from three phase 3 trials in patients with idiopathic pulmonary fibrosis. *Thorax*. 2016;71(5):429–435.
- 66. Conte E, Gili E, Fagone E, Fruciano M, Iemmolo M, Vancheri C. Effect of pirfenidone on proliferation, TGF-β-induced myofibroblast differentiation and fibrogenic activity of primary human lung fibroblasts. *Eur J Pharm Sci.* 2014;58:13–19.
- Du J, Paz K, Flynn R, et al. Pirfenidone ameliorates murine chronic GVHD through inhibition of macrophage infiltration and TGF-beta production. *Blood*. 2017;129(18):2570–2580.
- Stahnke T, Kowtharapu BS, Stachs O, et al. Suppression of TGF-beta pathway by pirfenidone decreases extracellular matrix deposition in ocular fibroblasts in vitro. *PLoS One*. 2017;12(2):e0172592.
- Didiasova M, Singh R, Wilhelm J, et al. Pirfenidone exerts antifibrotic effects through inhibition of GLI transcription factors. *FASEB J*. 2017;31(5):1916–1928.
- Yu W, Guo F, Song X. Effects and mechanisms of pirfenidone, prednisone and acetylcysteine on pulmonary fibrosis in rat idiopathic pulmonary fibrosis models. *Pharm Biol.* 2017;55(1):450–455.
- Fujiwara A, Shintani Y, Funaki S, et al. Pirfenidone plays a biphasic role in inhibition of epithelial-mesenchymal transition in non-small cell lung cancer. *Lung Cancer*. 2017;106:8–16.
- 72. Cantelli G, Crosas-Molist E, Georgouli M, Sanz-Moreno V. TGFbetainduced transcription in cancer. *Semin Cancer Biol.* 2017;42:60–69.

- Lin RL, Zhao LJ. Mechanistic basis and clinical relevance of the role of transforming growth factor-beta in cancer. *Cancer Biol Med.* 2015;12(4):385–393.
- Bellomo C, Caja L, Moustakas A. Transforming growth factor beta as regulator of cancer stemness and metastasis. *Br J Cancer*. 2016;115(7):761–769.
- 75. Felipe Lima J, Nofech-Mozes S, Bayani J, Bartlett JM. EMT in breast carcinoma-A review. *J Clin Med.* 2016;5(7):E65.
- 76. Li L, Qi L, Liang Z, et al. Transforming growth factor-β1 induces EMT by the transactivation of epidermal growth factor signaling through HA/CD44 in lung and breast cancer cells. *Int J Mol Med.* 2015;36(1):113–122.
- Chen W, Zhou S, Mao L, et al. Crosstalk between TGF-beta signaling and miRNAs in breast cancer metastasis. *Tumour Biol.* 2016;37(8):10011–10019.
- Li Z, Liu X, Wang B, et al. Pirfenidone suppresses MAPK signaling pathway to reverse epithelial-mesenchymal transition and renal fibrosis. *Nephrology (Carlton)*. Epub 2016 Jun 1. doi: 10.1111/nep.12831.
- Pang MF, Georgoudaki AM, Lambut L, et al. TGF-β1-induced EMT promotes targeted migration of breast cancer cells through the lymphatic system by the activation of CCR7/CCL21-mediated chemotaxis. *Oncogene*. 2016;35(6):748–760.
- Ye Y, Liu S, Wu C, Sun Z. TGFβ modulates inflammatory cytokines and growth factors to create premetastatic microenvironment and stimulate lung metastasis. *J Mol Histol*. 2015;46(4–5):365–375.
- Chen L, Fu H, Luo Y, et al. cPLA2α mediates TGF-β-induced epithelial-mesenchymal transition in breast cancer through PI3k/Akt signaling. *Cell Death Dis.* 2017;8(4):e2728.
- Ruff M, Leyme A, Le Cann F, et al. The disintegrin and metalloprotease ADAM12 is associated with TGF-beta-induced epithelial to mesenchymal transition. *PLoS One*. 2015;10(9):e0139179.
- Desai K, Aiyappa R, Prabhu JS, et al. HR+HER2- breast cancers with growth factor receptor-mediated EMT have a poor prognosis and lapatinib downregulates EMT in MCF-7 cells. *Tumour Biol.* 2017;39(3):1010428317695028. doi: 10.1177/1010428317695028.
- Nooshinfar E, Safaroghli-Azar A, Bashash D, Akbari ME. Melatonin, an inhibitory agent in breast cancer. *Breast Cancer*. 2017;24(1):42–51.
- Zhou YC, Liu JY, Li J, et al. Ionizing radiation promotes migration and invasion of cancer cells through transforming growth factor-betamediated epithelial-mesenchymal transition. *Int J Radiat Oncol Biol Phys.* 2011;81(5):1530–1537.
- Mediavilla-Varela M, Boateng K, Noyes D, Antonia SJ. The antifibrotic agent pirfenidone synergizes with cisplatin in killing tumor cells and cancer associated fibroblasts. *BMC Cancer*. 2016;16:176.
- Choi SH, Nam JK, Jang J, Lee HJ, Lee YJ. Pirfenidone enhances the efficacy of combined radiation and sunitinib therapy. *Biochem Biophys Res Commun.* 2015;462(2):138–143.
- Kozono S, Ohuchida K, Eguchi D, et al. Pirfenidone inhibits pancreatic cancer desmoplasia by regulating stellate cells. *Cancer Res.* 2013;73(7):2345–2356.
- Takai K, Le A, Weaver VM, Werb Z. Targeting the cancer-associated fibroblasts as a treatment in triple-negative breast cancer. *Oncotarget*. 2016;7(50):82889–82901.
- Polydorou C, Mpekris F, Papageorgis P, Voutouri C, Stylianopoulos T. Pirfenidone normalizes the tumor microenvironment to improve chemotherapy. *Oncotarget*. 2017;8(15):24506–24517.
- Wang X, Wei S, Zhao Y, et al. Anti-proliferation of breast cancer cells with itraconazole: Hedgehog pathway inhibition induces apoptosis and autophagic cell death. *Cancer Lett.* 2017;385:128–136.
- 92. Flores-Contreras L, Sandoval-Rodríguez AS, Mena-Enriquez MG, et al. Treatment with pirfenidone for two years decreases fibrosis, cytokine levels and enhances CB2 gene expression in patients with chronic hepatitis C. *BMC Gastroenterol*. 2014;14:131.
- Szabó R. Antiviral therapy and prevention against hantavirus infections. Acta Virol. 2017;61(1):3–12.
- Kast RE. Ribavirin in cancer immunotherapies: controlling nitric oxide augments cytotoxic lymphocyte function. *Neoplasia*. 2003; 5(1):3–8.

- Dixit NM, Layden-Almer JE, Layden TJ, Perelson AS. Modelling how ribavirin improves interferon response rates in hepatitis C virus infection. *Nature*. 2004;432(7019):922–924.
- Nakatsuka K, Atsukawa M, Shimizu M, Takahashi H, Kawamoto C. Ribavirin contributes to eradicate hepatitis C virus through polarization of T helper 1/2 cell balance into T helper 1 dominance. *World J Hepatol.* 2015;7(25):2590–2596.
- Siddiqui N, Sonenberg N. Signalling to eIF4E in cancer. *Biochem Soc Trans*. 2015;43(5):763–772.
- Kerekatte V, Smiley K, Hu B, Smith A, Gelder F, De Benedetti A. The proto-oncogene/translation factor eIF4E: a survey of its expression in breast carcinomas. *Int J Cancer*. 1995;64(1):27–31.
- Qin X, Jiang B, Zhang Y. 4E-BP1, a multifactor regulated multifunctional protein. *Cell Cycle*. 2016;15(6):781–786.
- Wang R, Ganesan S, Zheng XF. Yin and yang of 4E-BP1 in cancer. Cell Cycle. 2016;15(11):1401–1402.
- Muller D, Lasfargues C, El Khawand S, et al. 4E-BP restrains eIF4E phosphorylation. *Translation (Austin)*. 2013;1(2):e25819.
- Grzmil M, Huber RM, Hess D, et al. MNK1 pathway activity maintains protein synthesis in rapalog-treated gliomas. *J Clin Invest*. 2014;124(2):742–754.
- 103. Armengol G, Rojo F, Castellví J, et al. 4E-binding protein 1: a key molecular "funnel factor" in human cancer with clinical implications. *Cancer Res.* 2007;67(16):7551–7555.
- 104. Ramalingam S, Gediya L, Kwegyir-Afful AK, et al. First MNKs degrading agents block phosphorylation of eIF4E, induce apoptosis, inhibit cell growth, migration and invasion in triple negative and Her2-overexpressing breast cancer cell lines. *Oncotarget*. 2014;5(2):530–543.
- Pyronnet S. Phosphorylation of the cap-binding protein eIF4E by the MAPK-activated protein kinase Mnk1. *Biochem Pharmacol*. 2000;60(8):1237–1243.
- Lu C, Makala L, Wu D, Cai Y. Targeting translation: eIF4E as an emerging anticancer drug target. *Expert Rev Mol Med.* 2016;18:e2.
- 107. Fan W, Wang W, Mao X, et al. Elevated levels of p-Mnk1, p-eIF4E and p-p70S6K proteins are associated with tumor recurrence and poor prognosis in astrocytomas. *J Neurooncol.* 2017;131(3):485–493.
- Grzmil M, Seebacher J, Hess D, et al. Inhibition of MNK pathways enhances cancer cell response to chemotherapy with temozolomide and targeted radionuclide therapy. *Cell Signal*. 2016;28(9):1412–1421.
- 109. Li S, Fu J, Lu C, et al. Elevated translation initiation factor eIF4E is an attractive therapeutic target in multiple myeloma. *Mol Cancer Ther*. 2016;15(4):711–719.
- 110. Karaki S, Andrieu C, Ziouziou H, Rocchi P. The eukaryotic translation initiation factor 4E (eIF4E) as a therapeutic target for cancer. Adv Protein Chem Struct Biol. 2015;101:1–26.
- 111. Hu A, Sun M, Yan D, Chen K. Clinical significance of mTOR and eIF4E expression in invasive ductal carcinoma. *Tumori*. 2014;100(5):541–546.
- 112. DeFatta RJ, Turbat-Herrera EA, Li BD, Anderson W, De Benedetti A. Elevated expression of eIF4E in confined early breast cancer lesions: possible role of hypoxia. *Int J Cancer*. 1999;80(4):516–522.
- Sorrells DL, Black DR, Meschonat C, et al. Detection of eIF4E gene amplification in breast cancer by competitive PCR. *Ann Surg Oncol.* 1998;5(3):232–237.
- Norton KS, McClusky D, Sen S, et al. TLK1B is elevated with eIF4E overexpression in breast cancer. J Surg Res. 2004;116(1):98–103.
- 115. Heikkinen T, Korpela T, Fagerholm R, et al. Eukaryotic translation initiation factor 4E (eIF4E) expression is associated with breast cancer tumor phenotype and predicts survival after anthracycline chemotherapy treatment. *Breast Cancer Res Treat*. 2013;141(1):79–88.
- 116. Ramon Y, Cajal S, De Mattos-Arruda L, Sonenberg N, Cortes J, Peg V. The intra-tumor heterogeneity of cell signaling factors in breast cancer: p4E-BP1 and peIF4E are diffusely expressed and are real potential targets. *Clin Transl Oncol.* 2014;16(11):937–941.
- 117. Li BD, Gruner JS, Abreo F, et al. Prospective study of eukaryotic initiation factor 4E protein elevation and breast cancer outcome. *Ann Surg.* 2002;235(5):732–738.

- McClusky DR, Chu Q, Yu H, et al. A prospective trial on initiation factor 4E (eIF4E) overexpression and cancer recurrence in node-positive breast cancer. *Ann Surg.* 2005;242(4):584–590.
- 119. Yin X, Kim RH, Sun G, Miller JK, Li BD. Overexpression of eukaryotic initiation factor 4E is correlated with increased risk for systemic dissemination in node positive breast cancer patients. *JAm Coll Surg.* 2014;218(4):663–671.
- 120. Li Z, Sun Y, Qu M, Wan H, Cai F, Zhang P. Inhibiting the MNK-eIF4Ebeta-catenin axis increases responsiveness of aggressive breast cancer cells to chemotherapy. *Oncotarget*. 2017;8(2):2906–2915.
- 121. Volpin F, Casaos J, Sesen J, et al. Use of an anti-viral drug, ribavirin, as an anti-glioblastoma therapeutic. *Oncogene*. 2017;36(21):3037–3047.
- 122. Bhat M, Yanagiya A, Graber T, et al. Metformin requires 4E-BPs to induce apoptosis and repress translation of Mcl-1 in hepatocellular carcinoma cells. *Oncotarget*. Epub 2016 Jul 18. doi: 10.18632/ oncotarget.10671.
- 123. Pons B, Peg V, Vázquez-Sánchez MA, et al. The effect of p-4E-BP1 and p-eIF4E on cell proliferation in a breast cancer model. *Int J Oncol.* 2011;39(5):1337–1345.
- 124. Shi F, Len Y, Gong Y, et al. Ribavirin inhibits the activity of mTOR/ eIF4E, ERK/Mnk1/eIF4E signaling pathway and synergizes with tyrosine kinase inhibitor imatinib to impair Bcr-Abl mediated proliferation and apoptosis in Ph+ leukemia. *PLoS One.* 2015;10(8):e0136746.
- Zhang Y, Zheng XF. mTOR-independent 4E-BP1 phosphorylation is associated with cancer resistance to mTOR kinase inhibitors. *Cell Cycle*. 2012;11(3):594–603.
- Kentsis A, Volpon L, Topisirovic I, et al. Further evidence that ribavirin interacts with eIF4E. RNA. 2005;11(12):1762–1766.
- 127. Kentsis A, Topisirovic I, Culjkovic B, Shao L, Borden KL. Ribavirin suppresses eIF4E-mediated oncogenic transformation by physical mimicry of the 7-methyl guanosine mRNA cap. *Proc Natl Acad Sci* USA. 2004;101(52):18105–18110.
- 128. Pettersson F, Yau C, Dobocan MC, et al. Ribavirin treatment effects on breast cancers overexpressing eIF4E, a biomarker with prognostic specificity for luminal B-type breast cancer. *Clin Cancer Res.* 2011;17(9):2874–2884.
- 129. Pettersson F, Del Rincon SV, Emond A, et al. Genetic and pharmacologic inhibition of eIF4E reduces breast cancer cell migration, invasion, and metastasis. *Cancer Res.* 2015;75(6):1102–1112.
- Isakovic AM, Dulovic M, Markovic I, et al. Autophagy suppression sensitizes glioma cells to IMP dehydrogenase inhibition-induced apoptotic death. *Exp Cell Res.* 2017;350(1):32–40.
- 131. Ogino A, Sano E, Ochiai Y, et al. Efficacy of ribavirin against malignant glioma cell lines. *Oncol Lett.* 2014;8(6):2469–2474.
- 132. Criscitiello C, Gelao L, Viale G, Esposito A, Curigliano G. Investigational platelet-derived growth factor receptor kinase inhibitors in breast cancer therapy. *Expert Opin Investig Drugs*. 2014;23(5):599–610.
- 133. Yu JM, Sun W, Hua F, et al. BCL6 induces EMT by promoting the ZEB1-mediated transcription repression of E-cadherin in breast cancer cells. *Cancer Lett.* 2015;365(2):190–200.
- 134. Pinto MP, Dye WW, Jacobsen BM, Horwitz KB. Malignant stroma increases luminal breast cancer cell proliferation and angiogenesis through platelet-derived growth factor signaling. *BMC Cancer*. 2014;14:735.
- 135. Weigel MT, Banerjee S, Arnedos M, et al. Enhanced expression of the PDGFR/Abl signaling pathway in aromatase inhibitor-resistant breast cancer. *Ann Oncol.* 2013;24(1):126–133.
- 136. Attar-Schneider O, Drucker L, Gottfried M. Migration and epithelial to mesenchymal transition of lung cancer can be targeted via translation initiation factors eIF4E and eIF4GI. *Lab Invest*. 2016;96(9):1004–1015.
- Nasr Z, Robert F, Porco JA Jr, Muller WJ, Pelletier J. eIF4F suppression in breast cancer affects maintenance and progression. *Oncogene*. 2013;32(7):861–871.
- Robichaud N, del Rincon SV, Huor B, et al. Phosphorylation of eIF4E promotes EMT and metastasis via translational control of SNAIL and MMP-3. *Oncogene*. 2015;34(16):2032–2042.

- Cuny GD, Suebsuwong C, Ray SS. Inosine-5'-monophosphate dehydrogenase (IMPDH) inhibitors: a patent and scientific literature review. *Expert Opin Ther Pat.* 2017;27(6):677–690.
- 140. Guidicelli G, Chaigne-Delalande B, Dilhuydy MS, et al. The necrotic signal induced by mycophenolic acid overcomes apoptosis-resistance in tumor cells. *PLoS One*. 2009;4(5):e5493.
- 141. Decarlo L, Mestel C, Barcellos-Hoff MH, Schneider RJ. Eukaryotic translation initiation factor 4E is a feed-forward translational coactivator of transforming growth factor β early pro-transforming events in breast epithelial cells. *Mol Cell Biol.* 2015;35(15):2597–2609.
- 142. Dai D, Chen H, Tang J, Tang Y. Inhibition of mTOR/eIF4E by anti-viral drug ribavirin effectively enhances the effects of paclitaxel in oral tongue squamous cell carcinoma. *Biochem Biophys Res Commun.* 2017;482(4):1259–1264.
- 143. McHutchison JG, Gordon SC, Schiff ER, et al. Interferon alfa-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis C. Hepatitis Interventional Therapy Group. *N Engl J Med.* 1998;339(21):1485–1492.
- 144. Crabol Y, Catherinot E, Veziris N, Jullien V, Lortholary O. Rifabutin: where do we stand in 2016? *J Antimicrob Chemother*. 2016;71(7): 1759–1771.
- 145. Kunin CM. Antimicrobial activity of rifabutin. *Clin Infect Dis*. 1996;22(suppl 1):S3–S13.
- 146. Li J, Huang Y, Gao Y, Wu H, Dong W, Liu L. Antibiotic drug rifabutin is effective against lung cancer cells by targeting the eIF4E-β-catenin axis. *Biochem Biophys Res Commun.* 2016;472(2):299–305.
- 147. Nakayama S, Sng N, Carretero J, et al. β-catenin contributes to lung tumor development induced by EGFR mutations. *Cancer Res.* 2014;74(20):5891–5902.
- 148. Togashi Y, Hayashi H, Terashima M, et al. Inhibition of β-Catenin enhances the anticancer effect of irreversible EGFR-TKI in EGFRmutated non-small-cell lung cancer with a T790M mutation. *JThorac Oncol.* 2015;10(1):93–101.
- 149. Wei Z, Gao W, Wu Y, Ni B, Tian Y. Mutual interaction between BCL6 and miRNAs contributing to the pathogenesis of various cancers. *Clin Transl Oncol.* 2015;17(11):841–846.
- 150. Wu Q, Liu X, Yan H, et al. B-cell lymphoma 6 protein stimulates oncogenicity of human breast cancer cells. *BMC Cancer*. 2014;14:418.
- 151. Walker SR, Frank DA. Targeting BCL6 and STAT3 in triple negative breast cancer: the one-two punch? *Oncoscience*. 2015;2(11):912.
- 152. Chang CC, Ye BH, Chaganti RS, Dalla-Favera R. BCL-6, a POZ/ zinc-finger protein, is a sequence-specific transcriptional repressor. *Proc Natl Acad Sci U S A*. 1996;93(14):6947–6952.
- 153. Evans SE, Goult BT, Fairall L, et al. The ansamycin antibiotic, rifamycin SV, inhibits BCL6 transcriptional repression and forms a complex with the BCL6-BTB/POZ domain. *PLoS One*. 2014;9(3): e90889.
- 154. Zhu Z, Wang S, Zhu J, Yang Q, Dong H, Huang J. MicroRNA-544 down-regulates both Bcl6 and Stat3 to inhibit tumor growth of human triple negative breast cancer. *Biol Chem.* 2016;397(10):1087–1095.
- 155. Goodman CR, Sato T, Peck AR, et al. Steroid induction of therapyresistant cytokeratin-5-positive cells in estrogen receptor positive breast cancer through a BCL6-dependent mechanism. *Oncogene*. 2016;35(11):1373–1385.
- 156. An H, He L. Current understanding of metformin effect on the control of hyperglycemia in diabetes. *J Endocrinol*. 2016;228(3):R97–R106.
- 157. Zou YF, Xie CW, Yang SX, Xiong JP. AMPK activators suppress breast cancer cell growth by inhibiting DVL3-facilitated Wnt/beta-catenin signaling pathway activity. *Mol Med Rep.* 2017;15(2):899–907.
- 158. Xu H, Aldrich MC, Chen Q, et al. Validating drug repurposing signals using electronic health records: a case study of metformin associated with reduced cancer mortality. *J Am Med Inform Assoc.* 2015;22(1): 179–191.
- Sośnicki S, Kapral M, Węglarz L. Molecular targets of metformin antitumor action. *Pharmacol Rep.* 2016;68(5):918–925.

- Gongol B, Marin T, Peng IC, et al. AMPK exerts its anti-inflammatory effects through PARP-1 and Bcl-6. *Proc Natl Acad Sci USA*. 2013;110(8): 3161–3166.
- 161. Queiroz EA, Puukila S, Eichler R, et al. Metformin induces apoptosis and cell cycle arrest mediated by oxidative stress, AMPK and FOXO3a in MCF-7 breast cancer cells. *PLoS One.* 2014;9(5):e98207.
- 162. Silvestri A, Palumbo F, Rasi I, et al. Metformin induces apoptosis and downregulates pyruvate kinase M2 in breast cancer cells only when grown in nutrient-poor conditions. *PLoS One*. 2015;10(8):e0136250.
- 163. Wu L, Zhou B, Oshiro-Rapley N, et al. An ancient, unified mechanism for metformin growth inhibition in *C. elegans* and cancer. *Cell*. 2016;167(7):1705.e13–1718.e13.
- 164. Ben Sahra I, Regazzetti C, Robert G, et al. Metformin, independent of AMPK, induces mTOR inhibition and cell-cycle arrest through REDD1. *Cancer Res.* 2011;71(13):4366–4372.
- 165. Gui DY, Sullivan LB, Luengo A, et al. Environment dictates dependence on mitochondrial complex I for NAD+ and aspartate production and determines cancer cell sensitivity to metformin. *Cell Metab.* 2016;24(5):716–727.
- 166. Chae YK, Arya A, Malecek MK, et al. Repurposing metformin for cancer treatment: current clinical studies. *Oncotarget*. 2016;7(26):40767–40780.
- Coyle C, Cafferty FH, Vale C, Langley RE. Metformin as an adjuvant treatment for cancer: a systematic review and meta-analysis. *Ann Oncol.* 2016;27(12):2184–2195.
- Gadducci A, Biglia N, Tana R, Cosio S, Gallo M. Metformin use and gynecological cancers: a novel treatment option emerging from drug repositioning. *Crit Rev Oncol Hematol.* 2016;105:73–83.
- Zhang HH, Guo XL. Combinational strategies of metformin and chemotherapy in cancers. *Cancer Chemother Pharmacol*. 2016;78(1): 13–26.
- Cazzaniga M, Bonanni B. Breast cancer metabolism and mitochondrial activity: the possibility of chemoprevention with metformin. *Biomed Res Int.* 2015;2015:972193.
- 171. Qu C, Zhang W, Zheng G, Zhang Z, Yin J, He Z. Metformin reverses multidrug resistance and epithelial-mesenchymal transition (EMT) via activating AMP-activated protein kinase (AMPK) in human breast cancer cells. *Mol Cell Biochem*. 2014;386(1–2):63–71.
- 172. Soo JS, Ng CH, Tan SH, et al. Metformin synergizes 5-fluorouracil, epirubicin, and cyclophosphamide (FEC) combination therapy through impairing intracellular ATP production and DNA repair in breast cancer stem cells. *Apoptosis*. 2015;20(10):1373–1387.
- 173. Honjo S, Ajani JA, Scott AW, et al. Metformin sensitizes chemotherapy by targeting cancer stem cells and the mTOR pathway in esophageal cancer. *Int J Oncol.* 2014;45(2):567–574.
- 174. Harada K, Ferdous T, Harada T, Ueyama Y. Metformin in combination with 5-fluorouracil suppresses tumor growth by inhibiting the Warburg effect in human oral squamous cell carcinoma. *Int J Oncol.* 2016;49(1):276–284.
- 175. Tian Y, Tang B, Wang C, et al. Metformin mediates resensitivity to 5-fluorouracil in hepatocellular carcinoma via the suppression of YAP. *Oncotarget*. 2016;7(29):46230–46241.
- 176. Engels CC, de Glas NA, Sajet A, et al. The influence of insulin-like Growth Factor-1 Receptor expression and endocrine treatment on clinical outcome of postmenopausal hormone receptor positive breast cancer patients: a Dutch TEAM substudy analysis. *Mol Oncol.* 2016;10(4):509–516.
- 177. Hadad S, Iwamoto T, Jordan L, et al. Evidence for biological effects of metformin in operable breast cancer: a pre-operative, window-of-opportunity, randomized trial. *Breast Cancer Res Treat*. 2011;128(3):783–794.
- 178. Niraula S, Dowling RO, Ennis M, et al. Metformin in early breast cancer: a prospective window of opportunity neoadjuvant study. *Breast Cancer Res Treat*. 2012;135(3):821–830.
- 179. Kalinsky K, Crew KD, Refice S, et al. Presurgical trial of metformin in overweight and obese patients with newly diagnosed breast cancer. *Cancer Invest*. 2014;32(4):150–157.

- Bonanni B, Puntoni M, Cazzaniga M, et al. Dual effect of metformin on breast cancer proliferation in a randomized presurgical trial. *J Clin Oncol.* 2012;30(21):2593–2600.
- Cabello P, Pineda B, Tormo E, Lluch A, Eroles P. The antitumor effect of metformin is mediated by miR-26a in breast cancer. *Int J Mol Sci.* 2016;17(8):E1298.
- 182. Łabuzek K, Suchy D, Gabryel B, Bielecka A, Liber S, Okopień B. Quantification of metformin by the HPLC method in brain regions, cerebrospinal fluid and plasma of rats treated with lipopolysaccharide. *Pharmacol Rep.* 2010;62(5):956–965.
- Lalau JD, Lemaire-Hurtel AS, Lacroix C. Establishment of a database of metformin plasma concentrations and erythrocyte levels in normal and emergency situations. *Clin Drug Investig.* 2011;31(6):435–438.
- 184. Pantziarka P, Bouche G, Sukhatme V, Meheus L, Rooman I, Sukhatme VP. Repurposing drugs in oncology (ReDO)-propranolol as an anticancer agent. *Ecancermedicalscience*. 2016;10:680.
- 185. Montoya A, Amaya CN, Belmont A, et al. Use of non-selective betablockers is associated with decreased tumor proliferative indices in early stage breast cancer. *Oncotarget*. 2017;8(4):6446–6460.
- 186. Cardwell CR, Pottegård A, Vaes E, et al. Propranolol and survival from breast cancer: a pooled analysis of European breast cancer cohorts. *Breast Cancer Res.* 2016;18(1):119.
- 187. Rico M, Baglioni M, Bondarenko M, et al. Metformin and propranolol combination prevents cancer progression and metastasis in different breast cancer models. *Oncotarget*. 2017;8(2):2874–2889.
- 188. Botteri E, Munzone E, Rotmensz N, et al. Therapeutic effect of β-blockers in triple-negative breast cancer postmenopausal women. *Breast Cancer Res Treat*. 2013;140(3):567–575.
- Hwa YL, Shi Q, Kumar SK, et al. Beta-blockers improve survival outcomes in patients with multiple myeloma: a retrospective evaluation. *Am J Hematol.* 2017;92(1):50–55.
- 190. Choy C, Raytis JL, Smith DD, et al. Inhibition of β 2-adrenergic receptor reduces triple-negative breast cancer brain metastases: the potential benefit of perioperative β -blockade. *Oncol Rep.* 2016;35(6): 3135–3142.
- 191. Childers WK, Hollenbeak CS, Cheriyath P. β-blockers reduce breast cancer recurrence and breast cancer death: a meta-analysis. *Clin Breast Cancer*. 2015;15(6):426–431.
- 192. Melhem-Bertrandt A, Chavez-Macgregor M, Lei X, et al. Beta-blocker use is associated with improved relapse-free survival in patients with triple-negative breast cancer. J Clin Oncol. 2011;29(19):2645–2652.
- 193. Elefteriou F. Role of sympathetic nerves in the establishment of metastatic breast cancer cells in bone. J Bone Oncol. 2016;5(3): 132–134.
- 194. Jang HI, Lim SH, Lee YY, et al. Perioperative administration of propranolol to women undergoing ovarian cancer surgery: a pilot study. *Obstet Gynecol Sci.* 2017;60(2):170–177.
- 195. Wang T, Li Y, Lu HL, Meng QW, Cai L, Chen XS. β-Adrenergic receptors: new target in breast cancer. Asian Pac J Cancer Prev. 2015;16(18):8031–8039.
- 196. Wilson JM, Lorimer E, Tyburski MD, Williams CL. β-Adrenergic receptor suppress Rap1B prenylation and promote the metastatic phenotype in breast cancer cells. *Cancer Biol Ther.* 2015;16(9): 1364–1374.
- 197. Işeri OD, Sahin FI, Terzi YK, Yurtcu E, Erdem SR, Sarialioglu F. beta-Adrenoreceptor antagonists reduce cancer cell proliferation, invasion, and migration. *Pharm Biol.* 2014;52(11):1374–1381.
- 198. Kang F, Ma W, Ma X, et al. Propranolol inhibits glucose metabolism and 18F-FDG uptake of breast cancer through posttranscriptional downregulation of hexokinase-2. J Nucl Med. 2014;55(3):439–445.
- 199. Campbell JP, Karolak MR, Ma Y, et al. Stimulation of host bone marrow stromal cells by sympathetic nerves promotes breast cancer bone metastasis in mice. *PLoS Biol.* 2012;10(7):e1001363.
- 200. Sloan EK, Priceman SJ, Cox BF, et al. The sympathetic nervous system induces a metastatic switch in primary breast cancer. *Cancer Res.* 2010;70(18):7042–7052.

- 201. Shaashua L, Shabat-Simon M, Haldar R, et al. Perioperative COX-2 and & beta-adrenergic blockade improves metastatic biomarkers in breast cancer patients in a phase-II randomized trial. *Clin Cancer Res.* Epub 2017 May 10. doi: 10.1158/1078-0432.CCR-17-0152.
- Wagstaff AJ, Ibbotson T, Goa KL. Capecitabine: a review of its pharmacology and therapeutic efficacy in the management of advanced breast cancer. *Drugs*. 2003;63(2):217–236.
- 203. Walko CM, Lindley C. Capecitabine: a review. *Clin Ther*. 2005;27(1):23-44.
- Daniele G, Gallo M, Piccirillo MC, et al. Pharmacokinetic evaluation of capecitabine in breast cancer. *Expert Opin Drug Metab Toxicol*. 2013;9(2):225–235.
- 205. Rossi D, Alessandroni P, Catalano V, et al. Safety profile and activity of lower capecitabine dose in patients with metastatic breast cancer. *Clin Breast Cancer*. 2007;7(11):857–860.
- 206. Zielinski C, Gralow J, Martin M. Optimising the dose of capecitabine in metastatic breast cancer: confused, clarified or confirmed? *Ann Oncol.* 2010;21(11):2145–2152.
- 207. Ambros T, Zeichner SB, Zaravinos J, et al. A retrospective study evaluating a fixed low dose capecitabine monotherapy in women with HER-2 negative metastatic breast cancer. *Breast Cancer Res Treat*. 2014;146(1):7–14.
- Nishijima TF, Suzuki M, Muss HB. A comparison of toxicity profiles between the lower and standard dose capecitabine in breast cancer: a systematic review and meta-analysis. *Breast Cancer Res Treat*. 2016;156(2):227–236.
- 209. Azuma Y, Hata K, Sai K, et al. Significant association between handfoot syndrome and efficacy of capecitabine in patients with metastatic breast cancer. *Biol Pharm Bull*. 2012;35(5):717–724.
- 210. Kast RE. Erlotinib augmentation with dapsone for rash mitigation and increased anti-cancer effectiveness. *Springerplus*. 2015;4:638.
- 211. Harbeck N, Saupe S, Jäger E, et al; PELICAN Investigators. A randomized phase III study evaluating pegylated liposomal doxorubicin versus capecitabine as first-line therapy for metastatic breast cancer: results of the PELICAN study. *Breast Cancer Res Treat*. 2017;161(1):63–72.
- 212. Emet M, Ozcan H, Ozel L, Yayla M, Halici Z, Hacimuftuoglu A. A review of melatonin, its receptors and drugs. *Eurasian J Med.* 2016;48(2):135–141.
- Kast RE. Agomelatine or ramelteon as treatment adjuncts in glioblastoma and other M1- or M2-expressing cancers. *Contemp Oncol (Pozn)*. 2015;19(2):157–162.
- Liu J, Clough SJ, Hutchinson AJ, Adamah-Biassi EB, Popovska-Gorevski M, Dubocovich ML. MT1 and MT2 melatonin receptors: a therapeutic perspective. *Annu Rev Pharmacol Toxicol.* 2016;56:361–383.
- 215. Hardeland R, Poeggeler B. Melatonin and synthetic melatonergic agonists: actions and metabolism in the central nervous system. *Cent Nerv Syst Agents Med Chem.* 2012;12(3):189–216.
- 216. Perlemuter G, Cacoub P, Valla D, et al. Characterisation of agomelatineinduced increase in liver enzymes: frequency and risk factors determined from a pooled analysis of 7605 treated patients. *CNS Drugs*. 2016;30(9):877–888.
- Vriend J, Reiter RJ. Breast cancer cells: modulation by melatonin and the ubiquitin-proteasome system – a review. *Mol Cell Endocrinol*. 2015;417:1–9.
- 218. Hill SM, Belancio VP, Dauchy RT, et al. Melatonin: an inhibitor of breast cancer. *Endocr Relat Cancer*. 2015;22(3):R183–R204.
- Cos S, Alvarez-García V, González A, Alonso-González C, Martínez-Campa C. Melatonin modulation of crosstalk among malignant epithelial, endothelial and adipose cells in breast cancer. *Oncol Lett.* 2014;8:487–492.
- Sánchez-Barceló EJ, Cos S, Mediavilla D, Martínez-Campa C, González A, Alonso-González C. Melatonin-estrogen interactions in breast cancer. *J Pineal Res*. 2005;38(4):217–222.
- 221. Mediavilla MD, Sánchez-Barceló EJ, Tan DX, Manchester L, Reiter RJ. Basic mechanisms involved in the anti-cancer effects of melatonin. *Curr Med Chem.* 2010;17:4462–4481.

- 222. Lopes J, Arnosti D, Trosko JE, Tai MH, Zuccari D. Melatonin decreases estrogen receptor binding to estrogen response elements sites on the OCT4 gene in human breast cancer stem cells. *Genes Cancer*. 2016;7(5–6):209–217.
- 223. Nooshinfar E, Bashash D, Safaroghli-Azar A, et al. Melatonin promotes ATO-induced apoptosis in MCF-7 cells: proposing novel therapeutic potential for breast cancer. *Biomed Pharmacother*. 2016;83:456–465.
- 224. Lopes JR, da Silva Kavagutti M, Medeiros FA, De Campos Zuccari DA. Evaluation of melatonin effect on human breast cancer stem cells using a three-dimensional growth method of mammospheres. *Anticancer Agents Med Chem.* Epub 2016 Sep 22.
- 225. Cos S, Fernández R, Güezmes A, Sánchez-Barceló EJ. Influence of melatonin on invasive and metastatic properties of MCF-7 human breast cancer cells. *Cancer Res.* 1998;58(19):4383–4390.
- 226. Mao L, Summers W, Xiang S, et al. Melatonin represses metastasis in Her2-positive human breast cancer cells by suppressing RSK2 expression. *Mol Cancer Res.* 2016;14(11):1159–1169.
- 227. Mao L, Yuan L, Slakey LM, Jones FE, Burow ME, Hill SM. Inhibition of breast cancer cell invasion by melatonin is mediated through regulation of the p38 mitogen-activated protein kinase signaling pathway. *Breast Cancer Res.* 2010;12(6):R107.
- Jardim-Perassi BV, Arbab AS, Ferreira LC, et al. Effect of melatonin on tumor growth and angiogenesis in xenograft model of breast cancer. *PLoS One*. 2014;9(1):e85311.
- 229. Su SC, Hsieh MJ, Yang WE, Chung WH, Reiter RJ, Yang SF. Cancer metastasis: mechanisms of inhibition by melatonin. *J Pineal Res.* 2017;62(1). doi: 10.1111/jpi.12370.
- 230. Gonçalves Ndo N, Colombo J, Lopes JR, et al. Effect of melatonin in epithelial mesenchymal transition markers and invasive properties of breast cancer stem cells of canine and human cell lines. *PLoS One*. 2016;11(3):e0150407.
- Rudra DS, Pal U, Maiti NC, Reiter RJ, Swarnakar S. Melatonin inhibits matrix metalloproteinase-9 activity by binding to its active site. *J Pineal Res.* 2013;54(4):398–405.
- 232. Borin TF, Arbab AS, Gelaleti GB, et al. Melatonin decreases breast cancer metastasis by modulating Rho-associated kinase protein-1 expression. *J Pineal Res.* 2016;60(1):3–15.
- Alvarez-Garcia V, Gonzalez A, Martinez-Campa C, Alonso-Gonzalez C, Cos S. Melatonin modulates aromatase activity and expression in endothelial cells. *Oncol Rep.* 2013;29(5):2058–2064.
- Alvarez-Garcia V, Gonzalez A, Alonso-Gonzalez C, Martinez-Campa C, Cos S. Antiangiogenic effects of melatonin in endothelial cell cultures. *Microvasc Res.* 2013;87:25–33.
- 235. Cos S, Alvarez-Garcia V, Gonzalez A, Alonso-Gonzalez C, Martinez-Campa C. Melatonin modulation of crosstalk among malignant epithelial, endothelial and adipose cells in breast cancer (Review). *Oncol Lett.* 2014;8(2):487–492.
- 236. Wang J, Xiao X, Zhang Y, et al. Simultaneous modulation of COX-2, p300, Akt, and Apaf-1 signaling by melatonin to inhibit proliferation and induce apoptosis in breast cancer cells. *J Pineal Res.* 2012;53(1):77–90.
- 237. Martínez-Campa C, Menéndez-Menéndez J, Alonso-González C, González A, Alvarez-García V, Cos S. What is known about melatonin, chemotherapy and altered gene expression in breast cancer. *Oncol Lett.* 2017;13(4):2003–2014.
- Ma C, Li LX, Zhang Y, et al. Protective and sensitive effects of melatonin combined with adriamycin on ER+ (estrogen receptor) breast cancer. *Eur J Gynaecol Oncol.* 2015;36(2):197–202.
- 239. Gao Y, Xiao X, Zhang C, et al. Melatonin synergizes the chemotherapeutic effect of 5-fluorouracil in colon cancer by suppressing PI3K/ AKT and NF-κB/iNOS signaling pathways. *J Pineal Res.* 2017;62(2). doi: 10.1111/jpi.12380.
- 240. Pariente R, Pariente JA, Rodríguez AB, Espino J. Melatonin sensitizes human cervical cancer HeLa cells to cisplatin-induced cytotoxicity and apoptosis: effects on oxidative stress and DNA fragmentation. *J Pineal Res.* 2016;60(1):55–64.

- Uguz AC, Cig B, Espino J, et al. Melatonin potentiates chemotherapyinduced cytotoxicity and apoptosis in rat pancreatic tumor cells. *J Pineal Res.* 2012;53(1):91–98.
- Ruiz-Rabelo J, Vázquez R, Arjona A, et al. Improvement of capecitabine antitumoral activity by melatonin in pancreatic cancer. *Pancreas*. 2011;40(3):410–414.
- Zhelev Z, Ivanova D, Bakalova R, Aoki I, Higashi T. Synergistic cytotoxicity of melatonin and new-generation anticancer drugs against leukemia lymphocytes but not normal lymphocytes. *Anticancer Res.* 2017;37(1):149–159.
- Alonso-González C, González A, Martínez-Campa C, et al. Melatonin enhancement of the radiosensitivity of human breast cancer cells is associated with the modulation of proteins involved in estrogen biosynthesis. *Cancer Lett.* 2016;370(1):145–152.
- 245. Lu JJ, Fu L, Tang Z, et al. Melatonin inhibits AP-2beta/hTERT, NF-kB/COX-2 and Akt/ERK and activates caspase/Cyto C signaling to enhance the antitumor activity of berberine in lung cancer cells. *Oncotarget*. 2016;7(3):2985–3001.
- 246. Calvo JR, González-Yanes C, Maldonado MD. The role of melatonin in the cells of the innate immunity: a review. *J Pineal Res*. 2013;55(2): 103–120.
- Devore EE, Warner ET, Eliassen AH, et al. Urinary melatonin in relation to postmenopausal breast cancer risk according to melatonin 1 receptor status. *Cancer Epidemiol Biomarkers Prev.* 2017;26(3):413–419.
- 248. Basler M, Jetter A, Fink D, Seifert B, Kullak-Ublick GA, Trojan A. Urinary excretion of melatonin and association with breast cancer: meta-analysis and review of the literature. *Breast Care (Basel)*. 2014;9(3):182–187.

Publish your work in this journal

Breast Cancer - Targets and Therapy is an international, peerreviewed open access journal focusing on breast cancer research, identification of therapeutic targets and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/breast-cancer---targets-and-therapy-journal

Dovepress



Yusuke Shiozawa, MD, PhD Curriculum Vitae

WAKE FOREST SCHOOL OF MEDICINE Curriculum Vitae Elements

NAME:	Yusuke Shiozawa, M.D., Ph.D.
CURRENT ACADEMIC TITLE:	Assistant Professor
ADDRESS:	Department of Cancer Biology Wake Forest School of Medicine Medical Center Boulevard Winston-Salem, NC 27157-1010 (336) 716-8743 (office) (336) 716-0255 (fax) E-mail: yshiozaw@wakehealth.edu
EDUCATION:	
1994-2000	Juntendo University School of Medicine Tokyo, Japan M.D.
2002-2006	Juntendo University School of Medicine Department of Pediatrics Tokyo, Japan Ph.D. Research Advisor: Yuichiro Yamashiro, M.D., Ph.D. Nobutaka Kiyokawa, M.D., Ph.D. Thesis: <i>Human osteoblasts support hematopoietic cell</i> <i>development in vitro</i>
2016-2017	Wake Forest School of Medicine Clinical and Translational Science Institute Translational Research Academy 2016 Cohort
POSTDOCTORAL TRAINING:	
2000-2002	Resident, (Department of Pediatrics) Juntendo University School of Medicine Tokyo, Japan
2002-2006	Staff Doctor, (Pediatric Hematology/Oncology) Juntendo University School of Medicine Tokyo, Japan
2004-2006	Predoctoral Research Fellow, National Research Institute for Child Health and Development, Department of Pediatric Hematology and Oncology Research Research Advisor: Nobutaka Kiyokawa, M.D., Ph.D. Research Project: <i>Human osteoblasts support hematopoietic</i> <i>cell development in vitro</i>
2006-2010	Postdoctoral Research Fellow, University of Michigan School of Dentistry,

Department of Periodontics & Oral Medicine. Research Advisor: Russell S. Taichman, D.M.D, D.M.Sc. Research Project: *Human prostate cancer metastases target the hematopoietic stem cell niche to establish footholds in mouse bone marrow*

PROFESSIONAL LICENSURE:

2000-present	Japanese Medical License
EMPLOYMENT:	
Academic Appointments:	

Wake Forest School of Medicine:

2015-Present	Assistant Professor,
	Department of Cancer Biology

University of Michigan School of Dentistry:

2010-2014	Research Investigator, Department of Periodontics & Oral Medicine
2014-2015	Assistant Research Scientist, Department of Periodontics & Oral Medicine

OTHER PROFESSIONAL APPOINTMENTS AND INSTITUTIONAL SERVICE:

Institutional Committee Service:

Wake Forest School of Medicine:

2016, 2017	The Cancer Center Protocol Review Committee
2016-2019	The Radiation Safety Committee
2017	Application reviewer for Medical Student Research Program

University of Michigan School of Dentistry:

2014, 2015	Judge for Dental School Research day
	0

Departmental Committee Service:

Wake Forest School of Medicine:

2015, 2016, 2017	Molecular and Cellular Bioscience PhD Track 4 Recruiting
	committee (Interview, Application Review)
2016, 2017	Molecular and Cellular Bioscience MD/PhD Track 4 Recruiting
	committee (Interview, Application Review)

University of Michigan School of Dentistry:

2011, 2012 Research track faculty promotion committee

EXTRAMURAL APPOINTMENTS AND SERVICE:

Journal Reviewer:

2008-Present	Journal of Gene Medicine
2008-Present	Biochemical Pharmacology (reviewed 16 times)
2010-Present	Journal of Bone and Mineral Research (reviewed 4 times)
2010-2016	Journal of Pediatric Biochemistry (reviewed 1 time)
2010-Present	Journal of Pediatric Epilepsy
2010-Present	Journal of Pediatric Neuroradiology
2010-Present	Journal of Pediatric Intensive Care
2011-Present	Journal of Dental Research (reviewed 1 time)
2012-Present	Asian Journal of Andrology (reviewed 1 time)
2012-Present	Clinical Cancer Research (reviewed 1 time)
2013-Present	Pediatric Oncall
2014-Present	The journal Minerva Pediatrica (reviewed 4 times)
2014-Present	Oncotarget (reviewed 5 times)
2015-Present	PLOS ONE
2015-Present	Cancer Research Frontiers (reviewed 1 time)
2015-Present	International Journal of Urology
2017-Present	Journal of Child Science

Ad-hoc Journal Reviewer:

2009	Blood
2009	American Journal of Pathology
2009	Journal of Clinical Investigation
2009	Acta Pediatrica
2011	International Medical Case Reports Journal
2012	Clinical & Experimental Metastasis
2012	Current Psychiatry Reviews
2012	Molecular Cancer Therapeutics
2013	African Journal of Biotechnology
2013	Expert Opinion On Biological Therapy
2013	Hematology/Oncology and Stem Cell Therapy
2013	International Journal of Nutrition and Metabolism
2013	Letters in Drug Design & Discovery
2013	Journal of Experimental & Clinical Cancer Research
2013	Tissue Engineering
2013	Journal of Immunoassay and Immunochemistry
2013	Topics in Anticancer Research (BSP-Patent Book Series)
2014	Expert Opinion On Biological Therapy
2014	International Journal of Endocrinology
2014	BioMed Research International
2014	Oncotarget
2014	Current Pharmaceutical Design

2014	Stem Cells International
2014	International Journal of Nanomedicine
2014	Cancer informatics
2015	Current Cancer Drug Targets (reviewed 2 times this year)
2015	Frontiers in Drug Design and Discovery (eBook Series)
2015	Evidence-Based Complementary and Alternative Medicine
2015	Medical Journals-Journal of Bone Marrow Research
2015	Expert Opinion On Drug Discovery
2015	BBA - Molecular Cell Research
2015	Medical Journals-Angiology (reviewed 2 times this year)
2015	Medical Journals-Integrative Oncology
2015	Cancers (reviewed 2 times this year)
2015	Calcified Tissue International and Musculoskeletal Research
2015	Cancer Biology & Medicine
2015	Medical Journals-Palliative Care & Medicine
2015	Jacobs Journal of Dentistry and Research
2015	Neoplasia
2015	Medical Journals-Journal of Pain Management & Medicine
2015	Cancer Growth and Metastasis
2015	Medical Journals-Journal of Pain Management & Medicine
2015	Gastroenterology Research and Practice
2015	Stem Cells and Cloning: Advances and Applications
2016	Medical Journals-Integrative Oncology
2016	Anesthesiology
2016	Medical Journals-Nuclear Medicine & Radiation Therapy
2016	The Journal of Cellular Biochemistry
2016	Medical Journals-Nuclear Medicine & Radiation Therapy
2016	Medical Journals-Journal of Prostate Cancer
	(reviewed 2 times this year)
2016	Stem Cell International (reviewed 2 times this year)
2016	International Journal of Molecular Sciences
	(reviewed 2 times this year)
2016	Cancer Growth and Metastasis
2016	Journal of Urology and Renal Diseases
2016	Journal of Spine & Neurosurgery
2017	International Journal of Molecular Sciences
	(reviewed 3 times this year)
2017	Nature Communications
2017	Journal of Clinical & Experimental oncology
	(reviewed 3 times this year)
2017	Marine Drugs
2017	Molecular Medicine Reports
2017	Urologic Oncology: Seminars and Original Investigations
2017	BBA - Molecular Cell Research
··	

Funding Agency Reviewer:

The NY Stem Cell Research Program, Washington DC (Ad hoc Reviewer, October 15-17, 2008)

Cancer Research UK Program (Ad hoc Reviewer (remote, online), February, 2009)

The Fund for Scientific Research (F.R.S.-FNRS)

Yusuke Shiozawa, M.D., Ph.D. Department of Cancer Biology

(Ad hoc Reviewer (remote, online), September, 2013)

- The Fund for Scientific Research (F.R.S.-FNRS) (Ad hoc Reviewer (remote, online), September, 2014)
- The Fund for Scientific Research (F.R.S.-FNRS) (Ad hoc Reviewer (remote, online), September, 2015)
- Prostate Cancer Foundation, 2017 Young Investigator Award (Ad hoc Reviewer (remote, online), April, 2017)

Editorial Boards:

2011-Present	Journal of Clinical Medicine and Research
2011-Present	Case Reports in Hematology (edited 61 times)
2012-Present	International Journal of Experimental Dental Science
2012-Present	Case Reports in Pediatrics (edited 13 times)
2013-Present	JSM Dentistry
2013-Present	Genetic Disorders & Gene Therapy
2013-Present	Cancer Studies
2014-Present	Pharmacologia
2014-Present	Austin Journal of Dentistry
2014-Present	Journal of Stem Cell Research and Transplantation
	(reviewed 4 times)
2014-Present	Journal of Hematology, Blood Transfusion and Disorders
2014-Present	Journal of Stem Cells Research, Development and Therapy
2014-Present	Journal of Clinical Studies and Medical Case Reports
2014-Present	Journal of Cancer Science and Clinical Research
2014-Present	Journal of Dentistry and Clinical Research
2014-Present	Journal of Stem Cells Research, Reviews & Reports
2014-Present	Sarcoma Research - International (reviewed 5 times)
2014-Present	International Journal of Hematology Research
2015-Present	Medical Oncology
2015-2016	Journal of Molecular Cytotherapy
2016-Present	Journal of Cancer Biology and Treatment (edited 2 times)
2016-Present	JSM Bone Marrow Research

Ad-hoc Journal Editor for Special Issue:

Session Chair:

2014

19th Annual Prostate Cancer Foundation Scientific Retreat Session 1: Field Cancerization & the Tumor Microenvironment October 25-27, 2012 San Diego, CA

PROFESSIONAL MEMBERSHIPS:

2008-Present	American Association for Cancer Research
2008-2016	American Society of Hematology
2011-Presnet	International Bone and Mineral Society

Yusuke Shiozawa, M.D., Ph.D. Department of Cancer Biology

2015-Presnet	Metastasis Research Society
2017-Present	Cancer and Bone Society
2017-Present	Society for Basic Urologic Research

HONORS AND AWARDS:

2006	Pediatric Oncology Research Fellowship
2007	Best Paper Award from the Japan Cytometry Society
2009	Department of Defense Prostate Cancer Physician Research Training Award
2009	4th place in the 2009 University of Michigan Cancer Center Fall Research Symposium Poster Session (out of 78 posters)
2011	Prostate Cancer Foundation Young Investigator Award
2017	Accepted Participant of The U.S. Bone and Joint Initiative (USBJI) and Bone and Joint Canada (BJC) Young Investigator Initiative Program

INVITED PRESENTATIONS AND SEMINARS:

- 1. Invited Presentation. <u>Annexin II regulates prostate cancer metastasis.</u> Joint prostate cancer P01 meeting. Atlanta, GA. January 28, 2007.
- 2. Invited Presentation. <u>How metastatic prostate cancer parasitize the osteoblastic hematopoietic</u> <u>stem cell niche.</u> Joint prostate cancer P01 meeting. Seattle, WA. January 27, 2010.
- 3. Invited Seminar. <u>The hematopoietic stem cell niche is the target of metastatic cells from solid</u> <u>tumors.</u> Case Western Reserve University, Cleveland, OH. February 23, 2012.
- Invited Presentation. <u>Prevalence of prostate cancer metastasis following intravenous inoculation</u> provides clues into the molecular basis of dormancy in the bone marrow microenvironment. 5th Annual Prostate Cancer Program Retreat. Fort Lauderdale, FL. March 20, 2012.
- 5. Invited Presentation. <u>The bone marrow microenvironment controls the phenotype of disseminated</u> <u>tumor cells.</u> Joint prostate cancer P01 meeting. Ann Arbor, MI. April 12, 2012.
- 6. Invited Seminar. <u>Solid tumors target hematopoietic stem cell niche is during metastasis.</u> University of South Florida, Tampa, FL. July 9, 2012.
- 7. Invited Seminar. <u>Solid tumors target hematopoietic stem cell niche is for metastasis.</u> University of Oklahoma Health Science Center, Oklahoma City, OK. July 31, 2012.
- 8. Invited Seminar. <u>Prostate cancer target hematopoietic stem cell niche during metastasis.</u> The Cedars-Sinai medical center, Los Angeles, CA. Dec 3, 2012.

- 9. Invited Presentation. <u>The bone marrow niche controls the "stemness" of disseminated tumor cells.</u> 6th Annual Prostate Cancer Program Retreat. Fort Lauderdale, FL. March 20, 2013.
- 10. Invited Seminar. <u>The critical roles of bone marrow microenvironment in prostate cancer bone</u> <u>metastasis.</u> Oregon Health & Science University, Portland, OR. April 22, 2013.
- 11. Invited Presentation. <u>The bone marrow niche controls the "stemness" of disseminated tumor cells.</u> 13th International Conference on Cancer-Induced Bone Disease. Miami, FL. November 8, 2013.
- 12. Invited Seminar. *Bone as a Target: Solving the Mystery of Cancer Metastasis.* Penn State Hershey Children's Hospital, Hershey, PA. July 9, 2014.
- 13. Invited Seminar. <u>Bone as a Target: Solving the Mystery of Prostate Cancer Metastasis.</u> Wake Forest School of Medicine, Winston-Salem, NC. July 15, 2014.
- 14. Invited Seminar. *Bone as a Target: Solving the Mystery of Prostate Cancer Metastasis.* Cleveland Clinic, Cleveland, OH. July 28, 2014.
- 15. Invited Seminar. <u>My Ongoing Journey To Uncover The Mystery Of Prostate Cancer Bone</u> <u>Metastasis.</u> Jikei University School of Medicine, Tokyo, Japan. January 5, 2015.

GRANTS:

Current:

W81XWH-14-1-0403, Idea Development Award (Shiozawa, PI, 25% effort)

Department of Defense

03/01/2015-02/28/2018 \$75,000/year direct cost

The roles of the bone marrow microenvironment in controlling tumor dormancy

The goal of this project is to provide significant insight into the mechanisms whereby the bone marrow microenvironment is involved in regulation of tumor dormancy. Aim 1 allows us to identify and explore how disseminated tumor cells (DTCs) stay dormant for long periods of time within the bone marrow microenvironment through the GAS6 pathway. Aim 2 will determine whether exogenous BMP2 triggers the activation of dormant DTCs through the bone marrow microenvironment. This work will inform greater understanding of microenvironmental aging's effects on metastatic growth, and could result in valuable new treatment approaches.

Cancer Center, Tumor Progression and Recurrence Program Pilot Funding (Shiozawa, PI)

Wake Forest Baptist Comprehensive Cancer Center

The roles of renin-angiotensin system in cancer-induced bone pain

In this pilot project we will generate preliminary data for future external grant applications, which will determine the mechanisms of cancer-induced bone pain development.

Multi-Investigator Prostate Cancer Research Internal Pilot Funding (Shiozawa, PI) 06/01/2017-05/31/2018

Wake Forest Baptist Comprehensive Cancer Center

\$20,000/year direct cost Development of theranostic strategies for prostate cancer bone metastases In this pilot project we will develop a theranostic strategy that targets pain signaling in bone metastatic disease.

W81XWH-17-1-0541, Idea Development Award (Shiozawa, PI, 20% effort, Peters, Partnering PI)

09/01/2017-08/31/2020 \$333,333/year direct cost

Department of Defense

Molecular crosstalk: bone metastatic prostate cancer and nociceptive neurons

The proposed studies will focus on fundamental mechanisms behind the novel concept that nerves in the bone marrow microenvironment have a crucial role in prostate cancer bone metastasis. Aim 1 will provide the framework to identify the extent to which the interaction between disseminated tumor cells (DTCs) and nociceptive neurons affects cancer-induced bone pain. Aim 2 will determine the molecular mechanisms of cancer-induced bone pain. Aim 3 will define how DTC/nociceptive neuron interactions influence tumor outgrowth. We believe that the insights derived from our investigations will lead to new strategies for reducing cancer-induced bone pain and also the outgrowth of DTCs.

Cancer Center, Tumor Progression and Recurrence Program Push F	Pilot Funding (Shiozawa, PI)
	09/01/2017-08/31/2018
Wake Forest Baptist Comprehensive Cancer Center	\$50,000/year direct cost
The roles of the stem cell factor/its receptor c-Kit axis in cancer-induced bone pain	

In this pilot project we will generate preliminary data for future external grant applications, which will determine the mechanisms whereby the stem cell factor/its receptor c-Kit axis controls cancer-induced bone pain development.

06/01/2017-05/31/2018 \$10,000/year direct cost Wake Forest School of Medicine CTSI Ignition Pilot Funding (Shiozawa, PI)

Wake Forest School of Medicine CTSI

09/20/2017-12/31/2017 \$5,000/year direct cost

Feasibility study for *in vivo* gene manipulation with adeno-associated viral vectors

In this pilot project, we will optimize and demonstrate the feasibility of the generation of nerve-specific c-Kit knockdown mice using adeno-associated virus (AAV) vectors in vivo. Our other studies will continue to investigate the influence of SCF on neurite outgrowth and neuropeptide expression in sensory nerves.

Pending:

None

PAST GRANT HISTORY:

Pediatric Oncology Research Fellowship (Shiozawa, PI) 2006-2007 (¥4,000,000/year direct cost)

Department of Defense Role and Function of the Chemokine CXCL16 and its Receptor CXCR6 in Prostate Cancer (Shiozawa, PI) 01/01/2009-07/31/2011 (\$60,000/year direct cost)

Prostate Cancer Foundation

Molecular dissection of the premetastatic niche in prostate cancer (Shiozawa, PI) 01/01/2012-12/31/2014 (\$75,000/year direct cost)

National Cancer Institution

1U54CA163124, Tumor Microenvironment Network U54 Mechanisms of Prostate Cancer Dormancy In The Bone Marrow Niche (Taichman, PI: Shiozawa, Co-PI) 09/16/2011-07/31/2016 (\$87,233/year direct cost to YS)

Wake Forest School of Medicine CTSI Wake Forest School of Medicine Internal Pilot Funding Molecular Crosstalk of Bone Metastatic Prostate Cancer and Nociceptive Neurons (Shiozawa, PI) 07/01/2016-06/30/2017 (\$10,000/year direct cost)

PATENTS:

 Publication number: WO2012139131 A1, Application number: PCT/US2012/032807, "Autologous human adult pluripotent very small embryonic-like (HVSEL) stem cell regeneration of bone and cartilage", Taichman RS, Krebsbach PH, Havens AM, Mishra A, Rodgerson DO, Wang J, Shiozawa Y, Jung Y., Publication date: October 11, 2012.

The invention provides methods and compositions for the repair or regeneration of osteochondral tissue. The methods and compositions provide an effective amount of isolated differentiable human Very Small Embryonic Like Stem Cells (hVSELs) sufficient for regeneration or repair of an osteochondral tissue. The compositions can be administered directly to the tissue or administered systemically.

BIBLIOGRAPHY:

Peer-Reviewed Journal Articles:

- Shiozawa Y, Fujita H, Fujimura J, Suzuki K, Sato H, Saito M, Shimizu T, Yamashiro Y. A fetal case of transient abnormal myelopoiesis with severe liver failure in Down syndrome: Prognostic value of serum markers. Pediatr Hematol Oncol. 2004;21:273-278. PMID: 15202167. PMCID: N/A.
- Taguchi T, Kiyokawa N, Takenouch H, Matsui J, Tang WR, Nakajima H, Suzuki K, Shiozawa Y, Saito M, Katagiri YU, Takahashi T, Karasuyama H, Matsuo Y, Okita H, Fujimoto J. Deficiency of BLNK hampers PLC-gamma2 phosphorylation and Ca2+ influx induced by the pre-B-cell receptor in human pre-B cells. Immunology. 2004;112:575-82. PMID: 15270728. PMCID: PMC1782531.
- Matsui J, Kiyokawa N, Takenouchi H, Taguchi T, Suzuki K, Shiozawa Y, Saito M, Tang WR, Katagiri YU, Okita H, Fujimoto J. Dietary bioflavonoids induce apoptosis in human leukemia cells. Leuk Res. 2005;29:573-81. PMID: 15755510. PMCID: N/A.
- Shiozawa Y, Kiyokawa N, Saito M, Fujimoto J, Hata JI, Yamashiro Y. Granulocytic sarcoma of the spine in a child without bone marrow involvement: a case report and literature review. Eur J Pediatr. 2005;164:616-20. PMID: 16012854. PMCID: N/A.
- 5. **Shiozawa Y**, Kiyokawa N, Fujimura J, Suzuki K, Yarita Y, Fujimoto J, Saito M, Yamashiro Y. Primary Malignant Lymphoma of the Central Nervous System in an Immunocompetent Child A Case Report. **J Pediatr Hematol Oncol**. 2005;27:561-4. PMID: 16217262. PMCID: N/A.
- Taguchi T, Takenouchi H, Matsui J, Tang WR, Itagaki M, Shiozawa Y, Suzuki K, Sakaguchi S, Ktagiri YU, Takahashi T, Okita H, Fujimoto J, Kiyokawa N. Involvement of insulin-like growth factor-I and insulin-like growth factor binding proteins in pro-B-cell development. Exp Hematol. 2006;34:508-18. PMID: 16569597. PMCID: N/A.
- Jung Y, Wang J, Song J, Shiozawa Y, Wang J, Havens A, Wang Z, Sun YX, Emerson SG, Krebsbach PH, Taichman RS. Annexin II expressed by osteoblasts and endothelial cells regulates stem cell adhesion, homing, and engraftment following transplantation. Blood. 2007;110:82-90. PMID: 17360942. PMCID: PMC1896130.
- Taguchi T, Takenouchi H, Shiozawa Y, Matsui J, Kitamura N, Miyagawa Y, Katagiri YU, Takahashi T, Okita H, Fujimoto J, Kiyokawa N. Interleukin-7 contributes to human pro-B-cell development in a mouse stromal cell-dependent culture system. Exp Hematol. 2007;35:1398-407. PMID: 17656007. PMCID: N/A.
- Wang J, Shiozawa Y, Wang J, Wang Y, Jung Y, Pienta KJ, Mehra R, Loberg R, Taichman RS. The role of CXCR7/RDC1 as a chemokine receptor for CXCL12/ SDF-1 in prostate cancer. J Biol Chem. 2008;283:4283-94. PMID: 18057003. PMCID: N/A.
- 10. **Shiozawa Y**, Havens AM, Pienta KJ, Taichman RS. The Bone Marrow Niche: Habitat to hematopoietic and mesenchymal stem cells, and unwitting host to molecular parasites. **Leukemia**. 2008;22:941-50. PMID: 18305549. PMCID: N/A.
- Havens AM, Pedersen EA, Shiozawa Y, Ying C, Jung Y, Sun YX, Neeley C, Wang J, Mehra R, Keller ET, McCauley LK, Loberg RD, Pienta KJ, Taichman RS. An *In Vivo* Mouse Model for Human Prostate Cancer Metastasis. Neoplasia. 2008;10:371-80. PMID: 18392141. PMCID: PMC2288546.

- Havens AM, Chiu E, Taba Jr. M, Wang J, Shiozawa Y, Jung Y, Taichman LS, D'Silva NJ, Gopalakrishnan R, Wang CY, Giannobile WV, Taichman RS. Stromal Derived Factor-1alpha (CXCL12) Levels Increase in Periodontal Disease. J Periodontol. 2008;79:845-53. PMID: 18454663. PMCID: PMC2582372.
- 13. **Shiozawa Y**, Sakaguchi S, Sakakibara O, Yagishita K, Saito M, Yamashiro Y. Urolithiasis in an acute lymphoblastic leukemia child during induction chemotherapy. **Pediatr Hematol Oncol**. 2008;25:359-63. PMID: 18484482. PMCID: N/A.
- 14. Jung Y, Song J, Shiozawa Y, Wang J, Wang Z, Williams B, Havens A, Schneider A, Ge C, Francaschi RT, McCauley LK, Krebsbach PH, Taichman RS. Hematopoietic Stem Cells Regulate Mesenchymal Stromal Cell Induction into Osteoblasts Thereby Participating in The Formation of the Stem Cell Niche. Stem Cells. 2008;26:2042-51. PMID: 18499897. PMCID: PMC3513687.
- Sun YX, Pedersen EA, Shiozawa Y, Havens AM, Jung Y, Wang J, Pienta KJ, Taichman RS. CD26/dipeptidyl peptidase IV regulates prostate cancer metastasis by degrading SDF-1/CXCL12. Clin Exp Metastasis. 2008;25:765-76. PMID: 18563594. PMCID: N/A.
- Shiozawa Y, Havens AM, Jung Y, Ziegler AM, Pedersen EA, Wang J, Wang J, Lu G, Roodman GD, Loberg RD, Pienta KJ, Taichman RS. Annexin II/Annexin II receptor axis regulates adhesion, migration, homing, and growth of prostate cancer. J Cell Biochem. 2008;105:370-80. PMID: 18636554. PMCID: PMC3614912.
- 17. **Shiozawa Y**, Takenouchi H, Taguchi T, Saito M, Katagiri YU, Okita H, Shimizu T, Yamashiro Y, Fujimoto J, Kiyokawa N. Human Osteoblasts Support Hematopoietic Cell Development in vitro. **Acta Haematol**. 2008;120:134-45. PMID: 19039204. PMCID: N/A.
- 18. Jung Y^{*}, Shiozawa Y^{*}, Wang J^{*}, Wang J, Wang Z, Pedersen EA, Lee CH, Hall CL, Hogg PJ, Krebsbach PH, Keller ET, Taichman RS. Expression of PGK1 By Prostate Cancer Cells Induces Bone Formation. Mol Cancer Res. 2009;7:1595-604. PMID: 19825988. PMCID: PMC2771860. (^{*} Authors contributed equally)
- Shiozawa Y, Pedersen EA, Taichman RS. GAS6/Mer axis regulates the homing and survival of the E2A/PBX1 positive B-cell precursor acute lymphoblastic leukemia in the bone marrow niche. Exp Hematol. 2010;38:132-40. PMID: 19922767. PMCID: PMC2815170.
- Shiozawa Y, Pedersen EA, Patel LR, Ziegler AM, Havens AM, Jung Y, Wang J, Zalucha S, Loberg RD, Pienta KJ, Taichman RS. GAS6/AXL axis regulates prostate cancer invasion, proliferation, and survival in the bone marrow niche. Neoplasia. 2010;12:116-27. PMID: 20126470. PMCID: PMC2814350.
- 21. Taichman RS, Wang Z^{*}, Shiozawa Y^{*}, Jung Y, Song J, Balduino A, Wang J, Havens A, Kucia M, Ratajczak MZ, Krebsbach PH. Prospective Identification and Skeletal Localization of Mesenchymal Stem Cells Capable of Multilineage Differentiation *In Vivo.* Stem Cells Dev. 2010;19:1557-70. PMID: 20446812. PMCID: PMC3089003. (*Authors contributed equally)
- 22. **Shiozawa Y**, Jung Y, Ziegler AM, Pedersen EA, Wang J, Wang Z, Song J, Wang J, Lee CH, Sud S, Pienta KJ, Krebsbach PH, Taichman RS. Erythropoietin Couples Hematopoiesis with Bone Formation. **PLoS One**. 2010;27:e10853. PMID: 20523730. PMCID: PMC2877712.
- 23. Jung Y, **Shiozawa Y**, Wang J, Patel LR, Havens AM, Song J, Krebsbach PH, Roodman GD, Taichman RS. Annexin 2 is a Regulator of SDF-1/CXCL12 Function in the Hematopoietic Stem Cell Endosteal Niche. **Exp Hematol**. 2011;39:151-66. PMID: 21108988. PMCID: PMC3026087.

24. Shiozawa Y, Pedersen EA, Havens AM, Jung Y, Mishra A, Joseph J, Kim JK, Patel LR, Ying C, Ziegler AM, Pienta MJ, Song J, Wang J, Loberg RD, Krebsbach PH, Pienta KJ, Taichman RS. Human prostate cancer metastases target the hematopoietic stem cell niche to establish footholds in mouse bone marrow. J Clin Invest. 2011;121:1298-312. PMID: 21436587. PMCID: PMC3069764.

• Comment in

Payton S. Prostate cancer: molecular insights into bone metastasis. Nat Rev Urol. 2011 Jun 10;8(6):291.

Keller ET. Prostate cancer cells metastasize to the hematopoietic stem cell niche in bone. Asian J Androl. 2011 Jul;13(4):622-3.

Schuettpelz LG, Link DC. Niche competition and cancer metastasis to bone. J Clin Invest. 2011 Apr;121(4):1253-5.

- 25. Kim J, Jung Y, Sun H, Joseph J, Mishra A, Shiozawa Y, Wang J, Krebsbach PH, Taichman RS. Erythropoietin mediated bone formation is regulated by mTOR signaling. J Cell Biochem. 2012;113:220-8. PMID: 21898543. PMCID: PMC3237787.
- 26. D'Souza S, Kurihara N, Shiozawa Y, Joseph J, Taichman R, Galson DL, Roodman GD. Annexin Il interactions with the annexin II receptor enhance multiple myeloma cell adhesion and growth in the bone marrow microenvironment. Blood. 2012;119:1888-96. PMID: 22223826. PMCID: PMC3293642.
- 27. Joseph J, Shiozawa Y, Jung Y, Kim JK, Pedersen E, Mishra A, Zalucha JL, Wang J, Keller ET, Pienta KJ, Taichman RS. Disseminated Prostate Cancer Cells Can Instruct Hematopoietic Stem and Progenitor Cells To Regulate Bone Phenotype. Mol Cancer Res. 2012;10:282-92. PMID: 22241219 . PMCID: PMC3307952. (Highlighted in this issue)

Comment in

Weckermann D. Words of wisdom. Re: Disseminated prostate cancer cells can instruct hematopoietic stem and progenitor cells to regulate bone phenotype. Eur Urol. 2012 Jul;62(1):181.

- 28. Mishra A, Wang J, **Shiozawa Y**, McGee S, Kim JK, Jung Y, Joseph J, Berry JE, Havens AM, Pienta KJ, Taichman RS. Hypoxia stabilizes GAS6/AXI signaling in metastatic prostate cancer. **Mol Cancer Res**. 2012;10:703-12. PMID: 22516347. PMCID: PMC3378814.
- 29. **Shiozawa Y**^{*}, Taichman RS^{*}. Getting Blood from Bone: An Emerging Understanding of the Role That Osteoblasts Play In Regulating Hematopoietic Stem Cells Within Their Niche. **Exp Hematol**. 2012;40:685-94. PMID: 22640993. PMCID: PMC3419274. (^{*}Co-corresponding authors)
- Sun HL, Jung Y, Shiozawa Y, Taichman RS, Krebsbach PH. Erythropoietin Modulates the Structure of Bone Morphogenetic Protein 2-Engineered Bone. Tissue Eng Part A. 2012;18:2095-105. PMID: 22703029. PMCID: PMC3463277.
- 31. Jung Y^{*}, Shiozawa Y^{*}, Wang J, McGregor N, Dai J, Park SI, Berry JE, Havens AM, Joseph J, Kim JK, Patel LR, Carmeliet P, Daignault S, Keller ET, McCauley LK, Pienta KJ, Taichman RS. Prevalence of Prostate Cancer Metastases Following Intravenous Inoculation Provides Clues into the Molecular Basis of Dormancy in the Bone Marrow Microenvironment. Neoplasia. 2012;14:429-39. PMID: 22745589. PMCID: PMC3384430. (^{*}Authors contributed equally)
- 32. Havens AM^{*}, Shiozawa Y^{*}, Jung Y^{*}, Sun H^{*}, Wang J, McGee S, Mishra A, Taichman LS, Danciu T, Yavanian G, Leary E, Krebsbach PH, Rodgerson D, Taichman RS. Human Very Small Embryonic-Like (VSEL) Cells Generate Skeletal Structures, *In Vivo.* Stem Cells Dev. 2013;22:622-30. PMID: 23020187. PMCID: PMC3564465. (^{*}Authors contributed equally)

- 33. Taichman RS, Patel LR, Bedenis R, Wang J, Weidner S, Schumann T, Yumoto K, Berry JE, **Shiozawa Y**, Pienta KJ. GAS6 Receptor Status Is Associated With Dormancy And Bone Metastatic Tumor Formation. **PLoS One**. 2013;8:e61873. PMID: 23637920. PMCID: PMC3634826.
- 34. Jung Y, Kim JK, Shiozawa Y, Wang J, Mishra A, Joseph J, Berry JE, McGee S, Lee E, Sun H, Wang J, Jin T, Zhang H, Dai J, Krebsbach PH, Keller ET, Pienta KJ, Taichman RS. Recruitment of mesenchymal stem cells into prostate tumours promotes metastasis. Nat Commun. 2013;4:1795. PMID: 23653207. PMCID: PMC3649763.
- 35. **Shiozawa Y**, McGee S, Pienta MJ, McGregor N, Jung Y, Wang J, Yumoto K, Berry JE, Pienta KJ, Taichman RS. Erythropoietin supports the survival of prostate cancer, but not growth and bone metastasis. **J Cell Biochem**. 2013;114:2471-8. PMID: 23696192. PMCID: PMC3978144.
- 36. Kim JK, Jung Y, Wang J, Joseph J, Mishra A, Krebsbach PH, Pienta KJ, Shiozawa Y, Taichman RS. TBK1 mediates prostate cancer dormancy through mTOR inhibition. Neoplasia. 2013;15:1064-74. PMID: 24027431. PMCID: PMC3769885.
- 37. Havens AM, Sun H, Shiozawa Y, Jung Y, Wang J, Mishra A, Jiang Y, O'Neill D, Krebsbach PH, Rodgerson DO, Taichman RS. Human and Murine Very Small Embryonic-Like (VSEL) Cells Represent Multipotent Tissue Progenitors, *In Vitro* and *In Vivo*. Stem Cells Dev. 2014;23:689-701. PMID: 24372153. PMCID: PMC3967374.
- Yumoto K, Berry JE, Taichman RS, Shiozawa Y. A novel method for monitoring tumor proliferation in vivo using fluorescent dye DiD. Cytometry Part A. 2014;85:548-55. PMID: 24700602. PMCID: PMC4143457.
- Jung Y, Wang J, Lee E, McGee S, Berry JE, Yumoto K, Dai J, Keller ET, Shiozawa Y, Taichman RS. Annexin 2-CXCL12 Interactions Regulate Prostate Cancer Growth in the Bone Marrow. Mol Cancer Res. 2015;13:197-207. PMID: 25139998. PMCID: PMC4297714.
- Seib FP, Berry JE, Shiozawa Y, Taichman RS, Kaplan DL. Tissue engineering a surrogate niche for metastatic cancer cells. Biomaterials. 2015;51:313-9. PMID: 25771021. PMCID: PMC4367489.
- Zalucha JL, Jung Y, Joseph J, Wang J, Berry JE, Shiozawa Y, Taichman RS. The Role of Osteoclasts in Early Dissemination of Prostate Cancer Tumor Cells. J Cancer Stem Cell Res. 2015;3:e1005. PMID: 26097863. PMCID: PMC4469294.
- 42. **Shiozawa Y**, Berry JE, Eber MR, Jung Y, Yumoto K, Cackowski FC, Yoon HJ, Parsana P, Mehra R, Wang J, McGee S, Lee E, Nagrath S, Pienta KJ, Taichman RS. The Marrow Niche Controls The Cancer Stem Cell Phenotype Of Disseminated Prostate Cancer. **Oncotarget**. 2016;7:41217-32. PMID: 27172799. PMCID: PMC5173053.
- Sharma S, Xing F, Liu Y, Wu K, Said N, Pochampally R, Shiozawa Y, Lin HK, Balaji KC, Watabe K. Secreted Protein Acidic and Rich in Cysteine (SPARC) Mediates Metastatic Dormancy of Prostate Cancer in the Bone. J Biol Chem. 2016;291:19351-63. PMID: 27422817. PMCID: PMC5016675.
- 44. Cackowski F, Eber MR, Rhee J, Decker A, Yumoto K, Berry JE, Lee E, **Shiozawa Y**, Jung Y, Aguirre-Ghiso JA, Taichman RS. Mer Tyrosine Kinase Regulates Disseminated Prostate Cancer Cellular Dormancy. **J Cell Biochem**. 2017;118:891-902. PMID: 27753136. PMCID: PMC5296338.
- 45. Kast RE, Skuli N, Cos S, Karpel-Massler G, **Shiozawa Y**, Goshen R, Halatsch ME. The ABC7 regimen: a new approach to metastatic breast cancer using seven common drugs to inhibit

epithelial-to-mesenchymal transition and augment capecitabine efficacy. **Breast Cancer (Dove Med Press)**. 2017;9:495-514. PMID: 28744157. PMCID: PMC5513700.

Invited Reviews:

- 1. Havens AM, Pedersen EA, **Shiozawa Y**, Taichman RS. Innovative Mouse Models for Metastatic Disease. **Drug Discov Today Dis Models**. 2009;6:27-31. PMID: N/A. PMCID: N/A.
- 2. Pedersen EA, **Shiozawa Y**, Taichman RS. Hematopoietic Stem Cells, Cancer Stem Cells, and the Hematopoietic Stem Cell Niche. **IBMS BoneKEy**. 2010;7:32-46. PMID: N/A. PMCID: N/A.
- Shiozawa Y, Pienta KJ, Taichman RS. Hematopoietic stem cell niche is a potential therapeutic target for bone metastatic tumors. Clin Cancer Res. 2011;17:5553-8. PMID: 21676926. PMCID: PMC3593121.
- 4. Mishra A, **Shiozawa Y**, Pienta KJ, Taichman RS. Homing of Cancer Cells to the bone. **Cancer Microenviron**. 2011;4:221-35. PMID: 21826451. PMCID: PMC3234327.
- Patel LR, Camacho DF, Shiozawa Y, Pienta KJ, Taichman RS. Mechanisms of cancer cell metastasis to the bone: a multistep process. Future Oncol. 2011;7:1285-97. PMID: 22044203. PMCID: PMC3258525.
- 6. McGee SJ, Havens AM, **Shiozawa Y**, Jung Y, Taichman RS. Effects of erythropoietin on the bone microenvironment. **Growth Factors**. 2012;30:22-8. PMID: 22117584. PMCID: PMC3295609.
- 7. Pedersen EA, Shiozawa Y, Mishra A, Taichman RS. Structure and function of the solid tumor niche. Front Biosci (Schol Ed). 2012;4:1-15. PMID: 22202039. PMCID: PMC3594847.
- 8. **Shiozawa Y**^{*}, Taichman RS^{*}. Cancer stem cells and the bone marrow microenvironment. **BoneKEy Rep**. 2012;1:48. PMID: 23745167. PMCID: PMC3671623. (^{*}Co-corresponding authors)
- Pedersen EA, Shiozawa Y, Pienta KJ, Taichman RS. The Prostate Cancer Bone Marrow Niche: More than Just "Fertile Soil". Asian J Androl. 2012;14:423-7. PMID: 22367179. PMCID: PMC3602965.
- 10. Yu C, **Shiozawa Y**, Taichman RS, McCauley LK, Pienta KJ, Keller ET. Prostate cancer and parasitism of the bone hematopoietic stem cell niche. **Crit Rev Eukaryot Gene Expr**. 2012;22:131-148. PMID: 22856431. PMCID: PMC3615416.
- 11. **Shiozawa Y**^{*}, Nie B, Pienta KJ, Morgan TM, Taichman RS^{*}. Cancer Stem Cells and their Role in Metastasis. **Pharmacol Ther**. 2013;138:285-93. PMID: 23384596. PMCID: PMC3602306. (^{*} Co-corresponding authors)
- 12. Noda K, **Shiozawa Y**, Komatsu Y. Molecular etiology of craniosynostosis. **Genet Disord Gene Ther**. 2013;1:101. PMID: N/A. PMCID: N/A.
- 13. Yumoto K, Eber MR, Berry JE, Taichman RS, **Shiozawa Y**. Molecular Pathways: Niches in metastatic dormancy. **Clin Cancer Res**. 2014;20:3384-9. PMID: 24756372. PMCID: PMC4079737.
- 14. Eber MR, Yumoto K, Berry JE, Taichman RS, **Shiozawa Y**. The roles of microenvironment in tumor dormancy. **Cancer Stud Open Access**. 2014;1:103. PMID: N/A. PMCID: N/A.

- Shiozawa Y^{*}, Eber MR, Berry JE, Taichman RS^{*}. Bone marrow as a metastatic niche for disseminated tumor cells from solid tumors. BoneKEy Rep. 2015;4:689. PMID: 26029360. PMCID: PMC4440229. (*Co-corresponding authors)
- 16. Dai J, Hensel J, Wang N, Kruithof-de Julio M, **Shiozawa Y**. Mouse models for studying prostate cancer bone metastasis. **BoneKEy Rep**. 2016;5:777. PMID: 26916039. PMCID: PMC4757481.
- 17. Tsuzuki S, Park SH, Eber MR, Peters CM, **Shiozawa Y**. Skeletal complications in cancer patients with bone metastases. **Int J Urol**. 2016;23:825-32. PMID: 27488133. PMCID: In Progress.
 - The Figure 1 is chosen as the Cover Figure of Int J Urol Vol. 23 No. 10

Invited Book Chapters:

- Havens AM, Shiozawa Y, Taichman RS. (2009) Blood-Bone Axis and Bone Marrow Microenvironment. In Bronner F and Farach-Carson MC (Eds.), *Bone and Cancer*, Topics in Bone Biology volume 5 (pp. 1-17). London: Springer. Topics in Bone Biology. 2009;5:1-17.
- 2. Forest AE, **Shiozawa Y**, Pienta KJ, Taichman RS. (2013) The Hematopoietic Stem Cell Niche and Bone Metastasis. In Jandial R (Ed.), *Metastatic Cancer: Clinical and Biological Perspectives*, Molecular biology intelligence unit. Texas, Landes Bioscience.
- 3. Mehta G, **Shiozawa Y**, Taichman RS. (2015) Hematopoietic Stem Cells and their Niches. In Ma PX (Ed.), *Biomaterials and Regenerative Medicine Book*. (pp. 44-63). Cambridge: Cambridge University Press.
- 4. Shiozawa Y^{*}, Taichman RS, Keller ET^{*}. (2013) Detection and Isolation of Human Disseminated Tumor Cells in the Murine Bone Marrow Stem Cell Niche. In Turksen K (Ed.), Stem cells and Niche: Methods and Protocols, Methods in Molecular Biology volume 1035 (pp. 207-15). New York: Humana Press. Methods Mol Biol. 2013;1035:207-15. PMID: 23959994. PMCID: N/A. (^{*} Cocorresponding authors)
- 5. Kawahara R, **Shiozawa Y**. (2015) Hematopoiesis. *Reference Module in Biomedical Sciences*. (pp. 1-4). Elsevier. 6-Mar-15 doi: 10.1016/B978-0-12-801238-3.05054-6.
- 6. Schrader J, White M, Silberstein P, **Shiozawa Y**. (2015) Hemophilia. *Reference Module in Biomedical Sciences*. (pp. 1-3). Elsevier. 9-Apr-15 doi: 10.1016/B978-0-12-801238-3.05056-X.
- 7. Miler SF, Thomas CY, **Shiozawa Y**. (2016) Molecular involvement of the bone marrow microenvironment in bone metastasis. In Ahmad A (Ed.), *Introduction to Cancer Metastasis.* In Press. Philadelphia, Elsevier.

Miscellaneous:

Letters to the Editor

 Shiozawa Y, Taichman RS. Erythropoietin and Bone Remodeling. [e-Letter], Blood (<u>http://bloodjournal.hematologylibrary.org/content/117/21/5631.full/reply#bloodjournal_el_2584</u>). Published July 25, 2011.

Editorial

- 1. Shiozawa Y, Taichman RS. Dysfunctional Niches as a Root of Hematopoietic Malignancy. Cell Stem Cell. 2010;6:399-400. PMID: 20452309. PMCID: N/A.
- 2. Shiozawa Y, Morgan TM, Taichman RS. Welcome to Journal of Cancer Stem Cell Research. Journal of Cancer Stem Cell Research. 2013;1:e1001. PMID: N/A. PMCID: N/A.
- 3. **Shiozawa Y**, Taichman RS. HSCs & Niche Relations Marked by CD166. **Blood**. 2014;124:471-2. PMID: 25061166. PMCID: N/A.
- 4. Shiozawa Y, Taichman RS. Elucidating which cell erythropoietin targets in bone. Nat Rev Endocrinol. 2015;11:263-4. PMID: 25752274. PMCID: N/A.

Abstracts/Scientific exhibits/Presentations at national meetings:

- 1. **Shiozawa Y**, Saito M, Miyata Y, Saito Y, Sakaguchi S, Sakakibara O, Fujimura J, Suzuki K, Yamashiro Y. Clinical evaluation of Cefpirome as an empiric therapy for febrile neutropenia among pediatric patients. **Pediatr Blood Cancer**. 2004;43:462-463. 36th Congress of the International Society of Pediatric Oncology. Oslo, Norway, September 16-19, 2004. Poster.
- Shiozawa Y, Saito M, Kiyokawa N, Fujimoto J, Hata J-I, Yamashiro Y, Yanai T, Yamataka A. Three cases of mesoblastic nephroma: two prenatal diagnosed cases and an adolscent case. Pediatr Blood Cancer. 2004;45:519. 37th Congress of the International Society of Pediatric Oncology. Vancouver, Canada, September 21-24, 2005. Poster.
- Shiozawa Y, Havens AM, Zalucha S, Ziegler AM, Pederson EA, Wang Y, Jung Y, Wang JH, Ying C, Wang J, Lu G, Roodman GD, Pienta KJ, Loberg RD, Taichman RS. Prostate cancer cells as parasites of the hematopoietic stem cell niche. Clin Exp Metastasis. 2008;25 (suppl):22. The symposium Skeletal Complications of Malignancy V, Philadelphia, PA, USA, October 25–27, 2007. Oral.
- Sakakibara O, Saito M, Sakaguchi S, Shiozawa Y, Fujimura J, Nagata S, Shimizu T, Yamashiro Y, Asahara T, Takahashi A, Nomoto K. Ribosomal RNA-targeted reverse transcription-PCR is sensitive and useful system in pediatric febrile neutropenia: farewell to empiric therapy. Pediatr Blood Cancer. 2007;49:428. 39th Congress of the International Society of Pediatric Oncology. Mumbai, India, October 31- November 3, 2007. Oral.
- Wang Z, Jung Y, Kucia M, Song J, Shiozawa Y, Ratajczak MZ, Krebsbach PH, Taichman RS. Prospective *In Vivo* Identification of Osteogenic Stem/Progenitor Cells from Bone Marrow-Derived Lin⁻/Sca-1⁺/CD45⁻ Cells. Blood. 2007;110:1409. The American Society of Hematology 49th Annual Meeting, Atlanta, GA, USA, December 8-11, 2007. Poster.
- Del Prete D, Lu G, Esteve, Shiozawa Y, Taichman RS, Roodman GD. Role of Annexin II and Annexin II Receptor in Homing Multiple Myeloma Cells in the Bone Marrow. JBMR. 2008;23:s118. The American Society for Bone and Mineral Research (ASBMR) 30th Annual Meeting, Montreal, Canada, September 12-16, 2008. Poster.
- Saito M, Takata O, Onda K, Saito Y, Sakaguchi S, Shiozawa Y, Suzuki K, Fujimura J, Takanashi T, Shimizu T. The effectiveness of dose-escalated piperacillin-tazobactam monotherapy for pediatric febrile neutropenia in Japan. 40th Congress of the International Society of Pediatric Oncology. Berlin, Germany, October 2-6, 2008. Poster.

- Jung Y, Ziegler AM, Shiozawa Y, Wang Z, Song J, Wang J, Wang J, Pienta MJ, Lee CH, Havens AM, Pedersen EA, Krebsbach PH, Taichman RS. Erythropoietin Stimulates Bone Formation In Vivo By Targeting Both HSCs and MSCs. Blood. 2008;112:123-24. The American Society of Hematology 50th Annual Meeting, San Francisco, CA, USA, December 6-9, 2008. Oral.
- Shiozawa Y, Pedersen EA, Havens AM, Jung Y, Ying C, Ziegler AM, Pienta MJ, Song J, Wang J, Loberg RD, Krebsbach PH, Pienta KJ, Taichman RS. Solid Tumors Target the HSC Niche to Establish Metastatic Footholds in the Marrow. Blood. 2008;112:206. The American Society of Hematology 50th Annual Meeting, San Francisco, CA, USA, December 6-9, 2008. Oral.
- Taichman RS, Shiozawa Y, Pedersen EA, Havens AM, Jung Y, Ying C, Ziegler AM, Pienta MJ, Song J, Wang J, Loberg RD, Krebsbach PH, Pienta KJ. Solid Tumors Target the Hematopoietic Stem Cell Niche for Metastasis. Bone. 2010;47:S265. The IX International Meeting on Cancer Induced Bone Disease, Arlington, VA, USA, October 28-31, 2009. Oral.
- 11. Shiozawa Y, Pedersen EA, Havens AM, Jung Y, Ziegler AM, Pienta MJ, Song J, Wang J, Loberg RD, Krebsbach PH, Pienta KJ, Taichman RS. Metastases From Solid Tumors Target the Hematopoietic Stem Cell Niche to Establish Footholds in Marrow. Bone. 2010;47:S289. The IX International Meeting on Cancer Induced Bone Disease, Arlington, VA, USA, October 28-31, 2009. Poster.
- 12. Pedersen EA, Shiozawa Y, Havens AM, Jung Y, Ying C, Ziegler AM, Pienta MJ, Song J, Wang J, Loberg RD, Krebsbach PH, Pienta KJ, Taichman RS. The role of the Hematopoietic Stem Cell Niche in Bone Metastasis Regulation. Bone. 2010;47:S288-9. The IX International Meeting on Cancer Induced Bone Disease, Arlington, VA, USA, October 28-31, 2009. Poster.
- D'Souza S, Shiozawa Y, Galson DL, Taichman RS, Roodman DG. Annexin II and Annexin II Receptor Interaction Enhance Multiple Myeloma Growth in the Bone Marrow Microenvironment. Bone. 2010;47:S284-5. The IX International Meeting on Cancer Induced Bone Disease, Arlington, USA, October 28-31, 2009. Poster.
- 14. Patel LR, **Shiozawa Y**, Taichman RS, Pienta KJ. CD133+/CD44+ Cancer Stem Cells Represent a Disproportionately Large Fraction of Early Bone Disseminated Prostate Cancer Tumor Cells. The AACR 101st Annual Meeting, Washington DC, USA, April 17-21, 2010. Poster.
- 15. D'Souza S, Kurihara N, **Shiozawa Y**, Taichman RS, Galson D, Roodman DG. Annexin II Interactions with the Annexin II Receptor Enhance Multiple Myeloma Cell Adhesion and Growth In the Bone Marrow Microenvironment. **Blood**. 2010;116:130. The American Society of Hematology 52th Annual Meeting, Orlando, FL, USA, December 4-7, 2010. Oral.
- Ziegler A, Jung Y, Shiozawa Y, Wang J, Krebsbach P, Taichma RS. Erythropoietin Couples Hamatopoiesis with Bone Formation. IADR/AADR/CADR 89th general session and exhibition, San Diego, CA, USA, March 16-19, 2011. Poster.
- 17. Shiozawa Y, Jung Y, Mishra A, Joseph J, Kim JK, Wang J, Iura A, Marotte H, Koch AE, Taichman RS. CXCL16/CXCR6 chemokine axis participates in the growth of prostate cancer by recruiting mescenchymal stem-like cells from bone marrow. Congressionally Directed Medical Research Programs IMPaCT 2011, Orlando, FL, USA, March 9-12, 2011. Oral & Poster.
- Shiozawa Y, Taichman RS, Jung Y, Wang J, Wang J, Wang Z, Pedersen EA, Lee CH, Hall CL, Hogg PJ, Krebsbach PH, Keller ET. Expression of PGK1 By Prostate Cancer Cells Induces Bone Formation. Congressionally Directed Medical Research Programs IMPaCT 2011, Orlando, FL, USA, March 9-12, 2011. Poster.

- 19. **Shiozawa Y**, Taichman RS, Joseph J, Jung Y, Mishra A, Pedersen EA, Kim JK, Wang J, Pienta KJ. The Bone Phenotype Of Prostate Cancer May Be Due To The Ability Of Cancer To Alter Hematopoietic Stem Cell Function. Congressionally Directed Medical Research Programs IMPaCT 2011, Orlando, FL, USA, March 9-12, 2011. Oral & Poster.
- 20. Shiozawa Y, Patel LR, Mishra A, Jung Y, Joseph J, Kim JK. Wang J, Pienta KJ, Taichman RS. The CD133+/CD44+ population is Enriched in Disseminated Human Prostate Cancer Cells Isolated from Murine Marrow. The XI International Conference on Cancer-Induced Bone Disease, Chicago, IL, USA, November 30-December 3, 2011. Poster.
- 21. Taichman RS, Shiozawa Y, Pedersen EA, Havens AM, Jung Y, Lalit LR, Ying C, Pienta MJ, Song J, Wang J, Krebsbach PH, Pient KJ. Hematopoietic Stem Cell Niche is the Target of Metastatic Cells from Solid Tumors. The XI International Conference on Cancer-Induced Bone Disease, Chicago, IL, USA, November 30-December 3, 2011. Oral.
- 22. D'Souza S, Kurihara N, Shiozawa Y, Joseph J, Taichman RS, Galson D, Roodman DG. Annexin II Interactions with the Annexin II Receptor Enhance Multiple Myeloma Cell Adhesion and Growth in the Bone Marrow Microenvironment. Blood. 2011;118:3942. The American Society of Hematology 53th Annual Meeting, San Diego, CA, USA, December 10-13, 2011. Poster.
- 23. Zalucha JL, Joseph J, **Shiozawa Y**, Jung Y, Wang J, Taichman RS. Determining osteoclasts' role in hematopoietic mobilization and metastasis. The AADR/CADR 2012 Annual Meeting, Tampa, FL, USA, March 21-24, 2012. Oral.
- 24. Havens AM, Sun H, **Shiozawa Y**, Jung Y, Wang J, McGee S, Mishra A, Taichman LS, Danciu T, Jiang Y, Yavanian G, Leary E, Krebsbach PH, Rodgerson D, Taichman RS. Multilineage Tissue Regeneration by Human Very Small Embryonic-Like Cells. IADR/AADR/CADR General Session & Exhibition, Seattle, WA, USA, March 20-23, 2013. Poster.
- 25. Zalucha JL, Joseph J, Jung Y, Berry J, Wang J, **Shiozawa Y**, Taichman RS. The Role of Osteoclasts in Early Dissemination of Tumor Cells. The AADR/CADR 2014 Annual Meeting, Charlotte, NC, USA, March 19-22, 2014. Oral.
- 26. Jung Y, Berry JE, Shiozawa Y, Eber M, Yumoto K, Wang J, Jiang Y, Rodgerson D, Taichman RS. Efficacy of Human VSELs to Reverse Bone Loss in Osteoporosis. The 2015 IADR/AADR/CADR General Session in Boston, MA, USA, March 11-14, 2015. Oral.
- 27. Tsuzuki S, Eber MR, Miler SF, Park SH, Widner DB, **Shiozawa Y**. The effects of neuropeptides on the prostate cancer progression. IMPaCT Young Investigator Meeting, Baltimore, MD, USA, August 4-5, 2016. Poster.
- 28. Mamo M, Park SH, Eber MR, Widner DB, Tsuzuki S, Warren CC, Strickland TB, Peters CM, Shiozawa Y. The Interaction between Disseminated Prostate Cancer Cells and Sensory Nerves. The 2016 Annual Biomedical Research Conference for Minority Students (ABRCMS), Tampa, FL, USA, November 9-12, 2016. Poster.
- Williams SN, Eber MR, Tsuzuki S, Park SH, Shiozawa Y. Does histamine influence the osteoblastic activities that help regulate the progression of bone metastatic prostate cancer? The 2017 Annual Biomedical Research Conference for Minority Students (ABRCMS), Phoenix, AZ, USA, November 1-4, 2017. Poster.

GRADUATE STUDENTS/RESIDENTS/FELLOWS ADVISED:

Wake Forest School of Medicine:

Courses taught:

- Metastasis, Cancer Cell Biology (CABI 705), Lecture, (2015: 3h, 2016: 3h, 2017: 3h) (directed by Dr. William Gmeiner)
- Prostate Cancer: Oncogenesis and Progression, Carcinogenesis, DNA Damage and Repair (MCB721), Lecture, (**2015**: 1.5h, **2016**: 1.5h, **2017**: 1.5h) (directed by Dr. Hui-Wen Lo)
- Prostate Cancer Etiology and Treatment, Cancer Cell Biology (CABI 705), Lecture, (2015: 3h, 2016: 3h, 2017: 3h) (directed by Dr. William Gmeiner)
- Prostate cancer / Bone metastatic niche / Cancer dormancy, Advanced Topics in Cancer Biology (CABI 711), Facilitator, (2015: 1h, 2016: 1h) (directed by Dr. Ravi Singh (2015, 2016), and by Dr. Neveen Said (2017))
- Prostate cancer / Bone metastatic niche / Cancer dormancy, Advanced Topics in Cancer Biology (CABI 712), Facilitator, (**2016**: 1h, **2017**: 1h) (directed by Dr. Ravi Singh)
- Prostate cancer diagnosis and treatment, North Carolina Summer Undergraduate Prostate Cancer Research Program, Lecture, (**2016**: 1h, **2017**: 1h) (directed by Dr. Kounosuke Watabe)
- A small group discussion using the Problem-Based Learning (PBL) case-based method, Scientific Integrity (Grad 713), Facilitator, (**2016**: 8h) (directed by Dr. J. Charles Eldridge)
- A small group discussion using the Problem-Based Learning (PBL) case-based method, Scientific Integrity (Grad 714), Facilitator, (**2017**: 18h) (directed by Dr. J. Charles Eldridge)
- Bone metastasis and treatment, Molecular Pathogenesis of Cancer (MCB722), Lecture, (**2017**: 1.5h) (directed by Drs. Thaddeus J. Wadas and Bethany Kerr)
- Tumor Metastasis, Oncology Core Curriculum Lecture Series (CABI 707), Lecture, (**2017**: 1h) (directed by Dr. Susan Melin)

Student research:

Undergraduate Students:	
2015-2017	 Sloan Miler (Wake Forest University) Cancer Biology Research mentor (BIO 391) Undergraduate Student Stipend for Participation in International Conferences/Activities
2016-2016	Caroline Warren (Wake Forest University) Cancer Biology Research mentor
2016-2016	 Mahder Mamo (North Carolina A&T State University) Cancer Biology Research mentor 2016 North Carolina Summer Undergraduate Prostate Cancer Research Program

2016-2016	 Taryn Strickland (Johnson C. Smith University) Cancer Biology Research mentor The CRUISE program
2016-	Sara Nakamura-Peek (Wake Forest University) Cancer Biology Research mentor
2016-	L. Daniela Smith (Wake Forest University) Cancer Biology Research mentor
2017-2017	Claudia Thornton (Wake Forest University) Cancer Biology Research mentor
2017-2017	Aaron Sunil (Wake Forest University) Cancer Biology Research mentor
2017-	Ethan Green (Wake Forest University) Cancer Biology Research mentor
2017-2017	 Selena Williams (North Carolina A&T State University) Cancer Biology Research mentor 2017 North Carolina Summer Undergraduate Prostate Cancer Research Program
2017-	Amy Medford (Wake Forest University) Cancer Biology Research mentor
2017-	Molly Sohn (Wake Forest University) Cancer Biology Research mentor
2017-	Sarah Sharp (Wake Forest University) Cancer Biology Research mentor
Medical Students: 2017-2017	Mary Booker Cancer Biology Research mentor • The 2017 Medical Student Research Program, provided by Laura Scales Cancer Fund
Graduate Students: 2015-2016	Amanda Hyre Cancer Biology Research mentor (rotation)

2016-	D. Brooke Widner Cancer Biology Research mentor
Postdoctoral Fellows: 2015-	Shunsuke Tsuzuki Cancer Biology Research mentor
2016-	Sun Hee Park Cancer Biology Research mentor

University of Michigan School of Dentistry:

Courses taught:

- Bone marrow transplantation, Advanced Topics in Oral Pathology OP 711, Lecture, (**2010**: 1h, **2011**: 1h, **2012**: 1h, **2013**: 1h) (directed by Dr. Theodora Danciu)
- Bone marrow niche and prostate cancer bone metastasis, Advanced Cellular and Molecular Biology course MCDB 440 Cell Cycle Control and Cancer, Guest Lecture, (**2014**: 2h) (directed by Dr. Laura Buttitta)

Student research:

High School Students: 2013-2013	Jacob Scherba Periodontics and Oral Medicine Lab supervisor
Undergraduate Students: 2006-2010	Elisabeth A. Pedersen Periodontics and Oral Medicine Lab supervisor
2007-2008	Michael J. Pienta Periodontics and Oral Medicine Lab supervisor
2010-2014	Samantha J. McGee Periodontics and Oral Medicine Lab supervisor
2012-2015	 Matthew R. Eber Periodontics and Oral Medicine Lab supervisor 4th Prize Poster Award - 6th Annual Prostate Cancer Program Retreat (2013).
2013-2013	Felicia Scharf Periodontics and Oral Medicine Lab supervisor
2013-2015	 Lulia Kana Periodontics and Oral Medicine Lab supervisor Third place award in Undergraduate, DDS, DH, MS/Certificate - Basic Science and Research - the University of Michigan School of Dentistry 2014 Research Day (2014).
2013-2013	Alexis N. Witkin Periodontics and Oral Medicine Lab supervisor
2013-2013	James Rhee Periodontics and Oral Medicine Lab supervisor

Visiting Undergraduate Students (2012-2013	(from Brazil): Eric Spinetti Periodontics and Oral Medicine Lab supervisor
Undergraduate/Dental Students: 2006-2010	 Anne M. Zeigler Periodontics and Oral Medicine Lab supervisor Grand Prize - ADA Dentsply Award in the University of Michigan School of Dentistry 2010 Research Day (2010).
Undergraduate/Master Students: 2006-2013	 Aaron M. Havens Periodontics and Oral Medicine Lab supervisor Second prize - the Edward Hatton Competition of the American Associations for Dental Research (2013).
Dental hygiene/Dental Students: 2008-2015	 Janet L. Zalucha Periodontics and Oral Medicine Lab supervisor Audience Choice Award - the University of Michigan School of Dentistry 2011 Research Day (2011).
Dental Students: 2010-2013	 Alexandra Forest Periodontics and Oral Medicine Lab supervisor Grand Prize - ADA Dentsply Award - the University of Michigan School of Dentistry 2012 Research Day (2012). Young Investigator Award - the University of Michigan 1st symposium on Head & Neck cancer stem cells (2012). Third place award in basic science research - 2012 the 53rd annual American Dental Association/Dentsply Student Clinician Research Award Program (2013).
2013-2015	Baxter Jones Periodontics and Oral Medicine Lab supervisor
Master Students: 2009-2012	Lalit R. Patel Urology and Internal Medicine Lab supervisor
Graduate Students (rotation): 2014-2014	Ann Delker Periodontics and Oral Medicine Lab supervisor
Graduate Students: 2008-2011	Amy Hsiao
Biomeo	dical Engineering
--------	-------------------
Lab su	pervisor

Postdoctoral Fellows	
2010-2012	Jeena Joseph Periodontics and Oral Medicine Lab supervisor
2010-2011	Anjali Mishra Periodontics and Oral Medicine Lab supervisor
2010-2011	Jinkoo Kim Periodontics and Oral Medicine Lab supervisor
2012-2012	Monika Verma Periodontics and Oral Medicine Lab supervisor
2012-2012	Biao Nie Periodontics and Oral Medicine Lab supervisor
2012-2015	Kenji Yumoto Periodontics and Oral Medicine Lab supervisor
2013-2014	 Jacqueline Jones Periodontics and Oral Medicine (Dr. McCauley's lab) Lab supervisor DOD Postdoctoral Training Award (2014). Assistant Professor at Troy University.
2014-2015	Frank Cackowski Periodontics and Oral Medicine Lab supervisor