AD/A-000 802

「「「「いうい」

WEATHER EXTREMES AROUND THE WORLD

Pauline Riordan

Army Engineer Topographic Laboratories Fort Belvoir, Virginia

April 1974 '

DISTRIBUTED BY:

National Technical Information Service U. S. DEPARTMENT OF COMMERCE



۰.

Destroy this report when no longer needed. Do not return it to the originator.

The citation in this report of trade names of commercially available products does not constitute official endorsement or approval of the use of such products. A STREET ST

ii a

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

BLANK PAGES IN THIS DOCUMENT WERE NOT FILMED

EC	URITY CLASSIFICATION OF THIS PAGE (When Dete	Entered)	TUTA OCOOD
	REPORT DOCUMENTATION	PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
, 1	REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	ETL-TR-74-5	<u> </u>	
• •	T(TLE (and Subtitio) W/E ATTLEED EV/DENIES A DOLLNIN THE	SWADL IN	S. TYPE OF REPORT & PERIOD COVERED
	WEATHER EXTREMES AROUND THE		
	(Revision of NLABS Report 1 R-70-45-E	(0)	S PERFORMING ORG. REPORT NUMBER
7	AUTHOR(e)		S. CONTRACT OR OR # 11UNBER(+)
1	Pauline Riordan		
	PERFORMING OROANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK
(Geographic Applications Division, Grog	aphic Sciences	AREA & WORK UNIT NUMBERS
	Laboratory, U. S. Army Engineer Topog	raphic Laboratories	1T162112A129
1	Fort Belvoir, Virginia 22060		
۰.	CONTROLLING OFFICE NAME AND ADDRESS		April 1974
1	U, 5. Army Engineer Topographic Labor Fort Balvoir, Viscinia, 22060	atorns	13. NUMBER OF PAGES
1	ron Denvon, virginia 44000		
4,	MONITORING AGENCY NAME & AODRESS(II differen	t from Controlling Office)	IS. SECURITY CLASS. (of this report)
			Unclassified
			ISA DECLASSIFICATION DOWNOR ADING
	.4		SCHEOULE
b .	DISTRIBUTION STATEMENT (ut this Report)		
	Approved for public release; distribution	unlimited.	
7.	Approved for public release; distribution	unlimited. In Block 20, 18 different fro	
17.	Approved for public release; distribution	unlimited. In Black 20, If different fra	
7.	Approved for public release; distribution	unlimited. In Block 20, if different fro	
7.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES	unlimited. In Block 20, it different fre	- MOV 15 1974
7.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obstract entered SUPPLEMENTARY NOTES	unlimited. In Block 20, if different fro	
7.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obstract entered SUPPLEMENTARY NOTES	unlimitud. In Block 20, if different fro	
7.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obstreet entered SUPPLEMENTARY NOTES	unlimited. In Black 20, if different fro d Identify by black number)	
7. 6.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed antered SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse aids if necessary an Extrumes Price	n unlimited. In Black 20, it different fro d identify by black number) ipitation (meteorolo	т Короно
7. 8.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse aids if necessary an Extremes Pre-	unlimited. In Block 20, it different fro d Identify by block number) ipitation (meteorolo osplierie pressure	т Короно DDC DC NOV 15 1974 NOV 15 1974 DC DC D Humidity Fog
7. 0. 1	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioci entered SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side it necessary on Extremes Proc Weather Atm Weather Atm	unlimited. In Block 20, if different fro d Identify by block number) ipitation (meteorolo ospheric pressure r railiation)	gy) Humidity Fog Thunderstorms
7. •.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioci entered SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse aids if necessary an Extremes Price Weather Atm Weather Atm Weather Sola Methorological charts Wind	unlimited. In Block 20, if different fro d Identify by block number) ipitation (meteorolo ospheric pressure r railiation) 1 (meteorology)	gy) Humidity Fog Thunderstorms
7. 8. 1	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioci entered SUPPLEMENTARY NOTES Extremes Price Weather Atm Weather observations Sola Methorological charts Wind Temperature Weat	unlimited. In Block 20, if different fro d Identify by block number) ipitation (netcorolo ospheric pressure r railiation) 1 (metcorology) ther intelligence	gy) Humidity Fog Thunderstorms
7.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioci entered SUPPLEMENTARY NOTES Extremes Price Weather Atm Weather observations Sola Methorological charts Wine Temperature Weather Weat	a unlimited. In Block 20, if different fro d Identify by block number) ipitation (netcorolo ospheric pressure r railiation) 1 (metcorology) ther intelligence I Identify by block number)	gy) Humidity Fog Thunderstorms
7.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES Extremes Prec Weather Atm Weather observations Sola Methorological charts Wind Temperature Weat ABSTRACT (Continue on reverse side II necessary and ABSTRACT (Continue on reverse side II necessary and This report consists of a worldwide m America, with comments on the reliabilitions	a unlimited. In Block 20, if different fro In Block 20, if different fro ipitation (meteorolo pspheric pressure r ratilation) 1 (meteorology) ther intelligence I dentify by block number) ap of weather extrer ty of the records sho variations, greatest	r Report) DDC DC NOV 15 1974 NOV 15 1974 USGISTIC D gy) Humidity Fog Thunderstorms nes and a separate map for North own. Included are highest and and hast amounts of precipitation
7. 8. 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES Extremes Prec Weather Atm Weather Observations Sola Meteorological charts Wind Temperature Weather Wrai ABSTRACT (Continue on reverse side if necessary end ABSTRACT (Continue on reverse side if necessary end This report consists of a worldwide m America, with comments on the reliabilit lowest temperatures, largest temperature for various ilurations, maximum precipit	a unlimited. In Block 20, It different fro Dipitation (meteorolo Dispheric pressure r radiation l (meteorology) ther intelligence d formit by block number) ap of weather extrer ty of the records sho variations, greatest ation variability, greatest	gy) Humidity Fog Thunderstorms
7. 0.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES KEY WORDS (Continue on reveice aids if necessary an Extremes Prec Weather Atm Weather observations Sola Methorological charts Wind Temperature Wrai ABSTRACT (Continue on reverce eids if necessary and This report consists of a worldwide m America, with comments on the reliabili lowest temperatures, largest temperature for various ilurations, maximum precipit leighest and lowest atmospheric pressure.	d dentily by block number) ipitation (meteorolo ospheric pressure r radiation l (meteorology) ther intelligence (dentily by block number) ap of weather extrer ty of the records sho variations, greatest ation variability, gre	gy) Humidity Fog Thunderstorms
7. 7. 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES EXTREMES Pre- Weather Atm Weather observations Sola Meteorological charts Wind Temperature Wrai ABSTRACT (Continue on reverse olde II necessary and This report consists of a worldwide m America, with comments on the reliabili lowest temperatures, largest temperature for various ultrations, maximum precipit leighest and lowest atmospheric pressure, snowfall, highest wind speed, high dew p	a unlimited. In Block 20, It different fro i pitation (meteorolo ospheric pressure r radiation 1 (meteorology) ther intelligence I dentity by block number) ap of weather extrer ty of the records sho variations, greatest ation variability, gre , highest solar radiation i and most frequ	gy) Humidity Fog Thunderstorms
7. •.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES EXTREMES Prec Weather Atm Weather observations Sola Meteorological charts Wind Temperature Weat ABSTRACT (Continue on reverse side if necessary on This report consists of a worldwide m America, with comments on the reliabili lowest temperatures, largest temperature for various durations, maximum precipit lightest and lowest atmospheric pressure, snowfall, highest wind speed, high dew p (Co	a unlimited. In Block 20, it different from in Block 20, it different from ipitation (meteorology) ipitation (meteorology) ther intelligence i (meteorology) ther intelligence i (dentify by block number) ap of weather extrem ty of the records sho variations, greatest ation variability, greatest	gy) Humidity Fog Thunderstorms
7. 0.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioed entered SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side if necessary on Extremes Pre Weather Atm Weather Atm Weather Sola Meteorological charts Wine Temperature Weat ABSTRACT (Continue on reverse side if necessary on ABSTRACT (Continue on reverse side if necessary on America, with comments on the reliabili lowest temperatures, largest temperature for various iltrations, maximum precipit loighest and lowest atmospheric pressure, showfall, highest wind speed, high dew p (Co	a unlimited. In Block 20, if different fro block 20, if different fro pipitation (meteorolo ospheric pressure r radiation) I (meteorology) ther intelligence I fondify by block number) ap of weather extrer ty of the records sho variations, greatest ation variability, gre highest solar radiation, highest solar radiation oint, and most frequenting)	gy) Humidity Fog Thunderstorms
•	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioce entered SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse aide if necessary an Extremes Prec Weather Atm Weather observations Sola Methorological charts Wind Temperature Weal ABSTRACY (Continue on reverse aide if necessary and This report consists of a worldwide m America, with comments on the reliabili lowest temperatures, largest temperature for various iltrations, maximum precipit loighest and lowest atmospheric pressure, snowfall, highest wind speed, high dew p (Co	a unlimited. In Block 20, if different fro Block 20, if different fro pipitation (meteorolo ospheric pressure r raliation I (meteorology) ther intelligence (Identify by block number) ap of weather extrer ty of the records sho variations, greatest ation variability, gre highest solar raliati point, and most frequ- ntinued)	gy) Humidity Fog Thunderstorms mes and a separate map for North own. Included are highest and and heast amounts of precipitation atest thunderstorm frequency, ion largest hailstomes, greatest ent occurrence of deuse fog.
17. (6. 17. 10. 1	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioed entered SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse aide if necessary an Extremes Price Weather Atm Weather observations Sola Methorological charts Wind Temperature Weat ABSTRACT (Continue on reverse aide if necessary and This report consists of a worldwide m America, with comments on the reliabili lowest temperatures, largest temperature for various durations, maximum precipit tighest and lowest atmospheric pressure, showfall, highest wind speed, high dew p (Co	a unlimited. In Block 20, it different fro d Identify by block number) ipitation (meteorolo ospheric pressure r raliation) I (meteorology) ther intelligence r Identify by block number) ap of weather extrer ty of the records sho variations, greatest ation variability, gre tooint, and most frequentinued) .ere security CLAN	Balt Concernance of druse for.
•. •.	Approved for public release; distribution DISTRIBUTION STATEMENT (of the obvioest entered SUPPLEMENTARY NOTES EXTREMES Continue on reverce side if necessary an Extremes Price Weather Atm Weather Observations Sola Methorological charts Wine Temperature Weat ABSTRACT (Continue on reverce side if necessary and This report consists of a worldwide m America, with comments on the reliabili lowest temperatures, largest temperature for various durations, maximum precipit lighest and lowest atmospheric pressure, snowfall, highest wind speed, high dew p (Co rooms 1473 EDITION OF 1 NOV 65 IS OBSOL Reproduced by NATIONAL TECHNICAL	a unlimited. In Block 20, if different from a identify by block number) ipitation (meteorolo pspheric pressure r raliation) 1 (meteorology) ther intelligence I dentify by block number) ap of weather extrem ty of the records sho variations, greatest ation variability, gre- tooint, and most frequenting ntinued) .eve security CLAN	(WCLASSIFIED)
7. •	Approved for public release; distribution DISTRIBUTION STATEMENT (of the observed entered SUPPLEMENTARY NOTES EXTREMES Continue on reveice side if necessary an Extremes Prec Weather Atm Weather observations Sola Meteorological charts Wind Temperature Weat ABSTRACT (Continue on reverce side if necessary and ABSTRACT (Continue on reverce side if necessary and This report consists of a worldwide m America, with comments on the reliabilit lowest temperatures, largest temperature for various durations, maximum precipit lighest and lowest atmospheric pressure. snowfall, highest wind speed, high drw p (Co PORM 1473 EDITION OF 1 NOV 65 IS OBSOL Reproduced by ATTIONAL TECHNICAL INFORMATION SERVIC	a unlimited. In Block 20, 11 different fro d identify by block number) ipitation (meteorolo ospheric pressure r raliation) 1 (meteorology) ther intelligence 1 (meteorology) ther intelligence 1 (meteorology) ther intelligence t dentify block number) ap of weather extrem ty of the records sho + variations, greatest ation variability, gre variations, and most frequenting ntinued) .ere security CLAN L i E	(WCLASSIFIED)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)

.•

(Block 20 Cont'd)

Where appropriate, the value for the highest and lowest annual mean is also given. As far as possible, the records are taken from official sources, and all of them are documented. Conditions of site, instrumentation, observational procedure, and other factors such as environmental and meteorological conditions pertinent to the reliability of extremes are discussed.

49.543 - Ft Belvolf

UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE/When Date Entered)

PREFACE

In 1955, a map titled "World Weather Extremes" was prepared by the Cartography Branch of the Environmental Protection Research Division, Quartermaster Research and Engineering Center, Natick, Massachusetts, for distribution to visitors and other interested persons. This map showed the location of certain meteorological extremes which had been recorded in elimatological publications. In subsequent years the map was periodically revised as new records came to light and the number of requests for it increased. It has been reprinted in a number of publications, both military and civilian, and has been widely used for instruction and display purposes.

In 1969, a need for an updated map and some discussion of the records shown appeared in the Earth Sciences Laboratory at Natick in connection with a revision of Military Standard 210A, "Climatic Extremes for Military Equipment." Natick Laboratories had been assigned responsibility for certain portions of the standard. As an indication of the most extreme climatic conditions that have been recorded in various parts of the earth, the map gives an approximation of the absolute limits within which design criteria must be established.

In addition to the revised map of world weather extremes, a new map showing extremes in North America was added and the sources for records on both maps were reviewed and documented. When controversial records were included, the arguments for and against their acceptance were examined and summarized in the text. For many of the records, the text included such information as the conditions during occurrence of the extremes, problems of measuring or recording them, and the kinds of investigations carried out by the U. S. Weather Bnreau before they were accepted. Thus, for the first time a map of extreme weather occurrences was presented together with full documentation of the sources of the records and references to pertinent comments in the technical literature.

Weather records continue to be broken and the maps and text again have been updated. In addition, some further background material has been included and some changes have been made to improve textual clarity. This work has been carried out in the Environmental Criteria Branch of the Engineer Topographic Laboratories at Fort Belvoir, Virginia, which is the successor organization to the former Earth Sciences Laboratory at Natiek.

Valuable advice and information were received from Dr. M. A. Arkin, Environmental Data Service, National Oceanic and Atmospheric Administration (NOAA) for both this revision and the original report. His assistance is acknowledged with appreciation. The cartography in the original report was completed by Ms Kristin Gill and Ms Olive Lesueur,

iii

under the direction of Aubrey Greenwald, Jr., Chief of the Cartography Office, Earth Seicnees Laboratory. It has been reproduced, with the necessary updating for this report, by SP-4 Richard Vail under the direction of James Smith of the Topographie Products Development Branch of Engineer Topographic Laboratories. Their assistance and suggestions are appreciated as are the suggestions made for printing the records by categories on the reverse of the map sheets by Dr. William Robison and for setting up eoordinates for locating the places of occurrence of the records on the world map by Ms Jane Kroll, both of the Engineer Topographic Laboratories.

CONTENTS

٤.

ê

1

Section	Title	Page
	PREFACE	iii
I	INTRODUCTION	
	1. Subject 2. Documentation	1 1
п	COMMENTARY ON RELIABILITY OF THE RECORDS	17
	 Temperature Precipitation Other Extremes 	17 28 40
Ш	CONCLUSIONS	
	6. Conclusions	51
	Weather Extremes Around the World (map, following page 1)	
	Weather Extremes in North America (map, following page 1)	

v

WEATHER EXTREMES AROUND THE WORLD

I. INTRODUCTION

1. Subject. The world map and the map of North America were prepared primarily as aids in specifying extreme climatic conditions for design of military equipment. The two maps together contain approximately 100 records. The categories of extremes indicated in MIL-STD 210A were kept in mind in the selection of records, but selection was not limited to these categories. This report is primarily intended to review the validity of the records. A listing of the records, with documentation of their sources, is followed by a commentary on their reliability.

The representativeness and significance of weather records in general are commented upon, and for each of the main elements, certain factors bearing on reliability are noted. Among these are measuring instruments and problems, environmental conditions which favor occurrence of extremes, theoretical limits of occurrence, and the geographical areas and seasons in which the probability of extremes is greatest. Following this general treatment, pertinent information is given about individual records and their documentation. There is much information available about some of the extremes, especially controversial ones, but little or nothing about others. Similarly, records are more easily available to us for the United States and Canada than for other countries. To make room for showing these records, and because Americans would be especially interested in records of their own continent, the North America map has been included.

In addition to documenting each of the records, we have coordinated them, where possible, with determinations of the Environmental Data Service (EDS) part of the U.S. NOAA. In the United States, EDS is responsible for archiving and disseminating weather data, and in line with these functions it validates extreme occurrences, both at home and abroad.¹ Other countries follow a similar policy, but there is, as yet, no world agency that determines and establishes a register of weather extremes. As a result, a given record might be accepted by EDS but not by its equivalent agency in some other country.

2. Documentation. To facilitate finding the extremes for a particular element, the records are listed, with documentation, by elements (see the Table, pages 7-15). Records for each element are arranged in order of decreasing extremity beginning with the world's record if accepted as such.

¹ M. A. Arkin, Chief, Foreign Branch, Environmental Data Service, U. S. Environmental Science Services Administration, letter dated 10 Sep 69.



a series and a series of the series of the



· ·

LOCATION OF WEATHER EXTREMES BY ELEMENT - WORLD MAP

I. TEMPERATURE

A. High		C. Least Precip
	World: El Azizia, Libya B7 W. llemisphere: Death Valley, California, B3	Number
	Asia: Tirat Tsvi, Israel BS Australia: Cloncurry Oucensland D11	D. Low Average
	Europe: Seville, Spain B6	Warld
	S. America: Rivadavia, Argentina D5	Africa
	Antarctica: Esperanza, Antarctic Peninsula F5	N. Ame
	Sea surjace: Persian Gulj C8	Asia:
	Annual mean: Dallol, Ethlopia C8 Hattast summers in W. Hamisphere: Death Valley, California, B3	Austra
	Long hot spell: Marble Bar, W. Australia D10	Europe
B. Low		' E. Variability o
	Warldy Vactab Autorctica FIO	Averag
	N. Hemisphere: Verkhovansk & Oimekon, U.S.S.R. All	Relatit
	Greenland: Northice A5	Relati
	N. America (excluding Greenland): Snag, Yukon Territory A2 Europe: Ust 'Shchugor, U.S.S.R. B8	F. Hailstones
	S. America: Sarmiento, Argentina E4	Larges
	Africa: Ifrane, Morocco B6	-
	Australia: Charlotte Pass, New South Wates Ell Monthly and annual means: Plateau Station, Antarctica F8	G. Great Snowfe
C. Varia	stions	24-ho
C1 / 4/14		19-10
	Difference between high and low: Verkhoyansk, U.S.S.R. All	One si
	Mean annual range: Eastern Sayan Region of U.S.S.R. B10	one o.
	Largest 2-minute rise in 11.5.: Spearlish, South Dakota B3	III. OTHER ELEMENT
	Instantaneous rise: Eanburgh, Scotlana Bb Largest 24-hour fall in 11.5.: Browning, Montana B3	A. Thunderston
. PRECIPITAT	TION	Avera
A Great	t Painfall Amounts	Avera
Al Orea		B. Sea-Level (
	One-minute, World: Unionville, Maryland B4	
	20-minute, World: Curtea-de-Arges, Romania B7	Highe
	42-minute, World: Holt, Missouri B3	Lowe
	12-hour, World: Belouve, Lu Reunion I. Do 24-hour, World: Cilgos, La Réunion I. DR	C. Salar Padia
	24-hour, N. Hemisphere: Paishih. Taiwan Cll	C. Solar Radia
	24-hour: Dharampuri, India C9	Avera
	24-hour, Australia: Crohamhurst, Queensland D12	One-
	5-day, World: Cilaos, La Réunion I. D8	
	One month, World: Cherrapunji, India C10	D. Wind Speed
	12-months, World; Cherrapunji, India C10	Deab
	12-months, 0.5 Mr. Halaicaic, Radai, Hadair Ci	Peak
B. Great	t Average Yearly Precipitation	Peak
		5-mi
	World: Mt. Waialeale, Kauai, Hawaii Cl	24 - h
	Asia: Cherrapunji, India C10	Mean
	Ajrica: Debunascna, Cameroon C/ S America: Quibdo Calambia C4	Days
	N. America: Henderson Lake, British Columbia B2	F. Dew Point
	Europe: Crhvice, Yugoslavia B7	
	Australia: Tully, Queensland D11	Mean
	Average number of days with rain per year: Bahia Felix, Chile E4	

XTREMES BY ELEMENT -- WORLD MAP

C. Least Precipitation

Number of years without rain. Arica, Chile D4

D. Low Average Yearly Precipitation

World: Arica, Chile D4 Africa: Wadi Halfa, Sudan C8 N. America: Bataques, Mexico B3 Asia: Aden, South Yemen C8 Australia: Mulka, South Australia D11 Europe: Astrakhan, U.S.S.R. B8

E. Variability of Precipitation

Average variability: Debundscha, Cameroon C7 Relitive variability: Themed, Israel B8 Relative variability (1935–38): Lhasa, Tibet C10

F. Hailstones

Largest, U.S.: Coffeyville, Kansas B3

G. Great Snowfall Amounts

24-hour, N. America: Silver Lake, Colorado B3 19-hour: Bessans, France B7 One storm, N. America: Mt. Shasta Ski Bowl, California B2 One season, N. America: Rainter Paradise Ranger Station, Washington B2

III. OTHER ELEMENTS

A. Thunderstorms

Average number of thunderstorm days per year: Kampala, Uganda C8 Average number of thunderstorm days, 1916–1919: Bogor, Indonesia D10

B. Sea-Level Air Pressure

Highest, World: Agata, U.S.S.R. A10 Lowest, World: estimated at 19° N, 135° E, C11

C. Solar Radiation

Average daily December: South Pole, Antarctica F6 & 7 One-hour: Malange, Angola D7

D. Wind Speed

Peak gust, World: Mt. Washington, New Hampshire B4 Peak gust, Thule Air Base, Greenland A4 Peak gust, Miyakojima I., Ryukyu Islands C11 5--minute, World: Mt. Washington, New Hampshire B4 24-hour mean: Port Martin, Antarctica 1911 Mean monthly: Port Martin, Antarctica 1911 Days with hurricane force winds: Port Martin, Antarctica 1911

E. Dew Point

Meen afternoon, in June: Assab, Ethiopia C8.



WEATHER EXTREMES IN NORTH AMERICA

市市村 市井市からい

From Weather Extremes Around the World, 1974

Environmental Science Services Administration, correspondence dated M. A. Arkin, Chief, Foreign Branch, Environmental Data Service, U.S. D. M. Ludlum, Weather Record Book, U.S. and Canada, Weatherwise, Data Service. Worldwide Extremes of Temperature, Precipitation and Data Service, Temperature Extremes, revised May 1967. (L.S. 5821.) Data Service, Temperature Extremes, revised May 1967. (L.S. 5821.) U.S. Environmental Science Services Administration, Environmental D. F., Pedgley, Air Temperatures at Dallol, Ethiopia, Meteorol. Mag., U.S. Environmental Science Services Administration, Environmental **P.S. Environmental Science Services Administration, Environmental** Pressure Recorded by Continental Area, October 1968. (FSSA) London, Vol. 96, No. 1142, pp. 265-71, 1967. Reference Princeton, New Jersey, 1971. Weather Records – With Documentation hy Element P1680032.) Ibid. Ibid. Ibid. Ibid. Ibid. Canada, 113°F, Midale and Yellow Grass, Saskatchewan, 5 July 1937 Western Hemisphere, 134°F, Death Valley, California, 10 July 1913 Antarctica, 58°F, Esperanza, Antarctic Peninsula, 20 October 1956 South America, 120°F, Rivadavia, Argentina, 11 December 1905 Dallol, Ethiopia, has a 94°F annual mean (possibly the world's Anstralia, 128°F, (Joncurry, Queensland, 16 January 1889 World, 136°F, El Azizia, Libya, 13 September 1922 Persian Gulf had a 96°F sea surface, 5 August 1924 U.S. highest annual mean, 77°F, Key West, Florida Element (to nearest whole number) Europe, 122°F, Seville, Spain, 4 August 1881 Asia, 120°F, Tirat Tsvi, Israel, 21 June 1942 A. Ilighest Temperature highest)

7

Reproduced from best available copy.

0

6 October 1969.

Weather Records – With Docume	ntation by Flcment (Cont'd)
Element (to nearest whole number)	Reference
U.S. warmest winters, 70°F, average, Key West, Florida	U.S. Environmental Science Services Administration, Environmental Data Service, Temperature Extremes, revised May 1967. (1S. 5821.)
Western Hemisphere hottest summers, <u>98°F</u> , avera ₅ e. Death Valley, California	Ibid.
Marhle Bar, West Australia, recorded temperatures of 100°F or above on 162 consecutive days, 30 October 1923 to 7 <u>April 19</u> 24	ll. T. Ashton and J. V. Maher, Australian Forecasting and Climate, 6th cd. Melbourne, Australia, 1960.
B. <u>Lowest</u> World, <u>127°F</u> , Vostok, Antarctica, 24 August 1960	U.S. Environmental Science Services Administration, Environmental Data Service, <i>Temperature Extremes</i> , revised May 1967. (L.S. 5821.)
Northern Hemisphere, <u>-90°F</u> , Verkhoyansk, U.S.S.R., 5 and 7 Fehruary 1892; and Oimekon, U.S.S.R., 6 Fehrucry 1933	Ibùl.
Greenland, 87°F. Northire, 9 January 1954	Ibid.
North America, excluding Greenland, <u>-81°F</u> , Snag, Yukon Territory, 3 February 1947	Ibid.
U.S., -80°F. Prospect Creek, Alaska, 23 January 1971	D. M. Ludlum, op. cit.
U.S., excluding Alaska70°F. Rogers Pass, Montana, 20 January 1954	U.S. Environmental Science Services Administration, Euvironmental Data Service, <i>Temperature Extremes</i> , revised May 1967. (L.S. 5821.)
Europe, <u>-67°F</u> , Ust 'Shehugor, U.S.S.R. Jan (date not known)	U.S. Environmental Science Services Administration, Environmental Data Service, Worldwide Extremes of Temperature, Precipitation and Pressure Recorded by Continental Area. October 1968 (ESSA/ P1680032).
South America, <u>-27°</u> F, Sarmiento, Argentina, I June 1907	Ibid.
Africa, -11°F, Ifrane, Morocco, 11 February 1935	Ibid.

P. C. Dalrymple, Geographic Applications Division, U.S. Army Engineer R. R. Hamann, The Remarkable Temperature Fluctuations in the Black Topographic Laboratories, Fort Belvoir, Va., Personal Communication. Data Service, Temperature Extremes, revised May 1967. (L.S. 5821.) Hills Region, January 1943. Monthly Weather Rev. Vol., 71, No. 3. Data Service, Temperature Extremes, revised May 1967. (L.S. 5821.) Gershevsky and ed. by J. E. Williams, W. II. Freeman, San Francisco, U.S. Environmental Science Services Administration, Environmental I. II. Chuter, A Sudden Rise of Temperature at Edinburgh. Weather, U.S. Environmental Science Services Administration, Environmental S. P. Suslov, Physical Geography of Asiatic Russia, transl. by N. D. Reference Vol. 21, No. 11, pp. 159-62, 1970. pp. 29-32, March 1943. R. R. Hamann, op. cit. 1961. Reid. Ibid. Ibid. Ibid. Fastern Savan Region of U.S.S.R. has a 146F° (93.2 to -53.2) annual U.S. largest 24-hour temperature fall, $100F^{\circ}$ (44 to -56), Browning, Edinburgh, Scotland, had an instantaneous temperature rise, 21 F^o U.S. largest 2-minute temperature rise, 49F° (4 to 45), Spearfish, Verkhoyansk, U.S.S.R. has a difference of 183F° (93.5 to -89.7) Mateau Station, Antarctica, had a monthly mean of -100°F, July Australia, -8°F, Charlotte Pass. New South Wales, 14 June 1945 Rapid City, South Dakota, had three temperature rises and two U.S. coldest winters, -16°F average, Barter Island, Alaska L.S. coolest summers, 37°F average, Barrow, Alaska between highest and lowest recorded temperatures Element (to nearest whole number) U.S. lowest annual mean, 10°F, Barrow, Alaska 1968, and an annual mean of -70°F, 1966-68 **Temperature Variations** South Dakota. 22 January 1943 Nontana, 23-24 January 1916 (45 to 66), 8 April 1969 mean temperature range and 22 July 1947 Ċ

9

Weather Records -- With Documentation by Element (Contⁱd)

falls of 40F° or over during a period of 3 hours, 10 minutes, 22 January 1943

November 1971. (NOAA Tech Memo WR-28 and Western Region Tech J. I., II. Paulhus. Indian Ocran and Taiwan Rainfalls Set New Records, munication from Director to P. A. Siple, Scientific Advisor to Director B. W. Newman, Australia's Highest Daily Rainfall, Australian Meteorol R. J. Schmidli, Weather Extremes, Salt Lake (Jty, Utah, U.S. National Indian Gentral Meteorological Research Institute, Poona, India. Com-J. G. Potter, Record Precipitation on One Day in Canada, CDS #1-68, Occanic and Atmospheric Administration, National Weather Service, Earth Sciences Laboratory, U.S. Army Natick Laboratories, in letter of Army Research, transmitted from Dr. Siple to Dr. W. B. Brierly, Monthly Weather Rev, Vol. 93, No. 5, pp. 331-35, 1965. Canada, Meteorological Branch, Toronto, 1968. Reference Mag, No. 20, pp. 61-65, March 1958. Weather Records - With Documentation by Element (withd) J. L. H. Paulhus, op. cit. dated 17 January 1962. Memo No. 28.) Ibid. Ibid. Ibid. Ibid. Ibid. World, 20-minute 8.10 incluss, Curtea-de-Arges, Romania, 7 July 1889 Workl. 24-hour, 74 inches, Glaos, La Rémion L. 15-16 March 1952 Dharampuri, India. had a 21-Hour rainfall of 39 inches, 2 July 1941 World, 12-hour, 53 incles, Belonve, La Réunion 1., 28-29 February Anstralia, 24-hour. 36 inches, Crohamhmst, Qucensland, 3 Febru-Vorthern Hemisphere, 24-hour, 49 inches, Paishih, Taiwan, 10-11 World, I-minute, 1.23 inches, Unionville, Maryland, 4 July 1956 Yankeetown, Florida, had a 24-hour rainfall of 39 inches. 5-6 World, 42-minute, 12 inches, Holt, Nissouri, 22 June 1947 Canada, 24-bour, 19 inches, Uchnelet. British Columbia, Element (to nearest whole number) **Greatest Rainfall** Precipitation September 1963 September 1950 6 Octoler 1967 ary 1893 بر 1961 _

10

World, 5-day, 152 inches, Cilaos, La Rénnion I., 13-18 March 1952

Uata Service, Worldwide Extremes of Temperature, Precipitation and U.S. Environmental Science Services Administration, Environmental Weather and Crop Bull, Vol. 43, No. 9, pp. 6-8, 27 February 1956. L. H. Seamon and G. S. Bartlett, Climatological Extremes, Weekly Pressure Recorded by Continental Area, October 1968. (ESSA/ Reference Weather Records – With Documentation by Element (Cont'd) R. J. Schmidli. op. ci¹. R. J. Schmidli, op. cit. P1680032.) Ibid. Ibid. Ibid. Ibid. Ibid. Ibid. Ibid. Ibid. Africa, 405 inclues during a 32-year period, Debundscha, Cameroon U.S., 12-month, <u>624 inches,</u> Mt. Wajaleale, Kanai, Hawaii, 24 July 1947 to 27 July 1948 Nustralia, 179 inches during a 31-year period, Tully, Queensland Europe. 183 inches during a 22-year period, Crkvice, Yugoslavia Workl, 460 inclues during a 32-y ear period, Mt. Waialeale, Kauai, World, 12-month, 1042 inches, Cherrapunji, India, August 1860 South America. 354 inches during a 10-16 year period, Quibdo, North America, 262 inches during a 14 year period, Henderson Arica, Chile, had no rain for more than 14 consecutive years, Bahia Felix. (Itile, has an at trage of 325 days/year with rain <u>Vsia. 450 inclees during a 74-year period</u>, (Therrapunji, India Workl, 1-month, 366 inches, Cherrapunji, India, July 1861 Greatest Average Yearly Precipitation Element (to nearest whole number) Least Precipitation Lake, British Columbia to July 1861 ن (iolombia ÷ llawaii

11

October 1903 through December 1917

 10. Lowest Average Yearly Precipitation 11. Rinches, Iberth Valty, California 12. Lowest Average Ave
--

Weather kecords – With Docum	cutation by Element ((ont'd)
Element (to nearest whole number)	Reference
Lhass, Tibet, had a 108% relative variability of annual precipitation, 1935-1938	A. Lu, A Brief Survey of the Climate of Lhasa, Quarterly J Roy Meteorol Soc, Vol. 65, No. 281, pp. 297-302, 1939.
 F. B. 3stones U.S. largest hailstone, 17.5 inches circumference, Coffeyville, Kansas, 3 September 1970 	The "New Champ" Hailstone, Weatherwise, Vol. 24, No. 4, p. 151, 1971,
(;. (irratest Snowfall	
North America, 24-hour, <u>76 inches,</u> Silver Lake, Colorado, 14-15 April 1921	K. J. Schmidt, op. cit.
Alaska, 24-hour, <u>62 inclu</u> s, Thompson Pass, 29 December 1955	Ibid.
Bessuns, France, had a snowfall of <u>68 inclues</u> in 19 hours, 5-6 April 1969	M. Jail, Un remarquable effet de lombarde: Acs chutes de neige de Paques 1969 en llaute-Maurienne, <i>Rerue de Géog</i> raphie Al <i>pine</i> , Grenoble, Vol. 57, No. 3, pp. 613-21, 1959.
Canada, observation day. <u>44 inches</u> , Livingston Ranger Station, Alberta. 29 June 1963	R. J. Schmidli, op. cit.
North America, one storm, 189 inches. Mt. Shasta Ski Bowl, Cali- fornia, 13-19 February 195 <u>9</u>	lhid.
Alaska, oue storm, <u>175 inches</u> Thompson Pass, 26-31 December 1955	I bid.
North America, oue season, <u>1122 inches</u> , Rainer Paradise Ranger Station, Washington, 1971-72	New U.S. Record Snowfall, Weatherwise, Vol. 25, No. 4, p. 173, 1972. 🖉
. Maska, one stason, <u>974 inches.</u> Thompson Pass, 1952-1953	R. J. Schmidli, op. cit.
Canada, one season, 880 <u>inches.</u> Kemano, Kildala Pass, British Columbia, 1956-1957	lbid.

which is the second of the second second

Element (to nearest whole number)	Reference
greatest depth of snow on the ground, <u>451 inches</u> , Tamarack, ornia, 11 March 1911	Ibid.
Other Elements	
A. <u>Thunderstorms</u> Ipala, Uganda. averages <u>242</u> thunderstorm days per year	World Meteorological Organization, World Distribution of Thunder- storm Days. (OMM, No. 21.) Geneva, Switzerland, 1953.
or, Indonesia, averaged <u>322</u> thunderstorm days per year, 1916- 9	V. I. Arabadzhi, Klimat i grozy [Climate and Thunderstorms], <i>Priroda</i> , No. 2, pp. 65-66, 1966. (Translated by E. R. Hope as Canada Defence Research Board Translation T456R, April 1966.)
B. Sea-Level Air Pressure	
ld. highest, 32.01 inches, Agata. U.S.S.R., 31 December 1968	Extremes of Atmospheric Pressure, Weatherwise, Vol. 24, No. 3, pp. 130-31, 1971.
th America, highest, 31.52 inches, Medicine Hat, Alberta, 24 arv 1897	R. J. Schmidli, op. cit.
ld, lowest. <u>25.90 inches</u> , estimated by aerial recontaissance in ey.: of Typhoon Ida at 19°N, 135°E, 24 September 1958	Ibid.
th Am rica, lowest, <u>26.35 inches,</u> Matecumbe Key, Florida, ptember 1935	Ibid.
C. Solar Radiation	
th I ale has 955 langleys average daily insolation in December	U.S. Fnvironmental Science Services Administration, Environmental Data Service, <i>Climatological Data for Antarctica Staticus</i> , Nos. 1 and 9 Washington, D.C., GPO, 1962 and 1968.
ange, Angola, had 113 langleys in 1 hour, 7 November 1961	Quart Radiation Bull, Union of South Africa, Vol. 8, No. 2, p. 160, 1961.

Weather Pecords – With Docur	nèntation by Element (('ont'd)
Element (to nearest whole number)	Reference
D. Wind Speed	
Workl, highest peak gvst, 231 miles per hour, Mt. Washington, New Hampshire, 12 April 1934	U.S. Environmental Science Services Administration, Environmental Data Service, Local Climatological Data; Annual Summary with Com- parative Data, 1967, Mount Washington Observatory, Gorham, New Hampshire, Washington, D.C., GPO, 1968.
Thule Air Base, Greenland, had a peak gust of <u>207 miles per hour</u> , 8 March 1972	 R. Stansfield, The Severe Arctic Storm of 8-9 March 1972 at Thule Air Force Base, Greenland, Weatherwise, Vol. 25, No. 5, pp. 228-33, 1972.
Niyakojima I., Ryukyu Islands, had a peak gust of <u>190 miles per</u> hour, 5 September 1966	Y. Mitsuta and S. Yoshizumi, Characteristics of the Second Miyakojima Typhoon, Kyoto, Japan. Univ. Disaster Prevention Research Institute. Bulketin, Vol. 18, pt. 1, No. 131, pp. 15-34, 1968.
World, highest 5-minutes, <u>188 miles per hour</u> , Mt. Washington, New Hampshire, 12 April 1934	R. J. Schmidli, op. cit.
Port Martin, Antarctica, had a 24-hour mean of 108 miles per hour, 21-22 March 1951 and a mean monthly of <u>65 miles per hour</u> , March 1951.	F. Loewe. The Land of Storms, <i>Weather</i> , Vol. 27, No. 3, pp. 110-121, 1972.
U.S. highest annual mean, <u>35 miles per hour.</u> Mt. Washington, New Hampshire	M. A. Arkin, op. cit.
E. Dew Point (Humidity)	
Assab, Ethiopia has 81°F mean afternoon dew point in June	A. V. Dodd, Areal and Temporal Occurrence of High Dew Points and Associated Temperatures, Tecn Rpt ES-49, U.S. Army Natick Labora- tories, Natick, Massachusetts, August 1969.
F. Fog Frequency	
U.S. West Coast highest average, <u>2552 hours</u> per year, Cape Disappointment, Washington	M. A. Arkin, op. cit.
U.S. East Coast highest average, <u>1580 hours</u> per year, Moose Peak Lighthouse, Mistake Island, Maine	M. A. Arkin, op. cit.

15

15

II. COMMENTARY ON RELIABILITY OF THE RECORDS

To insure comparability of meteorological observations, various regulations concerning site, instrumentation, and procedure have been established by the World Meteorological Organization.² Observations taken in accordance with these regulations, over which some sort of quality control is exercised to correct observational errors that might appear, would be accepted by the national meteorological agencies.³ Also, for highest or lowest average records, the period of observation should be long enough to be representative. However, even after records have been accepted by appropriate agencies, their reliability is still sometimes questioned.

An extreme phenomenon observed and recorded in conformance with prescribed procedures and accepted by the appropriate meteorological service represents only the most extreme acceptable record that is available. It is not necessarily – or even probably – the most extreme value that could occur or ever has occurred. According to M. A. Arkin, "... record extremes must be taken with a grain of salt...."⁴ He explains that news of an extreme weather occurrence is not always widely disseminated, meteorological records are relatively short, stations are very few, and "... even the densest network of stations provide: only a very small sample of the weather" To this it can be added that records may be established which the observers do not recognize as records, and sometimes suspected records are not verified because of difficulties in validating.

3. Temperature. Air temperatures are measured at standard heights varying in different countries from 4 to 6 feet above ground level. The values obtained can be affected by radiation from the sun, sky, earth, and other surrounding objects, and precautions need to be taken to protect the measuring instruments from radiation. In addition, adequate ventilation must be provided to insure representation of the circulating air. In addition to these requirements concerning height, radiation protection, and ventilation, there are requirements pertinent to construction and accuracy (manufacturing tolerance) of the measuring instruments.

Daily maximum and minimum temperatures are recorded at more than 10.000 weather stations throughout the world.⁵ Maximum and minimum thermometers which stay at the highest or lowest point, respectively, reached during the reading period are used to record these temperatures. Maximum thermometers are generally of the mercury-in-glass type. But since mercury freezes at about -32°F, minimum thermometers contain

Preceding page blank

⁴ World Meleorological Organization, "Guide to Meleorological Instrument and Observing Practices," 2nd ed., Geneva, 1961. (WMO-No. 8, TP, 3,)

³ M. A. Arkin, *op. cit.*

⁴ M. A. Arkin, op. cit.

⁵ G. Hoffman, "Die höchsten auf die tiefsten Temperaturen auf der Erde," (The Highest auf Lowest Temperatures of the Earth). Umschau, Frankfurt am Main, Vol. 63, No. 1, pp. 16-18, 1 Jan 63.

other liquids such as ethyl alcohol; these are commonly called spirit thermometers because of the type of liquids used.

Besides the absolute maximum and minimum temperatures ever recorded in an area, several other kinds of extreme temperature records can be of interest. Among these are the highest and lowest mean daily, monthly, or yearly temperatures; the highest and lowest mean daily, monthly, or yearly maximum and minimum temperatures; the longest durations of very high or very low temperatures; the greatest variations in a given period (e.g., between summer and winter, day and night); and the fastest rises and falls in temperature during short periods of time.

a. High Temperatures. Factors favorable to occurrence of very high temperatures have been listed by the British meteorologist H. H. Lamb.⁶ Among these factors are strong heating of the surface, especially desert sand or bare rock, during times of high sun and very clear atmosphere; a long period of time during which the air has passed or remained over extremely warm surface; inhibition of vertice¹ convection or local eirculations by subsidence; passage of the air over mountains, especially when latent heat absorbed during condensation or rainfall on the ascent is released into the descending air; and advection from places where the air was already heated.

Lamb's investigations suggest that conjunction of most or all of these influences might occur during record high extremes, and that many of the peak temperatures "... are associated with the speeding up, and longer fetch, of warm air advection just ahead of a cold front" Further, he concludes that these extreme high temperatures "... may have been very local and associated with some locally forced turbulence – for instance, in air descending some sort of declivity or merely passing over town buildings – the circumstances being such as to caise adiabatically by a degree or two the temperature of already very warm air at a slightly higher 'evel "⁷

The highest possible temperature that could ocenr has been considered by the German meteorologist Hoffman.⁸ According to itim, because warm air is lighter, air near the hot ground surface seeks to rise above overlying layers which are cooler. Thus, an ascending air stream is established through which the hot air escapes so that the temperature of the lower air levels should not rise above a theoretically definable value. He puts this value at slightly over 55°C (131°F) and states that world weather observations confirm it. The bighest accepted record is 58°C (136°F), at El Azizia in North Africa, but elaims have been made of higher temperatures; e.g., 140°F at two

lbid.

⁶H. H. Lamb, "The Occurrence of Very High Surface Temperatures," *Meteorol. Mag.*, London, Vol. 87, No. 1028, _pp, 39-43, 1958.

⁸G. Hoffman, op. cit.

Mexican stations, Delta, Baja California and Riito, Sonora⁹ and 167°F in the Gobi Desert of interior Asia.10

vrly maximum temperatures, 122°F or above, The highest mean occur in the northern and wes' a contart ara; Death Valley, California; low-lying desert areas in Iran; and a small part f western Pakistan.¹¹ By contrast, the highest yearly maximum temperatures in some parts of Antarctica average below 0°F.

> World's highest temperature: 136°F El Azizia, Libya, 13 September 1922

El Azizia is located in the northern Sahara at 32°32' N, 13°01' E, elevation 367 feet.¹² At least 30 years of observations are available for the station,¹³ and the elimate has been described by Eredia.¹⁴ The world record temperature occurred during the month of September; other months during which very high maximum temperatures have occurred are August and June, with 133°F and 127°F, respectively.¹⁵

Although this record of 58°C has attained general acceptance as the world's highest temperature recorded under standard conditions,¹⁶.¹⁷ it has been questioned. A. Fantoli, who was Director of the Libyan Meteorological Service at the time the observation was reported, has written about the subject, 18 19 and summaries in English are available.²⁰²¹ Although he was unable to personally investigate the reliability of the reading at the time it occurred, he has since examined the evidence in some detail: the exposure; effect of instrument shelter or of instrumental error; records for pressure, wind, and humidity at three other stations in Tripoli for 9-12 September 1922; and synoptic charts for the area for 10-14 September 1922. He also

International Boundary and Water Commission, United States and Mexico, "Flow of the Colorado River and other Western Boundary Streams and Related Data," p. 58, Western Water Bulletin, 1964. 10

H. T. Ashton and J. V. Maher, "Australian Forecasting and Climate," 6th ed., Melbourne, Australia, 1960. II G. lloffmann, op. cit.

¹² Great Britain, Air Ministry, Metrorological Office, "Tables of Temperature, Relative Humidity and Precipitation for the World, Part IV, Africa, the Atlantic Ocean South of 35° N and the Indian Ocean," (M.O. 617d). London Her Majesty's Stationery Office, 1958. 13

Ibid.

F. Eredia, "Klima von Azizia (Climate of Azizia," Meteorol Zeitschrift, Itraunschweig, Vol. 42, p. 294, 1925.

¹⁵ Great Britain, Air Ministry, Meteorological Office, op. cit.

¹⁶ U. S. Environmental Science Services Administration, Environmental Data Service, *Temperature Extremes*, revised May 67 (L.S. 5821).

L. II, Seamon, and G. S. Itartlett, "Climatological Extremes," Weekly Weather and Crop Bulletin, Vol. 43, No. 9, pp 6-8, 27 Feb 56,

A. Fántoli, "La Piu Alta Temperatura del Mondo." (The Highest Temperature in the World). Rev Meteorol Aeronautica, Rome, Vol. 18, No. 3, pp 53-63, Jul-Sep 58. (English Simmary)

A. Fantoli, "I Valori Medi Della Temperatura in Libia," Boll. Sov. Geogr. Italiana, ser. 8, Vol. 7; pp 59-71, 1954. 20

J. Gentilli, "Libyan Climate," Geograph Rev, Vol. 45, No. 2, p 269. Apr 55.

Meteorol Abstracts, 10 (10), 1959. (Contains abstract of #18.)

reviewed temperature records at El Azizia for June to October 1914 1922, as well as the maximum temperatures (99°F to 118°F) at 10 places in Tripoli on 11-14 September 1922. From his investigations, Fántoli concluded that although there was an unusually violent and persistent ghibli²² at the firse, the probable or tentative maximum on 13 September 1922 was 56°C (132.8°F). This conclusion may have been influenced in part by the comparatively low maximum temperatures at nearby stations, a point also mentioned by other authors.²³ Lamb also investigated the El Azizia record, and from his investigations he has described the synoptic situation which caused the extreme heat.24 A cold front was advancing castward from Algeria, and advection of waren air from the Saharan interior was established some distance ahead of the front. Tripoli had strong southerly winds for two successive days before the front passed. Rainfall in mountains to the south in the Anderas area (17° N, 8° E) was thought to have contributed latent heat of condensation to the air mass.

Western Hemisphere's highest temperature: 134°F Death Valley, California, 10 July 1913

Death Valley, at 37° N, 117° W, is a desert below sea level, flanked by mountains. It has the hottest summers in the Western Hemisphere.²⁵ The temperature of 134°F, recorded : Greenland Ranch Station, is accepted by the U.S. Weather Burean as an official record. On the day this temperature occurred, winds were strong, there were sandstorm conditions, and a heat spell of 8 consecutive days with maximums of 127°F or higher was in progress.²⁶

A. Conrt²⁷ has examined the Death Valley record in detail: the surrounding environment, the equipment and procedures used for taking the observation, weather conditions at the time it occurred, and temperature frequencies at the station over a period of record from 1911 through 1947. He found that at the time of the record there were comparatively low maxima at other stations in the area, as had been the case at El Azizia. He found that aside from the period during which the record occurred in July 1913, temperatures of 127°F or higher were reached in only two other years from 1911 through 1947. During a later period, 1948 to 1967. a temperature of 129°F was

A ghiblt is a hot, dust-bearing wind in Tripolitania.

 ²³G. Hellmann, "Grenzwerte der Klimaclemente auf der Erde." [Limits of Climatic Elements of the Earth]. Sitzungsberichte Preuss. Akademie der Wissenschaft, Phys. mathem, Klasse, 1925, pp. 200-215, Uited by Mark Jefferson: "Limiting Values of Temperature and Rainfall Over the World," Geograph Rev. Vol. 16: pp. 324-26. 24 1926.

H. H. L. mb, op. cit.

²⁵ U. S. Environmental Science Services Administration, Temperature Extremes, op. cit.

²⁶ L. Williams, "A Contribution to the Philosophy of Climatic Design Limits for Army Materiel: Extreme Hot-Desert Conditions," Technical Report ETL/TR-72-5, H. S. Army Engineer Topographic Laboratories, Fort Belvoir, Va., 27 1972.

A. Court, "How Hot is Death Valley?" Geograph Rev. Vol. 39, No. 2, pp 214-20, Apr 49,

reported in July 1960²⁸; but according to Williams this value is doubtful.²⁹ From his analysis of the 1911 through 1947 data, Court determined that a temperature of 134°F has an expectancy of only once in 650 years.³⁰ He concluded that "constantly increasing accuracy in weather observations and higher standards of instrument exposure make it seem probable that no future official observation will exceed the present high temperature record for North America new held by Death Valley."31

Persian Gulf had a sea-surface temperature of 96°F 5 Angust 1924.

This sea surface temperature, measured by the SS Frankenfels, is "... among the highest recorded...."³² However, sea temperature in the Persian Gulf has been known to reach 98°F in August and September.³² Although the 96°F record is not an absolute maximum in the sense of being the highest known, it was suggested for inclusion here to indicate how hot the rea surface can become. The Persian Gulf also has very high average temperature ¹ summer, 88°F during July and August.³⁴

> Dallol, Ethiopia, has an annual mean temperature of 94°F. possibly the world's highest.

Places that are hot in summer and remain warm in winter have high annual means of daily maximum temperature. The highest occur at low elevations, away from coasts, and within the latitude belt between 12° and 20°N across Africa and possibly the southwestern Arabian penisula.³⁵ Dallol is within this belt at 14°19′ N. 40°11'E. It is located 258 feet below sea level on the edge of the Danakil Depression, a salt desert. By averaging the annual mean daily maximum temperature of 106°F³⁶ and the annual mean daily minimum of 83°F³⁷ an annual mean temperature of 94°F is obtained. All of these values are possible world records. The 94°F annual mean

²⁸ P. I. Tattelman, N. Sissenwine and R. W. Lenhard, "World Frequency of High Temperature," Environmental Research Papers, No. 305, Air Force Cambridge Research Laboratories, Bedford, Mass., 1969.

L. Williams, op. cit.

³⁰ A. Court, op. cit.

³¹ Ibid.

³² U. S. Environmental Science Services Administration, Temperature Extremes, op. cit.

Great Britain, Air Ministry, Meteorological Office, "Weather in the Indian Ocean to Latitude 30°S and Longitude 95°E including the Red Sea and Persian Gulf, ' volume II "Local Information," part 3, "The Persian Gulf and Gulf of Oman including the part of the Makran Coast west of Gwadar," (M. O. 451b), Ilis Majesty's Stationery Office, London, 1941. 34

U. S. Environmental Science Services Administration, Temperature Extremes, op. cit.

³⁵ D. E. Pedgley, "Air Temperatures at Dallol, Ethiopia." *Meteorol Mag.* London, Vol. 96, No. 1142, pp. 265-71, 36 Sep 67. 36 Ibid.

³⁷ Ihid.

exceeds the 88°F averaged at Lugh Ferrandi, Somalia from 1923 to 1935 and cited as probably the world's highest annual mean temperature.³⁸ The 106°F annual mean daily maximum temperature exceeds the 102°F recorded at Abecher, Chad for a 6-year period and the 101°F at Merowe, Sudan (30 years) and Araonne, Mali (8 years): these values are the highest listed in the British Meteorological Office's "Tables of Temperature, Relative Humidity and Precipitation for the World."³⁹

The Dallol temperatures were obtained from readings taken at a climatological station maintained at the base camp of an American prospecting company over a period of 6 years, October 1960 through November 1966. Maximum and minimum temperatures were recorded ". — using standard thermometers kept at a height of four feet in a ventilated screen "⁴⁰ The period of observation is short; but it is believed that the difference between these available 6-year values and a long-term mean would be very small. At nearby Khormaksar, whose temperature trends appear similar to Dallol's, the greatest difference between monthly means of daily maximum temperature for the years 1961 through 1966 and those for 1947 through 1966 was only 0.3°C.⁴¹

D. E. Pedgley has considered areas where annual mean daily maxima might equal or exceed Dallol's.⁴² According to him, in the lowest part of the Danakil Depression, which reaches a minimum of about 390 feet below sea level some 20 miles south of Dallol, the annual mean daily maximum might be a "fraction of a degree Fahrenheit greater" than at Dallol.⁴³ In the Abecher area where there is a "... descent of potentially warmer air resulting from a blocking by the Marra Mountains of the north-easterlies that blow for much of the year" there might also be low-lying places with values similar to Dallol's.

³⁸ U. S. Environmental Science Services Administration, *Temperature Extremes*, op. cit.

Great Britain, Air Ministry, Meteorological Office, op. cit.

D. E. Pedgley, op. cit.

⁴² *[bid.*

⁴² Ibid.

[&]quot; Ibid.

^{*} R. A. McCormiek, "An Estimate of the Minimum Possible Surface Temperature at the South Pole," *Monthly Weather Rev.*, Vol. 86, No. 1, pp. 1-5, Jan 50.

Extremely cold temperatures ocenr in interior high latitude localities with clear skies, which are conducive to maximum terrestrial radiation, and with topographic features which afford protection from wind. Geographic areas of extreme cold are the eastern Antarctic plateau (approximately 9,000 to 12,000 feet of in elevation): the central part of the Greenland leccap (approximately 8,200 to 9,800 feet in elevation); Siberia between 63° and 68°N and between 93° and 160°E (below 2,500 feet in clevation): and the Ynkon basin of northwestern Canada and Alaska (below 2,500 feet in elevation).

In Europe, the lowest temperature has been recorded at Ust'Shchngor in the Soviet Union at about 55°N, 50°E.⁴⁵ This station, located near the Siberian border at an elevation of 279 fect, reported a -67°F in January; the exact date is not eited, but it was the lowest in a 15-year period.⁴⁶

Annual mean minimum temperature can range from approximately 70°F in places like Dallol, Ethiopia, to a possible -130°F in the Antarctic. This difference of about 200 Fahrenheit degrees is considerably greater than the difference of about 125 Fahrenheit degrees between the highest and lowest annual mean maximum temperatures. Annual mean minimums for the world's coldest areas are -76°F in Siberia, below -85°F on the Greenland Leecap, and from about -94°F to -130°F in the Antarctic.⁴⁷ The -130°F value is considered possible as an annual mean minimum, even though -127°F is the lowest temperature recorded, because in the extensive cold area of East Antarccica only a few stations have taken temperature measurements and then only for short periods of time.⁴⁸

World's lowest temperature: -127°F Vostok, Antarctica, 24 August 1960

The theoretical minimum temperature that could be reached has been calculated by Shliakhov⁴⁹ and by McCormick.⁵⁰ Shliakhov estimated -80°C ±2°C at about 4 kilometers altitude (i.e., -112°F at about 13,123 feet) with a decrease of 0.5°C for every further 100-meter rise in height. This estimate was exceeded when the temperature fell to -87.4°C (-125°F) at Vostok. Antarctica (78°27'S, 106°52'E, elevation 11,220 feet) in 1958 and again in 1960 when the world record low temperature of

 ⁴⁵ U.S. Environmental Science Services Administration, Environmental Data Service, Worldwide Extremes of Temperature, Precipitation and Pressure Recorded by Continental Area, Oct 68 (ESSA/P1h80032).

⁴⁷ Ibid.

G. Hoffmann, op. cit.

⁴⁹ Ibid.

^{**} V. L. Shliakhov, "D. Minimal'nykh Temperarurakh v Antarktide [Oo Minimum Temperatures in Autarctica]," ____Meteorologija i Gidrologija, No. 4, pp. 5-7, 1958.

R. A. McCormick, op. cit.

-126.9°F occurred, also at Vostok. The basic error in Shliakhov's calculations, according to H. Wexler⁵¹ resulted from associating a zero radiation balance with a certain temperature – the theoretical minimum. McConnick's calculation for the possible minimum was for "..., virtually optimum conditions ... assumed to persist during the polar night (about 180 days) "⁵² The resultant value, -200°C (-328°F) is "... hardly realistic for the troposphere ...," where, apparently, wintertime radiative energy losses are compensated by advective gains. However, it might be more applicable to the ozonosphere where such "... energy-balancing processes are not equally operative ...," ⁵³ An ozonospheric temperature of -97°C (-143°F) was measured at about 24,000 meters (78,740 feet) above Halley Bay at 75°31'S, 26°36'W, altitude 98 feet, on 9 August 1959.⁵⁴ The lowest temperature recorded in the Earth's atmosphere is -153°C at 93 kilometers above Point Barrow, Alaska, in June 1966.⁵⁵ However, this is above the ozonosphere and June is not the time of polar night at Point Barrow.

Northern Hemisphere's Lowest Temperature: -90°F Verkhoyansk, USSR, 5 and 7 February 1892 and Oimekon, USSR, 6 February 1933

Very low winter temperatures occur in the Verkhoyansk-Oimekon cold zone, approximately between 63° and 68°N, and 93° and 160°E in the East Siberian taiga (northern coniferous forest). It is an area of extreme continentality, lying uear the eastern end of the world's largest landmass and blocked off by mountain ranges from the moderating influence of oceans. In winter, the high air pressures from the Asiatic anticyclone create clear weather which promotes strong radiation from the snow surface during the long nights. A value of -89.7°F was recorded at Verkhoyansk (67°34'N, 133°51'E, elevation about 350 feet) twice in 1892, and -89.9°F was recorded at Oimekon (63°28'N, 142°49'E, elevation 2,625 feet) in 1933.⁵⁶ Oimekon is situated in a valley 10 to 20 kilometers (about 3,280 feet) wide, below a platean which is, in turn, enclosed by mountains.⁵⁷ In the winter, cold air accumulates in the valley and there is hardly any wind. It is possible that the temperature there may have fallen

²¹ II. Wexler, "Note on Lowest Antarctic Temperature Estimated by Shliakhov," *Monthly Weather Review*, Vol. 87, 200, 40, 147, Apr 59,

⁵³ R. A. McCormick, op. cit.

⁵⁴ Ibid.

D. J. George, "Coldest Place on Earth," Weather, Vol. 16, No. 5, pp. 144-59, 1961.

⁵⁵ II. Faust, "Die niedrigsten Temperaturen in der Erdatmosphäre [The lowest temperatures in the Easth's atmosphere]," Weltraumfahrt, Frankfurt am Main, Vol. 20, No. 1/2, pp. 25-27, 1969.

 ⁵⁹ U.S. Environmental Science Services Administration, Temperature Extremes, op. cit.
 57

²⁷ A. A. Borisov, "Climates of the U.S.S.R." translated by R. A. Ledward and edited by C. A. Halstearl, Chicago, Aldine, 1965.

below the value recorded; temperatures of .95°F and down to .108°F have been claimed.^{58 59 60} Considerable controversy has arisen about the Verkhoyansk records due to problems concerning instrument corrections, and about both the Verkhoyansk and Oimekon records because of misleading references to incorrect values in the literature.^{61 62 63}

Greenland's lowest temperature: -87°F Northice, 9 January 1954

On the permanent icecap, which covers most of the interior of Greenland, temperatures are very low due both to loss of heat through radiation and to evaporative cooling from the snow and ice surface.⁶⁴ A temperature of -86.8°F has been recerded at Northice (78°04'N, 38°29'W, elevation 7,687 feet), a station established by the British North Greenland Expedition.⁶⁵ Since the period of record for Northice was only 20 months (November 1952 through June 1954) and temperatures below -75°F occurred 16 times, it is quite probable that temperatures there have been lower than -87°F at other times.⁶⁶

North America's lowest temperature (excluding Greenland): -81°F Snag, Yukon Territory 3 February 1947

Snag is located near the Alaskan border of Canada's Ynkon Territory, at 62°23'N, 140°23'W, elevation 2,120 feet. This part of Canada resembles the Verkhoyansk-Oimekon Area of eastern Siberia in its continentality. The lowest gradation on the minimum thermometer which recorded this extreme was -80°F, but a pencil mark was made at a distance about 4 degrees below -80°F.⁶⁷ However, subsequent

 ⁵⁸ N. A. Stepanova, "On the Lowest Temperatures on Earth," *Monthly Weather Review*, Vol. 86, No. 1, pp. 6-10, Jap 58, 000
 ⁵⁹ Jap 58, 000

L. II. Seamon and G. S. Bartlett, op. eit.

⁶⁰ E. A. Finn, Kak Byl Otkryt "Polius Kholoda" [How was the "Unld Pole" Discovered?]. Priroda, No. 7. pp. 85-88, Moscow, Jul 67.

N. A. Stepanniva, op. eit.

⁶² E. S. Rubinshtein, "O Prirode Poliusov Kholoda," [On the Nature of the Cold Poles]. Vsesoiuznoe Geografieheskoe Obshehestvo Izvestija, Vol. 91, No. 3, pp. 265-68, May-Jun 59.

⁶³ E. S. Rulinshtein, "K. Voprosu & Poliusakh Kholoda," [Lontribution to the Problems of the Earth's Cold Poles]. Meteorologiia i Gidrologiia, No. 12, pp. 28-30, Dec 1968, Translated by U.S. Weather Bureau, Washington, D.G., May 59,

⁶⁴ D. W. Hogne, "Environment of the Greenland Icecap," Tech Rpt ES-14, F.S. Army Natick Laboratories, Natick, Mass., Dec 64, 65

²⁵ R. A. Hamilton and G. Rollitt, "British North Greenland Expedition 1952:54: climatological tables for the site of the expedition's base at Britannia Sø(Lake) and the station on the inland-ice Nortbice." København, U. A. Reitzels Forlag, Medd Grønland, Vol. 158, pp. 4-83, 1957.

⁶⁶ R. S. Quiroz, "Lowest Temperature in Greenland," Monthly Weather Review, Vol. 86, No. 3, p. 99, Mar 58,

⁶⁷ A. Court and H. A. Salmela, "Improbable Weather Extremes and Measurement Needs," *Amer Meteorol Soc Bull*, Vol. 44, No. 9, pp. 571-75, Sep 63.

laboratory calibration of the thermometer indicated an instrumental error of ±3 degrees: and a value of -81.4°F was officially set by the Canadian Meteorological Service.68 69 70 On the preceding day, 2 February, the corrected minimum temperature was -80.1° F.⁷¹ The previous record Canadian low temperature was -78.5°F at Fort Good Hope in the Northwest territorics on 30 December 1910.72

U. S. lowest temperature: -80°F Prospect Creek, Alaska, 23 January 1971

This low temperature was recorded at a camp along the Alaskan nipeline in the Endicott Mountains southeast of Bettles at 66°48'N, 150°40'W, 1,100 feet elevation. An official figure of -79.8°F was established "... after subsequent recalibration of the thermometer at the Bureau of Standards in Washington "⁷³ It replaced the previous official United States low of .76°F recorded at Tanana, Alaska, in the Ynkon Valley at 65°10'N, 152°06'W, elevation 220 feet, in January 1886.⁷⁴ Among other low temperatures claimed for Alaska was one of .78°F recorded by an airways observer at Fort Yukon (66°34'N, 145°18'W, elevation 417 feet) on 14 January 1934 two days after the weather station closed there, which made the record unofficial.⁷⁵ Also, a minimum thermometer left at 15,000 feet on Mt. McKinley for 19 years indicated a temperature lower than -100°F at some time during its exposure.⁷⁶

> Platean Station, Antarctica, had a mean July temperature of -100°F in 1968 and a mean temperature of .70°F for 1966 through 1968

Platean Station is located at 79°15'S, 40°30'E, elevation 11,890 feet. It is in or near the coldest part of the Antarctic, which is believed to be close to the ridge line in East Autarctica.⁷⁷ During the winter of 1968, there were 118 days with temperatures below -100° F.⁷⁸ The July mean temperature for that year is believed to be

71 Ibid. 72

⁶⁷ A. Court and H. A. Salmela, "Improbable Weather Extremes and Measurement Needs," Amer Meteorol Soc Bull, Vol. 44, No. 9, pp. 571-75, Sep 63. 68 Ibid.

⁶⁹

N. A. Slepanova, op. cit.

⁷⁰ A. Thomson, "Lowest Temperature in Canada," Monihly Weather Rev. Vol. 116, No. 11, p. 2911, Aug 58,

N. A. Stepanova, op. cit.

Weatherwise, Apr 71, Vol. 24, No. 2, p. 94, 74

U. S. Environmental Science Services Administration, Temperature Extremes, op. cit. 75

L. S. Seamon and G. S. Bartlett, op. cit.

⁷⁶ Ibid. 77

^{11.} H. Lamb, "Differences in the Meteorology of the Northern and Southern Polar Regious," Meteorol Mag. Vol. 87, No. 1038, pp. 353-79, London, Dec 58.

⁷⁸ F. S. Army Natick Laboratories," Annual Record of Major Events 5 FY 1969," Natick, Mass., 1969.

a new world record, -99.8°F.⁷⁹ It exceeds a mean of -97.2°F (-71.8°C) during August 1958 at Sovietskaya⁸⁰ which has been quoted as "... probably the lowest mean monthly temperature on record...."⁸¹ Sovietskaya, a Russian station, at 78°24'S, 87°35'E, elevation 11,713 feet, reported an annual mean temperature of -71°F during the IGY period in 1957 and 1958.⁸² This is colder than Platean Station's -70°°, but the area is not considered to be as cold. The periods of record at Plateau Station and Sovietskaya are not long enough, however, to be conclusive. Some other low average temperatures are -67°F at Vostok during 1958 and 1959, and -59°F at Amundsen-Scott Station (90°S, elevation 9,186 feet) for the period 1957-1964.⁸³ During this same period at Amundsen-Scott, the July temperature averaged -74.5°F with the maximums averaging -69°F and the minimums, -80°F.⁸⁴

c. Temperature Variations. Rapid temperature changes can occur under certain weather conditions; e.g., changes in wind direction whether onshore or offshore in coastal areas especially during seasons of the year when land-sea temperature differences are greatest. Foehns (warm, drying winds descending the lee sides of mountain ranges) can bring substantial temperature rises in short periods of time; increases of 20 to 30 degrees in an hour are not uncommon along the eastern side of the Rocky Mountains in the United States and Canada.⁸⁵ Advection of cold air masses can cause temperature falls which can be further intensified by loss of heat through radiation. Still other weather situations can cause extreme fluctuations in temperature.

> U. S. Greatest 2-minute temperature rise: 49F°, from -4°F to 45°F Spearfish, Sonth Dakota, 22 January 1943

Rapid City, Sonth Dakota, had three temperature rises and two falls of 40F° or more during a period of 3 hours and 10 minutes, 22 January 1943

Spearfish is located at 44°30′N, 104°W, elevation 3,637 feet, and Rapid City is at 44°N, 103°W, elevation 3,234 feet. They are in the Black Hills, a dome-shaped mass colminating in peaks over 7,200 feet above sea level, which slopes abruptly on the

⁷⁹ P. C. Dalrymple, Geographic Applications Division, U. S. Army Engineer Topographic Laboratories, Fort Belvoir, Va. Personal Communication.

⁸⁰ J. Alt. Quelques considérations générales sur la météorologie de l'Antarrtique, La Météorologie, No. 57, pp. 17-42, 1960.

D. J. George, op. cit.

⁸² N. A. Stepanova, op. cit.

⁸³ U. S. Environmental Science Services Administration, Temperature Extremes, op. cit.

^{84 [}bid.

⁸⁵ D. M. Ludhum, Weather Record Book, United States and Canada. Princeton, N.J., Weatherwise, 1971.

east and gradually on the west. This region lies mostly between 43° and 45°N, and 103° and 104°30'W, and the effects of the topography on winds from the west cause rapid temperature changes to occur there rather frequently.

On 22 January 1943, very rapid and pronounced fluctuations in temperature took place in the Black Ilills. The phenomenon, investigated by Hamann, was "... essentially the result of the wavering motion of a pronounced quasistationary front separating Continental Arctic air from Maritime Polar air"⁸⁶ Local chinook⁸⁷ effects possibly contributed to the unusual conditions. At Spearfish, the temperature rose from -4°F to 45°F and returned to -4°F between 0732 and 0927 hours (7:32 and 9:27 a.m.)⁸⁸ During this period, in which there were many abrupt changes, the record 2-minute rise occurred. At Rapid City, the temperature rose from 5°F to 54°F between 0920 and 0940 hours (9:20 and 9:40 a.m.), fell to 11°F at 1030 hours, rose to 55°F at 1045 hours, fell to 10°F at 1130 hours, rose to 34°F at 1150 hours, fell to 16°F at 1215 hours and rose to 56°F at 1240 hours.⁸⁹ Changes were so rapid that buildings were experiencing winter on one side and spring around the corner. The phenomenon also caused unusually large daily ranges in temperature, over 50 degrees at some places, and sharp contrasts between some nearby places. For example, at Lead, South Dakota, the temperature was 52°F, while at Deadwood, less than 3 miles away, it was -16°F.⁹⁰

In Canada, similar conditions have occurred. At Pineher Creek, Alberta, on 6 January 1966, the temperature rose 45F° (from -12 to 33) and fell 40F° (from 33 to -7) during a 2-hour period.⁹¹ Several hours later it again rose 46F° (from -10 to 36) within an hour.

4. Precipitation. Precipitation is measured by the depth to which it covers a horizontal unit area during a given period. Precipitation is caught in gages whose diameter represents the horizontal unit area; the more representative the catch is of actual fall over the entire observation area, the more useful the measurement. For this reason, site, form, and exposure of the gage are important, and precautions need to be taken to prevent precipitation from splashing out of the gage or being blown ont by wind. In hot, dry areas, evaporation can be a problem. Various requirements to cover these points have been established by the World Meteorological Organization.⁹²

⁸⁶ R. R. Hamann, The Remarkable Temperature Fluctuations in the Black Hills Region, January 1943, Monthly Weather Review, Vol. 71, No. 3, pp. 29-32.

A name given to foehns in the Western United States and Canada.
 88 manual states and Canada.

⁸⁸ R. R. Hamann, op. cit.

⁸⁹ Ibid. 90

⁹⁰ *Ibid*. 91

D. M. Ludlum, op. cit.

World Meteorological Organization, op. cit.

Gages are of two main kinds – the ordinary or non-recording gages and recording gages. The former provide a means of collecting and measuring precipitation, and the latter incorporate mechanisms for recording the amount of fall during a given period or the rate of fall at any instant. Descriptions of the many varieties of recording and nonrecording gages can be found in Middleton and Spilhaus, Meteorological Instruments.⁹³

Even with the most efficient instruments, functioning perfectly and in the most favorable sites and exposures, there are still problems in obtaining representative precipitation values. The standard gage diameter is 8 inches in the United States. Therefore, the horizontal unit area covered by the measured precipitation is about 50 square inches, and measuring sites are often several miles apart. Furthermore, intense rainfall is often very localized and may be missed by the observation network. An example is described by Lautzenheiser and Fay.⁹⁴ During a rainfall in August 1959, two gages in the town of Island Falls, Mainc, neither in the official Weather Bureau network, each measured 6.35 inches. The next nearest gage, about 12 miles away, measured 3.06 inches. The nearest official network gages, at Houlton approximately 20 to 25 miles northeast of Island Falls, measured 0.59 and 0.25 inch, respectively.

a. Greatest Precipitation. There are three general types of precipitation: convective, cyclonic, and orographic. Each has its characteristics and geographical distributions, as well as different expectations for record extremes. Convective rain results from overturning of cooler air by warmer air from below and takes the form of heavy, localized showers, such as thundershowers. Convective showers tend to be most frequent in warm areas and seasons and are responsible for many of the extreme short-period rainfalls.

Cyclonic precipitation results from mechanisms associated with lowpressure centers (cyclones) and with zones of convergence of different air masses (fronts). The most severe cyclonic storms, hurricanes or typhoons, bring very heavy and prolonged vain and are responsible for most of the extreme amounts that occur over a period of several hours or days. Certain parts of the world, mostly occanic and coastal areas, lie along the tracks usually taken by these storms. The storms are prevalent in different areas at different times of the year; their tracks are mapped by month in the U. S. Navy's Marine Climatic Atlas of the World.⁹⁵

W. E. K. Middleton and A. F. Spilhaus, Meteorol Instruments, 3rd ed. Univ. of Toronto Press, Toronto, 1953.

R. E. Lautzenheiser and R. Fay, Heavy Rainfall at Island Falls, Maine, 28 Aug 59, Monthly Weather Review, Vol. 94, No. 12, pp. 711-714, Dec 66.

⁹⁵ H. L. Crutcher and O. M. Davis, U. S. Navy Marine Climatic Atlus of the World, v. 8, The World. (NAVAIR 50-1C-54). U. S. Naval Weather Service Command, 1969.

Orographic precipitation results from upward deflection of air when it strikes higher ground. This often occurs in conjunction with convective and cyclonic types and tends to increase the precipitation amounts produced by them; the increase is greatest on steep slopes. Precipitation can also increase when sharply narrowing valleys between slopes act as funnels on up-valley winds. Highlands in the path of moisturecarrying winds from warm seas have abundant and frequent precipitation; such areas have the highest average yearly rainfalls. Among them are the east- and south-facing slopes of the Himalayas, the western slopes of the Andes in Colombia, and mountain ranges along the northwest coast of North America. Annual mean precipitation values often differ slightly in different sources because of differences in the period of record. (This is also true of the average values for other meteorological elements.) The number of years on which the precipitation means included on the weather extremes maps are based are given in the table, pages 7-15. Generally, a longer period of record would be more reliable than a short one.

> World's greatest 1-minute rainfall 1.23 inches Unionville, Maryland, 4 July 1956

The U. S. Weather Bureau's investigation of this record is described by H. H. Engelbrecht and G. N. Brancato.⁹⁶ The extreme fall occurred during an afternoon of intense thunderstorms in the foothills of northern Virginia and adjacent north-central Maryland. At Unionville, the total precipitation during the storm was 3.60 inches, of which 2.84 inches fell during a 50-minute period from 1450 to 1540 hours (2:50 to 3:40 p.m.). Rainfall was measured with a recording rain gage located in satisfactory exposure. Some 13 points pertaining to functioning of the gage were considered in evaluating this record by Engelbrecht, then State Climatologist for Maryland, and T. E. Hostrander, who was Substation Inspector. An enlarged photograph of the recording rain gage chart revealed that at chart time 3:23+ the pen was at 2.47 inches on the chart scale and at chart time 3:23-, it was at 3.70 inches. It was concluded that "... 1.23 inches of precipitation occurred in an estimated period of one-minute or less"⁹⁷ This exceeded the previous world record 1-minute rainfall of 0.69 inch at Jefferson, Iowa, which in turn had exceeded the carlier record of 0.65 inch at Opid's Camp, California.

> World's greatest 42-minute rainfall: 12 inches Holt, Missõuri, 22 June 1947

⁹⁶ H. H. Engelbrecht, and G. N. Brancato, World Record One-Minute Rainfall at Unionville, Maryland, Monthly Weather Review, Vol. 87, No. 8, pp. 303-306, Aug 59.

[&]quot; Ibid.

G. A. Lott has examined the meteorological data available for this storm and considered the factors responsible for its remarkable intensity.⁹⁸ According to him, the storm occurred "... as a local intensification in a long, narrow, warm sector convective system (the leading edge of which may be interpreted as an instability line) a short distance ahead of a surface cold front" He further states that "... a unique factor was the tightening of the pressure gradient north of an instability-line Low, causing an extraordinarily strong low-level flow of unstable air into the pre-existing convective system"

> World's greatest 12-hour rainfall: 53 inches on 28-29 February 1964 Belouve, La Réunion I

World's greatest 24-hour rainfall: 74 inches on 15-16 March 1952 and

World's greatest 5-day rainfall: 152 inches on 13-18 March 1952 Cilaos, La Rénnion I

La Réunion Island is located in the Indian Ocean at approximately 21°S, 55°30′E. It is about 30 by 40 miles in extent and very mountainous, with steep slopes and narrow valleys. Sea surface temperature is highest during the tropical cyclone season, reaching 81°F in March.⁹⁹ The record-producing rainfall at Cilaos occurred during a tropical storm as did, presonably, that at Belonve and another very heavy rainfall (62.33 inches in 1 day and 136.83 in 5 days) at Aurere, in April 1958.¹⁰⁰ All three of these storms broke the previous 24-hour world record, 45.99 inches at Bagnio in the Philippines in 1911, and the Cilaos storm broke the previons 5-day world record of 150 inches at Cherrapunji, India, in August 1841. The values given for Anrere and Cilaos were obtained from a survey of about 6 years of official published data for those places, and the February 1964 rainfall at Belonve was reported in a communication from the French Meteorological Service.¹⁰¹ Since these record-breaking amounts "… were the result of an incomplete survey of a short period of record, there is a good chance that a more thorough survey of a longer period of record would disclose other, and perhaps, even greater, amounts of similar magnitude.¹⁰²

⁹⁸ G. A. Lott, The World-record 42-minute Holt, Missouri, Rainstorm, *Monthly Weather Review*, Vol. B2, No. 2, pp. 50-59, Feb 54,

J. L. H. Paulus, Indian Ocean and Taiwan Rainfalls Set New Records, Monthly Weather Review, Vol. 93, No. 5, pp. 331-35, May 1965, 100

Thid.

¹⁰¹ *lbid*.

¹⁰² Ibid.

Northern Hemisphere's greatest 24-honr rainfall: 49 inches Paishih, Taiwan 10-11 September 1963

Paishih is located at 24°33′N, 121°13′E at 5,368 feet on the island of Taiwan. Taiwan, like La Rénnico, is very mountainous and is surrounded by warm ocean water, 82°F in August and September during the tropical storm season. The record rainfall, 49.13 inches, occurred during typhoon Gloria, and it was measured in a recording gage, thus adding to the reliability of the observation.¹⁰³ Rain of similar intensity fell at nearby stations during the same storm. At one of these places, Paling, total rainfall during the typhoon was greater than at Paishih, and for some durations, intensity might have been greater. However, no further information on Paling is available.

> Dharampuri, India, had a 24-hour rainfall of 39 inches 2 July 1941 – possibly the world's greatest on flat terrain

Information on this rainfall extreme was obtained in 1962 by the late Paul Siple, then Scientific Advisor to the Director of Army Research.¹⁰⁴ Dr. Siple obtained it from the Director of the Indian Central Meteorological Research Institute of Poona, India. Dharampuri is at latitude 20.5°N in the Surat District of Gujarat. According to Siple.

> Institute personnel believe this to be a maximum rain on essentially flat terrain. Actually, it was a monsoon rain, and the Western Ghats (mountains) must bave had some effect. However, the Poona meteorologists minimized the orographic influence, for the location of the rainfall was not in the immediate vicinity of high mountains.

Yankeetown, Florida, had a 24-hour rainfall of 39 inches 5 September 1950

Yankeetown is located on the west coast of Florida at approximately 29°2'N. 82°40'W and would be at or near sea level. The 38.70 inches that fell there is

¹⁰³ Ibid.

¹⁰⁴ Indian Central Metcorol Research Inst. Poona. India. Communication from Director to P. A. Siple, Scientific Advisor to Director of Army Research, transmitted from Dr. Siple to Dr. W. B. Brierly, Earth Sciences Laboratories, U. S. Army Natick Laboratories, in letter dated 17 Jan 62.

North America's greatest 24-hour rainfall and could be a rival claim to Dharampuri's for rainfall on nonmountainous terrain. Although this is not an official record, it is one of the "few well-accepted unofficial extremes" included by Schmidli. 105

Australia's greatest 24-honr rainfall: 39 inches Crohamlmist, Queensland 3 February 1893

This record, discussed by Newman, is officially accepted and apparently quite reliable.¹⁰⁶ It occurred during a cyclonic storm in which comparable heavy rain fell at nearby stations.

World's greatest annual mean precipitation: 460 inches Mt. Waialeale, Kauai, Ilawaii

According to official sources, "... normal annual precipitation at Mt. Waialeale is 460 inches, the highest recorded annual average in the world ** 107 This amount is based on data for the period from 1931 through 1960. An average of 472 inches has been quoted in other sources, based on data for 1912 through 1949.108 Mt. Waialeale also had the greatest precipitation for 1 year of any place in the United States, 624 inches between 24 July 1947 and 27 July 1948.¹⁰⁹ This mountain is located on the island of Kanai, at 22°04'N, 159°30'W. Conditions pertinent to the record rainfall are described by Henning.¹¹⁰ According to him, the storage rain gage is at an clevation of 1,547 meters (5,075.5 feet), and measurements are made at 3-month intervals.

> Asia's greatest annual mean precipitation: 450 inches Cherraphnji, India

When the summer monsoon depressions (moderately vigorous, warmcored cyclonic disturbances accompanied by heavy rain) from the Bay of Bengal reach the Himalayas, the rainfall is further increased by orographic lifting.¹¹¹ As a result of these monsoon disturbances, which are still not fully understood, the eastern Himalayan

¹⁰⁵ R. J. Schmidli, op. cit.

¹⁰⁶ B. W. Newman, Australia's Highest Daily Rainfall, Australian Meteorol Mag. No. 20, pp. 61-65, Mar 1958. 107 U. S. Environmental Science Services Administration, Environmental Data Service, Local Climatoligical Data;

aunual summary with comparative data. 1967, Lihue. Ilawaii. Washington, D. C., GPO, 1968, 108,

L. II. Seamon and G. S. Bartle(1, op. cit.

¹⁰⁹ R. J. Schmidli, op. cit.

¹¹⁰ D. Henning, Mt. Waialcale, Wetter und Leben, Vienna, Vol. 19, No. 5/6, pp. 93-100, 1967.

¹¹¹ J. M. Walker. The Monsoon of Southern Asia: a review. Weather, Vol. 27, No. 5, pp. 178-89, 1972.

foothills are very wet regions. Cherraphuji is located in this area at 25°02'N, 91°08'E., 4,309 feet in elevation. The annual precipitation record there is based on a 74-year period.¹¹² In addition to the Asian record for annual mean precipitation, Cherraphuji also holds records for the world's greatest rainfalls for various durations of from 15 days to 2 years.¹¹³ Among these are the greatest rainfalls in 1 month, 366 inches in July 1861, and in 12 months, 1042 inches from August 1860 to July 1861. The greatest amount for a calendar year at Cherraphuji is 905.1 inches and the least, 282.6 inches.¹¹⁴ Another place in this Himalayan region, Mawsynram, Assam, could be a rival to Cherraphuji.¹¹⁵

Sonth America's greatest annual mean rainfall 3.54 inches Quibdó, Colombia

Quibdó is situated at an elevation of 120 feet at 5°41′N, 76°40′W. A rainfall average of 413 inches at Quibdó, based on data from 1931 through 1946 taken from Colombian sources,^{116–117} was cited on the 1964 revision of the *Weather Extremes* map. Earlier maps cited a value of 342 inches at Buena Vista, Colombia.¹¹⁸ The value of 354 inches, shown on the current revision, was obtained from Environmental Data Service.¹¹⁹

North America's greatest annual mean precipitation 262 inches Itenderson Lake, British Cohi**n**ibia

This station was at a fish hatchery, now closed, at the head of Henderson Lake on the west coast of Vanconver Island at 49°08'N 125°08'W, elevation 12 feet. The topography there contributes to extreme rainfall through orographic lifting reinforced by convergence.¹²⁰ Mountains to the north and northwest of the station are

¹¹² U. S. Environmental Science Services Administration, Worldwide Extremes of Temperature, Precipitation and Pressure Recorded by Continental Area, op. cit.

¹¹³ J. L. H. Paulhus, op. cit.

¹¹⁴ U. S. Environmental Science Services Administration, Worldwide Extremes of Temperature, Precipitation and Pressure Recorded by Continental Area, op. cit, 115

A. M. Riabchikov, Cherrapundzhi ili Manshuram-samoe dozbilivoe mesto ma Zemle? [Cherrapunji or Mawsynram — Which is the Rainiest Spot on Earth?] Moscow, Universitet, Vestudk, Ser 5, Geografiia, Vol. 25, No. 3, pp. 79-81, 1970.
 116 A. M. Riabchikov, Cherrapundzhi ili Manshuram-samoe dozbilivoe mesto ma Zemle? [Cherrapunji or Mawsynram — Which is the Rainiest Spot on Earth?] Moscow, Universitet, Vestudk, Ser 5, Geografiia, Vol. 25, No. 3, pp. 79-81, 1970.

¹¹⁶ Colombia, *Hol Agei*, between 1929 and 1947.

^{3.} II. Scamon and G. S. Bartlett, op. cit.

¹¹⁹ M. A. Arkin, Chief, Foreign Branch, Environmental Data Service, U.S. Environmental Science Services Administration, correspondence dated 6 Det 69.

¹²⁰ J. G. Potter, Record Precipitation on One Day in Canada, CDC #1-68. Canada, Meteorological Branch, Foronto, 1968.

at right angles to the main inflows of moist air. A direct onshore flow of moist air may be deflected by the mountains and converge in the Henderson Lake area. During these conditions, it is also likely that the outflow in the lowest level from Juan de Fuca Strait adds to the convergence. A nearby station, Ucluclet Brynnor Mines, had the greatest 1-day precipitation in Canada, 19.26 inches. Henderson Lake had the second greatest amount, 16.61 inches,¹²¹ as well as North America's greatest precipitation in 1 year. 320 (319.78) inches in 1931.122

Least Precipitation. Generally, areas of low precipitation occur in conb. tinental interiors, on lee sides of high momitains, on coasts adjacent to cool enrients, in zones of higher atmospheric pressure where the air is sal siding, and in high latitudes. Arid areas are found in east Africa and adjacent southwest Asia between 15° and 35°N. western South America between 5° and 30°S, eastern South America between 35° and 50°S, western Africa between 15° and 35°S, western and interior Australia, interior Asia, parts of western North America between 25° and 40°N, and in the polar regions, As with records of high average precipitation, those of low average precipitation vary according to the years on which they are based and tend to be more reliable for a longer period. In some very dry areas, e.g., Chile and Sudan, several years can pass with no precipitation.

Precipitation Variability. Variations in precipitation can be upward: c. i.e., occurrence of above-average amounts, or downward: i.e., occurrence of below} average amounts or even drought. Among the factors causing them are displacements of ocean currents and differences in strength of the monsuonal circulation from year to year. Coasts adjacent to cold enrients are generally dry, but if the enrient deviates even slightly, making room for warmer water, relatively abundant rainfall can occur, as in the coastal areas of Chile and Peru. Disastrons dronglets in northeast Brazil might be due to the opposite occurrence, invasion of warmer water by edd currents. The other factor, differences in strength of monsoonal circulations, is most prononneed along the borders of areas covered by these seasonal winds. During years of weak monsoon, less territory is covered by the rain-bearing winds and less rain is deposited by them.

> Debundscha, Cameroon, has 75 inches average variability of annual precipitation

The average and relative variability of anunal precipitation at 384 places throughout the world were tabulated by E. Biel.¹²³ and examined statistically by

121 D. M. Ludhum, op. ed.

122 . 15al.

¹²³ E. Bief. Die Veranderhehkent der Jahressumme des Niederschlags auf der Erde [The Variability of the Yearly Amount] of Precipitation of the Earth J. Geograph Lans Oesterreich, Leipzig, Vols, 11-15, pp. 151-80, 1929.

V. Conrad.¹²⁴ Places with greatest average variability (differences between mean value and individual yearly value averaged for a given number of years) were Debundscha. Cameroon, and Cherrapunji, India, with 75.28 and 66.02 inches, respectively. Debundscha is located at 4°01′N, 9°01′E at 30 feet elevation, and has the greatest average annual precipitation in Africa, 405 inches. Cherrapunji has the greatest average annual precipitation in Asia, 450 inches.¹²⁵ Because the rainfall amounts at these places are so high, the average differences from year to year can be correspondingly high without being extreme in proportion to the mean. In addition to having a very high average variability of annual precipitation, Cherrapunji may hold the record for the highest actual amount of variability. The difference between the greatest amount of precipitation there during a calendar year, 905.1, and the least amount, 282.6, is over 600 inches.¹²⁶

Themed, Israel, has a 94 percent relative variability of annual precipitation

Themed is a desert station on the Sinai Peninsula. Its relative variability record is based on the ratio between the mean deviation from the arithmetic mean of annual precipitation for a period of years beginning in 1921 and ending in 1947 and the arithmetic mean for those years.¹²⁷ Other places with high values based on this same measure of variability are Walvis Bay at 22°53′S, 14°26′E in South-West Africa with 87 percent,¹²⁸ and Malden Island in the Line Islands of the Equatorial Pacific with 71 percent.¹²⁹ Walvis Bay, like Themed, is a very dry area and because the rainfall amounts c these places are so low, a small variation in actual amount can become a large percentage of the mean value. Malden Island's average variability of 20.24 inches from its mean annual precipitation of 28.6 inches is thought to be due to displacement of ocean currents.¹³⁰

The relative variability of monthly rainfall can have much higher values. A value of 193 percent for September has been cited at Beersheba, Israel (31°14′N, 34°47′E) where rain fell in only one of 30 Septembers from 1921 through 1950.¹³¹

¹²⁴ V. Conrad. The Variability of Precipitation. *Monthly* II *eather Rev.* Vol. 69, No. 1, pp. 5-11, Jan 41, 125

¹²⁵ L. S. Environmental Science Services Administration, Rorbbeide Extremes of Temperature, Precipitation and Pressure Recorded by Continental Area, op. cit, 126

[&]quot; Ihid.

¹²⁷ J. Katsuelson and S. Kotz, On the 1 pper Limits of Some Measures of Variability, Archiv fur Meteorologie, Geophysik and Rioklimatologie, Ser. B, Vol. 8, No. 1, pp. 103-107, 1957.

^{1 28} Ibid.

¹²⁹ E. Biel, op. cit.

¹³⁰ fbid.

⁽a) 11 J. Katsuelson and S. Kotz, op. cit.

This almost reaches the theoretical upper limit of 200 percent determined analytically by Schumann and Mostert.¹³²

Lhasa, Tibet, had a 108 percent relative variability of annual precipitation, 1935 tbrongh 1938

Lhasa is located on the Tibetan plateau at 29°40'N, 91°07'E, elevation 12,090 feet. It is in an approximately east-west valley flanked on both sides by monntains of 15,000 to 16,000 feet. Its elimate has been described by Ginn-Tze Hsii, who established a meteorological station there in 1934,¹³³ and also by A. Ln¹³⁴ and H. Flohn.³⁵ Data for 1935 through 1938 from Ln's paper¹³⁶ were calculated by Coorad's methods¹³⁷ to obtain the average variability of 67.7 inches and a relative variability of 108 percent.

In 1936, the animal precipitation reported for Lhasa, 198.3 inclus, was more than 10 times greater than the average amount. However, there is some doubt as to the authenticity of the amount of precipitation recorded for 1936. Flohn considers the value questionable and attributes it to a possible misplacement of a decimal point by a partly educated weather observer.¹³⁸ On the basis of Ln's data and data for most of the years from 1941 to 1955. Floha found no amounts that even approached that of 1936. The highest was 22.9 inches. The fréquency of precipitation, as indicated by the number of rainy days during 1936, was not numsual. Furthermore, Gyantse, in the same climatic region as Lhasa at 28°56'N 89°36'E, clevation 10,486 feet, had no similar extreme variation in a 38-year period. In 1936, Gyantse had only 13 inches more than the average precipitation.

However, Hsñ was at Lhasa in 1936, presumably as the station observer or supervisor, and he wrote abont the unusually heavy rain and its causes¹³⁹ and informed Lu¹⁴⁰ of the difference between the rainfall in 1936 and in other years.

¹³² T. E. W. Schumann and J. S. Mostert, On the Variability and Reliability of Precipitation, Amer Meteorol Soc Bull, Vol. 30, No. 110, 1949.

¹³³ Gimi-Tze Hsti, A Noticon the Climatic Conditions of Lhasa, Amer. Meteorol Soc. Bull, Vol. 22, No. 2, pp. 68-70, Feb 11.

¹³⁴ A. Lu, V Brief Survey of the Climate of Lhasi, *Quarterly J Roy Meteorol Soc*, Vol. 65, No. 281, pp. 297-302, Jul 39.

 ^{1,35} H. Flohn, Beitrage zur Klimakunde von Hochasien, [Contributions Towards a Climatology of the Central Asian Platean], Erdkunde, Vol 12, No. 1, pp. 294-308, Dec 59, 136

¹³⁰ A. Un. op. cit. 137

¹³⁷ V. Courad, op. cit.

¹³⁸ II, Flohn, op. cit.

¹³⁹ Ginu-Tze IIsu, op. -it.

¹⁴⁰ A. L.n. op. cit.

Normally, rainfall at Lhasa comes from thundershowers, but in 1936, only 23 percent was of this type and 69 percent was from nighttime rain, possibly caused by interaction between a strong sonthwest monsoon and cold air masses from the north.¹⁴¹ In another paper. In described a similar extreme variation at Omei Shan, China, to the east of Lhasa, at approximately 29°30'N, 103°30'E, elevation 10.023 feet.¹⁴² The annual mean there is 73 inches, but during the Second Polar Year from August 3932 to August 1933 Omei Shan had 319 inches, "... the largest amount ever recorded in China in a 13-month period "¹⁴³ This difference of 246 inches between the annual mean and the Angust 1932 to Angust 1933 value is even greater than the difference between Lhasa's ammal mean and its 1936 precipitation. Flohn's paper also mentioned Omei Shan, but his figures differ somewhat from Lu's with a 25-inch annual mean and 300 inches in the Second Polar Year, making a difference of 275 inches.¹⁴⁴

> d. Hail.

U.S. largest hailstone 17.5 inches circumference Coffeyville, Kansas, 3 September 1970

Hailstones are pieces of ice which fall as precipitation either separately as spheres or cones or agglomerated into irregular limps. They originate in convective clouds of the commonimum stype and are usually associated with thunderstorms. The hailstone at Coffeyville, 37°02'N, 95°37'W is the largest officially recorded in the United States. It was irregular in shape and weighed 1.67 pounds.¹⁴⁵ It fell during a severe storm with hundreds of other large stones and was preserved and sent to the National Center for Atmospheric Research where it was photographed. The previous official record for largest hailstone in the United States, which appeared on the earlier Weather Extremes maps, was for a hailstone that weighed 1.5 pounds and measured 17 inches in cirenmference. It was recorded at Potter, Nebraska, on 6 July 1958,

Claims have been made of large bailstones in many places. One hailstone weighing 972 grams (2.14 pounds) was picked up and photographed in Strasbourg. France, near the German border on 11 August 1958, and a heavier one of 1.9 kilograms (4.18 pounds) was collected in Kazaklistan U.S.S.R. in 1959.⁹⁴⁶ In a review ¹⁴⁷ of a

^{141 [}hid.

¹⁴² A. Lu, Precipitation in Tiber, Geograph Rev. Vol. 37, No. 1, pp. 80-93, Jan 17, 143

Ibid. 144

H. Flohu, op. cit. 145

Amer Meteorof Soc Bull, Vol. 52, No. 2, title page, 1971.

J. P. Verdon, Extrêmes chinatiques mondules, [World Chinatic Extremes] - France - Direction de la Mitéorologie Nationale: Bulletin d'Information, April 1972, 147

K. Bognar, Remnüscences: Amer Meteorof Soc Bull, Vol. 52, No. 11, pp. 1402303, 1971

Hungarian publication,¹⁴⁸ if is noted that haifstones reported in the publication weighed 3.5 to 4.0 kilograms (8 to 9 pounds) compared to a weight of less than 1 kilogram for the one at Coffeyville. These Hungarian hailstones fell at Söjtör village in Zala country in August 1786, but "..., probably a few hailstones were clustered together..., "¹⁴⁹ Hailstones weighing up to 7.5 pounds have been reported from India and up to 10 pounds from China.⁶⁵⁰ Hailstones in these countries are said to have killed people and animals and destroyed villages.^{151–152}

e. Snow. There are two main ways of measuring snowfall depth: by direct measurement of fresh snow on open ground with a graduated ruler or scale, and by a snow gage. Precantions need to be taken against drifting or blowing snow, or, if the open ground method is used, against measurement of old snow. When there are strong winds, the snow gage is more accurate. Greatest amounts of snowfall – as of rainfall – occur in areas where there is moist air and a mechanism for lifting it. They tend to occur in the middle latitudes rather than at the very high latitudes where there is less moisture.

North America's greatest 24-hour snowfall 76 inches Silver Lake, Colorado 14-15 April 1921

Silver Lake is located at approximately 10°N, 105°40′W, at 10,220 feet elevation in the Colorado Rockies. The snowfall there in April 1921 established several records: 75.8 inches in 24 bours, prorated from a measured fall of 87 inches in 27.5 hours, 95 inches in 32.5 hours, 98 inches in 72 hours, and 100 inches in 85 hours.¹⁵³ According to Paulhus, the measurement was examined thoroughly before being accepted by the U.S. Weather Bureau.¹⁵⁴ ··· . . There was no evidence to indicate that the measurement was any less reliable than that of other heavy snowfalls, and it appears that a 'srowfall of this magnitude is meteorologically possible " The maximum amount of snow that can fall in 24 hours has been estimated as approximately 72 inches for snow with a density of 0.10 under normal packing conditions and correspondingly greater for lesser density.¹⁵³ The density of the snow at Silver Lake was 0.06.¹⁵⁶ During the storm.

¹⁵⁴ J. H. Field, The Meteorology of India, I Roy Soc. 198, Vol. 82, p. 7B4406, 1933-34, 152

Thid.

¹⁵⁵ L. F. Brooks, Un Maximum Snowfalls, Amer Meteorol Soc Bull, Vol. 19, No. 2, p. B7, Feb. 193B.

(56 J. L. H. Paullus, op. ett.

 ¹⁴⁸ V. Réffils. Weather Phenomena and Basec Wrought by Weather between 1700-1800 in Hongary, Hungarian Academy of Sciences, Budapest, 1974.

K Bognar, eye cat.

¹⁵⁰ L. H. Ladlant, The Hadstorm, Il enther, Vol. 16, No. 5, pp. 152-62, 1961.

¹³⁷ D. M. Ludhum, ego cit.

¹⁵³ J. L. H. Paullius, Record Snowtall of April 14-15, 1924 at Silver Lake, Colorado, Monthly Roother Rev. Vol. 54, No. 2, pp. 3B-10, Feb 53, 154

thunder occurred in various parts of the region, indicating widespread convective activity, and the combined convective and orographic influences produced excessive amounts of snow at several places. In addition to the record at Silver Lake, a fall of 62 inches in 22 hours was reported at Fey's Ranch, Colorado: both of these exceeded the previons U.S. record of 60 inches in 24 hours at Giant Forest, California, in January 1933.¹³⁷

> Alaska's greatest snowfall in 24 honrs, 62 inches, 29 December 1955 one storm, 175 inches, 26-31 December 1955 one season 974 inches, 1952-1953 Thompson Pass

This station is in south-central Alaska, at 61°07′N, 145° 14′W, elevation 2.700 ft. It ranks near the top among places holding North American records for the greatest amounts of snowfall in 24 hours, during a single storm and during a season. Besides this, Thompson Pass has extremely frequent occurrences of these intense snow-storms. During the 20 winters from 1951-52 through 1970-71, there were 32 snow-falls of 30 inches or more during a 24-hour observation period reported at this station, and 12 of these snowfalls were at least 40 inches.¹⁵⁸

Bessins, France, had a snowfall of 68 inches (67.7) in 19 honrs, 5-6 April 1959

Bessans is located at 1710m (5,610 feet) in the French Alps near the Italian border. Very intense snowstorms occur in this area as the result of a South East wind, known locally as "la lombarde." These storms, of which the one at Bessans is a good example, are also very localized; the meteorological and environmental conditions contributing toward them are examined by M. Jail.¹⁵⁹ Parts of Norway also have very high 24-hour snowfalls, as do parts of nortbwestern Japan and some other world areas. In this connection, it should be mentioned that the predominance of North American records in our listing (see the table, pages 7-15) is due to their availability rather than to greater amounts of snowfall on this continent.

5. Other Extremes. Besides temperature and precipitation, several other meteorological and climatic conditions are shown on the maps. For each of these, the most extreme occurrences have their own particular set of causes, limits, and distributions in time and space, and for each there are problems in obtaining accurate measurements.

¹⁵⁷ 158 J. L. H. Pardbos, op. cit.

¹⁵⁸ Climatological Data, Monthly, Haska, Asheville, N. C., L.S. Weather Bureau, 1951-65, U.S. Environmental Data Services, 1965-71. 159

⁵⁹ M. Jail, Remanphable effet de kombarde: les ebutes de neige de Paques 1969 en Haute-Maurienne, Reche de Geographie Alpine, Frenuble, Vol. 57, No. 3, pp. 613-21, 1969.

In evaluating reliability of the records, all of these factors should be taken into consideration.

a. Thunderstorms. A thunderstorm is defined as "... in general, a local storm invariably produced by a common inhibit cloud, and always accompanied by LIGHENING and THUNDER, usually with strong gusts of wind, heavy rain, and sometimes HAH... It is usually of short duration, seldom lasting over two hours for any one storm ..., "¹⁶⁰ Criteria for recording thunderstorms can vary between storms actually occurring at a station, or merely thunder heard or lightning seen from the station.

Thunderstorms are most prevalent in warm weather and, in some places, during the rainy season. However, although thunderstorms frequently produce heavy rainfall, there are places where seasons of rainfall and thunderstorm maxima do not coincide. In addition, some very rainy places have few thunderstorms, while places with very frequent thunderstorms can have relatively small amounts of rainfall.¹⁶¹

Kampala, Uganda averages 242 thnuderstorm days per year

¹⁶⁰ Glossery of Environmental Terms (Terrestrial), MIL-STD 1165, Washington, D. C., Dept of Defense, 25 Mar 68, 161.

¹⁶⁾ W. H. Portig and J. R. Gerhardt, Research in Tropical Meteorology, Second Interna Technical Report, Austra, Texas, Eury of Texas, Electrical Engineering Research Laboratory. (Sponsored by U. S. Army Research and Development Laboratory, Fort Monimonth, New Jersey), 1962.

¹⁶² World Meteorological Organization, World Distribution of Thomberstoric Days (OMM, No. 21), Geneva, Switzerland, 1953, 264

¹⁶³ F. F. Umib, Lopographic Influences on Fluinderstorm Activity Near Lake Victoria, Il cather, Vol. 25, No. 9, pp. 104-10, 1970.

¹⁶⁴ W. H. Portig, Research Meteorologist, U. S. Army Tropic Lest Center, Fort Clayton, Canal Zone, Correspondence dated 23 Nov 70.

breeze front.¹⁶⁵ Thus, the Kampala area is subject to frequent thunderstorm activity. and nearby stations such as Entebbe and Kisumu also average very high numbers of thunderstorm days per year.166

Bogor, Indonesia, averaged 322 thunderstorm days per year 1916-1919

Bogor, formerly Buitenzorg, is located on the island of Java at 6°30'S, 106°48′E. The mean annual number of thunderstorm days recorded there changed from 151 in the years 1841-1857 to 322 in 1916-1919; it ranged from 4 to 41 for the years 1953 through 1962.¹⁶⁷ These differences might reflect changes in the criteria for recording thunderstorms. For instance, "... during a certain period " Bogor recorded lightning seen in addition to thunder heard and thunderstorm at the station.¹⁶⁸ Also, the interpretation of 322 in the Bogor record is doubted low W. 11. Portig¹⁶⁹ and others who consider that it represents the "... mean numbers of occurring thunderstorms and not the mean number of days on which thunder was heard* According to Portig, the "... absolute occurring maximum " is indicated by statistical curves. to be "... approximately 250 or 260 thunderstorm days annually, although such a location has not as yet leen found ** 170

h. Air Pressure: The pressure value for a given unit area of surface verv nearly represents the actual weight of a vertical column of air of the same unit area. extending upward from that surface to the top of the atmosphere. At sea level, this column of air averages about 14.7 pounds (29.92 inches of mercury) per square inch of exposed surface, but the weight varies with latitude and with changes in daily weather. It falls with altitude from an average of 31.3 inches in the earth's deepest depression. 1.300 feet loclow sea level, to less than half that amount (about 14.9 inches) at 18,000 feet. At the top of the world's highest monntain, 29.028-foot Mt. Everest, it would be about 9 inches.

Air pressure is usually measured by mercurial or aneroid harometers. The former halances pressure of the atmosphere against the weight of a column of merenry. The aneroid or elastic type contains a hollow metal chamber, partly emptied of

Ibid.

¹⁶⁵ F. E. Lamb. op. cit. 166 [bid.

¹⁶⁷

V.I. Arabadzhi, Khinat i grozy, Climate and Thunderstori is. Priroda, No. 2, pp. 65-66, Feb 66. Translated by E. R. Hope as Canada, Defence Research Roard, Translation T156R, Apr 66,

¹⁶⁸ W. H. Portig. op. ed.

¹⁶⁹ W. H. Portig, Thuoderstorm Frequency and Amount of Precipitation in the Tropics, Especially in the African and Indian Monsoon Regions. Trehie für Meteorologie, Geophysik und Bioklämatologie, Ser. II, Vol 13, No. 1. 170 pp. 21-35, 1963,

air and scaled, which expande and contracts as the pressure changes. Mercurial barometers are generally more accurate, but the aneroids are smaller and more portable. Since mercury is affected by temperature and gravity, adjustments must be made in the readings of mercurial barometers to allow for these factors. Also, in order to standardize readings of any kind of barometer when made at different times and places, adjustments are required to compensate for errors in the individual instrument used (index error) and for differences in altitude. For the altitude correction the station pressure readings are usually equated to sea level pressure, and these are the values which are recorded. There are possible sources of error peculiar to each type of barometer, as well as errors that could result from making the various types of adjustments. For example, different values can be obtained when different methods are used for reducing the actual or station pressure to its sea level equivalent.

(1) High Pressure. The highest station pressures occur at the lowest elevations; i.e., in depressions below sea level such as Death Valley. California, where the lowest point is -280 feet; the Qattara depression in northwestern Egypt, with a minimum elevation of -436 feet; and the shores of the Dead Sea between Israel and Jordan, at -1,286 feet. In the Dead Sea area, a station pressure of 31.91 inches was reported at Sedom (the biblical Sodom) on 21 February 1961.¹⁷¹ Another depression, the Torfan in central Asia at approximately 45°N, 90°E, is thought to have station pressures similar to or even higher than those of the Dead Sea area.¹⁷² However, after the adjustment to sea level is made for pressure readings at places below sea level, the values recorded are lower. At or near sea level, high pressures occur along Aretic coasts. Values of 31.37 inches (31.38 sea level equivalent) and of 31.40 inches (31.41 sea level) have heen recorded at Clyde in northeastern Canada and at Myggbukta in East Greenland on 18 January 1958 and 15 January 1940, respectively.^{173–174} At stations above sea level the pressure readings are actually lower, but when equated to their sea level convalents, the values recorded are higher.

World's highest sea level air pressure: 32.01 inches Agata, U.S.S.R., 34 December 1968

The highest recorded pressures occur in Siberia during winter. Agata is located in this area at 66°53′N, 93°28′E, elevation 855 feet. The high

¹⁷¹ A. Coart, Improbable Pressure Extreme: 1070 mb. Duer Meteorol Soc Bull, Vo. 50, No. 1, pp. 248-50, Apr 69.

¹⁷⁷ R. Scherbag, Newe Methoder der Bettermidyse und Betterprognose, [New Methods of Weather Analysis and Weather Forecast]. Revin, 1948.

pressnre record there was the culmination of an intense anticyclone which originated in East Siberia on 22 December and continued until 2 January when it declined somewhat as it moved south and west into European Russia.¹⁷⁵ On 31 December, seven stations in north-central Siberia had air pressures above 31.565 inches (1.070 millibars). The weather was clear and calm with temperatures between -40° and -50°C. Previous world records for high pressure were 31.84 inches at Barnaul, Siberia, in 1900 and 31.75 inches at Irkutsk, Siberia, in 1893.

(2) Low Pressure. The lowest station pressures occur at the highest altitudes; e.g., in the Himalayas of Asia, the Andes in South Ameria, and the Rockies in western North America. Beginning at about 4,000 feet, the average station pressures are lower than the most extreme low pressures that occur at sea level. At the approximate upper limit of weather stations, 15,000 feet, standard pressure is 16.90 inches and the extreme lowest is estimated at about 14.76 inches.¹⁷⁶ At sea level, the lowest pressures and most rapid falls in pressure occur during hurricanes and tornadoes. "... Accurate pressure measurements by a recorder directly in the center of a tornado have never been made"¹⁷⁷ but various estimates are available. According to one estimate, the drop in pressure can be as great as one fifth of an atmosphere, i.e., about 6 inches.¹⁷⁸ Other estimates are for "... a reduction of no more than one fourth of the pre-existing pressure ..." which could occur "... within 15 seconds"¹⁷⁹ and a drop of 1.5 pounds per square inch (3 inches) and a time change of 0.5 psi (1 invh) per second in pressure of tornadoes with 200-mile per hour winds.¹⁸⁰

> World's lowest air pressure at sea level (excluding tornadues) 25.90 inches estimated by aerial reconnaissance in eye of Typhoon 1da at 19°N, 135°E, 24 September 1958

Reliability of the estimated record has been evaluated by Jordan.¹⁸¹ It was based on a dropsonde observation made from the 700-millibar level by a U. S. Air Force reconnaissance aircraft of the 54th Weather Reconnaissance Synadron,

¹⁷⁵ B. D. Giles, Extremely High Atmospheric Pressures, Beather, Vol. 25, No. 1, p. 19-24, 1970.

A. Conrt and H. A. Sadniela, Improbable Weather Extremes and Measurement Needs, Amer Meteorol Soc Bull, Vol. 34, No. 9, pp. 573-75, Sep 53,
 477

D. M. Ludhum, op. cit. 178

¹⁷⁶ "Tornadoes," in: Encyclopedia of Atmospheric Science and Astrogeology, Reinhold, N. Y., 1967, 179

A. Court and H. A. Salmela, op. cit.

 ¹⁸⁰ T. T. Fujita, Estimate of Maximum Wind Speeds of Tornadoes in Three Northwestern States, Chicago Univ., III. Dept. of the Grophysical Sciences, Research Paper 92, NOAA-71000308, Dec 70.
 181

¹⁸¹ C. L. Jordan, A Reported Sca-Level Pressure of 877 mb, Monthly Weather Review, Vol. 87, No. 9, pp. 365-66, Sep 59.

about 600 miles northwest of Gnam. In the observation, the 700-millibar height was determined from the pressure and radio altimeters aboard the aircraft, and the 850-millibar height and sea level pressure were computed from temperature and humidity data measured by the aircraft instruments at flight level and by the dropsonde instrument at levels below the aircraft. According to Jordan, the sounding was found to be hydrostatically consistent and the vertical temperature distribution was realistic.¹⁸² A source of major error in this type of measurement would be in the aircraft altimetry, but on reconnaissance aircraft, calibration checks are a routine part of each flight. Dropsonde observations from two other aircraft, made about 8 hours earlier and 23 hours later, respectively, both reported sea level pressures lower than the previously accepted lowest pressure of 26.185 inches from the SS Supperea. Lower values than the Suppercu's were also reported in 1953 during Typhoon Nina, with dropsonde sea level pressures of 886 millibars (26.17 inches) and 883 millibars (26.04 inches).¹⁸³ However, the record taken on the Sapoerea, 460 miles east of Luzon, Philippine Islands, on 18 August 1927 was an actual measurement rather than an estimate, and it remains the lowest measured pressure at sea level. 184

c. Solar Radiation. Measurements of the flux of solar radiation penetrating to the lower layers of the atmosphere can be subdivided into several main classes. Values considered here are for global solar radiation received on a horizontal surface. "... This includes both radiation received direct from the solid angle of the surfs disc and also radiation that has been scattered or diffusely reflected in traversing the atmosphere "¹⁸⁵ Such measurements are usually made with pyranometers, and, as in all radiation measurements, considerable care and attention to detail is required to insure accuracy. Several points to be evaluated in estimating accuracy of radiation measurements are listed in the WMO Guide.¹⁸⁶

> Sonth Pole has 955 Langleys average daily insolation in December

The world's highest daily amounts of solar radiation are received on the Antarctic Platean during summer when there are 24 hours of continuous daylight.¹⁸⁷ The North Polar area also has continuous daylight during its summer, but at that time of the year the earth is about 3 million miles further from the sun. Consequently,

186 *[bid.* 187

^{182 [}bid.

^{183 [}bid.

¹⁸⁴ R. J. Schmidli, op. cit.

¹⁸⁵ World Meteorological Diganization, op. eit.

¹⁷ K. J. Hanson, Some Aspects of the Thermal Energy Exchange on the South Polar Snow Field and Arctic Ice Pack, Monthly Wenther Review, Vol. 09, No. 5, pp. 173-77, May 61.

about 7 percent more solar radiation impinges on the top of the Antarctic's atmosphere than on the Arctic's during midsummer.¹⁸⁸

The South Pole record was obtained by averaging daily values of hemisp'-rie global solar radiation, which were available from Annudsen-Scott Station for the Decembers of 1958 through 1965.¹⁸⁹ The resultant value was 954.6 langleys, Ammulsen-Scott is located at an elevation of 9,186 feet, at 90°S. However, the average daily insolation could be greater at other stations in the Antarctic.¹⁹⁰

> Malange, Angola had a solar radiation of 113 langleys in one hour 7 November 1961

Malange is located in western Africa at 9°33'S, 16°22'E, altitude 3,710 feet. A very high reading of 112 langleys in 1 hour was also obtained at Windhoek, South West Africa (22°34'S, 17°16'E, altitude 5.640 feet) on 20 December 1956.¹⁹¹ High hourly values of solar radiation can be expected at relatively low latitudes during the hours of the day and times of the year when the sun is strongest and when cloud cover is absent and the air is as clear as possible from dust and other impurities. Other things being equal, they would tend to occur at the higher elevations.

Wind Speed. Of all the elements, wind is most variable. To compare d. wind speeds reported from varions places and times, information should be known about the height and exposure of the measuring instrument (anemometer), the type of instrument and record, and the time interval covered by the measurement. Wind speed is usually measured by either rotating or pressure anemometers. In the former type, wind passage is measured by the rate of motion imparted to a freely rotating mechanism. and a timing mechanism is usually combined with the anemometer to indicate the rate of passage. Pressure anennometers measure the instantaneous speeds (actually averages for about 1 second) by means of pressure effects; i.e., force applied to a surface or surfaces. The time intervals of wind observations differ between countries and even between stations. In the U.S., the extreme values tabulated are: the greatest average speed over a 5-minute interval, maximum speed: the speed of the fastest mile of wind from the record of rotating anemometers, extreme speed; and the highest reading of the instantaneous recording anemometers, strongest gust.¹⁹² In addition, extreme average

¹⁸⁸ Ibid.

¹⁸⁹ 1. S. Environmental Science Services Administration, Environmental Data Service, Elimatological Data for Autarctic Stations, Nos. 1 and 9, Washington, D. C., 13PD, 1912 and 1968. 190 Personal communication, M. Kuhu, University of Inushruck, Ionsbruck, Austria.

¹⁹¹

Quart Radiation Bull, Union of South Africa, Vol. 3, No. 172, p. 17, 1957.

¹⁹² A. Court, Wind Extremes as Design Factors, J Franklin Inst. Vol. 256, pp. 39-56, Jul 53.

values are obtained from records of the total wind passage in each hour from recording rotation anemoneters at the principal weather stations.¹⁹³

Winds are strongest at the times and places of maximum temperature and pressure gradients. They increase with altitude and during thunderstorms, hurricanes, and tornadoes. Gnsts are sudden brief increases in the speed of wind, usually less than 20 seconds in duration.¹⁹⁴ and are known to exceed 200 miles per hour in hurricanes and tornadoes.¹⁹⁵ A tornado is defined as "... a slightly funnel-shaped, upwardspiraling wind column of destructive velocity "¹⁹⁶ Because wind measuring instruments are mable to withstand the full impact of a tornado, estimates of the wind speed are made on the basis of various criteria such as the amount of force that would be required to cause the resultant damage. According to Ludhum, the maximum winds in a tornado probably range between 150 and 225 miles per hour.¹⁹⁷ Fujita gives a similar range of values, 175 to 225 miles per hour based on his investigation of tornadoes occurring in the states of Washington, Oregon, and Idaho during the years 1950-1969.¹⁹⁸

> World's highest surface wind (excluding tornadoes) 231 miles per hour, peak gust, and 188 miles per hour, 5-minute wind speed, on 12 April 1934 and

U. S. highest annual mean wind speed 35 miles per hour (35.4) Mt. Washington, New Hampshire

Winds are stronger at the summit of this 6.288-foot monutain, at 44° 16'N, 71° 18'W, than they are at the same elevation in the free air some distance away. "... This is probably due to 'uplift' over the slope or to the Bernoulli effect introduced by the surrounding monutains. Windspeeds of 100 mph are not uncommon"¹⁹⁹ The speed of the peak gust was measured by a heated rotation anemonieter, but, in such strong winds, no apparatos can record the airflow except approximately, and actual

197 Ibid.

¹⁹³ 19. Ibul.

R. E. Huselike, ed., Glossary of Meteorology, Boston, Mass., Am Meteorol Soc, 1959, 195

C. F. Riley, Jr. and P. F. Walker, Atmospheric Pressure Criteria for Aircraft and Airborne Equipment, WADC Tech Note 55-98, Chapter II, "Wind," Wright-Patterson Air Force Base, Ohio, 1955. 196

¹⁹⁷ FI, M. Emilinm, op. cit.

T. F. Enjita, op. cu.

 ^{(19) &}lt;sup>11</sup> F. Cupus *Querna* (19) F. S. Environmental Science Services Administration, Environmental Data Service, *Local Climatological Data:* animal summary with comparative data, 1967, Mount Washington Observatory, Gorham, New Hampshire, Washington, D. C., U. S. Goyi, Print, Off., 1960.

velocity may be in error by 10 to 40 miles per hour.^{200 201} The 231-mile-per-hour value is documented in the official records.²⁰² A value of 225 miles per hour, after anemometer calibration, is given in some sources²⁰³ and was cited in previous editions of the Weather Extremes map. Direction of the wind during the gust was from the southeast, the direction from which the most severe storms in the Mt. Washington area usually come. Its force, due to the reduced air density on the mountain top, was equal to that of about a 180-mile-per-hour wind at sea level,²⁰⁴

> Thule, Greenland, had a 207 miles per hour peak gust 8 March 1972

This peak wind speed of 180 knots (207 miles per honr) was recorded at 2155 honrs (9:55 p.m.) at an off-base survival shelter at 76°31′20″N, 68°19′00″W, elevation 990 freet.²¹² It was observed by two site dispatchers on an anomometer capable of

²⁰⁹ C. F. Brooks, The Worst Weather in the World, Appalachia, pp. 194-202, Dev 40.

²⁰¹ Note on the Precipitation, Snowfall and Wind Records, Mount Washington Observatory, News Bull, No. 22, pp. 23-24, Jun 53.

 ²⁰² L. S. Environmental Science Services Administration, Environmental Data Service, *Local Climatological Data:* animal summary with comparative data, 1967, Mount Washington Observatory, Gorham, New Hampshire, Washington, D. G. L. S. Goyt, Print, Dff., 1968.
 ²⁰³

²⁰⁵ Hid.

R. J. Schmidli, op. cif.

²⁰⁶ C. F. Brooks. The Worst Weather in the World. Appalachia, 10, 194-202, Dec 10, 207

V. Court, Wind Extremes as Design Factors, J Franklin Just, Vol. 256, pp. 39-56, Jul 53.

²⁰⁸ F. S. National Oreanic and Atmospheric Administration, Environmental Data Service. Local Climatological Data: annual summary with comparative data, 1950, Blue Hill Observatory, Mitton, Massaclassetts, Washington, D. C., GPU, 1971.

²¹⁰⁷ L. H. Seamon and G. S. Bartlett, op. cit. 210

Note on the Precipitation, Snowfall and Wind Records, op. ed.

²¹⁴ F. Lanscher, I ber die Verteilung der Windgeselnwindigkeit ant der Fide. On the Distribution of Wind Velocity Over the Earth. Arch Meteorol Geophys Bioklimatol, Series B, Vol. 2, No. 5, p. 436, 1951.

²¹² J. R. Stanstield, The Severe Arctic Storm of 8-9 March 1972 at Thule Air Force Base, Greenland, Weatherwise, Vol. 25, No. 5, pp. 228-33, 1972.

registering winds of me to 276 miles per hour. It occurred at a considerably lower elevation than the peak gust on Mt. Washington and, consequently, could have exceeded it in force. The gust was observed during a severe Arctic storm in which the site experienced winds of 146 miles per hour or greater for 4 hours. Similar, though somewhat lesser, speeds were observed throughout the area. The U.S. Air Base at Thule had a speed of 110 miles per hour. High winds are common in this northwest Greenland area due to intense winter storms and local topographic conditions. Winds blowing down from the dome-shaped Greenland icceap are accelerated by gravity and by compression and funneling as a result of the local orientation of mountains and valleys.

> Miyakojima Island, Rynkyn Islands had a 190 mile-per-honr µeak gust and a maximum wind speed of 136 miles µer honr 5 September 1966

These wind speeds were observed at the Miyakojima Weather Station and have been officially reported by the Rynkyn Meteorological Agency. They occurred during a typhoon whose maximum wind zone passed over the island. The peak gust of 85.3 meters per second (190 miles per hour) was recorded at 0(31 hours (6:31 a.m.) and the maximum wind of 60.8 meters per second (136 miles per hour) at 0731 hours (7:31 a.m.)²¹³. The direction of both was Northeast. Miyakojima is a small, flat island located at about 24° 47′N, $125^{\circ}17′E$. Since the record there occurred at sea level, there would have been no reduction in the force of the wind. Typhoons of similar severity occur about once or twice a year in the northwest Pacific area.²¹⁴

> Port Martin, Autarctica, had a 24-hour mean wind speed of 108 miles per 5, sur-21-22 March 1954 a mean monthly wind speed of 65 miles per hour-March 1954

These wind speeds were recorded at a station maintained by the Expéditions Polaires Françaises at 66° 19'S. 111°21'E on the coast of Adélie Land from February 1950 to January 1952.²¹⁵ The wind measuring installations did not fully comply with international standards, being somewhat sheltered by snowdrifts, or the values would have been slightly higher.²¹⁶ Winds of Intricane force (74 miles per boar or greater) were recorded at the Jation on 10 consecutive days in March 1953 and on 122

214 Ibid.

215 F. Locwe, The Land of Storm, Weather, Vol. 27, No. 3, pp. 116621, 1972.

21.6 *Ibid*.

²¹³ Y. Mitsula and S. Yoshizumi, Characteristics of the Second Miyakojima Lyphoon, *Kyoto, Japen, Luiv, Director Prevention Research Bulletin*, Vol. 38, pt. 1, No. 131, pp. 15-34, 1968.

days during calendar year 1951. The hurricane force winds are extremely steady both in their direction, south-southeast, and in their strength, which varies very little and appears lacking in normal gustiness. However, they can within a minute drop from full hurricane strength to near-calm and just as suddenly return. Extreme winds were also reported by Mawson's expedition at nearby Cape Denison, 67°1'S, 142°41'E in 1912 and 1913,²¹⁷ Various explanations of the causes of strong winds in this part of Autarctica have been advanced;²¹⁸ but as yet the reasons for them are not definitely established,²¹⁹

e. **Dewpoint (Humidity).** Dewpoint is "... the temperature to which a given parcel of air must be cooled at constant pressure and constant watervapor content in order for saturation to occur ... "²²⁰ This temperature is measured by instruments in which a mirror or other element is cooled to the saturation point. High dewpoints indicate high lumidity, and low dewpoints indicate low lumidity. As far as could be ascertained, there are no accepted extreme records of high or low dewpoints. The high values cannot be higher than the temperature of the body of water from which the vapor originates. They occur in proximity to water bodies with high surface temperatures such as parts of the Gult of California, the Red Sca, the Persian Gulf, and possibly in some tropical swamplands.

Assab, Ethiopia, has 84°F Average afternoon dewnoint in June

The dewpoint value for Assab was taken from a footnole in a report by A. Dodd:²²¹ ··· . . . Recently available data furnished by the National Weather Records Center indicate that very high dew points occur also in the Red Sea littoral. Assab and nearby Ras Audalighe on the Red Sea coast of Eritrea (Ethiopia) had *average* afternoon dew points higher than 84°F "

f. Fog.

U. S. West Coast highest average fog frequency 2552 hours per hear Cape Disappointment, Washington

²¹⁷ Heid.

²¹⁸ K. B. Mather and G. S. Miller, The problem of the Katabatic Winds on the Coast of Terre Adélie. *Polar Record*, Vol. 13, pp. 125-32, 1967.

²¹⁹ F. Loewe, op. cit.

²²⁰ B. E. Huschke, ed., op. cit.

²²¹ A. V. Dodd, Areat and Temporal Occurrence of High Dev Printwood Associated Temperatures, Tech Rpt LS-49, U. S. Arny Natick Laboratories, Natick, Mass. Aug 69.

U. S. East Coast highest average fog frequency 1580 hours per year Moose Peak Lighthouse, Mistake Island, Maine

Fog is a cloud at the surface of the earth, consisting of a multitude of miunte water droplets suspended in the atmosphere; according to international definition, it reduces horizontal visibility below 1 kilometer, or 0.62 miles,²²² It can occur. if there are sufficient condensation nuclei present, whenever the air is cooled to its dewpoint or the dewpoint is raised by the addition of moisture to the air. Fog occurs most often on monutains where the surface of the earth is high enough to be in the clouds and on coasts where land and water temperatures differ and moisture is present. The records for Cape Disappointment and Moose Peak Lighthouse were obtained from M. A. Arkin of the U.S. Environmental Data Service. They are based on fog signal operation and on low visibility operation of radio beacons at light stations, lightships, and other Coast Guard units during a period of 10 years or longer. Also, for a 4-year period at Willapa, Washington, the average was 3,863 hours per year, and for one of those years the total was 7,613 hours. Mountain stations with very frequent occurrences of fog are Mt. Washington, New Hampshire, averaging 308 days a year on which there is some tog, and Stampede Pass, Washington, averaging 252 days,²²³

III. CONCLUSIONS

Conclusions. From the foregoing examples and discussions, several points 6. seencto stand ont:

For each meteorological and climatological element there are certain cona. ditions or combinations of conditions that favor extreme values and are most likely to occur within particular geographical areas or seasons of the year: also, there are upper and lower limits of the values that can possibly occur. The records shown on the world and North American maps appear to be within the limits of possibility and to have occorred in places and at times in which the necessary conditions could be expected. However, occurrences of equal or greater extremity could have happened at other places in the same areas or at other times in the same places without being recorded or withont being publicized. There is no routine exchange of such information between differcut constries.

b. For each meteorological and climatological element, there are factors of site, instrumentation, and observational procedure which can affect the measured values.

^{222 11.} M. Ludlum, op. cit.
223 Ibid.

To insure uniformity of measurements, certain standardized equipment and procedures are recommended by the World Meteorological Organization. However, standardized equipment can malfunction and errors can be made in observation and recording, even inder standard procedures. Further, improvements in the reliability of measurements continue to be made as knowledge of both technology and the physical environment increases. Some of the earlier records might have different values if measured with the newest instruments and procedures; even now, extremes are sometimes not measured exactly, or at all, because they exceed the scale of standard meteorological instruments or their rate of response.²²⁴ For these reasons, it is not to be assumed that any one of the values shown on the maps in this report is correct to the tenth of a decimal place; but those for which a claim of record extreme is made were obtained under conditions that were acceptable to the Environmental Data Service.

c. Considerable research remains to be done both on the general subject of extreme weather and chimate and on the records of individual extremes. There is more to be known about the causes of extreme conditions as well as their absolute limits, frequencies, and distributions in time and place. Further search would yield additional records of extremes which are not included on the present maps.

²²⁴ A. Court and H. A. Salmela, op. cit.