

AD-A031 945

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

SPECTRAL PROPERTIES AND OSCILLATION THEOREMS FOR MIXED BOUNDARY--ETC(U)

SEP 76 J W LEE, A PINKUS

DAAG29-75-C-0024


UNCLASSIFIED

MRC-TSR-1683

NL

1 OF 1  
AD  
A031945





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ADA031945

MRC Technical Summary Report #1683

SPECTRAL PROPERTIES AND  
OSCILLATION THEOREMS FOR MIXED  
BOUNDARY-VALUE PROBLEMS OF  
STURM-LIOUVILLE TYPE

J. W. Lee and A. Pinkus

**Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706**

September 1976

(Received May 14, 1976)



*See Form  
1473*

Approved for public release  
Distribution unlimited

Sponsored by  
U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

SPECTRAL PROPERTIES AND OSCILLATION THEOREMS FOR MIXED  
BOUNDARY-VALUE PROBLEMS OF STURM-LIOUVILLE TYPE

J. W. Lee\* and A. Pinkus\*\*

Technical Summary Report # 1683  
September 1976

ABSTRACT

This paper presents analogues of the spectral and oscillatory properties exhibited by classical Sturm-Liouville systems for certain mixed boundary value problems that include periodic boundary conditions as a special case. These mixed boundary-value problems have Green's functions which are sign consistent for all even and/or odd orders.

AMS (MOS) Subject Classification: 34B25

Key Words: Mixed boundary conditions  
Green's function  
Tchebycheff systems

Work Unit No. 1 (Applied Analysis)

ADDITIONAL INFO	
AMS	Whole Section <input checked="" type="checkbox"/>
DIC	Full Section <input type="checkbox"/>
DECLASSIFIED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DESCRIPTION, AVAILABILITY CODES	
Dist. A FIL. OR W. SPECIAL	
A	

\* Department of Mathematics, Oregon State University, Corvallis, Oregon 97331.

\*\* Mathematics Research Center, University of Wisconsin, Madison, Wisconsin 53705.



SPECTRAL PROPERTIES AND OSCILLATION THEOREMS FOR MIXED  
BOUNDARY-VALUE PROBLEMS OF STURM-LIOUVILLE TYPE

J. W. Lee<sup>\*</sup> and A. Pinkus<sup>\*\*</sup>

§1. Introduction

This paper presents analogues, for certain mixed boundary-value problems, of the spectral and oscillatory properties exhibited by classical Sturm-Liouville systems. The usual analysis of the Sturm-Liouville eigenvalue problem is based on special ad hoc methods. In contrast, Gantmacher and Krein showed in [ 3 ] (see also references therein) that these fundamental spectral properties are direct consequences of the total positivity of the Green's function for the problem. They further showed that the total positivity of the Green's function is itself the mathematical expression of certain basic physical properties of vibrating mechanical systems, which are typically modeled by Sturm-Liouville systems. Subsequent to the work in [ 3 ], extensive studies have revealed several important classes of boundary-value problems with separated boundary conditions whose Green's functions are totally positive or sign regular. As in the classical Sturm-Liouville problem, these boundary-value problems exhibit a rich oscillation theory. Some principal contributors in this area are

---

\* Department of Mathematics, Oregon State University, Corvallis, Oregon 97331.

\*\* Mathematics Research Center, University of Wisconsin, Madison, Wisconsin 53705.

Gantmacher and Krein [ 3], Karlin [ 6], [ 7], [ 8], Karon [ 12], Krein [ 13], and Krein and Finkelstein [ 14].

In [ 9], Karlin and Lee studied the sign consistency properties for periodic boundary-value problems. These results were later used by Lee in [ 15] and [ 16] to develop the spectral properties of these periodic problems. The analysis in [ 15] and [ 16] extends easily to the case of anti-periodic boundary-value problems. Recently, Karlin and Pinkus [ 10] (see also Melkman [ 18]) showed that an important class of mixed boundary-value problems have Green's functions which are sign consistent for all even and/or odd orders. Special examples of this correspond to the above mentioned periodic and anti-periodic boundary-value problems. In this paper, the spectral properties and oscillation theorems associated with the mixed boundary-value problems are developed.

§ 2. Terminology and Preliminary Results

For given  $w_i(x) > 0$ ,  $x \in [0, 1]$ , and  $w_i \in C^{n-i}[0, 1]$ ,  $i = 1, \dots, n$ ,  
define

$$(2.1) \quad L = D_n \cdots D_1$$

where

$$(D_j u)(x) = \frac{d}{dx} \left[ \frac{u(x)}{w_j(x)} \right], \quad j = 1, \dots, n.$$

Thus  $L$  is a differential form of Polya type  $W$  (disconjugate) on  $[0, 1]$ . The differential equation  $Lu = 0$  has a basis of solutions

$$(2.2) \quad \begin{cases} u_1(x) = w_1(x) \\ u_2(x) = w_1(x) \int_0^x w_2(t_1) dt_1 \\ \vdots \\ u_n(x) = w_1(x) \int_0^x w_2(t_1) \int_0^{t_1} w_3(t_2) \cdots \int_0^{t_{n-2}} w_n(t_{n-1}) dt_{n-1} \cdots dt_1 \end{cases}$$

which constitute an extended complete Tchebycheff (ECT) system on  $[0, 1]$  (see [11]). Evidently,

$$D^{j-1} u_i(0) = w_i(0) \delta_{ij}, \quad i, j = 1, \dots, n,$$

where  $D^j = D_j \cdots D_1$ ,  $j = 1, \dots, n$ , and  $D^0 = I$ , the identity operator. The function

$$(2.3) \quad \phi_n(x; \xi) = \begin{cases} 0 & , x < \xi \\ w_1(x) \int_{\xi}^x w_2(t_1) \int_{\xi}^{t_1} w_3(t_2) \cdots \int_{\xi}^{t_{n-2}} w_n(t_{n-1}) dt_{n-1} \cdots dt_1 & , \xi \leq x \end{cases}$$



with  $0 \leq x, \xi \leq 1$  is the fundamental solution for  $Lu = 0$  determined by zero initial data at zero, and the characteristic jump discontinuity

$$D^{n-1} \phi_n(\xi+; \xi) - D^{n-1} \phi_n(\xi-; \xi) = w_n(\xi)$$

which corresponds to the requirement that the ordinary  $(n-1)$ -st derivative of  $\phi_n(x; \xi)$  exhibits a jump of  $1/p_0(\xi)$  at  $x = \xi$ , where  $L = p_0(x)(d^n/dx^n) + \dots$ .

Let  $A$  be an  $m \times n$  matrix and  $r \leq m, n$ . Then  $A \begin{pmatrix} i_1, \dots, i_r \\ j_1, \dots, j_r \end{pmatrix}$  denotes the determinant of the matrix obtained from  $A$  by deleting the rows and columns except for those labeled  $i_1, \dots, i_r$  and  $j_1, \dots, j_r$ , respectively. The matrix  $A$  is said to be sign consistent of order  $r$  ( $SC_r$ ) if

$$\epsilon_r A \begin{pmatrix} i_1, \dots, i_r \\ j_1, \dots, j_r \end{pmatrix} \geq 0$$

for all  $1 \leq i_1 < \dots < i_r \leq m$  and  $1 \leq j_1 < \dots < j_r \leq n$ , where  $\epsilon_r = +1$  or  $-1$ , dependent only upon  $r$ .

The following notation will be used. Given a matrix

$$C = \|A, B\|_{n \times 2n}$$

with both  $A$  and  $B$   $n \times n$  matrices, define

$$(2.4) \quad C^{(p)} = \left\| c_{ij}^{(p)} \right\|_{i=1, j=1}^{n, 2n}$$

by

$$c_{ij}^{(p)} = \begin{cases} a_{ij}(-1)^{j+p+n}, & i = 1, \dots, n; j = 1, \dots, n \\ b_{i, 2n+1-j}, & i = 1, \dots, n; j = n+1, \dots, 2n \end{cases}$$

where  $p \neq 0$  or  $1$ .

The matrix  $C = \|A, B\|$  is said to satisfy Postulate J with respect to  $p$  if

- (i)  $A$  and  $B$  are  $n \times n$  matrices, and
- (ii)  $C^{(p)}$  has full rank and is  $SC_n$ .

Let  $K(x, s)$  be a real-valued kernel defined on  $J \times J$ , where  $J$  is a real interval, and let

$$J_r = \{ \underline{x} : \underline{x} = (x_1, \dots, x_r), x_1 < \dots < x_r, x_j \in J, j = 1, \dots, r \}.$$

The function

$$K_{[r]}(\underline{x}, \underline{s}) = K \begin{pmatrix} x_1, \dots, x_r \\ s_1, \dots, s_r \end{pmatrix} = \det \|K(x_i, s_j)\|_{i,j=1}^r$$

defined on  $J_r \times J_r$  is called the compound kernel of order  $r$  induced by  $K$ . The kernel  $K(x, s)$  is said to be  $SC_r$  if  $\epsilon_r K_{[r]}(\underline{x}, \underline{s}) \geq 0$  for  $\underline{x}, \underline{s} \in J_r$ , where  $\epsilon_r = \pm 1$ , dependent only on  $r$ .

Let  $f_1, \dots, f_r$  be defined on  $J$ . Then

$$(f_1 \wedge \dots \wedge f_r)(\underline{x}) = \det \|f_i(x_j)\|_{i,j=1}^r$$

for  $\underline{x} \in J_r$ . Under suitably mild measurability and integrability assumptions, in particular when  $f_1, \dots, f_r$  are bounded and continuous on  $J$  and  $K(x, s)$  is bounded and continuous on  $J \times J$ , the basic composition



formula [7, p. 17] yields

$$(2.5) \quad K_{[r]}(f_1 \wedge \cdots \wedge f_r) = Kf_1 \wedge \cdots \wedge Kf_r$$

where

$$(Kf)(x) = \int_J K(x, s)f(s)ds ,$$

$$(K_{[r]}g)(\underline{x}) = \int_{J_r} K_{[r]}(\underline{x}, \underline{s})g(\underline{s})d\underline{s} ,$$

for  $g$  defined on  $J_r$  and  $d\underline{s} = ds_1 \cdots ds_r$ .

A family of real, continuous functions  $\{f_1, \dots, f_r\}$  is a Tchebycheff (T) system on  $J$  if  $f_1 \wedge \cdots \wedge f_r$  maintains a fixed strict sign on  $J_r$ , or equivalently, never vanishes on  $J_r$ . Linear combinations of  $f_1, \dots, f_r$  are called  $f$ -polynomials. If  $f$  is a real continuous function on  $J$ , an isolated zero  $x_0$  of  $f$  in the interior of  $J$  is called a nodal zero or node if  $f$  changes sign at  $x_0$ . All other zeros, including zeros at the endpoints of  $J$ , are called nonnodal zeros. We shall use this concept for  $J = (0, 1)$ , the open interval, and thus endpoints will not in fact concern us. Let  $Z(f)$  denote the number of zeros of  $f$  in  $J$  where nonnodal zeros in the interior of  $J$  are counted twice and all other zeros are counted once. Let  $N(f)$  denote the number of (distinct) nodes of  $f$  in  $J$ . If  $\{f_1, \dots, f_r\}$  is a T-system on  $J$ , then any  $f$ -polynomial satisfies  $Z(f) \leq r - 1$ .

The following version of Jentzsch's theorem will be used. We are concerned with the eigenvalue problem

$$(2.6) \quad \phi(x) = \lambda \int_J K(x, s)\phi(s)d\mu(s)$$

for the kernel  $K(x, s)$  where  $d\mu(s) = w(s)ds$  with  $w(s) > 0$  and continuous on  $\bar{J}$ .

Theorem A. Assume the kernel  $K(x, s)$  in (2.6) is nonnegative and continuous on  $\bar{J} \times \bar{J}$  and that  $K(x, s) > 0$  for all (resp., almost all) points  $(x, s)$  in some neighborhood of the diagonal  $\{(x, x) : x \in J\}$ .

Then the kernel  $K(x, s)$  has a positive eigenvalue  $\lambda_0$  which is a simple root of the Fredholm determinant and which is strictly smaller in modulus than all other eigenvalues of  $K(x, s)$ . Furthermore, the corresponding eigenfunction may be chosen positive (resp., positive almost everywhere) on  $J$ .

Remark 2.1. Jentzsch's original proof in [4] may be modified in a straightforward manner to obtain Theorem A. The reasoning in [1] is, however, simpler.

Remark 2.2. Jentzsch's theorem (Theorem A) generalizes directly to kernels defined on the simplices  $J_r$ ,  $r = 2, 3, \dots$ .

There is a fundamental relation, Schur's theorem (Theorem B below), see [21] or [3], between the eigenvalue problem (2.6) for the kernel  $K(x, s)$  and the corresponding eigenvalue problem

$$(2.7) \quad \Phi(\underline{x}) = \Lambda \int_{J_r} K_{[r]}(\underline{x}, \underline{s})\Phi(\underline{s})d\mu(\underline{s})$$

for the  $r$ th compound kernel  $K_{[r]}(\underline{x}, \underline{s})$ , where  $d\mu(\underline{s}) = d\mu(s_1) \cdots d\mu(s_r)$ .

Theorem B. Let  $K(x, s)$  be continuous on  $\bar{J} \times \bar{J}$ . Let  $\lambda_0, \lambda_1, \lambda_2, \dots$   
be the set (possibly empty) of eigenvalues of  $K(x, s)$  where each eigenvalue  
is listed according to its multiplicity as a root of  $D(\lambda)$ , the Fredholm  
determinant of  $K(x, s)$ . Then

$$\lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_r}, \quad 0 \leq i_1 < i_2 < \cdots < i_r$$

are the totality of eigenvalues of  $K_{[r]}(\underline{x}, \underline{s})$ , and each such eigenvalue  
automatically occurs to its multiplicity as a root of  $D_{[r]}(\lambda)$ , the  
Fredholm determinant of  $K_{[r]}(\underline{x}, \underline{s})$ .

Remark 2.3. The assumption that  $K(x, s)$  be continuous on  $\bar{J} \times \bar{J}$   
can be substantially relaxed; however the result as stated is adequate for  
our purposes.



### § 3. Sign Consistent Green's Functions

Karlin and Pinkus [10] consider an important class of boundary value problems with mixed boundary conditions. Examples of these boundary conditions include the separated boundary conditions commonly used in Sturm-Liouville problems as well as periodic and antiperiodic boundary conditions. In this section we summarize the sign consistency results of [10] for certain Green's functions and present some refinements necessary for our analysis of the associated eigenvalue problem. Assume henceforth that  $n > 1$ .

Consider the differential operator  $(L, \mathcal{B})$  specified by a differential form  $L$  of Pólya type  $W$ , see (2.1), and a set of mixed boundary conditions

$$(3.1) \quad U_i(u) = \sum_{j=1}^n a_{ij} D^{j-1} u(0) + \sum_{j=1}^n b_{ij} D^{j-1} u(1) = 0, \quad i = 1, 2, \dots, n.$$

Subsequently,  $\mathcal{B}$  will denote either the set of boundary conditions (3.1), or the set of functions in  $C^n[0,1]$  satisfying these conditions. It is a well-known fact (see e.g. [2, Chap. 7] or [20, Chap. 1]) that the differential operator has a Green's function  $G(x, s)$  iff

$$\Delta = \det \| \| U_i(u_j) \| \|_{i,j=1}^n \neq 0$$

where  $\{u_j\}_{j=1}^n$  are as given in (2.2). If  $\Delta \neq 0$ , then

$$(3.2) \quad G(x, s) = \frac{1}{\Delta} \det \left\| \begin{array}{c|c} \| U_i(u_j) \| \|_{i,j=1}^n & \| U_i[\phi_n(\cdot; s)] \| \|_{i=1}^n \\ \hline u_1(x), \dots, u_n(x) & \phi_n(x; s) \end{array} \right\|_{(n+1) \times (n+1)}$$





Theorem 3.1. If the Green's function  $G(x, s)$  for  $(L, \mathcal{B})$  exists and  
 $C^{(p)}$  is  $SC_n$ , then  $G(x, s)$  is  $SC_{2\ell-p}$  for  $\ell = 1, 2, \dots$ . Further-  
more  $G_{[2\ell-p]}(x, s) \neq 0$  iff the following holds: There exists at  
least one set of indices  $1 \leq j_1 < \dots < j_s \leq n$ ,  $1 \leq k_1 < \dots < k_{n-s} \leq n$   
such that

$$C^{(p)} \begin{pmatrix} 1, & \dots & \dots & \dots & \dots & \dots & n \\ j_1, \dots, j_s, 2n+1-k_{n-s}, \dots, 2n+1-k_1 \end{pmatrix} \neq 0$$

and

(a) if  $\min\{s, n-s\} > 2\ell - p$ , then

$$M_{\mu} + (2\ell - p) \geq \mu, \quad \mu = 2\ell - p + 1, \dots, n$$

while

(b) if  $\min\{s, n-s\} < 2\ell - p$ , then

$$x_{\mu-s} < s_{\mu} < x_{\mu+n-s}, \quad \mu = 1, \dots, 2\ell - p$$

wherever these inequalities are meaningful.

Remark 3.1. Kalafaty in [5] obtains results bearing on Theorem 3.1. It is essentially shown in [5] that if  $C^{(p)}$  is  $SC_n$  then  $G(x, s)$  is  $SC_{2\ell-p}$ ,  $\ell = 1, 2, \dots$ ; however, the precise conditions determining when

$G_{[2\ell-p]}(x, s) \neq 0$  are not treated.

The next result is a consequence of Proposition 3.1 and Theorem 3.1.

Proposition 3.2. Assume  $(L, \mathcal{B})$  has a Green's function  $G(x, s)$ ,  $C^{(p)}$   
is  $SC_n$ , and

$$(3.5) \quad C^{(p)} \begin{pmatrix} 1, & \dots & \dots & \dots & \dots & \dots & n \\ j_1, \dots, j_s, 2n+1-k_{n-s}, \dots, 2n+1-k_1 \end{pmatrix} \neq 0,$$

for some  $1 \leq s \leq n-1$ , and some choice  $\{j_{\ell}\}_1^s, \{k_{\ell}\}_1^{n-s}$  as above. Then

$G_{[2\ell-p]}(x, x) \neq 0$ .

Proof. Since the Green's function exists, (3.4) is valid for some  $0 \leq s \leq n$  and  $M_{\mu} \geq \mu$  for  $\mu = 1, \dots, n$ , by Proposition 3.1. If  $s \neq 0$  or  $n$ , then the result follows from Theorem 3.1, (a) and (b). Assume this is not the case. Thus (3.4) is valid only for  $s = 0$  or  $s = n$ . Assume its validity for  $s = 0$ . Since (3.5) must hold for some  $1 \leq s \leq n - 1$ , it is easily shown that (3.5) is maintained for  $s = 1$ . Appealing again to Theorem 3.1, the result follows.

Remark 3.2. Assume that the Green's function  $G(x, s)$  exists and  $C^{(p)}$  is  $SC_n$ . By Theorem 3.1,  $\sigma_{2\ell-p} G_{[2\ell-p]}(\underline{x}, \underline{s}) \geq 0$  where  $\sigma_{2\ell-p} = \pm 1$ , independent of  $\underline{x}$  and  $\underline{s}$ . In fact, from [10, Remark 3], we see that  $\sigma_{2\ell-p}$  is also independent of  $\ell$ . Explicitly,

$$\sigma_{2\ell-p} = (-1)^{np+n(n-1)/2} \sigma_n(C^{(p)}) \operatorname{sgn}(\Delta)$$

where  $\sigma_n(C^{(p)})$  is the constant sign of the  $n \times n$  minors of  $C^{(p)}$ . If  $C^{(0)}$  is  $SC_n$ , then this further reduces to  $\sigma_{2\ell} = +1$ . If both  $C^{(0)}$  and  $C^{(1)}$  are  $SC_n$ , then

$$\sigma_{2\ell-p} = (-1)^{np} \sigma_n(C^{(p)}) \sigma_n(C^{(0)}) .$$

§ 4. Existence of Eigenvalues

Consider the eigenvalue problem

$$(4.1) \quad \begin{cases} \frac{1}{w_{n+1}} Lu = \lambda u, & 0 \leq x \leq 1 \\ u \in \mathcal{B} \end{cases}$$

where  $w_{n+1}(x) \neq 0$  and continuous on  $[0,1]$ , and  $\mathcal{B}$  is specified as in (3.1). Assume  $(L, \mathcal{B})$  has a Green's function  $G(x, s)$ . Then (4.1) is equivalent to the eigenvalue problem

$$(4.2) \quad u(x) = \lambda \int_0^1 G(x, s) u(s) w_{n+1}(s) ds .$$

The boundary conditions  $\mathcal{B}$  whose matrix is denoted by  $C = \|A, B\|$ , as in Section 3, may be equivalent to a set of initial conditions. In this case the Green's function always exists, is a Volterra kernel, and hence has no eigenvalues. This situation is exceptional for boundary conditions satisfying Postulate J: If the Green's function for  $(L, \mathcal{B})$  exists and the boundary conditions satisfy Postulate J for  $p = 0$  or  $1$ , then  $G(x, s)$  has an infinite number of eigenvalues provided  $\mathcal{B}$  is not equivalent to a set of initial conditions. Thus we shall exclude below the case when  $\mathcal{B}$  is equivalent to a set of initial conditions. The following easily proven result is pertinent.

Proposition 4.1. A set of boundary conditions  $\mathcal{B}$  with matrix  $C = \|A, B\|$  of rank  $n$  is equivalent to a set of initial conditions iff  $A = 0$  or  $B = 0$ .

This result should be considered with Proposition 3.2.



The main result of this section follows.

Theorem 4.1. Assume that the boundary conditions  $\mathcal{B}$  satisfy Postulate J for some  $p$ , are not equivalent to a set of initial conditions, and that  $(L, \mathcal{B})$  has a Green's function  $G(x, s)$ . Then the eigenvalue problem (4.1), equivalently (4.2), has an infinite number of eigenvalues  $\lambda_0, \lambda_1, \lambda_2, \dots$ . Furthermore, if  $|\lambda_0| \leq |\lambda_1| \leq |\lambda_2| \leq \dots$ , where each eigenvalue is listed according to its multiplicity as a root of the Fredholm determinant of the kernel  $G(x, s)$ , then the following holds.

I. If  $C^{(0)}$  is  $SC_n$ , then

$$(1) \quad 0 < |\lambda_0| \leq |\lambda_1| < |\lambda_2| \leq |\lambda_3| < \dots < |\lambda_{2\ell}| \leq |\lambda_{2\ell+1}| < \dots;$$

$$(2) \quad \lambda_{2\ell} \lambda_{2\ell+1} > 0, \quad \ell = 0, 1, 2, \dots;$$

$$(3) \quad \text{also } \lambda_{2\ell} \text{ is not real iff } \lambda_{2\ell+1} = \bar{\lambda}_{2\ell}.$$

II. If  $C^{(1)}$  is  $SC_n$ , then

$$(1) \quad 0 < \hat{\sigma}_1 \lambda_0 < |\lambda_1| \leq |\lambda_2| < \dots < |\lambda_{2\ell-1}| \leq |\lambda_{2\ell}| < \dots \quad \text{where}$$

$$\hat{\sigma}_1 = \text{sgn } G(x, s) w_{n+1}(s);$$

$$(2) \quad \lambda_{2\ell} \lambda_{2\ell-1} > 0, \quad \ell = 1, 2, \dots;$$

$$(3) \quad \text{also } \lambda_{2\ell} \text{ is not real iff } \lambda_{2\ell-1} = \bar{\lambda}_{2\ell}.$$

Proof. Assume  $C^{(p)}$  is  $SC_n$  and  $w_{n+1} > 0$ . Thus  $\sigma_{2\ell-p} G_{[2\ell-p]}(\underline{x}, \underline{s}) \geq 0$

for all  $(\underline{x}, \underline{s}) \in J_{2\ell-p} \times J_{2\ell-p}$ ,  $J = [0, 1]$ , for some  $\sigma_{2\ell-p} = \pm 1$

independent of  $\underline{x}$  and  $\underline{s}$ . Since  $\mathcal{B}$  is not equivalent to a set of initial

conditions, the hypothesis of Proposition 3.2 is easily seen to hold

and so  $\sigma_{2\ell-p} G_{[2\ell-p]}(\underline{x}, \underline{x}) > 0$ .

Jentzsch's theorem (Theorem A) implies that  $\sigma_{2\ell-p} G_{[2\ell-p]}(\underline{x}, \underline{s})$  has a positive simple eigenvalue, strictly smaller in modulus than all other eigenvalues of  $\sigma_{2\ell-p} G_{[2\ell-p]}(\underline{x}, \underline{s})$ . Schur's theorem (Theorem B) then implies that  $G(x, s)$  has at least  $2\ell - p$  eigenvalues,  $\ell = 1, 2, \dots$ . Thus  $G(x, s)$  has infinitely many eigenvalues and again by Theorems A and B

$$0 < \sigma_{2\ell-p} \lambda_0 \cdots \lambda_{2\ell-p-1} < |\lambda_0 \lambda_1 \cdots \lambda_{2\ell-p-2} \lambda_{2\ell-p}|,$$

for  $\ell = 1, 2, \dots$ . Thus,  $|\lambda_{2\ell-p-1}| < |\lambda_{2\ell-p}|$  and  $\lambda_{2\ell-p} \lambda_{2\ell-p+1} > 0$

where we have used the fact that  $\sigma_{2\ell-p} \sigma_{2\ell-p+2} = 1$  (see Remark 3.2).

Note that for  $p = 0$ ,  $\sigma_{2\ell} = +1$  and thus  $\lambda_0 \lambda_1 > 0$ . For  $p = 1$ ,  $\sigma_1 \lambda_0 > 0$ .

Thus both (1) and (2) of Cases I and II obtain. The fact that complex eigenvalues occur in conjugate pairs follows from the fact that the Fredholm determinant of  $G(x, s)$ ,

$$D(\lambda) = 1 + \sum_{r=1}^{\infty} \frac{(-\lambda)^r}{r!} \int_{J_r} G_{[r]}(\underline{x}, \underline{x}) d\mu(\underline{x}),$$

is an entire function with real coefficients. The proof is complete. (When  $w_{n+1} < 0$  replace  $\lambda$  by  $-\lambda$  and apply the results just proven.)

The following examples illustrate the breadth of applicability of Theorem 4.1.

#### Examples.

(a) Periodic Boundary Conditions. For the periodic boundary conditions







(d) Consider the eigenvalue problem

$$(4.3) \quad \begin{cases} -e^x \left( \frac{d}{dx} e^{-2x} \left( \frac{d}{dx} e^x \right) \right) u = \lambda u \\ u(0) = u(1), u'(0) = u'(1) . \end{cases}$$

The differential equation simplifies to  $-u'' + u = \lambda u$  while

$$C^{(p)} = \begin{vmatrix} (-1)^{p+1} & 0 & 0 & -1 \\ (-1)^p & (-1)^p & -1/e & 1 \end{vmatrix} .$$

$C^{(0)}$  is not  $SC_2$  but  $C^{(1)}$  is. Furthermore the Green's function exists.

The eigenvalues for (4.3) are

$$1, 1 + 4\pi^2, 1 + 4\pi^2, \dots, 1 + 4\pi^2 n^2, 1 + 4\pi^2 n^2, \dots .$$

Thus equality can hold in  $\lambda_{2\ell-1} \leq \lambda_{2\ell}$  for each  $\ell$  in Theorem 4.1,

Case II. Replacing the periodic boundary conditions by antiperiodic ones provides a similar example for Case I.

(e) Consider the eigenvalue problem

$$(4.4) \quad \begin{cases} -u'' = \lambda u \\ -u'(0) + u(1) = 0, u'(0) + u'(1) = 0 . \end{cases}$$

Here

$$C^{(p)} = \begin{vmatrix} 0 & (-1)^{p+1} & 0 & 1 \\ 0 & (-1)^p & 1 & 0 \end{vmatrix} ,$$

$C^{(0)}$  is  $SC_2$  while  $C^{(1)}$  is not. The characteristic equation determining

the eigenvalues of (4.4) is

$$(4.5) \quad 1 + \cos \sqrt{\lambda} - \sqrt{\lambda} \sin \sqrt{\lambda} = 0 .$$



Hence

$$\cos \frac{\sqrt{\lambda}}{2} \left( \cos \frac{\sqrt{\lambda}}{2} - \sqrt{\lambda} \sin \frac{\sqrt{\lambda}}{2} \right) = 0,$$

$$\cos(\sqrt{\lambda}/2) = 0 \quad \text{or} \quad \sqrt{\lambda} \tan(\sqrt{\lambda}/2) = 1.$$

The first equation yields eigenvalues

$$v_n = (2n+1)^2 \pi^2, \quad n = 0, 1, 2, \dots,$$

with eigenfunctions,

$$(4.6) \quad v_n(x) = -(2n+1)\pi \cos(2n+1)\pi x + \sin(2n+1)\pi x.$$

The second equation,  $\sqrt{\lambda} \tan(\sqrt{\lambda}/2) = 1$ , has only real zeros because it is the characteristic equation for the self adjoint eigenvalue problem on  $[0, \frac{1}{2}]$

$$\begin{cases} -\varphi'' = \lambda \varphi \\ \varphi'(0) = 0 \\ \varphi'(\frac{1}{2}) + \varphi(\frac{1}{2}) = 0. \end{cases}$$

The zeros  $\mu_0, \mu_1, \mu_2, \dots$  of  $\sqrt{\lambda} \tan(\sqrt{\lambda}/2) = 1$  satisfy

$$2n\pi < \sqrt{\mu_n} < (2n+1)\pi, \quad n = 0, 1, 2, \dots,$$

$$0 < \sqrt{\mu_n} - 2n\pi \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty,$$

and corresponding eigenfunctions are

$$(4.7) \quad u_n(x) = \sqrt{\mu_n} \cos \sqrt{\mu_n} x + \sin \sqrt{\mu_n} x.$$

The totality of eigenvalues of (4.4) is

$$0 < \mu_0 < v_0 < \mu_1 < v_1 < \dots < \mu_n < v_n < \dots$$

which shows that equality need never hold for Theorem 4.1, Case I, even when only  $C^{(0)}$  is  $SC_n$ .

Under the hypotheses of Theorem 4.1 stronger results are obtained when both  $C^{(0)}$  and  $C^{(1)}$  are  $SC_n$ , as is clear from Example (c) (separated boundary conditions). We now prove the rather surprising result that if both  $C^{(0)}$  and  $C^{(1)}$  are  $SC_n$  of rank  $n$ , then the boundary conditions are, in fact, equivalent to separated boundary conditions.

Proposition 4.2. If the matrix  $C$  is of rank  $n$  and both  $C^{(0)}$  and  $C^{(1)}$  are  $SC_n$ , then the boundary conditions  $B$  are equivalent to separated boundary conditions.

Remark 4.1. By the term "equivalent to separated boundary conditions" we mean that the boundary conditions may be rewritten as separated boundary conditions. For example, the boundary conditions  $u(0) = 0$ ,  $u(0) + u(1) = 0$  can be rewritten as  $u(0) = 0$ ,  $u(1) = 0$ .

We prove Proposition 4.2 via two lemmas.

Lemmas 4.1. If the conditions of Proposition 4.2 hold, then there is exactly one  $s$ ,  $0 \leq s \leq n$ , such that

$$(4.8) \quad C \begin{pmatrix} 1, & \dots & \dots & , n \\ j_1, \dots, j_s, k_1, \dots, k_{n-s} \end{pmatrix} \neq 0$$

for some choices of indices  $\{j_i\}_{i=1}^s$  and  $\{k_i\}_{i=1}^{n-s}$  satisfying

$$1 \leq j_1 < \dots < j_s \leq n < k_1 < \dots < k_{n-s} \leq 2n.$$

Proof. Since  $C$  has rank  $n$  there exists at least one  $s$ ,  $0 \leq s \leq n$ , and corresponding indices for which (4.8) holds.



Assume there exists an  $s' > s$  and indices  $\{j'_i\}_{i=1}^{s'}$ ,  $\{k'_i\}_{i=1}^{n-s'}$

ordered as above such that

$$(4.9) \quad C \begin{pmatrix} 1, & \dots, & n \\ j'_1, \dots, j'_{s'}, k'_1, \dots, k'_{n-s'} \end{pmatrix} \neq 0.$$

Let  $C_\ell$  denote the  $\ell$ th column vector of the matrix  $C$ ,  $\ell = 1, \dots, 2n$ .

Then (4.8) and (4.9) imply

$$\{C_{j'_m}\}_{m=1}^{s'} \in \text{span}\{C_{j_1}, \dots, C_{j_s}, C_{k_1}, \dots, C_{k_{n-s}}\}.$$

Since  $s' > s$ , and  $\{C_{j'_m}\}_{m=1}^{s'}$  is a linearly independent set, there

exists an  $m_0 \in \{1, \dots, s'\}$  such that

$$C_{j'_{m_0}} \notin \text{span}\{C_{j_1}, \dots, C_{j_s}\}$$

and, hence, the matrix with column vectors

$$\{C_{j_i}\}_{i=1}^s, C_{j'_{m_0}}, \{C_{k_i}\}_{i=1, i \neq i_0}^{n-s}$$

must be nonsingular for some  $i_0 \in \{1, \dots, n-s\}$ . Ordering and renumbering,

it follows that there exist indices

$$1 \leq \hat{j}_1 < \dots < \hat{j}_{s+1} \leq n < \hat{k}_1 < \dots < \hat{k}_{n-s-1} \leq 2n$$

such that

$$(4.10) \quad C \begin{pmatrix} 1, & \dots, & n \\ \hat{j}_1, \dots, \hat{j}_{s+1}, \hat{k}_1, \dots, \hat{k}_{n-s-1} \end{pmatrix} \neq 0.$$

Then for  $\ell_i = 3n + 1 - k_{n-s+1-i}$ ,  $i = 1, \dots, n-s$ , and  $\hat{\ell}_1 = 3n + 1 - \hat{k}_{n-s-1}$ ,

$i = 1, \dots, n-s-1$ , we have

$$C^{(1)} \begin{pmatrix} 1, & \dots & \dots & , n \\ j_1, \dots, j_s, \ell_1, \dots, \ell_{n-s} \end{pmatrix} = (-1)^s C^{(0)} \begin{pmatrix} 1, & \dots & \dots & , n \\ j_1, \dots, j_s, \ell_1, \dots, \ell_{n-s} \end{pmatrix},$$

$$C^{(1)} \begin{pmatrix} 1, & \dots & \dots & , n \\ \hat{j}_1, \dots, \hat{j}_{s+1}, \hat{\ell}_1, \dots, \hat{\ell}_{n-s-1} \end{pmatrix} = (-1)^{s+1} C^{(0)} \begin{pmatrix} 1, & \dots & \dots & , n \\ \hat{j}_1, \dots, \hat{j}_{s+1}, \hat{\ell}_1, \dots, \hat{\ell}_{n-s-1} \end{pmatrix},$$

with all determinants nonzero by (4.8) and (4.10). This shows that

$C^{(0)}$  and  $C^{(1)}$  cannot both be  $SC_n$  and proves the lemma.

Lemma 4.2. If (4.8) holds for exactly one  $s$ ,  $0 \leq s \leq n$ , then the boundary conditions  $\mathcal{B}$  determined by  $C = \|A, B\|$  are equivalent to separated boundary conditions.

Proof. Let  $\{j_i\}_{i=1}^s$  and  $\{k_i\}_{i=1}^{n-s}$  be ordered indices such that (4.8) holds.

Since the interchange of rows and the addition of linear combinations of one row

to another in  $C$  in no way affects the boundary conditions  $\mathcal{B}$ , we may assume

that the matrix which gives rise to the determinant in (4.8) is the identity,

i.e.,  $C_{j_i} = e_i$  ( $i = 1, \dots, s$ ) and  $C_{k_i} = e_{s+i}$  ( $i = 1, \dots, n-s$ ) where

$e_i$  is the standard  $i$ th coordinate basis vector in  $n$ -space. Let

$j \notin \{j_1, \dots, j_s\}$ ,  $1 \leq j \leq n$ . Then

$$C_j = \sum_{i=1}^s \alpha_i C_{j_i} + \sum_{i=1}^{n-s} \beta_i C_{k_i},$$

and if  $\beta_i \neq 0$  for some  $i$ , the uniqueness of  $s$  is contradicted.

Thus,  $\beta_i = 0$ ,  $i = 1, \dots, n-s$ ,

$$C_j = \sum_{i=1}^s \alpha_i C_{j_i} = \sum_{i=1}^s \alpha_i e_i,$$

and hence  $c_{lj} = 0$  for  $l = s + 1, \dots, n$ . Similarly if  $k \notin \{k_1, \dots, k_{n-s}\}$ ,  $n + 1 \leq k \leq 2n$ , the first  $s$  components of  $C_k$  are zero. Hence the boundary conditions  $\mathcal{B}$  are equivalent to boundary conditions specified by a matrix of the form  $\left\| \begin{array}{c|c} A_1 & 0 \\ \hline 0 & B_1 \end{array} \right\|$  with  $A_1$   $s \times n$  of rank  $s$  and  $B_1$   $(n-s) \times n$  of rank  $n-s$ , i.e.,  $\mathcal{B}$  is equivalent to separated boundary conditions.

Remark 4.2. Assume  $C = \|A, B\|$  is such that  $C^{(p)}$  is  $SC_n$ . Let  $\tilde{C} = \|-A, B\|$  and  $\tilde{\tilde{C}} = \|A, -B\|$ . Then  $\tilde{C}^{(q)}$  and  $\tilde{\tilde{C}}^{(q)}$  are  $SC_n$  where  $p \neq q$ ,  $p, q \in (0, 1)$ . Furthermore, if  $C = \|I, B\|$ , then  $C^{(p)}$  is  $SC_n$  iff  $(-1)^p B$  is totally positive, i.e.,  $SC_k$  for all  $k = 1, \dots, n$  with  $\varepsilon_k = 1$ ,  $k = 1, \dots, n$ . If  $C = \|A, I\|$ , set  $\hat{A} = \|\hat{a}_{ij}\|$ ,  $\hat{a}_{ij} = a_{ij}(-1)^{i+j}$ . Then  $C^{(p)}$  is  $SC_n$  iff  $(-1)^p \hat{A}$  is totally positive.



§ 5. Oscillation Properties of Eigenfunctions

Throughout this section, we shall assume the hypotheses of Theorem 4.1 hold for the eigenvalue problem (4.1). Let

$$\lambda_0, \lambda_1, \lambda_2, \dots$$

be the eigenvalues of (4.1) enumerated as in Theorem 4.1, and

$$(5.1) \quad u_0, u_1, u_2, \dots$$

be a corresponding sequence of independent eigenfunctions and/or generalized eigenfunctions. Suppose  $\lambda_\ell$  is a simple eigenvalue and  $u_\ell$  a corresponding eigenfunction. If  $\lambda_\ell$  is real, choose  $u_\ell$  real. If  $\lambda_\ell$  is nonreal, then  $\bar{\lambda}_\ell$  is also a simple eigenvalue, and  $\bar{u}_\ell$  is chosen for its eigenfunction. If  $\lambda_\ell$  is not simple, it must be real with multiplicity two, say  $\lambda_\ell = \lambda_{\ell+1}$ . Either  $u_\ell$  and  $u_{\ell+1}$  are both eigenfunctions (chosen real) for  $\lambda_\ell = \lambda_{\ell+1}$  or, if a generalized eigenfunction occurs,  $u_\ell$  and  $u_{\ell+1}$  may be chosen real satisfying

$$(5.2) \quad \begin{cases} \lambda_\ell \int_0^1 G(x, s) u_\ell(s) w_{n+1}(s) ds = u_\ell(x) \\ \lambda_\ell \int_0^1 G(x, s) u_{\ell+1}(s) w_{n+1}(s) ds = u_\ell(x) + u_{\ell+1}(x) \end{cases}$$

(see [15, p. 597 eqn. (1.9)]). Throughout this section the sequence (5.1) is assumed to satisfy the preceding conditions.

Theorem 5.1. Let the hypotheses of Theorem 4.1 hold. Then there is a (complex) constant  $\hat{\sigma}_{2\ell-p}$  such that

$$\hat{\sigma}_{2l-p}(u_0 \wedge u_1 \wedge \cdots \wedge u_{2l-p-1}) > 0$$

on  $J_{2l-p}^0$ ,  $J^0 = (0, 1)$ . Furthermore, if  $\lambda_{2j-p}$  is nonreal, let  
 $v_{2j-p} = \operatorname{Re}(u_{2j-p})$  and  $v_{2j-p+1} = \operatorname{Im}(u_{2j-p})$ , while if  $\lambda_k$  is real,  
let  $v_k = u_k$ . Then

$$\{v_0, v_1, \dots, v_{2l-p-1}\}$$

is a T-system on  $(0, 1)$  satisfying the boundary conditions  $\mathcal{B}$ .

Proof. By Schur's Theorem, the eigenvalue of  $G_{[2l-p]}(\underline{x}, \underline{s})$  of minimum modulus is  $\lambda_0 \lambda_1 \cdots \lambda_{2l-p-1}$ . By Jentzsch's Theorem this eigenvalue is real, simple, of sign  $\sigma_{2l-p}$ , and has a real eigenfunction which does not vanish on  $J_{2l-p}^0$ . Since

$$\lambda_0 \lambda_1 \cdots \lambda_{2l-p-1} G_{[2l-p]}(u_0 \wedge \cdots \wedge u_{2l-p-1}) = u_0 \wedge \cdots \wedge u_{2l-p-1},$$

where  $G_{[2l-p]}$  denotes the integral operator with kernel  $G_{[2l-p]}(\underline{x}, \underline{s})$ , and  $u_0 \wedge \cdots \wedge u_{2l-p-1} \neq 0$  because  $u_0, \dots, u_{2l-p-1}$  are linearly independent, it follows that

$$(5.3) \quad \hat{\sigma}_{2l-p}(u_0 \wedge u_1 \wedge \cdots \wedge u_{2l-p-1}) > 0$$

on  $J_{2l-p}^0$  for some constant  $\hat{\sigma}_{2l-p}$ . A short calculation using (5.3) yields

$$\tilde{\sigma}_{2l-p}(v_0 \wedge v_1 \wedge \cdots \wedge v_{2l-p-1}) > 0,$$

where  $\tilde{\sigma}_{2l-p} = (-2i)^r \hat{\sigma}_{2l-p}$ , and  $r$  is the number of pairs of complex conjugate eigenvalues in  $\{\lambda_0, \dots, \lambda_{2l-p-1}\}$ . Consequently

$\{v_0, v_1, \dots, v_{2l-p-1}\}$  is a T-system and the theorem is proven.

The real sequence  $v_0, v_1, v_2, \dots$  (see Theorem 5.1) of real and imaginary parts of the eigenfunctions  $u_0, u_1, u_2, \dots$  belonging to the Green's function  $G(x, s)$  exhibits oscillation properties analogous to those of the classical Sturm-Liouville eigenvalue problem. These properties emanate from the fact that  $\{v_j\}_{j=0}^{2\ell-p-1}$  is a T-system on  $(0, 1)$  for  $\ell = 1, 2, \dots$ , and from certain orthogonality properties of  $v_0, v_1, \dots$ .

Let  $G^*(x, s) = G(s, x)$  be the Green's function for the adjoint problem to (4.1). Then  $G^*(x, s)$  has precisely the same eigenvalues (to multiplicity) as  $G(x, s)$ , and the generalized eigenfunctions of  $G^*(x, s)$ , say  $\{u_\ell^*\}_0^\infty$ , which we choose according to the same conventions set forth at the beginning of this section for  $\{u_\ell\}_0^\infty$ , must satisfy

$$(5.4) \quad \int_0^1 u_\ell(s) \overline{u_m^*(s)} w_{n+1}(s) ds = 0 \quad \text{if } \lambda_\ell \neq \bar{\lambda}_m.$$

Let  $\{v_j^*\}_0^\infty$  be obtained from  $\{u_j^*\}_0^\infty$  just as  $\{v_j\}_0^\infty$  was obtained from  $\{u_j\}_0^\infty$ . Since  $G^*(x, s)$  clearly has the same sign consistency properties as  $G(x, s)$ , the argument of Theorem 5.1 establishes that  $\{v_0^*, v_1^*, \dots, v_{2\ell-p-1}^*\}$ ,  $\ell = 1, 2, \dots$ , is a T-system on  $(0, 1)$  satisfying the boundary conditions adjoint to  $\mathcal{B}$ . Also (5.4) and the fact that nonreal eigenvalues occur in conjugate pairs yields the orthogonality relation:

If  $|\lambda_\ell| \neq |\lambda_m|$ , then

$$(5.5) \quad \int_0^1 v_\ell(s) v_m^*(s) w_{n+1}(s) ds = 0$$



where  $v_l(s)$  [resp.,  $v_m^*(s)$ ] is the real or the imaginary part of an eigenfunction of  $G(x, s)$  [resp.,  $G^*(x, s)$ ] belonging to  $\lambda_l$  [resp.,  $\lambda_m$ ].

Theorem 5.2. Assume that the hypotheses of Theorem 4.1 hold. Then the following obtain.

Case I. If  $C^{(0)}$  is  $SC_n$ , then

(a) for each  $0 \leq k \leq 2l - 1$  the zeros of

$$v = \sum_{j=k}^{2l-1} a_j v_j \quad (\sum a_j^2 > 0, a_j \text{ real})$$

satisfy

$$2\left[\frac{k}{2}\right] \leq N(v) \leq Z(v) \leq 2l - 1;$$

consequently,  $v_{2l-2}$  and  $v_{2l-1}$  have either  $2l - 2$  or  $2l - 1$  nodes in  $(0, 1)$  and no other zeros, and

(b) the nodes of  $v_{2l-2}$  and  $v_{2l-1}$  strictly interlace.

Case II. If  $C^{(1)}$  is  $SC_n$ , then

(a) for each  $0 \leq k \leq 2l$  the zeros of

$$v = \sum_{j=k}^{2l} a_j v_j \quad (\sum a_j^2 > 0, a_j \text{ real})$$

satisfy

$$2\left[\frac{k-1}{2}\right] + 1 \leq N(v) \leq Z(v) \leq 2l;$$

consequently  $v_{2l-1}$  and  $v_{2l}$  have either  $2l - 1$  or  $2l$  nodes in  $(0, 1)$  and no other zeros. Also  $v_0$  has no zeros in  $(0, 1)$ , and

(b) the nodes of  $v_{2l-1}$  and  $v_{2l}$  strictly interlace.

Remark 5.1. In case  $G(x, s)$  has real spectrum, in particular if the eigenvalue problem is self-adjoint,  $v_j = u_j$  for  $j = 0, 1, 2, \dots$  and Theorem 5.2 describes the oscillation properties of the eigenfunctions of  $G(x, s)$ . See also Theorem 5.3 below.

Proof. We shall prove only Case I. The proof of Case II follows in an entirely analogous manner. Recall that  $N$  and  $Z$  count zeros in  $(0, 1)$ .

The inequality  $Z(v) \leq 2l - 1$  holds because  $\{v_0, v_1, \dots, v_{2l-1}\}$  is a T-system (Theorem 5.1) on  $(0, 1)$ . The inequality for  $N(v)$  follows

from the orthogonality relation (5.5): Suppose that  $v = \sum_{j=k}^{2l-1} a_j v_j$   
 ( $\sum a_j^2 > 0$ ,  $a_j$  real) has nodes

$$0 < \xi_1 < \dots < \xi_r < 1, \quad r = N(v).$$

1. If  $r$  is odd,  $r = 2m - 1$  say, form a nontrivial  $v^*$ -polynomial

$$v^*(x) = \sum_{j=0}^{2m-1} b_j v_j^*(x)$$

which has nodal zeros (sign changes) at  $\xi_1, \dots, \xi_{2m-1}$ . Since  $\{v_0^*, \dots, v_{2m-1}^*\}$  is a T-system,  $v^*$  has no other zeros in  $(0, 1)$ . By construction

$$\int_0^1 v(s) v^*(s) w_{n+1}(s) ds \neq 0.$$

However, if  $k > r = 2m - 1$ , then since  $|\lambda_{2m-1}| < |\lambda_{2m}|$  (Theorem 4.1) the orthogonality conditions (5.5) imply that the above integral is zero, a contradiction. Thus,  $k \leq r = N(v)$ .

2. If  $r$  is even,  $r = 2m$  say, form the  $v^*$ -polynomial

$$v^{*(n)}(x) = \sum_{j=0}^{2m+1} b_j^{(n)} v_j^*(x)$$

which has nodal zeros at  $\xi_0^{(n)}, \xi_1, \dots, \xi_{2m}$  ( $0 < \xi_0^{(n)} < \xi_1$ ) and satisfies

$$\sum_{j=0}^{2m+1} |b_j^{(n)}| = 1. \quad v^{*(n)} \text{ has no additional zeros in } (0, 1) \text{ by the}$$

Tchebycheff property of  $\{v_0^*, \dots, v_{2m+1}^*\}$ . Let  $\xi_0^{(n)} \downarrow 0$  as  $n \uparrow \infty$ .

A simple compactness argument implies the existence of a  $v^*$ -polynomial

$$v^*(x) = \sum_{j=0}^{2m+1} b_j v_j^*(x), \quad \sum_{j=0}^{2m+1} |b_j| = 1,$$

which has nodal zeros at  $\xi_1, \dots, \xi_{2m}$ . Since any sign change of  $v^*$  in  $(0, 1)$  is the limit of zeros of the  $v^{*(n)}$ 's as  $n \rightarrow \infty$  through an appropriate subsequence, and  $Z(v^*) \leq 2m$ , it follows that  $v^*$  has only the zeros  $\xi_1, \dots, \xi_{2m}$  in  $(0, 1)$ . As above, a contradiction follows if  $2m + 1 = r + 1 < k$ . Thus,  $k - 1 \leq N(v)$ .

Thus if  $N(v)$  is odd,  $N(v) \geq k$ , while if  $N(v)$  is even,  $N(v) \geq k - 1$ . This is equivalent to  $N(v) \geq 2[k/2]$ . The remaining part of Case I(a) follows from the first part and the fact that  $Z(v) - N(v)$  is always a nonnegative even integer.

The proof of Case I(b) is set forth as a sequence of lemmas. The arguments follow those used by Gantmacher and Krein in [3, pp. 215-217] with suitable modifications required by our weaker hypotheses. As an immediate consequence of Case I(a), we have, since  $Z(v) - N(v)$  is always even,



Lemma 5.1. The zeros of the v-polynomial

$$v = av_{2l-2} + bv_{2l-1} \quad (a^2 + b^2 > 0)$$

satisfy

$$2l - 2 \leq N(v) \leq Z(v) \leq 2l - 1.$$

Hence, v has only nodal zeros in (0, 1).

In what follows, let  $\phi$  denote either  $v_{2l-2}$  or  $v_{2l-1}$  while  $\psi$  denotes  $v_{2l-1}$  or  $v_{2l-2}$ .

Lemma 5.2. Let  $0 < \xi_1 < \dots < \xi_r < 1$ ,  $r = N(\phi)$ , be the nodes of  $\phi$  in (0, 1). Then the function

$$h = \psi/\phi$$

is strictly monotone on  $I_i = (\xi_i, \xi_{i+1})$ ,  $i = 0, 1, \dots, r$ ;  $\xi_0 = 0$ ,  $\xi_{r+1} = 1$ .

Proof. From Lemma 5.1,  $h$  cannot be constant on any interval of positive length. Thus, if  $h$  is not strictly monotone on  $I_i$ ,  $h$  has a relative extremum at some point  $x_i$  of  $I_i$ . However, this would imply that the v-polynomial  $\psi(x) - h(x_i)\phi(x)$  has a nonnodal zero at  $x_i$ , contradicting Lemma 5.1. Thus  $h$  is strictly monotone.

Lemma 5.3.  $h(x)$  has a zero in each  $I_i$ ,  $i = 1, \dots, r - 1$ .

Proof. Since  $h(x)$  is monotone in each  $I_i$ ,  $i = 0, 1, \dots, r$ , the limits

$$\lim_{x \rightarrow \xi_i^-} h(x) = l_i^- \quad \text{and} \quad \lim_{x \rightarrow \xi_i^+} h(x) = l_i^+$$

both exist as extended real numbers for  $i = 1, \dots, r$ . We shall show that none of the  $\{l_i^+\}_{i=1}^r$  and  $\{l_i^-\}_{i=1}^r$  is finite.

Neither  $l_i^-$  nor  $l_i^+$  is finite when  $\xi_i$  is not a zero of  $\psi$ .

We are concerned with one of the following four cases which may occur only if  $\xi_i$  is a zero of  $\psi$ .

- (i) Exactly one of  $l_i^-$  and  $l_i^+$  is finite.
- (ii) Both  $l_i^-$  and  $l_i^+$  are finite and unequal.
- (iii)  $l_i^- = l_i^+$  (finite) and  $h$  is monotone near  $\xi_i$ .
- (iv)  $l_i^- = l_i^+$  (finite) and  $h$  is monotone in opposite senses for  $x < \xi_i$  and  $x > \xi_i$  but near  $\xi_i$ .

We show that (i) - (iv) are incompatible with Lemma 5.1. If (i) or (ii) holds choose  $c$  in the open interval determined by  $l_i^-$  and  $l_i^+$ , while if (iii) holds set  $c = l_i^- = l_i^+$ . Then the  $v$ -polynomial  $v = \psi - c\phi$  has a nonnodal zero at  $\xi_i$  as is easily seen from the fact that  $\phi(\xi_i) = \psi(\xi_i) = 0$  and  $\phi$  changes sign at  $\xi_i$ .

To contradict (iv), assume  $h \leq c = l_i^+ = l_i^-$  in  $I_{i-1}$  and  $I_i$ . Consider the polynomial  $v_\epsilon = \psi - (c - \epsilon)\phi$ ,  $\epsilon > 0$ . The polynomial  $v = \psi - c\phi$  has at least  $2l - 2$  nodes in  $(0, 1)$  one of which is at  $\xi_i$ . Since  $v_\epsilon = v + \epsilon\phi$ , for  $\epsilon > 0$  sufficiently small,  $v_\epsilon$  must maintain at least  $2l - 3$  nodes in  $(0, 1)$  bounded away from  $\xi_i$ . However, by construction it is easily seen that for  $\epsilon > 0$ , sufficiently small,  $v_\epsilon$  has two "new" nodes (one less than  $\xi_i$  and one greater than  $\xi_i$ ) as well as the one at  $\xi_i$ . Thus for  $\epsilon > 0$ , sufficiently small,  $v_\epsilon$  has at least  $2l$  nodes, a contradiction.

Lemma 5.4. The nodes of  $v_{2\ell-2}$  and  $v_{2\ell-1}$  strictly interlace in  $(0,1)$ .

Proof. If  $N(v_{2\ell-1}) \neq N(v_{2\ell-2})$  then since  $|N(v_{2\ell-2}) - N(v_{2\ell-1})| = 1$ , the result is an immediate consequence of Lemma 5.3. Assume

$N(v_{2\ell-2}) = N(v_{2\ell-1})$  and  $\ell > 1$ . If  $v_{2\ell-2}$  and  $v_{2\ell-1}$  have a common node  $\xi$ , then if  $\eta$  is an adjacent node of  $v_{2\ell-1}$  (such a node exists because  $\ell > 1$ ),  $v_{2\ell-2}$  must have a node  $\zeta$  between  $\xi$  and  $\eta$  by Lemma 5.3. But  $v_{2\ell-1}$  must then have a node between  $\zeta$  and  $\xi$ , contradicting the definition of  $\eta$ . Thus the lemma holds for  $\ell > 1$ .

Assume  $\ell = 1$  so that  $N(v_0), N(v_1) \leq 1$ . If  $v_0$  and  $v_1$  have a common node  $\xi$ , then  $h = v_0/v_1$  has equal, infinite (see the proof of Lemma 5.3) left and right limits at  $\xi$ . Thus there exists a constant  $c$  such that  $v_0 - cv_1$  has three nodes, a contradiction.

This completes the proof of Theorem 5.2.

Examples.

(a) Consider the eigenvalue problem

$$\begin{cases} -u'' = \lambda u \\ u(0) + u(1) = 0 \\ u'(0) + u'(1) = 0. \end{cases}$$

These boundary conditions are antiperiodic and the problem has eigenvalues  $\lambda_n = (2n+1)^2 \pi^2$ ,  $n = 0, 1, 2, \dots$ , each with multiplicity two and corresponding eigenfunctions  $\cos(2n+1)\pi x$  and  $\sin(2n+1)\pi x$ . The interlacing properties guaranteed by Theorem 5.2, Case I, are easily verified in this case.



(b) Consider the eigenvalue problem

$$\begin{cases} -u'' + u = \lambda u \\ u(0) - u(1) = 0 \\ u'(0) - \frac{1}{2}u'(1) = 0. \end{cases}$$

Then (see example (d) of Section 4)

$$C^{(p)} = \begin{vmatrix} (-1)^{p+1} & 0 & 0 & -1 \\ (-1)^p & (-1)^p & -\frac{1}{2e} & \frac{1}{2} \end{vmatrix},$$

and so  $C^{(1)}$  is  $SC_2$  but  $C^{(0)}$  is not. The eigenvalues of this problem are  $\lambda_n = 1 + 4\pi^2 \left[ \frac{n+1}{2} \right]^2$ ,  $n = 0, 1, 2, \dots$ . The eigenvalue

$\lambda_0 = 1$  has eigenfunction  $u_0(x) = 1$  and the double eigenvalue

$\lambda_{2n} = \lambda_{2n-1}$ ,  $n \geq 1$ , has an eigenfunction and generalized eigenfunction given, respectively, by

$$u_{2n-1}(x) = \cos 2\pi nx$$

$$u_{2n}(x) = (x+1)\sin 2\pi nx.$$

The interlacing properties of Theorem 5.2, Case II, are easily confirmed.

This example also shows that generalized eigenfunctions can occur in Theorem 5.2.

(c) In the eigenvalue problem (4.4),  $C^{(0)}$  is  $SC_2$  while  $C^{(1)}$  is not.

The eigenfunctions given in (4.6) and (4.7), when properly ordered,

must have interlacing zeros as in Theorem 5.2, Case I. In this case,

Theorem 5.2 seems to be the easiest way to verify the interlacing of the nodes.

For completeness, we include the following consequence of Theorems 4.1, 5.1 and 5.2 which pertains to the case of separated boundary conditions (Proposition 4.2), (see [6], [12], and [13]).

Theorem 5.3. Assume that the hypotheses of Theorem 4.1 hold for both  
 $p = 0$  and  $p = 1$ . Then the eigenvalues of  $G(x, s)$  are all real and  
simple, and the eigenfunctions  $\{u_\ell\}_0^\infty$  satisfy:

- (1)  $\{u_0, u_1, \dots, u_\ell\}$  is a T-system on  $(0, 1)$  satisfying the boundary  
conditions  $\mathcal{B}$  for each  $\ell = 0, 1, 2, \dots$
- (2) For  $0 \leq k \leq \ell$ , the zeros of

$$u = \sum_{j=k}^{\ell} a_j u_j \quad (\sum a_j^2 > 0, a_j \text{ real})$$

satisfy

$$k \leq N(u) \leq Z(u) \leq \ell.$$

- (3) The eigenfunction  $u_\ell$  has exactly  $\ell$  nodes in  $(0, 1)$  and no other  
zeros there, and the nodes of  $u_\ell$  and  $u_{\ell+1}$  strictly interlace in  
 $(0, 1)$ .

§ 6. Composite Differential Operators

Many important differential operators are naturally expressed as compositions of lower order operators. The applicability of the results of Sections 3-5 to such composite operators may be inferred by an examination of its factors, as is shown in Theorem 6.1. Theorem 6.2 refines the results of Theorem 6.1 in the important special case where the differential term is self-adjoint.

Let  $(L, \mathcal{B})$  be an  $n$ th order differential form of Pólya type  $W$  with boundary conditions specified by the matrix  $C = \|A, B\|$ , as in the previous sections. Moreover, we shall also assume that

$$w_j \in C^{n+m+1-j}[0, 1], \quad j = 1, \dots, n.$$

Likewise, let  $(L^+, \mathcal{B}^+)$  be an  $m$ th order differential form where

$$L^+ = D_m^+ \cdots D_1^+, \quad (D_j^+ u)(x) = \frac{d}{dx} \left[ \frac{u(x)}{w_j^+(x)} \right], \quad j = 1, \dots, m,$$

and  $w_j^+(x) > 0$ ,  $x \in [0, 1]$ ,  $w_j^+ \in C^{m+1-j}[0, 1]$ , with associated boundary conditions determined by the matrix

$$C^+ = \|A^+, B^+\|_{m \times 2m}$$

of rank  $m$ .

Let  $N = L^+L$ , and  $\mathcal{B}^+\mathcal{B}$  be the set of boundary conditions

$$\left\{ \begin{array}{l} \sum_{j=1}^n a_{ij} D_j^{j-1} u(0) + \sum_{j=1}^n b_{ij} D_j^{j-1} u(1) = 0, \quad i = 1, \dots, n \\ \sum_{j=1}^m a_{i-n, j}^+ D_j^{n+j-1} u(0) + \sum_{j=1}^m b_{i-n, j}^+ D_j^{n+j-1} u(1) = 0, \quad i = n+1, \dots, n+m \end{array} \right.$$



specified by the matrix

$$E = \begin{pmatrix} A & 0 & B & 0 \\ 0 & A^+ & 0 & B^+ \end{pmatrix}_{(n+m) \times 2(n+m)},$$

where  $D_{n+j} = D_j^+$ ,  $j = 1, \dots, m$ , and  $D^j = D_j \cdots D_1$ ,  $j = 1, \dots, n+m$ .

Theorem 6.1. Assume  $C^{(p)}$  and  $C^{+(p)}$  satisfy Postulate J for the same  $p$  and that either  $\mathcal{B}$  or  $\mathcal{B}^+$  is not equivalent to a set of initial conditions. If  $(N, \mathcal{B}^+ \mathcal{B})$  has a Green's function, then the conclusion of Theorems 4.1, 5.1 and 5.2 obtain for the eigenvalue problem

$$\begin{cases} \frac{1}{w} Nu = \lambda u \\ u \in \mathcal{B}^+ \mathcal{B} \end{cases}$$

where  $w(x)$  is continuous and non-vanishing on  $[0, 1]$ . Also,

$(N, \mathcal{B}^+ \mathcal{B})$  has a Green's function  $H(x, s)$  iff both  $(L, \mathcal{B})$  and  $(L^+, \mathcal{B}^+)$

have Green's functions, say  $G(x, s)$  and  $G^+(x, s)$ , respectively, in

which case

$$(6.1) \quad H(x, s) = \int_0^1 G(x, t) G^+(t, s) dt.$$

Proof. The proof is a result of the application of the basic composition formula [7, p. 17]. From (6.1),

$$H_{[2l-p]}(\underline{x}, \underline{s}) = \int_{I_{2l-p}} G_{[2l-p]}(\underline{x}, \underline{t}) G_{[2l-p]}^+(\underline{t}, \underline{s}) dt.$$

Since  $G(x, s)$  and  $G^+(x, s)$  are  $SC_{2l-p}$ ,  $l = 1, 2, \dots$ , so is  $H(x, s)$ .

The fact that  $H_{[2l-p]}(\underline{x}, \underline{x}) \neq 0$  follows in a similar fashion.

Remark 6.1. It should be noted that one can also prove directly that under the above assumptions,  $E^{(p)}$  is  $SC_{n+m}$  and of rank  $n+m$ .

An important case of Theorem 6.1 is when  $L^+ = L^*$  and  $B^+ = B^*$ , the adjoint differential form and boundary conditions to  $L$  and  $B$ , respectively. Let  $C = \|A, B\|$  denote the boundary conditions associated with  $(L, B)$ , and  $C_* = \|A_*, B_*\|$  denote the adjoint boundary conditions associated with  $(L^*, B^*)$ . While the analysis is rather lengthy, an explicit form of  $C_*$  may be exhibited and, as is shown in [17] (see also [19]), if  $C^{(p)}$  is  $SC_n$  and of rank  $n$ , then  $C_*^{(p)}$  is  $SC_n$  and of rank  $n$ . From Theorem 6.1 we obtain the following.

Theorem 6.2. Assume  $(L, B)$  has a Green's function  $G(x, s)$ , and  $C^{(p)}$  satisfies Postulate J. Then the self-adjoint differential operator  $(L^* L, B^* B)$  has a Green's function  $H(x, s)$  which is  $SC_{2l-p}$  for  $l = 1, 2, \dots$ . Thus the conclusions of Theorems 4.1, 5.1 and 5.2 apply and since  $H(x, s)$  is symmetric and positive definite, the spectrum of the associated eigenvalue problem is positive.

§7. Extensions

The spectral results for the eigenvalue problem (4.1) developed in Sections 4 and 5 can be cast in a more general setting as described below. No proofs will be given because the reasoning used in Sections 4 and 5 can be applied with inessential changes.

Let  $J = (0, 1)$  and  $K(x, s)$  be a real, continuous kernel on  $\bar{J} \times \bar{J}$  for which

$$(7.1) \quad \begin{cases} \varepsilon_{2\ell-p} K_{[2\ell-p]}(\underline{x}, \underline{s}) \geq 0 & \text{for } \underline{x}, \underline{s} \in J_{2\ell-p} \\ \varepsilon_{2\ell-p} K_{[2\ell-p]}(\underline{x}, \underline{x}) > 0 & \text{for } \underline{x} \in J_{2\ell-p} \end{cases}$$

where  $p = 0$  or  $1$ ,  $\varepsilon_{2\ell-p} = +1$  or  $-1$  dependent only on  $2\ell - p$ , and  $\ell = 1, 2, \dots$ . Consider the eigenvalue problem

$$(7.2) \quad \phi(x) = \lambda \int_J K(x, s) \phi(s) d\mu(s)$$

for the kernel  $K(x, s)$  where  $d\mu(s) = w(s)ds$  and  $w$  is a positive, continuous weight function on  $\bar{J}$ .

Theorem 7.1. Assume (7.1) holds for  $p = 0$  or  $1$ . Then the eigenvalue problem (7.2) has an infinite sequence of eigenvalues  $\lambda_0, \lambda_1, \lambda_2, \dots$ .

Moreover, if  $|\lambda_0| \leq |\lambda_1| \leq |\lambda_2| \leq \dots$  where each eigenvalue is listed according to its multiplicity as a root of the Fredholm determinant of  $K(x, s)$ , then the following holds.

I. If  $p = 0$ , then

$$(1) \quad 0 < |\lambda_0| \leq |\lambda_1| < |\lambda_2| \leq |\lambda_3| < \dots < |\lambda_{2\ell}| \leq |\lambda_{2\ell+1}| < \dots;$$



$$(2) \quad \varepsilon_{2\ell} \varepsilon_{2\ell+2} \lambda_{2\ell} \lambda_{2\ell+1} > 0 \quad \text{for } \ell = 0, 1, 2, \dots \quad \text{where } \varepsilon_0 = +1;$$

$$(3) \quad \text{also } \lambda_{2\ell} \text{ is nonreal iff } \lambda_{2\ell+1} = \overline{\lambda_{2\ell}}.$$

II. If  $p = 1$ , then

$$(1) \quad 0 < \varepsilon_1 \lambda_0 < |\lambda_1| \leq |\lambda_2| < \dots < |\lambda_{2\ell-1}| \leq |\lambda_{2\ell}| < \dots;$$

$$(2) \quad \varepsilon_{2\ell-1} \varepsilon_{2\ell+1} \lambda_{2\ell-1} \lambda_{2\ell} > 0, \quad \ell = 1, 2, \dots;$$

$$(3) \quad \text{also } \lambda_{2\ell} \text{ is nonreal iff } \lambda_{2\ell-1} = \overline{\lambda_{2\ell}}.$$

Parts (2) and (3) of Cases I and II yield the following interesting result guaranteeing that the spectrum of the kernel  $K(x, s)$  is real.

Corollary 7.1. If in Theorem 7.1 - I,  $\varepsilon_{2\ell} \varepsilon_{2\ell+2} < 0$  for  $\ell = 0, 1, 2, \dots$

(i.e., if successive compounds in (7.1) alternate in sign) then  $K(x, s)$

has only real spectrum. Likewise if  $\varepsilon_{2\ell-1} \varepsilon_{2\ell+1} < 0$  for  $\ell = 1, 2, \dots$

in Theorem 7.1 - II, then  $K(x, s)$  has only real spectrum.

On the basis of Theorem 7.1 and the reasoning of Section 5, we can establish the analogues of Theorems 5.1 and 5.2 for the kernel  $K(x, s)$ . Let  $\{u_i\}_0^\infty$  be the set of eigenfunctions of  $K(x, s)$  corresponding to  $\{\lambda_i\}_0^\infty$  and chosen as in Section 5. Let  $\{v_i\}_0^\infty$  be the real sequence constructed from the real and imaginary parts of the eigenfunction  $\{u_i\}_0^\infty$  as in Theorem 5.1. Then

$$\{v_0, v_1, \dots, v_{2\ell-p-1}\}$$

is a T-system on  $(0, 1)$  for  $\ell = 1, 2, \dots$ , and when  $p = 0$  (resp.,

$p = 1$ ) the conclusions of Theorem 5.2 - I (resp., II) hold for the sequence

$$\{v_i\}_0^\infty.$$

#### REFERENCES

- [1] Anselone, P. M., and J. W. Lee, Spectral Properties of Integral Operators with Nonnegative Kernels, *Linear Algebra and Appl.* 9 (1974), 67-87.
- [2] Coddington, E. A., and N. Levinson, The Theory of Ordinary Differential Equations, McGraw-Hill, New York, 1955.
- [3] Gantmacher, F. R., and M. G. Krein, Oszillationsmatrizen, Oszillationskerne und kleine Schwingungen mechanischer Systeme, Akademie Verlag, Berlin, 1960.
- [4] Jentzsch, R., Über Integralgleichungen mit positivem Kern, *J. Math. Crelle* 141 (1912), 235-244.
- [5] Kalafaty, P., Sur les fonctions de Green des equations differentielles ordinaires, *C. R. (Dokl.) Acad. Sci. URSS* 26 (1940), 526-530.
- [6] Karlin, S., Total Positivity, Interpolation by Splines, and Green's Functions of Differential Operators, *J. Approximation Theory* 4 (1971), 91-112.
- [7] Karlin, S., Total Positivity, Vol. I, Stanford University Press, Stanford, Calif., 1968.
- [8] Karlin, S., Total Positivity, Vol. II, in preparation.
- [9] Karlin, S., and J. W. Lee, Periodic Boundary-value Problems and Cyclic Totally Positive Green's Functions with Applications to Periodic Spline Theory, *J. Differential Equations* 8 (1970), 374-396.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1683 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) SPECTRAL PROPERTIES AND OSCILLATION THEOREMS FOR MIXED BOUNDARY-VALUE PROBLEMS OF STURM-LIOUVILLE TYPE ✓	5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) J. W. Lee and A. Pinkus	8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 ✓	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin ✓ Madison, Wisconsin 53706	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709	12. REPORT DATE September 1976	
	13. NUMBER OF PAGES 42	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Mixed boundary conditions Green's function Tchebycheff systems		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  This paper presents analogues of the spectral and oscillatory properties exhibited by classical Sturm-Liouville systems for certain mixed boundary value problems that include periodic boundary conditions as a special case. These mixed boundary-value problems have Green's functions which are sign consistent for all even and/or odd orders.		



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1683	2. GOVT ACCESSION NO. MRC-TSR-1683	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) SPECTRAL PROPERTIES AND OSCILLATION THEOREMS FOR MIXED BOUNDARY-VALUE PROBLEMS OF STURM-LIOUVILLE TYPE.		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) J. W./Lee and A./Pinkus		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1243 p.
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE Sep 1976
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 42
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 9. Technical summary rept.,		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Mixed boundary conditions Green's function Tchebycheff systems		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) → This paper presents analogues of the spectral and oscillatory properties exhibited by classical Sturm-Liouville systems for certain mixed boundary value problems that include periodic boundary conditions as a special case. These mixed boundary-value problems have Green's functions which are sign consistent for all even and/or odd orders.		

mt