SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered) READ INSTRUCTIONS BEFORE COMPLETING FORM REPORT DOCUMENTATION PAGE I. HEPOHT NUMBER 2. GOVT ACCESSION NO. 3 .- RECIPIENT'S CATALOG NUMBER TITLE (and Subilite) TYRE OF REPORT A RERIOD COVERED FINAL REPORT STUDIES ON MICROBIAL PROPAGATION IN THE Feb 1973 - Oct 1976 AIRBORNE STATE 6. PERFORMING ORG. REPORT NUMPER AUTHORIAL 8. CONTRACT OR GRANT NUMBER(+) R.L. (DIMMICK, H. WOLOCHOW and N00014-75-C-1133 M.A. /CHATIGNY - 🗵 PERFORMING ORGANIZATION NAME AND ADDRESS 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Naval Biosciences Laboratory Sch.Pub.Hlth, University of California Berkeley, California 94720 11. CONTROLLING OFFICE NAME AND ADDRESS 12. REPORT DATE Office of Naval Research (Code 443) П 29 Nov 800 N. Quincy Street 13. NUMBER OF PAGES Arlington, Virginia 22217 14. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office) 15. SECURITY CLASS. (of this report) Unclassified 154. DECLASSIFICATION/DOWNGRADING 16. DISTRIBUTION STATEMENT (of this Report) Scientific Officer, ONR Branch Office-1/0; ACO-1/0; NRL Code 2627 1/0; ONR Çode 1021P-6/0; DDC-12/0; NASA Office of Planetary Quarantine.  $3^*/0$ DUNDERT: CON SECTION A Α. thease: bettau 17. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, If different the Report 18. SUPPLEMENTARY NOTES Reprints of appendices included with report. 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Aerosol. Planetary quarantine. Bacterial growth, Jupiter, Particulate ejection Serratia marcescens, Airborne particles, ABSTRACT (Continue on reverse side if necessary and identify by block number) The purpose of this project was to conduct studies to determine whether bacteria, in or on small particles dispersed as aerosols, could maintain growth processes to the extent that more than one division step could occur. The question is important with respect to estimating the probability of contaminating the biological zone of Jupiter if space vehicles were to penetrate the planet. -(Gentinued) DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE UNCLASSIFTED 5/N 0102-014-6601 : SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered) 990 G

UNCLASSIFIED

LUNRITY CLASSIFICATION OF THIS PAGE (When Data Entered)

An additional task was to determine whether metallic surfaces, simulating landing vehicle surfaces, would discharge attached microbes as a result of temperature changes.

The following findings, in chronological order, resulted from this study:

A. In the airborne state:

- 1. Bacteria ingested labelled glucose and produced labelled CO<sub>2</sub>.
- 2. Bacteria ingested labelled thymidine and produced labelled DNA.
- 3. Phage was able to penetrate bacteria and, in one instance, additional phage appeared to have been formed.
- Bacteria doubled in numbers when enclosed in droplets in the 1 - 3 μm diameter range.
- 5. Bacteria almost trebled in numbers when enclosed in droplets in the 3 6 µm range.

B. On a simulated Lander structure bacteria were found to be ejected from surfaces as a result of mechanical stress caused by temperature changes.

We conclude that, in an appropriate environment, bacteria could live indefinitely in the airborne state.

# UNCLASSIFIED

# NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

# FINAL REPORT

In fulfillment of Contract

#N00014-75-C-1133

R. L. Dimmick, H. Wolochow and M. A. Chatigny

STUDIES ON MICROBIAL PROPAGATION

IN THE AIRBORNE STATE

Naval Biosciences Laboratory 100 8113 School of Public Health, University of California 926 Berkeley, California 94720

Final Report N00014-75-C-1133 Page 2

## STUDIES ON MICROBIAL PROPAGATION

# IN THE AIRBORNE STATE

R.L. Dimmick, H. Wolochow and M.A. Chatigny

Naval Biosciences Laboratory School of Public Health, University of California Berkeley, California 94720

# 1. Objectives of Project

The purpose of this project was to investigate the possibility that microbes, airborne in or on small particles (1  $\mu$ m to 100  $\mu$ m diameter) could undergo cellular division under any circumstances. The study was initiated to gain additional information about the probability that the gaseous atmosphere of Jupiter could be contaminated by spacecraft from Earth. Projects associated with the major objective were to investigate:

- (a) The possibility that, if cellular division occurred in air, it could be induced to continue for more than one generation,
- (b) The possibility that airborne microbes could absorb nutrients from air,
- (c) The possibility that anaerobic microbes could sustain functions similar to aerobes,
- (d) The possibility that small microbial particles could be "shed" from metallic surfaces that were undergoing extreme temperature shifts,
- (e) The possibility the the "lifetime" of a particle in the turbulent atmosphere of Jupiter might be sufficient to allow continued propagation.

## 2. Approach

Since the aerobiological literature of the past 30 years contained no reports of workers having observed evidence of division of cells in aerosol particles, the initial approach was to search for indirect evidence that might be immediately productive and that might provide information

Final Report N00014-75-C-1133 Page 3

to guide planning of direct, but more difficult experiments.

Proposed steps in an "If --, then --" sequence (with cells in the airborne state), were:

1. Find evidence of metabolism;

and the second second

- 2. Find evidence of formation of new DNA;
- 3. Find evidence of competent genetic structure and function;
- 4. Find direct evidence of cell division;
- 5. Find indirect evidence for continued propagation (gravitational forces limits the time available).
- 6. Find direct evidence for continued propagation.

Major Accomplishments: Studies with Airborne Aerobic Bacteria

- 1. Active metabolic functions have been demonstrated (Appendices A, B).
- 2. Formation of new DNA has been demonstrated (Appendix C).
- 3. Marginally significant evidence for genetic integrity as shown by formation of phage has been found (Appendix D).
- 4. At least one generation of new cells has been demonstrated to form in small particles (average diameter 2  $\mu$ m) (Appendix E).
- Slightly more than two generations of new cells has been shown to form in large particles (4 to 6 µm diam.)(Appendix F).
- 6. Cup-shaped, metallic objects have been shown to release attached microbes in the form of small particles (Appendix G).

Minor Accomplishments, or those that have not shown affirmative data, include:

1. A method was developed to increase the coagulation rate of airborne particles (Appendix H).

Final Report N00014-75-C-1133

Page 4

- 2. Preliminary data from experiments in a large aerosol chamber indicated that particles decreased according to laws of stirred settling.
- 3. No evidence of metabolism of vegetative cells of anaerobes as aerosols in an anaerobic gas  $(N_2)$ .
- 4. No evidence of division of anaerobic cells as aerosols in an anaerobic gas  $(N_2)$ .
- 5. Contradictory evidence that anaerobic spores become heat sensitive as aerosols in anaerobic gas  $(N_2)$ .
- 6. No evidence that airborne, aerobic cells can "feed" on other airborne particles or vapors.
- 7. A micro-aerophylic bacteria that grew at pH 10.6 in an ammonium atmosphere was isolated from soil. It failed to survive the 16th transfer and was not recovered.

During the project, 154 sets of aerosol runs (2 - 3 days per set) were conducted, most of which furnished negative or inconclusive data. In support of the runs, about 600 batches of cultures were produced and 4,850 samples were assayed.

### CONCLUSIONS

1. Under special conditions of high humidity, growth temperature (30°C), suitable medium, and the use of cells in the latter stages of logarithmic growth, the bacterial species <u>Serratia marcescens</u> can sustain more than two cellular replications in the airborne state. Thus, the null hypothesis that no microbial species can replicate in the airborne state has been shown to be false.

2. Certain metallic structures undergoing dynamic thermal stress can discharge attached, viable, microbes in the form of small particles.

3. If the turbulent atmosphere of Jupiter is treated as a stirred settling chamber, that is, the boundaries are neither expanding nor contracting so the net vector velocities of air movement are zero, then stirred settling theory predicts the mean half-life of particles in the size range of 1 - 10 µm diameter would be between 5 to 10 years.

Final Report N00014-75-C-1133

Page 5

## DISCUSSION

There are a number of factors associated with any attempts to estimate the probability that Jupiter would be contaminated by microbes unintentionally released from spacecraft entering the Jovian atmosphere. The factor of interest to this project was whether growth (in the sense of continued cellular division) is possible under any circumstance.

Prior to these studies, the best estimate of that probability was either nil or so close to nil that there was no practical difference. Our work has shown that this probability is actually 1, but the environment under which propagation can occur is not likely to be found on earth, and even less likely on Jupiter. It is important to note, however, that because of practical restrictions imposed by gravitational forces and chamber size, our studies were limited to 4-day periods at best. But growth usually ceased after 6 to 8 hours, indicating that essential nutrients had been consumed. In most instances, however, there was no additional death of airborne cells.

In view of the above, we believe that aerobic, airborne cells could be "fed", either continuously at a slow rate, or intermittently. by either nutrients in the vapor state, or by very small particles that would collide with the larger particles at a greater rate than if bacterial size particles were used. We might reasonably expect to find that the division time of airborne cells is longer than that of cells <u>in vitro</u>, so we might look for that process in cells 3 or 4 days (aerosol time) old. Limited studies of this nature are being continued.

Because it is possible for many aerobic cells to produce an ultimate waste product consisting of volatile products (CO<sub>2</sub> and water), whereas anaerobic cells, generally, do not easily rid themselves of waste materials, the possibility of demonstrating airborne growth of anaerobic microbes -- assuming there might be a species so oriented -seems remote.

Final Report: NASA N00014-75-C-1133 Page 6

#### INDEX

- Title: STUDIES ON MICROBIAL PROPAGATION IN THE AIRBORNE STATE R.L. Dimmick, H. Wolochow and M.A. Chatigny
- APPENDIX A. EVIDENCE FOR METABOLIC ACTIVITY OF AIRBORNE BACTERIA R.L. Dimmick, Patricia Ann Straat, H. Wolochow, G.V. Levin, M.A. Chatigny and J.R. Schrot Journal of Aerosol Science (1975) <u>6</u>:387-393
- APPENDIX B. POSSIBILITY OF GROWTH OF AIRBORNE MICROBES IN OUTER PLANETARY ATMOSPHERES R.L. Dimmick and M.A. Chatigny
  - In: Chemical Evolution of the Giant Planets (1976) Academic Press, Inc., New York pp. 95-106.
- APPENDIX C. EVIDENCE FOR FORMATION OF NEW DNA IN AIRBORNE, BACTERIAL CELLS Patricia Ann Straat, H. Wolochow, R.L. Dimmick and M.A. Chatigny Prepared for submission to: Applied and Environmental Microbiology
- APPENDIX D. PRODUCTION OF PHAGE IN INFECTED AIRBORNE BACTERIA H. Wolochow, R.L. Dimmick, Patricia Straat and M.A. Chatigny Prepared for submission to: Applied and Environmental Microbiology
- APPENDIX E. STUDIES ON PROPAGATION OF MICROBES IN THE AIRBORNE STATE R.L. Dimmick, H. Wolochow, Patricia Straat and M.A. Chatigny In: 50th Technical Progress Report, Naval Biosciences Laboratory, School of Public Health, University of California, Berkeley, Ca. 94720. pp. 330-338.
- APPENDIX F. EVIDENCE FOR PROPAGATION OF BACTERIA IN PARTICLES SUSPENDED IN GASEOUS ATMOSPHERES R.L. Dimmick, M.A. Chatigny, H. Wolochow and P. Straat. In: Proceedings, COSPAR Symposium (In Press)

Final Report: NASA N00014-75-C-1133

Page 7

APPENDIX G. Part I: RELEASE OF BACTERIAL SPORES FROM THE INNER WALLS OF A STAINLESS STEEL CUP SUBJECTED TO THERMAL STRESSES AND MECHANICAL SHOCK. H. Wolochow, M. Chatigny and J. Hebert <u>In</u>: 48th Technical Progress Report, Naval Biomedical Research Laboratory, School of Public Health, University of California, Berkeley, California. pp. 363-385

> Part II: RELEASE OF BACTERIAL SPORES FROM INNER WALLS OF A STAINLESS STEEL CUP SUBJECTED TO THERMAL STRESS. H. Wolochow, M.A. Chatigny and J. Hebert In: 49th Technical Progress Report, Naval Biomedical Research Laboratory, School of Public Health, University of California, Berkeley, California. pp. 358-374.

APPENDIX H. A SIMPLE METHOD FOR ESTIMATION OF COAGULATION EFFICIENCY IN MIXED AEROSOLS R.L. Dimmick, Alvin Boyd and H. Wolochow Journal of Aerosol Science (1975) <u>6</u>:375-377.

# Final Report: NASA N00014-75-C-1133

#### DISTRIBUTION LIST:

Dr. Richard S. Young (3 copies, with reprints) Chief, Planetary Biology NASA Headquarters, Code SL Washington, D.C. 20546

Naval Research Laboratory (Code 2627)(6 copies, without reprints) DODAAD Code N00173 Washington, D.C. 20375

Defense Documentation Center (12 Copies without reprints) Bldg 5, Cameron Station Alexandria, Virginia 22314

Dr. Arthur J. Emery, Jr. (1 copy, without reprints) Program Director Microbiology Department of the Navy Office of Naval Research (Code 443) 800 Quincy Street Arlington, Virginia 22217

Director, Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91101 (1 copy, without reprints)

Elmer G. Keith, Resident Representative (1 copy, without reprints) Department of the Navy, Office of Naval Research University of California 553 Evans Hall Berkeley, California 94720

Special Assistant, Code 1021P (6 copies, without reprints) International Programs and Special Activities Mr. R.H. Imus Rm 1000, Ballston Tower #1 Arlington, Va. 22217

Activity Commanding Officer (1 copy without reprints) Naval Biosciences Laboratory Naval Supply Center Oakland, California 94625