



The Use of a Photodensitometric Technique to Evaluate 9 ADA 03464 the Efficacy of Antiviral Compounds against Swine Influenza (Influenza A/New Jersey) Infection in Mice (11) 28 Dec 76 WILLIAM C./HALL, JOHN D. WHITE E GEORGE H. SCOTT

Running Head: PULMONARY PHOTODENSITOMETRY

U.S. Army Medical Research Institute of Infectious Diseases

Deferrent are

mit

Fort Detrick, Frederick, Maryland 21701 3M161102BS\$3

genterin rept.;] 28 Dec. 1976

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on the Revision of the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Research Council. The facilities are fully accredited by the American Association for Accreditation of Laboratory Animal Care.

405039

Approved for public release; distribution unlimited.

	Data Entered)	
REPORT DOCUMENTATI	ION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
T. REPORT NUMBER	2. GOVT ACCESSION	NO. 3. RECIPIENT'S CATALOG NUMBER
and an a routine	want	
TITLE (and Subtitio)	TILE (and Subtitie) if you man do tw bantate bes , alaval	
the Use of a Photodensitometric Evaluate the Efficacy of Antivi	rechnique to	Interim
Against Swine Influenza (Influenza A/New Jersey) Infection in Mice AUTHOR(*)		6. PERFORMING ORG. REPORT NUMBER
		8. CONTRACT OR GRANT NUMBER(+)
William C. Hall, John D. White, Scott	and George H.	icniticantly reduces contactor if treatment was colayed until tere effective in reducing WUM
PERFORMING ORGANIZATION NAME AND ADD	RESS LAND DO IS LOT	10. PROGRAM ELEMENT, PROJECT, TASK
U.S. Army Medical Research Institute of Infectious Diseases SGRD-UIP-A Fort Detrick, Frederick, MD 21701 1. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Medical Research and Development		61102B 3M161102BS03/00 016
		12. REPORT DATE 28 December 1976
Command, Office of The Surgeo	on General	13. NUMBER OF PAGES
4. MONITORING AGENCY NAME & ADDRESS(I di	liferent from Controlling Offic	e) 15. SECURITY CLASS. (of this report)
		Ibalagatitad
		15a. DECLASSIFICATION/DOWNGRADING
7. DISTRIBUTION STATEMENT (of the abstract on	ntered in Block 20, if differen	t from Report)
7. DISTRIBUTION STATEMENT (of the abstract on	stered in Block 20, if differen	t from Report)
7. DISTRIBUTION STATEMENT (of the abetract on	stered in Block 20, if differen	t from Report)
7. DISTRIBUTION STATEMENT (of the abetract on 8. SUPPLEMENTARY NOTES Reprints bearing assigned AD nu To be submitted for publication 1. KEY WORDS (Continue on reverse elde il necess	umber will be forw in Antimicrobial	varded upon receipt. Agents and Chemotherapy.
7. DISTRIBUTION STATEMENT (of the abstract on 6. SUPPLEMENTARY NOTES Reprints bearing assigned AD nu To be submitted for publication 9. KEY WORDS (Continue on reverse side if necession Swine influenza, mice, photoder	ntered in Block 20, if differen umber will be forw in Antimicrobial may and identify by block num nsitometry, virus	t from Report) All JAN 19 JAN
7. DISTRIBUTION STATEMENT (of the abstract on 8. SUPPLEMENTARY NOTES Reprints bearing assigned AD nu To be submitted for publication 9. KEY WORDS (Continue on reverse elde if necesso Swine influenza, mice, photoder	umber will be forw in Antimicrobial way and identify by block num	t from Report)
 DISTRIBUTION STATEMENT (of the abstract on Reprints bearing assigned AD nu To be submitted for publication KEY WORDS (Continue on reverse elde if necess Swine influenza, mice, photoder AUSTRACT (Continue on reverse elde if necess During studies of swine in technique was devised to quanti different time intervals with a is based on the premise that as the optical density of lung sec increase in cell numbers in inf instillation of the virus, mice 	amber will be forw mber will be forw in Antimicrobial any and identify by block num hsitometry, virus fluenza virus (A/ lfy the pulmonary intiviral chemothe the severity of ctions also increa ected lungs. Sev were killed and	t from Report) A gents and Chemotherapy. Agents and Chemotherapy. (ber) New Jersey) infection a lesions in mice treated at rapeutic agents. The technique microscopic change increases, ses due to edema and to an ren days after intranasal the lungs were perfused with
 DISTRIBUTION STATEMENT (of the abstract on Reprints bearing assigned AD nu To be submitted for publication KEY WORDS (Continue on reverse elde if necess Swine influenza, mice, photoder Adstract (Continue on reverse elde if necess During studies of swine in technique was devised to quanti different time intervals with a is based on the premise that as the optical density of lung sec increase in cell numbers in inf instillation of the virus, mice D FORM 1473 EDITION OF 1 NOV 65 IS C 	mber will be forw in Antimicrobial ary and identify by block num insitometry, virus filuenza virus (A/ lfy the pulmonary intiviral chemothe the severity of ctions also increas fected lungs. Sev were killed and	t from Report) A from Report arded upon receipt. Agents and Chemotherapy. Agents and Chemoth
 DISTRIBUTION STATEMENT (of the abstract on SUPPLEMENTARY NOTES Reprints bearing assigned AD nu To be submitted for publication KEY WORDS (Continue on reverse side if necess Swine influenza, mice, photoder Adstract (Continue on reverse side if necess During studies of swine in technique was devised to quanti different time intervals with a is based on the premise that as the optical density of lung sec increase in cell numbers in inf instillation of the virus, mice D 1000 1000 1000 1000 00 1000 00 1000 00	amber will be forw amber will be forw in Antimicrobial ary and identify by block num asitometry, virus ary and identify by block num asitometry, virus afluenza virus (A/ ify the pulmonary antiviral chemothe the severity of ctions also increas fected lungs. Sev a were killed and bosoLETE	t from Report) A from Report) A gente and Chemotherapy. Agents and



20. Abstract

2% glutaraldehyde at constant pressure. Lungs were processed in a routine manner, sectioned at standard levels, and stained with hematoxylin and eosin. Using standard photomicrography equipment, multiple optical density measurements were made of lung sections in a carefully controlled systematic manner and a mean optical density determined for each lung. The optical density of lungs of mice pretreated with amantadine, rimantadine, or ribavirin was significantly reduced compared with infected-untreated controls (P < 0.01). If treatment was delayed until 15 h after infection, amantadine and ribavirin were effective in reducing pulmonary optical density but rimantadine was with out effect. These findings correlated well with mean lung weight of each group; however, the sensitivity of the optical density technique was greater. The method offers promise as a reliable means of objectively quantifying the pulmonary response to a variety of infectious, toxic, and therapeutic agents.

REPORT DOCUMENTATION PAGE



ABSTRACT

During studies of swine influenza virus (A/New Jersey) infection a technique was devised to quantify the pulmonary lesions in mice treated at different time intervals with antiviral chemotherapeutic agents. The technique is based on the premise that as the severity of microscopic change increases, the optical density of lung sections also increases due to edens and to an increase in cell numbers in infected lungs. Seven days after intranasal instillation of the virus, mice were killed and the lungs were perfused with 2% glutaraldehyde at constant pressure. Lungs were processed in a routine manner, sectioned at standard levels, and stained with hematoxylin and eosin. Using standard photomicrography equipment, multiple optical density measurements were made of lung sections in a carefully controlled systematic manner and a mean optical density determined for each lung. The optical density of lungs of mice pretreated with amantadine, rimantadine, or ribavirin was significantly reduced compared with infected-untrested controls (P < 0.01). If treatment was delayed until 15 h after infection, amantadine and ribevirin were effective in reducing pulmonary optical density but rimentadine was without effect., These findings correlated well with mean lung weight of each group; opever, the consitivity of the optical density technique was greater. The hethod offers promise as a reliable means of objectively quantifying the pulmonary response to a variety of infactious, toxic, and therepeutic agents.

633

162



Animal models for evaluation of potential antiviral compounds have frequently been developed using human disease pathogens adapted to produce high mortality. The efficacy of the compounds is thus measured by survival rate of treated infected animals compared with untreated controls. These model systems are unnatural and unrealistic when the purpose intended for the drug is to treat diseases of man and enimals which are characterized by high morbidity and low mortality (6). During studies of swine influenza (influenza A/New Jersey), (Scott, G. H., manuscript in preparation), it was observed that the mouse-adapted virus did not have lethal virulence for adult mice and thus resembled natural influenza infections of man (4). It was necessary to seek smother end-point to measure the effectiveness of antiviral compounds on the respiratory disease in mice.

Numerous systems have been proposed to grade morphologic changes occurring in the lungs of animals infected with various respiratory disease-producing agents. Many are not completely objective. The majority of the reliable procedures are morphometric where various measurements can be unde of pulmonary structure to obtain consistent quantitative values (2). These techniques require considerable time to perform and thus are inefficient when numerous specimens must be examined.

A technique, described for studying emphysems in pigs and rats, was based on the postulate that emphysematous lung sections are less optically dense than normal lung and that differences could be quantified by a densitometer (3). A similar procedure has been employed to determine degree of Scythrocyte aggregation in vitro (1). Likewise, pulmonary disease characterized by edems and cell infiltration into

alveolar spaces and the interstitium should increase optical density. If this premise were correct, the effect of drugs, toxins, or chemicals, on the pulmonary response could be quantitatively assessed and compared in an unbiased, completely objective manner. This technique was utilized to compare the efficacy of antiviral chemotherapeutic agents in mice infected with swine influenza virus.

and there Ber Mary and Ling

de.

505 They are

A A A A

at the

Contraction of the second A start and a start and a start

> E. Anger States and

MATERIALS AND METHODS

<u>Virus</u>. Influenza, A/NJ/8/76 virus, (H_{sw} 1N1), was passed six times in embryonated eggs; adaptation to mice was accomplished in nine serial passages by intranasal (1.n.) instillation of supernatant from a suspension of homogenized lungs from mice infected 3 to 4 days previously. After the 9th passage, allantoic cavities of 10-day-old embryonated eggs were inoculated with supernatant from infected lungs. After incubation for 48 h at 35 C, infected allantoic fluid was harvested and clarified by centrifugation at 1200 x g at 4 C for 15 min. Aatibiotics were added to the clarified fluid to achieve a final concentration of 250 units of penicillin/ml and 100 ug of streptomycin/ ml; aliquots of the suspension were stored frozen at -60 C. The titer of the infected allantoic fluid was $10^{7.7}$ median egg infectious doses (EID_{s0}) per ml.

<u>Experimental design</u>. Eight week old (25 g) outbred, female Swiss mice [Cr1:COBS CD-I (ICR) ER] obtained from the Sendai-free colony of Charles River Laboratories were divided into eight groups each containing 10 mice. Three groups were prophylactically treated with amentadine, rimentadine or ribevirin in the drinking water at a domage of 0.25 mg/ml (approximately 60 mg/kg/mouse/day) 24 h prior to intranssal instillation of $10^{6.1}$ EID₅₀ of influenza virus in 0.05 ml. Three other groups were treated therapeutically with each of the drugs beginning 15 h after virus inoculation. In all groups, treatment was then continued for 7 days after inoculation of the virus. Two control groups of mice were utilized, one was infected but not treated and the other was neither infected nor treated.

<u>Mecropsy procedures</u>. Seven days after infection, all mice were killed by cervical dislocation. In half the mice from each group, lungs were dissected from surrounding tissue and weighed to the nearest milligram. In the remaining mice, the cutaneous tissue of the neck was dissected free from the trachea, and the thorax was opened. A 21-guage hypodermic needle attached to a flask containing 27 glutaraldehyde, pH 7.2, was inserted into the trachea and the lungs were perfused in situ for 30 sec. The pressure of the perfusate was 20 cm of water. To maintain pulmonary distension a small wascular clamp was placed on the trachea. Lungs were then dissected free and suspended in 27 glutaraldehyde.

Perfused lungs were subsequently fixed in 10% buffered neutral formalin and embedded whole in paraffin. Paraffin blocks were sectioned horizontally through the lung lobes and three 6 μ m sections were taken at 1.5 mm intervals (Fig. 1). The sections were deparaffinized and stained with hematoxylin and eosin (5). Staining was controlled so that variation was not observed among slides.

Densitometry. The sections of lung were placed on the machanical stage of a Zeiss Ultraphot II (Carl Zeiss, Oberkochen/Wuerttemberg, Germany) and voltage was adjusted to 15: a filter with peak absorption at 560 and 645 nm was placed between the light source and specimen, and the image was projected onto a viewing screen using the 2.5 X objective. Total magnification was 27 X. Twenty to 35 measurements were made on each section directly from the same location on the viewing screen using a photodensitometer (Model 501 M, Photovolt Corp., New York, NY). Optical densities were measured through 0.5 mm diameter holes spaced 0.85 mm apart in a rectangle of brass foil which was placed on the microscope slide (Fig. 2). The photovolt meter was adjusted to 0 optical density at a place on the slide where there was no tissue. The lung at each perforation was viewed independently for artifacts or extraneous tissue and, if present, the field was eliminated from the sample. Essentially, only alveoli and small bronchioles were measured. The above procedure was repeated for each of the three lung sections and the mean optical density of each lung was calculated.

7

•

Statistics. Data on lung weight and optical density were analyzed by one way analysis of variance using the least significant difference test to determine differences between groups. A linear regression analysis was used to determine correlation of mean optical density and mean lung weight of each group (7). RESULTS

Lesions of swine influenza in untreated infected controls consisted of multifocal areas of bronchiolitis surrounded by thickening of alveoli, minimal to moderate inflammatory cell infiltration, and edema in alveolar spaces. In contrast, lungs of uninfected control mice were free of lesions. These morphologic changes were reflected by an almost 3-fold increase in pulmonary optical density in the untreated infected control group compared to the unifected controls (Table 1, Fig. 3). Consistent readings were obtained by repeating the procedure. In general, with the exception of mice treated with rimantadine 15 h after infection, the mean optical density of treated mice was midway between the values of uninfected and infected-untreated controls and differed significantly from the latter (P < 0.01). By analyzing lung weights of the various groups, there was significant reduction of lung weight only for mice given rimentadine prior to virus inoculation and mice give ribavirin 15 h subsequent to infection corpored with infected-untreated controls (P < 0.01). However, the trend was the same as that observed by densitometry, with the lung weights of infected-treated animals falling between values observed for both control groups. The linear regression analysis of the means of pulmonary optical density against means of lung weight gave a coefficient of correlation of 0.92.

DISCUSSION

·....

This photodensitometric technique provided an excellent means of quantifying the effects of antiviral agents on the pulmonary lesions of mice infected by swine influenza virus. The virus used in this study was characterized by low mortality for adult Swiss mice; thus the infection was similar to many naturally-occurring influenza outbreaks (4). In all likelihood, evaluation of the efficacy of the chemotherapeutic agents was performed under conditions simulating the natural disease. It is important to note that in these studies, lungs from all mice were sampled at the same time after infection when the pneumonia was at a similar stage. Thus, differences in optical density were a reflection of differences in severity of disease rather than qualitative differences in inflammation.

With influenza A/New Jersey infection, good correlation between mean optical density and mean lung weight for the various groups (r = 0.92) suggests that both methods have merit in the assessment of antiviral activity. The advantage of the optical density technique over lung weight in this experiment is the increased sensitivity observed with the former method. Reasons for this are: 1) The variability of optical density measurements within groups was less than lung weight; 2) optical density measurements were independent of lung size; and 3) structures not contributing to the pathologic process such as blood, large airways, tissues artifacts, etc. were selectively eliminated by viewing each field prior to obtaining an optical density measurement. In addition, this technique utilized photomicrography equipment already found in most laboratories, thus precluding the expense of more elaborate apparatus.

9

1

We feel this technique offers promise as a means of evaluating the response of lungs to a variety of infectious and toxic agents and will provide an objective, unbiased measure of the therapeutic value of various drugs for treating various respiratory diseases.

ACKNOWLEDGEMENTS

÷ .

The authors appreciate the assistance given by Dr's. Elwell and Machotka on the necropsies and Mr. Hauer and Dr. Higbee on the statistics.

LITERATURE CITED

- Brooks, K. M., and R. C. Robbins. 1968. An in vitro method for determination of degree of intravascular aggregation of blood cells. Lab. Invest. 19:580-583.
- Dungworth, D. L., L. W. Schwartz, W. S. Tyler, and R. F. Phalen 1976. Morphological methods for evaluation of pulmonary toxicity in animals. Annu. Rev. Pharmacol. 16:381-399.
- Glauser, S. C., and E. M. Glauser. 1968. Densitometric analysis of normal and emphysematous lung tissue. Arch. Environ. Health 16:862-864.
- 4. Loosli, C. G. 1974. Influenza virus epidemics: a continuing problem. Geriatrics 29:103-120.
- Luna, L. G. 1968. Manual of histologic staining methods of the Armed Forces Institute of Pathology, 3rd Ed. McGraw-Hill Book Co., New York.
- 6. <u>Schabel, F. M.</u> 1965. Virus chemotherapy-why and how. Ala. J. Med. Sci. 2:149-154.
- Snedecor, G. W., and W. G. Cochran. 1967. Statistical methods,
 6th ed. Iowa State University Press, Ames.

TABLE 1. <u>Prophylactic and therapeutic effects of amantadine</u>, <u>rimantadine</u>, <u>or ribavirin on lung optical density and weight in influenza A/New Jersey-</u> <u>infected mice</u>.

Group	Mean value + SEM		
	Optical density X 10 ⁻³	Lung weight (mg)	
Prophylaxis + Therapy			
Amentadine	69 ± 4 ^{a,b}	310 ± 14 ^b	
Rimentadine	68 + 5 ^{a,b}	272 ± 17 ^{a,b}	
Ribevirin	71 ± 8 ^{a,b}	332 ± 42^{b}	
Therapy			
Amentadine	63 ± 2ª	324 ± 24 ^b	
Rimentadine	84 ± 11 ^b	352 ± 17 ^b	
Ribevirin	. 56 ± 3ª	266 ± 19 ^{a,b}	
Controls			
Infected-untrested	98 ± 11°	390 ± 36°	
Uninfected	38 ± 1 ^c	142 ± 2°	

^aDiffers significantly from infected-untrested control (P < 0.01). ^bDiffers significantly from uninfected control (P < 0.01). ^cControls differ significantly from each other (P < 0.01).

a tart hall some to

FIGURE LECENDS

....

in the

A state

he ward

FIG. 1. Schematic drawing of lateral view of mouse lung showing the level of sectioning.

FIG. 2. Perforated brass cover on microscope slide through which light was transmitted for optical density measurements.

FIG. 3. Effect of antiviral agents on lungs of mice infected with swime influenza.