

AD-A034 886

STATE UNIV OF NEW YORK AT BUFFALO DEPT OF ENGINEERIN--ETC F/G 20/1
TIME HARMONIC ACOUSTIC RADIATION FROM A SUBMERGED ELASTIC SHELL--ETC(U)
SEP 76 R P SHAW N00014-75-C-0302

UNCLASSIFIED


97

NL

1 of 1
ADA034886



END
DATE
FILMED
7-77



ADA 034886

(20)
B.S.

Department of Engineering Science,
FACULTY OF ENGINEERING AND APPLIED SCIENCE
State University of New York at Buffalo



Report No. 97

**TIME HARMONIC ACOUSTIC RADIATION FROM A
SUBMERGED ELASTIC SHELL DEFINED
BY NONCONCENTRIC CYLINDERS**

Handwritten signatures and initials, including "S" and "H/B".

by

Richard Paul Shaw
Professor



**COPY AVAILABLE TO DDC DOES NOT
PERMIT FULLY LEGIBLE PRODUCTION**

September, 1976

This research was supported by the Office of Naval Research under Contract N00014-75-C0302. Approved for public release. Distribution is unlimited.

ABSTRACT

The acoustic radiation from an elastic cylindrical shell defined by nonconcentric cylindrical surfaces and submerged in an acoustic fluid is studied. The driving mechanism is a time harmonic internal pressure. In particular, the influence of the nonconcentricity parameter, δ , on the resulting radiation field is examined. The solution is obtained by means of an expansion in powers of this parameter about the concentric solution, $\delta = 0$. Each order of solution is solved in closed form by means of boundary integral equation methods. Results are given for the zeroth and first order solutions and it is seen that a j^{th} driving mode will introduce nearest neighbor ($(j-1)$ and $(j+1)$) modes in the first order solution.

ADVISOR	DATE	✓
NTIS	NO. 8	
CLASSIFICATION		
UNCLASSIFIED / AVAILABLE	YES	
6. 1/19 81		
A		

I. INTRODUCTION

We consider the radiation into an infinite acoustic fluid from a submerged elastic shell subjected to a time harmonic internal pressure with some prescribed spatial distribution. The shell is bounded by surfaces of separable geometry, i.e., such that the governing (Helmholtz) partial differential equations have modal solutions, that are nonconcentric, i.e., the origin of the coordinate system in which the outer boundary is "separable" is displaced from that for the inner boundary by a nonconcentricity distance, δ . If δ were zero, the entire problem could be solved by separation of variables but with δ nonzero, the problem is not separable.

The purpose of this investigation is to study the influence of the nonconcentricity on the resulting radiation field by making an expansion of the solution in powers of δ . Clearly the zeroth order will give the classical separation of variables solution. Higher order solutions will then represent the influence of nonconcentricity on this basic solution.

Rather than work with the governing differential equations, we convert to a boundary integral equation representation which eventually allows us to solve a set of algebraic equations for each mode in each order to obtain the exact solution.

To simplify the algebra, the submerged elastic body will be taken as an infinite cylinder in plane strain thereby reducing the problem to a two dimensional case. A specific example of nonconcentric circular boundaries is calculated as an illustration. Analogous three dimensional problems are also being studied - the axisymmetric three dimensional case requires essentially no more effort than this two dimensional problem.

The full equations of elasticity shall be used rather than approximating set of shell equations since the use of the boundary integral equation method

reduces the problem by one dimension anyway thereby accomplishing one of the main purposes of shell theory.

This type of problem has been studied previously for acoustic radiators, both by the boundary integral equation method [1] and by shifting and superposition theorems [2].

II. GENERAL FORMULATION

A. ACOUSTIC FLUID EQUATIONS

The infinite acoustic fluid exterior to the elastic body is described in terms of an acoustic (excess) pressure field, p , which satisfies a Helmholtz equation

$$(1) \quad \nabla^2 p + k^2 p = 0$$

where $k = \omega/c$ is the acoustic wave number, ω is the frequency of the time harmonic dependence (taken in the form $\exp[-i\omega t]$) and c is the acoustic sound speed. This equation is readily replaced by a Helmholtz (or Weber, for the two dimensional case) integral equation; e.g., [3]

$$(2) \quad \epsilon p(\bar{r}) = \oint_{\Gamma} \left[-p(\bar{r}_0) \frac{\partial G(\bar{r}, \bar{r}_0)}{\partial n_0} + G(\bar{r}, \bar{r}_0) \frac{\partial p(\bar{r}_0)}{\partial n_0} \right] dS_0$$

where $G = \frac{i}{4} H_0^{(1)}(kR)$ in two dimensions and $G = \frac{1}{4} \exp[ikR]/R$ in three dimensions. R is the distance between the field point \bar{r} and the integration variable, \bar{r}_0 , n_0 is the outward normal from the fluid, Γ is the line or surface bounding the fluid, i.e., the elastic solid-acoustic fluid interface with length or area element dS , and $\epsilon = 0, 1/2, 1$ depending on whether the field point is exterior to the fluid, on the interface or interior to the fluid. The radiation condition is automatically satisfied by the choice of G . Since only Hankel functions of the first kind will occur, the superscript (1) will be dropped.

The fluid velocity field is related to the pressure field through the acoustic velocity potential ϕ , which could also have been used as the dependent variable

$$(a) \quad \bar{v} = -\nabla \phi$$

$$(3) \quad (b) \quad p = \rho_f \frac{\partial \phi}{\partial t} = -i\omega \rho_f \phi$$

where ρ_f is the acoustic fluid density.

B. ELASTIC SHELL EQUATIONS

The elastic shell involves a finite domain with an inner and an outer boundary (treating the infinite cylinder as a two dimensional plane strain problem). It is best described by the displacement potentials Φ and $\bar{\Psi}$ defined by the displacement field \bar{u} as;

$$(4) \quad \bar{u} = \nabla \Phi + \nabla \times \bar{\Psi} \quad ; \quad \nabla \cdot \bar{\Psi} = 0$$

These potentials also satisfy Helmholtz equations

$$(5) \quad \begin{aligned} (a) \quad & \nabla^2 \Phi + k_D^2 \Phi = 0 \\ (b) \quad & \nabla^2 \bar{\Psi} + k_R^2 \bar{\Psi} = 0 \end{aligned}$$

where $k_D = \omega/c_D$, $k_R = \omega/c_R$ and $c_D = \sqrt{(\lambda+2\mu)/\rho_s}$ is the dilatational wave speed, $c_R = \sqrt{\mu/\rho_s}$ is the rotational wave speed in an elastic medium of density ρ_s and Lamé parameters, λ and μ , e.g., [4].

In the two dimensional plane strain case considered here, the vector displacement potential $\bar{\Psi}$ has only one nonzero component and will be written as a scalar, ψ .

The displacement potentials are also related to the dilatation Δ and the rotation $\bar{\omega}_R$

$$(6) \quad \begin{aligned} (a) \quad & \Delta = -k_D^2 \Phi \\ (b) \quad & \bar{\omega}_R = +\frac{1}{2} k_R^2 \bar{\Psi} \end{aligned}$$

which may also be used as dependent variables.

Clearly these displacement potentials also satisfy Helmholtz integral equations analogous to equation (2) with two major changes. The wavenumber k must be changed to either k_D or k_R and the surface Γ now consists of two parts, the inner and the outer boundaries. We note that the direction of n_0 changes

sign between these boundaries since it is required to be "outward" from the elastic medium. Then

$$(7) \quad \varepsilon \Phi(\bar{r}) = \oint_{\Gamma_o} \left[-\Phi(\bar{r}_o) \frac{\partial G_D}{\partial n_o} + G_D \frac{\partial \Phi(\bar{r}_o)}{\partial n_o} \right] dS_{OUTER} \\ + \oint_{\Gamma_i} \left[-\Phi(\bar{r}_o) \frac{\partial G_D}{\partial n_o} + G_D \frac{\partial \Phi(\bar{r}_o)}{\partial n_o} \right] dS_{INNER}$$

$$(8) \quad \varepsilon \Psi(\bar{r}) = \oint_{\Gamma_o} \left[-\Psi(\bar{r}_o) \frac{\partial G_R}{\partial n_o} + G_R \frac{\partial \Psi(\bar{r}_o)}{\partial n_o} \right] dS_{OUTER} \\ + \oint_{\Gamma_i} \left[-\Psi(\bar{r}_o) \frac{\partial G_R}{\partial n_o} + G_R \frac{\partial \Psi(\bar{r}_o)}{\partial n_o} \right] dS_{INNER}$$

where G_D, G_R are equivalent to the previous G with k_D, k_R replacing k .

\bar{r} in any of these boundary integral equations will be placed on (or infinitesimally near to) either the inner or the outer surface. Thus each of the above boundary integral equations will produce two separate integral equations.

C. BOUNDARY CONDITIONS

We supplement these equations with five boundary conditions requiring zero shear stress, continuity of the normal displacement or velocity field and continuity of the normal stress field at the outer boundary of the elastic shell and zero shear stress and continuity of normal stress at the inner boundary. (Actually, those at the outer boundary might better be called interface conditions.)

For plane strain in a two dimensional orthogonal curvilinear coordinate system (α, β) , we have the infinitesimal strains, e.g., [5]

$$\begin{aligned} (a) \quad \epsilon_{\alpha\alpha} &= (1/h_1) \partial u_\alpha / \partial \alpha + (1/h_1 h_2) u_\beta \partial h_1 / \partial \beta \\ (9) \quad (b) \quad \epsilon_{\beta\beta} &= (1/h_2) \partial u_\beta / \partial \beta + (1/h_1 h_2) u_\alpha \partial h_2 / \partial \alpha \\ (c) \quad \epsilon_{\alpha\beta} &= (h_2/h_1) \partial [u_\beta/h_2] / \partial \alpha + (h_1/h_2) \partial [u_\alpha/h_1] / \partial \beta \end{aligned}$$

and the dilatation and rotation

$$\begin{aligned} (a) \quad \Delta &= [\partial (u_\alpha h_2) / \partial \alpha + \partial (u_\beta h_1) / \partial \beta] / [h_1 h_2] \\ (10) \quad (b) \quad \omega_R &= [\partial (u_\beta h_2) / \partial \alpha - \partial (u_\alpha h_1) / \partial \beta] / [2 h_1 h_2] \end{aligned}$$

where h_1 and h_2 are the metric coefficients (scale factors) for these coordinates, ($h_3 = 1$).

We can also write out equation (4) specifically:

$$\begin{aligned} (a) \quad u_\alpha &= (1/h_1) \partial \Phi / \partial \alpha + (1/h_2) \partial \Psi / \partial \beta \\ (11) \quad (b) \quad u_\beta &= (1/h_2) \partial \Phi / \partial \beta - (1/h_1) \partial \Psi / \partial \alpha \end{aligned}$$

We shall choose the boundaries to be surfaces of constant α ; the α direction is then normal to these surfaces. Continuity of normal stress requires

$$-p = \sigma_{\alpha\alpha} = \lambda \Delta + 2\mu \epsilon_{\alpha\alpha}$$

which reduced to

$$(12) \quad -p = -\rho_s \omega^2 \Phi + 2\mu \left\{ \frac{-1}{h_1^2 h_2} \frac{\partial h_2}{\partial \alpha} \frac{\partial \Phi}{\partial \alpha} + \frac{1}{h_2^3} \frac{\partial h_2}{\partial \beta} \frac{\partial \Phi}{\partial \beta} - \frac{1}{h_2^2} \frac{\partial^2 \Phi}{\partial \beta^2} - \frac{1}{h_1 h_2^2} \frac{\partial h_2}{\partial \alpha} \frac{\partial \Psi}{\partial \beta} - \frac{1}{h_1^2 h_2} \frac{\partial h_1}{\partial \beta} \frac{\partial \Psi}{\partial \alpha} + \frac{1}{h_1 h_2} \frac{\partial^2 \Psi}{\partial \alpha \partial \beta} \right\}$$

Similarly, vanishing of the shear stress requires

$$(13) \quad 0 = \frac{2}{h_1 h_2} \frac{\partial^2 \Phi}{\partial \alpha \partial \beta} - \frac{2}{h_1^2 h_2} \frac{\partial h_1}{\partial \beta} \frac{\partial \Phi}{\partial \alpha} - \frac{2}{h_1 h_2^2} \frac{\partial h_2}{\partial \alpha} \frac{\partial \Phi}{\partial \beta} + \frac{2}{h_1^2 h_2} \frac{\partial h_2}{\partial \alpha} \frac{\partial \Psi}{\partial \alpha} - \frac{2}{h_2^3} \frac{\partial h_2}{\partial \beta} \frac{\partial \Psi}{\partial \beta} + \frac{2}{h_2^2} \frac{\partial^2 \Psi}{\partial \beta^2} + h_1^2 \Psi$$

and continuity of normal velocity requires

$$(14) \quad \frac{\partial p}{\partial \alpha} = \rho_f \omega^2 \left[\frac{1}{h_1} \frac{\partial \Phi}{\partial \alpha} + \frac{1}{h_2} \frac{\partial \Psi}{\partial \beta} \right]$$

D. SOLUTION PROCEDURE

There will in general be ten equations, two boundary conditions plus two integral equations (on ϕ , ψ) evaluated at the inner boundary and three boundary conditions plus three integral equations (on p , ϕ and ψ) evaluated at the outer boundary. Treating p at the inner boundary as known, these equations describe the behavior of ϕ , $\partial\phi/\partial\alpha$, ψ , $\partial\psi/\partial\alpha$ on the inner boundary and p , $\partial p/\partial\alpha$, ϕ , $\partial\phi/\partial\alpha$, ψ , $\partial\psi/\partial\alpha$ on the outer boundary as ten unknown functions of the remaining independent variable β . To emphasize the use of a different origin, we shall use the symbols (α, β) for the inner boundary coordinate system and the symbols (α', β') for the outer. We also use a subscript o to represent an integration coordinate. To simplify the algebra as much as possible, the inner boundary forcing pressure will be taken proportional to the j th eigenfunction of the separated Helmholtz equation for that surface. Finally the inner boundary will be a surface of constant $\alpha = a$ and the outer of constant $\alpha' = b$, as shown in Figure 1.

We recognize that the inner boundary variables may be expanded as a series in terms of the inner boundary eigenfunctions; similarly for the outer boundary variables. The ψ and $\partial\psi/\partial\alpha$ series will be in terms of the antisymmetric modes if p , ϕ , etc. are in terms of the symmetric modes and vice-versa as required by the boundary conditions. Since each boundary condition involves only a single boundary, they must be separable, i.e., each mode of the series expansion must satisfy the boundary conditions independently of the other modes thereby providing five algebraic equations on the unknown coefficients of each mode of the expansion.

The boundary integral equations, however, have one set of terms wherein the integration and field point lie on the same surface, thereby allowing a direct integration to be carried out independently of δ , and another set of "mixed"

terms where the integration and field point lie on different surfaces, thereby involving δ explicitly.

We can carry out the integration of the first type quite simply since the Greens' function possesses an eigenfunction expansion in terms of the same functions (either completely inner or completely outer boundaries) as did the other dependent variables. These integrals then pose no real difficulty. Integrals of the mixed (second) type would pose no difficulty if δ were zero, i.e., in the concentric cases, the eigenfunctions in terms of β must be identical on surfaces of constant α . This then suggests a power series expansion in terms of δ about δ equal to zero of all of the algebraic coefficients of all of the dependent variable series expansions as well as that of the Greens' function for the mixed integrals. The equations for the boundary conditions and for the first type of integral do not involve δ explicitly and expressions for each order of δ are identical to the original expression. The mixed integrals involve a product of two series in powers of δ and thereby introduce a coupling of orders. The zeroth order solution will, of course, be that obtainable directly by separation of variables for the concentric case. The first order solution is coupled to the zeroth order solution through the mixed integrals -- this in the first order solution will be found to introduce "nearest neighbor modes" to each mode of the zeroth order solution, etc.

By taking the forcing function to excite only a single mode in the zeroth order solution, we find ten algebraic equations to solve for the ten unknown coefficients in the zeroth order series expansion, two sets of ten algebraic equations for the first order nearest neighbor modes, etc.

We shall illustrate this by an example.

III. SPECIFIC EXAMPLE - CIRCULAR CYLINDER

A. BOUNDARY CONDITIONS AND EIGENFUNCTION EXPANSIONS

Let us consider nonconcentric circular boundaries such that $(\alpha, \beta) = (r, \theta)$ and $(\alpha', \beta') = (r', \theta')$ with $r = a$ the inner and $r' = b$ the outer boundaries. The scale factors are $h_1 = 1$ and $h_2 = r$ and the boundary conditions become

$$(15) \quad -p = -\rho_s \omega^2 \Phi + 2\mu \left[\frac{-1}{r} \frac{\partial \Phi}{\partial r} - \frac{1}{r^2} \frac{\partial^2 \Phi}{\partial \theta^2} - \frac{1}{r^2} \frac{\partial \Psi}{\partial \theta} + \frac{1}{r} \frac{\partial^2 \Psi}{\partial r \partial \theta} \right]$$

$$(16) \quad 0 = \mu \left[\frac{2}{r} \frac{\partial^2 \Phi}{\partial r \partial \theta} + \frac{2}{r^2} \frac{\partial^2 \Psi}{\partial \theta^2} - \frac{2}{r^2} \frac{\partial \Phi}{\partial \theta} + \frac{2}{r} \frac{\partial \Psi}{\partial r} \right] + \rho_s \omega^2 \Psi$$

$$(17) \quad \frac{\partial p}{\partial r} = \rho_f \omega^2 \left[\frac{\partial \Phi}{\partial r} + \frac{1}{r} \frac{\partial \Psi}{\partial \theta} \right]$$

We have the appropriate eigenfunction expansions

$$(18) \quad \left[p, \Phi, \frac{\partial \Phi}{\partial r} \right]_{r=a} = \sum_{n=0}^{\infty} \left[A_{1n}, A_{3n}, A_{4n} \right] \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta)$$

$$(19) \quad \left[\Psi, \frac{\partial \Psi}{\partial r} \right]_{r=a} = \sum_{n=0}^{\infty} \left[A_{5n}, A_{6n} \right] \begin{Bmatrix} \cos \\ -\sin \end{Bmatrix} (n\theta)$$

$$(20) \quad \left[p, \frac{\partial p}{\partial r'}, \Phi, \frac{\partial \Phi}{\partial r'} \right]_{r'=b} = \sum_{n=0}^{\infty} \left[B_{1n}, B_{2n}, B_{3n}, B_{4n} \right] \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta')$$

$$(21) \quad \left[\Psi, \frac{\partial \Psi}{\partial r'} \right]_{r'=b} = \sum_{n=0}^{\infty} \left[B_{5n}, B_{6n} \right] \begin{Bmatrix} \cos \\ -\sin \end{Bmatrix} (n\theta')$$

which reduce the boundary conditions to algebraic equations

$$(22) \quad -A_{1n} = -\rho_s \omega^2 A_{3n} + 2\mu \left\{ -\frac{1}{a} A_{4n} + \frac{n^2}{a^2} A_{3n} + \frac{n}{a^2} A_{5n} - \frac{n}{a} A_{6n} \right\}$$

$$(23) \quad 0 = 2\mu \left\{ \frac{1}{a} A_{4n} - \frac{n^2}{a^2} A_{5n} - \frac{n}{a^2} A_{3n} + \frac{1}{a} A_{6n} \right\} + \rho_s \omega^2 A_{5n}$$

$$(24) \quad -B_{1n} = -\rho_s \omega^2 B_{3n} + 2\mu \left\{ -\frac{1}{b} B_{4n} + \frac{n^2}{b^2} B_{3n} + \frac{n}{b^2} B_{5n} - \frac{n}{b} B_{6n} \right\}$$

$$(25) \quad 0 = 2\mu \left\{ \frac{1}{b} B_{4n} - \frac{n^2}{b^2} B_{5n} - \frac{n}{b^2} B_{3n} + \frac{1}{b} B_{6n} \right\} + \rho_s \omega^2 B_{5n}$$

$$(26) \quad B_{2n} = \rho_f \omega^2 \left[B_{4n} - \frac{n}{b} B_{5n} \right]$$

We now consider the eigenfunction expansion of the Greens' function:

$$(27) \quad G = \frac{i}{4} H_0(kR) = \frac{i}{4} \sum_{m=0}^{\infty} \epsilon_m \cos[m(\theta - \theta_0)] J_m(k\alpha) H_m(k\beta)$$

where $\epsilon_0 = 1$, $\epsilon_m = 2$ ($m \neq 0$), $R^2 = \alpha^2 + \beta^2 - 2\alpha\beta\cos(\theta - \theta_0)$ and β must be greater than α . [These β, α have no relationship to the general coordinates used before but are merely convenient labels here for the field and integration points.] θ or θ_0 may be primed or unprimed depending on the particular cases and α, β may be either a or b . To keep α from being identically equal to β , we choose field points just "off" the integration surface. In all cases, we shall choose them to be "outside" of the region over which the original integration is performed, i.e., for the infinite acoustic fluid we choose r' at b^- while for the elastic solid we choose r' at b^+ and r at a^- for field points at the outer and inner boundaries respectively.

B. BOUNDARY INTEGRAL EQUATION - FLUID

The boundary integral equation for the infinite acoustic fluid becomes

$$(28) \quad 0 = \int_0^{2\pi} \left[\left\{ \sum_{n=0}^{\infty} -B_{2n} \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta'_0) \cdot \sum_{m=0}^{\infty} \epsilon_m \cos [m(\theta' - \theta'_0)] J_m(kr') H_m(kr'_0) \right\} \right. \\ \left. + \sum_{n=0}^{\infty} B_{1n} \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta'_0) \cdot \sum_{m=0}^{\infty} k \epsilon_m \cos [m(\theta' - \theta'_0)] J_m(kr') H'_m(kr'_0) \right] b d\theta'_0 \quad \left. \begin{array}{l} r'_0 = b^- \\ r'_0 = b \end{array} \right.$$

where we have identified α with $r'_0 = b^-$ as smaller in magnitude than β which is identified with $r'_0 = b$. This tells us essentially which term to differentiate in the Greens' function series, and leads to a simple relationship

$$(29) \quad 0 = \sum_{n=0}^{\infty} \left\{ B_{2n} J_n(kb) H_n(kb) - k B_{1n} J_n(kb) H'_n(kb) \right\} \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta')$$

which by the orthogonality of the eigenfunctions reduces, for $J_n(kb) \neq 0$, to

$$(30) \quad B_{2n} = B_{1n} \cdot k \cdot H'_n(kb) / H_n(kb)$$

C. BOUNDARY INTEGRAL EQUATION - SOLID

We may now consider the boundary integral equations for the elastic shell. First, we examine the outer boundary equations with $r'=b^+$, i.e., we place the field point just outside of the outer boundary of the elastic body. Then, for example, we have

$$(31) \quad 0 = \int_0^{2\pi} \left\{ -\Phi(b) \frac{\partial G_0}{\partial r_0'} + G_0 \frac{\partial \Phi(b)}{\partial r_0'} \right\}_{r_0'=b, r'=b^+} b d\theta_0' \\ + \int_0^{2\pi} \left\{ +\Phi(a) \frac{\partial G_0}{\partial r_0} - G_0 \frac{\partial \Phi(a)}{\partial r_0} \right\}_{r_0=a, r'=b} a d\theta_0$$

The first integral leads directly to terms involving $\left\{ \frac{\sin}{\cos} \right\} n\theta'$ as in the acoustic fluid case above. The second integral does not. We cannot mix θ and θ' in equation (31) as long as they have different origins. To avoid this dilemma, we expand about $\delta=0$ such that, for example,

$$(32) \quad G = \frac{i}{4} H_0(kR) = \sum_{l=0}^{\infty} \delta^l G^{(l)} [s=0] = \frac{i}{4} \sum_{l=0}^{\infty} \delta^l [H_0(kR)]_{s=0}^{(l)}$$

The δ terms may be removed from the integration leaving an integral over $G_D^{(l)}(\delta=0)$, i.e., some Greens function with θ and θ' measured from the same origin.

Since this is simply a Taylor Series in δ , we have (see reference [] for details)

$$(33) \quad G^{(l)} [s=0] = \frac{1}{l!} \left[\frac{\partial^l H_0(kR)}{\partial \delta^l} \right]_{s=0}$$

We cannot use the eigenfunction, equation (27), expansion of G directly since it applies only when θ and θ' are measured from the same origin. Instead we can use, for example,

$$(34) \quad [H_0(kR)]_{s=0}^{(l)} = \left[\frac{\partial H_0(kR)}{\partial \delta} \right]_{s=0} = -k [H_1(kR) \frac{\partial R}{\partial \delta}]_{s=0}$$

where

$$(35) \quad R^2 = \lambda + \lambda_0^2 - 2\lambda\lambda_0 \cos(\theta_0 - \theta) + 2\delta[\lambda_0 \cos \theta_0 - \lambda \cos \theta] + \delta^2$$

and

$$(36) \quad \frac{\partial R}{\partial \delta} = [\delta + \lambda_0 \cos \theta_0 - \lambda \cos \theta] / R$$

Here λ and λ_0 are distances measured along θ and θ_0 respectively from origins which are a distance δ apart on the x axis. We then have the first order kernel

$$(37) \quad [H_0(kR)]_{\delta=0}^{(1)} = -k [H_1(kR) \cdot [\lambda_0 \cos \theta_0 - \lambda \cos \theta] / R]_{\delta=0}$$

For $\lambda = r$, $\lambda_0 = r_0'$ we have

$$(38) \quad \int_0^{2\pi} \left\{ \frac{\sin}{\cos} \right\} (n\theta_0') [H_0(kR)]_{\delta=0}^{(1)} d\theta_0' \\ = \pi k H_n(kr_0') \left[\left\{ \frac{\sin}{\cos} \right\} ((n+1)\theta) J_{n+1}(kr) - \left\{ \frac{\sin}{\cos} \right\} ((n-1)\theta) J_{n-1}(kr) \right]$$

We may set r_0' equal to b and r equal to a after any appropriate differentiations are carried out.

Similarly for $\lambda = r'$ and $\lambda_0 = r_0$

$$(39) \quad \int_0^{2\pi} \left\{ \frac{\sin}{\cos} \right\} (n\theta_0) [H_0(kR)]_{\delta=0}^{(1)} d\theta_0 \\ = \pi k J_n(kr_0) \left[\left\{ \frac{\sin}{\cos} \right\} ((n-1)\theta') H_{n-1}(kr') - \left\{ \frac{\sin}{\cos} \right\} ((n+1)\theta') H_{n+1}(kr') \right]$$

Clearly we must also expand all of the other dependent variables in powers of δ - or equivalently the coefficients of their eigenfunction expansions:

$$(40) \quad \begin{aligned} (a) \quad A_{jm} &= \sum_{l=0}^{\infty} \delta^l A_{jm}^{(l)} \quad ; \quad j = 1, 3, 4, 5, 6 \\ (b) \quad B_{jm} &= \sum_{l=0}^{\infty} \delta^l B_{jm}^{(l)} \quad ; \quad j = 1, 2, 3, 4, 5, 6 \end{aligned}$$

Solving each order of δ separately, we have

(i) Zeroth order equation

$$(43) \quad 0 = \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[-B_{3n}^{(0)} \frac{\partial G_D}{\partial r_0'} + B_{4n}^{(0)} G_D \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0') b d\theta_0' \right\}_{r_0'=b, r'=b^+} \\ + \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[+A_{3n}^{(0)} \frac{\partial G_D^{(0)}}{\partial r_0} - A_{4n}^{(0)} G_D^{(0)} \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0) a d\theta_0 \right\}_{r_0=a, r'=b^+}$$

(ii) First order equation:

$$(44) \quad \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[-A_{3n}^{(1)} \frac{\partial G_D^{(1)}}{\partial r_0} + A_{4n}^{(1)} G_D^{(1)} \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0) a d\theta_0 \right\}_{r_0=a, r'=b^+} \\ = \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[-B_{3n}^{(1)} \frac{\partial G_D}{\partial r_0} + B_{4n}^{(1)} G_D \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0') b d\theta_0' \right\}_{r_0'=b, r'=b^+} \\ + \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[A_{3n}^{(1)} \frac{\partial G_D^{(0)}}{\partial r_0} - A_{4n}^{(1)} G_D^{(0)} \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0) a d\theta_0 \right\}_{r_0=a, r'=b^+}$$

We see that the first order solution is coupled to the zeroth order solution.

Furthermore, the form of $G_D^{(1)}$ is such that an j mode zeroth order term will produce $(j+1)$ and $(j-1)$ mode first order terms. A similar equation holds on ψ .

The inner boundary integral equations are handled in the same manner with r' equal to a^- , i.e., the field point just short of the inner boundary. For example, we now have

$$(45) \quad 0 = \int_0^{2\pi} \left\{ -\Phi(b) \frac{\partial G_D}{\partial r_0'} + G_D \frac{\partial \Phi(b)}{\partial r_0'} \right\}_{r_0'=b, r'=a^-} b d\theta_0' \\ + \int_0^{2\pi} \left\{ \Phi(a) \frac{\partial G_D}{\partial r_0} - G_D \frac{\partial \Phi(a)}{\partial r_0} \right\}_{r_0=a, r'=a^-} a d\theta_0$$

Now the first integral is a mixed type. Using the same expansions as before we have

(i) Zeroth order equation

$$(46) \quad 0 = \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[-B_{3n}^{(0)} \frac{\partial G_D^{(0)}}{\partial r_0'} + B_{4n}^{(0)} G_D^{(0)} \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0') \right\} b d\theta_0' \\ + \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[A_{3n}^{(0)} \frac{\partial G_D}{\partial r_0} - A_{4n}^{(0)} G_D \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0) \right\} a d\theta_0$$

$r_0' = b, r = a^-$

$r_0 = a, r = a^-$

(ii) First order equations

$$(47) \quad 0 = \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[-B_{3n}^{(0)} \frac{\partial G_D^{(1)}}{\partial r_0'} + B_{4n}^{(0)} G_D^{(1)} - B_{3n}^{(1)} \frac{\partial G_D^{(0)}}{\partial r_0'} + B_{4n}^{(1)} G_D^{(0)} \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0') \right\} b d\theta_0' \\ + \int_0^{2\pi} \left\{ \sum_{n=0}^{\infty} \left[A_{3n}^{(1)} \frac{\partial G_D}{\partial r_0} - A_{4n}^{(1)} G_D \right] \left\{ \frac{\sin}{\cos} \right\} (n\theta_0) \right\} a d\theta_0$$

$r_0' = b, r = a^-$

$r_0 = a, r = a^-$

with a similar equation on ψ , etc.

We now choose $p(a)$ to represent a single mode, i.e., $A_{1n} = 0, n \neq j$ and $A_{1j} = P_j$. The zeroth order boundary integral equations and boundary conditions will also only involve this one mode. The first order boundary integral equations will be satisfied only with the $(j+1)$ and $(j-1)$ modes; the boundary conditions will therefore be applied only for these modes as well.

D. ZEROETH ORDER SOLUTION

The boundary conditions, equations (15-17), apply for each order separately and may be used directly as may the acoustic fluid boundary integral equation (30). The outer boundary shell equation on ϕ , equation (46), integrates to

$$(48) \quad 0 = \sum_{n=0}^{\infty} \left\{ b \left\{ -B_{3n}^{(0)} k_D J_n'(k_D b) + B_{4n}^{(0)} J_n(k_D b) \right\} + a \left\{ A_{3n}^{(0)} k_D J_n'(k_D a) - A_{4n}^{(0)} J_n(k_D a) \right\} \right\} H_n(k_D b) \cdot \left\{ \frac{\sin}{\cos} \right\}_{n\theta'}$$

Since $H_n(k_D b)$ cannot be zero and $\left\{ \frac{\sin}{\cos} \right\}_{n\theta'}$ are orthogonal, we have

$$(49) \quad 0 = k_D \left\{ -b B_{3n}^{(0)} J_n'(k_D b) + a A_{3n}^{(0)} J_n'(k_D a) \right\} + b B_{4n}^{(0)} J_n(k_D b) - a A_{4n}^{(0)} J_n(k_D a)$$

Similarly for ψ , with $H_n(k_R b)$ not equal to zero;

$$(50) \quad 0 = k_R \left\{ -b B_{5n}^{(0)} J_n'(k_R b) + a A_{5n}^{(0)} J_n'(k_R a) \right\} + b B_{6n}^{(0)} J_n(k_R b) - a A_{6n}^{(0)} J_n(k_R a)$$

The inner boundary shell equations give similar results provided $J_n(k_D a)$ and $J_n(k_R a)$ are not zero.

$$(51) \quad 0 = k_D \left\{ -b B_{3n}^{(0)} H_n'(k_D b) + a A_{3n}^{(0)} H_n'(k_D a) \right\} + b B_{4n}^{(0)} H_n(k_D b) - a A_{4n}^{(0)} H_n(k_D a)$$

$$(52) \quad 0 = k_R \left\{ -b B_{5n}^{(0)} H_n'(k_R b) + a A_{5n}^{(0)} H_n'(k_R a) \right\} + b B_{6n}^{(0)} H_n(k_R b) - a A_{6n}^{(0)} H_n(k_R a)$$

In summary, our equations are

$$(53) \text{ (a) } -A_{1n} = (-\rho_s \omega^2 + 2\mu n^2/a^2) A_{3n} + 2\mu [-A_{4n}/a + n A_{5n}/a^2 - n A_{6n}/a]$$

$$(b) \quad 0 = -n\mu A_{3n}/a^2 + n\mu A_{4n}/a^2 + \left[\frac{\rho_s \omega^2}{2} - \mu n^2/a^2 \right] A_{5n} + \mu A_{6n}/a$$

$$(c) \quad 0 = B_{1n} + [-\rho_s \omega^2 + 2\mu n^2/b^2] B_{3n} + 2\mu [-B_{4n}/b + n B_{5n}/b^2 - n B_{6n}/b]$$

$$(d) \quad 0 = -n\mu B_{3n}/b^2 + \mu n B_{4n}/b + \left[\frac{\rho_s \omega^2}{2} - \mu n^2/b^2 \right] B_{5n} + \mu B_{6n}/b$$

$$(e) \quad 0 = -B_{2n} + \rho_f \omega^2 [B_{4n} - n B_{5n}/b]$$

$$(f) \quad 0 = -B_{2n} + k H_n'(kb) B_{1n} / H_n(kb)$$

$$(g) \quad 0 = k_0 [-b J_n'(k_0 b) B_{3n} + a J_n'(k_0 a) A_{3n}] + b J_n(k_0 b) B_{4n} - a J_n(k_0 a) A_{4n}$$

$$(h) \quad 0 = k_R [-b J_n'(k_R b) B_{5n} + a J_n'(k_R a) A_{5n}] + b J_n(k_R b) B_{6n} - a J_n(k_R a) A_{6n}$$

$$(i) \quad 0 = k_0 [-b Y_n'(k_0 b) B_{3n} + a Y_n'(k_0 a) A_{3n}] + b Y_n(k_0 b) B_{4n} - a Y_n(k_0 a) A_{4n}$$

$$(j) \quad 0 = k_R [-b Y_n'(k_R b) B_{5n} + a Y_n'(k_R a) A_{5n}] + b Y_n(k_R b) B_{6n} - a Y_n(k_R a) A_{6n}$$

We have dropped the superscript (0) from the coefficients in these equations and trivially combined the last four equations to give real rather than complex expressions, although A_{in} , B_{in} are complex anyway due to the complex term in the sixth equation. If A_{in} is zero for $n \neq j$, all coefficients are zero for $n \neq j$. This then leaves only one mode, $n = j$, to be determined. These equations may readily be checked by the well known solution for the concentric case; they are solved here to indicate the procedure to be used for the higher order terms which have the same coefficients but a modified forcing function (inhomogeneous terms).

We may nondimensionalize these equations using a reference "potential", Φ_j , related to the reference pressure, P_j by $P_j = \omega^2 \rho_s \Phi_j = (\lambda + 2\mu) k_D^2 \Phi_j$. We use:

$$(54) \quad \alpha_D = k_D a, \quad \alpha_R = k_R a, \quad \beta_D = k_D b, \quad \beta_R = k_R b, \quad \gamma = k b$$

$$e = \rho_F / \rho_s, \quad \tau = 2\mu / (\lambda + 2\mu) = 2\alpha_D^2 / \alpha_R^2, \quad h_j = [\gamma H_j'(\gamma) / H_j(\gamma)]^{-1}$$

and

$$(55) \quad \begin{aligned} \tilde{A}_1 &= A_{1j} a^2 / (\lambda + 2\mu) \Phi_j = \alpha_D^2 ; & \tilde{B}_1 &= \left(\frac{\rho_s}{\rho_F}\right) B_{1j} b^2 / (\lambda + 2\mu) \Phi_j \\ & & \tilde{B}_2 &= \left(\frac{\rho_s}{\rho_F}\right) B_{2j} b^3 / (\lambda + 2\mu) \Phi_j \\ \tilde{A}_3 &= A_{3j} / \Phi_j ; & \tilde{B}_3 &= B_{3j} / \Phi_j \\ \tilde{A}_4 &= A_{4j} \cdot a / \Phi_j ; & \tilde{B}_4 &= B_{4j} \cdot b / \Phi_j \\ \tilde{A}_5 &= A_{5j} / \Phi_j ; & \tilde{B}_5 &= B_{5j} / \Phi_j \\ \tilde{A}_6 &= A_{6j} \cdot a / \Phi_j ; & \tilde{B}_6 &= B_{6j} \cdot b / \Phi_j \end{aligned}$$

leading to

$$(56) (a) -\alpha_D^2 = [\tau j^2 - \alpha_D^2] \tilde{A}_3 + \tau [-\tilde{A}_4 + j \tilde{A}_5 - j \tilde{A}_6]$$

$$(b) 0 = -j\tau \tilde{A}_3 + j\tau \tilde{A}_4 + [\alpha_D^2 - j^2\tau] \tilde{A}_5 + \tau \tilde{A}_6$$

$$(c) 0 = e \tilde{B}_1 + [j^2\tau - \beta_D^2] \tilde{B}_3 + \tau [-\tilde{B}_4 + j \tilde{B}_5 - j \tilde{B}_6]$$

$$(d) 0 = -j\tau \tilde{B}_3 + j\tau \tilde{B}_4 + [\beta_D^2 - j^2\tau] \tilde{B}_5 + \tau \tilde{B}_6$$

$$(e) 0 = -\tilde{B}_2 + \beta_D^2 \tilde{B}_4 - j \beta_D^2 \tilde{B}_5$$

$$(f) 0 = -\tilde{B}_2 + \tilde{B}_1 / h_j$$

$$(g) 0 = -\beta_D J_j'(\beta_D) \tilde{B}_3 + \alpha_D J_j'(\alpha_D) \tilde{A}_3 + J_j(\beta_D) \tilde{B}_4 - J_j(\alpha_D) \tilde{A}_4$$

$$(h) 0 = -\beta_R J_j'(\beta_R) \tilde{B}_5 + \alpha_R J_j'(\alpha_R) \tilde{B}_5 + J_j(\beta_R) \tilde{B}_6 - J_j(\alpha_R) \tilde{A}_6$$

$$(i) 0 = -\beta_D Y_j'(\beta_D) \tilde{B}_3 + \alpha_D Y_j'(\alpha_D) \tilde{A}_3 + Y_j(\beta_D) \tilde{B}_4 - Y_j(\alpha_D) \tilde{A}_4$$

$$(j) 0 = -\beta_R Y_j'(\beta_R) \tilde{B}_5 + \alpha_R Y_j'(\alpha_R) \tilde{A}_5 + Y_j(\beta_R) \tilde{B}_6 - Y_j(\alpha_R) \tilde{A}_6$$

Although this may appear to be a formidable system - ten equations in ten complex unknowns corresponding to a twenty by twenty matrix system - the equations are readily reduced, eliminating variables until a simpler system is reached, e.g., B_1 and B_2 are eliminated directly using the fifth and sixth equations:

$$(a) \quad \tilde{B}_2 = \beta_0^2 [\tilde{B}_4 - j \tilde{B}_5]$$

$$(57) \quad (b) \quad \tilde{B}_1 = h \beta_0^2 [\tilde{B}_4 - j \tilde{B}_5]$$

A_3, A_5 can be written in terms of A_4 and A_6 with similar equations for the B_3, B_5

$$(a) \quad \tilde{A}_3 = K_3 + C_{34} \tilde{A}_4 + C_{36} \tilde{A}_6$$

$$(b) \quad \tilde{A}_5 = K_5 + C_{54} \tilde{A}_4 + C_{56} \tilde{A}_6$$

$$(58) \quad (c) \quad \tilde{B}_3 = D_{34} B_4 + D_{36} B_6$$

$$(d) \quad \tilde{B}_5 = D_{54} B_4 + D_{56} B_6$$

using the first four equations,

$$K_3 = \alpha_0^2 (\alpha_0^2 - \tau j^2) / [(\alpha_0^2 - \tau j^2)^2 - \tau^2 j^2]$$

$$\text{with } K_5 = j \tau \alpha_0^2 / [\quad " \quad]$$

$$C_{34} = C_{56} = - \tau \alpha_0^2 / [\quad " \quad]$$

$$C_{36} = C_{54} = (-j \tau^2 - j \tau \alpha_0^2 + j^3 \tau) / [\quad " \quad]$$

and

$$D_{34} = - \beta_0^2 (\tau - e h \beta_0^2) / [(\beta_0^2 - \tau j^2)^2 - j \tau (j \tau - e h j \beta_0^2)]$$

$$D_{36} = D_{54} = [j \tau (j^2 \tau - \beta_0^2) - \tau (j \tau - e h j \beta_0^2)] / [\quad " \quad]$$

$$D_{56} = - \beta_0^2 \tau / [\quad " \quad]$$

The last four equations (56 g-j) are rewritten as

$$\begin{aligned}
 \tilde{B}_3 &= \frac{\lambda_D}{\beta_D \Delta_D} \tilde{B}_4 + \frac{2}{\pi \alpha_D \beta_D \Delta_D} \tilde{A}_4 \\
 \tilde{A}_3 &= \frac{-2}{\pi \alpha_D \beta_D \Delta_D} \tilde{B}_4 + \frac{q_D}{\alpha_D \Delta_D} \tilde{A}_4 \\
 \tilde{B}_5 &= \frac{\lambda_R}{\beta_R \Delta_R} \tilde{B}_6 + \frac{2}{\pi \alpha_R \beta_R \Delta_R} \tilde{A}_6 \\
 \tilde{A}_5 &= \frac{-2}{\pi \alpha_R \beta_R \Delta_R} \tilde{B}_6 + \frac{q_R}{\alpha_R \Delta_R} \tilde{A}_6
 \end{aligned}
 \tag{60}$$

where

$$\begin{aligned}
 p(a,b) &= J_j(a) Y_j(b) - J_j(b) Y_j(a) \\
 q(a,b) &= J_j(a) Y_j'(b) - J_j'(b) Y_j(a) \\
 \lambda(a,b) &= J_j'(a) Y_j(b) - J_j(b) Y_j'(a) \\
 \Delta(a,b) &= J_j''(a) Y_j'(b) - J_j'(b) Y_j''(a)
 \end{aligned}$$

and $p_D = p(\alpha_D, \beta_D)$ etc. We then have a four by four system. Using

$K_D = 2/(\pi \alpha_D \beta_D S_D)$ and $K_R = 2/(\pi \alpha_R \beta_R S_R)$ we have,

$$\begin{aligned}
 & \left[\begin{array}{cccc}
 \kappa_D & 0 & [\lambda_D/\beta_D \Delta_D - D_{34}] & -D_{36} \\
 0 & \kappa_R & -D_{54} & [\lambda_R/\beta_R \Delta_R - D_{56}] \\
 [q_D/\alpha_D \Delta_D - C_{34}] & -C_{36} & -\kappa_D & 0 \\
 -C_{54} & [q_R/\alpha_R \Delta_R - C_{56}] & 0 & -\kappa_R
 \end{array} \right] \begin{bmatrix} \tilde{A}_4 \\ \tilde{A}_6 \\ \tilde{B}_4 \\ \tilde{B}_6 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \kappa_3 \\ \kappa_5 \end{bmatrix} \\
 & \tag{61}
 \end{aligned}$$

with a solution given explicitly in appendix A.

By backwards substitution, we can evaluate all of the unknown coefficients. These must, of course, agree with the well-known solution for the concentric case and are carried out in detail to serve as a procedural check for the higher order solutions.

E. FIRST ORDER SOLUTION

Again, we use the boundary conditions and acoustic fluid boundary integral equation directly for the first order coefficients. The outer boundary shell integral equation on ϕ , equation (44), integrates to

$$\begin{aligned}
 (62) \quad 0 = & \sum_{n=0}^{\infty} \left\{ \left[-B_{3n}^{(1)} \cdot b \cdot k_0 \cdot 2\pi J_n'(k_0 b) + B_{4n}^{(1)} \cdot b \cdot 2\pi J_n(k_0 b) \right] H_n(k_0 b) \right. \\
 & + \left. \left[A_{3n}^{(1)} \cdot a \cdot k_0 \cdot 2\pi J_n'(k_0 a) - A_{4n}^{(1)} \cdot a \cdot 2\pi J_n(k_0 a) \right] H_n(k_0 b) \right\} \\
 & \cdot \left\{ \frac{\sin}{\cos} \right\} (n\theta') \\
 & + \sum_{n=0}^{\infty} \left\{ \left[A_{3n}^{(0)} \cdot a \cdot \pi \cdot k_0^2 \cdot J_n'(k_0 a) - A_{4n}^{(0)} \cdot a \cdot \pi \cdot k_0 J_n(k_0 a) \right] \right. \\
 & \cdot \left. \left[\left\{ \frac{\sin}{\cos} \right\} ((n-1)\theta') \cdot H_{n-1}(k_0 b) - \left\{ \frac{\sin}{\cos} \right\} ((n+1)\theta') \cdot H_{n+1}(k_0 b) \right] \right\}
 \end{aligned}$$

and a similar equation on ψ with A_{5n} , A_{6n} , B_{5n} , B_{6n} and k_R replacing A_{3n} , A_{4n} , B_{3n} , B_{4n} and k_D .

The inner boundary shell integral equation on ϕ becomes

$$\begin{aligned}
 (63) \quad 0 = & \sum_{n=0}^{\infty} \left[\left\{ -B_{3n}^{(0)} \cdot b \cdot k_0^2 \cdot \pi H_n'(k_0 b) + B_{4n}^{(0)} \cdot b \cdot k_0 \cdot \pi H_n(k_0 b) \right\} \right. \\
 & \cdot \left. \left\{ \left\{ \frac{\sin}{\cos} \right\} [(n+1)\theta] \cdot J_{n+1}(k_0 a) - \left\{ \frac{\sin}{\cos} \right\} [(n-1)\theta] \cdot J_{n-1}(k_0 a) \right\} \right] +
 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{n=0}^{\infty} \left\{ -B_{3n}^{(1)} \cdot b \cdot 2\pi \cdot k_0 \cdot H_n'(k_0 b) + B_{4n}^{(1)} \cdot b \cdot 2\pi \cdot H_n(k_0 b) \right. \\
 & \quad \left. + A_{3n}^{(1)} \cdot a \cdot 2\pi \cdot k_0 \cdot H_n'(k_0 a) - A_{4n}^{(1)} \cdot a \cdot 2\pi \cdot H_n(k_0 a) \right\} \\
 & \quad \cdot \left\{ J_n(k_0 a) \cdot \left\{ \begin{matrix} \sin \\ \cos \end{matrix} \right\} (n\theta) \right\}]
 \end{aligned}$$

with again a similar equation on ψ .

Clearly if there is only one zeroth order mode, $n = j$, there are only two first order modes, $n = j+1$ and $n = j-1$. Consider the $j+1$ mode first.

The governing equations are

$$\begin{aligned}
 (64) \quad & -B_{3j+1}^{(1)} \cdot k_0 b \cdot J_{j+1}'(k_0 b) + B_{4j+1}^{(1)} \cdot b \cdot J_{j+1}(k_0 b) \\
 & + A_{3j+1}^{(1)} \cdot k_0 a \cdot J_{j+1}'(k_0 a) - A_{4j+1}^{(1)} \cdot a \cdot J_{j+1}(k_0 a) \\
 & = \frac{1}{2} \left[A_{3j}^{(0)} \cdot k_0^2 a \cdot J_j'(k_0 a) - A_{4j}^{(0)} \cdot k_0 a \cdot J_j(k_0 a) \right] \\
 & = \frac{1}{2} \left[B_{3j}^{(0)} \cdot k_0^2 b \cdot J_j'(k_0 b) - B_{4j}^{(0)} \cdot k_0 b \cdot J_j(k_0 b) \right]
 \end{aligned}$$

and

$$\begin{aligned}
 & -B_{3j+1}^{(1)} \cdot k_0 b \cdot H_{j+1}'(k_0 b) + B_{4j+1}^{(1)} \cdot b \cdot H_{j+1}(k_0 b) \\
 & + A_{3j+1}^{(1)} \cdot k_0 a \cdot H_{j+1}'(k_0 a) - A_{4j+1}^{(1)} \cdot a \cdot H_{j+1}(k_0 a)
 \end{aligned}$$

$$(65) \quad = \frac{1}{2} \left[B_{3j}^{(0)} \cdot k_0^2 b \cdot H_j'(k_0 b) - B_{4j}^{(0)} \cdot k_0 b \cdot H_j(k_0 b) \right]$$

and two similar equations on ψ . The $j-1$ mode is essentially the same with a sign change for the inhomogeneous (zeroth order) terms

$$\begin{aligned}
 (66) \quad & - B_{3j-1}^{(1)} k_0 b J_{j-1}'(k_0 b) + B_{4j-1}^{(1)} b \cdot J_{j-1}(k_0 b) \\
 & + A_{3j-1}^{(1)} k_0 a J_{j-1}'(k_0 a) - A_{4j-1}^{(1)} a \cdot J_{j-1}(k_0 a) \\
 & = \frac{-1}{2} [A_{3j}^{(0)} k_0^2 a J_j'(k_0 a) - A_{4j}^{(0)} k_0 a \cdot J_j(k_0 a)]
 \end{aligned}$$

and

$$\begin{aligned}
 (67) \quad & - B_{3j-1}^{(1)} k_0 b \cdot H_{j-1}'(k_0 b) + B_{4j-1}^{(1)} b \cdot H_{j-1}(k_0 b) \\
 & + A_{3j-1}^{(1)} k_0 a H_{j-1}'(k_0 a) - A_{4j-1}^{(1)} a \cdot H_{j-1}(k_0 a) \\
 & = \frac{-1}{2} [B_{3j}^{(0)} k_0^2 b H_j'(k_0 b) - B_{4j}^{(0)} k_0 b \cdot H_j(k_0 b)]
 \end{aligned}$$

and two similar equations on ψ .

We note that the homogeneous part of these equations plus the first order boundary conditions and the acoustic boundary integral equation are identical to the zeroth order equations with $j+1$ and $j-1$ for the first order coefficients replacing j for the zeroth order coefficients. The inhomogeneous terms are modified and now appear only in the four shell boundary integral equations. Then we can reuse the solution scheme for the zeroth order j mode coefficients to solve for the first order $(j+1)$ and $(j-1)$ mode coefficients with a minor change. We again may write equations (65) and (67) with real coefficients by combining with the previous equations - equivalently, we replace H_n by Y_n in these equations.

We may again nondimensionalize using an additional length scale factor of a for these first order terms which must, in final form, be multiplied by the length δ in order to be compared to the zeroth order terms; e.g.,

$$\tilde{A}_{3j+1}^{(1)} = A_{3j+1}^{(1)} \cdot a / \Phi_j$$

$$\tilde{A}_{4j+1}^{(1)} = A_{4j+1}^{(1)} \cdot a^2 / \Phi_j$$

$$\tilde{B}_{3j+1}^{(1)} = B_{3j+1}^{(1)} \cdot a / \Phi_j$$

$$\tilde{B}_{4j+1}^{(1)} = B_{4j+1}^{(1)} \cdot ab / \Phi_j$$

etc.

The same scale factor is used for both $A^{(1)}$ and $B^{(1)}$ to keep the same form of equations as in the zeroth order case. We then have

$$(a) \quad 0 = [\tau(j+1)^2 - \alpha_0^2] \tilde{A}_{3j+1}^{(1)} + \tau [-\tilde{A}_{4j+1}^{(1)} + (j+1) \tilde{A}_{5j+1}^{(1)} - (j+1) \tilde{A}_{6j+1}^{(1)}]$$

$$(b) \quad 0 = -\tau(j+1) \tilde{A}_{3j+1}^{(1)} + \tau(j+1) \tilde{A}_{4j+1}^{(1)} + [\alpha_0^2 - (j+1)^2 \tau] \tilde{A}_{5j+1}^{(1)} + \tau \tilde{A}_{6j+1}^{(1)}$$

$$(c) \quad 0 = \tilde{B}_{1j+1}^{(1)} + [\tau(j+1)^2 - \beta_0^2] \tilde{B}_{3j+1}^{(1)} + \tau [-\tilde{B}_{4j+1}^{(1)} + (j+1) \tilde{B}_{5j+1}^{(1)} - (j+1) \tilde{B}_{6j+1}^{(1)}]$$

(68)

$$(d) \quad 0 = -\tau(j+1) \tilde{B}_{3j+1}^{(1)} + \tau(j+1) \tilde{B}_{4j+1}^{(1)} + [\beta_0^2 - (j+1)^2 \tau] \tilde{B}_{5j+1}^{(1)} + \tau \tilde{B}_{6j+1}^{(1)}$$

$$(e) \quad 0 = -\tilde{B}_{2j+1}^{(1)} + \beta_0^2 [\tilde{B}_{4j+1}^{(1)} - (j+1) \tilde{B}_{5j+1}^{(1)}]$$

$$(f) \quad 0 = -\tilde{B}_{2j+1}^{(1)} + \tilde{B}_{1j+1}^{(1)} / h_{j+1}$$

$$(g) \quad L_1 = \frac{\alpha_0}{2} [\beta_0 J_j'(\beta_0) \tilde{B}_{3j}^{(0)} - \tilde{B}_{4j}^{(0)} J_j(\beta_0)]$$

$$= -\tilde{B}_{3j+1}^{(1)} \beta_0 J_{j+1}'(\beta_0) + \tilde{A}_{3j+1}^{(1)} \alpha_0 J_{j+1}'(\alpha_0) + \tilde{B}_{4j+1}^{(1)} J_{j+1}(\beta_0) - \tilde{A}_{4j+1}^{(1)} J_{j+1}(\alpha_0)$$

$$(h) \quad L_2 = \frac{\alpha_R}{2} [\tilde{B}_{5j}^{(0)} \beta_R J_j'(\beta_R) - \tilde{B}_{6j}^{(0)} J_j(\beta_R)]$$

$$= -\tilde{B}_{5j+1}^{(1)} \beta_R J_{j+1}'(\beta_R) + \tilde{A}_{5j+1}^{(0)} \alpha_R J_{j+1}'(\alpha_R) + \tilde{B}_{6j+1}^{(1)} J_{j+1}(\beta_R) - \tilde{A}_{6j+1}^{(1)} J_{j+1}(\alpha_R)$$

$$(i) \quad L_3 = \frac{\alpha_D}{2} [\tilde{B}_{3j}^{(0)} \beta_D Y_j'(\beta_D) - \tilde{B}_{4j}^{(0)} Y_j(\beta_D)]$$

$$= -\tilde{B}_{3j+1}^{(1)} \beta_D Y_{j+1}'(\beta_D) + \tilde{A}_{3j+1}^{(1)} \alpha_D Y_{j+1}'(\alpha_D) + \tilde{B}_{4j+1}^{(1)} Y_{j+1}(\beta_D) - \tilde{A}_{4j+1}^{(1)} Y_{j+1}(\alpha_D)$$

$$(j) \quad L_4 = \frac{\alpha_R}{2} [\tilde{B}_{5j}^{(0)} \beta_R Y_j'(\beta_R) - \tilde{B}_{6j}^{(0)} Y_j(\beta_R)]$$

$$= -\tilde{B}_{5j+1}^{(1)} \beta_R Y_{j+1}'(\beta_R) + \tilde{A}_{5j+1}^{(1)} \alpha_R Y_{j+1}'(\alpha_R) + \tilde{B}_{6j+1}^{(1)} Y_{j+1}(\beta_R) - \tilde{A}_{6j+1}^{(1)} Y_{j+1}(\alpha_R)$$

and a similar set of equations with $j-1$ replacing $j+1$ and $-L_1, -L_2, -L_3, -L_4$ replacing L_1, L_2, L_3, L_4 .

We see at this point that we have two separate ten by ten systems of complex algebraic equations - each very similar to those of the zeroth order case. We again may reduce the unknowns to yield a feasible arithmetic problem - consider the $j+1$ system as an example:

$$(69) \quad \begin{aligned} (a) \quad \tilde{B}_{2j+1}^{(1)} &= \beta_D^2 [\tilde{B}_{4j+1}^{(1)} - (j+1) \tilde{B}_{5j+1}^{(1)}] \\ (b) \quad \tilde{B}_{1j+1}^{(1)} &= h \beta_D^2 [\tilde{B}_{4j+1}^{(1)} - (j+1) \tilde{B}_{5j+1}^{(1)}] \\ (c) \quad \tilde{A}_{3j+1}^{(1)} &= C_{34} \tilde{A}_{4j+1}^{(1)} + C_{36} \tilde{A}_{6j+1}^{(1)} \\ (d) \quad \tilde{A}_{5j+1}^{(1)} &= C_{54} \tilde{A}_{4j+1}^{(1)} + C_{56} \tilde{A}_{6j+1}^{(1)} \\ (e) \quad \tilde{B}_{3j+1}^{(1)} &= D_{34} \tilde{B}_{4j+1}^{(1)} + D_{36} \tilde{B}_{6j+1}^{(1)} \\ (f) \quad \tilde{B}_{5j+1}^{(1)} &= D_{54} \tilde{B}_{4j+1}^{(1)} + D_{56} \tilde{B}_{6j+1}^{(1)} \end{aligned}$$

where the C's and D's are the same as before with $j+1$ replacing n . Clearly the K_3 and K_5 are no longer present since the inhomogeneity in the system has moved to the last four equations.

Those last equations are now

$$\begin{aligned}
 (a) \quad \kappa_D \tilde{A}_{4j+1}^{(1)} + \left(\frac{\rho_D}{\Delta_D \beta_D} - D_{34} \right) \tilde{B}_{4j+1}^{(1)} - D_{36} \tilde{B}_{6j+1}^{(1)} &= \frac{L_3 J_{j+1}'(\alpha_D) - L_1 Y_{j+1}'(\alpha_D)}{\beta_D \Delta_D} = R_1 \\
 (b) \quad \kappa_R \tilde{A}_{6j+1}^{(1)} - D_{54} \tilde{B}_{4j+1}^{(1)} + \left(\frac{\rho_R}{\Delta_R \beta_R} - D_{56} \right) \tilde{B}_{6j+1}^{(1)} &= \frac{L_4 J_{j+1}'(\alpha_R) - L_2 Y_{j+1}'(\alpha_R)}{\beta_R \Delta_R} = R_2 \\
 (70) \quad (c) \quad \left[\frac{\rho_D}{\alpha_D \Delta_D} - C_{34} \right] \tilde{A}_{4j+1}^{(1)} - C_{36} \tilde{A}_{6j+1}^{(1)} - \kappa_D \tilde{B}_{4j+1}^{(1)} &= \frac{L_3 J_{j+1}'(\beta_D) - L_1 Y_{j+1}'(\beta_D)}{\alpha_D \Delta_D} = R_3 \\
 (d) \quad -C_{54} \tilde{A}_{4j+1}^{(1)} + \left[\frac{\rho_R}{\alpha_R \Delta_R} - C_{56} \right] \tilde{A}_{6j+1}^{(1)} - \kappa_R \tilde{B}_{6j+1}^{(1)} &= \frac{L_4 J_{j+1}'(\beta_R) - L_2 Y_{j+1}'(\beta_R)}{\alpha_R \Delta_R} = R_4
 \end{aligned}$$

remembering that the C, D, p, q, r and s are all modified to have $j+1$ replace n .

The equations for the $(j-1)$ mode are identical to the above with the corresponding changes of $j-1$ for $j+1$ everywhere and with the sign of the R terms reversed. The solution procedure is the same as for the zeroth order, and is also given in appendix A.

To convert these first order coefficients to a form suitable for comparison to the zeroth order solution, they must be multiplied by (δ/a) , which is a small number in order to have the first order be a sufficient correction to the zeroth order solution. Higher orders may be calculated in the same manner -- only a new set of L_i need be calculated since the same solution procedure will hold for higher orders as did for the first order, i.e., the same basic program.

CALCULATIONS AND CONCLUSIONS

The primary application of this solution is to a thin steel shell submerged in water. The parameters used were $a = 300$ cm, $b = 303$ cm, $c = 146000$ cm/sec, $c_D = 496000$ cm/sec and $c_R = 270000$ cm/sec with $\rho_S = 7.7$ gm/cm³ and $\rho_F = 1.0$ gm/cm³. Results for the exterior pressure B_1 , are shown in Figures 2 and 3 for the case $j = 0$ and $j = 2$ respectively for a unit interior forcing pressure. Results for the first order solutions are given on the same figure as the zeroth order solution, but to a scale 50 times as great. Since the first order terms must be multiplied by δ/a to determine their contribution to the exterior pressure field, the actual exterior pressure will depend on δ . For $\delta = 0.3$ cm, $\delta/a = 10^{-3}$ and the first order contribution is generally negligible. There are regions, however, where the zeroth order solution is minimum, e.g. near 300 rad/sec for $j = 2$. In this region, the first order contributions may be comparable to the zeroth order, but this is for a very limited frequency band. Figure 4 shows results for a thick shell with $a = 200$ cm and the remaining parameters unchanged. In all cases, ω runs from 200 to 9000 rad/sec. Here the first order and zeroth order solutions are plotted on the same scale and are very similar. This implies the first order contribution, when multiplied by δ/a will be negligible for any example in which δ/a is small enough to allow a first order theory to be used.

The general conclusions to be drawn from this work are first that such an approach is applicable to radiating submerged "nonconcentric" elastic shells and second that nearest neighbor modes are excited in the fluid in the higher order solutions for a single mode forcing function.

Both of these conclusions have already been drawn in the acoustic shell case, reference [1]; we have shown here that they are applicable to the physically more meaningful elastic shell case as well, with some additional effort. A specific result based on these two examples is that the influence of nonconcentricity is essentially negligible in most cases where δ/a is small. Since higher order solutions may be computed with the same basic program, extensions to larger values of (δ/a) are clearly possible for thicker shells. Such analytical solutions may be useful both in their own right and even more importantly as checks on general numerical schemes which might not be otherwise properly constructed to include (or recognize) such effects as nearest neighbor modes.

REFERENCES

1. R. P. Shaw and G. Tai, "Time Harmonic Acoustic Radiation from Non-concentric Circular Cylinders", J.A.S.A. 56(5), pp. 1437-1443 (1974).
2. W. Thompson, "Acoustic Radiation from a Spherical Source Embedded Eccentrically Within a Fluid Sphere", J.A.S.A. 54(), pp. 1694-1707 (1973).
3. R. P. Shaw, "An Integral Equation Approach to Acoustic Radiation and Scattering", Topics in Ocean Engineering, II, edited: C. Bretschneider, Gulf Publishing Co., Houston, Texas, 1970. [Note: The equations for time harmonic case use inward rather than outward normal--this provides an overview of boundary integral equations].
4. A. E. H. Love, A Treatise on the Mathematical Theory of Elasticity, Chap. XIII, Dover Publishing Co., N.Y.C., N. Y. (Fourth Edition) 1944.
5. Love, ref. [4], has an excellent discussion of this except that his definition uses h_i as the inverse of the usual definition of metric coefficient. As a result the equations in Chap. 1 have h_i inverted from the h_i^* used in this paper.

APPENDIX A

Define

$$QASD = q_D / \alpha_D s_D - C_{34}$$

$$QASR = q_R / \alpha_R s_R - C_{56}$$

$$RBSD = r_D / \beta_D s_D - D_{34}$$

Then

$$D = \begin{vmatrix} K_D & 0 & RBSD & -D_{36} \\ 0 & K_R & -D_{54} & RBSR \\ QASD & -C_{36} & -K_D & 0 \\ -C_{54} & QASR & 0 & -K_R \end{vmatrix}$$

and

$$N_4 = \begin{vmatrix} R_1 & 0 & RBSD & -D_{36} \\ R_2 & K_R & -D_{54} & RBSR \\ R_3 & -C_{36} & -K_D & 0 \\ R_4 & QASR & 0 & -K_R \end{vmatrix}$$

$$N_6 = \begin{vmatrix} K_D & R_1 & RBSD & -D_{36} \\ 0 & R_2 & -D_{54} & RBSR \\ QASD & R_3 & -K_D & 0 \\ -C_{54} & R_4 & 0 & -K_R \end{vmatrix}$$

(Zeroth order)

N_4 :

$$R_1 = 0, R_2 = 0, R_3 = K_3, R_4 = K_5$$

N_6 :

such that

$$\tilde{A}_{4j}^{(0)} = N_4/D$$

$$\tilde{A}_{6j}^{(0)} = N_6/D$$

(First order)

D (j+1) = D with j+1 replacing j in all terms

N₄ (j+1):

R₁, R₂, R₃, R₄ defined in equation 70

N₆ (j+1):

and a similar set for j-1 contribution.

$$\tilde{A}_{3j}^{(0)} = K_3 + C_{34} \tilde{A}_{4j}^{(0)} + C_{36} \tilde{A}_{6j}^{(0)}$$

$$\tilde{A}_{5j}^{(0)} = K_5 + C_{54} \tilde{A}_{4j}^{(0)} + C_{56} \tilde{A}_{6j}^{(0)}$$

$$\tilde{B}_{4j}^{(0)} = K_D - K_3 + QASD * \tilde{A}_{4j}^{(0)} - C_{36} \tilde{A}_{6j}^{(0)}$$

$$\tilde{B}_{6j}^{(0)} = K_R - K_5 - C_{54} * \tilde{A}_{4j}^{(0)} + QASR * \tilde{A}_{6j}^{(0)}$$

APPENDIX B

A simple test calculation is the "transparent" shell, i.e. $\rho_F = \rho_S$, $\mu = 0$ and $c_D = c$. Since the pressure is applied at the inner surface, $r = a$, we have

$$p(r, \theta) = \frac{P_n H_n^{(1)}(kr)}{H_n^{(1)}(ka)} \cdot \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta)$$

On an "offset" circle, we have $r' = b$ given by $r = r' + \delta \cos \theta + \dots$

$$r = b + \delta \cos \theta + \dots$$

and

$$\begin{aligned} p(r'=b, \theta) &= \frac{P_n}{H_n^{(1)}(ka)} \left[H_n^{(1)}(kb) + k \cdot \delta \cdot \cos \theta \cdot [H_n^{(1)}(kb)]' + \dots \right] \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta) \\ &= p_n^{(0)}(b, \theta) + \delta p_{n+1}^{(1)}(b) \cdot \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} [(n+1)\theta] + \delta p_{n-1}^{(1)}(b) \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} [(n-1)\theta] + \dots \end{aligned}$$

where

$$p_n^{(0)}(b, \theta) = P_n \cdot \frac{H_n^{(1)}(kb)}{H_n^{(1)}(ka)} \cdot \begin{Bmatrix} \sin \\ \cos \end{Bmatrix} (n\theta)$$

$$p_{n+1}^{(1)}(b) = P_n \cdot \frac{k}{2} \cdot \frac{[H_n^{(1)}(kb)]'}{[H_n^{(1)}(ka)]} = p_{n-1}^{(1)}(b)$$

If we return to the integral equation approach, a direct substitution of these parameters into the zeroth order equations leads to

$$B_{in}^{(0)} = P_n \cdot \frac{H_n(\beta)}{H_n(\alpha)}$$

The first order equations lead to

$$B_{1, n+1}^{(1)} = -\frac{k}{2} \frac{H_{n+1}^{(1)}(kb)}{H_n^{(1)}(ka)} P_n$$

$$B_{1, n-1}^{(1)} = -\frac{k}{2} \frac{H_{n-1}^{(1)}(kb)}{H_n^{(1)}(ka)} P_n$$

We must bear in mind, however, that these are values for the pressure at the outer surface measured in terms of θ' and not in terms of θ . We need to convert $\left\{ \frac{\sin}{\cos} \right\} (n\theta)$. We need to convert $\left\{ \frac{\sin}{\cos} \right\} (n\theta)$ to $\left\{ \frac{\sin}{\cos} \right\} (n\theta')$ which modifies our first order results. We have the definition

$$\theta' = \theta + \sin^{-1} \left\{ \frac{\delta}{b} \sin \theta \right\}$$

and to a first order

$$\left\{ \frac{\sin}{\cos} \right\} (n\theta) = \left\{ \frac{\sin}{\cos} \right\} (n\theta') - \delta \cdot \frac{n}{2} \left[\left\{ \frac{\sin}{\cos} \right\} [(n+1)\theta] + \left\{ \frac{\sin}{\cos} \right\} [(n-1)\theta] \right] + \dots$$

Then

$$\begin{aligned} p(r'=b, \theta') &= B_{1, n}^{(0)} \left\{ \frac{\sin}{\cos} \right\} (n\theta') \\ &+ \delta \cdot B_{1, n+1}^{(1)} \left\{ \frac{\sin}{\cos} \right\} [(n+1)\theta'] + \delta \cdot B_{1, n-1}^{(1)} \left\{ \frac{\sin}{\cos} \right\} [(n-1)\theta'] + \dots \\ &= P_n \cdot \frac{H_n^{(1)}(kb)}{H_n^{(1)}(ka)} \cdot \left\{ \frac{\sin}{\cos} \right\} (n\theta) + \frac{\delta \cdot P_n \cdot k}{2 H_n^{(1)}(ka)} \left[-H_{n+1}^{(1)}(kb) \right. \\ &+ \left. \frac{n}{2b} H_n^{(1)}(kb) \right] \left\{ \frac{\sin}{\cos} \right\} [(n+1)\theta] + \frac{\delta P_n k}{2 H_n^{(1)}(ka)} \left[H_{n+1}^{(1)}(kb) + \frac{n}{2b} H_n^{(1)}(kb) \right] \left\{ \frac{\sin}{\cos} \right\} [(n-1)\theta] + \dots \end{aligned}$$

which agrees with the previous solution.




Figure 1. Geometry of Nonconcentric Cylinder




Figure 2: Exterior Pressure Coefficients for Thin Shell for $j = 0$ Mode.




Figure 3: Exterior Pressure Coefficients for Thin Shell for $j = 2$ Mode.




Figure 4: Exterior Pressure Coefficients for Thick Shell for $j = 0$ Mode.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing information must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) The Research Foundation of the State University of New York	2b. REPORT SECURITY CLASSIFICATION Unclassified 2b. GROUP
--	---

3. REPORT TITLE
Time Harmonic Acoustic Radiation from a Submerged Elastic Shell Defined by Nonconcentric Cylinders.

4. DESCRIPTIVE NOTES (Type of report and, inclusive dates)
Summary *rept.*

5. AUTHOR (Last name, middle initial, first name)
Richard P. Shaw 114 97

6. REPORT DATE 11 September, 1976	7a. TOTAL NO. OF PAGES 39	7b. NO. OF REFS 4
--------------------------------------	------------------------------	----------------------

8a. CONTRACT OR GRANT NO. 15 N00014-75-C0302	9a. ORIGINATOR'S REPORT NUMBER(S) Report No. 97 Department of Engineering Science, Aero- space Engineering and Nuclear Engineering
---	---

9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

10. DISTRIBUTION STATEMENT
Approved for Public Release. Distribution Unlimited

11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Office of Naval Research - Structural Mechanics
-------------------------	--

13. ABSTRACT
The acoustic radiation from an elastic cylindrical shell defined by nonconcentric cylindrical surfaces and submerged in an acoustic fluid is studied. The driving mechanism is a time harmonic internal pressure. In particular, the influence of the nonconcentricity parameter, δ , on the resulting radiation field is examined. The solution is obtained by means of an expansion in powers of this parameter about the concentric solution, $\delta=0$. Each order of solution is solved in closed form by means of boundary integral equation methods. Results are given for the zeroth and first order solutions and it is seen that a j^{th} driving mode will introduce nearest neighbor $((j-1)$ and $(j+1))$ modes in the first order solution.

406156

B

PART 1 - GOVERNMENT

Administrative & Liaison Activities

Chief of Naval Research
Department of the Navy
Arlington, Virginia 22111
Attn: Code 439 (2)
471

Director
ONR Branch Office
495 Summer Street
Boston, Massachusetts 02210

Director
ONR Branch Office
219 S. Dearborn Street
Chicago, Illinois 60604

Director
Naval Research Laboratory
Attn: Library, Code 2029 (ONRL)
Washington, D.C. 20390 (6)

Commanding Officer
ONR Branch Office
207 West 24th Street
New York, New York 10011

Director
ONR Branch Office
1030 E. Green Street
Pasadena, California 91101

U.S. Naval Research Laboratory
Attn: Technical Information Div.
Washington, D.C. 20390 (4)

Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314 (20)

Army

Commanding Officer
U.S. Army Research Office Durham
Attn: Mr. J.J. Murray
CRD-AA-IP
Box CM, Duke Station
Durham, North Carolina 27706

Commanding Officer
ANXMR-ATL
Attn: Mr. J. Bluhm
U.S. Army Materials Res. Agcy.
Watertown, Massachusetts 02172

Watervliet Arsenal
MAGGS Research Center
Watervliet, New York 12189
Attn: Director of Research

Redstone Scientific Info. Center
Chief, Document Section
U.S. Army Missile Command
Redstone Arsenal, Alabama 35809

Army R & D Center
Fort Belvoir, Virginia 22060

Technical Library
Aberdeen Proving Ground
Aberdeen, Maryland 21005

Navy

Commanding Officer and Director
Naval Ship Research & Development Cen.
Washington, D.C. 20007
Attn: Code 042 (Tech. Lib. Br.)
700 (Struc. Mech. Lab.)
720
725
727
012.2 (Dr. W.J. Sette)

Naval Weapons Laboratory
Dahlgren, Virginia 22448

Naval Research Laboratory
Washington, D.C. 20390
Attn: Code 8400
8410
8430
8440
6300
6305
6380

Undersea Explosion Research Div:
Naval Ship R & D Center
Norfolk Naval Shipyard
Portsmouth, Virginia 23709
Attn: Dr. Schauer
Code 780

Naval Ship R & D Center
Annapolis Division
Annapolis, Maryland 21402
Attn: Code A800, Mr. W.L. Williams

Technical Library
Naval Underwater Weapons Center
Pasadena Annex
3202 E. Foothill Blvd.
Pasadena, California 91107

U.S. Naval Weapons Center
China Lake, California 93557
Attn: Code 4520 Mr. Ken Bischel

U.S. Naval Ordnance Laboratory
Mechanics Division
RFD 1, White Oak
Silver Spring, Maryland 20910

U.S. Naval Ordnance Laboratory
Attn: Mr. H.A. Perry, Jr.
Non-Metallic Materials Division
Silver Spring, Maryland 20910

Technical Director
U.S. Naval Undersea R & D Center
San Diego, California 92132

Supervisor of Shipbuilding
U.S. Navy
Newport News, Virginia 23607

Technical Director
Mare Island Naval Shipyard
Vallejo, California 94592

U.S. Naval Ordnance Station
Attn: Mr. Garet Bornstein
Research & Development Division
Indian Head, Maryland 20640

Chief of Naval Operations
Department of the Navy
Washington, D.C. 20350
Attn: Code OP-077

Deep Submergence Systems
Naval Ship Systems Command
Code 39522
Department of the Navy
Washington, D.C. 20360
Attn: Chief Scientist

Director, Aero Mechanics
Naval Air Development Center
Johnsville
Warminster, Pennsylvania 18974

Naval Air Systems Command
Dept. of the Navy
Washington, D.C. 20360
Attn: NAIR 320 Aero. & Structures
5320 Structures
604 Tech. Library
52031F Materials

Naval Facilities Engineering
Command
Department of the Navy
Washington, D.C. 20360
Attn: NFAC 03 Res. & Development
04 Engineering & Design
14114 Tech. Library

Naval Ship Systems Command
Dept. of the Navy
Washington, D.C. 20360
Attn: NSHIP 031 Ch. Scientists for
0342 Ship Mats. & Struct.
2052 Tech. Library

Naval Ship Engineering Center
Prince George Plaza
Hyattsville, Maryland 20782
Attn: NSEC 6100 Ship Sys. Engr. & D
6102C Computer-Aided Shi
6105 Ship Protection
6110 Ship Concept Design
6120 Hull Div.
61200 Hull Div.
6128 Surface Ship Struct
6129 Submarine Struct.

Naval Ordnance Systems Command
Dept. of the Navy
Washington, D.C. 20360
Attn: NORD 03 Res. & Technology
035 Weapons Dynamics
9132 Tech. Library

Engineering Department
U.S. Naval Academy
Annapolis, Maryland 21402

Air Force

Commander WADD
Wright-Patterson Air Force Base
Dayton, Ohio 45433
Attn: Code WHEMDD
AFFDL (FDSS)
Structures Division
AFLC (MCEBA)
Code WWRC
AFML (MAAM)

Commander
Chief, Applied Mechanics Group
U.S. Air Force Inst. of Tech.
Wright-Patterson Air Force Base
Dayton, Ohio 45433

Chief, Civil Engineering Branch
MLRC, Research Division
Air Force Weapons Laboratory
Hickam AFB, New Mexico 87117

Air Force Office of Scientific Res.
1400 Wilson Blvd.
Arlington, Virginia 22209
Attn: Mech. Div.

NASA

Structures Research Division
National Aeronautics & Space Admin.
Langley Research Center
Langley Station
Hampton, Virginia 23365
Attn: Mr. R.R. Heldenfels, Chief

National Aeronautic & Space Admin.
Associate Administrator for Advanced
Research & Technology
Washington, D.C. 20546

Scientific & Tech. Info. Facility
NASA Representative (S-AK/DE)
P.O. Box 5700
Bethesda, Maryland 20014

Other Government Activities

Technical Director
Marine Corps Development & Educ. Command
Quantico, Virginia 22134

Director
National Bureau of Standards
Washington, D.C. 20234
Attn: Mr. B.L. Wilson, RM 219

National Science Foundation
Engineering Division
Washington, D.C. 20550

Director
STBS
Defense Atomic Support Agency
Washington, D.C. 20350

Commander Field Command
Defense Atomic Support Agency
Sandia Base
Albuquerque, New Mexico 87115

Chief, Defense Atomic Support Agcy.
Blast & Shock Division
The Pentagon
Washington, D.C. 20301

Director Defense Research & Engr.
Technical Library
Room 3C-128
The Pentagon
Washington, D.C. 20301

Chief, Airframe & Equipment
FS-120
Office of Flight Standards
Federal Aviation Agency
Washington, D.C. 20553

Chief of Research and Development
Maritime Administration
Washington, D.C. 20235

Mr. Milton Shaw, Director
Div. of Reactor Develo. & Technology
Atomic Energy Commission
Germantown, Maryland 20767

Ship Hull Research Committee
National Research Council
National Academy of Sciences
2101 Constitution Avenue
Washington, D.C. 20418
Attn: Mr. A.R. Lytle

PART 2 - CONTRACTORS AND OTHER
TECHNICAL CONSULTANTS

Universities

Professor J.R. Rice
Division of Engineering
Brown University
Providence, Rhode Island 02906

Dr. J. Tinsley Oden
Dept. of Engr. Mechs.
University of Alabama
Huntsville, Alabama 35894

Professor R.S. Rivlin
Center for the Application of Mathematics
Lehigh University
Bethlehem, Pennsylvania 18015

Professor Julius Miklowitz
Division of Engr. & Applied Sciences
California Institute of Technology
Pasadena, California 91109

Professor George Sih
Department of Mechanics
Lehigh University
Bethlehem, Pennsylvania 18015

Dr. Harold Liebowitz, Dean
School of Engrg. & Applied Science
George Washington University
725 23rd Street
Washington, D.C. 20006

Professor Eli Sternberg
Division of Engr. & Applied Sciences
California Institute of Technology
Pasadena, California 91109

Professor Burt Paul
University of Pennsylvania
Towne School of Civil and Mechanical Engrg.
Room 113 Towne Building
220 S. 33rd Street
Philadelphia, Penna. 19104

Professor S.B. Dong
University of California
Department of Mechanics
Los Angeles, California 90024

Professor Paul M. Naghi
Div. of Applied Mechanics
Etcheverry Hall
University of California
Berkeley, California 94720

Professor W. Nachbar
University of California
Department of Aerospace & Mech. Engrg
La Jolla, California 92037

Professor J. Baltrukonis
Mechanics Division
The Catholic Univ. of America
Washington, D.C. 20017

Professor A.J. Durelli
Mechanics Division
The Catholic Univ. of America
Washington, D.C. 20017

Professor H.H. Bleich
Department of Civil Engrg.
Columbia University
Amsterdam & 120th Street
New York, New York 10027

Professor R.D. Mindlin
Department of Civil Engrg.
Columbia University
S.W. Mudd Building
New York, New York 10027

Professor A.M. Freudenthal
George Washington University
School of Engrg. & Applied Science
Washington, D.C. 20006

Professor B.A. Boley
Department of Theor. & Appl. Mech.
Cornell University
Ithaca, New York 14850

Professor P.G. Hodge
Department of Mechanics
Illinois Institute of Technology
Chicago, Illinois 60616

Dr. D.C. Drucker
Dean of Engineering
University of Illinois
Urbana, Illinois 61801

Professor H.M. Newman
Dept. of Civil Engineering
University of Illinois
Urbana, Illinois 61801

Professor James Mar
Massachusetts Inst. of Tech.
Rm. 33-318
Dept. of Aerospace & Astro.
77 Massachusetts Ave.
Cambridge, Massachusetts 02139

Library (Code 0384)
U.S. Naval Postgraduate School
Monterey, California 93940

Dr. Francis Cozzarelli
Div. of Interdisciplinary
Studies and Research
School of Engineering
State Univ. of N.Y.
Buffalo, New York 14214

Professor R.A. Douglas
Dept. of Engrg. Mechs.
North Carolina State University
Raleigh, North Carolina 27607

Dr. George Herrmann
Stanford University
Department of Applied Mechanics
Stanford, California 94305

Professor J.D. Achenbach
Technological Institute
Northwestern University
Evanston, Illinois 60201

Director, Ordnance Research Lab.
Pennsylvania State University
P.O. Box 30
State College, Pennsylvania 16801

Professor J. Kempner
Dept. of Aero. Engrg. & Applied Mech.
Polytechnic Institute of Brooklyn
333 Jay Street
Brooklyn, New York 11201

Professor J. Klosner
Polytechnic Institute of Brooklyn
333 Jay Street
Brooklyn, New York 11201

Professor A.C. Eringen
Dept. of Aerospace & Mech. Sciences
Princeton University
Princeton, New Jersey 08540

Dr. S.L. Koh
School of Aero., Astro. & Engr. Sc.
Purdue University
Lafayette, Indiana 47907

Professor R.A. Schapery
Civil Engineering Department
Texas A & M University
College Station, Texas 77840

Professor E.H. Lee
Div. of Engrg. Mechanics
Stanford University
Stanford, California 94305

Dr. Nicholas J. Hoff
Dept. of Aero. & Astro.
Stanford University
Stanford, California 94305

Professor Max Amiker
Dept. of Aero. & Astro.
Stanford University
Stanford, California 94305

Professor Chi-Chang Chao
Div. of Engr. Mechanics
Stanford University
Stanford, California 94305

Professor H.W. Liu
Dept. of Chemical Engr. & Metal.
Syracuse University
Syracuse, New York 13210

Professor S. Bodner
Technion R & D Foundation
Haifa, Israel

Dr. S. Dhawan, Director
Indian Institute of Science
Bangalore, India

Professor Tsuyoshi Hayashi
Department of Aeronautics
Faculty of Engineering
University of Tokyo
BUNKYO-KU
Tokyo, Japan

Professor J.E. Fitzgerald, Ch.
Dept. of Civil Engineering
University of Utah
Salt Lake City, Utah 84112

Professor R.J.H. Bollard
Chairman, Aeronautical Engr. Dept.
207 Guggenheim Hall
University of Washington
Seattle, Washington 98105

Professor Albert S. Kobayashi
Dept. of Mechanical Engr.
University of Washington
Seattle, Washington 98105

Professor G.R. Irwin
Dept. of Mech. Engrg.
Lehigh University
Bethlehem, Pennsylvania 18015

Dr. Daniel Frederick
Dept. of Engr. Mechs.
Virginia Polytechnic Inst.
Blacksburg, Virginia 24061

Professor Lambert Tall
Lehigh University
Department of Civil Engrg.
Bethlehem, Pennsylvania 18015

Professor M.P. Wmuk
South Dakota State University
Department of Mechanical Engineering
Brookings, South Dakota 57006

Professor Norman Jones
Massachusetts Institute of Technology
Dept. of Naval Architecture & Marine Engrg.
Cambridge, Massachusetts 02139

Professor Pedro V. Marcal
Brown University
Division of Engineering
Providence, Rhode Island 02912

Professor Werner Goldsmith
Department of Mechanical Engineering
Division of Applied Mechanics
University of California
Berkeley, California 94720

Professor R.B. Testa
Dept. of Civil Engineering
Columbia University
S.W. Mudd Bldg.
New York, New York 10027

Dr. Y. Weitsman
Dept. of Engrg. Sciences
Tel-Aviv University
Ramat-Aviv
Tel-Aviv, Israel

Professor W.D. Pilkey
Dept. of Aerospace Engrg.
University of Virginia
Charlottesville, Virginia 22903

Professor W. Prager
Division of Engineering
Brown University
Providence, Rhode Island 02912

Industry and Research Institutes

Mr. Carl E. Hartbower
Dept. 4620, Bldg. 2019 A2
Aerojet-General Corporation
P.O. Box 1947
Sacramento, California 95809

Library Services Department
Report Section, Bldg. 14-14
Argonne National Laboratory
9400 S. Cass Avenue
Argonne, Illinois 60440

Dr. F.R. Schwarzl
Central Laboratory T.N.O.
Schoemakerstraat 97
Delft, The Netherlands

Dr. Wendt
Valley Forge Space Technology Cen.
General Electric Co.
Valley Forge, Pennsylvania 10481

Library Newport News Shipbuilding
& Dry Dock Company
Newport News, Virginia 23607

Director
Ship Research Institute
Ministry of Transportation
700, SHINKAWA
Mitaka
Tokyo, Japan

Dr. H.N. Abramson
Southwest Research Institute
8500 Culebra Road
San Antonio, Texas 78206

Dr. R.C. DeHart
Southwest Research Institute
8500 Culebra Road
San Antonio, Texas 78206

Mr. Roger Weiss
High Temp. Structures. & Materials
Applied Physics Lab.
8621 Georgia Ave.
Silver Spring, Maryland 20910

Mr. E.C. Francis, Head
Mech. Props. Eval.
United Technology Center
Sunnyvale, California 94088

Mr. C.N. Robinson
Atlantic Research Corp.
Shirley Highway at Edsall Road
Alexandria, Virginia 22234

Mr. P.C. Durup
Aeromechanics Dept., 74-43
Lockheed-California Co.
Burbank, California 91503

Mr. D. Wilson
Litton Systems, Inc.
AATD, Dept. 400
El Segundo
9920 W. Jefferson Blvd.
Culver City, California 90230

Dr. Kevin J. Forsberg, Head
Solid Mechanics
Orgn 52-20, Bldg. 205
Lockheed Palo Alto Research Lab.
Palo Alto, California 94302

Dr. E.M.Q. Roren
Head, Research Department
Det Norske Veritas
Post Box 6060
Oslo, Norway

Dr. Andrew F. Conn
Hydronautics, Incorporated
Pindell School Road, Howard County
Laurel, Maryland 20810