

AD-A043 342

STANFORD UNIV CALIF DEPT OF AERONAUTICS AND ASTRONAUTICS F/G 20/4
SINGULAR PERTURBATION METHODS APPLIED TO THE DYNAMICS AND RHEOL--ETC(U)
JUL 77 I CHANG

DAAG29-73-C-0034

UNCLASSIFIED


ARO-11032.3-E

NL

| OF |
ADA
043342



END
DATE
FILMED
9-77
DDC



NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

9. ARO- 11032.3-E

13



**Department of AERONAUTICS and ASTRONAUTICS
STANFORD UNIVERSITY**

AD A 043342

Singular Perturbation Methods Applied to the
Dynamics and Rheology of Suspensions
Final Report

By I-Dee Chang

July 10, 1977

U.S. Army Research Office

DAAG29 73 C 0034


(for the period 1 July 1973-30 April 1977)

Department of Aeronautics and Astronautics
Stanford University, Stanford, CA 94305

AD NO. _____
DDC FILE COPY.

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited



1. Statement of the problem studied.

Many fundamental problems pertaining to the dynamics of suspended particles remain unsolved. Of particular interest to our present investigation are problems belonging to the following four categories:

- a. Where the non-linear effect of the fluid is a decisive factor in determining the motion of the suspended particles. An example of such problem is the motion of a particle in a slow shear flow. It was found that a sphere, when moving relative to a fluid under uniform simple shear, experiences a lift force transverse to the direction of fluid and particle motion. If the fluid inertia effect is neglected, as in the case of Stokes approximation, the lift force will disappear and only streamwise drag remains. Thus the lift force, which may cause the sphere to drift across streamlines, is a direct consequence of inertia effects. Harper and Chang have shown both analytically and experimentally for a dumb-bell shaped particle that the resulting lift tends to orient the particle to the orientation of maximum dissipation.
- b. Where the inertia of the suspended particle itself is of importance. An example of this is the rotary Brownian motion of particles in a suspension.
- c. Where the hydrodynamic couplings between the suspended particles and between the particles and the container wall, are important.
- d. Where the collective motion of an assembly of particles is of interest. During the present program, methods have been developed to study these types of

ACCESSION TO:	<input checked="checked" type="checkbox"/> X	<input type="checkbox"/>	<input type="checkbox"/>
	NTS	With Section	But Section
DDP			
SPANDED			
JUSTIFICATION			
BY			
DISTRIBUTION/AVAILABILITY CODES			
DEL. DATE, TIME, AND METHOD			

A

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

problems analytically. In particular, we have concentrated our effort on three representative problems. In these three problems, small but significant physical perturbation effects characteristics in the four types of problem list in § 1 are present. The mathematical techniques required to deal with them are therefore of general interest:

- (i) The motion of a small spherical particle suspended in an oscillating shear flow.
- (ii) The motion of ellipsoidal particles in a viscous shear flow, and
- (iii) unsteady motion of a small particle in a compressible viscous flow.

The above investigations resulted in large amounts of analytical results and numerical data. Some part of these investigations have been published as technical reports and released. Much work is still in progress and will be published as soon as the work is complete.

2. Summary of most important results.

- a. The motion of a spherical particle suspended in an oscillating shear flow. The force on a small sphere immersed in an oscillating shear flow is investigated. The unsteady Navier-Stokes equation is solved by the method of matched asymptotic expansions. The inertia effects of the fluid produce a lift force on the sphere in the direction of increasing flow velocity. For a neutrally buoyant sphere freely suspended in the flow, this lift force will cause the sphere to drift sideways across the streamline. The motion of the sphere in a sinusoidal oscillating shear flow is examined in detail. The small streamwise slip velocity

between the sphere and the fluid is calculated first. From this, the lift force on the sphere is determined. It is found that the lift force oscillates at twice the frequency of the sheer flow field and has a d-c component which tend to make the sphere drifting slowly across the stream-line toward the flow region of faster velocity. The drift velocity of a sphere of 200μ in a shear flow oscillating at 1 Hz is estimated. The magnitude of this velocity is found to be quite reasonable.

- b. The motion of ellipsoidal particles in a viscous shear flow. This investigation is related to the intrinsic viscosity of a dilute suspension of particles in a homogeneous fluid. Jeffery in 1922 calculated the intrinsic viscosity for a dilute suspension of ellipsoidal particles in a linear shear flow using Stokes equations. However, his solution was indeterminate in the sense that the motion, which is periodic, depends upon the initial conditions of release of the particle. We attempt to carry on Jeffrey's work by including the inertia effect of the fluid and to show that the orbit of rotation changes adiabatically over a long period of time.

The slow gyration of an ellipsoid in a shear flow is studies by a singular perturbation method. The small parameter used in the analysis is the shear Reynolds number R_K defined by $R_K = \frac{\rho \kappa l}{\mu}$ where κ is the shear rate and l the characteristic dimension of the ellipsoid. It is found that as $R_K \rightarrow 0$, the Jeffrey solution gives the first order uniformly valid solution of the problem. In the next order

approximation, taking into account the fluid inertia effect, a new set of equations is needed.

In this outer region, which is of distance $O(l/R_k)$ from the body, the flow field is to first-order equivalent to that induced by a set of gyrating dipoles in a shear flow. To match terms it is necessary to introduce a term $O(R_k^{3/2})$ in the inner expansion. This is of smaller order than the second-order inner solution in the expansion which arises from the nonlinear terms through iteration and having the order $O(R_k)$. The order $O(R_k^{3/2})$ inner solution corresponds physically to that of an ellipsoid rotating freely in an unsteady linear flow field composed of arbitrary strain deformation and rotational motion.

A technical report (SUDAAR No. 492) has been released.

- c. Unsteady motion of a small particle in a compressible viscous flow. The motion of a small particle suspended in an incompressible viscous shear flow has been investigated under steady and unsteady conditions. Recent interest in acoustics calls for extension of this effort to the case when the flow is compressible. We consider therefore the problem of slow vibration of a class of axially symmetric bodies moving parallel to the body axis with the velocity $U = U_0 e^{i\omega t}$ in a quiescent compressible viscous fluid. It is expected that results obtained from the analysis will be useful in many realistic physical situations where in the flow fluctuations are coupled with the motion of the small particle.

There are three non-dimensional parameters in the problem. These are the flow Reynolds number

$$Re = \frac{\rho_0 U_0 a}{\mu_0}, \text{ the unsteady Reynolds number } \alpha^2 = \frac{\rho_0 \omega a^2}{\mu_0}$$

and the Mach number $M = \frac{U_o}{C_o}$, where a is the characteristic dimension of the particle, and quantities with subscript "o" refer to undisturbed flow conditions. Since the particle is small, we assume that Re and α^2 are small. It follows that the Mach number M is also small. We therefore solve the problem by finding the asymptotic solution of the exact compressible, viscous heat conduction Navier-Stokes equations, for small values of Re , α and M . The problem differs from the acoustic-type problem in that viscous effect is important and must be considered. It is found that the solution depends on four characteristic length scales. These are the geometric length scale a , the viscous length scale $\frac{\mu_o}{\rho_o U_o}$, "Stokes" length scale $\sqrt{\frac{\mu_o}{\rho_o \omega}}$

and the wave length λ . The condition $M \ll 1$ in particular implies $\frac{a}{\lambda} \ll 1$. In the inner region $r \ll \lambda$, the flow is nearly incompressible. This greatly reduces the complexity of the problem in the inner region. In the outer region, $r = O(\lambda)$, the flow must be treated as compressible. However, simplification is derived from the boundary condition and again the problem becomes tractable. In fact, to the order $O(\alpha)$, the flow field is characterized by the "Stokes drag" experienced by the particle rather by the detailed geometry of the particle. The solution of the far field is obtained by means of Fourier transform. It is found that sound waves similar to that generated by dipole dominate the far field. The acoustic intensity pattern shows maxima both in the forward and rearward direction of the motion. The details are however modified by the effects of viscosity and heat conduction.

3. Publications

- a. The motion of a small spherical particle suspended in an oscillating shear flow. By I-Dee Chang, SUDAAR No. 487 August, 1974.
- b. The motion of ellipsoidal particles in a viscous shear flow. By I-Dee Chang SUDAAR No. 492 April 1975
- c. Unsteady motion of a small particle in a compressible viscous flow. By I-Dee Chang (To appear).

4. Participating scientific personnel.

I-Dee Chang, professor, Department of Aeronautics and Astronautics, Stanford University, CA 94305.

Jacques Beser, Research assistant, Department of Aeronautics and Astronautics (Ph.D. Candidate).

Christoph Holliger, Research assistant, Department of Aeronautics and Astronautics (Earned M.S. degree June 1975)

Raymond C. C. Luh, Research assistant, Department of Aeronautics and Astronautics (Earned M.S. degree on June 1977)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER 124807	2. GOVT ACCESSION NO. 11032.3-E	3. RECIPIENT'S CATALOG NUMBER 1	
4. TITLE (and Subtitle) SINGULAR PERTURBATION METHODS APPLIED TO THE DYNAMICS AND RHEOLOGY OF SUSPENSIONS		5. TYPE OF REPORT & PERIOD COVERED FINAL REPORT 1 July 1973-30 April 1977	
		6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) 10 I-Dee/Chang		8. CONTRACT OR GRANT NUMBER(s) 15 DAAG29-73-C-0034 new	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Aeronautics and Astronautics Stanford University, Stanford, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 11032-E	
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE July 10, 1977	
		13. NUMBER OF PAGES 121 pages	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION/DCN GRADING SCHEDULE NA	
16. DISTRIBUTION STATEMENT (of this report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA			
18. SUPPLEMENTARY NOTES The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Suspensions, viscous flow, non-linear effects, low Reynolds number			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Non-linear hydrodynamic interaction on particles suspended in viscous flows was investigated analytically by the method of matched expansions. In par- ticular, three problems of fundamental importance were analyzed. The first problem involves the motion of a particle in an oscillating viscous shear flow. It was found that because of the inertia of the fluid, a lift force is produced on the particle. This force has a d-c component which tends to cause a sidewise drift of the particle. The second problem involves the extension of Jeffery's classical work on the motion of ellipsoidal particle (continued)			

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

continued from front:

→ immersed in a viscous shear flow. It was found that the first order outer solution corresponds to gyrating dipoles in a shear flow. The effect of the shear produces a higher order effect than the one caused by the second order non-linear effect in the near field. The third problem dealt with the motion of a spherical particle in a viscous compressible flow. The flow field was calculated in terms of a near field and a far field. The flow pattern is rather complex because of the combined effects of heat conduction and viscosity. ↗