ADA043536

からの

2

REPORT MEDDH-288(R1)

la for

AD

ANNUAL RESEARCH PROGRESS REPORT

U.S. ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Approved for public release. Distribution unlimited.





Prepared for: US ARMY MEDICAL RESEARCH & DEVELOPMENT COMMAND WASHINGTON, D.C. 20314 DESTROY THIS REPORT WHEN IT IS NO LONGER NEEDED, DO NOT RETURN IT TO THE ORIGINATOR.

APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED

FINDINGS IN THIS REPORT ARE NOT TO BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, UNLESS SO DESIGNATED BY OTHER AUTHORIZED DOCUMENTS.

IN CONDUCTING THE RESEARCH DESCRIBED IN THIS REPORT THE INVESTIGATORS ADHERED TO THE "GUIDE FOR LABORATORY ANIMAL FACILITIES AND CARE", AS PROMULGATED BY THE COMMITTEE ON THE GUIDE FOR LABORATORY ANIMAL FACILITIES AND CARE OF THE INSTITUTE OF LABORATORY ANIMAL RESOURCES, NATIONAL ACADEMY OF SCIENCES NATIONAL RESEARCH COUNCIL.



REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2. GOVT ACCESSION NO	. 3. RECIPIENT'S CATALOG NUMBER
(a)	
4. TITLE (and Subtitio)	5. TYPE OF REPORT & PERIOD COVER
Annual Barrance Denant) (Rept for	1 Julie 75- 30 Sept 76.
Annual Research Progress Report .	6. PERFORMING ORG. REPORT NUMBE
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	
7. AUTHOR(.)	8. CONTRACT OR GRANT NUMBER(*)
(2)7-1P.	(12) \$1
	2201
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT. PROJECT, TA
US Army Institute of Surgical Research (16)	3A161102B71R-01
Fort Sali Houston, Texas 70234	SAIGIPTA91C (cont'd
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
US Army Medical Research & Development Command	30 September 76 1
washington, DC 20317. $(1)$	NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office)	15. SECURITY CLASS, (of this report)
	15. DECLASSIFICATION/DOWNGRADIN
	SCHEDULE
Approved for public release. Distribution unlimit	ted.
<ul> <li>Approved for public release. Distribution unlimitation of the sector of t</li></ul>	oon Report)
16. DISTRIBUTION STATEMENT (of Inte Report) Approved for public release. Distribution unlimit 17. D.STRIBUTION STATEMENT (of the ebetract enterned in Block 20, if different fo	on Report) DDC
16. DISTRIBUTION STATEMENT (of Inte Report) Approved for public release. Distribution unlimi 17. D.STRIBUTION STATEMENT (of the ebetract enterned in Block 20, if different fo	DDC
<ul> <li>Approved for public release. Distribution unlimited of the electron of the electr</li></ul>	om Report) DDC DCOCIO
<ul> <li>Approved for public release. Distribution unlimited in the second second</li></ul>	
<ul> <li>Approved for public release. Distribution unlimited of the electron of the electr</li></ul>	
<ul> <li>Approved for public release. Distribution unlimited in the second second</li></ul>	om Report) DDC DC AUG 31 197
<ul> <li>ISTRIBUTION STATEMENT (of this Report)</li> <li>Approved for public release. Distribution unlimited on the second of th</li></ul>	DDC DDC DC AUG 31 197 B
<ul> <li>IS. DISTRIBUTION STATEMENT (of this Report)</li> <li>Approved for public release. Distribution unlimited</li> <li>17. D.STRIBUTION STATEMENT (of the abstract enteroid in Block 20, if different to</li> <li>18. SUPPLEMENTARY NOTES</li> <li>18. KEY WORDS (Continue on reverse eide if necessary and identify by block numbers)</li> </ul>	non Report) DDC DC AUG 31 197 AUG 31 197 B
<ul> <li>IS. SUPPLEMENTARY NOTES</li> <li>IS. KEY WORDS (Continue on reverse eide if necessary and identify by block number</li> </ul>	red. DDC DC AUG 31 197 B
<ul> <li>Is distribution statement (of this Report)</li> <li>Approved for public release. Distribution unlimited on the second of the second o</li></ul>	non Report) DDC DCOCIDI AUG 31 197 SUGGIGIT " B
<ul> <li>ISTRIBUTION STATEMENT (of this Report)</li> <li>Approved for public release. Distribution unlimited in the provided of the source of</li></ul>	red. DDC DC AUG 31 197 AUG 31 197 B
<ul> <li>IS. DISTRIBUTION STATEMENT (of this Report)</li> <li>Approved for public release. Distribution unlimited in the statement of the ebstract entered in Block 20, if different for the statement of the ebstract entered in Block 20, if different for supplementary notes</li> <li>IS. KEY WORDS (Continue on reverse eide if necessary and identify by block number of the supplementary and the supplementary and identify by block number of the sup</li></ul>	activities of the US Army
<ul> <li>16. DISTRIBUTION STATEMENT (of Inte Report)</li> <li>Approved for public release. Distribution unlimited in the provided of the electron of the elect</li></ul>	activities of the US Army 1976. These activities
<ul> <li>16. DISTRIBUTION STATEMENT (of Inte Report)</li> <li>Approved for public release. Distribution unlimited in the statement of the abstract entered in Block 20, if different in the statement of the abstract entered in Block 20, if different in the supplementary notes</li> <li>18. SUPPLEMENTARY NOTES</li> <li>19. KEY WORDS (Continue on reverse elde if necessary and identify by block number in the supplementary include patient care, clinical investigation and is a supplementary of the supplementary of the supplementary in the supplementary of the supplementary of the supplementary investigation and its supplementary of the sup</li></ul>	activities of the US Army 1976. These activities aboratory research in the and (3) general traima
<ul> <li>IS. DISTRIBUTION STATEMENT (of this Report)</li> <li>Approved for public release. Distribution unlimited in the provided of the electron of the elect</li></ul>	activities of the US Army 1976. These activities aboratory research in the and (3) general trauma. hent of burned patients and
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report)</li> <li>Approved for public release. Distribution unlimited in the second of the second o</li></ul>	activities of the US Army 1976. These activities aboratory research in the and (3) general trauma. went of burned patients and burned wound infection.

-

UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered) 1

- ---

3816111028805-00 3A1611028719-02 3A162110A821-00

-

. .

•



DEPARTMENT OF THE ARMY US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

SGRD-US

SUBJECT: Annual Research Report FY 1976

TO: See Distribution

Annual report(s) of the US Army Institute of Surgical Research for FY 1976 are forwarded under provisions of the OTSG Regulation 70-31, dated 2 April 1969.

BASIL A. PRUITT, JR, MD Colonel, MC Commander & Director

ACCESSIC	1 1		٦
NTIS	1	te 🔿 Hoa 🐧	Л
DDC	E,	3 . S. [	1
LNT	ñ	Ē	
4 ° 1 c			ļ
ţεγ			1
ťΝ	~		
-		ال ياريا	
Λ.		ł	
H	ļ	I	
1		1	



### FOFEWORD

The effectiveness of categorical health care in terms of clinical results and research productivity is we'i illustrated by the material within this volume produced by the concerted effort of enthusiastic, hard-working, and generally optimistic laboratory and clinical staff. Not only has average duration of hospital stay for our burn patients decreased over the past five years but improved functional results have also been achieved in many patients with severe burns, and clinical and laboratory investigations have led to improved diagnosis of inhalation injury, effective prophylaxis against Curling's ulcer, identification of basic defects in host resistance, and improved techniques of metabolic and nutritional support. This clinical-laboratory symbiosis, which has broad medical applicability, is dependent upon several factors which are in them-selves mutually reinforcing.

The Army Medical Department has, by concentrating burn care at this Institute, ensured an adequate and dependable density of patients. The Medical Research and Development Command, by farsighted personnel management, has provided for stability of senior staff, assignment of selected clinical staff and the implacement of a multicisciplinary research team. Planned construction by the Medical Department has also established sophisticated research laboratory facilities. The staff, in turn, has identified significant problems for study, organized interdisciplinary research teams, established dedicated clinical study areas, promoted active involvement of allied scientists and, as noted above, applied research techniques in an innovative fashion to the successful resolution of clinical problems. This symbiosis, which must exist for effective, clinically relevant research, benefits the individual staff member by facilitating his scientific and clinical productivity and benefits the Army by the merit of the end product and the productivity of the Institute. Most importantly, the burned or otherwise injured soldier remains the focal point of all the care and research activities of the Institute and benefits from these combined individual and organizational efforts.

Basil a. Preitt, Jr., MD, FACS

BASIL A. PRUITT, JR., MD, FAC Colonel, MC Commander & Director

### TABLE OF CONTENTS

# PAGE NO.

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

Clinical Operation, Center for Treatment of Burned Soldiers H.D.Peterson, J.Dorethy, J.Z. Kaplan, T.S.Lescher, W.S.McDougal, R.H.Merrill, A.Rosenthal, J.K.Sims, D.K.Teegarden, G.W.Welch, D.W.Wilmore, M.G.Michael, H.M. Diaz, D.N. Herndon, N.F.Jesse, J.P.Peterson, B.A.Pruitt, Jr.	1
Clinical Operation, Surgical Study Branch for Treatment of Injured Soldiers D.W. Wilmore, L.H. Aulick,J.P.Peterson,D.H. Dahlquist	32
Anesthesiology G.W.Welch, J.K.Sims, A.K.Morris	37
Clinical Operation, Metabolic Branch, Renal Section, for Treatment of Soldiers with Renal Failure W.D.Myers, R.H. Merrill	47
Detection of Endotoxin in Burned Soldiers with Sepsis R.B.Lindberg,V.C.English,A.D.Mason,Jr., B.A.Pruitt,Jr.	52
Antibiotic Sensitivity of Current Military Burn Patient Flora R.B.Lindberg, A.A.Contreras, D.T. Zamora	62
Emergence of Methicillin-Resistant <u>Staphylococcus</u> aureus Type 84 and 84,85 in Burned Military Personnel R.B.Lindberg,R.L.Latta,B.A.Pruitt,Jr.,A.D.Mason,Jr.	81
Sensitivity to Sultany on of <u>Pseudomonas aeruginosa</u> Recovered from Burned Soldiers R.B.Lindberg, C.C.English, B.A.Pruitt, Jr., A.D. Mason, Jr.	97
Pathogenesis of Burn V ound Infection: Bacterial Flora of Wounds of Military Personnel Receiving Treatment with Sulfamylon or Silver-Sulfadiazine R.B.Lindberg, A.A.Contreras, R.L.Latta, D.T.Zamora	110
Bacteriophage Types of <u>Pseudomonas aeruginesa</u> Found in Burned Soldiers R.L.Latta, R.B.Lindberg, A.D.Mason, Jr., B.A.Pruitt, Jr.	134

i

# PAGE NO.

Bacteriophage Types of <u>Serratia marcescens</u> from Burn Wounds of Military Personnel R.B.Lindberg,V.C.English,B.A.Pruitt,Jr.,A.D. Mason,Jr.	150
Five Per Cent Aqueous Sulfamylon Soaks Used in Topical Treatment of Burned Soldiers H.D.Peterson,B.A.Pruitt,Jr.	153
Development of Prophylactic Topical Therapy for Use on Burn Wounds of Military Patients: The Search for New Formulations R.B.Lindberg,V.C.English,R.L.Latta,G.T.Daye, B.A.Pruitt,Jr.	157
The Role of Fungi in Burn Wound Infection:Observations on Biopsy and Autopsy Tissues from Seriously Burned Soldiers R.B.Lindberg	169
Non-Fermentative Gram-Negative Bacilli in Burned Soldiers: New Potential Oppportunistic Pathogens V.C.English,R.B.Lindberg	177
Studies of Disturbance of Protein Turnover in Burned Troops- Use of an Animal Model W.L.Brown,E.G.Bowler,A.D.Mason,Jr.	189
Evaluation of Gastrointestinal Absorption and Nutritional Efficacy of Standard High Protein Diets in Burned Soldiers D.W.Wilmore,J.P.Peterson	194
A Therapeutic Trial of Antacid in Prevention of the Clinical Complications Associated with Gastric Mucosal Disease in Burned Soldiers D.K.Teegarden,J.C.McAlhany,Jr, B.A.Pruitt, Jr.	197
Pulmonary Complications in Thermal Injury V.Lam,J.K.Sims,G.W.Welch,P.A.Petroff,Jr.	201
Evaluation of Dopamine (3,4-Dehydroxyphenylethylamine) for Treatment of Septic Shock in Burned Troops G.W.Welcn, C.Drueck,III,R.W.J.Baird,D.W.Wilmore	206
Prolongation of Skin Allograft Survival by Immunosuppressive Therapy (Upjohn ATC) in Soldiers with Massive Thermal Injury H.D.Peteron,D.W.Wilmore,B.A.Pruitt,Jr.	215

The Effect of Dextrose and Amino Acid Solutions on Nitrogen Excretion and Nitrogen Balance Following Thermal Injury in Burned Soldiers W.S.McDougal, D.W.Wilmore	218
Excision of Eschar in Burned Soldiers H.D.Peterson,B.A.Pruitt,Jr.	238
Evaluation of Synthetic Sheeting as Operating Room Drape Material for Use in a Military Burn Unit B.A.Pruitt,Jr., R.B.Lindberg, A.D.Mason,Jr.	244
PROJECT, NO. 3A16101A91C-00, IN-HOUSE LABORATORY INDEPE RESEARCH	NDENT
Studies of the Effect of Variations of Temperature and Humidity on Energy Demands of the Burned Soldier in a Controlled Metabolic Room D.W.Wilmore,A.D.Mason,Jr, L.H.Aulick,B.A.Pruitt,Jr.	251
Development and Analysis of An Animal Model for the Post Thermal Injury Hypermetabolic Response Found in Burned Soldiers D.N.Herndon, D.W.Wilmore, A.D.Mason, Jr., R.V. Condon, Jr., E.W.Hander, B.A.Pruitt, Jr.	262
Evaluation of Calcium Metabolism in Burned Troops R.H.Merrill,W.D.Myers,H.R.Jacobson	290
Laboratory Investigation of the Mechanisms of Acquired Leukocyte Dysfunction Following Thermal Injury A.T.McManus,Jr, T.J.Lescher,A.D.Mason,Jr., B.A.Pruitt,Jr.	293
Echocardiographic Evaluation of Left Ventricular Performance in Severely Burned Military Population	297
PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT MILITARY MEDICINE	OF
Renal Function in the Burned Soldier, I.Histology W.D.Myers, T.Rizzo, Jr., R.H.Merrill, J.McPhaul	300
Localization of Infection within the Urinary Tract W.D.Myers,H.A.Gersh,T.Rizzo,Jr.,M.Forland,V.Thomas	307

	PAGE NO.
To Determine * Role of the Prostaglandins (PG) in the Response to Volume Expansion in the Dog - A Model of Changes in Injured Troops D.B.Olin,R.H.Merrill	312
To Determine Whether the Prostaglandins are Important in Development of Acute Renal Failure - A Rabbit Model of Renal Failure in Soldiers D.B.Olin,R.H.Merrill	324
The Effects of Calcium on the Renin-Angiotensin System - Use of an Animal Model of Hypertension R.H.Merrill, W.D.Myers, T.F.Lescher	327
The Use of Lyophilized Venous Homografts for Peripheral AV Fistula in Burned Soldiers R.H.Merrill,H.D.Peterson,W.D.Myers,T.J.Lescher	330
Determination of Arterial and Tissue Oxygen Tension in Normal Main to Develop a Technic to Measure Tissue Oxygen Tension in Burned Soldiers C.L.Slade,G.W.Welch,P.A.Petroff	333
Serum Pepsinogen Levels in the Thermally Injured Soldier: A Possible Predictor of Gastroduodenal Disease D.K.Teegarden,I.M. Samloff	337
Alterations in the Bacterial Flora of the Stomach and Upper Small Intéstine Following Thermal Injury in a Military Population D.K.Teegarden,W.S.McDougal,T.M.Sasaki,B.L. Davison,R.B.Lindberg	341
Evaluation of Altered Metabolism of Phenylalanine and Tyrosine in Burned Patients D.N.Herndon,D.W.Wilmore,A.D.Mason,Jr.,J. DuBois, B.A.Pruitt, Jr.	346
Evaluation, Donor Site Healing and Mesh Graft Spread with Hyperbaric Oxygen H.D.Peterson, B.A.Pruitt, Jr.	371
PROJECT NO. 3A162110A821-00, COMBAT SURGERY	
Low Flow Vein Anastomosis in Dogs Anticoagulated with AncrodA Model for Blood Vessel Repair in Injured Troops C.L.Slade, W.A. Andes	376

# PAGE NO.

A Prospective Comparison Study of Sulfamylon and Silver Sulfadiazine in the Treatment of Burned Troops H.D.Peterson, A.D. Mason, Jr., B.A.Pruitt, Jr.	379
Evaluation of the Effect of Fresh Frozen Plasma on Leukocyte Chemotaxis in Burned Soldiers J.W.Taylor, J.M.Long,III,A.D.Mason,Jr.,B.A. Pruitt,Jr.	383
Evaluation of Gastric Physiologic Disturbances Associated with Thermal Injury in a Military Population D.K.Teegarden, B.A. Pruitt, Jr.	386
The Zinc Requirements of the Burned Rat and the Influence of Zinc on LDH Activity, Growth Rate and Wound Healing: A Model of Burned Soldiers D.J.Johnson, Y.Villarrezl, H.L.Walker, A.D.Mason, Jr.	392
The Relationship Between Limb Blood Flow and Metabolism Following Thermal Injury in Soldiers L.H.Aulick,D.W.Wilmore,A.D.Mason,Jr, B.A. Pruitt,Jr.	398
Intrarenal Blood Distribution: A Noninvasive Technique for Determining the Intrarenal Distribution of Blood Flow W.S.McDougal,W.D.Myers,R.J.McAuley,R.Lull	420
Glucagon and heat Production in Normal Man and Septic Burn Patients in a Military Population L.H.Aulick,D.W. Wilmore,A.D.Mason,Jr.	427
Aerosolized Gentamicin in the Prophylactic Treatment of Inhalation Injury B.A.Levine,H.D.Peterson,R.B.Lindberg,B.A.Pruitt,Jr.	441
Publications - 1 July 1975-30 September 1976	445

Simony and a contraction										
RESEARCH	AND TECHNOLOG	WORK UNIT S	UMMARY	DA	OP, 63	80 B	76 10 (	MARY [®] 1	BPORT C	WINDL STREEL
3. DATE PREV SUMPRY 75 07 01	6 KIND OF SUMMARY D. CHANGE	S. SUNMARY SCTY [®] , U	A. WORK SECURITY	P RECA	ioing ^e A	104. 246	PN INSTA'N VT.	CONTRACTOR	TA-	. LEVEL OF SUE
10. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TARK					10	
A PRIMARY	61102A	351611102	BS05		00		A20	WORK UNIT	UMBER	
S. CONTRIBUTING	61102A	3A161102B	71R		00 01		005			
C. CONTRIBUTING	0120111	0.11011022	-	<u>ل</u> ــــــــــــــــــــــــــــــــــــ	<u>.</u>		415			
11. TITLE (Procode with	Security Classification Code								×>	2
(U) Clinica	al Operation,	Center for	Treatment	of B	urned	Solo	liers (4	14)		
12 SCIENTIFIC AND TE	CHHOLOGICAL AREAS									
003500 Clir	nical Medicine	2								
13. START DATE		14. ESTINATED COM	PLETION DATE	IN FUNE	NNG AGEN	CY		16. PERFORMA	CE METI	100
50 07		Cont		<u>na</u>			<u> </u>	C. In-	Hous	e
17. CONTHACT/GRANT	Not Applica	able		10. RES	URCES ES	TIMATE	& PROFESSI	OWAL MAN YES	L FUN	D6 (Jn thr seands)
A DATES/EFFECTIV 1		EXPIRATION:			76		51.6		6	25
h HUNDER.				FISCAL	T T		<u>  11.1</u>		1	54
G TYPE:		4 AMOUNT:		YEAR						
& KIND OF AWARD		f. CUM. AMT.			77		54.6	) 	7	76
IN. RESPONSIBLE DOD		Ĺ		-200. PERI -		RGANIZA	TION			
HAME!" US Army	Institute of	Surgical H	lesearch	H-ME:U	S Arm	y Ins	stitute	of Surgi	cal	Research
ADDRESS!* FORT	low Noveton D			ADDRES		inica	al Divis	sion		
FOLCE	ball houston,	lexas /02.			FOF	t Sai	n Housto	on, Texas	3 78	234
				PRINCIP	AL INVEST	GATOR	(Funish SEAN i	l U.S. Annumic p	u i i tu tianj	,
RESPONSIBLE INDIVIDU	AL			HANE:	Hug	h D.	Peterso	on, COL,	MC	
NAME Basil A.	Pruitt, Jr,	COL, MC		TELEP	HONE:	512	-221-330	)1		
TELEPHONE: 512	2-221-2/20			SOCIAL	SECURIT	ACCOU	NT NUMBER:			
21. GENERAL USE				ASSOCIA	TE HAVEST	GATOR	l -			
FOREIGN INT	TELLIGENCE NOT	CONSIDERE	D	NAME						
22. KEYBORDA (Presede	RACH WIR Somethy Classif	tellan Cada)		HAME:						DA
(U) Heteroo	raft; (U) Res	suscitation	; (U) Homo	graft	; (U)	Air	evacuat	ion; {U	The	rmal
1njury; (U)	Inhalation 3	Lnjury; (U)	Topical t	herap	<u>y; (U</u>	<u>) Au</u>	<u>tograft</u>			,
23. (U) Th	ne Clinical Di	vision of	the US Arm	y Ins	titut	e of	Surgica	al Reseau	ch c	, ontinues
its role as	a major spec	cialized cl	inical tre	atmen	t cen	ter :	for the	mally in	niure	d military
personnel.	Its main ob	jectives ar	e the inve	stiga	tion	and r	nodifica	ation of	riew	diagnosti
and therape	eutic methods	for optimu	in care of	the b	urn p	atie	nt as we	ell as di	usen	ination
of the scie	entific advand	ces to mili	tary and c	ivili	an me	dica:	l treat	nent cent	ers.	
24. (U) Th	nermally inju	red patient	s both from	m the	Cont	inen	tal Uni	ted State	es an	đ
throughout	the world are	e evacuated	l to the US	Army	Inst	itut	e of Su	gical Re	esear	ch for
intensive i	inpatient the	rapy. Care	fully cont	rolle	d cli	nica	l evalua	ation of	the	efficacy
of many tre	eatment modal:	lties is ur	dertaken.							
25. (U) 75	5 01 - 75 12	During 197	75, 254 bur	n pat	ients	wer	e admit	ted to th	ne Ir	stitute of
Surgical Re	esearch. Spec	cific atter	tion to th	e ear	ly di	agno	sis and	treatmen	nt of	inh <b>ala-</b>
tion injury	y involved the	e use of fi	beroptic b	ronch	oscop	e ar	i xenon	scan for	the	diagnosi:
and treatme	ent of inhala	tion injuri	les. In a	doubl	e-bli	nd s	tudy st	eroids in	n inh	alation
injuries we	ere found to l	oe non effi	cacious.	Early	flui	d re	suscita	tion was	eval	uated with
ongoing at	cention to the	e role of d	colloid in	the f	irst	24 h	ours po	st burn.	As	in
previous ya	ears, pulmonar	ry infectio	on with gra	m-neg	ative	org	anisms a	continues	5 CO	be a
frequently	observed com	plication of	or thermal	injur	y and	int	ensive :	Investig	atior	in _
methods of	prevention a	nd treatmer	nt continue	s. C	ontin	ued	evaluat	lon of t	ne ro	le of
topical che	motherapeuti	c agents is	s proceedin	g wi	n an	atte	mpt to	lineat	e the	true
role of top	pical therapy	in the cas	e or the 1	arge	therm	al i	njury.	General	prin	ciples of
management	previously de	escribed at	this unit	rema	in ba	sica	TIA nuc	hanged.	Seve	ral new
clinical a	pproaches to t	the treatme	ent of the	exten	sive	then	mal inj	iry and .	lts	
complication	ons are presential. mainton	ntly Laing	evaluated.	The	se in	CLu	e earli	r excis	ion b	oth forma.
warmer and Available to contract	More rigorous	s pursuit c	of the pati	ent's	nutr	eran. itio	nal requ	ie patier lirement:	1 <b>C, 1</b> 3.	

DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 1498A I NOV 65 AND 1495 1 1 MAR 68 OR ARMY USE TARE OBSOLETE

### ANNUAL PROGRESS REPORT

# PROJECT NO. JANGE 102871R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY REPORT TITLE: CLINICAL OPERATION, CENTER FOR TREATMENT OF BURNED SULDIERS

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 January 1975 - 31 December 1975

Investigators:

Hugh D. Peterson, DDS, MD, Colonel, MC James Dorethy, MD, Major, MC Jerold Z. Kaplan, MD, Major, MC Thomas S. Lescher, MD, Major, MC W. Scott McDougal, MD, Major, MC Richard H. Merrill, MD, Lieutenant Colonel, MC Arthur Rosenthal, MD, Major, MC James K. Sims, MD, Major, MC David K. Teegarden, MD, Major, MC Gary W. Welch, MD, Lieutenant Colonel, MC Douglas W. Wilmore, MD Marbeth G. Michael, Lieutenant Colonel, ANC Hilda M. Diaz, Captain, AMSC David N. Herndon, MD, Captain, MC Norris F. Jesse, Captain, AMSC John P. Peterson, Captain, AMSC Bast! A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

### ABSTRACT

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

REPORT TITLE: CLINICAL OPERATION, CENTER FOR TREATMENT OF BURNED SOLDIERS'

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 January - 31 December 1975

Investigators: Hugh D. Peterson, DDS, MD, Colonel, MC James Dorethy, MD, Major, MC Jerold Z. Kaplan, MD, Major, MC Thomas S. Lescher, MD, Major, MC W. Scott McDougal, MD, Major, MC Richard H. Merrill, MD, Lieutenant Colonel, MC Arthur Rosenthal, MD, Major, MC James K. Sims, MD, Major, MC David K. Teegarden, MD, Major, MC Gary W. Welch, MD, Lieutenant Colonel, MC Douglas W. Wilmore, MD Marbeth G. Michael, Lieutenant Colonel, ANC Hilda M. Diaz, Captain, AMSC David N. Herndon, MD, Captain, MC Norris F. Jesse, Captain, AMSC John P. Peterson, Captain, AMSC Basil A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

Two hundred and fifty-four patients with thermal injury were admitted to the Clinical Division of the United States Army Institute of Surgical Research during the calendar year of 1975. The Institute's main emphasis continues to be providing optimal care to military personnel and authorized civilians with major thermal injuries. Clinical investigation has continued into the physiology, biochemical and bacteriologic aspects of the mal injury. Continued evaluation of the role of topical chemotherapeutic agents is proceeding with an attempt to delineate the true role of topical therapy in the care of patients with extensive thermal injury. Several new clinical approaches to the treatment of the extensive thermal injury and its complications are presently being evaluated. These include earlier excision both tangential and at the level of the investing fascia, maintenance of a warm microenvironment to decrease energy needs, and improved nutritional support. This report summarizes the activity of the Clinical Division of the United States Army Institute of Surgical Research in 1975 and cites the recognizable complications which have contributed to mortality and morbidity in burn patients.

Thermal injury Resuscitation Air evacuation Topical therapy

•

Heterograft Autograft Homograft Inhalation injury

### CLINICAL OPERATION, CENTER FOR TREATMENT OF BURNED SOLDIERS

The Clinical Division of the United States Army Institute of Surgical Research continued through the year of 1975 to have as its primary objective the provision of clinical care for the thermally injured soldier. The number of admissions increased from 244 in 1974 to 245 patients in 1975.

In 1975 there were a total of 87 aeromedical evacuation flights, 86 of these were CONUS flights (within the Continental United States) with 108 patients evacuated. All patients within a radius of 200 miles of Brooke Army Medical Center requiring air evacuation were transported by helicopter. There were 18 flights for that purpose. There was one flight outside of the Continental United States that being to Alaska.

### CLINICAL MANAGEMENT

In depth description of the management of patients with thermal injury as practiced by this institute are found in previous Annual Reports and in numerous scientific publications. Therefore the following remarks will be limited to new and current methods of clinical therapy.

1975 was the first year in which silver sulfadiazine was used exclusively as the initial primary topical chemotherapeutic agent. The initial success noted during the period of comparison treatment with Sulfamylon in 1974 did not continue to the degree first noted. During 1975 there was a remarkable increase in the resistance of the Klebsiella cultured from the wounds to silver sulfadiazine. A complete change of the predominant organism in the burn wound from Enterobacter cloacae to Klebsiella was seen. This increased resistance was heralded by earlier time of positive blood culture, earlier onset of sepsis and earlier fime to death in the large burns. There has also been a decrease in the 1050 in the 15 to 40 age group so that it is now less than the 63.5 noted in 1974. The premises of the earlier comparison study were again born out through 1975 that being that Sulfamylon offers a far better bacterial control of the gram negative flora of the burn wound, but may exert deleterious systemic effects on patients with burns greater than 30% in the early post burn period.

Early pulmonary complications characterized by hyperventilation, respiratory alkalosis with progression to a fluid overload pattern and the appearance of "shock lung" were rare in 1975. The major pulmonary complications in 1975 were fluid overload secondary to vigorous resuscitation which responded rapidly to diuresis and fluid restriction and pneumonitis secondary to both inhalation injury and gram negative infection per se. The lack of significant early pulmonary complications except those related to severe inhalation injury bespeaks the benign properties of silver sulfadiazine as an initial topical agent, however, during 1975 this positive quality began to be outweighted by silver sulfadiazine's lack of efficacy against the gram negative population of the burn wound and a re emergence of Pseudomonas organisms in the burn wound of terminal patients. However it remains our opinion that Sulfamylon does offer better bacteriologic control of the burn wound but it may cause marked hyperventilation in patients with burns greater than 30% and it is our opinion that it should be reserved for use as the topical agent for treatment of heavily colonized or infected burn wounds.

The ¹³³Xenon lung scan, fiberoptic bronchoscopy, and pulmonary function testing continued to be used for the documentation of inhalation injury. As has been noted in the past we have obtained falsely positive xenon scans due to chronic lung disease or acute upper respiratory infection which existed prior to burning. However, when the ¹³³Xenon lung scan is correlated with bronchoscopy falsely positive diagnoses are rare. At present we are evaluating nebulized gentamicin in a double blinded fashion in patients with positive findings of inhalation injury. To date no results are available. At present we administer large doses of steroids systemically and aerosolized racemic epinephrine in the treatment of upper airway inflammation to reduce or limit edema formation and avoid intubation. Attempts are being made to correlate pulmonary function with the status of the acute upper airway before and after such treatment in an effort to document its efficacy.

In 1975 as in previous years the two major fluid and electrolyte problems seen at the Institute of Surgical Research were hypernatremia and hyponatremia. Hypernatremia reflecting a water deficit was found to be the most common electrolyte abnormality in the burn patient. Hyponatremia is the most common electrolyte change in the burned child and is often manifested by seizures. It is caused either by excessive administration or too rapid administration of electrolyte free fluids or the "leeching" effect of silver nitrate dressings when used. These findings again call for caution in the administration of nonelectrolyte fluid in small children and for the careful monitoring of all patients in silver nitrate dressings and for careful observation of all patient's electrolyte status during resuscitation.

Interest has continued during 1975 in primary excision both to the ievel of the investing fascia and tangential. Twenty-one formal excisions to the level of the investing fascia were done while 42 tangential excisions were carried out in 1975, 23 of these being excision of burns of the hand. The excisions to the level of the investing fascia fell into two categories; excisions of massive burns with immediate coverage with cutaneous allograft or xenograft, all of which have done poorly and excision of localized full thickness injury in patients with moderate sized burns 40-60%, all of which have done well in 1975. Evaluation of tangential excision continued throughout 1975. Tangential excisions also fall into two groups. The first being tangential excision of burns of the hands with immediate autografting. This is carried out on deep second degree burns of the hand that have no significant areas of third degree involvement. The functional results have been excellent and this treatment has much to recommend it over scalpel excision of the hands. When the hands that were tangentially excised had deep second degree burns, the results were uniformly excellent. In the few cases where full thickness injury was recognized and excised, the results were indistinguishable from classical conservative treatment. Tangential excision was also employed in patients having full thickness wounds with tenacious eschar. The eschar was tangentially excised until freely bleeding yellow fat was encountered. These areas were either covered with biologic dressing or with Sulfamylon soaks and allowed to granulate. The wound matured and good granulation tissue formed in seven to 14 days permitting subsequent "mesh" grafting.

Interest has continued in the metabolic status of the burn patient during 1975. Glucose flow through the extravascular compartment has been evaluated and found to be significantly elevated in 17 burn patients between the 6th and the 16th post burn day when compared to normal individuals. This shows an increased rate of gluconeogenesis in the second post burn week that was directly related to the extent of injury and with time returned in a curvilinear manner towards normal with closure of the burn wound. In contrast to this glucose flow decreased towards normal in the patients with gram negative infection. The elevated glucose flow in patients with extensive burn injuries drops precipicously towards normal with onset of gram negative sepsis. Simultaneous measurements of glucose flow and oxygen consumption demonstrated a close relationship between gluconeogenesis and heat production.

Glucogon administration in normal man was also evaluated. Increased heat production occurred with glucogon administration in a hypothermic elderly burn patient. This increased heat response is closely related to and appears concomitantly with translocation of glucose to the periphery, but cannot be duplicated by glucose infusion alone. The increased caloric needs of the burn patient appear to reflect the increased energy requirements of accelerated gluconeogenesis.

During 1975 intragastric instillation of antacids was used as prophylaxis against the complications of Curling's ulcers. Sufficient antacid was instilled on a q i to 2 hour basis to maintain the pH of the stomach contents at 7.0 or above. This has been attended by a marked decrease in clinical complications of stress ulcers so that only one major GI bleeder required operation in 1975. We have made antacid prophylaxis a routine in the early post burn course with excellent results.

### EDUCATION

During the period of this report, _ irgical residents from Brooke Army Medical Center, 1 from Fitzsimons, 3 . om David Grant USAF Medical, Center, Travis AFB, California and 1 civilian physician from Northwestern and 1 resident from the University of Wisconsin at Milwaukee participated as active members of the medical staff for periods of 2 months to 1 year as part of their surgical training. A physician from the Navy of Sweden,

two Army and one Navy reservist completed their tours of active duty training with our unit and one senior medical student from the University of Texas in San Antonio, one from Northwestern, and one from the University of Missouri School of Medicine spent from two weeks to one month in observer training. Twenty-five Air National Guard officers and paramedical personnel from one unit were given a two hour tour and briefing. Approximately 93 civilian and military physicians and 130 nurses, students and paramedical personnel visited and were briefed during 1975. Forty-two foreign visitors from the following countries: Beigium, Japan, France, Norway, Ireland, Israel, Sweden, England, Switzeriand, Germany, Denmark, Korea, Republic of Zaire, Holland, Uruguay and the Republic of Vietnam, received Griefings on the overall mission of the Institute of Surgical Research and the care of the thermally injured patient.

Numerous scientific presentations concerning various aspects of thermal injury were made by members of the Clinical Division at local, state, regional and national meetings as listed at the end of this section.

### STATISTICAL RESUME

During the year 1975, 245 thermally injured patients were admitted to the institute of Surgical Research. There were 254 dispositions during 1975 and the subsequent data will be based upon those dispositions. The patients ranged in age from 6 weeks to 85 years with 208 males and 46 females. The average age of the patient was 29 years with an average burn size of 42.1% and a 17.9% average third degree component. The average burn index was 29.7. Out of 254 dispositions, 197 had third degree burns. Forty patients were less than 15 years of age with an average age of 4.75 years. The average total burn size in the pediatric age group was 32.2% with a 14.2% third degree. The burn index in children was 22.5 Of the 40 pediatric patients admitted, 24 had some third degree burn (60%).

The mortality in the pediatric age group was 20%. In the group of pediatric burn patients who died, the average age was 5.5 years and the average burn size was 64.8% with 44.6% third degree injury. The overall mortality for the year 1975 was 37% or 94 patients out of the 254 dispositions. Of the patients expiring 73 were male and 21 were female. The average age of the patients who died was 31.8 years, the average total burn size was 61.3% with 32.8% being third degree involvement, with an average burn index of 46.3%. There was a 6% decrease in raw mortality but the mortality in general remains higher than during the period when many patients were air evacuated from Southeast Asia. Of the 94 patients who expired 89 or 94.9% had some areas of third degree burn. Autopsies were performed in 72 patients or 76,6% of all deaths. The average post burn day of death was 16.6 which was decreased from the 21.4 mean time to death in 1974 but still significantly longer than the 11.8 days in 1973. Table 1 identifies the source of admission of patients during calendar year 1975. The majority of the patients were

† rea	A	AD	AF	AFD	N	ND	VAB	0ther	TOTAL
lst Army	4	0	1	0	0	0	5	4	14
3rd Army	5	5	3	3	7	2	7	14	46
5th Army	18	11	9	9	7	2	16	63	135
Sth Army	2	0	1	3	2	2	5	8	23
1DW	1	0	0	0	C	0	0	0	1
\laska	0	0	1	0	0	Ò	0	1	2
Germany	5	3	2	0	1	0	0	0	11
Ecuador	0	n	0	0	0	0	0	2	2
lawali	0	0	0	0	0	0	0	1	1
Mexico	0	0	0	1	0	0	0	4	5
San Salvador	0	1	0	0	0	0	0	0	1
Okinawa	1	1	1	0	1	0	0	0	1
Korea	1	0	0	0	0	0	0	0	1
Costa Rico	0	0	0	0	0	0	0	1	1
Panama	0	1	0	0	0	0	0	1	2
Honduras	0	0	0	0	0	0	0	4	4
Virgin Island	0	0	0	0	0	0	0	١	۱
	37	22	18	16	18	6	33	104	254

Table 1. Source of Admission, 1975

D - Dependent Other: Civilian Emergency

US Public Health Service Beneficiary

Bureau of Employees Compensation Beneficiary

	. <u>9</u>		254	1014
0.0\$	0	1.6%		Mortar Shell Exp.
82.43	Ŧ	6.78	17	Smoking Cisthës Ignited
11.1\$	-	3.5\$	و	Welding Accidents
42.9\$	15	13.8\$	35	Bucar, Propane or Natural Gas Exp.
41.28	7	6.72	17	Others
50.0\$		0.8%	2	Chemi ca i
4.8\$	-	8.3\$	21	Hot Liquid
7.18		5.5\$	14	Electrical
34.8%	8	9.0%	23	Open Flames
54.5\$	6	4.38	11	Aircraft Accidents
13.3\$	2	5.9\$	15	Motor Vehicle Accidents
61.5\$	16	10.2%	26	Structural Fires
36.78	22	23.6%	8	Gasoline & Kerosene
t Hortality	Deaths	° Disposition	Number of Patients	Causes

-

٠

-

Table 2. Burn Etiology, 1975 - 254 Dispositions

					.								
1				Pei	r Cent B	L.					Total	Total	•
(K'N)	9 9	<b>R-</b> 0	<b>R</b> - <b>R</b>	9-2	8-9	8	8-9	2	8	8	<b>5963</b>	Petho	hertality
0	-	-	-		-	(1)(	0	•	0	0	•	-	16.7
<b>1-2</b>	~	~	-	•	(1)	0	0	0	•	0	۲	-	14.3
2-3	. 0	~		~	-	0	(i) 1	•	•	0	•	-	12.5
ī	0	0	0	0	0	0	•	Ē	0	0	-	-	100.0
1	-	2	0	0	0	0	•	ò	0	0	m	0	0
5-10	0	8	-	-	0	-	(1)1	0	0	(i) 1	٢	2	28.6
10-15	~	0	7	2(1)	•	-	-	0	(1)1	0	đ	64	25.0
15-20	4	ų	7	-	(1)†	-	5(4)	<b>3(1)</b>	0	2(2)	27	•0	<b>29.6</b>
20-30 20-30	m	12	•	13(3)	•0	13(5)	17(10)	7(5)	7(6)	•	8	8	32.6
9-9.	ŝ	-	7	7	3(1)	2(2)	2(1)	(i)(	0	(1)	27	•	22.2
95-9 <del>1</del>	~	-	m	7(4)	(I) <del>1</del>	7(5)	3(2)	4(4)	(1)	0	8	11	51.5
<b>3</b> 8	0	0	3(3)	(1)9	4(2)	2(1)	(1)	2(2)	(i) (i)	(i) (i)	8	12	6.0
R-9	0	0	2(1)	2(2)	(1)	0	<b>(</b> C)C	(i) (i)	2(2)	0	1	9	6-06
8- R	•	2(1)	-	0	(i)(	0	0	¢	(1)	0	Ś	~	0.0)
	•	0	-	ġ	0	0	0	(1)	0	0	~	-	6.0
Totaî		×	R	8	*	*	*	2	5	~	ž		
ri de	•	-	-	=	•	4	ដ	¥	13	~		\$	
\$ Marcallty	•	2.8	13.4	*	28.6	8	67.6	<b>M.</b> 2	92.3	8			37.6
The second se													

Table 3. Age, Body Surface levelwamment & Mortellity, 1975

ないない。ないない

No. of Contraction

** <b>B</b> um	0-14	10-20	20-30	5-5	5-8	8-8 8	60-70	8-8	8-8	90-100	Total
					(1972	~					
lo. Burned	5	8	43	42	8	23	22	16	=	S	100
beaths		2	7	13	15	13	21	51	=	Ś	103
Hortality	2.1		16.3	2	41.7	УЯ .5	95.4	93.8	8	8	94 is
	1	1		1	(1973	~			;	: : : :	
to. Surned	8	X	\$	X	ä	×	8	16	12	•	273
beaths	0	0	7	٠	21	2	17	5	12	•	113
Nortality	   0 		15.2	34.6	· · · · ·		, <b>3</b> 7	; ; §	8	. 8	
					<b>K</b> (I)	č					
to. Burned	¥	*	28	X	×	28	IJ	5	16	v	226
beachs	0	0	*	5	5	17	17	•	13	•	97
t toreality	; •		14.3	31.2	6 <b>8</b>	8.7	<b>3</b>	8	8	8	
					(1975	0					
lo. Jurned	8	×	3	5	¥	2	¥	J	13	v	ş
hache	•	-	<b>8</b> 7	1	•	Ŧ	8	¥	8	~	*
t Hertallty	0	2.8	13.8	X.2	¥.6	8	67.6	<b>\$</b> -7	Q.3	8	¥

Table 4. Per Cent Body Surface Involvement and Hortality, 1972 - 1975

	Surviv	ors (bu <b>rns</b>	over 302)		Spoths	
Year	No. Cases	Average 1 Total	<u>Burn</u>	No. Cases -	Average ( Yotal	L turn,
1955	20	39.5	20.3	21	55.6	38.1
1956	22	41.0	17.3	20	<b>57.8</b>	37.8
1957	19	38.4	24.1	17	57.1	38.8
1958	15	42.3	21.6	23	56.5	35.3
1959	29	43.1	20.6	24	63.1	<b>38.</b> 1
1960	17	44.2	20.1	30	57.8	37.3
1961	18	44.2	25.0	97	58.0	39.7
1962	18	42.7	21.4	54	59.1	46.2
1963	28	45.8	19.6	57	<b>69</b> .9	41.0
1964	40	41.8	14.8	37	65.0	42.4
1965	47	43.8	21.0	33	66.0	33.4
1966	68	41.5	14.9	59	59.9	31.3
1967	103	42.7	13.3	51	59.9	32.3
1968	143	44.2	12.6	38	54.6	24.6
1969	113	43.2	11.1	70	58.7	26.4
1970	<del>9</del> 2	39.4	10.7	70	51.9	32.6
1971	63	41.9	14.0	68	60.8	38.0
1972	62	42.0	17.2	103	56.7	35.9
1973	47	43.7	19.6	113	60.3	36.2
1974	55	43.9	12.2	97	60.8	35.5
1975	80	46.1	14.7	94	61.3	32.8

Table 5. Per Cent Burn Versus Survivel, 1955-1975

							3	T Cont	Wm						
		9			¥	•		ş	8		8	8		6-10	δ
				Į								,	•	F	•
	<b>.</b>	No.	t hortaìlty	<b>₽</b> 5.	No.	1 Hortallty		No. Deaths	<b>\$</b> Hortallty	ਕ ਛ	No.	1 Hortallty		Deaths	Hortality
	ś	-		*	-	5	×	n	61.1	23	ä	78.3	55	\$	<b>3</b>
The second				S.	:	18.0	123	IJ,	31.7	275	137	<b>19.8</b>	2	421	85.2
61	Ĩ	X													

D
-
•
š .
-
<u> </u>
÷ .
,
_
<b>1</b>
•
-
•
÷.
1
•
-
_
-
2
2
2
2
2
2
ty Reta
ty Rates
ty Rates
ty Rates,
ty Rates,
ty Rates,
ty Rates, I
ty Rates, 19
ty Rates, 194
ty Rates, 196
ty Rates, 1962
ty Rates, 1962-
ty Rates, 1962-
ty Rates, 1962-1
ty Rates, 1962-19
ty Rates, 1962-191
ty Rates, 1962-196
ty Rates, 1962-1963
ty Rates, 1962-1963
ty Rates, 1962-1963
ty Rates, 1962-1963 a
ty Rates, 1962-1963 an
ty Rates, 1962-1963 and
ty Rates, 1962-1963 and
ty Rates, 1962-1963 and
ty Rates, 1962-1963 and 1
ty Rates, 1962-1963 and 1
ty Rates, 1962-1963 and 19
ty Rates, 1962-1963 and 196
ty Rates, 1962-1963 and 196
ty lates, 1962-1963 and 1964
ty Rates, 1962-1963 and 1964
ty Rates, 1962-1963 and 1964-
ty lates, 1962-1963 and 1964-1
ty lates, 1962-1963 and 1961-19
ty lates, 1962-1963 and 1964-191
ty Rates, 1962-1963 and 1961-1972

from the Continental United States. Table 2 summarizes the burn etiology in 1975. Table 3 summarizes the effect of age and total body surface burn on mortality.

Table 4 lists the mortality rates in increments of 10% total body surface from the year 1971 through 1975. Table 5 presents the survival and mortality rate of patients with greater than 30% body surface burns in the calendar years 1955-1975. It should be noted that there are no marked changes. Table 6 shows a comparison of burn mortality in the pretopical antimicrobial years 1962-1963 with the cummulative index since 1965 when Sulfamylon and later silver sulfadiazine have been used. As previously reported the improvement is primarily in that group of burn patients whose injury is 30-60% with little or no change in those patients with greater than 60% burn injury.

The average hospital stay in 1975 was 49.8 days. When convalescent leave for active duty military personnel was excluded the average hospital stay was 42 days. The average post burn day of admission to the Institute of Surgical Research was 2.9. This figure reflects the decrease in the average post burn day of admission from 11.2 days in 1970 to 9 days in 1971 and 7 days in 1972 and 5 days in 1973, and was the same as the 2.8 days in 1974. The decrease in the average admission time is because the patient population originates in the Continental United States and also reflects the rapid aeromedical evacuation currently practice ) at the Institute of Surgical Research.

During the year 1975 1,201 operations were performed on 211 patients, or an average of approximately 5 operations per patient. Four hundred and ninety anesthetics whre performed on 1½ patients or three anesthetics per patient participating. The total of 670 ward procedures were performed. Three hundred and thirty-one autografting procedures were carried out on 114 patients or approximately 2 1/2 procedures per patient. Thirty-one patients had 76 applications of allograft for an average of approximately 2.2 applications per patient participating.

Porcine cutaneous xenograft was applied to 17 patients 28 times or an average of 1 1/2 times per patient. There was a marked decrease in the use of porcine cutaneous xenograft during 1975. Evaluation of this material on wounds ready to accept autograft and wounds not clean enough to autograft revealed that it is clearly inferior to cutaneous allograft. Cadaver allograft was aseptically harvested from 60 cadavers, an increase from the 33 in '974. Escharotomies were performed on 80 patients or 31.5% of all dispositions. Twenty-five patients required major amputations usually for treatment of severe electric injury. Tracheostomy was performed in 16 patients or 6.3% of all dispositions. This is marked diminution from the 16.4% in 1974. This probably reflects both the decrease in number of early pulmonary complications and a tendency to leave endotracheal tubes in place for longer periods.

One hundred and ninety-two patients or 75.6% of all dispositions

had at least one blood culture drawn during their hospitalization and 115 patients had one or more positive blood cultures. For further information of the bacteriologic data the reader is referred to the succeeding chapters. Suppurative thrombophlebitis occurred in 8 patients or 3% of all dispositions. In keeping with the high index of suspicion of suppurative thrombophlebitis, 35 "cutdown" sites were explored. An important aspect of burn therapy is monitoring the flora of the burn wound and this was done with frequent wound biopsies. Eighty-four patients or 33% of all dispositions in 1975 had at least one burn wound biopsy performed.

A total of 740,885 cc of blood were administered to 164 patients or 64.6% cf all dispositions. An average of 4,518 cc of blood were given to each of the 164 patients. Thirteen patients or 5.1% of all dispositions had a major fracture upon admission. Two patients were also admitted with hip dislocations and two with traumatically open knee joints. The aforementioned 25 major amputations were all related to electrical injuries with the exception of two patients.

There were frequent ophthalmologic injuries in the 254 dispositions in 1975 which included corneal burns, corneal abrasions, corneal lacerations but mainly thermal injuries to the lids. Significant long term sequelae of eyelid burns occurred in the form of all degrees of ectropion. Of the surviving patients in 1975 there was only one with permanent visual impairment, that occurring secondary to a corneal scar from a deep corneal burn. Again in 1975 chondritis in the thermally injured ear has been a major problem. In 1975 27 patients or 10.6% of all dispositions had chondritis. This is double the incidence in 1974 and continues to underline our failure in understanding this process. The current treatment has emphasized keeping pressure off of burned ears and keeping them covered with a topical antimicrobial agent, however a good portion of the chondritis seems to be unresponsive to these procedures.

Gastrointestinal complications played a lesser role in 1975. Although some GI bleeding was noted in 60 patients, the bulk of these had only gualac positive stools and frank ulceration was diagnosed in only 33 patients or 12.9% of all dispositions, the majority of these were diagnosed at post mortem examination and many were considered to be terminal changes. There was only one GI bleed severe enough to require surgery during 1975 but there were three ulcer perforations, all occurring as terminal events. This lack of significant GI complications may well reflect the effectiveness of prophylactic antacid neutralization of gastric acid in all early post burn patients. Superior mesenteric avery syndrome was diagnosed radiographically in three patients; all were managed conservatively and had favorable outcomes. This stands in contrast to 1974 when three were diagnosed, and two were operated upon. with unfavorable outcomes. It would seem from this small series that a vigorous trial at conservative therapy is indicated prior to operative intervention. The only other significant GI complication occurred in a patient with idiopathic atony of the right colon which eventually reached 16 centimeters in diameter and required a cecostomy for decompression. Major renal complications in our burn population occurred in 43 patients or 17% of all dispositions. This was usually a terminal event often related to sepsis expressed as some degree of renal failure. In fact the bulk of the renal problems in the burn patients appear to be secondary to sepsis. Hemodialysis was carried out in nine patients with no survivors.

Cardiac complications continued to play a significant role in burn patient morbidity and mortality. Again as in 1974 nine patients sustained acute myocardial infarction and again all succumbed from the insult. As in 1974 there have been several episodes of acute bacterial endocarditis with staph aureus coagulase positive being the causative organism. Four cases of endocarditis were diagnosed in 1975 as compared to five in 1974 however there were multiple patients that had blood cultures positive for Staphylococci which cleared on therapy in whom the source of bacteria remained undefined. This lesser incidence of staphylococcal endocarditis may reflect the increased awareness of the lethal potential of Staphylococcal septicemia in the burn patient and the beneficial effect of starting antibiotic therapy with the first positive culture.

Pulmonary problems continued to be a significant cause of morbidity and mortality in 1975 as they have in the past. However, early pulmonary problems appear to have decreased in frequency except for those of inhalation injury. There were 60 patients with inhalation injuries documented on ¹³⁵Xenon lung scan and bronchoscopy or an incidence of 23.6% of all dispositions. It is again apparent that inhalation injury significantly increases the mortality in patients with 30 to 65% burns and has little if any effect on the mortality of patients with burns of more than 65%.

### SUMMARY

During 1975 245 patients were admitted to the U.S. Army Institute of Surgical Research and 254 dispositions were made. Silver sulfadiazine was continued as the initial burn therapy however, it appears to be sorting out a more resistant Klebsiella as documented by earlier time of positive blood culture, earlier onset of sepsis and earlier time to death. Again there were fewer significant pulmonary complications and the patients had a much more benign early post burn course. Infection continues to be the most common cau e of mortality in the thermally injured patient. Clinical research efforts are oriented towards evaluation and control of the septic process with ongoing research on the efficacy of various topical agents, the role of hydrotherapy in the management of burn wounds, the use of systemic antibiotics, immunosuppression with excision and allografting in massive burns and other means to improve the survival of the burned soldier.

*Autonsy not performed

Patient	Age	Sex	<b>t</b> bu Total	Ē	PBD Death	Cause of Death
-	6	x	95	16	21	Septicemia (staphylococcal and Pseudomonas), severe "adult hyaline mombrane disease"
8	36	T	95	62.5	-	Severe inhalation injury
ŝ	8	x	92.5	80.5	2	Severe inhalation injury
4	18	T	30.5	55	8	Gram-negative sepsis (Enterobacter, Pseudomonas)
2	15	ţı.	8	8	7	Sepsis (Klebsiella); acute renal fallure
Q	:	X	&	ጽ	*	Severe hyaline membrane disease, severe bronchopneumonia
7	68	<b>'</b> 2-	85	8	-	*Cardiovascular collapse
30	63	X	85	54	-	inhalation injury and cardiovascular collapse
Ø	28	x	84.5	2	-	Severe inhalation injury
0	٩	x	83	I.	~	Severe inhalation injury
=	26	x	82.5	47	15	Sepsis (Enterobacter, Klebsiella and Pseudomonas)
12	22	T	82	£	Ś	Severe inhalation injury
13	25	<b>ای</b> .	82	43	41	Septicemia (Klebsiella, Enterobacter), gastric ulcer with hemorrhage
41	53	x	80.5	57.5	12	*Severe inhalation injury
15	27	٤.	80.5	8	0	*Septicamia (Klebsiella)
16	27	x	2	8	9	*Severe inhalation injury; septicemia (Pseudomonas)
11	\$	ż	0	22	15	Severe bronchopneumonia with abscess formation organism Staphylococcus aureus coagulase positive

Table 7. Causes of Death, 1975

No.

•

٩,

Table 7. Causes of Death, 1975	cause of Death	Aspiration of gastric contents	Burn wound sepsis organisms Klebsiella, liver failure	Arteriosclerotic heart disease manifested by acute myocardial infarction	Arteriosclerotic heart discase manifested by acute myocardial Infarction	Acute bacterial endocarditis organism Staphylcroccus aureus coagulase positive	Severe pneumonla secondary to inhaiation injury	*Severe inhalation injury	Severe inhalation injury	Hultiple pulmonary emboli	Cardiovascular collapse	Inhalation injury, sepsis (Enterobacter and Klebsiella)	Sepsis organism Pseudomonas	Cardiovascular collapse	Sepsis organism Klebsiella	*Severe inhalation injury	Burn wound sepsis, organism Pseudomonas	Acute bacterial endocarditis organism Staphylococcus	Nultiple pulmonáry emboli	
	PBD Death	4	41	4	19	65	r F	œ	2	m	7	13	37	6	12	2	17	\$	11	
	٤'n	7	91	67	15	61	34.5	61	58	43	29	17	38.5	33.5	54	o	1.5	x	53.5	
	\$ Bui Total	75.5	ጽ	78.5	78	78	77.5	76.5	76	75.5	75.5	73	73	73	11	יר	70.5	5.69	69.5	read
	Sex	HL	x	r	I	Σ	Ŀ	I	T	x	LL.	x	LL.	r	x	æ	T	T	٤.,	perfo
	Age	23	<u>.</u> ¥	65	141	18	28	34	45	28	85	29	ñ	21	58	23	42	28,	25	p\$y not
	Patient	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	*Auto

. . .

Pacient	ŧ	ž	Zotal	۽	De the	Cause of Beath
×	65	x	69.5	32.5	ຄ	Severs lobar pneumonía
37	17	z	\$	36.5	*	*Sepsis organism Staphylococcus
38	6 <b>4</b>	X	68	33.5	13	*Staphylococcal septicemia and severe inhalation injury
39	26	I	68	8	ĩ	Pseudomonas burn wound învasion
2	23	۱L.	68	ជ	23	Mycotic burn wound invasion, mycotic abscass left temporal lobe, severe pneumonia
41	18	L.	67	<del>1</del> 3	25	*Burn wound invasion organism Pseudomonas
42	-24	x	67	39.5	23	Severe inhalation injury, sepsis Enterobacter and Klobsiella
43	54	r	67	32	m	Severe inhalation injury
44	26	I	8	<b>Ş</b>	6	Diffuse bronchopneumonia and cardiomyopathy, secondary to Friedreich's ataxia
45	9	LL.	66	01	18	Aspiration gastric contents
<b>46</b>	61	x	3	5.5	9	Severe inhalaticn injury and bronchopneumonia organism Klebsiella
47	20	I	65.5	9 <b>4</b>	17	Sepsis organism Kiebsiella
48	8	z	65	42	٥,	Severe pneumonia organism Pseudomonas
617	21	I	65	40.5	15	Sepsis (Klebsiella), aspiration gastric contents
50	2	x	64	59	71	*Septicemia organism Klebsiella
15	20	L.	63	14	15	Inhalation injury
52	28	LL.	62	0	15	Septicemia, Staphylococcal
53	18	u.	61.5	8	38	Staphylococcal sepsis secondary to acute bacterial endocarditis
* Aut	ou J.sdo:	it perf	ormed			

# Table 7. Causes of Bach, 1975

Paclor	r Age	ž	Total	ε'n	Peet D	Cause of Death
Ł	\$	x	61	31	5	⁴ Sepsis organism E. coli
55	67	z	3	m	9	Acute renal failure and acute carabral adams
X	19	x	3	0	6	Sepsis organism, Klebsiella
23	33	54	57.5	13	16	inhaiation injury and acute necrotizing bronchopneumonia
<b>5</b> 8	24	x	57.5	3.5	0	#Sapsis organism Klebsiella
23	28	T	51	4	-	Multiple trauma including multiple fractures, lacaration right Illofemoral vein, messive intra-abdominal hemorrhage
3	2	æ	55	30.5	1	*Inhale2ion injury, bronchopneumonia
61	33	T	54.5	£3	28	Sepsis, organism Klebsiella
62	21	x	54.5	11	œ	Inhalation injury and severe bilateral bronchopneumonia
63	27	T	53	4	•0	Burn wound invasion organism Klebsiella and Pseuduronas
3	47	X	53.5	x	٢	*Severe inhalation injury and bronchopneumonia
65	47	Ξ.	5	ĩ	٢	Şepsit organism Pseudomonas
38	6/52	۹¢.	15	ĩ	10	Burn wound invasion multiple organic s, hematogenous pneumonia
67	\$	ł.	50.5	35.5	23	Sepsis and inhelation injury
8	27	X	8	31.5		Severe inhalation injury
<b>\$</b>	41	¥.	8	8		*Severs inhalation injury
ዩ	ず	E	8	\$	12	Sepsis organism Enterobacter
×	58 Autopsy n	vt ber tr tr	5. 64 Sommed	<b>t</b> e .5	2	Arterfoscierotic hear: divesse menifested by acute myocardiel infarct

•

, , ,

Table 7. Causes of Death, 1975

*Autopsy not performed

Pat i en t	Age	Sex	t Bur Totel	°.	PBD Death	Cause of Death
72	łş5	r	6 <del>1</del>	30	34	Burn wound sepsis organism Staphylococcus and Pseudomonas
73	-	r	48	22.5	ø	Septicemia organism Klebsiella
74	18	x	47	26	29	*Septicemia organism Staphylococcus
75	33	T	46.5	m	42	Inhalation injury, severe hyaline membrane disease secondary to Inhalation injury and respirator
76	52	r	42	32	20	*Seµticemia organism Enterobacter and Staphyiococcus aureus coagulase positive
11	68	x	4,2	12.5	4	Cardiovascular collapse secondary to severe arteriosclerotic heart disease
78	72	T	41.5	30	14	Diffuse bronchopneumonia and recent cerebro-vascular accident
Ŕ	48	T	39	31	-	Inhalation injury and cardiovascular collapse
8	45	x	36.5	34.5	17	Sepsis organism Klebsie}la
18	¢3	T	36	7	33	Gastrointestinal hemorrhage secondary to cecal ulcer and constrictive pericarditis
<b>B</b> 2	61	x	35	61	29	Severe arteriosclerotic heart disease and aspiration pneumonia
83	24	x	34.5	6.5	1	inhalation injury severe, bilateral pneumonia, acute bacterial endocarditis organism Staphylococcus aureus coagulase pusitive
84	28	T	34	25.5	4	*Disseminated intravascular coagulopathy and respiratory failure
85	9	٤.,	33	61	13	Sepsis organism Klebsiella
86	ጽ	>	33.5	0	10	lnhalation injury with severe bilateral bronchopneumonia secondary to Inhalation injury

~ -

-

Table 7. Cavses of Death, 1975

-

Patient	Age	Sex	<b>2</b> Bur Total	_ <b>ش</b>	PBD Death	Cause of Death
87	13	x	32	26	27	*Staphylococcal septicemia
88	38	I	8	8	24	*inhalation injury and septicemia organisms Klebsiella and Pseudomonas
ଛ	12	x	ጽ	ጽ	123	Severe cardiomyopathy etiology unknown
96	57	T	28	0	Ś	Severe inhalation injury
16	8	T	25	20	21	Cardiovascular insufficiency secondary to severe arteriosclerotic heart disease
92	Ť.	r	20	17	15	Sepsis, organism Klebsiella and renal failure
<b>3</b> 3	. 2	LL.	20	10.5	6	*inhalation injury with resultant respiratory failure
<b>4</b> 6	11	Ŀ	18.5	3.5	34	Multiple pulmonary emboli

-----

•

Table 7. Causes of Death, 1975

のないで、ため

ľ

### PRESENTATIONS

Long JM III: Treatment of Burns, Residents School of Aerospace Med, Brooks AFB TX 7 Jan 75.

Agee RN: Modern Burn Therapy. Medical Aspects of Advanced Warfare Course, USAF Sch of Aerospace Med, Brooks AFB, TX 14 Jan 75.

Erickson DR: The Dying Patient. Christian Medical Society of San Antonio, San Antonio, TX 14 Jan 75.

Slade CL: Treatment of Burns. Off Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 14 Jan 75.

Wilmore DW: Thermogenesis Following Thermal Injury. Workshop on Thermogenesis. University of Washington, Seattle, WA 17 Jan 75.

Berry DM: Nursing Research in the U.S. Army. ANC Officers, BAMC Fort Sam Houston, TX 20 Jan 75.

Peterson HD: Recent Advances in Burn Therapy. San Antonio Society of Insurance Underwriters, San Antonio, TX 20 Jan 75.

The following presentations were made to the Physical Therapy students Academy of Health Sciences, Fort Sam Houston, TX 27 Jan 75: Michael MG: Nursing Care of Burned Patients Long JM III: Current Therapy of Burns Fuller P: Introduction to Current Concepts of Burn Nursing Jesse NF: Physical Therapy for the Burn Patient Shaw AL: Occupational Therapy for the Burn Patient

Long JM III: Principles of Intravenous Hyperalimentation. Eastern Virginia Society of Hospital Pharmacists and Norfolk area nurses. Norfolk, VA 29 Jan 75.

Long JM III: Comparison of Carbohydrate and Fat as Caloric Sources for Total Intravenous Feeding. South Texas Chapter of the American College of Surgeons mtg, Galveston, TX 31 Jan 75.

Agee RN. Classification of Burns. Students Intensive Care Nursing Course, Brooke Army Medical Center, Fort Sam Houston, TX 3 Feb 75.

Taylor JW: Burn Wound Therapy. Students Intensive Care Nursing Course, Brooke Army Medical Center, Fort Sam Houston, TX 5 Feb 75.

Erickson DR: Complications of Burns. Students Intensive Care Nursing Course, Brooke Army Medical Center, Fort Sam Houston, TX 6 Feb 75.

The following presentations were given at the inservice Program for the Brooke Army Medical Center Nursing Service on 5 and 13 Feb 75: Berry DM: Nursing Research Michael MG: Current Concepts of Burn Nursing Podgornoff WC: Intermediate and Convalescent Care Zuber D: Care During the Acute State of Injury

Long JM [1]: Hyperalimentation. Students Intensive Care Nursing Course, Brooke Army Medical Center, Fort Sam Houston, TX 12 Feb 75.

Fuller P: Current Concepts of Burn Nursing. Students Intensive Care Nursing Course, Brooke Army Medical Center, Fort Sam Houston, TX 13 Feb 75.

Wilmore DW: Applied Science in the Care of Critically 111 Patients. Engineer Conference for Texas High School Seniors, Univ of Texas, Austin, TX 14 Feb 75.

Long JM III; Burn Injury: A Challenge to Patient, Doctor and Community. North Austin Rotary Club. Austin, TX 17 Feb 75.

Wilmore DW: Use of Intravenous Fat Emulsions. San Antonio Surgical Society, San Antonio, TX 18 Feb 75.

Long JM III: 1) Clinical Aspects of Intravenous Hyperalimentation. Loma Linda area Pharmacists, Nurses and Physicians. 2) What's New in Burn Therapy. Surgical Grand Rounds. Loma Linda University. 3) Problems in the Care of Burned Patients. Nurses of the Intensive Care Unit and Burn Unit. Loma Linda, CA 19 Feb 75.

Long JM III: Clinical Aspects of Intravenous Hyperalimentation. Inservice seminar. Santa Rosa Medical Center, San Antonio, TX 21 Feb 75.

Taylor JW: Treatment of Burns

Podgorn off WC: Nursing Care of Burns. Nursing students San Antonio College, San Antonio, TX 25 Feb 75.

Wilmore DW: Uses and Abuses of Parenteral Nutrition. Columbia Univ, NY 7 Mar 75.

The following presentations were given to the Overseas Industry Research Center Study Group on 10 Mar 57:

Pruitt BA Jr and Welch GW: Early Care of the Burn Patient, Including Fluid Resuscitation

Peterson HD: Wound Care - Topical Therapy

Levine NS: Wound Care - Excision

Taylor JW: Wound Care - Wound Closure

Long JM III; Metabolic Consequences of Thermal Injury and Nutrition of the Burn Patient

Petroff PA and Hander EW: Pulmonary Changes Following Thermal Injury Czaja AJ: Gastrointestinal Complications of Thermal Injury Merrill RH and Myers WD: Renal Changes Following Thermal Injury Andes WA: Hematologic Changes Following Thermal Injury Rizzo TA Jr: The Burn Autopsy and Burn Wound Biopsies
Michael MG: Orientation to the Nursing Care of the Burned Patient. 156th Aeromedical Evac Flt, NC Air National Guard, Charlotte, NC 12 Mar 75.

Long JM III: Clinical Aspects of Intravenous Hyperalimentation. South Illinois Soc of Hosp Pharmacists and Champaign-Urbana area nurses. Champaign, IL 14 Mar 75.

Wilmore DW: Metabolism Following Injury. Univ of Michigan Sch of Med, Ann Arbor, Mi 14 Mar 75.

Taylor JW: Burns. Boy Scouts of America. San Antonio, TX 17 Mar 75.

The following presentations were made at the American Burn Assn Anl mtg in Denver CO 20-22 Mar 75:

Levine NS: Limitations in Applying Tangential Excision to Large Burns

Taylor JW: Scalp as a Donor Site

Czaja AJ: Acute Liver Disease and Cholestatic Jaundice After Cutaneous Thermal Injury.

McAlhany JC: Perforated Curling's Ulcer: Incidence, Diagnosis and Therapy

Agee RN: Xenon¹³³ Lung Scan for Early Diagnosis of Inhalation Injury

Wilmore DW: Attempts to Modify the CNS Directed Stress Response to Thermal Injury

Petroff PA: Ventilatory Patterns Following Burn Injury and Effect of Sulfamylon

Long JM III: Effect of Carbohydrate and Fat Intake on Nitrogen Excretion During Total Intravenous Feeding

Michael MG: Inservice in a Folder

Baskin TW: Acute Bacterial Endocarditis A Silent Source of Sepsis in the Burn Patient

Wilmore DW: Use of Nutrition and Environmental Control for Burn Patients.

Wilmore DW: Recent Care of the Thermally Injurad. R.N. Club, Fort Sam Houston, TX 18 Mar 75.

Long JM III: Problems and Potentials of Intravenous Therapy. Illinois State Soc of Hospital Pharmacists, Chicago, IL 27 Mar 75.

Long JH III; 1) Clinical Use of Total Intravenous Feeding and, 2) Team Approach to IV Hyperalimentation. Physicians, Nurses, and Pharmacists. Clearwater, FL 1 Apr 75

Wilmore DW: Metabolism and Nutrition in the Septic Patient. Wayne State University, Detroit, MI 5 Apr 75.

Long JM [11: Inc menous Hyperalimentation. Louisiana Soc of Hosp Pharmacists and nurses. New Orleans LA 4 Apr 75. Peterson HD: Burn Complications. Nursing Inservice Program. BAMC Fort Sam Houston, TX 29 Apr 75.

Peterson HD: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX '30 Apr 75.

Long JM III: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX 2 May 75.

Long JM III: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX 5 May 75.

Long JM [11: Intravenous Hyperalimentation. Gastroenterology Service, BAMC, Fort Sam Houston, TX 12 May 75.

Long JM III: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX 16 May 75.

Long JM III: Intravenous Hyperalimentation. Central Texas: Society of Hospital Pharmacists, San Antonio, TX 20 May 75.

Wilmore DW: 1) The Role of the Central Nervous System Following Injury; 2) Catecholamines: Mediator of the Hypermetabolic Response Following Injury; 3) Nutritional Support in Injured Patients. International School of Medical Science, Erice, Italy 19-28 May 75.

Long JM [11: What's New in Burn Therapy. Staff of Harrisburg Polyclinic Hospital, Harrisburg, PA 4 Jun 75.

Long JM III: Use of IV Fat Emulsion in Burns and Trauma. AMA Symposium on Intravenous Fat Emulsions. Chicago, IL 6 Jun 75.

Wilmore DW: Essential Fatty Acid Deficiency in Burn Patients. AMA Symposium of Fat Emulsions. Chicago, IL 6 Jun 75.

Long JM III: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX 10 Jun 75.

Long JM [11: Practical Aspects of Total Intravenous Feeding. IV Therapy Seminar at the University of Indiana Medical Center, Indianapolis, IN 11 Jun 75.

Peterson HD: Current Therapy in Burns. Regional Meeting of San Antonio Society of Insurance Underwriters. San Antonio, TX 13 Jun 75. Wilmore DW: Metabolism Following Thermal Injury, Univ of Virginia, Richmond, VA 13 Jun 75.

Long JM III: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX 17 Jun 75.

Long JM III: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX 24 Jun 75.

Long JM III: Priorities for Care of Acute Thermal Injury. Academy of Health Sciences Physician's Assistant Program students, USA ISR, BAMC, Fort Sam Houston, TX 1 Jul 75.

Rosenthal A: Treatment of Burns. Officers Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 21 Jul 75.

Rosenthal A: Treatment of Burns. Officers Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 4 Aug 75.

Peterson HD: Plastic Surgery in the Military. Senior Army Chaplains BAMC CPE Chaplains Course, Fort Sam Houston, TX 7 Aug 75.

Stade CL: Classification of Burns - Initial Care. Students Intensive Care Nursing Course, Brooke Army Medical Center, Fort Sam Houston, TX 12 Aug 75.

Slade CL: Burn Wound Therapy. Students Intensive Care Nursing Course, BAMC, Fort Sam Houston, TX 14 Aug 75.

Rosenthal A: Treatment of Burns. Officers Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 18 Aug 75.

Erickson D: Complications of Burns. Students Intensive Care Hursing Course, BAMC, Fort Sam Houston, TX 19 Aug 75.

Slade CL: Hyperalimentation Therapy. Students Intensive Care Nursing Course, BAMC, Fort Sam Houston, TX 21 Aug 75.

Hayward C: Nursing Care of the Burn Patient. Students Intensive Care Nursing Course, BAMC, Fort Sam Houston, TX 26 Aug 75.

Peterson HD: Burns in the Multiple Trauma Patient. Sixth International Congress of Plastic and Reconstructive Surgery, Paris France. 26 Aug 75.

Pruitt BA Jr: Early Care of the Extensively Burned Patient. Beneral Surgery staff, residents, interns, Wilford Hall Med Ctr, Lackland AFB, TX 29 Aug 75. Pruitt BA Jr: Gram-Negative Infections in the Burn Patient. Symposium on Gram-Negative Infections, Baylor College of Medicine, Houston, TX 30-31 Aug 75.

Rosenthal A: Treatment of Burns. Officers Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 4 Sep 75.

Pruitt BA Jr: The Early Treatment and Complications of Burn Injury in Aged and Other High Risk Patients. Stanford Univ Sch of Med, Stanford CA 4 Sep 75.

Peterson HD: Burn Assessment and Management. Interns Brooke Army Medical Center, Fort Sam Houston, TX 8 Sep 75

Wilmore DW: The Neurohumorel Control of the Post-traumatic Changes. European Injury Meeting, Glasgow, Scotland 12 Sep 75.

Rosenthal A: Treatment of Burns. Officers Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 22 Sep 75.

The following presentations were made to the student nurses of San Antonio College on 23 Sep 75:

Rosenthal A: The Treatment of Burns

Taylor D: Nursing Care of the Burn Patient

Pruitt BA'Jr: Discussor for three papers: 1) Sepsis--Prevention and Management; 2) Respiratory and Gastrointestinal Complications; 3) The Future for Burn Care. American Burn Assn mtg, Kansas City, MO 25-27 Sep 75.

The following presentations were made and meetings attended by Pruitt BA Jr in connection with the ACS mtg San Francisco, CA 12-17 Oct 75: 1) Mtg, Editorial Subcommittee for Manual on Control of Infections; 2) Mtg, ACS Board of Governors; 3) Mtg ACS Executive Committee, Pre-and Postop Care; 4) Mtg Surgical Biology Club III; 5) Mtg ACS Committee on Trauma (Publications); 6) ABA Board of Trustees mtg; 7) ACS Pre and Postop Care Committee Mtg; 8) Mtg US Chap Internatl Soc of Surgery; 9) Panel Postgraduate Course No. 13 "Management of Facial Injury" presentation: The Acutely Burned Face; 10) ABA Ad Hoc Comm on Standards; 11) Adjourned mtg, ACS Board of Governors.

Wilmore DW: Alterations in Glucose Kinetics Following Thermal Injury. ACS Mtg, San Francisco, CA 15 Oct 75.

Rosenthal A: Treatment of Burns, Officers Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 16 Oct 75.

Peterson HD: Prospective Comparison Sulfamylon and Silver Sulfam diazine. American Soc of Plastic and Reconstructive Surgeons, Inc., Toronto, Canada 21 Oct 75. Wilmore DW: Parenteral Nutrition in Burn Patients. Massachusetts General Hospital, Boston, MA 29 Oct 75

Peterson HD: Plastic Surgery in the Military. Plastic Surgery Service, Wilford Hall USAF Hospital, Lackland AFB TX 30 Oct 75.

Pruitt BA Jr: Burns. Department of Surgery, Fort Leonard Wood Army Hospital, MO 6 Nov 75.

Peterson HD: Plastic Surgery. Third Annual Institute of the Assn of Operating Room Technicians of San Antonio, TX 8 Nov 75.

Pruitt BA Jr: Burns. Panel Member, Emergency Medical Service Patient Care Systems Design and Implementation Symposium, Grand Rapids, MI 11-13 Nov 75.

Pruitt BA Jr: 1) Prevention and Treatment of Sepsis in Burn Patients 2) Respiratory and Gastrointestinal Complications of Burn Patients; and 3) Unsolved Problems in Burn Care. ABA Seminar, Birmingham, AL 14-16 Nov 75.

Rosenthal A: Treatment of Burns. Officers Basic Course, Academy of Health Sciences, Fort Sam Houston, TX 17 Nov 75.

Peterson HD: Modern Burn Therapy. Medical Aspects of Advanced Warfare Course, Brooks AFB, TX 19 Nov 75.

Peterson HD: Recent Advances in Burn Management. General Surgery Service, University of Mississippi and Treatment of the Burned Hand, Plastic Surgery Service, Univ of Mississippi, Jackson, MS 21 Nov 75.

Pruitt BA Jr: Importance of Burn Units, Role in Education and Prevention and Patient Care Dedication of Baptist Medical Center Burn Center, Oklahoma City, OK 22 Nov 75.

Wilmore DW: Energy and Protein Metabolism in Disease. Ross Conference, Amiele Island, FL 22 Nov 75.

Pruitt BA Jr: Opportunistic Infection in Burn Patients. Charlotte Surgical Society, Visiting Professor Charlotte Memorial Hospital, Charlotte, NC, 24-26 Nov 75.

Pruitt BA Jr: Burn Wound Care. Surgical Staff, Wilford Hall Madical Center, Lackland AFB, TX 5 Dec 75.

The following presentations were made at the mtg of the Massachusetts Chapter of the APTA Workshop, Framingham, MA 4-6 Dec 75:

Peterson HD; 1) Early Medical Management of Burn Patients 2) Nutritional Management. Prevention and Treatment of Infection in the Burn Patient. 3) Skin Grafts and Wound Coverage for Burn Patients. 4) Reconstructive Surgery and Management for Burn Patients. 5) inhalatm ion injuries of Burn Patients.

Roberts ML: 1) Nursing Care in the Acute Phase. 2) Continued Nursing Care

Jesse F and Shaw'A: Acute Phase of Burn Treatment. 2) Long Term Care Phase of Burn Patients.

And the Second

Rosenthal A: Burns - Assessment and Management, Interns BAMC Fort Sam Houston, TX 8 Dec 75.

Rosenthal A and Taylor D: Management of Burns. Physical Therapy students, Academy of Health Sciences, Fort Sam Houston, TX 9 Dec 75.

Pruitt BA Jr: Pulmonary Complications of Thermal Injury. Plastic Surgical Symp of Nassau Surgical Soc, New York, 10 Dec 75.

Pruitt BA Jr: 1) Resuscitation and Early Care of the Severely Burned Patient; 2) Home Kemedies and the History of Burn Care; 3) Wound Care and Opportunistic Infections in Burn Patients. Visiting Professor at State Univ of New York at Stony Brook, Stony Brook, NY 11-12 Dec 75.

RESEARCH	AND TECHNOLOG	Y WORK UNIT S	UMMARY	I. ADENCY	ACCESSION	2 DATE OF SU	MARY	REPORT CO	WTROL STMBOL
S OATE POEN SUMPY	A PIND OF SUMMARY	T		DAUA		10 10 0		DD-DK	E(AR)636
75 07 01		8. SUMMANT 2017-	WORK SECURITY	7. REGRADI	NG" D. (	NSB'H INSTR'N	SA SPECIFIC C	ACCESS	LEVEL OF SUM
12 01 0T	H. TERM.	U	<u> </u>	NA		NL	E YES C	<b>) HO</b>	A. WORK UNIT
10 NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK ARE	EA NUMBER		WORK UNIT	NUMBER	
. PRIMARY	61102A		2 <u>B71R</u>	<u> </u>	1	168			
B. CONTRIBUTING				₋					
11 TITLE (Brace, ) with	Constitut Classification Code			<u></u>			<u></u>		la nor e h
of Indured		(0) Clini	cal Operati	lon, Su	irgical	Study Br	anch, Fo	or Tre	atment
DL LIIJULEU	SOLULEIS (44	1)							
003500 014	ad and the dial								
13. START DATE	nical Medicin	10		Tie gunnu	ADDICH				
<b>CD DD</b>	1				I AGENCT	I	16 TERFORMA	NCE METHO	0
17. CONTRACT/GRANT		/6	09				<u> </u>	In-Hou	.se
A DATES/EFFECTIVE	Not Applica	ble		H. RESOUR	CES ESTIMAT	TE A PROFESSI	ONAL MAN YRS	& FUND	i (In those ands)
		BAFINA HON:			76	1.	4	83	
C TVPE:		4		PISCAL TO	7T	·	3	20	_
A HIND OF AWARD									
IS. RESPONSIBLE DOD C	RGANIZATION	I.CUM. AMT.						1	
		L				IZATION .	L		L
HANE US Army	Institute of	Surgical R	esearch	HAME * U	IS Army	Institut	e of Su	rgical	Research
40000F881 ⁰					Surgic	al Study	Branch		
Fort	Sam Houston,	Texas 782	34	ADORESS -	Fort S	am Housto	on, Texas	s 782	34
				PRINCIPAL	INVESTIGATO	)R (Fumiah 36AN 1 	IUS. Academic	ine i (hution)	
HALSPONSIELL INDIVIDU	Druitt .Tr	COL MC		NAME."	Dougia	S W. WIIN	nore, M.I	D.	
HANE DUGIE A	-221_2720			TELEPHO	NE: 512	-221-5/12	2		
TELEPHONE: JLA	-221-2120			SOCIAL SE	CURITY ACC	OUNT NUMBER			
FORETCN TH			<b>5</b> 5	ASSOCIATE	INVESTIGATO	₩8 			
FOREIGN IN	TEPPTGENCE NO	JI CONSIDER	ED	HAME: L	. HOwa	rd Autics	., Ph.U.	, MAJ,	MSC
				HAME: E	Basil A	. Pruitt,	Jr., C	OL, MC	DA

(U) Trauma; (U) Combat casualties; (U) Laboratory animals; (U) Thermally injured soldiers: (U) Wound healing; (U) Gastrointestinal pathology 23 TECHNICAL OBJECTIVE,* 24. APPROACH, 25. PROGRESS (Pumlish Individual paragraphs identified by number precede test of each with Socurity Classification Code )

23. (U) Clinical and laboratory investigations pertaining to severe physical trauma which has been sustained by soldiers in the field.

24. (U) Planned clinical and laboratory studies relating to acute and chronic effects of injury. Studies conducted in this branch have included both clinical investigation involving patients and normal individuals and laboratory studies involving animal models.

25. (U) 75 07 - 76 09 The Surgical Study Branch continued investigations of the metabolic response to thermal injury and the effects of physical trauma on wound healing, susceptibility to infection, and gastrointestinal and liver function. The effects of nutritional support on posttraumatic metabolic response and organ system alteration is also being evaluated. The metabolic kitchen provides support for these investigations, in addition to preparation of specialized diets for patients with feeding problems, and the provision of diets containing known constituents to be used in a wide variety of metabolic balance studies. In addition, members of the branch provide nutritional, gastrointestinal and physiological consultation for all institute patients and other individuals outside the institute with complex gastrointestinal, metabolic, nutritional and temperature regulation problems.

### TERMINATION REPORT

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BROMEDICAL SCIENCE-SURGERY

**REPORT TITLE: CLINICAL OPERATION, SURGICAL STUDY BRANCH FOR TREATMENT OF INJURED SOLDIERS** 

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Douglas W. Wilmore, MD Louis H. Aulick, PhD, Major, MSC John P. Peterson, Captain, AMSC David H. Dahlquist, SP5, EM

Reports Control Symbol MEDDH-288(R1)

Unclassified

### ABSTRACT

### PROJECT NO. 3A161132B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

### REPORT TITLE: CLINICAL OPERATION, SURGICAL STUDY BRANCH FOR TREATMENT OF INJURED SOLDIERS.

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Douglas W. Wilmore, MD Louis H. Aulick, PhD, Major, MSC John P. Peterson, Captain, AMSC David H. Dahlquist, SP5, EM

Reports Control Symbol MEDDH-288(R1)

This branch studies the physiologic alterations which occur following severe injury and interrelates the changes in circulation, metabolism, temperature regulation, and hormonal control with alterations in the wound and the whole patient. The design and construction of a whole leg plethysmograph for measurement of leg blood flow and development of an on-line technique for continuous oxygen consumption measurement have been major technical advances in support of this investigative effort. Branch members provide consultative services to departments of the Brooke Army Medical Center and the University of Texas Health Science Center at San Antonio. Branch members are actively engaged in teaching programs on a local, national, and international level.

Surgical study Surgical physiology

### CLINICAL OPERATION, SURGICAL STUDY BRANCH FOR TREATMENT OF INJURED SOLDIERS

The three major activities of the Surgical Study Branch are: 1) delivery of medical and surgical care to the thermally injured soldier admitted to this institute; 2) clinical and laboratory research in problems related to care and rehabilitation of burn patients, and application of this knowledge to other critically ill patients; 3) the education of medical and paramedical personnel in the care of the seriously injured.

Members of this branch round daily to evaluate the patient problems in the Clinical Division of the US Army Institute of Surgical Research. The branch chief serves as coordinator of all clinical research, and all branch members provide consultative service and care in the areas of post-traumatic metabolism and nutrition, energy balance, gastrointestinal and hepatic dysfunction, peripheral circulation, and temperature regulation. Branch members provide consultative services for General Surgical, Medical, Endocrinology, Obstetrics and Gynecology, Gastroenterology, Pediatrics, and Dermatology Services at the Brooke Army Medical Center and the University of Texas Health Sciences Center at San Antonio. Techniques and modalities developed in this unit are currently applied to the care delivered to seriously injured patients who remain hospitalized until all wounds are healed.

Clinical and laboratory research in this division may be placed in the following categories: 1) temperature regulation; 2) posttraumatic metabolism; 3) energy balance; 4) nutritional support of critically 11 soldiers; 5) relationship between peripheral circulation and substrate flow to the burn wound; 6) description of hepatic dysfunction which occurs following injury, with specific emphasis on the interrelation between hepatic glucose production, splanchnic blood flow, and energy production of the burn patient. Two technical developments have aided in these areas: 1) construction of a whole leg plethysmograph, and 2) development of an on-line continuous measurement system for total body oxygen consumption (by Mr. Edwin W. Hander). In addition, the development of a hypermetabolic animal model by Dr. David Herndon has greatly broadened the scope of physiological and nutritional studies which can be accomplished. Branch members participate actively in teaching activities of this unit, the Brooke Army Medical Center, and are affiliated on the staff of the Medical School of the University of Texas at San Antonio. In addition, Branch members have actively participated in local, national, and international meetings to present and discuss their research findings and increase the scientific interchange in these areas of study.

PUBLICATIONS AND/OR PRESENTATIONS:

See report of Clinical Division, USAISR

### TERMINATION REPORT

# **PROJECT NO.** 3A161102B71R-01, RESEARCH IN BLOMEDICAL SCIENCE-SURGERY **REPORT** TITLE: ANESTHESIOLOGY

US ARMY INSTITUTE OF SUKGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 January 1975 - 31 December 1975

Investigators:

Gary W. Welch, MD, Ph.D, Lieutenant Colonel, MC James K. Sims, MD, Major, MC Allister K. Morris, MD, Major, MC

Reports Control Symbol MEDDH-288(R1)

Unclassified

### ABSTRACT

PROJECT NO. 3A161102B71R-01, RESEARCH IN BLOMEDICAL SCIENCE-SURGERY REPORT TITLE: ANESTHESIOLOGY

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 January 1975 - 31 December 1975

Investigators: Gary W. Welch, MD, Ph.D., Lieutenant Colonel, MC James K. Sims, MD, Major, MC Allister K. Morris, MD, Major, MC

Reports Control Symbol MEDDH-288(R1)

In the period covered in this report, 490 anesthetics were administered to 142 patients, an average of 3.45 anesthetics per patient. Although most patients received only two or three anesthetics, several received over 10 anesthetics.

The most commonly used anesthetic was ketamine followed by nitrous oxide and ethrane. Regional anesthetics were administered to only a small per cent.

Three cases of difficult anesthetic management are presented.

Anesthesia

### ANESTHESIOLOGY

### PREOPERATIVE EVALUATION

With the exception of patients with electrical injury, most burn patients scheduled for surgery are several days to weeks postinjury. In the immediate postburn period, patients have the determination of electrolytes, arterial blood gases, hematocrit, and other hematologic parameters determined as indicated. Blood pressure, central venous pressure and urine output are recorded on an hourly basis. During the past year, adult patients with a thermal injury greater than 40% have had the Swan-Ganz thermal dilution flow directed pulmonary artery catheter inserted, allowing frequent determination of the cardiac index. Daily chest x-rays are also obtained. Hence, all patients coming to the operating room have abundant physiologic data available upon which anesthetic decisions may be made. In addition, a preoperative electrocardiogram is obtained in all electrical injuries to determine the possible presence of myocardial injury.

### PREOPERATIVE PREPARATION

All adult patients are kept NPO past midnight. Exceptions to this are patients with a tracheostomy or nasotracheal tube in place who may be receiving tube feeding. Because the airway is protected in these patients, nutritional support may be continued up to the time of surgery depending upon the age and size of the patient.

### PREOPERATIVE ASSESSMENT AND FLUIDS

Premedication varies from none to heavy doses of narcotic. The usual premedication is a combination of either demerol or morphine and glycopyrolate. In the past year, intramuscular administration of diazepam has been replaced by oral administration. Two factors account for this: 1) it is a painful injection; 2) blood levels are nearly the same for both routes of administration. Pediatric cases may be given a small dose of ketamine prior to coming to the operating room. All fluids being given to the patient are changed to  $D_5RL$  on arrival in the operating room.

### TYPES OF ANESTHESIA

### A. General Anesthesia

1. Ketamine

Ketamine, a phencyclidine derivative, produces a dissociative state upon intravenous administration. It has two major advantages. The first is maintenance of ventilation during the anesthetic state, and the second advantage is the lack of depression of the cardiovascular system. Hence, patients with hypotension or compromised cardiovascular system can be induced relatively safely with ketamine. Although the airway is usually protected, obstruction or regurgitation and aspiration may occur. Ketamine is particularly useful in procedures around the face or for difficult surgical positions. Thirtytwo per cent of anesthetics given over the past year were ketamine.

### 2. Nitrous oxide with barbiturate or narcotic supplementation

Used approximately 28% of the time, this technique allows for rapid emergence by reversal of the narcotic at the end of the procedure. Patients to be intubated while awake or ventilated postoperatively are often anesthetized using a narcotic for sedative effect. Most frequently, this technique is used in conjunction with a nondepolarizing muscle relaxant. In the past year, pancuronium has been used exclusively because of its rapid onset and lack of effect on blood pressure. One possible side effect of pancuronium is a tachycardia following administration of intubating doses (see Complications).

### 3. Ha'othane with or without nitrous oxide in oxygen

Although previous research in this unit has shown halothane to be a safe anesthetic in the thermally injured patient, the use of halothane has decreased over the past year. The main advantage of halothane over other inhalational agents is its ability to provide a rapid induction with adequate relaxation of the vocal cords, allowing easy intubation. During the past year, we evaluated serum bromide levels in one patient requiring multiple halothane anesthetics. Serum bromide levels rose after every anesthetic, an indication of increased halothane biotransformation. In the past year, 8% of the anesthetics given were halothane.

### 4. Enflurane (Ethrane^R)

Enflurane has been available for approximately the past three years. It is a nonflammable fluoronated hydrocarbon which is metabolically stable, less than 2% being biodegraded. Although allowing rapid induction, it is not as smooth an agent as halothane and often relaxation with pancuronium is needed for intubation. To this date, no substantial cases of hepatotoxicity or nephrotoxicity have been reported. In 1975, 27% of the anesthetics given consisted of enflurane.

### 5. Subanesthetic ketamine

In the past year, the use of low-dose ketamine has been sporadic. For the most part, debridements have been done under general anesthesia in the operating room. The use of subanesthetic ketamine is not without potential problem because at least two patients have experienced alrway difficulty during their procedures in the Hubbard tank.

### B. REGIONAL ANESTHESIA

Any patient with a localized lesion who is not septic and is not burned at the site of injection is a candidate for regional anesthesia. That few patients meet such criteria is evidenced by a 2% use of regional for our anesthetics.

### MONITORING TECHNIQUES

A. CIRCULATION

1. Precordial and/or esophageal stethoscope.

2. Peripheral monitoring of pulse.

3. Blood pressure by either cuff, infrasonde^R, or direct intraarterial line.

- 4. Central venous pressure.
- 5. Swan-Ganz catheter.
- 6. EKG.
- 7. Sponge weight,
- 8. Urine output.
- B. RESPIRATION
  - 1. Rate.
  - 2. Auscultation.
  - 3. Blood gases as indicated.

### C. TEMPERATURE

Prolonged cases and pediatric cases have their temperature monitored during the procedure. Room temperature is maintained at 75-80 degrees Fahrenheit, and the K-thermia heating blanket is reserved either for infants weighing less than 10 kg or to cool febrile patients. Heat lamps are also utilized to warm the patients when needed.

### COMPLICATIONS

Case No. 1

### Tachycardia following pancuronium administration

This 23 year old patient sustained a 451 thermal injury following a leep accident. The areas of injury included the head, buttocks, arms, hands, thighs, and legs. Eight days following injury. he was taken to the operating room for excision of the eschar of his right leg. He was induced with pentothal and given 8 mg pancuronium for intubation. Shortly after intubation, the patient's heart rate was noted to be 200 beats per minute. The operating room monitor did not demonstrate P waves, and, consequently, a Hewlett-Packard EKG machine was brought to the operating room in an attempt to make a diagnosis. It was felt at this time that PAT was the probable diagnosis, and attempts were made to obtain a normal rhythm by carotid massage. This was unsuccessful. Because pancuronium is occasionally associated with tachycardias, it was decided to attempt conversion by reversing the muscle relaxant with edriphonium. Ten milligrams of edriphonium were given with reversion to a normal sinus tachycardia with a rate of 110 beats per minute. Plans for surgery were cancelled, and the patient returned to Ward 14A. During the next several days, the patient continued to have episodes of PAT and eventually he was placed on digoxin. propranolol, and quinidine. This resulted in control of his cardiac rhythm. These medications were continued until discharge, at which time the quinidine and propranolol were discontinued.

### Comment

Pancuronium is a nondepolarizing muscle relaxant with a steroidal nucleus. It is rapid in onset, although not as rapid as succinylcholine, with a duration of about 20 minutes. It has been noted to have anticholinergic properties and tachyarrhythmias have been associated with its use. Although the arrhythmia recurred and required propranolol, quinidine, and digoxin for eventual control, pancuronium appears to have been instrumental in initiating PAT in this patient. Subsequent surgical procedures were uneventful.

### Case No. 2

### Hypotension secondary to bowel necrosis

This patient was a 21 year old male who suffered a 30% thermal injury following an automobile accident. At the time of his arrival in this unit, he had had a left below-knee amputation, the neurovascular bundle in his left axilla was exposed, and he had subcutaneous crepitation. His initial therapy progressed well, but then he developed a myocardiopathy and congestive failure. On the 119th hospital day, the patient suddenly became cyanotic, hypotensive, and acidotic. Abdominal x-rays were consistent with visceral perforation, and the patient was taken to the operating room for exploratory laparotomy. The patient was induced and maintained with ketamine. His blood pressure required continuous support with a dopamine infusion. No urine output was obtained during surgery. At the end of the procedure, the patient was returned to Ward 14A, where he expired shortly thereafter. The operative procedure consisted of right hemicolectomy, terminal ileoectomy, transverse colostomy, and terminal ileostomy. These were done to resect necrotic large bowel.

### Comment

Most patients would tolerate a surgical procedure of this magnitude better than this patient did. The patient's underlying myocardiopathy, however, made adequate support of the cardiovascular system nearly impossible.

### Case No. 3

### Intraop rative myocardial infarction

This patient, a 58 year old male, sustained a 49.5% chemical injury when he fell into a pool of alkaline solution. Three days postinjury, he was scheduled for excision. During his preoperative interview, he admitted to occasional chest pain for which he took no therapy. He claimed he was otherwise healthy and able to carry out his normal activities. For the two days prior to his surgery, he had had cardiac arrhythmias, and he was placed on quinidine and digoxin.

On the day of surgery, the patient was taken to the operating room. He was induced with ketamine and halothane and intubated. For the first hour, the patient's blood pressure remained at 100/50. At that point, it was necessary to remove the blood pressure cuff to debride the arm. Upon reapplication of the cuff, the patient's blood pressure was noted to be 80/40. Throughout the case, the EKG remained stable and CVP varied from 7 to 9 cm of water. Estimated blood lcss was 4000 cc, and the patient received 7 units of blood and 2700 cc  $D_{cRL}$ . Upon cessation of the anesthetic, his blood pressure fell to 70 systolic, where it remained on his return to the intensive care unit. A STAT EKG revealed an acute myocardial infarction. Initially, he required pressors for maintenance of his blood pressure, but he was successfully weaned from dopamine. Two days later, it was necessary to return to the operating room for excision of skin from the patient's back and amputation of his left leg. During the procedure, he lost approximately 3500 ml of blood, and he received 6 units of whole blood and one unit of packed cells in addition to 1400 ml  $D_{c}RL$ . Urine output and blood pressure were satisfactory throughout surgery. Postoperatively, the patient remained stable until the 5th postoperative

day, when he developed ventricular instability. Treatment with lidocaine was effective; however, the patient became unresponsive and died that day.

### Comment

At autopsy, this patient had severe coronary artery thrombosis and myocardial fibrosis, probably from previous ischemic episodes. Although the preoperative EKG was normal, the patient's history of chest pain would appear to indicate previous ischemic episodes.

The episodes of arrhythmia the patient manifested prior to surgery may have been secondary to myocardial ischemia.

This patient illustrates the risk of anesthesia and surgery in a patient with an acute myocardial infarct. St 'ies have shown a greater than 50% reinfarction rate, 50% of which are facal, when a surgical procedure is performed within six months of an infarct. Although aware of this, the nature '' the patient's injury required repeated surgical procedures and nothing was available to obviate the risk.

	No. of Patients	No. Patients Anesthetized (ISR Only)	No. Patients Anesthetized No. Patients (x100)	Total Anesthetics (ISR Only)	Anesthetics No. Patients Anesthetized (x100)	Average Per Cent Burn
1966	311	181	58.2	713	3.94	30
1967	389	ż39	61.4	670	2.80	28
1968	389	259	66.6	794	3.07	30
1969	294	189	64.3	601	3.18	36
1970	321	198	61.7	497	2.51	30
1971	301	179	59.5	475	2.65	31
1972	301	183	60.8	575	3.14	34
1973	273	141	51.6	377	2.67	38.5
1974	226	123	54.4	380	3.09	41.57
1975	254	142	55.9	<b>49</b> 0	3.45	42.1

# TABLE 1. OVERALL PATIENT DATA, USAISR (1966-1975)

Procedure	1971	1972	1973	1974	1975
Debridement and/or homograft	15.5	19.7	21:5	22.60	25.0
Âutograft	52.9	51.3	52.6	56.9	51.0
Orthopedics	13.0	8.9	8.0	8.1	8.0
Ear (chondrectomy)	4.0	3.1	2.6	1.60	1.0
Eye and lid	3.8	0.7	1.8	1.60	2.0
Intra-abdominal	1.7	7.8	2.1	3.70	2.0
Tracheostomy & bronchoscopy	4.6	6.6	6.6	1.80	∢۱
Other	4.4	1.9	4.8	3.70	11.0

2

## TABLE 2. NATURE OF SURGERY, USAISR (Per Cent)

RESEARCH AND TECHNOLOGY WORK UNIT SUMMARY					CV ACCESSI DA 6956	<b>641®</b> 5	2. DATE OF SUI 76 10	01	DD-DR	AE(AR)636
& DATE PREV SUM'RY	4. KIND OF SUMMARY	S SUMMARY SCTY	A. WORK SECURITY	7 REGR	ADING	SA DI	EB'H INSTR'N	BA SPECIFIC	DATA -	LEVEL OF SUM
75 07 01	H. TERM	U	U	N/	1	1	NL			A. WORK UNIT
10 NO /CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK /		DER		WORK UNIT	NUMBER	
	61102A	3A161102	B71R		01		141			
S. CONTRIBUTING							•	·····;·····	~~~~~	. <u> </u>
C. CONTRIBUTING		,		1				· · · ·	12	• • • • • • • • • • • • • • • • • • •
11 TITLE (Procode with	Security Classification Code	(U) Clini	cal Operat	ion,	Metabo	olid	c Branch	, Renal	Secti	on, for
Treatmont o	f Soldiere Wi	th Renal Fa	ilure (44)					•		
12 SCIENTIFIC AND TE	CHHOLOGICAL AREAS	un numar re								
003500 Clin	ical Medicine									
13 START DATE		14. ESTIMATED CON	PLETION DATE		HIG AGENC	Y		IS. PERFORM		OD
52 07			75 12	EG	ì		1	C. In-	House	
JZ OT	Not Applicab	10			L				1.	
	NOL Appircan	10		10 RES	IPRECEDINI	INATI	C A PROPESS	IONAL MAN YR	E E PUN	
A DATES/EFFECTIVE		EXPINATION.		1.	76		1 1.0		28	
P HONDER		•		FISCAL	CUNNERT					
& TYPE:		4 ANOUNT		1						
& KINC OF AWARC.		I.CUM. AMT.		L						
I. RESPONSIBLE DOD	DRGANIZATION			20 PIR	FORMING OR	GANIZ	MOITA			
NAME:" US Army	Institute of	Surgical	Research		US Arm	уI	nstitute	e of Sur	gical	Research
-				i i	Meta	bol	ic Branc	ch		
ADDRESS.* Fort	Sam Houston,	Texas 782	34	ADDRES	• Fort	Sa	m Housto	on, Texa	s 782	234
				1						
				PRINCIP	AL INVEST	-	R (Fumish 36 AN	il U S. Asadomic	Inclination	
RESPONSIBLE INDIVIDU				NAME	• Will	iam	D. Myer	cs, DTC,	MC	
HANT Basil A	. Pruitt. Jr.	COL, MC		TELEP	HONE 5	12-	221-6532	2		
512	-221-2720	,		SOCIAL	SECURITY	ACCO	UNT NUMBER			
TELEPHONE. JIL							n4			
DODDIGN THE		CONSTREES	Л		Rich	ard	H. Meri	cill. L1	C. MC	
FOREIGN INT	EDDIGENCE NOI	CONSIDERE	0		100.011				••	DA
T VIVIO and /Procedo	RACH with Somelly Classifi	cation Code)		HANE						
IL KETTORG (FROM BACH VIN JONNY CHANNEL (VV)										
(U) Hemodialysis; (U) Peritoneal dialysis; (U) Renal laliure; (U) Soldiers										
23 TECHNICAL OBJECT	IVE, T 24 APPROACH, 28	PROGRESS (Fumine i	nelviousi persgraphe in manuta and au		tation	. fo	r + ho + l	hormally	เม่นสะเดียง เป็นการ์เป็น	red
23. (U) TO	provide diag	mostic sup	port and co	JIISUI	Lation	-1.	de beth	noritor	ه [جم	nd
soldier and	l to initiate	treatment	as is nece	ssary	10 10	CTU	de both	perruo.	lear a	
hemodialysi	.s. In additi	on, the re	nal section	n 15	TUADTA	ed	in clin:	LCAL TES	earch	h ]
activities	designed to i	improve pat	ient care	and o	ur und	lers	tanding	or rena	u pro	DIEMS
in the thermally injured soldier. The renal section also supports the clinical and										

research endeavors of the Nephrology Service at Brooke Army Medical Center in an effort to improve care of military personnel. 24. (U) The renal section provides consultation to the physicians involved in direct care of the thermally injured patient. Anal function is assessed by various techniques and therapeutic interventions in the form of hemodialysis and peritoneal dialysis are

and therapeutic interventions in the for. of hemodialysis and peritoneal dialysis are available. In addition, the renal section is involved in several clinical and laboratory research protocols.

25. (U) 75 01 - 75 12 Ten patients were hemodialyzed by the ISR Renal Section for a total of 59 patient treatments. One peritoneal dialysis was carried out. Of the 10 patients, only one survived. Femoral vein catheterization with unipuncture hemodialysis has markedly improved acute hemodialysis access. A video tape has been made of the technique and presented at the American Society of Artificial Internal Organs Meeting. Other areas of clinical investigation are also underway, with particular emphasis on pathophysiology of acute renal failure. The second course in Nephrology was sponsored by ISR Nephrologists and awarded credit by the American Medical Association for continuing education.

### TERMINATION

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

### **REPORT TITLE:** CLINICAL OPERATION, METABOLIC BRANCH, RENAL SECTION, FOR TREATMENT OF SOLDIERS WITH RENAL FAILURE

### US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 January 1975 - 31 December 1975

### Investigators:

William D. Myers, MD, Lieutenant Colonel, MC Richard H. Merrill, MD, Lieutenant Colonel, MC

Reports Control Symbol MEDDH-188(R)

UNCLASSIFIED

### ABSTRACT

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

### REPORT TITLE: CLINICAL CPERATION, METABOLIC BRANCH, RENAL SECTION, FOR TREATMENT OF SOLDIERS WITH RENAL FAILURE

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 January 1975 - 31 December 1975

Investigators: William D. Myers, MD, Lieutenant Colonel, MC Richard H. Merrill, MD, Lieutenant Colonel, MC

Reports Control Symbol MEDDH-288 (R1)

The Renal Service performed 59 hemodialysis treatments, 567 urinalysis, 379 fibrinogen degradation product determinations, 13 creatinine clearance, and 9 glofil clearances. The glomerular histology protocol was extended to include scanning electronmicroscopy; a protocol was initiated to develop a noninvasive technique of determining intrarenal blood distribution; and another protocol was initiated to determine the site of infection within the urinary tract. The first annual Brooke Army Medical Center Nephrology Seminar was also held.

Renal Failure Hemodialysis Peritoneal Dialysis Soldiers

### CLINICAL OPERATION, METABOLIC BRANCH, RENAL SECTION FOR TREATMENT OF SOLDIERS WITH RENAL FAILURE

The Renal Section is composed of a Nephrologist, and two hemodialysis technicians who also function as research technicians. The primary mission of the Renal Section is to support the operation of the Clinical Division of the Institute of Surgical Research Burn Unit, providing both consultation for patients with renal and metabolic problems and hemodialysis in cases of renal failure. A secondary mission of the unit has been to support the Nephrology Service of Brooke Army Medical Center. The USAISR Hemodialysis Team provides backup support when necessary and assists in treatment of cases of renal failure occurring at the main hospital of Brooke Army Medical Center. The Chief of the Metabolic Branch directs the BAMC Nephrology Service, and the Chief of the Renal Section directly participates in Nephrology patient care at BAMC.

Ten patients were hemodialyzed for a total of 59 dialyses. One patient was peritoneally dialyzed. Of the 11 patients dialyzed only one survived. Ten patients died of burn wound sepsis. In addition to dialysis, renal technicians performed 567 urinalysis, 379 determinations of fibrinogen degradation products, 13 creatinine clearances, 9 glofil clearances, plus assisted the burn unit in bedside care.

The thrust of research by the Renal Section has been toward definition of the type of renal insufficiency seen in the thermally injured patient. Histologic studes including transmission electron microscopy have failed to support earlier suggestions that a glomerular alteration may be responsible for the altered renal function. This study has been extended to include scanning electron microscopy since current studies of other investigators have suggested a role of glomerular surface alteration in acute renal failure.

Efforts are also being made to better define the renal hemodynamics in the thermally injured patient. Currently a noninvasive technique for determining intrarenal blood distribution is being investigated in a rat model using dimercaptosuccinic acid labled with  $T_c$  and microspheres labeled with Sr

Using fluorescene labeled immunoglobulin an investigation is being made into the role of retrograde urinary tract infection in renal parenchymal infection and systemic sepsis in the thermally injured patient. It would be particularly helpful if this technique could be used to localize sites of yeast infection since bladder irrigation would correct yeast infections limited to the lower tract without exposing patients to the toxicity of systemic therapy. USAISR Nephrologists initiated the first Brooke Army Medical Center Nephrology Seminar, a three day event devoted to kidney transplantation held and videotaped at the Academy of Health Sciences. Speakers included Transplant Surgeons and Nephrologists of national status. The seminar was well received and future seminars are to be sanctioned by the American Medical Association and the American College of Physicians for continuing education. Chronic Renal Failure will be the topic of next years seminar.

### PRESENTATIONS:

Merrill, RH: <u>History of Transplantation</u> First Annual Brooke Army Medical Center Nephrology Seminar 29-31 January 1975.

Myers, WD: Long Term Prognosis First Annual Brooke Army Medical Center Nephrology Seminar 29-31 January 1975.

Myers, WD: Principles of Acid-base Physiology ICU Advanced Nursing Course 19 Feb 75.

Myers WD: Flank Pain, Hematuria, and Renal Mass University of Texas Health Science Center at San Antonio combined Nephrology Conference 28 May 75

Myers WD: Principles of Acid-base Physiology ICU Advanced Nursing Course Aug 1975.

Merrill WH: Iron Absorption in Hemodialyzed Patients, Southeastern Dialysis and Transplantation Association Meeting, Dallas, Texas, August 1975.

Myers, WD: All About Kidney Transplantation ICU Advanced Nursing Course Seminar on Transplantation. October 1975

Myers WD: <u>Treatment of Glomerulonephritis</u> Brooke Army Medical Center-Selectéd Topics in Internal Medicine Lecture Series December 1975.

				1. AGENC	Y ACCESSION	2. DATE OF SU	MARY [®]	EPORT (	OWTROL STRENDL
RESEARCH	AND TECHNOLOG	Y WORK UNIT S	UMMARY	DA O	D 6951	76 10 0	1	DD-DI	R&E(AR)626
L DATE PREV SUMPRY	4. KIND OF SUMMARY	L SUMMARY SCTY	A. WORK SECURITY	7. REGRAS		ISS'N INSTR'N	SA SPECIFIC D	ATA	LEVEL OF SUN
75 07 01	H. TERM	υ	U	NA		NL		HO	A. WORK UNIT
18. NO./CODES: ⁸	PROGRAM ELEMENT	PROJECT	NUMBER	TASK AP	REA NUMBER		WORK UNIT	NUMBER	
	61102A	3A161102	<b>371</b> R	01		084			
L. CONTRIBUTING									
CONTRIBUTING								h	
(U) Detection	on of Endotox	,• in in Burne	ed Soldiers	With	Sepsis	(44)			
18. SCIENTIFIC AND TEC	CHHOLOGICAL AREAS						**************************************		···-
003500 Clin:	ical Medicine								
IS. START DATE		14. ESTIMATED COM	LETION DATE	IS. FUNDI	NG AGENCY	يى بور عالية الخرجيرينات	14. PERPORMAN	CE NETI	+0.0
71 03		76 09		DA	1	1	C. In-H	ouse	
IT. CONTRACT/ARANT	Not Applic	able		-	ACES ESTIMAT			A FUN	DE (In those ands)
& DATES/EFFECTIVE		EXPIRATION:		T	76		.5	1	7
N. NUNGERI [®]				PISCAL	7т		.2		5
& TYPE		4 ANOUNTI		YEAR C	UNNERY				
& KI D.OF AWARD		I.CUM. AMY.							
In. AREPONDIALE GOD O	REAVISATION			30. PERFO	RMING ORGANI	TATION			
AMMES US Army	Institute of	Surgical H	Research	HAME:*US	5 Army I	institute	of Surg	ical	Research
_		-			Clinica	l Divisi	on		
ABONESS!" Fort	Sam Houston,	Texas 7823	34	ADDRESS	Fort S	am Houst	on, Texa	s 78	3234
				PRINCIPAL	INVESTIGATO	R (Fumish BEAN (	I U.S. Asedenic je	a ti fu ti anj	
RESPONSIBLE INDIVIDU	AL			HANEI	Robert	B. Lindb	erg, PhD		
NAME Basil I	A. Pruitt, Jr	, COL, MC		TELEPHO	ms: 512-	221-2018			
TELEPHONE: 512-	-221-2720			SOCIAL S	ECURITY ACCO				
EI. GENERAL USE				-		N1			
FOREIGN INT	EILIGENCE NOT	CONSIDERED	)	NAME: V	/irginia	C. Engl	ish, MS		
				NAME: I	Basil A.	Pruitt,	Jr, COL	, MC	DA

II, KEY CALL (Placedo ZACE with Security Classification Code)

(U) Endot'Oxin; (U) Sepsis; (U) Assay; (U) Burns; (U) Humans E. TECHNICAL OBJECTIVE,* 34. APPROACH. 35. PROGREMS (Purilies Individual paragraphic identified by number Procedu inst of each with Society Clearification Code )

23. (U) Detect endotoxin in tissues of burn patients dying with sepsis; special interest directed toward liver, in view of current extensive findings of liver damage as part of pathological picture in septic shock, a factor in battle wounds of all types.

24. (U) Liver and other tissues are triturated and extracted with glacial acetic acid to neutralize inhibition and permit detection of endotoxin.

25. (U) 75 07 - 76 09 65 autopsies of patients dying with extensive burn injuries yielded liver-tissue for examination. 61 of these showed endotoxin by Limulus Amaebocyte Lysate test. 93% of patients had detectable endotoxin in the liver; median value was .01 ug/gm. Normal livers failed to reveal endotoxin and bacterial content of reacting livers was 10³ or less. This level will not produce lysate coagulation.

A valiable to contractors upon originator's approval

PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DO FORMS 1498A 1 NOV 65 AND 1498-1 1 MAR 68 (FOR ARMY USE) ASE OBSOLETE

### TERMINATION

**PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE - SURGERY REPORT TITLE: DETECTION OF ENDOTOXIN IN BURNED SOLDIERS WITH SEPSIS** 

And × ひょうひょうか

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 June 1976

Investigators:

Robert B. Lindberg, PhD Virginia C. English, MA Arthur D. Meson, Jr, MD Basil A. Pruitt, Jr, MD,Colonel,MC

Report Control Symbol MEDDH-288(R1)

UNCLASSIFIED

### ABSTRACT

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE - SURGERY

REPORT TITLE: DETECTION OF ENDOTOXIN IN BURNED SOLDIERS WITH SEPSIS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 June 1976

Investigators: Robert B. Lindberg, PhD Virginia C. English, MA Arthur D. Mason, Jr, MD Basil A. Pruitt, Jr, MD, Colonel, MC

Report Control Symbol MEDDH-288(R1)

Endotoxin can be demonstrated in plasma or serum by the Limulus ampebocyte lysate reaction, after extraction with acetic acid and neutralization of the supernate. A derivative of this procedure was applied to homogenates of liver from patients dying with severe burns, in most • instances with clinical and laboratory indication of severe sepsis. 94% of 65 patients had a positive reaction for endotoxin in the liver. The median concentration was .044 mcg/gm of liver. Seven patients, or 11.4% of reactors, had no positive blood cultures. The sequestration is, in at least 50% of cases, not associated with bacterial contamination of the specimen; the remainder of reactors may include some in whom the bacterial contribution to endotoxin level occurred.

Endotoxin Sepsis Assay Burns Humans

### DETECTION OF ENDOTOXIN IN BURNED SOLDIERS WITH SEPSIS

Detection of nanogram or lesser amounts of endotcxin (ET) is now feasible with the Limulus amoebocyte lysate reaction, and the availability of this technic has resulted in renewed interest in assessing the pathogenesis of gram-negative shock. Earlier reports have described the failure to detect circulating endotoxin the the blood of burned patients with sepsis (1). Those efforts led to the conclusion that circulating ET would be detected in approximately one third of the patients in whom gram-negative sepsis was plausibly established. The procedure thus is not a valid diagnostic aid in the diagnosis of sepsis, but other problems regarding ET and its mode of action can be approached by the LAL reaction. The study of liver tissue in patients dying with endotoxic shock was a problem to which LAL technic could be applied, and the results of this effort have been set down in this on-going study.

Since the endotoxin being detected with the LAL is a lipopolysaccharide embodied in the bacterial cell wall, bacteria in tissue could contribute a quantity of endotoxin directly, without representing any function of selective absorption of circulating endotoxin by hepatocytes, macrophages, or any other cell. This possibility was tested in several ways. One thousand bacterial cells in one ml of a broth culture will give a positive LAL reaction in some instances - i.e, when the cells are gram-negative bacilli or any other form rich in endotckin. However, tests of liver tissues with varying numbers of Enterobacteriaceae, including Klebsiella, Enterobacter and Serratia spp have shown a wide variation in the minimum number of bacteria that will give a positive gelation with Limulus lysate. Positive reactions were obtained with a minimum of 5 x  $10^{3}$  cells per gram of liver tissue, but the median reactive level with 20 liver samples was  $4.5 \times 10^{2}$ . In cultures, the reaction at 1000 cells per ml is the equivalent of .ool  $\mu g$  or even .0001  $\mu g$  in standard endotoxin. Such a minimal reaction thus represents a lower cut-off point for interpreting the results of tests on clinical material: i.e., tissues with low bacterial counts and with ET levels at 10 to 1000 times the maximum that such bacterial level could produce may be interpreted as containing endotoxin beyond a level to be accounted for by bacterial content.

Endotoxin level was determined in 65 liver samples. Of these patients, 49 had at least one positive blood culture, and 16 had negative blood cultures or did not have a blood culture drawn. Results of ET determinations are shown in Table 1. There were 61 samples in which a detectable level of ET was found. Among these, 31, or 50% gave levels of .01  $\mu$ g/gm or higher. This meant a larger proportion than had previously been seen contained

^{1.} Lindberg RB, English VC, Pruitt BA, Jr, Mason AD, Jr: Detection of endotoxin in burned soldiers with sepsis. USA Institute of Surg Res Ann Res Prog Rpt FY 1973, BAMC, Ft. Sam Houston, Texas, Section 6.

amounts of endotoxin which bordered on the levels which might reflect bacterial contamination rather than sequestered ET. However, the bacterial counts, shown in the third column range from high to low over the whole range of endotoxin levels observed. This supports the view that the bacterial count was not contributing consistently to the ET level demonstrated in these liver samples.

Endotoxin ug/gm	No. Patients	Average Bacterial Count/gm
5.12*	]	10 ⁶ •7
•64	1	10 ^{3.0}
•32	3	10 ^{2.8}
•16	2	10 ^{3.6}
•08	4	10 ³ •1
•04	4	102.8
•02	8	10 ³ • ⁴
•01	8	10 ^{2•9}
•008	1	10 ^{3.0}
•005	11	10 ^{2•9}
.0025	14	10 ^{3.1}
•001	4	10 ^{2•3}
≪.0001	4	102.8
Total tested	65	
Median level of ET:	.044 µg∕gm	

Table 1.	Endotoxin Levels	in Liver	Tissues	From
	Burn Autopsies,	ISR, 1975	;	

*This level was technically correct, but there is a possibility that direct fecal contamination of the sample occurred. It was omitted in calculating the median ET level. Endotoxin accumulated in livers of several patients who were clinically septic but in whom no positive blood cultures were collected. Table 2 presents data on patients in whom endotoxin was demonstrable and in whom all blood cultures were negative. There were seven patients in this category. Four had gram-negative bacteria in the liver. Multiple species were found in two, <u>Enterobacter aerogenes</u> in a third and <u>Escherichia coli</u> in a fourth. Three other liver samples were all negative for bacterial growth. Evidently the accumulation of endotoxin can occur in the absence of bacteria in the liver.

There were 4 patients in whom no detectable endotoxin could be found in the liver. One of these had no blood cultures collected. The other 3 had from 4 to 10 positive blood cultures. Multiple species were recovered from these samples. The results are shown in Table 3. The succession of species in blood cultures was not uncommon in a lethal sepsis. The blood stream strains were not always recovered from the liver in situations where sepsis had prevailed for a few days. Endotoxin was not detectable in any of these liver samples.

In autopsy liver sampels positive for endotoxin, 12 species of bacteria were recovered. Predominant were <u>Klebsiella pneumoniae</u>, <u>E. coli</u>, <u>Enterb-</u> <u>bacter clcacae</u> and <u>Pseudomonas aeruginosa</u>. There was no indication that any species was associated with more ET in liver tissue than any other gram negative bacillus.

### STABILITY OF LIMULUS AMOEBOCYTE LYSATE

A large pool of Limulus amoebocyte lysate was collected in 1971, and has been used since that time. The lysate, in 20 ml amounts, was forzen and stored at  $-70^{\circ}$ C. Since the usefulness of ET assay depends on active lysate, periodic tests of potency of this lysate, have been carried out. It is of general interest to determine the duration of potency under these storage conditions. Results of a series of texts on samples from a pool of lysate designated &-2a no.1 are summarized in Table 4.

In a series of 17 tests during 1975, variations in sensitivity occurred. A reading at 0.001 mcg/ml could be considered valid when the intensity of clot formed was rated at 2, which was a less than stable clot. However, only 4 out of 10 texts were read at this level. Ten tests gave 3 or 4 intensity reactions with a succession of endotoxin standards. Endotoxin standards themselves may vary; in the comparison series reported here; one control, in November 1975 gave the weakest results. That run gave a 2 reaction at 0.0025  $\mu$ g/ml; at 0.00125 mcg. of ET of this preparation, the reaction was only at a weak clot level, grade 1 in this instance, the vial of lysate was regarded as sub-standard in potency, although other ET preparations reacted with stronger clots in this particular vial of lysate. The over-all result was affirmation of the survival of amoebocyte lysate activity for at least 4 years in storage at  $-70^{\circ}$ C.

<b>54</b>	41	3	8	21	و	6	Autopsy
4	-	10	12	2	_	-	No. of Blood Cultures
•01	•02	.0025	•006	•04	•02	•08	Endotoxin Level mcg/gm
negative	5 × 10 ⁴	negative	negative	1.2 × 10 ⁴	$1 \times 10^2$	2 × 10 ⁴	Liver Bacterial Count per gm
no growth	E colt	no growth	no growth	Staph. aureus, E aerogenes, Strep GpD	Prov stuartii, X pneumoniae, Ps aeruginosa, Ps maltophilia	E coli, K pneumoniae, Ps aeruginosa, Prov stuartii, Pseudo maltophilia, Strep GpD	Becteria recovered in liver

# Table 2. Endotoxin in Livers of Patients with Negative Blood Cultures, 1975

### Table 3

# BLOOD CULTURE SURVEY OF PATIENTS WHO YIELDED NO DETECTABLE ENDOTOXIN FROM AUTOPSY LIVER

AUTOPSY	NO. NEG.AM.	NUMBER OF BLOOD CULTURES AM	ORGANISMS FOUND
NUMBER	BLD.CULTURES	YIELDING SPECIFIED ORGANISM	IN LIVER

     A=16=75	12	10 Positive bld. cultures giving rise to:	COUNT: 2 x 10 ²
A-10-75	13	4 E. cloacae	E. aerogenes
		4 r. aeruginosa 3 K. pneumoniae	Nonnemolytic Gp D Streptococcus sp.
			K. pneumoniae

A-47-75	0	0	not examined

		10 Positive bld. culture giving rise to:	5
A-19-75	5	4 E. cloacae 2 P. aeruginosa	$1.1 \times 10^3$
		5 K. pneumoniae 1 S. aureus	K. pneumoniae S. aureus

4 		4 Positive bld. cultures giving rise to:	
A-11-75	6	3 E. cloacae 1 K. pneumoniae	5.1 x 10' P. aeruginosa K. pneumoniae

Abbr: No. = number; AM = antemorteum; NEG.= negative; bld - blood

### Table 4 THE INTEGRITY OF LIMULUS LYSATE STOCK

### LYSATE U-2# # 1

DATE	DEGRE	E OF RE	ACTION /	T ET CONC	ENTRATIO	N IN MCG.		
	0.01	0.005	0.0025	0.00125	0.0006	0.0003		
1-14-75	3	3	2	· 2	1	1		
Fresh endotoxin dilution from stock prepared for following tests:								
3-11-75	4	2	3	2	ŀ	0		
Fresh endotoxi	n dílı	ution fr	om stocl	k prepared	for fol	lowing tests:		
7-22-75	4	3	3	3	3	2		
7-23-75	4	3	3	3	3	2		
7-29-75	4	3	3	3	2	1		
Fresh endotoxi	n diti	ution fr	om stoc	k prepared	l for fol	lowing tests;		
7-30-75	4	4	4	4	4	3		
7-31-75	4	4	4	4	4	3		
8-4-75	L.	4	4	4	4	3		
8-6-75	3	4	3	2	1	ī		
Fresh endotoxi	n dil	ution fr	om stoc	k prepared	l for fol	lowing tests:		
8-11-75	4	4	3	3	2	1		
8-25-75	4	4	4	4	3	3		
Fresh endotoxi	n dil	ution fr	om stoc	k prepared	for fol	lowing tests;		
9-13-75	4	4	4	4	3	1		
9-15-75	4	4	4	4	3	2		
9-29-75	4	4	4	3	3	1		
Fresh endotoxi	n dil	ution fr	om stoc	k prepared	for fol	lowing tests:		
11-10-75	<u> </u>	4	3	2	•	•		
New vial of ivsate thawed and fresh endotoxin dilution prepared								
for following tests:								
11-28-75	4	3	2	1	1	1		
1-5-76	ů.	ź	2	i	Ó	0		
		toned In	Baues	·	70 0	C though hafara		

Note: All lysate stored in Revco freezer at  $\sim 70$  C S thawed before use. 0 = No reaction; I = positive but weak, unstable clot; 2  $\approx$  slightly more stable clot than 1 +; 3  $\approx$  Tight clot, but slightly disrupted on tipping tube; 4 = clot forms in less than 1 hour; does not disrupt on tippint tube.

All lysates assayed by using E. coli Oll1:B4 lipopolysaccharide (Difco) as endotoxin source.

### · DISCUSSION

the process of the part of the second state of the second state of the second state of the second state of the

The observations summarized here add further evidence that endotoxin is indeed sequestered in the livers of burn patients dying with sepsis and gram negative infection. Although here were cases in the lower range (below .005  $\mu$ g/ml) in which the bacteria present in the sample might have contributed to the endotoxin demonstrated, such samples were few in number. 94% of the patients exhibited demonstrable endotoxin in the liver. The mean value of .044 mcg/ml was at least 4 times higher than the level that might be caused by presence of gram negative bacteria at 10⁴/gm of tissue.

The nature of the effect of endotoxin on hepatic cells remains to be elucidated. The complex derangement of essential functions that occurs with shock is such that this sequestering phenomenon may be an important factor in this aspect of burn injury.

### PRESENTATIONS

Lindberg RB, English VE, Mason AD, Jr, Pruitt BA, Jr, "Sequestration of Endotoxin (ET) in Liver of Burn Patients with Gram-megative Sepsis" presented at the Am. Assoc. Immunol., Anaheim, Calif., April 13, 1976.

### PUBLICATIONS

Lindberg RB, English VE, Mason AD, Jr, Pruit: BA, Jr: Sequestration of endotoxin in liver of burn patients with gram negative sepsis. Fed Proc 35:738, 1976 (Abst).
				-	CY ACCESS	in the	2. DATE OF SUM	MARY	REPORT	CONTAGL STREET
RESEARCH	AND TECHNOLOG	r work unit s	UMMARY	DA (	DA 6397	, ]	76 10	01	DD-D	R&//(AR)636
S. DATE PREV SUM'RY	4. KIND OF SUMMARY	S. SUMMARY SCTV	4. WORK SECURITY	7. REOR	ADING	-	ETH HETR'N	Sh SPECIFIC	DATA-	LEVEL OF SUN
75 07 01	H. TERM	ប	្រប	N	A	NI	<u>ل</u> م	CA YES		A WORK UNIT
10 NO /CODES *	PROGRAM ELEMENT	PROJECT	NUMBER	TASK	AREA NUME	ER		WORK UN	TNUMPE	h
& PRIMARY	61102A	3A161102B	71R	0	1		132			
b. CONTRIBUTING									<u>_</u>	
C. CONTRIBUTING								···		<u></u>
11. TITLE (Procode with	Security Classification Code	• •								
(U) Antibio	tic Sensitivi	ty of Curre	ent Militar	y Bu	m Pati	lent	: Flora	(44)		
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS									
003500 Clin	ical Medicine			<b>1</b>						
13 START DATE		14 ESTINATED COM	PLETION DATE	IL FUN	DING AGENC	¥		H. PLAFOR	NANCE ME	THOD
54 07		, 76	09	DA				<u>c.</u> II	n-Hous	e
17 CONTRACT/GRAMT	Not Applica	ble		18. RES	OURCES EST	IMATE	-	OFAL MAN Y	RS & PU	HDE (In Broundarda)
A DATES/EFFECTIVE:		EXPIRATION:		1	76	•	.6		2	2
N NUMBER.*				FISCAL	71		.2			8
G TYPE.		4 ANOUNT:		YEAR	CONNENT		1			
& KIND OF AWARD		LCUM. ANT.				_	1			
IS RESPONSIBLE DOD	ORGANIZATION	L		20. PER	FORMINGOR	GANIZ	ATION	Τ		
HAME!" US Army	Institute of	Surgical I	Research	HANE!	US Art	ay J	Institut	e of Su	urgica	l Research
-		-		1	Mic	- obi	iology B	ranch	-	
ADDRESS. Fort	Sam Houston,	Texas 782	234	ADDRES	• For	t s	Sam Hous	ton, Te	exas	79234
				PRINCIP	AL INVESTO	SATOR	(Fumish BEAN I	I U S. Atadam	ic (matikalia	•
RESPONSIBLE INDIVIDU	JAL			HAME	Robei	ct X	8. Lindb	erg, Ph	۱D	
HANE Basil A	. Pruitt, Jr,	COL, MC		TELE	ноне 51	122	221-2018			
TELEPHONE: 51	2-221-2720			\$0CIA	L SECURITY	Ar.co	UNT NUNDER-			
21 GENERAL USE				ABOCIA	TE INVESTI	ATOR	15			
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	2	HAME	Anthor	ny P	A. Contr	eras, N	1S	
				NAME						DA
22 KEYWORDS (Freede	BACH with Somethy Class"	(II)	Providenci	a:						

(U) Antibiotic sensitivity: (U) Pseudomonas: (U) Burns: (U) Chemotherapy;

23. (U) No single function is more important in microbiologic investigations in a burn unit than continued monitoring of infecting strains for antibiotic sensitivity. Resistance to antibiotics is often capricious--there is little predictability except by assessing the individual antibiotic. In military medicine or mass casualty management these predictions are of particular significance.

24. (U) MIC by tube dilution is applied to appropriate, selected isolates, including all septicemic strains.

25. (U) 75 07 - 76 09 Ninety-eight percent of strains tested were recovered from blood, most often from patients diagnosed as septic. Renewed methicillin resistance with staph aureus occurred after falling markedly in the preceding year. The newly cpidemic strain of <u>Klebsiella pneumoniae</u> was affected only by Colymycin, Minocin and Vibramycin. This was time for epidemic <u>Enterobacter cloacae</u> as well. <u>Pseudomonas aeruginosa</u> was sensitive to Keflin and partly sensitive to gentamycin, but sensitive to carbenicillin.

liable to contractora upon originator's approva

DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 14984 I NOV 68 AND 14"SI I MAR 68 (FOR ARMY USE) ARE OBSOLETE

# TERMINATION

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: ANTIBIOTIC SENSITIVITY OF CURRENT MILITARY BURN PATIENT FLORA

# US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1375 - 30 September 1976

Investigators:

Robert B. Lindberg, PhD Anthony A. Contreras, MS Daniel T. Zamora, SP6

Reports Control Symbol MEDDH-288(R1)

Unclassified

### ABSTRACT

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: ANTIBIOTIC SENSITIVITY OF CURRENT MILITARY BURN PATIENT FLORA

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975-30 September 1976

Investigators: Robert B. Lindberg, PhD Anthony, A. Contreras, MS Daniel T. Zamora, SP6

Reports Control Symbol MEDDH-288(R1)

A high level of septicemia in severely burned patients presented 517 strains of essentially opportunistic invaders from blood cultures for sensitivity testing in 1975. MIC showed a return of methicillin resistance in a staphylococcus population that had in the previous two years become more sensitive to methicillin and other categories of antibiotics. For these strains, keflin susceptibility increased, clindamycin remained highly active, and they were all sensitive to vancomycin. Klebsiella pneumoniae, appearing on an epidemic scale to supplant Enterobacter cloacae, was inhibited in a significant degree only by colymycin, minocin and, in a preliminary trial, by vibramycin. Results were similar for Enterobacter cloacae. These enteric opportunists were major causes of sepsis, and essentially did not respond to chemotherapy. Together with Pseudmonas aeruginosa they represent a cross-resistant treatment-refractory bacterial population which appears most often in terminal septic episodes. Extended investigation of chemotherapy is called for if they are to be reduced in importance as a cause of death in severely burned patients.

Antibiotic sensitivity Pseudomonas Burns Chemotherapy Providencia

### ANTIBIOTIC SENSITIVITY OF CURRENT MILITARY BURN PATIENT FLORA

Since bacterial sepsis has continued to be the major cause of death in burn injury, the antibiotic susceptibility of the burn flora in septicemia has remained a primary concern in the management of severe burns. The profile of sensitivity of those strains recovered in the Institute of Surgical Research from blood cultures, together with a small number of wound, sputum and urine cultures tested, furnishes a guide to current antibiotic therapy. Such treatment must frequently be instituted before the sensitivity of a given blood stream isolate could possibly be determined. A resume of sensitivity of the major offending species of bacteria enhances the potential for selecting the optimal antibiotic prior to acquisition of data on a specific strain. Bacteria found on burn populations may fluctuate rapidly in sensitivity to antibiotics so that ongoing summaries are important as guides in a valid program antibiotic therapy. The constitution of an optimal routine test battery of antibiotics was aided by sorting out long-term trends of sensitivity or resistance as seen in these summaries. Cross-resistance in several species of opportunistic pathogens found in burn patients at the Institute of Surgical Research has been far greater than that reported from other general hospitals. This difference permits more clearcut demonstrations of patterns of resistance than is the case with most organisms from hospital-acquired infections.

### **TECHNIC AND SOURCES OF STRAINS**

In this study, a Minimum Inhibitory Concentration (MIC) procedure has been used. It has been described in a previous report. 10 mm capped plastic tubes are prepared to contain 1.0 ml of antibiotic at concentrations from 1.56 ug/ml to 50 ug/ml in Trypticase Soy Broth (TSB). 1.0 ml of a 4 to 6-hour TSB culture, containing from 10[°] to 10[°] organisms per ml is added to each tube of antibiotic and incubated for 18 to 20 hours. The final concentrations of antibiotic employed extended from 0.78 ug/ml to 25.0 ug/ml. MIC was taken as the lowest concentration of antibiotic which prevented visible growth at 18 to 20 hours. When needed, MIC could be determined by sub-culturing a loopful of fluid from tubes at concentrations above the visible growth leve! to blood agar plates.

Ten species of microorganisms were tested, with a total of 527 strains (Table 1). In previous years, up to 40% of strains examined were recovered from sources other than blood cultures, but in 1975, 519 out of 527 strains (98%) were from blood cultures. Thus, the strains examined in the period covered by this report are those which, on an empirical basis, were most

1. Lindberg RB, Contreras AA, Smith HOD, Jr, Plowey EC, Mason AD, Jr: Antibiotic sensitivity of flora from military burn patients. USA Inst Surg Res Progress Rpt FY 1973, BAMC, Fort Sam Houston, Texas. Section 7.

probably virulent, since they had invaded the blood stream. The data collected are, of course, of maximum pertinence to the essential question of the control of sepsis in burned patients.

The species most frequently encountered in blood cultures were, in order of occurrence, <u>Staphylococcus aureus</u>, <u>Klebsiella pneumoniae</u>, <u>Pseudomonas aeruginosa and Enterobacter cloacae</u>. In previous reports, <u>Providencia stuartii</u> had been found to be a major cause of sepsis in burned patients at the Institute of Surgical Research. From 1970 through early 1974, it rose from being a conspicuous opportunistic invader to the predominant species numerically; however, in 1975, only one strain was recovered from a blood culture.

Species	Total No	Tested		Source	s of Strai	ns
opeeres	Patients	Strains	Blood	Lung	Wound	Other
S.aureus	43	153	152	1	0	0
S.epidermidis	6	6	6	0	0	0
Strep.spp. (alpha,						
non-hemolvtic)	7	13	13	0	0	0
Ps.aeruginosa	45	95	91	2	1	1
K.pneumoniae	58	159	156	2	0	1
Entero, cloacae	31	62	62	0	0	0
E, coli	8	9	8	1	0	0
S.marcescens	7	22	21	1	0	0
Prot.mirabilis	4	7	7	0	0	0
Prov.stuartii	1	1	1	0	0	0
Totals		527	517	7	1	2

### Table 1. Sources and Species of Basteria Tested for Antibiotic MIC, 1975

The antibiotics used in the test battery were selected in accordance with current trends in the field of antimicrobial usage, and also with reference to the effectiveness displayed in vitro against this particular bacterial population. The antibiotic battery used in 1975 is shown in Table 2.

^{2.} Lindberg RB, et al: <u>Providencia stuartii</u>: A new opportunistic pathogen recovered from burns. <u>In</u> Research in Burns: Proceedings of 4th International Congress on Research in Burns. (in press).

Gram-Positive	<u>Gram-Negative</u>
Gentamicin	Gentamicin
_incoin*	Minocin
Methicillin	Kanamycin
Dxacillin	Ampicillin*
Nafcillin	Keflin
/ancomvcin**	Colvmycin
Keflin	Tobramycin**
Minocin	
Clindamycin	
Tobramycin**	
/ibramycin**	

 Table 2. Antibiotics Used in Testing Gram-Positive and Gram-Negative Bacteria in 1975

- Lincocin was deleted in mid-1975; Ampicillin was deleted in October 1975.
- ** Vancomycin was introduced in August 1975; Tobramycin was introduced in November 1975; Vibramycin was introduced in mid-1975.

# RESULTS OF SENSITIVITY TESTS ON PRINCIPAL SPECIES OF BURN FLORA BACTERIA

Sensitivity of bacterial populations tested in this series are summarized in the following tables. Results are shown in cumulative increments, so that each increment of antibiotic inhibits all strains that were inhibited by lesser amounts. Although the most precise assessment of sensitivity in relation to achievable effective blood level varies with each antibiotic, an approximation of sensitivity demarcation level may be indicated at 6,5 ug/ml₃ for grampositive organisms and 12.5 ug/ml for gram-negative bacilli.

Staphylococcus aureus - The cumulative sensitivity of 153 strains of S. aureus tested in 1975 is shown in Table 3. The group of methicillintype antibiotics has been of particular interest due to the extremely high incidence of methicillin-resistance which was present in staphylococci from 1969 to 1973. In that year, sensitive strains increased in incidence, and sensitivity rose still further in 1974, but in 1975, resistant forms reappeared; only 21.8% of isolates were inhibited at 6.25 ug/ml of methicillin. Gentamicin was lower in inhibitory activity than in the previous year, and a marked decrease also occurred with minocin. The most effective antibiotic in the test

^{3.} Finland M Charging patterns of susceptibility of common bacterial pathogens to antimicrobial agents. Ann Int Med 76: 1009, 1972.

ug/ml	ດ	Ţ	8	Ps	c	<b>VB*</b> *	ζ	s	ū	10**	5
× 25	8	<b>ខ</b> ៍	100	100	<b>1</b> 8	8	18	ĩ	18	8	100
25	93.2	96.6	<b>88</b> .0	93.0	96.4	100	<b>9</b> .3	93.0	98.6	88.8	18
12.5	79.0	96.6	46.3	85.4	93.4	100	9 <b>8</b> .6	69.4	98.6	87. 8	ī
6. 25	38.8	96.6	21.8	73.6	<b>8</b> 5.6	100	97.2	<b>16</b> .5	98.0	88. 8	78.3
3, 12	24.3	72.8	10.5	52.0	71.2	9 <b>8.</b> 8	89.7	34.7	95.3	55.5	26.4
1.56	18.2	47.4	0	27.0	46.4	81.6	76.7	28.4	94.7	11.1	19.3
< 0.78	5.4	13.5	•	6.9	22.8	<b>4</b> .5	52.7	23.6	<b>88</b> ,0	0	Ĩ.
					5	87		Ē	151	•	106

Table 3. Cumulative Inhibitory Levels for 153 Strains of Staphylococcus aureus - ISR, 1975

U: Nafcillin (Unipen); Va: Vancomycin; Kf: Cephalothin (Keflin); M: Minocin; Cl: Clindamycin; G: Gentamicin; L: Lincocin; Sc: Methicillin (Staphcillin); Ps: Oxacillin (Prostaphlin); To: Tobramycin; Vb: Vibramycin.

* Lincocin discontinued in test battery: 1975

** Vancomycin, Tobramycin and Vibramycin added to test battery: 1975

battery was vancomycin; this drug was first tried in 1975. All but one of 87 strains tested were inhibited at 3.12 ug/ml. Keflin was similarly effective, as was clindamycin. Lincocin was deleted from testing during 1975, although it was still effective in vitro. Tobramycin was tested late in 1975; it has thus far been highly active. There were no instances of complete cross-resistance in this group of strains, although cross-resistance was common.

Despite this encouraging proportion of sensitive strains, success in treating staph/lococcal sepsis was not great. Vancomycin was selected as the drug of choice for clearing the blood stream in staphylococcemia, in hopes of preventing the development of endocarditis.

Retrospective comparison over a period of several years offers a major insight on patterns of resistance and susceptibility of staphylococci to antibiotics. The data for the period from 1968 through 1975 are summarized in Table 4. Strains inhibited at 6.25 ug/ml are regarded as sensitive; above that level, , resistant. The progression of increasing methicillin-resistance in the period from 1969 to 1972 is dramatically shown. No comparable incidence of methicillin-resistance has been found in other descriptions of this phenomenon. The re-appearance of a sensitive population in 1973 was described in a previous report; however, the reemergence of resistant strains in 1975 was a new episode, which resembled the earlier experience. The linkage of resistance to methicillin to the aminogly coside antibiotics was suggested by the progression of events with gentamicin. This was a less consistent fluctuation, but the 8 year experience encompassed the extremes between 32% of strains sensitive and 92% sensitive. After 2 years of increasing effectiveness (1973-1974) which paralleled the behavior of methicillin, the strains recovered in 1975 shifted abruntly to a low level of gentamicin sensitivity. Minocin behaved in a similar manner. Other effective antibiotics have not been in use long enough for similar observations to be made.

<u>Staphylococcus epidermidis</u> - Only 6 strains of <u>S. epidermidis</u> were recovered from blood cutlures, each in a single positive culture. Two of the 6 patients died, but neither of these was classified as death due to <u>S. epidermidis</u>. Sensiti /ity varied widely, but from half to two-thirds of the 6 strains were sensitive to the antibiotics used with gram-positive cocci. The highest degree of sensitivity - less than 0.78 ug/ml - was observed with gentamicin, keflin, minocin and clindamycin.

Alpha and non-hemolytic Streptococci - A small but significant number of strains of alpha-hemolytic and non-hemolytic streptococci have been recovered from blood cultures annually. In 1975, 13 strains of streptococci were recovered from 7 different patients. Five of these patients died. The sensitivity of these 13 strains is summarized in Table 5. Most strains were resistant to gentamicin, and all were methicillin and oxacillin resistant. Keflin was virtually ineffective, as was clindamycin. Penicillin G was the most effective antibiotic tested, and half of the strains were sensitive to minocin. This heterogeneous population of streptococci were mainly group D by biotype. Although they were numerically incons 'cuous, their minimal spectrum of antibiotic sensitivity indicates that their esence in burn patients with sepsis presents a potentially

		Annual	Dasis, Ion	( - 1908-19	70			
Antibiotic		Year ar	nd % of Str	ains Inhib	ited at 6.2	5 ug/ml o	f Antibiotic	
	1968	1969	1970	1971	1972	1973	1974	1975
Gentamicin	I	52.0	32.0	50.0	35.0	67.9	92.2	38.8
Lincocin	64.7	48.5	29.8	28.4	26.0	44.3	93.9	96.6
Methicillin	84.6	25.7	18.0	15.5	13.1	50.0	65.2	21.8
Oxacillin	80.0	33.0	22.4	20.1	18.8	69.7	82.6	73 6
Nafcillin	90.0	41.0	33.9	33.0	26.0	62.3	83.3	35.6
Vancomycin	I	I	۱	I	I	1	ŧ	100.0
Keflin	١	۱	I	56.4	22.6	72.1	90.4	97.2
Minocin	ł	I	ı	1	1	84.1	96.0	46.5
Clindamycin	1	ł	ł	1	I	40.7	95.8	98.0
Tobramycin	ŧ	1	1	ļ	1	I	t	88.8
Vibramycin	1	ı	1	ł	I	I	I	78.3

.

2. .

Table 4. Comparative Sensitivity of Staphylococcus aureus on an Annual Basis, ISR - 1968-1975

n-hemolytic od Cultures
22
No Bloc
la and frum
. Alph trains
ы М С
abl f 1

ないない。これにはないないで、これにないないで、ためため、ためにないないとないとないとなっていた。

ŝ
197
•

Concentration ug/ml	υ	ا¥ ب	ntibioti Sc	c and Ps	s of Str U	ains I Kf	nhibit M	C g	Pen G	
>25	100	100	100	100	100	100	<u>5</u>	90 100	100	1
25	23.0	41.6	7.6	84.6	100	69.2	92.3	92.3	100	
12.5	23.0	8.3	0	30.7	100	23.0	76.0	69.2	100	
6.25	23.0	0	0	0	69.2	7.6	53.8	23.0	100	
3.12	15.3	0	0	0	15.3	0	46.1	7.6	83.3	
1.56	15.3	0	0	0	15.3	0	46.1	7.6	0	
<b>≮</b> 0.78	15.3	0	0	0	0	0	46.1	7.6	0	
Total Tested	13	12	13	13	13	13	13	13	ø	

intractable infection as a possibility.

Pseudomonas aeruginosa - Although Ps aeruginosa has been essentially controlled since 1964 by use of appropriate antibacterial topical therapy, it remains a conspicuous part of the burn yound flora, and was third in frequency of species recovered from septicemia in burn patients in 1975. Many of these septicemic episodes can be classified as terminal events but even so, the species can scarcely be described as controlled in this burn population. Ps. aeruginosa has been frequently found to be resistant to most antibiotics, and in the ISR experience, this is indeed the situation. Results of sensitivity tests with strains from 45 patients with positive Pseudomonas blood cultures are shown in Table 6. Gentamicin-resistant strains were in the maiority: minocin was effective in a small percentage of strains but keflin, ampicillin and kanamycin were completely ineffective. Colymycin remained the most effective antibiotic in vitro. Disappointingly, tobramycin and vibramycin were minimally effective. Carbenicillin constituted a special case, since inhibition at high levels of antibiotic still left it acceptable for use. At 156 ug/ml, 88% of strains were inhibited. Two experimental antibiotics, ticarcillin and amikin, were tested extensively in 1974; these drugs were not available for further test in 1975 due to administrative procedures in securing approval of these agents. Both were highly effective in vitro against Ps. aeruginosa. No other antibiotics tested have shown such a level of inhibitory action.

On comparing sensitivity of <u>Ps.aeruginosa</u> in 1975 with results reported in previous years, only one significant change in proportion of sensitivity was found: a drop in propertion of strains sensitive to gentamicin. During 1969-1974, from 62% to 84% of isolates were sensitive to gentamicin; in 1975, only 40% of isolates were sensitive. During the same period, sensitivity to carbenicillin increased gradually from a low of 30% of strains tested in 1971 to 88% in 1975.

Klebsiella pneumoniae - Among Enterobacteriaceae species, K. pneumoniae was most commonly encountered in septicemia in 1975. This constituted a marked change in identity of opportunistic invading bacteria since 1974, when Enterobacter cloacae was a major cause of septicemia. The source of Klebsiella spp is, first of all, endogenous, since they are ubiquitous in the gut. The uniformity of biotype and antibiogram in strains recovered during 197^s, however, suggested that a small number of epidemic strains may be represented, with patient to patient transfer constituting the infection s ince.

Cumulative sensitivity for 159 strains <u>K</u> pneumoniae from 58 patients with septicemia is cummarized in Table 7. The susceptibility of spectrum was not encouraging. Minocin was effective with half the strains; colymycin was active with 73% of strains tested and vibramycin, tested here for the first time, was the only other effective antibiotic. The well-established phenomenon of resistance transfer factor is undoubtedly involved in this increasingly numerous opportunistic pathogen and selective enhancement of a resistant population tends to occur. Table 6. <u>Pseudomones aeruginosa</u> Cumulative Inhibitory Levels for 56 Strains ISR - 1975

* • • •

22

	,		<	ntibiotic and	a of Srai	tichit	<b>ited</b>				
c centration g/mi	υ	2	×	duv	ĸ	D	4 9	•	•	Conc. ug/ml	ප
23 25	108 1.2	100 35.7	100	8 0	8 o	7 6	00 100 .3 18.5		100 100	1250 1250	100 98.7
12.5	<b>9</b> .0 <del>4</del>	16.8	•	•	0	98	.3 18.3		20	625	93.8
6.25	29.4	4.2	•	0	0	4	.2 18.5		10	312	92.5
3.12	23.1	2.1	•	0	0	22	.6 14.8	-	0	156	69.8
1.56	9.4	2.1	•	•	0	35.	7 11.1		0	78	81.4
< 0.75	1.0	6	0	0	0	10	.5 3.7	_	0	39	50.6
		8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	3   							61	9.6
No. Tested	£	95	8	26	96		5 27		10	•	2.4
										Total	5
G: Centamicin;	M: Minocin;	K: Kanto	Amp:	Ampicillin;	Kf: Keflin;	о З	olymycin;	To: T	branvc	.c	

Vb: Vibramycin; Cb: Carbenicillin (testad up to 1250 ug/ml).

* Tobramycin and Vibramycin tes  $r_{1}$  was highn in the last 3 months of 1975 at the time when ampicillin was deleted from the test  $g_{2}r_{1}r_{2}$ .

Concentration			Antil	piotic and 8	of Strains	Inhibited		1	
ug/ml	ດ	Z	~	Ainp	Ę	Co	То	٩٧	1
> 25	100	100	100	100	100	100	100	100	
25	10.8	81.2	5.6	4.7	26.5	73.2	13.2	100	I
12.5	6.3	52.2	5.6	3.9	22.1	73.2	9.6	94.4	
6.25	3.8	21.9	3.7	3.9	7.5	72.6	9.6	88.8	
3,12	3.8	12.2	0.6	3.9	3.7	71.9	9.6	88.8	
1.56	0	6.4	0	0.7	0.6	63.0	2.4	58.3	
<0.78	0	0	0	0	0	46. <i>4</i> i	0	0	
Total Tested	157	155	159	126	158	157	83	36	

Table 7. <u>Klebsiella</u> pneumoniae: Cumulative Sensitivity for 159 Strains ISR - 1975

۰,

がいいと

**N** 

In view of the predominance of K. pneumoniae in sepsis in 1975, a retrospective comparison of sensitivity of strains collected from sensis in ISR patients is appropriate. Table 8 presents this data. Prior to 1974 numerous strains from sputum were tested; in 1974 and 1975, Klebsiella septicemia became far more frequent in occurrence and only blood stream isolates were tested. Sensitivity to Kantrex occurred at the 30% level until 1973, when a sharp increase in sensitivity occurred, but when Klebsiella septicemia strains increased in 1974, sensitivity to Kantrex abruptly fell to less than the 5 to 6% level. Keflin was consistent in inhibiting only about one-fifth of the strains tested, with the exception of the 1973 group. Change in sensitivity to gentamicin was especially disturbing; this useful aminoglycoside, which had been the most effective antibiotic prior to 1974, became virtually ineffective by the time the 1975 collection was tested. An inversion of effectiveness of colymycin occurred in the same interval; this antibiotic was much more effective after 1974 than it had been previously. Cf 2 new antibiotics, vibramycin has been highly effective in vitro; tobramyc., was virtually inactive. The loss of sensitivity that accompanied a marked rise in frequency of Klebsiella sepsis strongly suggests the incursion of an epidemic strain or strains which essentially reside in the burn ward, but this is, at present, only an hypothesis. Further study of these epidemic isolates is being carried on.

Antibiotic		Year a	and % of Stra	ins Sensitive	
	1971	1972	1973	1974	1975
Kantrex	31.2	32.5	72	8.2	5.6
Keflin	24.1	18.6	60.8	13.8	22.1
Gentamicin	71.8	60.4	83.3	15.2	6.3
Colymycin	31.2	48.8	32	95.8	73,2
Ampicillin	0	4.6	8	1.3	3,9
Minocin	NT*	NT	83.3	90.4	52.2
Vibramycin	NT	NT	NT	NT	94.4
Tobramycin	NT	NT	NT	NT	9.6
Total Strains Tested	32	43	25	73	159

Table 8. Comparison of Antibiotic Sensitivity of Strains of <u>Klebsiella pneumoniae</u> from Burned Patients ISR - 1971-1975

### * Not tested

Enterobacter cloacae - In the previous year, 1974, Entero, cloacae increased abruptly in incidence in septicemia and became the predominant species of gram-negative bacillus in the blood of patients who developed sepsis. This incidence decreased abruptly after April 1975, and since June 1975, Entero, cloacae sepsis has been seen infrequently. Sixty-two strains from blood cultures on 31 patients were tested in 1975, and their antibiotic sensitivity is summarized in Table 9. The strains were broadly resistant to antibiotic; only minocin and colymycin were effective at 12.5 ug/ml with 75% of strains tested. The organisms were resistant to gentamicin, ampicillin, kanamycin and keflin. Only 6 strains were tested with vibramycin; its encouraging result will require further confirmation, as will the negative results with 6 strains tested with tobramycin.

The behavior of this epidemic outbreak of a common enteric species resembles the experience of previous years with Providencia stuartii. In such instance a species with no history of causing severe invasive sepsis in burn patients appeared in the burn ward and rapidly assumed a state of predominance. Entero. cloacae was not quite as resistant to antibiotics as was Providencia, but the strains were sufficiently resistant to present a major therapeutic problem when they invaded the burn patient. Although the chronologic sequence of events was consistent with the presence of a continually transmitted opportunistic strain in the burn ward, the same species was recovered from several patients at the time of their arrival at the Institute of Surgical Research, and the species was recognized in at least 2 other burn research centers. Available data does not permit categorization of the outbreak as a self-contained epidemic in a single burn ward, although patientto-patient transmission certainly occurred extensively.

The evident cessation of the Enterobacter sepsis outbreak has no ready explanation. No control measures had been instituted prior to its sudden drop in incidence, nor did the strain actually disappear. In the first half of 1975, 22 patients showed Entero. cloacae in one or more blood cultures. From June to December, only 9 patients had positive blood cultures with this species. Since 1975, Entero. cloacae has been recovered but rarely from wounds, sputum, or blood. This episode illustrates a basic pattern of opportunistic epidemic infection in burn wards; whether the capability of setting up such epidemics is species-related may be determined with continued systematic monitoring of the burn population.

Infrequent Enterobacteriaceae - Escherichia, Serratia, Proteus and Providencia - Species of Enterobacteriaceae less frequently encountered are summarized in Table 10. Escherichia coli and Serratia marcescens were involved in 8 and 7 patients respectively. The organisms were not benign; 5 patients, who exhibited E. coli septicemia, died, as did 6 out of 7 patients with Serratia sepsis. Centamicin, keflin, and especially colymycin were the most effective antibiotics for E. coli. Serratia strains were totally and unequivocally resistant to the established test battery of antibiotics. Vibramycin was available only for testing of 5 strains; 3 of these were sensitive to 12.5 ug/ ml. It is fortunate that this highly resistant species has not shown further epidemic spread; however, the propensity of Serratia for causing nosocomial infections has been documented on numerous occasions in different hospitals, and its behavior in the ISR burn population will continue to be observed in detail.

DISCUSSION

2 Stra
vity of 6 1975
Sensiti s, ISR -
Cultures
cloacae
Enterobacter
.6
ole

; ζ

Concentration ug/ml	υ	Ŵ	K	biotic and 8 Amp	of Strains Kf	Inhibited Co	То	٩
> 25 25	100	100 91.9	100 8.0	100 5.3	100 4.8	100 82.2	100 33.3	100 100
12.5 6.25 3.12 1.56 <0.78	8.3 6.6 3.3 3.3	75.8 61.2 24.1 6.4 0	8.0 0.0 0.0	3.5 0 0 0		77.4 75.8 74.1 74.1 46.7	16.6 16.6 16.6 0	100 100 66.6 50.0 0
Total Tested	60	62	61	56	62	62	9	9

ains Recovered tr Tab

	Escheri	<u>chia, Serratia</u>	, Prote	us, P	rovide	ncia				}
Species	No. Patients	No.Strains		Anti	biotic a	and & Inh	ibited a	at 12.5 u	g/mt	
	Positive		G	z	∽	Атр	Kſ	င့	4	<u></u>
Escherichia coji	8	9	55	22	22	28	55	100	33	33
Serratia marcescens	7	22	0	0	0	0	0	0	0	60
Proteus mirabilis	4	7	50	0	28	43	59	0	0	c
Providencia stuartii			0	0	0	0	0	0	0	0

# Table 10. Infrequent Enterobacteriaceae in Burn Sepsis:

* Only 5 strains of Serratia tested with vibramycin.

The most significant species involved in bacterial sepsis in burned patients in 1975 were <u>S. aureus</u>, <u>K. pneumoniae</u>, <u>Ps. aeruginosa</u> and <u>Entero.</u> <u>cloacae</u>. When septicumia occurred in burn patients with these or with any of the 7 other species of bacteria recovered from the blood in 1975, a relatively high mortality could be predicted, even when an antibiotic active against the offending organism could be found. A retrospective survey of the type set down here aids in selecting the most promising antibiotic at the first indication of invasive infection. STaphylococci, having gone through a remarkably high level of methicillin resistance, regained up to a 68% sensitivity level during 1973 and 1974, only to have methicillin-resistance reappear. The high level of sensitivity to keflin and vancomycin indicated that they are the most promising antibiotics for use at this time. These fluctuations in sensitivity of a microbial species to antibiotics appeared to be unrelated to specific changes in antibiotic usage.

The increase of incidence of K. pneumoniae made this opportunistic invader the principal pathogen in septicemia in 1975. A progressive increase in resistance of Klebsiella to a broad spectrum of antibiotics has occurred during the past 5 years. Kantrex and gentamicin have virtually lost effect on K. pneumoniae; keflin has, during this time, been effective only with 20% of isolates, and colymycin, present the most effective antibiotic, has occupied this status only during the past 2 years. An encouraging pattern appeared with vibramycin, which in a small series tested was highly active. The ease with which resistance factors are acquired by Klebsiella, and its capacity for intensive colonization of burn patients, militate against complacency regarding any antibiotic as being a definitive solution for control of Klebsiella in the burn ward. In 1974, an experimental antibiotic, amikacin (Bristol Laboratories), was tested against gram-negative strains pertinent to the burn wound. It inhibited 75% of strains of Klebsiella at 3.12 ug/ml and was obviously a promising drug for use against such infections. It is to be released for use in 1976.

Entero. cloacae exhibited a sensitivity pattern closely paralleling Klebsiella in 1975. Minocin was more effective against this species than against Klebsiella, but, as with that genus, colymycin was the only other antibiotic in general use that was active against Enterobacter. Vibramycin was completely inhibiting in a preliminary trial at the 6.25 ug/mi level.

Ps. aeruginosa, despite the long history of successful control of surface infection by topical therapy, remains a major part of the flora in sepsis in burns. In 1975, the strains recovered from septicemia were somewhat more resistant to antibiotic than they had been earlier. Gentamicin resistance increased in 1975; only 40% of the strains tested were inhibited at the 12.5 ug/ ml level. Colymycin remained an effective antibiotic in vitro; and carbenicillin was slightly more effective in 1975 than it had previously been.

The armamentarium of antibiotics available for control of gram-negative sepsis in burns remained disturbingly limited in terms of predominant organisms in 1975. Newly developed agents, such as amikacin and vibiamycin merit prompt, careful consideration if this life-threatening problem is to be better controlled.

# PRESENTATIONS

Lindberg RB: Epidemic methicillin-resistant <u>Staphylococcus</u> <u>aureus</u> type 84 in a burn unit. Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC October 11, 1975.

Lindberg RB: Bacteriologic and clinical aspects of methicillin-resistant staphylococcus epidemics in a burn unit since 1968. 2th Annual meeting, American Burn Association, San Antonio, Texas, April 2, 1976.

PUBLICATIONS

None

			· · · · · · · · · · · · · · · · · · ·	1. AGER	CY ACCEN	low	1. DATE OF SU	MARY	REPORT	ONTROL ATMENT
RESEARCH	AND TECHNOLOG	Y WORK UNIT S	UMMARY	na (	DE 695	5	76 10	01	DD-DI	&E(AR)636
S BATE PREV SUMRY	4. KIND OF SUMMARY	S. SUMMARY SCTY	. WORK SECURITY	7. RESR	ADING	<b>PA</b> 0	68"H HISTA"H	OL SPECIFIC	DATA-	. LEVEL OF SUM
75 07 01	H. TERM	U	U	N	A	I N	t.	CONTRACTO		A VORK UNIT
IN NO./CODES:"	PROGRAM ELEMENT	PROJECT	NUMBER	TASK		968		WORK LINI		
	611024	331611028	718		01		304			
. CONTRIBUTING				<u> </u>	<u> </u>		S.7.848-5.2			
E. CONTRIBUTING					······					
11. TITLE (Procedo with	Security Classification Code	(U) Emer	gence of Me	thic	illin-	Res	istant S	taphylo	coccu	s Aureus
Type 84 in	Burned Milita	ry Personn	el (44)				-			
12. SCIENTIFIC AND TE	CHNOLOGICAL AREAS									
003500 Clin	ical Medicine	1								+
IL START DATE		14. ESTMATED COM	PLETION DATE	THE PUN		€¥		14. PERPOR		H00
69 06		76 09		DA	1		ł	C. Ir	-Hous	e
17. CONTRACT/ORANT	Not Applic	able		18. RES	OURCES ES	TIMAT		IONAL MAN YF		OE (In they and)
& DATES/EFFECTIVE:		EXPIRATION:			76		5		16	
N NUMBER:®				FISCAL	7T		1 .ĭ			, 1
G TYPE:		ANOUNT:		YEAR	CONNENT		1		-1	
& KIND OF AVARD:		f. CUM. AMT.								
IN RESPONSIBLE DOD	DREANIZATION	[	- T	38. PER	CHAIN & O	RGANI I	ATION			1
HANE!" US ATTIV	Institute of	Surgical	Research	HANEI	ns Am	w T	netitute	of Su	rgical	Research
· · · · · · · · · · · · · · · · · · ·	****************	Ouryzour .	1600az 01.		Mic	ny – ny –	iningy I	ranch	92002	160004.0
ABORENSI* FORT	Sam Houston	Toyac 782	34	ADDRES	H ^e Rort	. 52	m Yougto	DE GENERAL	ag 78	234
1010	built insusconty	16403 /02	54		E OL I	. 50		MI IGM	15 70	234
				PRINCIP	-		R (Fundah BEAN	if U.S. Anodomi	e jostikulas,	
	AL			-	• Robe	ert	B. Lindl	oera, Pl	nD	
HANG Basil	A. Pruitt, Jr	, COL, MC		TELEP	HONE: 5]	12-2	21-2018			
TELEPHONE 512	-221-2720			-	SECURITY	-				
EI. GENERAL USE	······································			-	TE INVEST		4			
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	NAME						
				NAME						DA
IL XEVECHOS (Procedo	BACH with Security Classifi	cation Co.(a)		<b>4</b>						

(U) Staphylococcus; (U) Burns; (U) Septicemia; (U) Burn infection; (U) Humans BE TECHNICAL OBJECTIVE,* 24. APPROACH, 25. PROGRESS (Purnis' Individual paragraphs identified by sumber Pr. edu test of each with press/first Classification Code,

23. (U) Continued high level of <u>Staphylococcus aureus</u> septicemia persisted despite intensive use of antimicrobial therapy. A persisting monotype epidemic, with recurring methicillin resistance, required characterizing as a valuable model for understanding nosocomial outbreaks and the possibility of controlling them, a major objective in military medicine.

24. (U) Staph phage typing, MIC sensitivity testing and antibiograms were used in epidemic tracing.

25. (U) 75 07 - 76 09 Continued predominance of phage type 84 in epidemic pattern in burn patients occurred despite a reappearance of methicillin sensitive strains in the burn ward. In 1975, a reappearance of methicillin resistance took place. Cephalosporium and glycopeptides especially Vancomycin, remained relatively effective in controlling staphylococcal sepsis. The variability of staphylococcal populations required continued monitoring in view of the unusual fluctuations described.

### TERMINATION

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIE. CE-SURGERY

### REPORT TITLE: EMERGENCE OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS TYPE-84 AND 84,85 IN BURNED MILITARY PERSONNEL

## US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 June 1976

Investigators:

Robert B. Lindberg, PhD Ruth L. Latta, BS Basil A. Pruitt, Jr, MD, Colonel, MC Arthur D. Mason, Jr, MD

Reports Control Symbol MEDDH-288(R1)

**UNCLASSIFIED** 

# ABSTRACT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

### REPORT TITLE: EMERGENCE OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS TYPE 84 AND 84,85 IN BURNED MILITARY PERSONNEL

US Army Institute of Surgical Research Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 20 June 1976

Investigators: Robert B. Lindberg, PhD Ruth L. Latta, BS Basil A. Pruitt, Jr, MD, Colonel, MC Arthur D. Mason, Jr, MD

Reports Control Symbol MEDDH-188 (R1)

Phage typing and MIC sensitivity testing were used to characterize the <u>Staphylococcus aureus</u> found extensively in burn patients. A previous epidemic of Phage type 84 was supplanted in mid-1974 by type 84,85. Extremely high methicillin-resistance was observed, subsiding abruptly after several years, only to re-emerge in 1975. Cross-resistance was observed with cephalothins, and with aminoglycosides as well. Staphylococcal seeding of the burn wound is an obvious source of propagation, and selective pressures, endogenous and exogenous, contribute to the continued persistence of this ubiquitous species. Currently adequate in vitro sensitivity prevails, but the possibility of re-emergence of resistance factors is shown by this study.

Staphylococcus Burns Septicemia Burn Infections Humans

### EMERGENCE OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS TYPE 84 AND 84,84 IN BURNED MILITARY PERSONNEL

The continuing problem of emergence of antibiotic resistant Staphylococci in hospital populations has prompted detailed study of the behavior of <u>Staphylococcus</u> <u>aureus</u> in the patient population of ISR. The organism is ubiquitous, and eradication of seeding by <u>Staph</u> <u>aureus</u> is an unrealistic goal. Control of infection by antibiotic remains the only feasible approach to prevention of staphylococcal sepsis in severe burns. In previous reports (1) description of the appearance of a uniquely methicillin-resistant population of staphylocci was described. Fluctuations of sensitivity were observed over several years, and in 1973 and 1974, susceptible strains reappeared after a 4-year period of maximum methicillin-resistance. However, these organisms do not of necessity remain static, and once more, methicillin-resistance was on the rise in 1975. The necessity for continuing close observation of this major pathogenic species in burn patients is only too obvious.

### Phage Types of S. aureus Over the Period 1968-1975

The incidence of predominant staphylococcal phage types recovered from burn patients in the ISR, from 1968 through 1975, is outlined in Figure 1. The types listed are, of course, not the total number of phage types observed, but they are the major ones that have been identified. The percentage of patients harboring a given phage type is shown as the fraction of staphylococcus positive patients observed; it includes the entire block outline. The solid black area is the proportion of all strains of staphylococci reacting in that type identity for that year.

Type 84 was present from 1968 to 1970, but was a relative minority of the strains typed. It became the predominant type in 1971, and continued in that status through 1974. In the last year, 1975, a striking drop in its incidence occurred. From a status of recovery of type 84 in over 80% of patients in 1971 and 1973, it fell to 67%, of patients in 1974; then in 1975, it was found in only 9% of patients.

Type 47, 54, 75, 84, 85 is presented because of its part in the continuing rise and disappearance of phage types in the burn patients. 15, 1970 it was the predominant form, but in 1971, it disappeared almost completely. A small number of strains were recovered through 1974, but in 1975, none were found.

The close relationship of type 84 with 84,85 has been noted in earlier reports. 84,85 was found in numbers comparable to type 84 in 1968-1970; it then fell to negligible numbers until 1974, when it was found in 34% of patients. In 1975, however, it supplanted type 84 as the predominant type; this transition was abrupt, and unprecedented.

8.2

L. Lindberg RB, Latta RL, Pruitt BA, Jr, Mason AD, Jr: Emerger so of methicillin-resistant Staphylococcus aureus Type 84 and 84,85 in Burned Military Personnel. US Army Ann Historical Rept 1976: pp93-105.



1

Non-typable strains fluctuated in incidence but not to an extent that suggested any marked invasion of a strain not typable by the current typing set. The relatively large proportion (up to 56% of patients harboring non-typables in 1973) with the low proportion of all isolates in this category reflects the fact that any patient may harbor non typable strains but they are seldom predominant in the patient population.

### S. aureus Phage Types in the ISR Burn Ward, 1975

Admissions to the ISR totalled 245 patients in 1975. Among these, 135 patients contributed staphylococci from various sources, and 14 patients admitted in late 1974 were also sources of isolates. 844 strains were thus collected from 149 patients. Sources included wounds, sputum, biopsy, catheter tips, urine, and other sites, plus autopsy tissues. Phage types observed, by patient and in proportion of strains, are shown in Table 1. This was the first time that type 84,85 had predominated since these phages were included in the typing set. It had appeared in 1974, but was found only in half as many patients as harbored type 84. The drop in occurrence of type 84 was marked and dramatic; only 8.7% of patients harbored this type in 1975. Non-typable (NT) strains were found in 40.3% of patients; this number has been even higher in some years, but the proportion of total strains that are NT is always low.

Phage type 94, formerly designated as WH-1, reached a significant level of incidence for the first time 1974, when 10.2% of patients harbored it. This proportion rose to 13.4% in 1975, and more significantly, the proportion of total strains of type 94 rose from 2.3% to 9.5%. Although this strain was not a major problem, it merits close observation since the sequence of increasing incidence resembles that which has been seen with other types in former years.

Phage Type	Patients	Strains
	Per (	Cent
84,85	75.2	61.3
Non-typeable	40.3	16.9
94	13.4	9.5
84	8.7	3.8
29	<b>2</b> <i>k</i>	0.9
85	5•4	0.8
834,85	2.7	0.5

Table 1	<ul> <li>Predo</li> </ul>	minan	t Ph	age	Types	of S	Staphylococ	cus
aureus	Strains	from	149	ISR	Burn	Ward	Patients.	1975

Types 29, 85, and 83A, 85 were found in very small numbers. Their presence confirmed the versatility of the typing system. The total of typed stains and unequivocal NT strains made up 93.7% of all 844 strains collected.

### S. aureus in Septicemia, ISR, 1975

è

The epidemiologic pattern of staphylococcal septicemia and especially of endocarditis and thrombophlebitis in ISR patients suggests, on the basis of clusters of episodes, that specific strains of staphylococci may at times set up circumscribed local epidemics. One clue to the validity of this hypothesis would be differences in incidence of specific phage types from septic episodes in contrast to the overall population of staphylococci. The bacteriologic aspect of septicemia was scrutinized with this possibility in mind. There were 44 patients with staphylococcemia in 1975, and 142 strains of <u>S. aureus</u> were recovered in blood cultures.

The order of frequency with which phage types were recognized is shown in Table 2. Phage type 84,85 was overwhelmingly preponderant; its incidence closely paralleled that found in the total staphylococcus population. In 1974, it was recovered from 31% of patients with staphylococcemia. The proportion of non-typable strains decreased from the level which was found in 1974. Only 9.1% of patients had non-typable strains in blood culture. In contrast to the 6.1% of patients harboring such strains in 1975, the low level of NT strains in blood in 1975 suggests that this specific type 84,85 exhibited selective ability to invade the blood stream.

	Patients	Strains
Phage Type	Per	Cent
84,85	81.8	77.5
NT	9.1	4.9
94	l. c	9.2
29,52	4.5	1.4
84		1.4
83A,85	2•3	0.7
Other	4.5	4.9

# Table 2. Phage Types of <u>Staphylococcus</u> <u>aureus</u> Strains from Blood Stream of 44 ISR Burn Ward Patients, 1975

						Pha	ige Type	
Month	Patient	No. of	84,85	NT	94	84	83A,85	Other
	NO.	Strains		I	L	No. c	of Strain	)\$
	240 74		1		74 ·			· · · · · · · · · · · · · · · · · · ·
Jan	2	0	0					
	221 74	3	3	1	1			alkadaa aha ahada aha
Feb	237 174		1					
	29	2	2	4				
	39	5		1	Î			79 -5
Har	41			1				
!	1.8		<u> </u>			-		
Ane	52	i	1					
	69		1		2	[		
	* 79					-		والمعصور الأملاب والمتحدث فالمتحاد فالأكان
	63	4	4					
May	80			l				
	83	10			10			
	89	4	2			2	[	
Jun	102		3					
	*89	-5-	3					
Jul	113	2	2					42E,47,53,54.75.
	119	1 i	1					83A,81,84 -2
	112	1	1					ويستيد المانات ويستشفها فيادانها بالعالم
	*113							
Aug	144	6	6					
	151	4	4					
	165	3	3					
<b>Sec.</b>	+151	2	2					
260	*163		8					
	189	2	2					
	191		!					
Oct	192	8	7				1	
	197	4	4					
	198		<u> </u>					
	*196	2	2					29,52 -1
	<b>*1</b> 98		1					
NOV	205	2	2					29,52 -1
	216	2	2	,				
<u> </u>	×21K		┝━╤━┥	<u> </u>				
0.00	218	{	1					
Vec	232	2		2				
	241	L i	1					

# Table 3. Phage Types of <u>Staphylococcus</u> aureus from Blood Stream of ISR Burn Ward Patients, 1975

- -- .

Types 94 and 29,52 were each, recovered from 4.5% of patients. The high proportion of strains of type 94. In contrast to the low percentage of type 29,52, reflects septicemia with an unusally large number of positive blood cultures from two patients with bacteremia due to type 94.

Type 84 was found in only 2.3% of patients which meant one patient harbored this formerly predominant type. In 1974, 66.6% of blood culture strains were type 84. One patient harbored type 83A, and two others had one strain each of other types.

The chronologic sequence in which staphylococcal sepsis episodes occurred furnished a basis for recognition of circumscribed epidemic outbreaks. Such a sequence is set down in Table 3. The pattern is one of continuous occurrence of episodes of the epidemic type; other bacteremia showed no indication of clustering or of individual outbreaks.

### S. aureus from Lung Tissue of Burn Patients at Autopsy

Since pneumonia constitutes one of the principal lethal complications in severe burns, the identity of Staphylococci involved in such episodes was of particular interest. The strains recovered from lung tissue at autopsy were regarded as most significant in their relation to pulmonary involvement. Twentysix patients, dying with sev: re burns, yielded 69 strains of <u>S. aureus</u> from ung tissue. The type distribution observed is summarized in Table 4. Seventeen, or 35.4% of the patients had type 84,85 present in the lung. Non-typable strains were recovered from 26.9% of the patients' lung samples. Type 84, type 85, and type 94 were recovered from 4, 2, and one patient respectively. The distribution, in consideration of the small sample size, is to be regarded as consistent with that seen in the blood stream isolates. There was no basis for assuming that a distinctive 'ung infecting strain had been part of the infection.

Ohung Tur	Patients	Strains
Phage Type	Per	Cent
84,85	65.4	58.0
NT	26.9	20.3
84	15.4	13.0
85	7.7	4.3
94	3.8	1.4
Other	7.7	2.9

Table 4.	Phage 1	ypes	of S	tuphy	00000	us aureus	Strains
from Lot		ies of	ISR	Surn	Ward	Patients,	1975

A comparison of the incidence of the principal staphylococcus types in blood, lung and other sites was of importance in its bearing on the relation of specific epidemic type to invasiveness or virulence. Figure 2 summarizes this information. The percent of patients harboring each type in the three categories is represented by the clear outline. The respective sites are differentiated to show the percentage of strains. It was ovious that the specific type 84,85 was most frequently encountered in blocd; in contrast, septicemic strains were least frequently found among non-typable strains. Type 94 was most common among other sources, while type 84 was most frequently found in lung samples. Type 29 was found neither in blood nor lung, while type 85 was found only in lung tissue and other sources. The samples suggest that a tissue predilection differing with phage type may exist.

### Chronologic sequence of S. aureus Phage Types, 1975

The existence of epidemic strains as represented by annual totals does not convey valid information regarding progression of this process. To elucidate the progress of these staphylococcal episodes, the phage typing data for all strains charted by months, to exhibt patterns of appearance and disappearance of various types Figure 3 summarizes this information. The predominant type was 84,85, in every month. Non-typable strains were second in frequency in each month with one episode, in June, when type 94 was equally common. Type 94 exhibited a capacity for establishing limited increase in frequency, in four additional months. In two months, outbreaks of type 84 were concentrated. The remaining types were essentially scattered, but mainly in Jan-Mar and in Aug-Oct 1975.

### Antibiotic Sensitivity of S. aureus Strains, 1970-1975

The epidemic pattern of S, aureus in the ISR burn ward makes obvious the need for effective control of these infections. Sensitivity to antibiotic has varied widely in this Institute over extended pariods of time. Methicillin resistance and associated decreased sensitivity to centamiacin and cephalathin war more extensive than has been reported elsewhere. The changes that have occurred since 1968 are summarized in Table 5. In 1968, a high level of sensitivity to meth.cillin and its analogues was present in the staphylococcal population. In 1969, a drop in proportion of isolates sensitive to methicillin began; it reached its lowest point in 1972, when only 13% of staphylococci were sensitive to methicillin. Oxacillin and nafcillin, although somewhat more effective, also became Ineffective. Gentamicin and cephalothin also became less effective. The pattern changed to reappearance of censitive strains in 1973, and 1974. In 1975, methicillin resistance reappeared, although as yet oxacillin and nafcillin have not lost effectiveness, nor has cephalothin. Minocin resistance increased markedly in 1975, and gentamicin also dropped markedly in in vitro activity against staphylococci. Relatively new antibiotics have been added to the test battery, and among these. Vancomycin has been extremely active against strains thus far tested. Clindamycin, first tested in 1973, displayed a limited degree of effectiveness, but in 1974-1975, alm st all strains tested were inhibited. Tests with Vibramycin were in progress in late 1975; the results were promising for in vitro effectiveness.



-----

Total Patients- Strains Each Month	83A,85	85	29	<b>%</b>	\$	NT	84,85	Phage Type
17-71	1-1	ī	1-2	ĩ	3-16	F	13-34	Jan
15-51		1-3	1-2		-	5-15	10-29	Feb
14-78	•	1-1		3-12	2-5	[]-]	6-27	2
31-134	•			7-15	<del>[</del> -10	15-29	23-76	βpr T
23-110					3-34	8-17	19-56	Мау
16-45				1-2	<b>[</b> - <b>!</b>	5	11-27	
15-34					2-2	I.	11-25	In L
21-77	ī-	Ĩ-]	1-2			2-2	18-65	Aug
16-41			Ξ	-	2-2	5.5	10-31	Sep
18-86	ĩ	-	-			<u>]-5</u>	13-72	Oct
16-50						6-10]	12-34	Nov
18-57				1-1		8-22	13-41	Dec
	6-4	5-7	5-8	13-32	20-80	60-143	112-517	. Total Patients- Strains Each Type

**

----

بمر ب ب

...

1. Sec. 1.

• - ·

Figure 3. Monthly Distribution of Predominant <u>Staphylococcus aureus</u> Phage Types in ISR Burn Ward Patients, 1975

۰.

,

Antibiotic				Year				
	1968	1969	1970	1971	1972	1973	1974	1975
Gentamycin	-	52.0	32.0	56.0	35.6	67.9	92.2	38.8
Minocin	-	-	-	-	51.5	84.1	96.0	46.5
Methicillin	84.6	25.7	18.0	15.5	13.1	50.0	82.6	21.8
Oxacillin	80.9	33.0	22.4	20.1	18.8	<b>69</b> .7	65.2	73.6
Nafcillin	90.0	41.0	33.9	33.0	26.0	62.3	83.3	85.6
Cephalothin	-	-	_	56.4	22.6	72.1	90.4	97.2
Vancomycin	-	-		-	_`	-	-	100.0
Clindemycin	-	-		-	-	40.7	95.8	98.0
Vibromycin	-	-	-	-		-	-	78.0

Table 5. Antibiotic Sensitivity of Staph aureus: % of StrainsInhibited by 6.25 ug/ml or less

Cumulative sensitivity of Staphylococci to antibiotics illustrates with greater precision the nature of sensitivity to antibiotics. Figure 4 shows the sensitivity of Staphylococci to methicillin, oxacillin, nafcillin and cephalothin in 1975. It was evident that cephalothin was by far the most active antibiotic in this group; methicillin was minimally effective, while oxacillin and nafcillin were between these two extremes of activity. The 6.25 mcg/ml level denotes the upper limit of inhibition of strains to be regarded as sensitive. Figure 5 presents the graphic expression of sensitivity of vancomycin, clindamycin, minocin and gentamycin. Action of both minocin and gentamycin, the tetracycline and the aminoglycoside, was minimal; only 45% of strains fell below the 6.25 mcg/ml level. Clindamycin and vancomycin were the most active antibiotics available for control of staphylococcal sepsis. Of these two, vancomycin has been the drug of choice in the treatment of systemic staphylococcal disease, although it was slightly less effective in vitro than was clindamycin.

### DISCUSSION

The staphylococcal population in the ISR burn ward has exhibited a unique and unprecidented lability with reference to methicillin. From a typically sensitive population, an almost completely resistant population emerged by progressive change over a threee year period. Reversal of this pattern occurred abruptly in 1973; but again, resistance has moved upward in 1975. The degree of lability implied in these shifts has not been previously described. Cross-resistance with tetracycline, aminogiycoside and cephalothins was evident during these shifts. Response to antibiotics active in vitro continues to be a variable entity; this variation points up the need for more effective guidance to antibiotic therapy.

# PUBLICATIONS

Lindberg RB, Latta RL, Pruitt BA, Jr: Methicillin-resistant staphylococci in epidemics in a burn unit: Bacteriologic aspect. 15th Intercience Conference



### CUMULATIVE SENSITIVITY OF STAPHYLOCOCCUS AUREUS TO ANTIBIOTIC, 1975

Methicillin, Oxacillin, Nafcillin, Cephalothin

Figure 4.



# CUMULATIVE SENSITIVITY OF STAPHYLOCOCCUS AUREUS TO ANTIBIOTIC, 1975

-- -- . --- .

••• - ----

Figure 5.

on Antimicrobial Agents and Chemotherapy. 1975, Abstract 151.

Sieger BE, Long JM, Lindberg RB, Pruitt BA, Jr: Methicillin resistant staphylococci in thermally injured patients: Clinical aspects. 15th Interscience Conference on Antimicrobial Agents and Chermotherpay. 1975, Abstract 152.

### PRESENTATIONS

Lindberg RB, Latta RL, Mason AD, Jr, Pruitt BA, Jr: Bacteriologic and Clinical Aspects of Methicillin - Resistant Staphylococcal Epidemics in a Burn Unit Since 1968 at the American Burn Association, San Antonio, Texas, March 31 - April 1, 1976.

RESEARCH	AND TECHNOLOGY	WORK UNIT S	UMMARY	DA O	CV ACCESSI C 5970	) )	76 10 (	MARY ⁴	REPORT DD-D	CONTROL SYMPOL R&E(AR)636
S. DATE PREV SUM'RY	4 KIND OF SUMMARY	S. SUMMARY SCTY	. WORK SECUNITY	7. REGRA	CINO	- 01	B'N INSTRIN	BA SPECIFIC	DATA-	. LEVEL OF SUM
75 07 01	H. TERM	U	U	NA	<u> </u>	NI		la ves l	<b>—</b> 100	A YORK UNIT
10. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK A	REA NUME	27		WORK UNI	TNUMBE	N
& PRMARY	61102A	3A161102	371k	0	)1		267			
S. CONTRIBUTING				L						
C. CONTRIBUTING		[					aite e a star	·····		
11. TITLE (Procodo with )	Somethy Classifisation Code;	(I) Sensit	ivity To S	ulfam	ylon d	of F	seudomo	nas Aer	ugino	63
Recovered E	com Burned So	<u>ldiers (44)</u>	L							·····
003500 Clin	ical Medicine									
15 START DATE		14. ESTIMATED COM	LETIS BATE		NHE AGENC	Ŷ		1. PERPORM	ANCE ME	THOD
68 07		76	09	DA	1		1	C. In	-Hous	e
17. CONTR CT/GRANT	Not Applica	ole	·····		DURCES EST	MATE		DNAL MAN YR	s h FUI	NDE (In thousands)
A DATES/EFFECTIVE:		EXPIRATION:			76		.5		1	8
N NOMBER.*				FISCAL	ŹŤ		.2			6
G TYPE:		& ANOUNT:		YEAR	CURNERT		T		T	
& KIND OF AWARD:		I.CUM. AMT.								
IS RESPONSIBLE DOD C	DREANIZATION		1	30. PERP	OPINING OR	GANIZ	ATION			
HANE!" US Army	Institute of	Surgical 1	Research	N ANE.*	US Arı	ny I	Institut	e of Su	rgica	l Research
				İ	Mie	crob	oiology	Branch		
ADDRESS:* FOTT	Sam Houston,	Texas 782	34	ADDREN	• Fo	rt S	Sam Hous	ton, Te	xas	78234
				PRINCIP	AL INVESTI	SATOR	(Furnish MAN) 	US Asadoonio	: Inelikulie:	*
RESPONSIBLE INDIVIDU	AL			NAME:"	KODE:	<b>Γ</b> Γ Ι	8. LINGD	erg, Pn	IJ	
NAME: BASIL A	. Pruitt, Jr,	COL, MC		TELEP	HONE: J.	12-4	221-2010			
TELEPHONE: 512	-221-2/20			BOCIAL	- SECURITY	ACCU				
21. GENERAL USE			•	AMOCIA	TE INVESTI	NOTAL Maria		nalich	мъ	
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D		¥۷.	сдті		ngrran,	t.ru	DA.
Z REVEABLE /Prosede	LACK WID Somethy Closell	sation Code)		HANEI						DA
	(**)	- (11) Con	Tformal on a	(11)	oni an l	+h	aranu. 1	11) Huma	ne	
(U) PSeudom	ONAS; (U) BUI	$\frac{ns; (U) - SU}{PROGRESS (Pumies)}$	dividual peredraphe id	entified by	NUE HAT. Pro		I al anosh with St	outly Classifi	callen Ced	• )

23. (U) Extensively burned soldiers have been increasingly colonized with <u>Pseudomonas aeruginosa</u> during the past 18 months when initial treatment of burns has been primarily topical silver sulfadiazene. 5% sulfamylon soaks are used to prevent invasion of separating eschar, and hence, the environment offers continuous exposure to this drug. Sensitivity or resistance of opportunists to this agent merits detailed study, since control of infection is a primary objective of combat casualty care.

24. (U) Sensitivity is assessed as MIC, in an agar dilution plate. Values are evaluated with epidemic phage types.

25. (U) 75 07 - 76 09 In 1975, Sulfamylon sensitivity for <u>Pseudomonas aeruginosa</u> increased to correspond to the median range, with a small but significant resistant population which required 1.25% or greater for inhibition. This confirms the hypothesis of the cluster of resistant strains developing in a transient manner in topically treated wounds.
### TERMINATION

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

### REPORT TITLE: SENSITIVITY TO SULFAMYLON OF PSEUDOMONAS AERUGINOSA RECOVERED FROM BURNED SOLDIERS

### US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Robert B. Lindberg, PhD Virginia C. English, MS Basil A. Pruitt, Jr, MD, Colonel, MC Arthur D. Mason, Jr, MD

Report Control Symbol MEDDH-288 (R1)

UNCLASSIFIED

### ABSTRACT

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

### REPORT TITLE: SENSITIVITY TO SULFAMYLON OF PSEUDOMONAS AERUGINOSA RECOVERED FROM BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Robert B. Lindberg, PhD Virginia C. English, MS Basil A. Pruitt, Jr, MD, Colonel, MC Arthur D. Mason, Jr, MD

Report Control Symbol MEDDH-288 (R1)

Six hundred thirty-seven strains of <u>Pseudomonas aeruginosa</u> were monitored for sensitivity to Sulfamylon, using an agar plate MIC dilution method. The strains were more sensitive than the 1972 collection, and even the 1973 set. The median sensitivity level was 0.125% over a 9-year period this value had fluctuated between 0.008 and 0.316%, but the median for 3548 strains was 0.129%. Sensitivity was correlated with predominant phage types; NT-16 and H-1 were sensitive at 0.625%, but the predominant type, E-2, was primarily sensitive to 0.039%. Sulfamylon may be unique in permitting prolonged use with no concrete proof that resistant variants are selected by it.

Pseudomonas Burns Sulfamylon Topical therapy Humans

### SENSITIVITY TO SULFAMYLON OF PSEUDOMONAS AERUGINOSA RECOVERED FROM BURNED SOLDIERS

Understanding of the behavior of microorganisms in response to longterm exposure to an antibacterial agent is of basic importance in the control of nosocomial infections., Control of infection in burns involves primarily topical therapy; present practice in the Institute of Surgical Research involves use of both Sulfamylon' burn cream and silver-sulfadiazine cream as topical agents, with follow-up soaks of 5% Sulfamylon used to control invasive infection during the period when eschar is separating and intermittent debridement is carried out to ready the burn site for skin grafting. On the basis of experience with long-term use of other antibacterial agents, it would be expected that bacterial strains resistant to Sulfamylon might be derived by continued exposure of a heterogeneous population to this drug, and continued monitoring of Pseudomonas aeruginosa strains recovered from treated wounds has been carried out since the introduction of Sulfamylon as a generally used medication. The topical agents control other bacteria as well, but Ps aeruginosa is probably the greatest threat to the burn wound because of its specific proclivity for setting up invasive infection and wound sepsis. This scrutiny of Ps.aeruginosa from a bacterial population exposed to a broad spectrum antibacterial such as Sulfamylon is also merited due to the insight it gives on mechanisms of development of resistance in general, and equally important, a view of the absence of derived resistance among strains in which it does not appear.

Technics of testing sensitivity to Sulfamylon have been described (1). The test is essentially seeding of agar plates containing serial dilutions of Sulfamylon, and reading presence or absence of growth as evidence of inhibition.

### SENSITIVITY OF PSEUDOMONAS AERUGINOSA TO SULFAMYLON

There were 637 strains of <u>Ps. aeruginosa</u> tested in 1975. This was the largest number collected in one year; over the past 8 years, the average number tested was 363. Silver-sulfadiazine was first used extensively beginning in 1974 in the Institute of Surgical Research; since its use, the rate of colonization of burns by <u>Ps.aeruginosa</u> has increased to approximately twice what it had been when Sulfamylon was used exclusively.

The sensitivity of the strains in 1975 are set down in Table 1. For comparison, results of tests for the preceding 4 years are presented. The proportion of strains inhibited at levels up to 0.312% was not markedly changed from results recorded in 1974. For the previous 5 years, the sensitivity level has not changed markedly, although an increase in strains requiring 0.625% for inhibition occurred during one year, 1972.

^{1.} Lindberg RB, Calvert J, Brame RE, Dent R: Sensitivity of burn wound flora to Sulfamylon. USA Surg Res Ann Rpt FY 1965, BAMC, Ft Sam Houston, Tx. Section 15.

Table 1

Inhibiting Concentrations of Sulfamylon for Pseudomonas aeruginosa. 1971-1975

-

-

		Ŭ	oncent	ration o	f Sulf	amy lon	in % an	d Numbe	r inhibi	ted
Year		2.5	1.25	0.625	0.312	0.156	0.078	0.039	0.019	< 0.019
1/61		0	0	48	ή]	56	57	65	13	0
	80)	0  	0	17. 1	-14.6	_ 20.0_	20.4	23.2	- 4.7	
1972		0	29	212	9†	88	31	37	15	Ŋ
Z_of total (4	<u>- [</u> 3]	0  	- <u>6</u> .3	45.8	<u> 9</u> .9	1.61	- 6.7		- 3.2	  -    
1973		0	4	14	85	85	52	32	12	-
%_of total (2		0		4.9	2.8	2.5	-18.3	- 11.2	- 4.2 -	0.4 -
1974		0	S	59	78	67	97	86	11	4
%_of total (4	37) -	0  		13.5	_ <u>18.0</u>	22.2	22.2	<u>-19-7</u>	_ 2.5 _	6.0
52é1		-	13	113	108	155	68	147	28	4
<u>%_of total (6</u>	37)	0.16	2.0	<u> </u>	_16.9	24.3	10.7		. 4.5	
Total: 2102		-	51	9444	358	181	305	367	6/	14
% of total		0.4	2.4	21.2	17.0	22.9	14.5	17.5	3.8	0.66

•

Cumulative sensitivity to Sulfamylon offered a clearer picture of the behavior of <u>Ps aeruginosa</u> toward this drug. The information is presented in Table 2, for years from 1967 through 1975. Inhibition was complete at 0.625% and almost total at 0.312% from 1968 through 1970. From 1971, onward a slight increase in resistance to Sulfamylon has been manifest. There was a sharp rise in 1972, when over half the strains tested required 0.625% for inhibition, but this figure subsided to a level of between 80 and 85% inhibited by 0.312% from 1973 through 1975. The sensitivity has remained relatively constant since 1972.

Median sensitivity, the level at which 50% of strains are inhibited, offered another measure of the behavior of <u>Ps. aeruginosa</u> in relation to the long-term exposure. These figures are presented for 1967 through 1975 in Table 3. After continuous exposure to the drug for over 10 years, the Pseudomonas population was still, and for the past 3 years, has been remarkably consistent in its response to this drug. It is a cliche to speak of acquired resistance on prolonged exposure as being virtually a law of nature, but the experience with Sulfamylon is an exception to this common experience.

Graphic representation of cumulative sensitivity levels since 1971 makes more evident the consistency of behavior of <u>Ps</u> aeruginosa toward Sulfamylon (Fig. 1). Out of the years shown, in 3 the sensitivity curve was virtually identical. The marked shift in sensitivity in 1972 was returning to this norm during 1973, to account for the right shift in sensitivity during that year.

Throughout studies of the role of <u>Ps.</u> aeruginosa as a pathogen in burns, the importance of recognizing individual strains in relation to a given attribute has been demonstrated. Phage type identity of strains of Pseudomonas was correlated with sensitivity. It was shown that certain types, present in micro-epidemic populations of bacteria, were consistent in sensitivity.

Sensitivity to Sulfamylon was not uniformly distributed over the bacterial population. Some strains were quite specific and consistent in sensitivity to Sulfamylon. Other types were less consistent in their behavior with this drug. The results with 4 major strains which typed only with undiluted phage (the NT-strains) are shown in Figure 2. NT-16 and NT-16V (a varient pattern) showed a relatively wide range of sensitivity, as did NT-19V. NT-19 and NT-22, however, showed a preponderance of strains inhibited at one particular level, 0.312% for NT-19 and 0.156% for NT-22.

Strains reacting with Routine Test Dilution in phage typing behaved in a manner comparable to the NT-types. Table 4 shows the behavior of 2 predominant phage types. H-1 included 35 strains inhibited at the relatively high level of 0.625%. However, there were also 12 strains of H-1 more sensitive to Sulfamylon. The attribute is strain-related but not absolute. The E-2V strain was mainly more sensitive, with 15 strains out of 19 inhibited by 0.039° or less.

Phage type E-2 (type 24 in phage number) was highly sensitive to

	Table	2. Cur	nulative	Sensitivity	to Sulfa 1967-197	imylon of	Pseudo	monas ac	ruginos	σI
Year	No.of Strains	2.50	1.25	Per Cent 0.625	Concentr 0.312	ation an 0.156	d of Stra 0.078	iins Inhi 0.038	bited 0.019	د ٥.019
1957	1.74	100	100	96.8	87.6	81.7	61.3	46.4	15.6	I
1968	294	100	100	100	95.1	60.4	45.8	14.1	1.7	8
1969	385	100	100	100	96.5	50.0	26.9	7.7	0.5	I
0261	296	100	100	100	100	78.0	49.9	21.9	2.0	ı
1871	Ň	100	100	100	82.9	68.3	48.3	27.9	4.7	1
1972	463	100	100	93.7	48.0	38.0	19.0	12.3	4.3	- <b></b>
1973	285	100	100	98. 1	81.3	57.0	33.5	16.1	3.2	. 0.4
ħ261	437	100	100	0.66	85.5	67.5	45.3	23.1	2.4	0.9
1975	637	100	99.8	97.8	80.1	63.2	38.9	24.2	5.1	0.64

I

* * * *

Year	No.of Strains	Median Inhibitory Level % Concentration
1967	471	0.083
1968	294	0.136
1969	385	ď.176
1970	296	0.068
1971	280	0.125
1972	463	0.316
1973	285	0.111
1974	437	0.086
1975	637	0.125
Total: 9 Years	3548	0.129

### Table 3. Median Value of Pseudomonas aeruginosaSensitivity to Sulfamylon: 1967 - 1975

2

\$



The second second second second

- -



ŗ





Table 4.SULFAMYLON SENSITIVITY REACTION OF VARIOUSPREDOMINANT PHAGE TYPES - 1975

44 100

Sulfamylon. Eighty out of 83 strains were inhibited by 0.039% or less; of these 68 were inhibited at that level, as shown in Table 5. The consistency of behavior of an epidemic strain is clearly demonstrated here. Routine testing for antibiotic sensitivity does not segregate individual strains; the data shown here is unusual because it is not often collected. However, the actual pattern of antibiotic sensitivity in nosocomial infections may be equally consistent. The study here reported may well serve as a model for specific strain sensitivity of other species of bacteria, if adequate sorting criteria were collected. This strain-linked sensitivity component prompts investigation of other species for further clues to antibiotic action.

### PRESENTATIONS

Lindberg RB. Association of Drug Resistance and Extreme Virulence with Specific Bacteriophage Types of <u>Pseudomonas</u> aeruginosa. Am Assoc.Immunol. Anaheim, CA, April 1976.

### PUBLICATIONS

Lindberg RB, Latta RL, Pruitt BA, Jr, Mason AD, Jr: Association of Drug Resistance and Extreme Virulence with Specific Bacteriophage Types of Pseudomonas aeruginosa. Fed Proc 33: 763, 1976 (Abst).

### Table 5. SULFAMYLON SENSITIVITY REACTION OF PREDOMINANT PHAGE TYPE - 24



				1. AGEN	CY ACCES	ion ^a z	DATE OF SU	WARY	REPORT	CONTROL STREOL
RESEARCH AND T				DA	)B 695		76 IC 0		DD-DI	R&E(AX)636
75 07 01 H. 1	TERM	U	U .	NA	(Q140)	NL	r 48 Halb I 49 Ha		ACCESS	A WORK WET
IS. NO./CODES!" PROGRA		PROJECT	NUMBER	TASK /	REA NUM			WORK UNIT	NUMBER	3
- PRIMARY 6110	026	3A161102	B71R	01			191			
b. CONTRIBUTING				<b> </b>		ŧ				
e. CONTRIBUTINS	ani linetian Code	Pround Tref		l	Flor	ia of	Burn W	lounds o	f Mil	itary
(U) Parnogenesis Personnel Receivi	or Burn ing Sulf	amylon or	Silver Sulf	adia	zene 1	reat	ment (4	4)		
12. SCIENTIFIC AND TECHNOLOGIC	CAL AREAS									
IL STARY BATE	Medicille	14. ESTMATED COM	LETION DATE		HAG AGEN	CY		IS. PERFORM	NCE MET	NOD
65 07		76 09		DA			ł	C. In	-Hous	se
W. CONTRACT/GRANY NOC	Applica	ble		16. RES	DURCES ES	TINATE		ONAL MAN YRS	1 10	106 (In these acide)
A DATES/EFFECTIVE		EXPIRATION:			76	•	.5		1 1	.5
A TYPE:		4 AMOUNT:		PISCAL YEAR	711 2038287		+ <u>·</u> ²		+	8
. KIND OF AWARDI		I. CUM. AMY.			ł		1			
I. RESPONSIBLE DOD ORGANIZAT	NON			20 PER	ORMINGO	GANIZ/	TION			
"AME" US Army Inst	itute of	Surgical	Research		US Art	ny Ir	nstitute	e of Sur	gical	Research
ADDRESS!" Fort Cam H	ouston.	Texas 782	34	ADDRES	Mic: Forf®	robic t Sar	ology Bi n Housto	canch on, Texa	s 78	3234
FOIC Sam In	ouscom	ICAUD OF	•••				-			
				PRINCIP	AL INVEST	IGATOR	(Furnish BEAN)	II U S. Acodomic	jnelikusies	v
RESPONSIBLE INDIVIDUAL	itt .Tr.	COY. MC		NAME	ROD	ert J 512-3	8. Lina 221-2018	serg, Pn	• 0 •	
TELEPHONE: 512-221-	2720			SOCIAL	SECURITY	ACCOU	NT NUMBER			
21. GENERAL USE						GATOR	1		_	
FOREIGN INTELLIG	ENCE NO	CONSIDERE	D	NAME:	Anthe	ony i	A. Conti	reras, M SD6	S	۵ <b>۵</b>
REVERSE (Press EACE -IN John (Countries Code)										
(U) Burns; (U) Staph aureus; (U) Enterobacter cloacae; (U) Sepsis; (U) Humans										
23. TECHNICAL OBJECTIVE. \$ 24.	APPROACH, 28	PROGRESS (Purish I	ndfridual paragraphs id	entitled by	number pr	*****	it of each with \$	eeurity Classifie	allen Codi	•••
23. (U) The sev	verely b	urned patie	ent is extr	emely	SUSC	epti.	ble to (	opportu etablish	listi ad a	С R Л
bacterial invas.	on, and	topical po ve treatmer	t. Howeve	r, in	vasiv	e in	fection	with a	succ	ession .
of species of or	portuni	stic normal	flora has	come	to r	epre	sent th	e princi	ipal (	cause of
death in such in	jury.	Ongoing mor	nitoring is	esse	ential	to	obtain	a valid	pict	nre of
the constantly o	hanging	pattern of	E lethal in	fecti	on in	bur	ns, a m	ajor pro	brew	in
military medicin	ne.									
24. (11) Kound.	sputum.	blood, uri	lne, biopsy	and	cathe	ter	culture	s, plus	quan	titative
autopsies are us	sed and	collated.								
25. (U) 75 07 -	- 76 09	Rise in St	taphylococc	us au	ireus	of a	single	phage 1	type	has been
shown. Increase	e in <u>Pse</u>	udomonas a	eruginosa a	nd <u>K</u>	lebsie	lla	pneumon	iae occi	irred	; a 19/4
epidemic of Ente	erobacte	r <u>cloacae</u> (	declined, a	is dia	i all Indino	but G GG	a remna	mide ti	herap	v and to
lethal problem a	species,	Providenc.	burn wour	d in.	fectio	on mu	del in	animals	• •	2
ald in developin	ny a mor	e ferbaurn					-			
1										
		ove/								

,

----

- - - ---

.

_ - <del>-</del>~

^

DDI HAR 53 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 14984 I NOV 68 AND 1498-1 I MAR 68 (FOR ARMY USE) ARE OBSOLETE

### TERMINATION

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE -SURGERY

### REPORT TITLE: PATHOGENESIS OF BURN WOUND INFECTION: BACTERIAL FLORA OF WOUNDS OF MILITARY PERSONNEL RECEIVING TREATMENT WITH SULFAMYLON OR SILVER-.SULFADIAZINE

### US ARMY INSTITUTE OF SURGICA' RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Robert B. Lindberg, PhD Anthony A. Contreras, MS Ruth L. Latta, BS Daniel T. Zamora, SP6

Report Control Symbol MEDDH-288 (R1)

UNCLASSIFIED

### ABSTRACT

### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

### REPORT TITLE: PATHOGENESIS OF BURN WOUND INFECTION: BACTERIAL FLORA OF WOUNDS OF MILITARY PERSONNEL RECEIVING TREATMENT WITH SULFAMYLON OR SILVER-SULFADIAZINE

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Robert B. Lindberg, PhD Anthony A. Contreras, MS Ruth L. Latta, BS Daniel T. Zamora, SP6

Report Control Symbol MEDDH-288(R1)

Bacterial sepsis has continued as the major cause of morbidity and death in burns in 1975. Principal offending species included <u>Staphylococcus aureus</u>, <u>Klebsiella pneumoniae</u> and <u>Pseudomonas aeruginosa</u>. <u>K. pneumoniae</u>, although not a newcomer in the area of burn infection, increased to a striking degree as a cause of wound sepsis; it literally replaced <u>Enterobacter cloacae</u>, the previous epidemic species, in early 1975. <u>Ps. aeruginosa</u> was higher in incidence than was the case when topical Sulfamylon^{*} was used exclusively in the ISR; its recurrence has accompanied the routine use of silver-sulfadiazine as the primary topical agent now used in the ISR. Self-limited outbreaks, epidemic in^{*} pattern, occurred with <u>Serratia marcescens</u> and <u>Escherichia coli</u>; these episodes are reminders that such forms are potential epidemic agents, as far as any control measures for limiting their spread are currently concerned.

Burns Staphylccoccus aureus Enterobacter cloacae Sepsis Humans

### PATHOGENESIS OF BURN WOUND INFECTION: BACTERIAL FLORA OF WOUNDS OF MILITARY PERSONNEL RECEIVING TREATMENT WITH SULFAMYLON OR SILVER-SULFADIAZINE

The incidence of invasive burn wound sepsis has been strikingly lowered during the past 12 years, by use of effective topical chemotherapy on the burn wound. This reduction was primarily due to control of <u>Pseudomonas aeruginosa</u>. However, since 1969 a continued high level of sepsis in burned patients has prevailed, with a variety of opportunistic pathogens which include <u>Staphylococcus</u> <u>aureus</u> and <u>Enterobacteriaceae spp</u> as the offending organisms. Pseudomonas has not been eliminated, although it seldom causes primary invasive wound sepsis when adequate topical therapy is used. Pseudomonas wound infection, especially in terminally ill burn patients, is common but presently resembles more an opportunistic invasion rather than a specific infectious process. In burn deaths which occur more than 3 weeks post injury, extensive proliferation of gram-negative bacilli occurs in those patients who develop septic shock. This terminal overgrowth of bacteria in the wound usually is made up of several species. Use of appropriate antibiotics has not appeared to improve survival in these conditions.

A previous report ¹ has pointed out that the potentially infected burn surface, modified by such antibacterial agents as Sulfamylon or silver sulfadiazine, harbors a bacterial population which can vary widely in the species present, and in harboring a wide variety of phage types of <u>Ps. aeruginosa</u>. Both topical agents are used extensively, and as with any broad spectrum antimicrobial agent, the wound flora is unquestionably modified as to identity by the presence of these agents. A succession of opportunistic invaders has occurred over the past 7 or 8 years; the conspicuous new epidemic species in 1975 was <u>Klebsiella</u> pneumoniae.

### ANTEMORTEM BACTERIOLOGY OF BURN PATIENTS, 1975

The bacterial flora and yeasts recovered from 7 major sites in clinical specimens are shown in Table 1. Two hundred and forty-five patients were admitted in 1975, but 26 of these had no cultures taken. The information shown here represents organisms recovered from cultures from 219 patients. Blood cultures were the most frequent category of specimens and were collected on 92.6% of all patients cultured. This emphasis reflects the concern regarding sepsis as the major prot em in microbiology of burns. Out of all specimens submitted 47.8% were blood cultures. The largest number of organisms recovered came from sputum and Luken's tube cultures. Together with wound cultures,

^{1.} Lindberg RB, Contreras AA, Latta RL, Smith HOD, Jr, Zamora DT: Pathogenesis of burn wound infection: Bacterial flora of wounds of military personnel receiving treatment with Sulfamylon or silver-sulfadiazene. USA Inst Surg Res Ann Res Prog Rpt FY 1975, BAMC, Ft Sam Houston, Tx. p. 121.

			Source a	nd Numb	er of l	solates		
Organism	Wound surface	Blood	Lukens Sputum	Urine	T.V. Cath.	Foley Cath.	Biopsy	Totai Isolates
S. aureus epidermidis	277	174	267 71	27 28	57	17	42	861 181
						-	-	
Alpha hemol-strep.	24	0	249	6	1	1	0	281
Beta hemol-strep.	4	0	1	3	2	1	0	11
Non hemol-strep.	94	12	163	69	14	24	13	727
Corynebacterium sp.	0	0	2	0	0	0	0	2
Bacillus sp.	15	2	6	1	8	2	1	35
Pseudomonas so.	383	101	320	120	68	50	83	1125
Mima-Herellea op.	11	0	19	1	Õ	Ō	0	31
K. pneumoniae	209	179	395	164	71	53	46	1117
Foloro Berogenes	2	0	6	0	0	0	0	8
cloacae	107	55	43	58	17	22	11	313
Serratia marcescens	16	29	50	27	10	5	8	145
liquefaciens	0	0	Ő	1	Ó	0	0	1
E coli	72	8	111	48	6	32	17	294
Citrobacter sn	4	ŏ	5	0	1	1	0	11
Prot mirabilis	55	8	39	20	6	8	12	148
morganii	2	ŏ	- 3	0	0	1	0	6
rettoerii	ō	ŏ	Ó	3	0	0	0	3
vulgaris	ĩ	Õ	Ó	1	0	0	0	2
Prov. stuartii	24	3	30	10	5	7	2	81
Neisseria sp.	- 1	ō	11	Ō	0	0	0	12
Candida spp.	55	2	53	75	18	9	7	219
No.patients cultured	176	203	133	157	170	120	63	
No. of specimens	1257	3288	803	738	471	194	135	
Total isolates								5276
Total specimens	6886							
Total patients on whom one or more cultures	210							
were done	219							

### Table 1. Antemortem Bacteriology of Burn Patients, 1975

these cultures offered definitive information as to the source of invasive sepsis when it occurred. S. aureus was the principal gram-positive organism on burn patients. Among gram-negative species, <u>Ps. aeruginosa</u> and <u>K. pneumoniae</u> were recovered in equal numbers, and were by far the most common gramnegative bacilli. <u>Enterobacter cloacae</u>, which had been the predominant species in 1974, became uncommon in June 1975, after an unprecedented predominance in the burn ward for a year and a half.

Chronologic changes in proportion of prominent species are shown in Table 2. Comparison of strain incidence is shown in terms of percent of all isolates recovered in 1975. <u>Ps. aeruginosa</u> increased markedly in incidence; the incidence in the previous 5 years was an average of 12.8%, while in 1975, 21.4% of isolates were Pseudomonas. <u>K. pneumoniae</u> was recovered on a 5 year average, from 11% of clinical specimens prior to 1975; in that year, 21.4% of specimens were positive for Klebsiella. The other species which showed a striking change was <u>Providencia stuartii</u> which was present in only 1.5% of the clinical specimens. Its numbers had decreased steadily since its high point in 1972, when 23.1% of clinical specimens harbored this unusual opportunistic pathogen.²

Entero. cloacae was the most recent example of a species of normal intestinal flora which had assumed epidemic proportions as a burn wound invader, in 1974. In early 1975, after a virtual crescendo of cross-infection, it suddenly became relatively uncommon in burn patients.

The number of isolates of each species offers an indication of the relative frequency of occurrence of these opportunistic offenders, but does not indicate how many patients were involved with a given species. The sites cultured and the incidence of patients positive for a given site are summarized in Table 3. Not unexpectedly, staphylococci were most common on the burn itself. Blood, sputum and urine were the sites most often positive for K. pneumoniae. Pseudomonas was most common in biopsies, which are more significant than surface cultures in defining burn wound flora. In contrast to previous years, Entero. cloacae decreased from third to sixth rank in frequency in various specimen sources. Ps. aeruginosa w. far more ubiquitous than it had been in the recent past, and K. pneumoniae increased in incidence as an opportunistic invader over the previous 3 years.

### BURN WOUND BACTERIOLOGY

The source of bacterial sepsis in a severely burned patient is often obscure. The burn wound itself is frequently described as the entry point for bacteria, and in the specific instance of Pseudomonas burn wound sepsis, the pathognomonic, lesion can be demonstrated microscopically. However, oppor-

^{1.} Lindberg RB, Mason AD, Jr, Pruitt <u>BA, Jr: Providencia stuartii</u> as a major factor in burn wound infections. Am Soc Microbiol 1973 130 (abstr).

^{2.} Toplitz C, Davis D, Mason AD, Jr, Moncrief JA. Pseudomonas burn wound sepsis. J. Pathogenesis of experimental burn wound sepsis. J. Surg Res 4, 200, 1964.

		% of All h	solates From C	linical Specin	nens
Species	1970	1971	1972	1973	1974
S. aureus	12.6	15.0	13.8	19.6	18.6
Pseudomonas sp.	13.6	12.4	13.2	10.4	14.4
K. pneumoniae	11.5	9.7	11.5	10.1	12.8
E. coli	6.4	11.0	6.1	10.4	8.2
Prov. stuartii	21.0	15.2	23.1	15.7	6.6
Entero.cloacaè	Not	2.0	3.8	4.3	11.8
	Differentiated				
% of all isolates	65.1	63.3	67.7	66.2	72.4
No. of isolates	3293	3179	6696	5672	5689

### Table 2. Predominant Species Among Isolates from Clinical Specimens 1970-1975

Organism	Woun Surfa	d ice	Bloc	d	Luke	n's Jm	Urine	1	T.V. Cath.	Foley Cath.	Biopsy
S. aureus epidermidis	103 32	(1)	45 6	(2)	77 38	(3)	22 ( 19	6)	38 12	15 15	27 (3) 5
Alpha hemol-strep	21		0		88		5		1	1	0
Beta hemol-strep	3		0		3		3		2	1	0
Gp A strep.	0		0		3		0		0	.0	0
Non hemol-strep	46		5		59		42		14	24	13
Corynebacterium sp	. 0		0		2		0		C	0	0
Bacillus sp.	9		1		5		1		7	2	1
Pseudomonas sp.	101	(2)	39	(3)	87	(2)	56	(2)	50	33 (2)	41 (1)
Mima-Herel!ea gp.	9		1		9		1		0	0	0
K. pneumoniae	83	(3)	55	-(i)	92	(1)	78	(1)	48	43 (1)	28 (2)
Entero, aerogenes	2		0		3		0		0	0	0
cloacae	49	(4)	25	(4)	23	(6)	31	(4)	17	22 (3)	9 (5)
Serratia marcescens	i1	3	7		14		16		7	5	4
liquefaciens	- 0		0		0		1		0	0	0
E. coli	42		9		33		28		6	24	12
Citrobacter sp.	- 4		0		4		0		1	1	0
Prot.mirabilis	26		- 4		10		12		5	6	,
morganii	2		0		3		0		0	1	0
rettgerii	0	)	0		0		2		0	0	0
vulgaris	1		0	r i	0		1		0	0	0
Prov. stuartii	15		2		11		4		5	6	1
Neisseria sp.	1		0	)	11		0		C	0	0
Candida sp.	31		2	2	27	,	29		10	16	5
Total patients Samples	176	5	203	1	133	5	157		170	120	63

Table 3. Antemortem Burn Patients Cultured, 1975

124-12 115-102

•

( ) relative frequency of patients positive from this site.

tunistic species of Enterobacteriaceae, which so often cause lethal sepsis in severely burned patients, do not exhibit unequivocal sub-dermal invasion nor has it thus far been possible to create a burn wound sepsis model in animals for any bacterial species except Ps. aeruginosa. The wound surface is certainly the most plausible site to exhibit bacterial populations which are altered by topical agents, and observations summarized in this report indicate that this may well have occurred with the change in the Institute of Surgical Research, from Sulfamylon to silver-sulfadiazine as the principal topical agent in early treatment of the burn wound. The environment continued to include the effect of Sulfamylon on the burn, sin :e 5% Sulfamylon soaks have continued in extensive use as a means of controlling bacterial proliferation during the period of eschar separation in the burn. Burn wound surface flora was determined on the basis of cultures on 176 out of the 245 (71.8%) patients admitted. The proportion of patients cultured out of all patients admitted has tended to become less with the increasing awareness that, in the two extremes of minor injury and of massive burns, the identity of the bacterial flora in the burn is of little value as a guide to therapy.

Bacterial species recovered from burn wounds, either from the wound surface or from biopsy, are summarized in Table 4. The highest incidence of colonization occurred with <u>S. aureus</u> and <u>Ps. aeruginosa</u>, with each of these species recovered from over half of the patients cultured. The predominant form of Staphylococcus was type 84,85; however, in terms of overall incidence, there were 25% fewer patients with staphylococcal colonization. <u>Staphylococcus</u> epidermidis appeared in fewer than 20% of patients as had been the case in earlier years. Non-hemolytic streptococci, primarily group D, were also relatively infrequent in occurrence. The incidence of non-hemolytic streptococci is, on an annual basis, relatively consistent.

K. pneumoniae was the third most common species recovered from the burn wound, and was found with twice the frequency observed in wounds in 1974. In this respect it crowded out Entero. cloacae which was recovered from 27.8% of the patients in contrast to the 44.4% found positive in 1974. The other striking change in wound flora was the marked decrease in Prov. stuartii, which appeared on wounds of only 8.5% of patients, from an incidence of 41.6% in 1974. This erstwhile epidemic pathogen became almost a rarity in 1975. Other enteric species, Escherichia coli and Proteus mirabilis, have been recovered from tewer than 1 out of 5 burned patients.

One organism which, although usually rare on burn wounds is continually cought, is the group A beta-hemolytic streptococcus. There is a sound basis for this surveillance; although it can be readily controlled by penicillin, these streptococci can rapidly destroy autograft if they seed the wound. When it appears on the burn ward, in any individual, it must be eradicated by antibiotic prophylaxis, and survey of the population is called for. Table 5 illustrates the wide fluctuation in incidence of group A streptococci that has been observed since 1969. There were two years, 1972 and 1974, in which relatively large' numbers of patients were colonized with group A streptococci. In 1975, only 3 strains were recovered. None of these involved an actual wound infection, and each was eradicated promptly by administration of penicillin. However,

Species	No.of Strains Recovered	No.of Patients Positive on Burn Surface	% of Cultured Patients Positive
S. aureus	277	103	58.5
epidermidis	42	32	18.2
Strep, non-hemolytic*	94	46	26.1
K. pneumoniae	209	83	47.2
Enterobacter cloacae	107	49	27.8
E. coli	72	42	23.9
Proteus mirabilis	55	26	14.8
Providencia stuartii	24	15	8.5
Pseudomonas sp.	383	101	57.4

### Table 4. Burn Wound Surface Flore in 176 Patients ISR, 1975

* These strains primarily group D.

the potential emergence of streptococci in a burn population requires continuing monitoring of this species.

Year	No. of Strains of Group A Streptococci Recovered
1969	8
1970	2
1971	1
1972	56
1973	3
1974	30
1975	3

### Table 5. Incidence of Group A Streptococci on Burn Wounds

### **RESPIRATORY TRACT FLORA IN BURNS**

Pneumonia is a major threat in the severely burned patient. The bacterial species associated with pulmonary complications are those that also colonize the burn wound, although the proportional distribution of species may vary from that seen on the burn. There were 803 specimens of sputum cultured from 133 patients in 1975. Thus, 54.2% of patients admitted

had at least one sputum culture. An average of 6 specimens per sampled patient were cultured.

Trends in species occurring in pneumonia in the burned patient are shown in Table 6. Incidence of the major species of significance in the ISR since 1971 are summarized here.

Species	<u>% of Patients</u>	Exhibiting	Positive	Sputum or	n Culture
	1971	1972	1973	1974	1975
S. aureus	43.0	38.5	56.9	59.2	57.9
K. pneumoniae	45.?	58.8	60.0	68.4	69.2
Enterobacter cloacae E. coli	11.0 27.2	27.0 40.9	23.8 53.8	35.7 33.7 21.4	17.3 24.8
Providencia stuartii	33.0	56.5	40.8	59.2	8.3
Pseudomonas sp.	39.0	38.5	36.2	63.3	65.4
Patients cultured	94	122	130	98	133

### Table 6. Principal Species of Bacteria Recovered from Respiratory Tract of Burned Patients, 1971-1975

An increase in incidence of <u>S.aureus</u> in sputum occurred in 1973, and this trend was continued through 1975. This increase was reflected in a concomittant rise in rate of recovery of staphylococci from the lung of burn fatalities at autopsy.

A more gradual but consistent rise in rate of recovery of <u>K.pneumoniae</u> from sputum was evident. This species was the predominant organism in sputum over the entire 5 year period. Out of 72 autopsies in which complete cultures were made, 40 (55.5%) showed <u>K. pneumoniae</u> as predominant in at least one lung lobe. Out of the same series, 31 (43%) had one or more lobes predominant for <u>Ps. aeruginosa</u> and 20 (27.7%) had <u>S.aureus</u> predominant. The high rate of recovery of Klebsiella in sputum thus was paralleled by its predominance in the lung tissue at death. Its pathogenic role was more than ever predominant during 1975.

Ps. aeruginosa increased markedly from 36.2% positive in sputum in 1973 to 64% in 1974 and 1975. This change coincided with the period of introduction of silver sulfadiazine as a topical agent in early treatment of burns, and it closely paralleled the recovery rate of Pseudomonas from burn wounds. The recovery rate was consistent for 1974 and 1975.

Entero. cloacae went from the status of a very prominent species in burn sepsis in 1974 to a low incidence in sputum and in surface colonization during this period. The incidence in sputum in 1975 was half that found in 1974. Lung tissue at autopsy showed a comparable decrease in this species, which adds weight to the significance of the sputum culture as an index of invasive infection in the lung. Only 3 autopsies out of 72 had Entero. cloacae predominant in at least one of 4 lobes of the lungs; one of these was predominant in 3 lobes. In 1974, out of 74 autopsies exhibited a predominance of Entero. cloacae in lung tissue. In the 23 patients with positive sputa in 1975, there were only 43 specimens positive. Thus, even when Entero.cloacae appeared in sputum it was non-persistent.

### SEPTICEMIA AND BACTEREMIA IN BURNED PATIENTS

The continuing lethal problem in severe burns is sepsis, and in consequence the most significant data regarding relative importance of opportunistic invaders is the population recovered in blood cultures. The impression remains strong that the identity of blood stream invaders may be a fact secondary to the existence of an altered susceptibility of the host, so that ultimate control of this problem lies with preventing this collapse of defense capability. However, survival may also be favorably enhanced by control of offending strains of gram-negative opportunistic invaders.

In 1975, 203 out of 275 patients (Table 7) contributed blood cultures during hospitalization. One hundred of these individuals had at least one positive blood culture. There were 12 genera or species differentiated among bacteremic strains; only 4 of these were of major numberical importance. <u>K.</u> <u>pneumonlae</u> was recovered from the highest proportion of patients positive, and over three-fourths of patients with this species in the blood stream expired. Mortality accompanying gram-negative bacteremia has consistently been higher than that seen with staphylococcal sepsis. Fifty-seven per cent of patients with staphylococcemia expired in 1974; the same ratio was observed in 1975. Fatality rates for patients with gram-negative sepsis ranged from 69% to 78% for the 3 numerically important species.

JSepticemia in severely burned patients usually involves more than one species of organism in successive cultures. The mortality seen in patients in whom only one species was recovered furnish a more precise insight into the relationship of invading organism to outcome. Only 3 species of organism were recovered as one-species bacteremias. Table 8 summarizes the result in these cases. Survival prospects in patients exhibiting only staphylococcemia were relatively favorable. For Klebsiella and Enterobacter sepsis, the survival rate was far less. The other 9 species recovered from burn patients were found as part of a polymicrobial septicemia.

The total of blood cultures from which the septicemia data was drawn in 1975 was modified somewhat by a study conducted on the clinical service, in which multiple blood cultures were drawn during manipulations of the wound. Surgical debridement under anesthesia and hydrotherapy treatment were the 2 modalities studied. It has been assumed that septicemia might arise from manipulation of the colonized wound, which would make more probable the entrance of bacteria into the blood stream. These observations were directed

Species Recovered	No.Patients Positive	No. Isolates	% of all Patients Cultured Positive	No.Patients Who Expired	% of all Patients With Positive Cultures Who Expired
S coag .Pos .	45	174	22.1	26	57.7
S. coag.Neg.	6	14	3.0	مد	16.7
Non-hemol.Strep.	ъ	12	2.5	σ	100.0
Bacillus sp.	2	2	6.0	-1	50.0
Ps aeruginosa	39	101	19.2	27	69.2
K. pneumoniae	55	179	27.0	43	78.2
Entero, cloacae	29	55	14.3	20	68.9
Serratia marcescei	ns 7	29	3.4	6	85.7
E. coli	7	9	3.4	ω	42.5
Proteus mirabilis	4	æ	2.0	ω	75.6
Providencia stuart	tii 2	ω	0.9		50.0
Candida sp.	2	a) 2	0.9	2	100.00
No. Patients with	at least 1 p	positive blood o	culture: 100		

### Table 7. Blood Culture Isolates from 203 Burned Patients, 1975: Relation of Species Recovered to Mortality

Species	No.Patients With One Species Recovered	Average No. Positive Blood Cultures/ Patient	Deaths	% Mortality For One Species Septicemia
S. aureus	12	3.9	3	25.0
Ķ. pneumoniae	16	3.3	10	62.5
Entero.cloacae	3	1.9	3	100.0

### Table 8. Bacteremia with Only One Species of Bacteria Recovered: Burn Patients, 1975

toward answering that question.

Twenty-eight patients were sampled with from 3 to 8 successive cultures collected before, during and after manipulation of the burn wound. Of these 16 showed no positive blood cultures at any time during their course, including the manipulation. Twelve patients had one or more positive blood cultures. Of these, 5 patients, whose burns were large, died, with an average burn size of 63.0% of the body surface. The 7 survivors of at least one positive blood culture had body surface burns averaging 38.9%. Table 9 compares the results of intensive sampling before, during and after manipulation of the burn wound, with the blood stream flora indicated by cultures collected on the basis of clinical status, on days when no manipulative procedures were performed. Among the 5 fatal burn cases, the similarity of the 2 sets of conditions was obvious: organisms recovered in the manipulated patient were quite comparable to those recovered on days when no manipulation occurred. Among the 7 cases who survived, 4 had no blood cultures positive under routine clinical conditions, but did vield staphylococci (2 cases), and one patient each with a Klebsiella or a Pseudomonas strain recovered. Such recoveries occurred in a total of 5 procedures; a total of 18 series of cultures were negative under similar conditions. Three patients displayed bacteremia with staphylococci (2 patients), Klebsiella (2 patients) and 1 Serratia positive. Under manipulation, the staphylococcus appeared in 2 patients, while 1 transient Klebsiella bacteremia yielded a Pseudomonas after manipulation. The contribution of this special set of observations to the total picture of septicemia in severely burned patients was not large enough to justify exclusion of these patients from the total for 1975, and indeed, the implication that such manipulation markedly affected bacteremia was not strongly supported. A slight increase in variety of species could be discerned in the septicemic patients, when they were subjected to manipulation, but no persistent conversion from sterile to contaminated blood stream occurred.

Blood cultures in 62 patients, or 62% of those with bacteremia, were positive for more than one species in successive cultures. In some instances

Outcome & Patient No.	% Burn	Cultures Drawn Other Than On Day of Manipulation	Cultures Drawn During Manipulation	
Died		Organism Recovered	Organism Recovered No. o Sets	f Culture Drawn
1	57.5	Klebsiella; Staph , Pseudo .	Klebsiella; Pseudo; Serratia; Proteus	2
2	80.5	None collected	Klebsiella; Staph.	1
3	89.0	Staph; Klebsiella; Pseudo; Enterobacter; Serratia	Serratia; Proteus; Pseudo; E . coli	2
4	47.0	Staph; Klebsielia	Staph; Pseudo; Klebsiella No growth	6 1
5	71.0	Kleb ^{, e} lla	Klebsiella No growth	2
Survived				
6	54.5	No growth	Staph. No growth	1 6
7	63.0	No growth	Staph . No growth	2 7
8	38.0	No growth	Klebsiella No growth	1 1
9	44.5	Staph; Klebsiella	Staph; Klebsielia No growth	4 2
10	39.5	Staph; Serratia Enterobacter	Staph; Klebsiella	1
11	21.0	Klebsiella	Pseudo. No growth	1 3
12	12.0	No growth	Pseudo. No growth	1 4

### Table 9. Patlents with Positive Blood Cultures During Wound Manipulation

a species would appear, disappear, then reappear, and in faully ill patients a terminal period of mixed blood cultures was not uncommon. The extreme heterogeneity of species observed in blood cultures is shown in Table 10. As has been pointed out in previous summaries, the absence of a consistent pattern of mixed cultures is a conspicuous feature of sepsis in the severely burned patient. Staphylococci were associated with 19 different combinations of Enterobacteriaceae species; the most common pairing of S. aureus with Entero. cloacae or with Ps.aeruginosa occurred in only 4 of each out of 31 patients. The most common pair was Klebsiella and Pseudomonas, in 8 patients out of 19 with Klebsiella bacteremia. Recurrent proposals to immunize burned patients to protect them against bacterial sepsis have been submitted. The bacterial flora of the blood stream in these severely burned patients suggests that this desirable end may indeed be a fond hope; while mobilization of versatile defense mechanism against this plethora of opportunistic invaders is not inconceivable, it does not appear plausible. It is important that the ubiquitous role of staphylococci in this redundant accumulation of species not be overlooked in any regimen designed to control sepsis.

### **BIOPSIES AND BURN WOUND FLORA**

Ę

1.1.1.

Biopsy as a diagnostic procedure has become well established technic. It is used primarily to assess the status of potential invasive wound infection, and also to evaluate the degree of success that therapy has attained. The bacterial population found in sub-surface tissue of the burn area is not of necessity a reflection of the surface colonizing flora. The predominant organism in a biopsy sample of burn wound may give a clue as to the identity of the most significant part of the burn wound flora, as far as potential invasive infection is concerned.

Table 11 presents the bacteriologic results collected on biopsies on 65 patients during 1975. This was 25.7% of all patients admitted in 1975, and contrasts to the 71.8% of admissions on whom wound surface cultures were collected. The numerically important species were Ps. aeruginosa, K. pneumoniae, and S. aureus. Non-hemolytic streptococci, E. coli, and Entero. cloacae were present in biopsies on from 21% to 14% of the patients. When the survival rate in relation to species recovered from biopsy was assessed, the highest mortality rates were recorded for Klebsiella, E. coli, and non-hemolytic streptococci. Fatal outcome from Pseudomonas and Entero. cloacae were virtually identical in recovery at 55.6% to 56.1% of patients S. aureus was associated with a lethal sepsis in but 44.4% of patients.

During the past decade it has been observed that major burn wound infecting organisms vary in identity over a long period of time. Illustrative of this hypothesis are the sequences of Prov. stuartii, Entero. cloacae and K. pneumoniae that have successively dominated the nosocomial infection pattern in the Institute of Surgical Research since 1969. Table 12 summarizes the annual incidence of the major species in biopsy samples and also displays the mortality rate associated with each species. S. aureus has consistently been recovered in from 40% to 50% of potents biopsied. The behavior of this species, even when a succession of phage types has been present,

### Table 10. Blood Culture Isolates in Patients with More than One Species Recovered

Species	No. of Patients
S. aureus, Pseduomonas sp.	4
S. aureus, Entero. cloacae	4
S. aureus, K. pneumoniae	3
S. aureus, E. coli	1
S. aureus, K. pneumoniae, E. coli	2
S. aureus, K. pneumoniae, Entero. cloacae	2
S. aureus, Entero. cloacae, Pseudomonas sp.	2
S. aureus, S. epidermidis, Pseudomonas sp.	1
S. aureus, Pseudomonas spNon-hemolytic strep.	1
S. aureus, K. pneumoniae, Pseudomonas sp.	1
S. aureus, K. pneumoniae, Pseudomonas sp.,S. marcescens	1
S. aureus, K. pneumoniae, Pseudomonas sp.,Entero. cloacae	3
S. aureus, K. pneumoniae, Pseudomonas sp., Prov.stuartii	1
S. aureus, K. pneumoniae, Pseudomonas sp., Mima-Herellea gp.	1
S. aureus, K. pneumoniae, E. coli, Prot. mirabilis	1
S. aureus, K. pneumoniae, Entero. cloacae, S. marcescens	1
S. aureus, K. pneumoniae, Pseudomonas sp. S. marcescens, Prot.mirabilis	1
S. aureus, K.pneumoniae, Pseudomonas sp., Entero. cloacae, non-	
hemolytic strep	1
<ol> <li>S. aureus, K.pneumoniae, Pseudomonas sp., S.marcescens, Prot.mirabilis E. coli</li> </ol>	1
K. pneumoniae, Pseudomonas sp.	8
K. pneumoniae, S. marcescens	1
K. pneumoniae, Entero. cloacae	4
K. pneumoniae, S. epidermidis	1
K. pneumoniae, Pseudomonas sp., Candida sp.	1
K. pneumoniae, Pseudmonas sp., Non-hemolytic strep.	1
K. pneumoniae, Pseudomonas sp., Entero. cloacae	1
K. pneumoniae, E. coli, Candida sp., S. marcescens	1
K. pneumonia, Entero. cloacae, Pseudomonas sp., Non-hemolytic strep.	1
Entero, cloacae, Prov. stuartii	
Entero, cloacae, Pseudomonas sp.	6
E. coli, Pseudomonas sp.	1
E. coli, S. marcescens	
Bacillus sp., S. epidermidis, Pseudomonas sp.	1
Non hemolytic strep., Prot. mirabilis	r
Total patients with 2 or more species	62
<u>% of All</u>	Positives

No. of patients with 2 species	35	35
No, of patients with 3 species	14	14
No. of patients with 4 species	10	4
No. of patients with 5 species	2	2
No. of patients with 6 species	1	1
No. of patients with Entero cloacae	26	
S aureus	32	
Prov stuartii	2	
K pneumoniae	38	
Pseudomonas sp.	35	

Species	No . Patients Positive	8 of Patients Positive	NoPatients with Positive Cultures Who Expired	% of Patients Positive Who Expired
S. aureus	27	49.2	12	ti . titi
S. epidermidis	Ŋ	8.0	-	20.0
Non-hemol.strep	13	21.0	ω	61.5
Bacillus sp.	-	1.6	-	100.0
Ps. aeruginosa	11	65.1	13	56.1
K. pneumoniae	28	44.44	22	78.6
Entero.cloacae	6	14.3	ß	55.6
E. coli	12	0.61	6	75.0
Prot. mirabilis	S	4.7	œ	100.00
Prov. stuartii	F	1.6	F	100.00
Candida sp.	9	9.5	ŵ	50.0
S. marcescens	7	6.3	m	75.0

Table 11. Bacterial Flora of Biopsies  $\omega n$  Burn Wounds of

.

No. of specimens collected: 135 No. of samples per patient (average):

2.1

		190	59 as C	ompare	ed with	-1971 ו	1975					i
Species		010	of Patie	nts Po	sitive		% of	Patien	ts Pos	itive V	Tho Ex	pired
Species	1969	1971 <u> </u>	1972	1973	1974	1975	1969	1971	1972	1973	1974	1975
	413	44	41	51.9	50.4	42.9	22	38	22	50.9	45.6	44.4
J. adleds	3 i	17	32	17.0	14.1	44.4	50	31	19	55.5	78.9	78.6
	<b>I</b>	1	I	19.8	47.3	14.3	ł	ł	ι	61.9	59.3	55.6
Enter V. Sloacac	14	10	27	25.5	20.7	19.0	717	33	16	65.6	67.9	75.0
	<u>7</u>	110	56	36.8	32,6	1.6	14	58	36	51.0	70.5	100.0
	75	1	14	9.4	9.6	4.7	38	40	9	60.0	53.8	100.0
Prot. IntraDitis	30	30	32	32,1	23.0	65.1	39	57	20	50.0	67.7	56.1
rs, der ugitiosa		1										

# Table 12. Burn Wound Biopsy Flora: Species Incidence and Mortality

is consistent with the hypothesis that it is an endogenous part of the burn flora. Despite the availability of antibiotics effective against staphylococci. its incidence has remained constant. Although it is ubiquitous, it has not increased in rate of spread over the past 8 years. Prov. stuartij presents the most dramatic example of a species of Enterobacteriaceae which reached a high level of spread in born wounds, at the same time when it was a major cause of sepsis. It then diminished in tissues biopsied, from 56% of patients in 1972 to 32% of patients in 1974, after which it virtually disappeared from invaded burn wounds. When it does again appear, it is associated with a high mortality rate. Entero, cloacae was of so little consequence in burn patients that it was not differentiated as a species until 1973. In that year it was found in 20% of patients biopsied. In 1974 it was present in almost half of biopsied patients, and then, in 1975, it virtually disappeared. The 14% incidence rate in 1975 is made up of the samples collected in the first half of the year. Its implications with regard to patient survival have remained relatively constant; over half of the patients with Entero. cloacae in wound biopsy expire. K. pneumoniae became the predominant species in 1975; it was found in half the patients biopsied, and on a monthly, not annual basis, it became far more common in the second half of the year. A disturbing rise in mortality rate, among patients with Klebsiella-positive biopsies has occurred; among predominant organisms found in tissues, it was associated with a continuously rising mortality rate, which reached 78.6% of patients thus positive in 1975.

### **KLEBSIELLA PNEUMONIAE IN BURN PATIENTS**

The change from one species to another of opportunistic invasive coliform bacillus in burn ward patients has prompted close study of the offending species. The emergence of <u>K</u>. pneumoniae in 1975 was a classic example of an epidemic of a pathogen which had not previously shown marked invasive potential in burn wounds. The extent of involvement of burn patients with Klebsiella in various sites is whosn in Table 13. Autopsy data from burn wound and lung are also presented. This predonderance of Klebsiella at autopsy adds weight to the implication that the antemortem Klebsiella bacteremia was clinically significant in the patients' demise.

An association between incidence of Klebsiella sepsis and any increase in seeding of the burn patient with <u>K. pneumoniae</u> was sought. Table 14 summarizes the incidence of Klebsiella seeding associated with septicemia over the 1971-1975 period. Seeding of the wound surface fluctuated over a range of 25% to 47%; sputum cultures similarly were positive in from 45% to 69% of patients. The incidence in 1975 was the highest recorded, but it was high also in 1974. Biopsy cultures, however, were more consistent with the rise in incidence seen in blood cultures. The conservative interpretation of these findings would suggest that, while surface and respiratory tract flora varied, tissue invasion as reflected in biopsy findings more nearly paralleled the rise in incidence of bacteremia. The incidence of positive wound and sputum cultures was sufficiently high to indicate that prompt and consistent transfer of Klebsiella from patient to patient in the burn wards was occurring.

No.Isolates/ Total Specimens	% of Specimens Positive	No.Patients Positive/ Total Patients Cultured	<b>% of Cultured</b> Patients Positive
209/1257	16.6	83/176	u7.2
46/135	34.1	28/63	44.4
179/3288	5.4	55/203	27.1
395/803	49.2	92/133	69.2
164/738	22.2	78/157	49.7
53/194	27.3	45/120	37.5
71/471	15.1	45/170	26.5
136/261	52.1	48/75	64.0
156/287	54,4	69/75	80.0
	No. Isolates/ Total Specimens 209/1257 46/135 179/3288 395/803 164/738 53/194 71/471 136/261 156/287	No. Isolates/ Total Specimens& of Specimens Positive209/125716.646/13534.1179/32885.4395/80349.2164/73822.253/19427.371/47115.1136/26152.1156/28754.4	No. Isolates/ Total Specimens% of Specimens PositiveNo. Patients Positive/ Total Patients Cultured209/125716.683/17646/13534.128/63179/32885.455/203395/80349.292/133164/73822.278/15753/19427.345/12071/47115.145/170136/26152.148/75156/28754.469/75

## Table 13. Klebsiella pneumoniae: Isolates from Clinical and Autopsy Specimens, 1975

'ear	Sit	e and Per C	Cent Positiv	'e
	Wound Surface	Biopsy	Blood	Sputum
1971	25.2	17.0	5.0	45.0
1972	43.8	32.0	9.5	58.8
1973	33.3	17.0	6.6	60.0
1974	25.7	14.1	14.4	68.4
1975	47.2	44.4	27.0	69.2

Table 14. Per Cent of Patients Cultured Who Harbored Klebsiella pneumoniae, 1971–1975

The epidemiologic shift from Entero. croacae to K. pneumoniae has been demonstrated as the major event in septic infection in burns in 1975. As a corollary, the change in incidence of Entero. cloacae in tissues and fluids of burn patients has been summarized to complete this sequence, which was not only intrinsically dramatic and unexpected, but also the most well documented episode thus far observed, in which 2 closely related species acted in succession to set up epidemics in a susceptible patient population. This occurred abruptly and resulted in a new nosocomial infection of great lethality. Table 15 shows the reciprocal level of Entero. cloacae and K. pneumoniae in various sites in 1974 and 1975. The replacement Enterobacter by Klebsiella was especially notable in biopsy, blood and post mortem tissues. Among the latter, the reciprocal replacement of Enterobacter by Klebsiella was numerically very precise.

Table 15. Klebsiella pneumoniae and Enterobacter cloacae: Isolates from Clinical and Autopsy Specimens 1974-1975

	Enterobac	ter cloacae	Klebsiella	pneumoniae
Site	1974	1975	1974	1975
Wound	35	27.8	25.7	47.2
Biopsy	47.3	14.3	14.1	44.4
Blood	20.2	14.3	14.1	27.1
Spútum	35.2	17.3	68.4	69.2
Urine	32.6	19.7	27.9	49.7
I.V. Tip	20.5	10.0	13.1	26.5
PM*Lung	30.6	10.7	9.5	36.5
PM Liver	27.5	2.0	10.8	80.0
PM Wound	40.0	32.0	36.0	64.0

* PM: post mortem

### CATHETER TIP CULTURES

The etiology of intravascular infection in severely burned patients is in most instances associated with the indwelling catheter as a source of infection, either as an avenue for ingress of infecting organisms or as a nidus on which circulating bacteria may attach. I.V. catheter tips are routinely cultured as part of the ISR operation at time of removal from the patient. Table 16 summarizes the observations on organisms recovered from this source. Four hundred seventy-one catheter tips from 170 burn patients were cultured. Ninety-one, or 53%, of these patients had positive cultures for at least one I.V. tip. Numerically important species, in order of frequency of occurrence, were Ps. aeruginosa in 55% of patients, K. pneumoniae (49.5%) and S. aureus (41.8%). The high incidence of Ps. aeruginosa was consistent with the incidence on burn wound cultures and biopsies. It was evident that, in this period, a close correspondence between wounds and I.V. catheter flora existed, at a period when sepsis was predominantly due to Klebsiella. The Pseudomonas incidence was most plausibly a reflection of burn wound colonization flora.

Species	No. Patients Positi∨e	% of All Patients Positive	% of Patients With Positive Cultures
S. aureus	38	22.3	41.8
S. epidermidis	12	7.0	13.1
Alpha.hemolstrep	1	0.6	1.1
Beta hemolstrep	2	1.2	2.2
Non-hemolstrep	14	8.2	15.4
Bacillus sp.	7	4.1	7.7
Ps. aeruginosa	50	29.4	55.0
Mima-Herellea gp.	0	0	0
K. pneumoniae	45	26.5	49.5
Entero. cloacae	17	10.0	18.7
S. marcescens	7	4.1	7.7
E. coli	6	3.5	6.6
Prot. mirabilis	5	2.9	5.5
Prov. stuartii	5	2.9	5.5
Candida sp.	10	5.9	11.0
Citrobacter sp	1	0.6	1.1

Table 16. Bacterial Flora of I.V. Catheter Tips from 170 Burn Patients, 1975

No. catheter tips cultured: 471 Average catheter tips per patient: 2.7 No. of Patients with Positive Cultures: 91

DISCUSSION

The period reviewed in this summary coincided with a major epidemiologic change in the nature of burn wound infection in the ISR. S. aureus remained, as it had in recent years, an important part of the infection flora in burns in this Institute. But the most lethal organism, and the cause of a major part of sepsis in the severely burned, was K. pneumoniae.. This opportunistic enteric species succeeded Entero.cloacae as the predominant opportunist in setting up infection which could lead to sepsis. No other gramnegative species incited an epidemic of sepsis in 1975. Ps. aeruginosa, however, became more prominent than it had in the recent past, especially in burn wound tissue. The lessened anti-pseudomonal activity observed in wounds topically treated with silver sulfadiazine compared with burns topically treated with Sulfamylon would appear to be an explanation for this rise in incidence of this ubiquitous burn wound colonizer. The virtual disappearance of Prov. stuartii and Entero, cloacae as wound invading forms were the most striking epidemiologic events in 1975. The possibility that other enteric species may emerge as opportunistic pathogens in the future is real. Antibiotic susceptibility of most of these forms makes them potentially susceptible to chemotherapy, but experience with sepsis due to Enterobacteriaceae does not offer an encouraging prospect for future control of infection in burns.

### PUBLICATIONS

Lindberg RB, Pruitt BA, Jr, Mason AD, Jr: Topical Chemotherapy and Prophylaxis in Thermal Injury. Chemotherapy vol. 3, 1976, pp. 351-359.

### PRESENTATIONS

Lindberg RB. Topical chemotherapy and the treatment of burns. IX International Congress of Chemotherapy, London, England. July 26, 1975.

Lindberg RB: Summarizing comments and session on Topical Chemotherapy of Wounds. IX. International Congress of Chemotherapy, London, England. July 27, 1975.

Lindberg RB: The microbiology of severe burns. Birmingham Accident Hospital, Birmingham, England. Surgical and Medical Staff Seminar. July 30, 1975.
	AND TECHNOLOGY	Y WORK UNIT S	UMMARY	1. AGEN	OB CO	017   .	76 16		DD-D	CONTROL STRE R&E(AR)436
DATE PREV SUMPY	4 KIND OF SUMMARY	S. SUMMARY SCT	A WORK SECURITY	DA UB 0397		9/	TO IU UI		IC DATA- A LEVEL OF MW	
	אפושיוי א	11	tī	N	A	N	L		CCESS	A WGRK LM
/5_U/_UL	PROGRAM ELEMENT	PROJECT	NUMBER	TASK A	REA NUM			WORK UNIT	UNBE	LR
PRIMARY	611022	3A161102	B71R	1	)1	<u> </u>	188			
. CONTRIBUTING							2.20		<u>.</u>	
C. CONTRIBUTING			· · · · · · · · · · · · · · · · · · ·			ľ	<u> Markins</u>		<u> 22</u>	~~~X6? X
1. TITLE (Procode with )	Security Classification Code	,* 	nes lamai		For-3	4-	Burned	Soldiere	( ^ /	1
(U) Bacterio	opnage Types	or rseudom	onas Aerugi	nosa	round	TU	Barnea	SOTATEL8	(444	
003500 CIT	INNOLOGICAL AREAS	•								
IS START DATE		14 ESTIMATED COM	PLETION DATE	-	NO AGEN	: <b>v</b>		16. PERFORMA		THOD
62 06		76 09		DA	1		1	C. In-	Hous	se
T CONTRACT/GRANT	Not Applica	ble		10. RES	DURCES ES	TINATE	A PROFEM	ONAL MAN YRS	-	HOL (In thousand
A DATES/EPPECTIVE:		EXPIRATION		<u> </u>	PRECEDIN	•	5		1	5
N. NUMBER *				PISCAL	.7T		1 .2			6
G TY <b>PE</b> :		4 ANOUNT:		YEAR	CURRENT		1			
KIND OF AWARD:		f. CUM. AMT.					1		1	
AESPONSIBLE DOD (	DRGANIZATION	L		39, PER  -{	19 <b>1111116 0</b>	RGANIZ/	TICH	L		<b>_</b>
HAME!" US Army	Institute of	E Surgical	Research	HANE:	JS Arn	ny Ir	stitute	e of Surg	ica	l Resear
-				APOPE	Mic	robi	.ology H	Branch	_	~~~ <i>4</i>
Fort	Sam Houston,	Texas 782	34		Fort	: San	Housto	on, Texas	1 7	8234
				PRINCIP	AL INVEST	IGATOR	(Pursiak BRAN	ll U.S. Anadamie :	ne ti ne ti	-
	JAL			-	Robe	ert I	3. Lindh	erg, Phi	)	-
HANG Basil A	. Pruitt. Jr	, COL, MC		TELE	HONE	512-2	221-2018	3		
TELEPHONE.	12-221-2720				L SECURIT	ACCOL	NT NUMBER:			
21. GENERAL USE				A550C14	TE INVEST		•			
FOREIGN INT	TELLIGENCE NO	T CONSIDERE	D	NAME						
W KEYWARDS / Dave - 4	RACH all Samelly Class	fation Codes		NAME						DA
(U) Pseudon	ionas; (U) Ph	age typing;	(U) Burn (	wound	; (U)	тор	icai che	emothera	₽¥;	(u) Huma
23 TECHNICAL OBJECT	TIVE, * 24 APPROACH, 28	. PROGRESS (Furnish	Individual peragraphe to	tentified by	ince .	ot on	uelerenterte ⊢tal fo∵	Health Classifier	ing co	nosocomi
23. (U) <u>P</u> s	eudomonas ae:	ruginosa ha	is snown in	creas	rud b	ocen	LIAI IO:	i produc.	-114 -114	hurn
infontion -	and despite in	ntermittent	errective	cont rimar	TOT'	n euid. Acti	ve of o	ombat me	n. dici	ne and
Intection a	Annaly and a feature of the second se	uch infecti	rous te g b	r mar	y UD]					
patients.	Control of s	windre of r	TTAIN MON	+i+v	can i	t ne	achiev	ea or ao	cume	nted.
patients. only with a	Control of s a precise know	wledge of s	strain iden	tity	can i	t be	achiev	ea or ao	cume	nted.
patients. only with a	Control of s	wledge of s	plus modif	tity icati	can i .ons t	t be o es	achiev tablish	an iden	cume tifi	nted. .cation
patients. only with a 24. (U) Ph system in	Control of s a precise known age typing p the non-typab	wledge of s rocedures, le strains	plus modif were appli	tity icati ed to	can i ons t all	t be o es Pseu	achiev tablish đomonas	ed or do an iden isolate	cume tifi s.	nted. cation
patients. only with a 24. (U) Ph system in t	Control of s a precise kno nage typing p the non-typab	wledge of s rocedures, le strains	plus modif were appli	tity icati ed to	can i ons t all	t be o es Pseu	achiev tablish đomonas	ed or do an iden isolate	cume tifi s.	nted.
patients. only with a 24. (U) Ph system in the 25. (U) 75	Control of s a precise kno nage typing p the non-typab 5 07 - 76 09	wledge of s rocedures, le strains Strains ty	plus modif were appli yped with R	tity icati ed to TD ma	can i ons t all ude up	t be o es Pseu 26%	achiev tablish domonas of iso	ed or do an iden isolate lates; 7	cume tifi s. 4% c	nted. cation could be
patients. only with a 24. (U) Ph system in a 25. (U) 75 differentia	Control of s a precise kno nage typing p the non-typab 5 07 - 76 09 ated by lysis	wledge of s rocedures, le strains Strains ty with undi	plus modif were appli yped with R luted phage	tity icati ed to TD ma	can i ons t all de up s in l	c be o es Pseu 26% 974,	achiev tablish domonas of iso four e	an iden isolate lates; 7 pidemic	tifi s. 4% c type	ented. cation could be
patients. only with a 24. (U) Ph system in the 25. (U) 75 differentia predominat.	Control of s a precise kno nage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe	wledge of s rocedures, le strains Strains ty with undi ent periods	plus modif were appli yped with R luted phage in 1975, '	tity icati ed to TD ma . As	can i ons t all ide up s in l ach on	t be o es Pseu 26% 974, e wa	achiev tablish domonas of iso four e s a new	an iden isolate lates; 7 pidemic typen	tifi s. 4% c type ot p	cation could be sources
patients. only with a 24. (U) Ph system in the 25. (U) 7 differentia predominat. in 1974.	Control of s a precise kno hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy	wledge of s rocedures, le strains Strains t with undi nt periods stem deling	plus modif were appli yped with R luted phage in 1975, ' eated epide	tity icati ed to TD ma . As out ea emic p	can i ons t all de up s in l ach on patter	t be o es Pseu 26% 974, e wa ns a	achiev tablish domonas of iso four e s a new nd perm	an iden isolate lates; 7 pidemic typen itted pr	tifi s. 4% c type ot <u>r</u> ecis	cation could be sourcesent se
patients. only with a 24. (U) Ph system in the 25. (U) 75 differentia predominate in 1974. recognition	Control of s a precise kno hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent	wledge of s rocedures, le strains Strains ty with undi ent periods stem deline strains with	plus modif were appli yped with R luted phage in 1975, ' eated epide hich were s	tity icati ed to TD ma . As out ea emic p cought	can i ons t all de up s in 1 ach on patter for	t be o es Pseu 26% 974, e wa ns a furt	achiev tablish domonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot I ecis as t	cation could be present se co the
patients. only with a 24. (U) Ph system in the 25. (U) 75 differentia predomination in 1974. recognition relative r	Control of s a precise kno hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains t with undi stat periods stem delin strains wi	plus modif were appli yped with R luted phage in 1975, ' eated epide hich were s an pathogen	tity icati ed to TD ma . As out ea emic p cought aesis	can i ons t o all de up s in l nch on patter for	t be o es Pseu 26% 974, e wa ns a furt	achiev tablish domonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot p ecis as t	nted. cation could be s present se co the
patients. only with a 24. (U) Ph system in the 25. (U) 74 differentia predominat. in 1974. recognition relative re	Control of s a precise known age typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole on virule	wledge of s rocedures, le strains Strains ty with undi strains ty stem deline strains we ence in hum	plus modif were appli yped with R luted phage in 1975, ') eated epide hich were s an pathogen	tity icati ed to TD ma . As out ea mic p sought mesis	can i ons t all de up s in l ach on patter t for	t be o es Pseu 26% 974, e wa ns a furt	achiev tablish domonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot p ecis as t	cation could be s present se to the
patients. only with a 24. (U) Ph system in a 25. (U) 74 differentia predomination recognition relative r	Control of s a precise known hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains ty with undi strains ty stem deline strains with ence in hum	plus modif were appli yped with R luted phage in 1975, ') eated epide hich were s an pathogen	tity icati ed to TD ma . As out ea emic p cought mesis	can i ons t all de up s in 1 ach on patter t for	t be o es Pseu 26% 974, e wa ns a furt	achiev tablish đomonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot p ecis as t	cation could be s present se co the
patients. only with a 24. (U) Ph system in a 25. (U) 7 differentia predominat. in 1974. recognition relative r	Control of s a precise known hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains ty with undi stem deline stem deline strains with ence in hum	plus modif were appli yped with R luted phage in 1975, ') eated epide hich were s an pathogen	tity icati ed to TD ma a. As out ea mic p sought mesis	can i ons t all de up in 1 ach on patter for	t be o es Pseu 26% 974, e wa furt	achiev tablish đomonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot <u>p</u> ecis as t	cation could be s present se co the
patients. only with a 24. (U) Pl system in the 25. (U) 7 differentia predomination recognition relative r	Control of s a precise known hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains t with undi nt periods stem delin stem delin strains wi ence in hum	plus modif were appli yped with R luted phage in 1975, ') eated epije hich were s an pathogen	tity icati ed to TD ma . As out ea emic p sought mesis	can i ons t all de up s in l ach on patter for	t be o es Pseu 26% 974, e wa furt	achiev tablish đomonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot I ecis as t	cation could be s present se co the
patients. only with a 24. (U) Pl system in the 25. (U) 7 differentia predomination recognition relative relative relativ	Control of s a precise kno hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains t with undil nt periods stem deline stem deline strains with ence in hum	plus modif were appli yped with R luted phage in 1975, ' eated epide hich were s an pathogen	tity icati ed to TD ma . As out ea emic p sought mesis	can i ons t all de up s in l ach on patter t for	t be o es Pseu 26% 974, 974, ae wa ns a furt	achiev tablish đomonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot I ecis as t	cation could be so present se to the
patients. only with a 24. (U) Pl system in the 25. (U) 7 differentia predominat. in 1974. recognition relative r	Control of s a precise kno hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains t with undi: nt periods stem delin stem delin strains with ence in hum	plus modif were appli yped with R luted phage in 1975, ' eated epide hich were s an pathogen	tity icati ed to TD ma . As out ea emic p cought mesis	can i ons t all de up s in l ach on patter tor	t be o es Pseu 26% 974, e wa ns a furt	achiev tablish domonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot I ecis as t	nted. cation could be so present se to the
patients. only with a 24. (U) Pl system in to 25. (U) 7 differentia predominat. in 1974. recognition relative r	Control of s a precise known hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains ty with undi: nt periods stem deling stem deling ence in hum	plus modif were appli yped with R luted phage in 1975, ' eated epide hich were s an pathogen	tity icati ed to TD ma . As out ea emic p cought nesis	can i ons t all de up s in l hch om patter t for	t be o es Pseu 26% 974, e wa furt	achiev tablish domonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot I ecis as t	cation could be soresent se to the
patients. only with a 24. (U) Pl system in a 25. (U) 7 differentia predominat. in 1974. recognition relative r	Control of s a precise known hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole of virule	wledge of s rocedures, le strains Strains t with undi stim periods stem deline strains we ence in hum	plus modif were appli yped with R luted phage in 1975, ' eated epide hich were s an pathogen	tity icati ed to TD ma . As out ea emic p cought nesis	can i ons t all de up s in l ach on patter tor	t be o es Pseu 26% 974, e wa ns a furt	achiev tablish domonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot I ecis as t	cation could be present se to the
patients. only with a 24. (U) Pl system in 25. (U) 7 differentia predominat. in 1974. recognition relative r	Control of s a precise known hage typing p the non-typab 5 07 - 76 09 ated by lysis ed at differe The typing sy n of virulent ole on virule	wledge of s rocedures, le strains Strains ty with undi is with undi strains ty stem deline strains wi ence in hum	plus modif were appli yped with R luted phage in 1975, ' eated epide hich were s an pathogen	tity icati ed to TD ma out ea emic p cought nesis	can i ons t all de up s in l ach on patter t for	t be o es Pseu 26% 974, 974, e wa ns a furt	achiev tablish domonas of iso four e s a new nd perm her eva	an iden isolate lates; 7 pidemic typen itted pr luation	tifi s. 4% c type ot I ecis as t	cation could be soresent se to the

DD, LAR 61498 PREVIOUS EDITIONS OF THIS FORM ARS OBSOLETE DD FORMS 1488A 1 NOV 68 AND 1488 1 1 MAR 68 (FOR ARMY USE) ARE OBSOLETE

#### TERMINATION

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE - SURGERY

REPORT TITLE: BACTERIOPHAGE TYPES OF PSEUDOMONAS AERUGINOSA FOUND IN BURNED SOLDIERS

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROGKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 1 July 1975 - 30 June 1976

Investigators:

Ruth L. Latta, BS Robert B. Lindberg, PhD Arthur D. Mason, Jr, MD Basil A. Pruitt, Jr, MD, Colonel, MC

Reports Control Symbol MEDDH-288 (R1)

UNCLASSIFIED

#### ABSTRACT

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE - SURGERY REPORT TITLE: BACTERIOPHAGE TYPES OF PSEUDOMONAS AERUGINOSA FOUND IN BURNED SOLDIERS
US Army institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234
Period covered in this report: 1 July 1975 - 30 June 1976
Investigators: Ruth L. Latta, BA Robert B. Lindberg, PhD Arthur D. Mason, Jr, MD Basil A. Pruitt, Jr, MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

Continued application of dilute and concentrated bacteriophages of the ISR <u>Pseudomonas aeruginosa</u> phage typing system has made possible the characterization of 95% of isolates from burn patients in 1975. <u>Ps</u> <u>aeruginosa</u> infection involved increasing numbers of patients annually for the past 3 years. In 1975, 1561 strains were collected from 161 patients. Type NT-16, (21, 68, F7) was the most prominent epidemic type, continuing from 1974, when it appeared abruptly in July. The second most important type, E-2, (24) had previously been seen only as a rare type. Types predominant in septicemia and in lung tissue at autopsy included strains typical of the overall type distribution and also strains which were relatively more common in blood and in lung tissue than in the Pseudomonas population as a whole. Epidemic episodes extended over periods from 8 weeks to 2 years, depending on the type causing infection.

Pseudomonas Phage typing Burn wound Topical chemotherapy Humans

# BACTERIOPHAGE TYPES OF PSEUDOMONAS AERUGINOSA FOUND IN BURNED SOLDIERS

<u>Pseudomonas aeruginosa</u> has shown a progressive capacity for extension as a nosocomial pathogen in a variety of disease states, and on a worldwide basis. This extending involvement is particularly disturbing in view of the concomittant development of antimicrobial agent resistance of <u>Ps aeruginosa</u>; it has become increasingly apparent 'hat individual successes have not altered the potential gravity of Pseudomonas infections in hospitals.

Burn wound infections are one of the most serious problems caused by Ps aeruginosa, and their role in pathogenesis of infection during burn treatment has remained prominent despite development of effective topical therapeutic and systemic agents (1,2). A burn ward such as that of the US Army Institute of Surgical Research offers significant advantage in studying the Pseudomonas problem: The proportion of patients with severe burns, at high risk of developing invasive sepsis, is large, and proximity presents a dismayingly optimal circumstance for transmission, so that strains capable of setting up an epidemic situation are offered the opportunity to do so. Precise recognition of bacterial strains of an infecting species is essential if the epidemiology of an ongoing infection problem is to be assessed. This differentiation not only permits precise characterization of infections that occur, but makes possible the selection of strains which vary in virulence, invasiveness, and esponse to treatment. Phage typing was selected as the method of choice for this differentiation (3,4). and a definitive set of phage for this procedure was developed in this laboratory and has been in use for 15 years. The phages have been furnished to other workers throughout the world, in England, Denmark, Japan, Germany, and Australia, and the value of the system has now prompted the Center for Disease Control (Atlanta) to request a set of phages to be used as an international definitive typing set.

3. Lindberg RB, Latta RL, Brame RE, Moncrief JA: Definitive bacteriophage typing system for <u>Pseudomonas aeruginosa</u>. Bact Proc 1964, p.81.

4. Lindberg RB, Latta RL: Phage typing of <u>Pseudomonas aeruginosa</u>: Clinical and epidemiologic considerations. J Inf Dis (Suppl) 130:S33-S42, 1974.

^{1.} Lindberg RB, Moncrief JA, Switzer WE, Order SE, Mills W, Jr: The successful control of burn wound sepsis. J Trauma 5:601-616, 1965.

^{2.} Curreri PW, Lindberg RB, DiVincenti FC, Pruitt BA, Jr: Intravenous administration of carbenicillin for septicemia due to <u>Pseudomonas</u> <u>aeruginosa</u> following thermal injury. J Inf Dis (Suppl) 122:S40-S43, 1970.

#### METHOD

Previous reports have presented the recognition of the ongoing problem of the appearance of strains of <u>Ps aeruginosa</u> which did not react with the Routine Test Dilution of some bacteriophages. These non-typable (NT) strains react with undiluted typing fluids, and to further the objective of reproducibility and continuity of results, undiluted typing was performed on strains NT at RTD. The prefix "NT" has continued to be used as the designation for the type of such strains, with no evaluation made for individual strains as to whether or not such lysis constitutes a pyocin reaction as opposed to a phage lysis.

#### Phage typing of Pseudomonas aeruginosa from Burn Ward Patients, 1975

The number of patients from whom <u>Ps aeruginosa</u> was recovered has increased in recent years as shown in Table 1.

Year	No Patients admitted	No Patients from whom Ps aeruginosa recovered	No. Strains Ps aeruginosa Collected for Phage Typing		
1973	256	97	788		
1974	244	116	1085		
1975	245	161	1561		

Table I	1.	Pseudo	monas	aerug	inusa	recoveri	ies
		in ISR	, 1973	- 19)	75		

During 1975, 1561 strains of <u>Ps aeruginosa</u> from 161 patients on the burn wards were collected for phage type identification. Sources included blood, wound surface, biopsy, catheter, sputum, urine and other sources.

Predominent phage types of strains typable either at RTD or with undiluted phage, are shown in Table 2. The Phage Type Code (left column) shows the assigned letter-number designation which has been developed to facilitate characterizing any given phage sensitivity pattern. Recall that "NT" designates strains not typable with dilute, but typable with concentrated phage. The phage type code lists the phages reacting; the NT designation shows that the reaction occurs only with undiluted phage. These patterns are marked with an asterisk (*). The proportions of patients harboring one of the principal types is greater than 100%, because of multiple types being found on one patient. In terms of proportion of strains belonging to a given type, the 9 patterns which were distinguished included 84.7% of all strains tested. The remaining 15.3% of strains were distributed over a large number of minor types. Any types found in fewer than 8 persons were not included; this practice prevents the inclusion of an unwieldy number of strains of minor numerical importance.

Phage Type	Phage Type	Patients	Strains	
Code		Per Cent		
NT-16	Non-typeable * (21),68,(F7)	36.6	20.0	
E 2	24	33.5	17.0	
NT-19	Non-typeable * (21),24,44,1214,68	30.4	11.8	
NT-22	Non-typeable * 44,1214,68,(109),352,F8	29.2	18.0	
H 1	1214	14.9	7.4	
NT-24	Non-typeable * (31),68	11.2	5+3	
E 2 ^V	24 or Non-typeable * (24)	10.5	2.4	
NT-3	Non-typeable * (21),(24),(44),1214,(68),109,352,(F7),F8	4.9	2.0	
NT-23	Non-typeable ★ (21),68	4.9	0.9	

Table 2.	Predominant	Phage Types of 1,561	Pseudomonas Strains
	from 161	ISR Burn Ward Patien	ts, 1975

. * Phage Type using undiluted phage

() Variable reaction

Nine separate types of <u>Ps</u> <u>aeruginosa</u> played major roles in burn wound infection during 1975. Three types, representing 58.9 of the strains listed, reacted at the RTD; the remaining 6 types were recognized only in undiluted (NT) reaction. In the case of E-2 and E-2^V, two distinct levels of ractivity were observed with these two strains. They are entered as two distinctive types in view of morphologic differences in reaction. E-2^V strains fluctuated in their reaction with phage type 24, while E-2 strains were consistent in level of sensitivity and in exhibiting a characteric in plaque reaction in RTD typing.

The most prevalent strain of <u>Ps aeruginosa</u> 1975 was NT-16, reacting with 21, 68, F7 phages. This was a relatively new type, first observed

in July 1974. It rapidly occupied a predominant position on burn patients and in 1975, was the most pervasive strain, on 36.6% of the patients positive for Pseudomonas, and with one-fifth of all isolates typed.

Phage type E-2, reacting with phage 24, was the second most common type in 1975. This type had been observed as a rare type in earlier years, although at least since 1970 it had never been common enough to merit inclusion in types recovered. None had been observed during 1974.

Type NT-19, the third most frequent type, involved a high proportion of patients (30.4% of all those with Pseudomonas present), but presented fewer strains than did the next most common type. This too was a strain which increased strikingly from almost zero incidence; only two strains were collected in late 1974.

Type NT-22 was not seen at all in 1974. It appeared abruptly in July 1975, spread very rapidly, and became fourth in incidence for the year.

The first four categories included 66.8% of all strains collected, and thus embodied the major part of the epidemic spectrum of <u>Ps aeruginosa</u> for 1975.

'Type 12-14, or H-1, which had been observed in earlier years but not in 1974, first appeared in May 1975. 14.9% of patients harbored it, and 7.4% of all strains collected were of this type.

Type NT-24 was a new type, never previously observed. It appeared in September 1975, and thus spr_ad very rapidly to colonize the ward as the sixth most common strain for the year. The pattern is related to 31 and to 68, but these reactions were never previously linked.

Fnage type  $E-2^{V}$  was described above. It reached 10.5% of patients, and was a new previously unrecorded type.

The last strains recorded were NT-3 and NT-23, each was found in 4.9% of the patients. NT-3 was more common amoung total strains but still made up only 2% of isolates typed. In the previous year, 1974, NT-3 was found in 32.75% of patients, but declined rapidly in incidence in 1975. NT-23, (21) 68, was another new type, never previously observed.

Thus, there were 9 types predominant in 1974, of which 7 had not been seen or were of negligible incidence in 1974. This was an unusually high rate of incursion of new types into the ISR burn ward, but showed how rapdily new strains can pervade a burn population such as theo ne presented here. 84.8% of all strains fell in one of the 9 types listed, and 66.8% of all strains tested in 1975 were NT in contrast to 75% in 1974. In that year, the predominant population included 12 types; the pattern in 1975 was consistent with more permeation by individual epidemic strains. The validity of the typing system for characterizing Pseudomonas strains was re-affirmed, and the undiluted typing addition appeared to be valid.

# Phage Types of Pseudomonas Recovered in Blood of ISR Burn Patients, 1975

Aside from general seeding of most burn patients, the role of Pseudomonas in sepsis is of great importance, since such septicemia displays a very high mortality rate, despite use of antibiotics which may be effective in vitro. Forty-five burn patients with positive blood cultures contributed 91 strains for typing. Phage types observed are listed in Table 3. Note that the most common phage types, among the total of strains collected, are represented among the five most common blood stream types. In terms of totals of patients from whom these types were recovered, NT-16 was more common in the total than in blood stream isolates, by 36.6% to 15.5%. Similarly, NT-19 was recovered from 30.4% of all patients, but only from 4.4% of blood culture positives. There were no strains which were more common in blood than in the population at lare. The pattern of sepsis, however, in relation to chronologic sequence, is better shown by detailed presentation, as shown in Table 4. it was obvious that the successive episodes of septicemia were not scattered; instead, the various types were clumped within given time periods. reflecting the epidemic nature of the episode the patients are represented by arbitrary accession numbers. Those patients who had a given type recovered in successive months are marked with an asterïsk. The principal 5 phage type codes are listed, plus a column for those types occurring only in one or two patients.

Phage Type Code	No. of Patients	No. of Strains
NT-22	16	25
E 2	15	31
NT-16	7	13
E 2∛	4	4
н 1	3	6
822 NT-17 NT-19	2	3
NT-3 NT-23 NT-	1	1

			Table	e 3.			
Phage	Types	of	Pseudomonas	Strains	from	Blood	Stream
	of 4	5 15	R Burn Ward	Patients	s. 19 ⁴	75	

のないでは、「ない」となった。

į		_			Phag	e i	Type C	ode	
Month	Patient	No. of	Ε2	E 2 ^V	H 1	1	NT-16	NT-22	Other
	No.	Strains			No.	01	f Stra	ins	·····
	222 174					-			NT2-1
	240 174	3					3		
Jan	242 .74	í					í		
	3	2							NT17-2
	4	1				1			NT17-1
	6	1				Т	-1		
Feb	23	1							NT19-1
	2/	3	2				1		NT 1
						_			
	75	2	2						
Mar	41	í	1						822-1
	48	1	1						
	46	3			T	Т	2		
	53	2	2						
	56								
	5/								
Apr	63		1				4		
	68	i	17	ł					
	69	3	3					]	
	79	4	2	1			1		
	80		<u>[]</u>					L	
	*79	3				Т	3		
Hay	*50		1 !		1				
	09			1,					
lup	62	<u> </u>						L	
2011	172	<u> </u>	<u>+</u>	+	-				
Jul	137	i		1 '				1	
	1 144	3	Î		13	T		1	<u> </u>
	151	i			1				
Aug	156	2	1		2	1		ł	
	163	2	1						NT19-2
	1 108	<u> </u>	<u></u>						
	*163	4	1					1	822-2 NT23-1
Sep	176	2						2	
	179	1 ī	1					1	
	186	1						<u> </u>	
	+151	2	T					2	
Oct	189	?	1					?	
	193								
<u></u>	1 +104	+	<del></del>					+	+
	204								
Nov	207	i						l i	
	208	4						4	
	218				_			<u> </u>	I
Dec	221	1						1	
	227	1 1	1					1 1	

Table 4. Phage Types of 91 <u>Pseudomonas</u> Strains from Blood Stream of ISR Burn Ward Patients, 1975

Type NT-22 was the most frequent septicemic strain occurring from July through December. The strain was fourth in frequency from all sources. Type E-2 (phage 24) was the second most frequently observered, and involved 15 septicemic patients from February through May. Third in incidence was type NT-16, which was the most common strain recovered from all sources, its incidence, from January through May, paralleled that of type E-2. From April through July, the related type  $E-2^{V}$  was recovered in 4 patients. The implication that  $E-2^{V}$  was a loss variant of E-2 is strongly suggested by this sequence. Type H-1, found in 3 patients, appeared only in August, in blood cultures. The six least common types, B-22, NT-17, NT-19, NT-3, NT-23 and a single non-reactive strain, were found from January to March, then in August and September. Only one, NT-17, appeared in two cases in close succession.

There were, thus, 91 strains of 11 phage types recovered from blood. No apparent correlation of phage type (or strain) with septicemia was apparent. Instead, the invasive behavior was roughly a reflection of the overall incidence of <u>Ps aeruginosa</u> in the population.

# <u>Pseudomonas Phage Types Recovered From Lung Tissue of ISR Burn Patients</u> <u>at Post-Mortem, 1975</u>

Pseudomonas strains from lung tissue at autopsy furnish one of the principal sources of definitive information about the pneumonia which so frequently occurs as part of the pathologic problem in the burn patient. Phage types of these strains, in comparison with septicemic strains, present an insight into the source of systemic invasion in these patients. Phage types of Pseudomonas from lung tissue are summarized in Table 5. The frequency of occurrence of types was rot entirely that seen in the overall population nor was it identical with that of blood stream isolates. Type E-2 was far less common than it was in blood or wound type NT-22 was most common in both blood and lung, but was less common on wounds and other sites.

Comparative frequency of occurrence of principal phage types in blood, and lung and in the entire collection from all sources is summarized in Table 6.

As with septicemias, a more adequate picture of epidemiology is seen when the chronologic sequence of type incidence is shown. This pattern is presented in Table 7. The mos common phage types are shown individually; additional infrequent types are shown in the month of their occurrence.

The most common epidemic type, NT-22, was found in lungs of 16 patients between August and December. Type NT-16, found in 13 patients, was recovered in 13 patients from January through July. These two strains were in succession predominant for the entire year. Type NT-19, the third most frequently seen, was found in 9 patients from January through August, and then once more in December. NT-24 was recovered from lungs of 5 patients from August through November. The less frequent types included E-2,

Phage Type Code	No. of Patients	No. of Strains
NT-22	16	30
NT-16	13	25
NT-19	9	19
NT-24	5	9
E 2	3	7
H 1 NT-3 NT-17	2	7 2 5
Other	10	14

Table 5. Phage Types of <u>Pseudomonas</u> Strains from Lung Tissues of ISR Burn Ward Patients, 1975

no. of patients: 41

no. of specimens: 102

no. of strains isolated: 118

Table 6. Relative frequency of Occurrence of <u>Ps aeruginosa</u> Phage Types, ISR 1975

Туре	Frequenc	y & Sourc	9
	All Sources	Blood	Lung
NT-16	1	3	2
E-2	2	2	5
NT-19	3	6	3
NT-22	4	1	1
H-1	5	5	6
NT-24	6	Ó	4
E-2 [∨]	7	4	Q
NT-3	8	7	6
NT-23	9	7	0

			Phage Type Code						
Honth	Patient No.	No. of Specimens	E 2	H 1	NT-16	KT-19	NT-22	NT-24	Other
					N	o. of S			
	223 •74	3			1	2			041-1 NT3-1
120	240 174	3	ĺ		3		ĺ		
Jen	3	4			3	3	Į		NT17-1
	15	1				<u> </u>			F 5-1
	6	1					[		
Feb	25	4	1		1	2	ļ		
:	27	3				2	i i		NT -2
Mag	28								NI -2
nar	<u> </u>	4							1 1 7 2 1
Apr	53					]			W1 J=1
	57	2	L		2	L			L
	79	3	1		3				
May	83	1			1	1			
	85	1			1	L	<u> </u>		
Jun	109	2	2	L			مر ماریندا دارند.		NT -1
Ju1	137	4		4	2	<u> </u>			
Aug	144	4 4		3	1	<u>ь</u> .	I		NT23-3
Aug	166	3				2	2		G28-1
	163	4	T				4	1	
Sep	168	3					3		
	183	í	i i				l í		
	186	2	<u> </u>				2		
	151						1	2	
{	190						1 1	1.	
Oct	197	4					3		C23-1
	199	3					2		NT -1
Į	202	!					1	1	
	205		1				$\frac{1}{1}$	+	+
Nov	207	1						1	1
L	214	3	1				1	3	<u> </u>
	221	2						1	
Uec	240					1	5	1	
		· · · · · · · · · · · · · · · · · · ·							

# Table 7. Phage Types of 118 <u>Pseudomonas</u> Strains from Lung Tissues of ISR Burn Ward Patients, 1975

Note:

;

found in lungs of 3 patients in February, May, and June. Two patients each exhibited H-1, NT-3 and NT-17, and 10 additional types were each found in only one patient. Obviously, there was no one type causing pneumonia, but there were significant differences in septicemia and pulmonary phage types. The NT-16 epidemic period for both sources was the same, from January to May or July. Type NT-22 alsc appeared at the same time interval for both sources. But the E-2 type found extensively in blood cultures from February through May was seen in lungs of only 3 patients during this period. Type NT-19, prominent from January to August, was found in the blood of only two patients. Thus a major episode of bacteremia was caused by a type rare in the lung, while a pulmonary strain of extensive occurrence was rare in the blood.

An attempt to assess differences in incidence of phage types by location was made in view of the differences in incidence thus far noted. Figure 1 compares the incidence of strains from blood, lung tissue, and all other sources. The proportionate incidence of the major types is represented for each source by the distinctive bar form. Type NT-16 was most common in lung and all other sources, but far less often recovered from blood. Type E-2 was found equally often in blood and all other sources, but was relatively uncommon in the lung. NT-19 was relatively rare in blood, but almost as frequent in lung as in other sources. Type NT-22 was most common in lung, least frequent in sources other than blood. The remaining principal types, H-1, NT-24, E-2^V, NT-3 and NT-23, were all less common than the first 4. H-1 was most common in other sources, least common in blood. Type NT-24, surprisingly, was never found in blood culture. Conversely,  $E-2^{V}$  was not found in lung. NT-3 was least often found in blood, but both NT-3 and NT-23 were the least common of the major types recognized.

#### Monthly Distribution of Prevalent Phage Types, ISR, 1975

The occurrence of small monotype epidemics of <u>Ps aeruginosa</u> has been documented in previous years, but the continued change in types, and the appearance of previously unrecorded types makes this observation of succession of types a continuing important assessment. Figure 2 summarizes the monthly incidence of 11 major phage types of <u>Ps aeruginosa</u>. All but two strains involved at least 8 patients. Two strains involved only 3 patients each. To render the distribution more legible, the most prevalent type for each month is outlined in solid black, the next most prevalent by a double line, and third most common by a single line.

The first 5 types listed, NT-17, NT-18, NT-3, NT-19 and NT-16 were types that had been seen in 1974. In that year, NT-18 was 10th in frequency, while NT-3 was the most common type found in 1974. Conversely, only two strains of NT-19 were found in 1974. NT-16, which like NT-19 has been a major type in 1975, was also the seond most common type found in 1974. NT-17 disappeared after its appearance in January. NT-18







similarly was seen only briefly in early 1975; NT-3 disappeared after April 1975.

NT-19 was a prominent feature of the burn flora in 1975; it was recovered every month, although it was never a major type. Type NT-16 was recovered in 11 of 12 months in 1975, and was predominant during 3 months. Its frequency was diminishing during the last quarter of 1975, and it can be projected that it may disappear.

Type E-2 presented an acute 4 months outbreak in March through June, when it was the dominant type. The closely related E-2 was recovered over a 6 month period, but was only once the second most common type.

Type H-1 displayed a capability for rapid dominance of the patient, flora especially in August when it was the predominant type. It persisted over a 7 month period, then disappeared abruptly, as it had entered the burn ward population. NT-22 strains appeared abruptly in July, increased in incidence from 4 to 5 patients, then assumed the predominant status on the burn ward through December. Type NT-23 occurred during a 4 month period starting in August but was never prominent. The last type NT-24, appeared in September, then for the rest of the year was the second most common type. Numerically it would be regarded as an epidemic strain; only the preponderance of NT-22 overshadowed its frequency.

The pattern of major and lesser epidemics of <u>Ps aeruginosa</u> appearing for larger or shorter periods was evident in this period of observation, as it had been in previous years. The second most common epidemic type, NT-19, had appeared in only 2 isolates in 1974; it was conspicuous but never predominant throughout 1975. Type NT-16 had been the second most common type in 1974. Type E-2 typifies the previously unknown type that appeared with explosive suddenness. There is still no explanation for the disparate behavior among strains of <u>Ps aeruginosa</u> in a burn ward.

#### PUBLICATIONS

None

#### PRESENTATIONS

Lindberg RB: "Epidemiology of <u>Pseudomonas aeruginosa</u> strains in nosocomial infections." Presented at Am Pub Health Assoc. Symposium, New Orleans, 1975.

RESEARCH	I AND TECHNOLOG	Y WORK UNIT S	WORK UNIT SUMMARY			DA OC 6396		2 DATE OF SUMMARY 76 10 01		REPORT CONTROL SYMBOL DD-DR&E(AR)636	
A DATE PREV SUMRY	4 KIND OF SUMMARY	S. SUMMARY SCTY	. WORK SECURITY	7. REGR	DING	-	NEB'N INSTR'N OL SPEC		FIC DATA - P LEVEL OF SUI		
75 07 Úl	H. TERM	U	ប	MA		NI		T YES		A. WORK UNIT	
10 NO./CODES P	PROGRAM ELEMENT	PROJECT	NUMBER	TASK AREA NUMBER			WORK UP	IT NUMBER	1		
a. PhimARY	61102A	<u>3A161102</u>	<u>B71R</u>	ļ	01		243				
S. CONTRIBUTING	<u> </u>	<b> </b>		ļ				• • •	• • • •	· · ·	
11 TITLE (Procode with	Sourity Classification Code	) " (II) Bacto	ariophage 7	YDAR	of Se		ia Marc		From	Dutto	
Wounds of M	lilitary Perso	onnel (44)		.ypes					FIOM	burn	
003500 Clin	ical Medicine	1									
13 START DATE		14. ESTIMATED COM	LETION DATE	IS FUNDING AGENCY IS PERFORMANCE METHOD'						HOD	
67 07		76 0	9	DA			L	<u> </u>	In-Hou	ise	
Not Applicable					DUNCES ES		A PROFEM	HONAL MAN Y	RS L PUI	IDE (In thousands)	
h NUMBER *		EAFIRATION:		FIRCAL	76		.	6	1	7	
-		4 ANOUNT:		YEAR	eunhenv		+	3			
& KIND OF AWARD.		I. CUM. AMT.					1				
IS RESPONSIBLE DOD	ORGANIZATION	T	1		PO MINA OI	ANIZ/	ATION	<u> </u>			
NAME." US Army	Institute of	Surgical	Research		US Arm	y Ir	nstitute	of Su	rgical	Research	
ADDRESS:"Fort S	Sam Houston, 1	Exas 7823	4	ADDRES	Mic For	robi t Sa	iology I am Houst	Branch ton, Te	xas 7	/8234	
				PRINCIP	AL INVEST	GATOR	(Fumioh base	If U.S. Acade	nic metinute	a)	
	UAL			NAME	• Robe	rt I	B. Lind	perg, P	hD	•	
NAME: Basil	A. Pruitt, J	COL, MC		TELEPHONE: 512-221-2018							
TELEPHONE: 51	12-221-2720			SOCIA	L SECURITY	ACCOL					
21. GENERAL USE				-	TE INVEST	GATOR	5				
FOREIGN INT	TELLIGENCE NO	CONSIDERE	D	NAME	Virg	inia	a E. Eng	glish,	MA		
N VIVENDA ZANT	BACH -IB Bandle Bland	calles (cal.)		NAME			<u></u>			DA	
(U) Burns;	(U) Serratia;	(U) Bacte	riophage;	(U) H	umans						
23 TECHNICAL OBJEC	TIVE, 24 APPROACH, 21	PROGRESS (Fuminh I	ndividual paragrapha is	leniitted by	number. Pri	tede te	zi ol oach with	ecurity Class	Realien Cod	•)	
23. (U) A	potential ep:	idemic path	ogen of sig	ynifi	cance,	Sei	rratia i	narcess	iens, l	nas been	
observed in	periodic ou	coreaks in	burn patie	nts.	Trans	mis	sion ap	pears t	o be		
relatively	rapid. Since	e sucn oppo	different	an pr	esent	ser:	lous na:	zaros t develor	o seve	erely	
phage tunit	TETS' g Wegu:	a direct m	aior tol	in co	ntrol	SOU	ynt Dy ( such or	neverot out the s	ament ( stic ·	u d Dathogene	
Fugde cybri	iy ayatem, dS	a arreat m	ייד יר אי			UL S	auon op	eor cuit	our j	actiogens.	
24. (U) Ph distinctive	nages, recove e reactions ca	red from se apable of d	wage efflu ifferentia	ent, ting	were p strair	oropa as o	agated f <u>Serra</u>	to deri tia mar	ve hig cesser	gh titer <u>15</u> .	
25. (U) 7 differentia component p by reaction previous ex evaluation	distinctive reactions capable of differentiating strains of <u>Serratia marcessens</u> . 25. (U) 75 07 - 76 09 Although the previously reported phage typing set was capable of differentiating <u>Serratia</u> epidemic strains successfully, instability of a major component phage in the system necessitated further phage isolations and differentiation by reaction patterns. A more definitive typing system evaluated in terms of the previous experience, has resulted, and this improved system is undergoing intensive evaluation to confirm its differentiating capability.										
Paralledie to contract	lore ypon origenator's app 20 PREVIOUS	EDITIONS OF T	115 FORM ARE O	830L E	TE DI	ORMI	5 1498A 1 M	10V 88			
U U I MAR CO THO	AND 1498-	1 1 MAR 68 (FO'I	ARMY USELAR	50 50	ы. <b>6</b> . Т 9 <b>.</b>						

# TERMINATION

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# **REPORT TITLE:** BACTERIOPHAGE TYPES OF SERRATIA MARCESCENS FROM BURN WOUNDS OF MILITARY PERSONNEL

# US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Robert B. Lindberg, PhD Virginia C. English, MA Basil A. Pruitt, Jr, MD, Colonel, MC Arthur D. Mason, Jr, MD

Report Control Symbol MEDDH-288(R1)

UNCLASSIFIED

# ABSTRACT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: BACTERIOPHAGE TYPES OF SERRATIA MARCESCENS FROM BURN WOUNDS OF MILITARY PERSONNEL

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Robert B. Lindberg, PhD Virginia C. English, MA Basil A. Pruitt, Jr, MD, Colonel, MC Arthur D. Mason, Jr, MD

Report Control Symbol MEDDH-288(R1)

The phage typing set for <u>Serratia marcescens</u> has been re-assessed to avoid problems associated with phage viability and the specificity of typing patterns. New phages have been collected to remedy these deficiencies, and a comprehensive typing will be applied to strains of <u>S. marcescens</u> collected for this assessment.

The importance of a technic for differentiating strains of <u>S. marcescens</u> is apparent in view of the capacity of this enteric species to cause severe nosocomial infections, including pneumonia, infections in surgical wounds and ir thermally injured patients. To achieve precise strain recognition, a phage typing system was developed in the ISR laboratory.

The typing set of 7 phages was applied to differentiating Serratia isolates from ISR wards and from other hospitals. The system was effective, and phage types were distinguishable. The technical approach was valid. However, the phages selected included some which were exceptionally labile, and which could fail to survive storage at 4[°]C, the usual methods of holding bacteriophages. Further, the phage type pattern achieved with this typing set, although usable, revealed a coverage of wild strains that was less definitive in making type distinctions than had been desired.

In view of these factors, re-study and acquisition of additional phages has been carried out during this interval. New phages have been recovered, propagated and compared for identity with the existing phage collection. An improved typing set is in process of being selected and standardized. It will be assessed on <u>S. marcescens</u> collections now held for this purpose.

Burns Serratia Bacteriophage Humans

				1. ASENC	TY ACCENSION	12	DATE OF SUN	WE AN	REPORT	CONT	NOL STREOL	
RESEARCH	AND TECHNOLOGY	WORK UNIT S	UMMARY	DA O	E 6382		76 10	01	DD-D	R&E(	AR)636	
& DATE PREV SUM'RY	4 KIND OF SUMMARY	S. SUNMARY SCTY	. WORK SECURITY	7. REGRA	01NG BO	0168		Sh SPECIFIC	DATA	6 LE	VEL OF SUM	
75 07 01	H. TERM	U	IJ	NA		N	L	Z YES		•	WORK UNIT	
10 NG /CODES:"	PROGRAM ELEMENT	PROJECT	NUMBER	TASK A	TASK AREA NUMBER WORK UNIT NUMBER							
& PRIMARY	61102A	3A161102	B71R	01			317					
S. CONTRIBUTING							1809	· · · ·		dini k		
C. CUNTRIBUTING						Ŀ		(***********			-87 (Q) (A)	
11. TITLE (Procode with )	Security Classification Code)	U) Five (U)	Per Cent A	queou	is Sulfa	my	lon Soa	ks Used	in 1	lopi	.cal	
Treatment o	f Burned Sold	iers (44)										
12 SCIENTIFIC AND TEC	CHNOLOGICAL AREAS											
003500 Clin												
3 START DATE [14. ESTIMATED COMPLETION DATE				IL FUNDING AGENCY				16. PERFORMANCE METHOD				
71 10		76 0	9	DA			C. In-		-House			
1). CONTRACT/GRANT Not Applicable				N. RESOURCES ESTIMATE & PROFES			-	HONAL MAN YRS L FUNDS (In these ands)				
& DATES/EFFECTIVE:		EXPIRATION:		ſ	76		.4			16		
N NUMBER:*				FISCAL	<u>71</u>		.1			4		
G TYPE:		4 ANOUNT:		YEAR	CONNERT.							
& KIND OF AWARD.		f. CUM. AMT.										
IS RESPONSIBLE DOD O	RGANIZATION			30. PERP	ORNENG ORGA	HZA	TION					
HANE!" IIS ATMY	Institute of	Surgical	Research	NA4E:* U	JS Army	In	stitute	of Sur	gica	l Re	esearch	
00				Burn Study Branch								
ADDRESS *	Sam Hougton	Toyag 782	34	ADDRESS. [®] Fort Sam Houston, Texas 78234								
FOIL	bain nous con	10,400 ,02										
				PRINCIPAL INVESTIGATOR (Pumich BEAN II U.S. Accounts Institutions)								
RESPONSIBLE INDIVIOU	AL			HAME!" Hugh D. Peterson, COL, MC								
NAME: Basil A	. Pruitt, Jr,	COL, MC		TELEPHONE: 512-221-3301								
TELEPHONE: 512	-221-2720			SOCIAL SECURITY ACCOUNT NUMBER:								
21, GENERAS USE				A550CIA1	E INVESTIGAT	ORS						
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	NAMES	Daryl H	<b>२.</b>	Erickso	n, MD				
				-	Barry A	١.	Levine,	MAJ, N	IC		DA	
IL KEVBORDS (Procede	ACH with Somethy Classifi	tailon Code)										

(U) Burn; (U) Eschar separation; (U) 5% Sulfamylon acetate solution; (U) Humans

23. (U) The objective of this study is to demonstrate the efficacy of 5% Sulfamylon acetate solution as an effective topical antibacterial agent when used either as continuous soaks or wet to dry dressings during the latter stages of eschar separation and when used as continuous soaks covering mesh grafted areas in soldiers with burn injury.

24. (U) Five percent Sulfamylon acetate is used as a debriding agent by applying gauze sponges soaked in a solution to the burn wound and wrapping the area. The gauze sponges are applied saturated with the solution and they are allowed to dry and removed every six to eight hours. The five percent Sulfamylon acetate solution is also used to keep dressings covering mesh graft wet until there is adherence of the mesh graft. This markedly decreases the bacterial flora.

25. (U) 75 07 - 76 09 Five percent Sulfamylon acetate solution was used in 128 patients. In 119 cases it was used with wet to dry applications of coarse mesh gauze to facilitate debridement. In 106 cases it was used in conjunction with mesh grafting. There were two cases of severe respiratory insufficiency among the 128 patients treated giving a complication incidence of 1.6%.

## TERMINATION

PROJECT NO. 3A (61102871R-0), RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

**REPORT TITLE:** FIVE PER CENT AQUEOUS SULFAMYLON SOAKS USED IN TOPICAL TREATMENT OF BURNED SOLDIERS

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 1 July 1975 - 30 September 1976

Investigators:

Hugh D. Peterson, DDS, MD, Colonel, MC Basil A. Pruitt, Jr., MD, Colonel, MC

ALMAN MAN

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

REPORT TITLE: FIVE PER CENT AQUEOUS SULFAMYLON SOAKS USED IN TOPICAL TREATMENT OF BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Armv Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 Jul5 1975 - 30 September 1976

Investigators: Hugh D. Peterson, DDS, MD, Col, MC Basil A. Pruitt, Jr., MD, Col, MC

Reports Control Symbol MEDDH-288(R1)

Five per cent Sulfamylon acetate solution was used in 128 patients as the wetting agent in the wet to dry coarse mesh gauze debridement technique and for continuously wet dressings used to cover excised and grafted areas in the burned soldier. Fifty-two point two per cent of all admissions had the agent used at least for one of the above mentioned techniques. Most of these patients (84 per cent) had the five per cent Sulfamylon acetate solution used for both treatment modalities. Toxic reactions manifested by pulmonary and/or cerebral signs were clinically diagnosed as being related to the use of 5% Sulfamylon acetate solution in two cases. Five per cent aqueous Sulfamylon soaks offer excellent control of the bacterial population of either separating eschar or healthy granulation tissue.

Burn Eschar separation 5% Sulfamylon acetate solution Humans.

# FIVE PER CENT AQUEOUS SULFAMYLON SOAKS USED IN TOPICAL TREATMENT OF BURNED SOLDIERS

Five per cent Sulfamylon acetate solution was used in 128 patients or 52.2 per cent of all admissions. It was used on 119 patients as the fluid agent in wet-to-dry coarse mesh gauze dressings used for wound debridement. It was used as an antibacterial wetting agent for continuous wet dressings covering meshed (graft) or excised areas in 106 patients. Most of these patients (84 per cent) had the five per cent Sulfamylon acetate solution used for both treatment modalities.

Toxic reactions manifested by pulmonary and/or cerebral signs were clinically diagnosed as being related to the use of five per cent Sulfamylon acetate solution in two cases. The pulmonary problems continued to be principally hyperventilation with associated mental confusion in some patients, resolving with removal of the topical soaks.

Since five per cent Sulfamylon solution is generally used 2-3 weeks post burn except in the case of excisions, it appears that when not combined with Sulfamylon cream the patients tolerate it extremely well.

Five per cent aqueous Sulfamylon soaks offer excellent control of the bacterial population of either separating eschar or healthy granulation tissue, with minimal toxicity.

PRESENTATIONS AND/OR PUBLICATIONS

None.

RESEARCH	AND TECHNOLOGY	Y WOLK UNIT S	UMMARY	DA (	CT ACCESS OB 698	2	76 10	01	DD-DR&E(AR)636			
. BATE PREV SUNPRY	4. KIND OF SUMMARY	S. SUMMARY SCTY	L WORK SECURITY	7. REOR	ADING	-	"H MSTA'H	DA SPECIFIC D	DATA . D. LEVEL OF SUM			
75 07 01	H. TERM	U	U	NA		NL		VES C	HO	A. WORK UNIT		
IO. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TABK /		BER		WORK UNIT	NUMBER	2		
	61102A	3A161102	B71R	0	1		223					
S. CONTRIBUTING							XXXX					
C. CONTRIBUTING		L		<u> </u>				i de la contrata de l				
1. TITLE (Procedo with .	formity Classification Code	(U) Devel	opment of P	rophy	ylacti	C 1	opical	Therapy	For	Use on		
Burn wounds	or Military	Patients:	search for	Impro	oved F	ormu	lations	(44)				
	icel Modicine											
15 START DATE	Ical Medicine	IL ESTIMATED COM				:¥		14. PERFORMA		NOD		
65 06		76.09			1		1	C The	House			
T. CONTRACT/BRANT	Not Applicab	10 03		10. 868					-nous			
A DATES FFECTIVE:	Noc Apprican	EXPIRATION:			PREZE		6			20		
				FISCAL	70		.0		1 1	.u		
& TYPE:		4 ABOUNT:		YEAR	CURRENT		<b>*</b> *-		+	- <u>-</u>		
. KND OF AWARD		f. CUM. AMT.			]							
	MEANIZATION			30. PER	CAMING OF		TION					
HANE!" TIS Armer	Institute of	Surgical	Regearch	-	IS Arm	w Tr	stitute	of Sur	rical	Research		
NAME? US Army Institute of Surgical Research					Min Min	11 Y		ranch	JICAI	. Nesearch		
ADONESI" Fort Cam Houston Meyse 78334					ADDRESS!" Fort Sam Houston, Texas 78234							
Fort Cam Houston, Texas 78234					FOIL Sall Houseon, Texas 76234							
				PRINCIP	AL INVEST	GATOR	(Punish BEAN I	U.S. Academic )	me i i nu <b>iden</b>	ŧ)		
	AL			NAME	<ul> <li>Robe</li> </ul>	ert E	3. Lindb	ery, Phi	)			
HANE Basil A	. Pruitt, Jr,	COL, MC		терен	HONE: 51	.2-22	21-2018					
TELEPHONE: 512-221-2720					L SECURITY	ACCOU	NT NUMBER-					
TELEPRONE, CAR ALL ALL ALL ALL ALL ALL ALL ALL ALL A						GATOR	1					
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	NAME								
				NAME:					-	DA		
EL XEVEORDS (Presede	EACH with Josufity Classifi	sallen Code)						_	4-			
(U) Burns;	(U) Sulfamylo	n; (U) Sem	i-synthetic	: pen	icilli	.ns;	(U) Pse	udomonas	3; (U	) Rats		
RB. TECHNICAL OBJECT	IVE," 24 APPROACH, JL	PROGRESS (Pumish i	ndividual paragraphe Id	millied by	number pri	tede iez	i ol osch with S	scurity Classifies	illen Code	• )		
23. (U) As	sessment of v	virulence o	f new epide	mic	types	or F	seudomo	<u>nas aeri</u>	iginc	osa .		
has been ca	lled for by a	ppearance	in the burr	war	a or p	revi	ously u	inknown s	strai	ins varying		
in virulenc	e and in resp	onse to ch	emotherapy.	Th	18 <b>a</b> ss	essi	nent and	evaluat	tion	or new		
antimicrobi	al agents, an	essential	aspect in	prep	aring	tor	the man	agement	ort	nermal		
injury in t	roops in comb	Dat.										
<b>A</b> (m) =			human ta J	n-1	án mi-		00 8000	domont -	.f	ach		
24. (U) Su	rveillance of	infecting	types to I	.nciù	ue VII	ure: .ure:	ice asse	sament (	JI ea	ach		
epidemic fo	um and experi	mental the	rapy on cha	itten	yeu an	TTUR						
ar (m) 75	07 76 00	Desidente e	trains		in wir		oo from	158 +0	909	in seeded		
25. (0) /5	0/ - /6 09	Epidemic s	Liains vary	/ing	111 VII	domi		. 136 LU	900 milor	in seeded		
burned rats	were uncover	rea. Type	A-/1, a rec	curre	nt epi	t voi	to type,	, was vi.	ru ir	lt itt		
9.9% to 100	t of animals.	In its c	ompiete ioi	.m, 1	t was	tree	timent I	erracto	cy II Sreig	i animais.		
Pirbenicill	in was tested	I for contr	of or purn	seps	19. 8	æsu.	ts indi	.cated pa	11 110	11		
effectivene	<b>88.</b>											
A valiable to contract	ure upon originator's appr							inen äidelus Dierreten				

# TERMINATION

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: DEVELOPMENT OF PROPHYLACTIC TOPICAL THERAPY FOR USE ON BURN WOUNDS OF MILITARY PATIENTS: THE SEARCH FOR NEW FORMULATIONS

# US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

のいたいなかからわられば

The second s

Investigators:

Robert B. Lindberg, PhD Virginia C. English, MA Ruth L. Latta, BS George T. Daye, MA Basil A. Pruitt, Jr, MD, Colonel, MC

Report Control Symbol MEDDH-288 (R1)

UNCLASSIFIED

# ABSTRACT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: DEVELOPMENT OF PROPHYLACTIC TOPICAL THERAPY FOR USE ON BURN WOUNDS OF MILITARY PATIENTS: THE SEARCH FOR NEW FORMULATIONS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Robert B. Lindberg, PhD Virginia C. English, MA Ruth L. Latta, BS George T. Daye, MA Basil A. Pruitt, Jr, MD, Colonel, MC

Report Control Symbol MEDDH-288 (R1)

The ultimate criterian for improved chemotherapy of cutaneous burns is based on reliable animal models. The burned rat seeded with <u>Pseudomonas</u> <u>aeruginosa</u> offers the most plausible simulation of human burn infection, but detailed study of a series of wound and blood stream isolates from <u>Pseudomonas</u> sepsis revealed a spectrum of inherent variation in virulence from zero to 100%. Strain variation occurs with many types, but a major epidemic phage type was recovered over an 8 year period with no change in virulence. In contrast, successive isolates during prolonged clinical sepsis in a major burn revealed inherent, stable variations of a major order in virulence and response to therapy of specific types. Surface tension alteration was shown to lack significance in affecting virulence or invasiveness, a trial of a new semi-synthetic penicillin, pirbenicillin, showed it to be ineffective in control of experimental Pseudomonas burn wound sepsis.

Burns Pseudomonas Sulfamylon Semi-synthetic penicillins Rats

\$

# DEVELOPMENT OF PROPHYLACTIC TOPICAL THERAPY FOR USE ON BURN WOUNDS OF MILITARY PATIENTS THE SEARCH FOR NEW FORMULATIONS

The assessment of virulence in Pseudomonas aeruginosa has been based on the ability of strains of this organism to set up invasive burn wound sepsis in the white rat when a burn is seeded with a young culture. The disease which results is strikingly similar to burn wound sepsis in the human, and medication designed to control this syndrome in animals has been highly effective on human subjects. Apart from the practical aspect of an infection model in deriving new modes of therapy, the seeded burned rat model offers an important epidemiologic tool for determining the extent of virulence in Ps. aeruginosa, the relationship of virulence to spread and persistence of a given strain in a patient population, the detection of virulence-related attributes in pathogenic strains, and other basic criteria of importance in understanding wound infection. Since the occurrence of epidemic waves of Ps.aeruginosa infections due to specific phage types has been well documented, there is ample reason to assess strains which cause invasive sepsis. These will vary, not only as to identity, but also as to virulence. The clinical significance of differences in invasive potential has not yet been elucidated, but the fact that such differences can occur makes it of fundamental importance that they be documented and studied.

Earlier reports, in the middle 1960's, showed awide spread of virulence over a series of strains collected from burn aptients. However, the identity of epidemic producing strains continues to vary and prolonged use of topical therapy in control of wound sepsis has made further study of this epidemiologic phenomenon desirable. Selection of strains has been based primarily on their exhibiting virulent attributes. They were from blood culture, wound biopsy, or from tissues post mortem, including lung and liver isolates as well as wound strains. Table 1 summarizes the range of virulence demonstrated. There was a complete range of virulence: 4 strains out of 70 were completely non-virulent; 9 more were weakly virulent; none killed more than 7% of the animals seeded. There were 2 types in each of these categories. In the intermediate range, which included up to 60% mortality, 20 strains which included 4 phage types were found. Six phage types and 37 strains exhibited high virulence. Each lethality value was determined on not less than 4 groups of animals, so that cumulative results confirmed the validity of the virulence level.

A unique degree of virulence associated with a specific phage type, A-71, has been observed in detail, in part because of its appearance as an epidemic strain associated with a high rate of Pseudomonas sepsis. When the strain reached epidemic proportions in 1973, retrospective review of the culture collection uncovered a strain of the same type recovered 8 years earlier, in 1965. It was at that time unique, and was not seen again for 8 years. It has been found since its epidemic outbreak but so seldom that it would not, except for its history, have been observed further. Table 2 shows the persistence

)

Non-Virulent	<u>Types</u> a	nd & Die	d Out of To Moderate	otal Rats Se Virulence	eded High Virulence (60 - 100%)		
(No deaths)	(1 -	198	(20 -	- 59%)			
NT-16	NT-3	2.3	NT-4	25	NT-5	82.7	
NT-A	NT-A ¹	6.4	NT-18	58.4	NT-17	94.5	
			NT-B	48.1	NT-2	67.3	
			F-21	<b>29</b> .2	D-5	92.4	
					C-26	84.3	
					A-71	99,2	
4 Strains	9 Strai	ns	20 Stra	ains	37 Strains		
2 Types	2 Туре	S	4 Typ	Des	6 Types		

# Table 1. Virulence of 70 Strains of Ps.aeruginosaRepresenting 14 Phage Types Recovered from Burn<br/>Patients, 1973-1975

of virulence in this phage type. It is unique in our experience with hundreds of strains of Ps. aeruginosa in that the strain is so uncommon, (except in epidemics) that it could not have persisted in the burn ward, but has been recovered as a new entry into this population. Nonetheless, this strain is specifically associated with a uniquely high virulence for the burned rat. This has not been the case with other virulent epidemic strains, when they have been assessed for virulence. As a model for studying the constancy of virulence, strains from a burn patient who underwent at least 2 separate septicemic episodes during his illness were studied. Their behavior is summarized in Table 3. This patient was invaded first by type F-21. Three successive isolates, identical by biochemical and phage criteria, were assessed. Significant differences in lethality occurred: 1 isolate was lethal for only 15% of untreated rats, another for 25% and the third for 44%. The progression of response to treatment paralleled the rise in virulence, from 100% effectiveness with the least virulent, to 88% effectiveness in the most virulent strain.

A subsequent and more severe episode of Fseudomonas sepsidoccurred a week later in this patient's course. This time the infecting strain was a different type, NT-2. Virulence varied from 42% to 89% in strains from the blood; 3 wound strains were all of the same level of virulence, and were probably identical. Thus, within one patient, 2 separate strains yielded isolates with 5 different degrees of virulence. Response to therapy was consistent with the concept that the more virulent the strain was, the higher

Year	No. of Patients	No.Strains A−71 Tested	Virulend Died/Tot	ce al, %	Average Time to Death (Days)
1965	1	1	30/30	100	6.7
1973	15	16	449/452	99.3	5.56
1974	1	1	10/11	90.1	7.5
1975	1	1	10/10	190	6.4

Table 2. Virulence of Tested Animals of 19	Strains of
Ps. aeruginosa, Type A-71, Recovered from	18 Burned
Patients Between 1965 and 1975	

Table 3. Animal Model Virulence and Response to Therapy of <u>Ps. aeruginosa</u> from a Patient with Lethal Burn Wound Sepsis

Strain	Source	Туре	No.of Tests	<u>Mortality</u> Sulfamylo	<u>and Tre</u> n क्षDied	atment Resp No Treatme Controls	onse nt & Died
9-25-36	Biopsy	F-21	4	0/20	0	3/20	15
9-27-13	Biopsy	F-21	5	3/25	12	11/25	44
9-28-9	Wound Swab	F-21	4	1/20	5	5/20	25
10-3-10	Biopsy	NT-2	4	7/26	27	16/26	62
10-3-11	Biopsy	NT-2	5	8/25	32	17/25	68
10-3-12	Wound Swab	NT-2	5	5/26	19	16/26	62
10-4-22	Blood	NT-2	4	8/20	40	17/19	89
10-5-15	Blood	NT-2	4	0/20	Q	8/19·	42
Strain S	ources:	Type F- NT-2; NT-2:	21: Wor Wor Blo	unds: 25- unds: 3 ( od 4-	28 Sept. Oct 5 Oct		
Control	Strains:	8-28-3:	Treat:	0-20%, mo	rtality		

No Treat.: 90-100% mortality

the proportion of treatment failures. This has not always been the case, in other experiments, high and low virulent strains were each completely susceptible to topical Sulfamylon treatment.

The experimental behavior of <u>Ps. aeruginosa</u> strains recovered from burn patients is not simply a case of virulent versus non-virulent organisms. Some strains are completely virulent, with no non-virulent variants observed. Other strains vary widely in virulence even with the most meticulous retrieval and storage technics. Further, virulent and non-virulent strains vary in their response to topical therapy. The assessment of new drugs and formulations can be made simple by using a "standard" strain that behaves in a consistent - i.e., desired manner. But the natural history of Pseudomonas burn wound infection is far more complex, and involves strain-related differences that must be taken into account if valid guides to clinically useful therapeutic agents are to be achieved.

# EFFECT OF ADDITION OF TWEEN-80 TO INOCULUM ON VIRULENCE AND RESPONSE TO THERAPY

Variations occur in response of burned rats to seeding with a given strain of <u>Ps. aeruginosa</u>. These fluctuations can interfere with interpretation of the nature of an infecting strain, and with evaluation of a therapeutic regimen. One possibility, frequently discussed as a possible factor, is the degree of dispersal of cells of the inoculum in growth; <u>Ps.aeruginosa</u> may form non-homogeneous suspensions of bacteria when they are grown in broth. A means of achieving more uniform dispersal, and of improving the wetting qualities of the culture on the burn wound, is to grow the culture in broth to which a small amount of non-toxic wetting agent has been added. This experiment was conducted with 2 challenge strains, 8-28-3 and 12-4-4, to determine whether or not demonstrable differences in lethality or in response to therapy could be demonstrated.

Five different strains, varying in lethality and in response to topical Sulfamylon therapy were used. Strain 8-28-3 is highly virulent; strain 12-4-4 is virulent but permits more survival; and A-71 represents a most lethal treatment refractory strain, with A-71-2 virulent but somewhat treatment responsive, and A-71-3 low in virulence yet treatment refractory. Table 4 summarizes the results.

There was no significant increase in lethality of strains in animals not treated, between the effect of the Tween-80 cultures and the Trypticase Soy Broth (TSB) cultures. Among treated animals, there was a slight increase in lethality despite treatment, although it was not statistically significant. Among the 7 or 9 experiments, with a total of 17 sets run, 11 of 88 of the Tween-80 broth animals died, while in the same set the TSB inoculum resulted in death of 5 out of 90 animals. It may be concluded that the wetting agent does not enhance the virulence of the challenge strain, nor does it significantly alter the response to therapy. There may be slightly more animals which die when treated following Tween-80 inoculum than with the TBS inoculum.

		Treated with	h Sulfamylon 10%	Contr	ols
Exp.No.	Strain Substrate	Died/Total	Survival* Time	Died/Tota	Survival Time
-	8-28-3 Tween-80	4/15	18.27 + 6.2	15/15	6.33 + 2.47
	TSB**	1/15	22,00	14/15	6.93 7 4.3
2	8-28-3 Tween-80	1/8	i9.88 <u>+</u> 6.01	6/7	9.43 + 6.08
	TSB	8/0	22.00	7/7	10.14 + 2.85
ω	12-4-4 Tween-80	1/7	20.28 + 4.53	7/7	$9.86 \pm 1.85$
	TSB	0/7	22.00	6/6	8.33 + 1.20
4	4-24-14 Tween-80 (A-71) TSB	14/15 14/15	$8.27 \pm 4.73$ $6.87 \pm 5.3$	14/15 13/14	7.60 + 4.12 7.36 + 4.75
1					
ţ	(A-71-3) TSB	2/13 0/14		1/13	21.46 + 1.9
ת	1-78-13 Tween-80	1/15	7N C + 7 U 1C	٤/١	22 NN
I	(A-71-3) TSB	1/14	$21.50 \pm 1.87$	1/13	20.62 + 5.0
7	5-15-2 Tween-80	2/15	20.80 + 3.2	13/14	10.36 + 3.75
	(A-71-2) TSB	0/14	22.00 -	14/14	10.72 + 2.55
œ	5-28-16 Tween	8/14	14.08 + 6.95	13/13	7.08 + 2.8
	(A-71-2) TSB	14/15	9.73 + 4.86	13/14	9.43 + 4.4
9	3-28-3 Tween-80	0/13	22.00	13/13	7.00 + 1.73
	TSB	3/14	19.86 ± 4.88	13/14	$11.07 \pm 5.35$
					والمحافظ والمحافظ والمحافظ المحافظ المحافظ والمحافظ المحافظ المحافظ والمحافظ والمحافظ والمحافظ والمحافظ والمحافظ

Table 4. Effect of 0.1% Tween-80 in Inoculum on Virulence and Response to Therapy of Pseudomonas aeruginosa in Burned Rats

* Survival time: Terminated on day 22; Survival = 22.0 days ** TSB: Trypticase Soy Broth

Since the wetting agent makes it noticeably easier to apply the 1.0 ml inoculum to the freshly burned rat, this modification has been introduced into the routine rat challenge model.

# PIRBEN/CILLIN, A NEW SEMI-SYNTHETIC PENICILLIN TRIAL IN CONTROL OF EXPERIMENTAL PSEUDOMONAS BURN WOUND SEPSIS

Perbenicillin, a new injectable semi-synthetic penicillin has a broad spectrum of antibacterial activity. Its activity is comparable to that of carbenicillin although it is reported to have enhanced activity against fecal streptococci, Pseudomonas and Enterobacteriaceae that do not produce beta-lactamase. In view of the long standing need for a more effective systemic antimicrobial agent which can control Ps.aeruginosa invasive infection in burns, and in the hope that this agent might be of value in controlling other opportunistic invading organisms in burns, it was tested in the invasive Pseudomonas burn wound sepsis model, using 200 gm rats with a 20% scald burn, seeded promptly with 10° cells of an 18 hour broth culture. In vitro tests on Ps. aeruginosa were carried out using the MIC technic standardized in this laboratory.

The sensitivity to antibiotics of 3 bacterial species of major importance in burn wound infections are summarized in Table 5. Staphylococci were relatively susceptible to the antibiotic, <u>Ps. aeruginosa</u> were moderately sensitive, and <u>Klebsiella pneumoniae</u> were resistant. The remaining species of Enterobacteriaceae were tested in small numbers; their behavior is an indication of sensitivity but numbers were few.

Species	No. Inhibited and Concentration of Pirbenicillin (mcg/ml)												
	≻ 50	50	25	12.5	6.25	3.12	1.5	0.78	0.39 <	<b>(</b> 0.39			
S. aureus	0	0	0	0	5	15	5	1	0	0			
K.pneumoniae	28	2	2	0	0	0	0	0	0	0			
Ps.aeruginosa	1	1	5	3	0	1	0	0	0	0			
E. coli	2	0	0	0	0	0	0	0	0	0			
S. marcescens	2	J	0	0	1	0	0	0	0	0			

Table 5. Sensitivity of Bacterial Species Prominent in Current Burn Infections to Pirbenicillin

Graphic representation of cumulative sensitivity gives a more informative picture of the sensitivity status. These data are shown in figure 1. The results



CUMULATIVE SENSITIVITY OF KLEBSIELLA PHEUMONIAE, STAPHYLOCOCCUS AUREUS AND PSEUDOMONAS AERUGINOSA STRAINS TO PIRBENICILLIN

のないない。このではない

ŧ

Figure 1.

make more emphatic the inherent resistance of Klebsiella strains and the intermediate status of Pseudomonas. Only <u>Staphylococcus</u> aureus was sensitive to Pirbenicillin.

The use of Pirbenicillin in control of experimental burn wound sepsis was based on the encouraging results in clinical trials reported in communications from research laboratories of the manufacturer, Pfizer, Inc. Pharmacologic data indicated that the drug was tolerated in doses up to 250 mg/kg, via subcutaneous administration, in rats. After preliminary trials, dosages of 250 mg/kg and of 125 mg/kg, intraperitoneally on a 1 X/day basis, were selected. Dosage schedules of 5 and 10 days were carried out in groups of 5 or 6 rats each. Treatment was begun 24 hours a ter seeding of the rats. Animals were observed for 21 days after burning and seeding. The cirterion of effectiveness was survival. Challenge strains included 12-4-4, 8-28-3 and VA-134, to cover a spectrum of virulence and invasiveness.

Results are summarized in Table 6. The 3 challenge strains varied slightly in their response to Pirbenicillin, but none of the results was encouraging to a significant extent. With strain 12-4-4, the maximum dose presented a 70% mortality with a strain which, untreated, killed 85.6% of the animals. Strain 8-28-3 was more virulent in the untreated control, but it showed the most encouraging results in vivo: maximum dose for maximum time showed a 29.4% mortality; when the 250 mg/kg/day dose was run for 5 instead of 10 days, mortality rose to 64.7%. The reduced dose of 125 mg/kg/day actually showed the best survival of all; it was the same for the 5 day as for the 10 day interval. The most virulent challenge strain, VA-134, showed a slight response to the maximum dose, and virtually no effect with the minimum dose.

Dosage mg/kg	Treatment Period	No.of Tests	Strain and Outcome - Died/Total (% Mortality)							
			12-4-4	8-28-3	<u>VA-134</u>					
250	10 days	3	12/17 (70.5)	5/17 (29.4)	10/17 (58. <b>8)</b>					
	5 days	3	12/17 (70.5)	11/17 (64.7)	13/17 (76.4)					
125	10 days	3	11/17 (64.7)	9/17 (52.9)	12/17 (70.5)					
	5 days	3	12/17 (70.5)	8/17 (47.0)	16/17 (94.1)					
Control			13/15 (85.6)	13/13 (100)	16/16 (100)					

Table 6. Pirbenicillin in Prophylaxis of Burn Wound Infection in Rats Seeded with Pseudomonas aeruginosa

#### DISCUSSION

The results of animal trials of fribenicillin in control of Ps. aeruginosa burn wound sepsis were not encouraging. The drug had a slight effect in the creasing survival, but it was not as effective as Carbenicillin in similar circumstances and far less effective than Ticarcillin which was shown to have a high potential for controlling invasive Pseudomonas infection in the seeded burned rat. On the basis of this particular model and challenge system, there is little enthusiasm for a clinical trial of Pirbenicillin.

#### PRESENTATIONS

Lindberg RB: Hospital Epidemics of Drug-Resistant <u>Pseudomonas aeruginosa</u> Infections. Am. Soc. Microbiol., New York, 28 May 1975.

# PUBLICATIONS:

Lindberg RB, Latta RL, Pruitt BA, Jr: Transfer and Control of Hospital Epidemics of Drug-Resistant <u>Pseudomonas aeruginosa</u> (Abst). Am. Soc. Microbiol. 1975, p. 40 (C-83).

				1. AOEN	CY ACCES	HON	L DATE OF SU	DATE OF SUMMARYS REPORT CONTINUE STUDDE			
RESEARCH	AND TECHNOLOGY	y work unit s	UMMARY	גמ	0B 69	78	76 10	76 10 01		DD-DR&E(AR)436	
1. DATE PREV SUMAY	4. KIND OF SUMMARY	S. SUMMARY SCTY	. WORK SECURITY	7. REGRA	VDING#	10.00	BI'N INSTR'N	OL SPECIFIC	OATA-	. LEVEL OF SUIT	
75 07 01	H. TERM		n ,	N	NA		NT.	CONTRACTOR		A WORK UNIT	
10 NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK A	REA NUN	DER		WORK UNI	TNUMBE	R	
	_1102A	3A161102	B71R	01			219				
S. CONTRIBUTING				[	· · · · · · · · · · · · · · · · · · ·		804-99 <i>6</i> 7				
C. CONTRIBUTING											
IL TITLE (Protes with On Biopsy an	security Close illustion Code and <u>Autopsy Ti:</u> Chinological are s	"(U) The Ro ssues From	ole of Fung Seriouslý	i in Burne	Bvrn d Sol	Wour die:	nd Infec (5 (44)	tion:Ob	serva	tions	
003500 Clini	ical Medicine										
13. START DATE			IL PUNDING AGE			16. PERFORMANCE			HOD		
66 02 76 09				DA				C. In-House			
17. CONTRACT/ORANT Not Applicable				10. RESOURCES ESTIMAT			A PROFEM		S & PU	iDE (In disustandig	
& DAYES/EPPECTIVE: EXPIRATION:			- A						17		
NUMBER:*				PISCAL 7T					_	7	
& TYPE:		A ANOUNT:		VEAR	CONNENT						
. KIND OF AWARD:		f. CUM. AMT.			L			-			
IN MESHANINGLE DOD C	PHEANTEATION	L		<b>30</b> . PCN7 		RGANIZ	ATION	L			
HANE!" US ATTTY	Institute of	Surgical I	Research	"MME." US Army Institute of Surgical Research							
ADDAESS: Fort Sam Houston, Texas 78234				Microbiology Branch ADDREMS!* Fort Sam Houston, Texas 78234							
RESPONSIBLE INDIVIDUAL NAME: Basil A. Pruitt, Jr, COL, MC TELEPHONE: 512-221-2720				PRINCIPAL INVESTIGATOR (Puriles SEAN II U.S. Associate (Milliones) NAME: ROBERT B. Lindberg, PhD TELEPHONE: 512-221-2018 SOCIAL SECURITY ACCOUNT NUMBER:							
FOREIGN INT	ELLIGENCE NOT	CONSIDEREI	)	NAME: Anthony A. Contreras, MS NAME: DA							

(U) Fungi; (U) Mucor; (U) Rhizopus; (U) Fusarium; (U) Burns; (U) Phycomycosis; (U) Humans 23. TECHNICAL OBJECTIVE, * 24. APPROACH, 26. PROGRESS (Punish Individual paragrap, a identified by number a test of each with geourity Classification Sude )

23. (U) To determine species of fungi in burn patients and assess the significance of these opportunistic invaders in producing fungal burn wound pepsis, respiratory tract involvement and other infectious processes peculizi to infection in burned soldiers.

24. (U) Cultivation of biopsy and autopsy specimens under  $gp^{\mu}(a,p)$  conditions for recovery of fungi and yeasts.

25. (U) 75 07 - 76 09 The occurrence of fungal colonization diminished further in FY 1976. Of 135 biopsies on 63 patients, 5 patients yielded 6 fungal cultures, while 11 patients yielded 13 strains of Candida sp. In autopsies, 38.8% of patients yielded Candida sp.; only 23.6% of autopsies yielded fungi in culture. Fusarium and Aspergillus were genera more frequently recovered; 9 genera were recovered and 38 strains were isolated. No phycomycetes were found. Fungal sepsis remains a possibility but in 1975 it was relatively uncommon.

A vallable to contractore upon artamator's aprioval PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DO FORMS 1496A 1 NOV 65 AND 14961 1 MAR 88 "FOR ARMT USE) ARE OBSOLETE
#### TERMINATION

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE - SURGERY

REPORT TITLE: THE ROLE OF FUNGI IN BURN WOUND INFECTION: OBSERVATIONS ON BIOPSY AND AUTOPSY TISSUES FROM SERIOUSLY BURNED SOLDIERS

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 June 1976

Investigators:

Robert B. Lindberg, PhD

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

#### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES - SURGERY

REPORT TITLE: THE ROLE OF FUNGI IN BURN WOUND INFECTION: OBSERVATIONS ON BIOPSY AND AUTOPSY TISSUES FROM SERIOUSLY BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 June 1976

Investigators: Robert B. Lindberg, PhD

Reports Control Symbol MEDDH-288(R1)

Autopsy and biopsy tissues, including wound, lung, spleen, liver and burn wound in autopsy, are routinely cultured for fungi. Fungal sepsis has not been a serious clinical problem since 1973, but fungal colonization still occurs. The recovery rate of yeasts and fungi fell from 71 - 75% in recent years to 52% in 1975. Predominant genus was <u>Fusarium</u>. Phycomycetes, the most dangerous fungi when invasive infection occurs, were not recovered in 1975.

Fungi Mucor Rhizopus Fusarium Burns Phycomycosis_ Humans

# THE ROLE OF FUNGI IN BURN WOUND INFECTION: OBSERVATIONS ON BIOPSY AND AUTOPSY TISSUES FROM SERIOUSLY BURNED SOLDIERS

Fungal infection in burn wounds has followed an unpredictable course. Individual institutions have seen such invasive infections increase to a point at which they were a serious problem in morbidity and mortality. The incidence may then regress, with no valid basis in changes in wound management. In the Institute of Surgical Research burn wound invasion by fungi has become rare, although 4 to 5 years ago it was a problem involving 5 to 10 patients per year, in a pathological process that was either uncontrollable or controlled with ablation of involved tissues when such surgical intervention was possible.

#### Fungi in Biopsy Specimens

Biopsies were performed on 63 patients in 1975, and 135 specimens were collected. Results of cultures of tissues on Sabourauds agar are set forth in Table 1. Comparative results for 1972 - 1974 are set down. It was obvious that far fewer biopsies were collected in 1975 than had been the case in preceding years and the scale of recovery of fungi had correspondingly narrowed. Thirteen genera of fungi were recovered in 1972, 10 in 1973 and 7 in 1974; in 1975 4 genera were recovered from burn wounds. Had it not been for preceding years and the importance of the role played by fungi; the very procedure might have been discontinued. However, the pathologic evidence of a significant role being played by fungi is, on an overall view, too important to be ignored.

#### Fungi Recovered from Autopsy in Burned Patients

Fungi recovered from burn wounds and viscera (lung, liver and spleen) at autopsy are summarized in Table 2. There were 9 genera of fungi recovered, as always, these organisms were more common in tissues at autopsy. Presumably one reason for this discrepancy is the ease with which opportunistic microorganisms proliferate in tissues of the immunologically impaired host. There were 9 genera of fungi, including the <u>Candida spp</u>, recovered. The number of isolates was small, but <u>Fusarium</u> <u>spp</u> were the predominant genus, as they had been in the preceding two years. <u>Aspergillus spp</u> were the other major fungal form recovered. No genera with a pedigreee of ability to cause invasive burn wound infection were recovered.

The nature of the decrease in frequency of recovery of fungi in burned patients in the ISR is most clearly shown in a comparison of recoveries made since 1971. These totals are shown in Table 3.

It was obvious that fewer positive fungal cultures appeared in 1975 than in recent years. In the interval between 1971 and 1975, the fraction of patients cultured who yielded fungi or yeasts from tissues ranged from

Genus	Nc. 1972	Patients 1973	Posi 1974	tive 1975	No. 5 1972	Strains 1973	<b>Re</b> cov 1974	ered 1975
Aspergillus	8	8	8	1	11	17	5	2
Cephalosporium	5	4	5	1	15	5	5	1
Fusarium	19	12	11	2	33	23	17	2
Sepedonium	0	1	0	0	0	1	ΰ	0
Curvularium	3	2	1	0	3	2	3	0
Scopul ariopsis	11	0	0	0	11	00	0	0
Alterneria	3	2	3	1	7	2	3	1
Diplosporium	1	0	0	0	ŀ	0	0	0
Penicillium	1	1	3	0	1	1	3	Ð
Helminthosporium	1 O	L;	2	0	0	9	2	0
Rhi zopus	2	2	0	0	3	2	0	Ó
Mucor	2	2	0	0	2	2	0	0
Stemphyllium	1	0	0	0	1	0	0	0
Syncephal astrum	1	0	0	0	í	0	0	0
Geotrichum	1	0	2	0	1	0	4	0
Candida sp.	28	57	54	11	46	141	144	15
No. Patients Cul	ture	d	135	63				

# Table 1. Biopsy Specimens Positive for Fungi and Yeast - 1972 - 1975

Genus	Wounds Patients Positive	. Total Isolates	Visc Patient - Positive	era Total Isolates
Aspergillus	3	3	9	12
Penicillium	1	1	0	0
Cephalosporium	1	1	2	2
Fus <b>arium</b>	5	5	3	6
Scopulariopsis	1	2	C	0
Cyrvularium	2	3	1	1
Al terneri a	0	0	3	4
S <b>epedoniu</b> m	1	1	2	2
Mycelia sterilia	a 0	0	1	1
C <b>an</b> di da	12	21	23	48

# Table 2. Genera of Fungi Recovered from Viscera (Liver, Lung, Spleen) and Burn Wound at Autopsy 1975

Genus		<b>B</b>	Nugbe	r of P	atients	Positive	at Aut	opsy	•	
	1971	1972	1973	1974	1975	1971	1972	1973	1974	1975
Hucor	3	0	3	3	v	0	0	1	1	0
Rhi zopus	1	1	2	1	0	1	1	0	0	0
Absidia	1	1	0	0	0	0	0	0	0	0
Aspergillus	6	11	19	4	3	3	2	5	2	9
Penicillium	8	1	0	1	1	1	2	0	5	0
PaeciloRyces	0	0	0	0	0	1	2	0	0	0
Al terneri a	2	3	6	1	0	0	0	0	2	3
Cephalosporium	2	4	3	4	1	0	4	4	8	2
Fusarium	9	30	25	12	5	7	20	19	8	3
Helminthosporium	4	1	4	4	0	0	0	2	l,	0
Nigrospora	9	0	0	0	0	3	0	0	0	0
Scopul artopsis	5	3	1	2	1	1	2	0	2	0
Sepedonium	1	5	5	١	1	1	4	3	۱	2
Diplosporum	1	2	0	0	0	0	0	0	0	0
Geotrichum	0	0	3	0	0	2	0	0	۱	0
Fonsecasa	2	0	0	0	0	2	0	0	0	0
Curvul ari J	0	5	1	C	2	0	0	0	0	۱
Ni crosporum	0	1	0	0	0	0	0	0	0	0
Cladosporium	0	0	1	0	0	0	0	3	0	0
Candida	13	32	67	41	12	11	25	50	42	23
Number patients positive Number patients cultured Number genera present	46 61 17	65 89 16	80 91 14	67 80 15	39 74 9					

Table 3. Fungi Recovered from Autopsy Tissues, 1971-1975

73% to 83%. In 1975, 52% of the patients cultured yielded yeasts or fungi.

The absence of Phycomycetes in biopsy and autopsy in 1975 was significant since this group includes the Mucor, Rhizopus and Obsidia genera. These can and have caused invasive burn wound sepsis with very lethal consequences.

The trend toward lessened incidence of invasive fungal infection of burn wounds was apparent in 1973, and since that time this disease has kirtually disappeared in the ISR. The low recovery rate in both autopsy and biopsy in 1975 was gratifying, but there has been no specific step that can be regarded as controlling fungi. Contunued monitoring is especially pertinent since interpretation of clinical disease may be aided very much by adequate comprehensive autopsy and biopsy culturing.

PRESENTATION AND/OR PUBLICATION

None

RESEARCH	AND TECHNOLOG	Y WORK UNIT S	UMMARY	DAC	E 639	8	76 10 0	1	DD-D	REE(AR)636
& BATE PREV SUPRY	L KIT OF SUMMARY	S. SUMMARY SCTY	A. WORK SECURITY	7. NEGR	DING	-		A SPECIFIC	ATA-	. LEVEL OF DUE
75 07 01	H. TERM	U	U	1	i)	N	L	Q ves C	) #0	A WORK WET
10. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK /	REA NUM	DEA		WORK UNIT	NUMBE	R
	61102A	<u>3A16110</u>	2B71R		01		319			
L CONTRIBUTING		1 <del> </del>		ļ			840 X		22	
C. CONTINUUTING	Annutite Classification Code	Verter Nor-E			Noge	13 1 1 1 1 0	Paci 11	(	and a	Soldiers
New Potenti	al Opportunis	+ic Pathog	ermentative	Grau	u-neya	CIVE	DaCIII		meu	<b>SOTATELS</b>
IL SCIENTIF'C AND TE	CHHOLOGICAL AREAS	cie i ucitog								
003500 Clin	ical Medicine	•								
IL ETAN / DATE		TA BATIMATED COM	PLETION BATE	TIE PUN	NPS AGEN	EV		H. PRAPORM	NCE ME	THUD
74 07		76 09		DA		-	1	C. In-	louse	•
W. CONTRACT/GRANT	Not Applicabl	le		18. MES		THATE	A PROFUSO		1 10	NBC (In these and g
A DATENEPPECTIVE		EXPIRATION:			76	5		5		18
A NUMBER:*				PHICAL	72	r	نعــــــــــــــــــــــــــــــــــــ	L	<b>_</b>	3
& TYPE:		4 AMOUNT:		TEAR						
& KNID.OF AWARD:	DRAMIZATION	I.CUM. ANT	·	1 30. 547		BRAW'S			1	
	The marked data data = 4	[]	Pagarat		110 ×		معت في معالم من			
HAME!" US ATTRY	institute of	surgical	Research	HANE!"	USAI	nny 1		e or su	rgica	11 Kesearci
ABBRENS Port	Sam Houston	Toyad 792	34	ADDRES	MLC M ^e Rort	· Can	Hough	m. Texa	e 71	8234
FOLC	Sam nouscon,	16743 102	74		- FOI	. Jan		MIT ICAU		
				PRINCI	-		(Pumish BEAN	il V.S. Acodemic	jas itasile	-
	AL			HANE	• Rober	t B.	Lindbe	erg, PhD		
HAME Basil A	. Pruitt, Jr,	, COL, MC		TELE	PHONE:	512-	221-20	18		
TELEPHONE: 51	2-221-2720			80CIA	L SECURITY	Y ACCOU	NT NUMBER:			
SI. CONERAL USE				A860CI/		ISATOR	•			
FOREIGN INI	ELLIGENCE NO	r considere	:D	NAME	Virgi	inia	E. Engl	Lish, MS		
				HANE						DA
L REVOONDE (Proses			Burns; $(0)$		ative	gram	i-negat.	LVE DACL	/ 17/4 777}	Mang
(U) Pseudon	ionas; (U) AC	Inecopacter	; (U) ALCA.	Ligen		JJ E J		CEUMIT (U		<b>mor</b> 19
22 (II) Mc	et infection	e in severe	ly burned	പിർ	OTS AL	re di	le to o	oportuni	stic	flora
and there h	ave heen sev	eral specie	s recomiz	ed as	havin	nor ma	tior pa	thogenic	pot	ential
only after	they had been	n found in	cases of s	epsis	in b	urns.	Dete	ction of	pot	ential
pathogens of	apable of se	tting up no	socomial i	nfect	ion is	s me	rited a	s enlarg	ing	
knowledge r	egarding wou	nd infectio	on potentia	litie	s in :	inju	red sol	diers.	-	
			-			•				
24. (U) Sr of detected	ecial cultur groups.	e technics	and substr	ates	are u	sed a	as indi	cated by	bio	logy
25. (U) 7	5 07 - 76 09	Significa	nt numbers	of 9	speci	ės,	B of wh	ich were	oxi	dative and
one rerment	tative, were	recovered i	rog ruled o	y and	( DIOD:	oont:	ecineti	on of a	ROUR	Ce.
in some que	intitative ti	+bogens bay	ve thus hee	n evr	nogeđ.	cont		on or u	JULL	
Potential	ourn wound pa	choyens na	ve chus bee							
}										
1										
1										
1										
1										
ļ										
ł										
1										
l										
A volleble to contrac	lors youn originator's app	pro ve l								

# TERMINATION

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: NON-FERMENTATIVE GRAM-NEGATIVE BACILLI IN BURNED SOLDIERS: NEW POTENTIAL OPPORTUNISTIC PATHOGENS

# US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ÁRMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Virginia C. English, MA Robert B. Lindberg, PhD

Report Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

## **REPORT TITLE:** NON-FERMENTATIVE GRAM-NEGATIVE BACILLI IN BURNED SOLDIERS: NEW POTENTIAL OPPORTUNISTIC PATHOGENS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Virginia C. English, MA Robert B. Lindberg, PhD

Report Control Symbol MEDDH-288(R1)

Oxidative, non-fermentative gram-negative bacteria have been found in significant numbers as predominant organisms in tissues at autopsy and/or biopsy of burned patients. These species are seldom recovered from clinical specimens; their relatively slower growth and fastidious character make overgrowth by Enterobacteriaceae a major complicating factor in their detection. Eighteen patients harbored one or more of 10 species, with <u>Pseudomonas maltophilia</u>, <u>Pseudomonas putida</u>, <u>Acinetobacter calcoaceticus</u> and <u>Acinetobacter lwoffi</u> as predominant forms. Their presence as predominant organisms at autopsy suggests that they may play a significant, if unrecognized, role in burn morbidity and mortality. As relatively new opportunistic invaders, they should be sought with more effort than is presently routinely made.

Burns Oxidative gram-negative bacilli Pseudomonas Acinetobacter Alcaligenes Flavobactrum Humans

# NON-FERMENTATIVE GRAM-NEGATIVE BACILLI IN BURNED SOLDIERS NEW POTENTIAL OPPORTUNISTIC PATHOGENS

From the first recognition of infectious disease as caused by bacteria. there has been a continually expanding recognition of pathogenic microorganisms. Specific diseases were associated with causative agents, and even in non-specific situations such as wound infections, emphasis was placed on aerobic cocci and Clostridia as the principal offenders. From pre-bacteriologic time, however, records point to one non-fermentative bacillus, B. pyocyaneus, now designated Pseudomonas aeruginosa. The blue-green pigment seen on wound dressings, which incidentally often prompted a favorable prognesis, was and still is caused by this ubiquitous organism, which is innocuous for healthy individuals but which can act as a lethal invasive pathogen in severe burns. debilitated individuals, or in immunologically compromised individuals. An increasing number of gram-negative bacilli other than classic Enterobacteriaceae species have been recognized in recent years as capable of opportunistic pathogenic activity. Many of these are oxidative rather than rementative in their utilization of carbohydrate. Their pathogenic activity can involve generalized sepsis, especially in infants and children, but their primary interest in burn patients revolves around their role as opportunistic wound invaders.

#### METHODS

Routine diagnostic procedures on wound, blood, sputum and other body fluids do not exclude specific search for obscure species, but it has not been feasible to scrutinize exhaustively every clinical specimen. However, tissues from wound, lung, liver and spleen are cultured quantitatively in autopsies, as are biopsy specimens, and the highest dilution with discrete colonies is scrutinized in detail for unusual organisms. Colonization of burn wound, pneumonia or septicemia caused by organisms sequestered in spleen and liver could be revealed by such examination, and the extent of recovery of unusual species has validated this supposition. A significant number of oxidative organisms of species not previously observed in the Institute of Surgical Research patients have been recovered.

# UNUSUAL GRAM-NEGATIVE BACILLI RECOVERED IN 1975

Unusual, mainly oxidative, non-fermentative gram-negative bacilli recovered from high dilution plates at autopsy are summarized in Table 1. There were 18 patients, out of 76 autopsies, in whom one or more of these unusual species were recovered. Thus, 23% of autopsies of burn deaths showed one or more of these species predominant in wound and/or viscera. This is not an inconsequential rate of colonization and potential infection, and makes continued search for these organisms a highly desirable procedure

There were 10 species recovered. Their distribution by numbers of patients and by source is shown in Table 2. The predominant species was

Autopsy No.	Patient No.	Source	Organism
A-3-75	3	Spleen:	Acinetobacter lwoffi
A-4-75	5	Tissue Bx. # 4:	Acinetobacter lwofii
A-5-75	240*	Spleen:	Pseudomonas maltophilia
A-6-75	15	LUL, Liver, Spleen, RLL:	Pseudomonas maltophilia
		LUL, RUL, RLL, Spleen:	Pseudomonas putida
A7-75	6	Liver:	Flavobacterium, Type II-B
A-9-75	30	Spisen:	Moraxella oslensis; Acinetobacter calcoaceticus
ĺ		Tissue Bx. # 2	Acinetobacter lwoffi; Pseudomonas maltophilia
		Tissue Bx. # 8, Liver; LUL, RUL, Spieen, LLL:	Pseudomonas maltophilla
A-10-75	25	Liver :	Pseudomonas marktophilla
A-11-75	23	Spleen, RLL, Liver: LLL:	Pschriomonas maltophilia Pseudrmonas putida
A-12-75	28	Spleen:	Apyocyanogenic Pseudomonas aeruginosa
A-28-75	46	Spleen:	Apyocyanogenic Pseudomonas aeruginosa
		RLL:	Achromobacter, biotype 3, Pseudomonas maltophilia
3-20-75	ØC.	LLL, RUL:	Pseudomonas maltophilia
A-30-75	86	Liver:	Acinetobacter calcoaceticus
		LLL, RLL:	Pseudomonas maltophilia
A-31-75	80	Spleen:	Acinetobacter calcoaceticus Pseudomonas maltophilla
A-38-75	109	Liver, LLL, Spleen: RUL, LUL:	Achtomobacter sp. Pseudomonas putida
A-39-75	96	RLL, LUL, Spleen:	Achromobacter sp.
A-41-75	128	RUL:	Achromobacter sp.
A-45-75 A-34-75	137 85		Citrobacter diversus
	~ •		

 
 Table 1. Uncommonly Isolated Gram-Negative Nonfermenting Organisms from Postmortem Tissues - 1975

* Organism recovered in 1975 from a patient admitted in 1974.

Pseudomonas maltophilia, found in 9 patients. Antemortem sepsis has been suggested as a plausible source of sequestration of bacteria in viscera. Acinetobacter lwoffi, formerly classified as Mima polymorpha, and Acinetobacter calcoaceticus, the former Herellea vaginicola, were similar in rate of occurrence. Pseudomonas putida, and 2 species of Achromobacter, were each recovered from 3 autopsies. The total recovery rate was similar to earlier studies of this material. The smaller totals for 1975 are due to a more limited series of examinations.

Species	No.Patients Positive	Wound	Sites of Red Spleen	covery Liver	Lung
A. Iwoffi	3	2	1	······	<u></u>
Ps. maltophilia	9	2	5	5	4
Ps. putida	3		1		2
Flavobact. II,B	1			1	
Moraxella osloensis	1			1	
A. calcoaceticus	4	1	2	1	
Apyocyanınogenic Ps. aeruginosa	2		2		
Achromobacter Type	3 3		2	1	3
Achromobacter sp	3		2		3
Citrobacter diversus	1				1

# Table 2. Patients and Sites From Which Oxidative Organisms Recovered at Autopsy, ISR - 1975

It was observed that patients in whom a non-fermenting, e>otic orcianism was found were especially prone to harbor more than one species.  $S_{x}$  en patients, out of the 18 positive for an oxidative species, harbored more than one species. This sequence is shown in Table 3. <u>Ps. maltophilia</u>, in parallel with its overall incidence, was found in 5 of the 7 multiple infections. Liver and lung were each positive in 6 patients, the spleen in  $S_{x}$  while only one patient with multiple organisms had a mixed culture in samples of the burn wound.

Antemortem isolations were less common than those recovered at autopsy, but there were 5 species recovered from clinical specimens. Their occurrence, in connection with postmortem recoveries, and the total of isolates of each species of oxidative mircoorganisms, are presented in Table 4. Apyocyaninogenic Ps_aeruginosa, Ps. putida, Ps. maltophilia and Ps_acidovorans were recovered antemortem, the first 3 were also conspicuous in autopsy material Table 3. Patients Whose Postmortem Tissues Presented Multiple Nonfermenting Organisms

ent	Surface			
	Biopsy	LIVER	SPLEEN	FUNG
		P.maltophilia	P.maltophilia	P.maltophilia P.putida
4 d	lwoffii maltophilia	P.maltophilia	M.osloensis A.calcoaceticus	P.maltophilia
		P.maltophilia	P.maltophilia	P.maltophilia P.putida
1	- 1 1 1 1 1	1 1 1 1 1 1 1	Apyocyanogenic P.aeruginosa	Achromobacter
1	,         	A.calcoaceticus	•                   	P.maltophilia
<b> </b>   <b> </b>		A.calcoaceticus P.maltophilia		) ; ; ; ; ; ;
1	, 1 1 1 1 1	Achromobacter sp.	Achromobacter sp.	Achromobacter P.putida

ORGANISM	Number of patients	Number of Isolates	Antemorteum Sources	РО	STMORTEU	M SOURCES		
				Surface	spleen	liver	lung	
P.aeruginosa,								
Apyocyanogenic	4	ىرى	ľ – Bx.L.arm 1 – lukens	-	2	2		
P.putida	4	8	l - Foley tip				6	
P, mal tophilia	10	24	1 - Blocd	_	ч	4	14	
P.acidovorans		-	l - urine					
A. lwoffii	w	w		2				
Flavobacterium, GP II-B	-1	<u> </u>						
M. osloensis	_	1						
A. calcoaceticus	4	4		_	2	_		
Achromobacter sp.	4	œ			2		4	

# Table 4. Incidence of Nonfermenting Organisms Infrequently Recovered from Patients

The burn injuries incurred by the patients from whom oxidative organisms were recovered were relatively severe. The total burn and extent of third degree burn are shown in Table 5. Burn extent ranged from 46.5% to 93% of body surface. Five out of 8 patients had burns of over 60% total body surface. Sepsis in such injuries is not unusual. However, there was no unique quality about the injuries that would set them aside from injuries with more typical fermentative flora.

The strains of 3 species of Pseudomonas other than <u>aeruginosa</u> were examined for their sensitivity to Sulfamylon, since each of these had been recovered from patients whose wounds merited topical treatment to control invasive infection. Results are shown in Table 6. The strains were inhibited by 0.078% concentration.

Since a highly refined phage typing system for <u>Ps. aeruginosa</u> has been developed and serves to monitor this organism, a sample of strains of <u>Ps.</u> <u>putida</u>, <u>Ps.maltophilia</u> and <u>Ps.pseudoalcaligenes</u>, were tested with all <u>18</u> of the typing phages. No strains reacted at the routine test dilution, but 8 out of 10 of the <u>Ps.maltophilia</u> reacted with phage M-6 when it was used undiluted. The other 3 strains of 2 species did not type. It is of interest to note that <u>Pseudomonas</u> fluorescens strains, a species closely related to <u>Ps. aeruginosa</u>, were found to be susceptible, in a mall proportion of isolates, to phage 44.Crossspecies susceptibility in the genus Pseudomonas suggests a common cell structure that also makes pathogenic involvement with this genus more plausible.

The distribution of oxidative organisms was not typical of major epidemic pattern of spread, but the periods of positive recoveries came in clusters. Some episodes chronologically suggestive of patient-to-patient transmission occurred. The strains are slow growing and readily obscured by Enterobacteriacese in pure-culture, and their detection is probably far from adequate in laboratories not specifically geared to search for them. All of the autopsy and biopsy specime is were recovered as predominant organisms in tissues; this prompts the standard upposition that they played a significant role in the infection. They were audoubtedly present in mixed flora on the burn wound, and routine cultural procedures failed to detect them.

The history of recognition of opportunistic invaders in wounds has been a succession of situations in which the opportunist was initially dismissed as a contaminant and not of clinical significance. <u>Ps. aeruginosa</u> was at one time thus regarded, as were the various species of Enterobacteriaceae which are now recognized as major causes of sepsis in burn wound infections. The relatively obscure group of oxidative forms thus far accumulated in burns is not now a recognizable clinical problem, but their presence in large numbers in wounds and viscera at autopsy indicates that they can play a pathogenic role. <u>Ps. maltophilia</u> in particular merits closer observation. Monitoring such exotic species is required if a true picture of burn wound pathogenesis is to be obtained.

# PRESENTATION

# Table 5. Survey of Pseudomonas Maltophilia Recovered From ISR Patients

Total Patients: 10

Total Isolates: 24

# ANTEMORTEM ' ISOLATE: 1

Patient No.:	133
Percent Burn:	37%
Percent of 3 [•] :	1%
Source:	8100d

# POSTMORTEM ISOLATES: 23

Patient No.	Percent Burn	Percent of 3	Source
240*	67.0	18.5	Spleen
RG	<b>63.0</b>	14.5	Liver
15	77.5	53.0	LUL, Liver, Spleen, RLL
23	65.5	40.5	Spleen, RLL, Liver
25	62.5	30	Liver
30	51.0	31.5	A # 2, A # 8, liver, Spleen LLL, LUL, RUL
46	46.5	3.0	LLL, RLL, RUL
80	54.5	11.0	Spleen
86	93.0	62.5	LLL, RLL

* 1974 admission; culture received 1975.

Patient No.	Sensitivity to Sulfamylon [*]	Reaction to P. aeruginosa Bacteriophage system ^{**}
P_MALTOPHILIA:		
30	0 078	volocia P= 1
15	0.070	P=1
23	-	P-1
28	0.078	P-1
80	0.078	P-1
RG	0.078	P-1
183	0.156 (3 isol	ates) P-1
128	0.312	NT
<b>4</b> ó	0.078	P-1
241	0.078	NT
<u>P. PUTIDA</u> :		
22	0 156	NT
15	0.078 (2 isol	ates) NT
P. PSEUDOALCALI	GENES:	
241	0.039	Th

# Table 6. Special Studies on Pseudomonas SpeciesOther Than Aeruginosa

* Sensitivity procedure same as than for P. aeruginosa ** Typing system used which is presently used for typing

P. aeruginosa

Υ.

**** Inhibiting concentration expressed in gms/100 ml.

**** These reactions were obtained using concentrated phage; Code P-1 denotes a single reaction with phage M-6. English, VC. Isolation of <u>Vibrio alginolyticus</u> from wounds and blood of a burned patient. Presented at National meeting Am Soc for Med Technology, Chicago, III. 23 June 76

# PUBLICATIONS

English VC, Lindberg RB: Isolation of <u>Vibrio alginoluticus</u> from wounds and blood of a burn patient. Am J Med Tech (in press).

DECEADOW	AND TECHNOLOGY			1. AGEI	CY ACCES	1000 ¹⁰ 2.	DATE OF SUM	MARY	REPORT O	WTROL STREOL
RESEARCH	AND TECHNOLOGI	WORK UNIT S		DA	OA 69	80 7	76 10 01	•	DD-DR	<b>h</b> E(AR)636
3. DATE PREV SUMRY	4. KIND OF SUMMARY	B. SUMMARY SCTY	. WORK SECURITY	7. REOR	ADING	BA DIES	rh metr'n	Sh SPECIFIC I	ACCESS	LEVEL OF SUM
75 07 01	H. TERM	ប	U	N	1	N	G	Tes [	но	A. WORK UNIT
10. NO /CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TABK	AREA NUM	DER		WORK UNIT	NUMBER	
. PRMARY	61102A	3A161102	B71R		01		165			
S. CONTRIBUTING							80.777j		×	
C. CONTRIBUTING		•								x
1. TITLE (Procode with .	Security Classification Code)	*(V) Studie	s of Distu	rband	ce of	Prote	ein Turr	nover in	Burn	ed
Troops-Use	of an Arimal M	<u>lodel (44)</u>								
12. SCIENTIFIC AND TE	CHNOLOGICAL AREAS									
003500 Clin:	ical Medicine						ويتحرج والمحرجين			
13. START DATE		14. ESTIMATED COM	LETION DATE	IL FUN	DING AGEN	CY		H. PERFORM	NCE METH	00
65 07		10 03						<u> </u>	i-nous	e
17. CONTRACT/GRANT	Not Applicab	le		18. RES	OUNCES ES	TIMATE	A PROFESSIO	MAL MAN YRS	E FUNC	t (In thousands)
A DATES/EFFECTIVE:		EXPIRATION:			76	4		7	28	
N. NUMBER:*				FISCAL	7 <u>T</u>			3	13	
G TYPE:		4 AMOUNTI		YEAR	CUPACNT				1	
& KIND OF AWARD:		I. CUM. AMT.					[		1	
19. RESPONSIBLE DOD C	DRGANIZATION			30. PER	FORMING C	RGANIZA	TION			
HAHE!" US Army	Institute of	Surgical H	Research	HANE!	US Ar	my I	nstitute	of Sur	gical	Research
					Renal Branch					
ADDRESS!" Fort Sam Houston, Texas 78234 Fort Sam Houston, Texas 7823					234					
				ł						
				PRINCIP	AL INVEST	GATOR (	Fumiah BEAN H	U.S. Anademic	ine li fution)	
RESPONSIBLE INDIVIDUAL			NAME: Wanda L, Brown, MS							
NAME: Basil	A. Pruitt, Jr	, COL, MC		TELEP	HONE:	512-	221-4652	2		•
TELEPHONE: 51	2-221-2720			SOCIAL	SECURITY	ACCOUN	IT NUMBER:			
21. UENERAL USE				ASSOCIA	TE INVEST	GATORS				
FOPEIGN INT	ELLIGENCE NOT	CONSIDERED	)	-	Elean	or G	. Bowle:	c, PhM		
				HANE	Arthu	r D.	Mason,	Jr, MD		DA
22. KEYWORDS /Pretade	BACH with Security Classific	allen Code)								

(U) Protein; (U) Burn; (U) Trauma; (U) Turnover; (U) Rats; (U) Albumin; (U) Edema

23 TECHNICAL OBJECTIVE.⁴ Is APPROA M, 21 PROGRESS (Funda individual perceptus identified by masker present test of each with Security Classification code) 23. (U) To determine the cause of the dysproteinemia observed following burn injury and to determine if the more marked dysproteinemia seen in the presence of infection of the burn wound is an effect caused by some action of the bacteria. It is hoped that this will aid in understanding similar changes which are observed in burned soldiers.

24. (U) Incorporation of  $(2^{-14}C)$  glycine into the serum proteins of groups of control, burned, burned-infected, and treated burned-infected rats has been measured. Body albumin distribution was determined by radioimmunoassay using extracts of tissues obtained on the sixth day postburn. Measurements of water content of skin during the early postburn period are being made. Exchange rates of albumin between the intravascular and extravascular compartments will be measured.

25. (0) 75 07 - 76 09 Groups of burned rats (20% body surface burn) untreated, or treated one hour postburn by subcutaneous injection of hyaluronidase or saline into the burn wound area all had lowered plasma volumes at 6 hours postburn. When provide, with food and water ad lib, but given no further treatment, the plasma volumes of all groups had returned to normal at 24 hours and were greater than normal at 48 hours postburn. The water content of the burned skin was elevated at 6 hours and was. similar for all groups. The water content of the burned skin remained elevated at 24 and 48 hours postburn but was consistently lower in the skin of rats treated with hyaluronidase than in the skin of the salir; treated rats or of those which received no treatment.

#### TERMINATION

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: STUDIES OF DISTURBANCE OF PROTEIN TURNOVER IN BURNED TROOPS - USE OF AN ANIMAL MODEL

# US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Wanda L. Brown, MS Eleanor G. Bowler, PhM Arthur D. Mason, Jr., MD

Report Control Symbol MEDDH-288(R1)

Unclassified

#### ABSTRACT

#### PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

REPORT T E: STUDIES OF DISTURBANCE OF PROTEIN TURNOVER IN BURNED TROOPS - USE OF AN ANIMAL MODEL

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in report: 1 July 1975 - 30 September 1976

Investigators: Wanda L. Brown, MS Eleanor G. Bowler, PhM Arthur D. Mason, Jr., MD

Report Control Symbol MEDDH-288(R1)

The carcass and the unburned skin of burned-infected rats had reduced levels of albumin. This was in contrast to the tissues of burned rats without infection in which the albumin content was essentially normal. The albumin content of the burned skin was elevated in both groups of rats.

Groups of burned rats (20% body surface burn) which were untreated, or were treated at one hour postburn by subcutaneous injection of hyaluronidase or saline into the burn wound, all had lowered plasma volumes at 6 hours postburn. When the rats were given free access to food and water, but no further treatment, the plasma volumes of all rats returned to normal at 24 hours and were greater than normal at 48 hours postburn.

Water content of the burned skin of these rats were elevated at 6 hours postburn but was similar for all groups. The water content of the burned skin of all of the rats was still elevated at 24 and 48 hours postburn, but it was consistently lower in the burned skin of the rats which had been injected with hyaluronidase.

Protein Burn Trauma Turnover Rats Albumin Edema

#### STUDIES OF DISTURBANCE OF PROTEIN TURNOVER IN BURNED TROOPS - USE OF AN ANIMAL MODEL

From the results of earlier studies in which we used the incorporation of (2-¹⁴C)glycine into serum proteins of burned and burned-infected rats to estimate synthesis on the 6th day postburn, we concluded that decreased synthesis was not the cause of the prolonged hypoabluminemia typically observed following burn injury. Measurements of tissue albumin content demonstrated that the tissue albumin pool size of burned rats was greater than normal, and that the elevated albumin content of the burn wound accounted for the total increase in the pool size. Details of the procedures used and the results have been reported.

Since that report we have measured the albumin content of the individual tissues of a group of burned-infected rats. The mean albumin contents (mg/gm tissue) in the 3 groups of rats are shown in Table 1. In contrast to the burned rats which had essentially normal levels of albumin in carcass and in unburned skin, the burned-infected rats had reduced albumin content in those tissues. Despite this, the burned-infected wound area contained 3.5 times the amount of albumin in the unburned skin of the burned or control rats and contained 6 times the amount present in the unburned skin of burned-infected rats.

	Mg Albumin/gm Tissue (wet wt)					
	Control N=3	Burned N=5	Burned-Infected N=3			
Carcass	2.14 <u>+</u> 0.10	2.02 <u>+</u> 0.10	1.51 <u>+</u> 0.13			
Eschar		20.23 <u>+</u> 1.96	16.85 <u>+</u> 2.01			
Unburned Skin	4.88 <u>+</u> 0.16	4.98 <u>+</u> 0.14	2.81 <u>+</u> 0.30			
Tuta! Skin	4.88 <u>+</u> 0.16	10.33 ± 0.53	8.85 <u>+</u> 0.77			
		Mg Albumin/m	I			
Plasma	26.04 <u>+</u> 1.24	24.30 <u>+</u> 0.99	7.98 <u>+</u> 0.49			

Table 1. Plasma and Tissue Albumin Content of Rats

Values are means +S.E.M. Mean rat weights (6 days postburn) were Concrol, 196 gm; Burned, 209 gm; Burned-Infected, 149 gm. Preburn weights were 180-200 gm.

1. Brown WL, Bowler EG, Mason AD, Jr, Pruitt BA, Jr: Protein metabolism in burned rats. Amer J Physiol 231.476-482, 1976.

Both the wet weight and dry weight of the skin from the burned area were greater than that of an equivalent area of skin from unburned rats. The cumulative result of these changes is shown in Table 2. The skin of the whole 20% body surface burned area contained 4.68 ml more water and 216 mg more albumin than did an equivalent area of skin of control rats.

	Wet Weiç .: C .	Dry Weight Gm	Gm Water in Tissue	Mg Albumin in Tissue
W.nole eschar (20% BS burn)	i4.16	4.18	9.99	256.95
Control skin of equal area*	8.42	3.11	5.31	41.10
Difference (Burn-Control)	+5.74	<u>+</u> 1.07	<u>+</u> 4.68	<u>+</u> 215.85

Table 2. Comparison of Burned and Unburned Skin of 200 gm Rats

* Control rats were placed in the burning mold and the outline of the area was marked on the skin with ink. That area, which was equivalent to that of a 20% BS burn, was excised, weighed, and dried to constant weight. Values are means of 3 rats each group.

Additional groups of rats with a 20% body surface burn were untreatedor were given a subcutaneous injection of hyaluronidase (150 units in 1 ml saline) or of 1 ml 0.15 M NaCl into the burn wound at one hour postburn. Rats of all 3 groups showed lowered plasma volumes at 6 hours postburn. When given free access to food and water, but no further treatment, the plasma volumes of all 3 groups returned to normal by 24 hours and were greater than normal at 48 hours postburn.

Water content of the burned skin of these rats was elevated at 6 hours postburn and was similar for all groups. The water content of the burned skin of all of the rats was still elevated at 24 hours and at 48 hours, but it was consistently lower in the burned skin of the rats which had been injected with hyaluronidase. This data will be reported later.

#### PRESENTATIONS

Brown WL: Albumin in burned rats. 8th Annual Meeting American Burn Association, San Antonio, Tx. 2 April 1976.

#### PUBLICATIONS

Brown WL, Bowler EG, Mason AD, Jr., Pruitt BA, Jr.: Protein metabolism in burned rats. Amer J Physiol 231: 476-482, 1976.

				-	-		-		-	
RESEARCH AND TECHNOLOGY WORK UNIT SUMLARY			DA	OC 691	78	76 10 0	1	DD-D	RAE(AR)436	
& BATE PREV SUPPT	4. SHO OF SUBBARY	1. SUMMARY SCTT	L TOPIC SECURITY	7. 16.00	Aller		IN NUTR'S	Manc	DATA-	. LEVEL OF SUS
75 07 01	H. TERM	σ	U	NA			SL	[? ves [	],,,,	A THE UST
18. 10./COOTS:*	PROGRAM ELEMENT	PROJECT	NUMPER	TARK /	AREA HUR	-	IDRE UNT HUNDER			
- PREMARY	611027	3A161102E	71R	01 300						
a. Contracting	L!	[		L				8		
C CONTRACTING								4.170 mil	ciar i	
Efficacy of	Standard Hig	h Protein 1	ition of Ga Diet in Bur	stro	intest Soldie	tina ers	l Absorp (44)	tion an	d Nut	ritional
12 SCIENTIFIC AND THE	DINOLOGICAL ANEAT						_	_		
003500 Clin	1cal Medicine									
11. START DATE		14. ESTIMATED COM	LETION DATE	118. 70000		CY		K PERFORMANCE WETHOD		
69 07		76 09		DA			<u> </u>	C. I	n-Hou	se
T. CONTRACT/BRANT	Not Appli	cable		18. 765	ounces es	THATE	-		-	IDE (In these and y
& DATES/EFFECTIVE:		EXPIRATION:	j		76	1 <b>4</b>	.7		2	3
L SUNDER:			ļ	FIECAL	7T		.3			7
C TYPL:		4 ANDWRT:		YEAR	CUMMENT				T	
& XIND OF ATARD:		L.CUM. AMT.					<u> </u>		<u> </u>	
IS RESPONSIBLE DOD O	MEAN ZATION			IR. PER		NGAN ZI	1784			
NAME.* US Army Institute of Surgical Research       NAME.* US Army Institute of Surgical Research         Burn Study Branch       Burn Study Branch         ADDREM.* Fort Sam Houston, Texas 78234       Port Sam Houston, Texas 78234										
PRESPONSIBLE WOIVIOUAL     PREMICIPAL HIVESTIGATOR (Provide MEAN HULE Account pointume)       NAME:     Basil A. Pruitt, Jr, COL, MC       TELEPHONE:     512-221-5712       Social Security Account Number										
FOREIGN INT	ELLIGENCE NOT	CONSIDEREI	>	A 500CIAT HANE: HANE:	John	F. I	eterson	, 1LT, 1	ISC	DA
II, KEYNDROS (Protodo 1	TVCE AIM BOORINA CHOREIED	ellen (selle)								

(U) Gastrointestinal absorption; (U) High protein diet; (U) Trace elements; (U) Humans

23. (U) To evaluate the nutritional efficacy of standard high protein diets, nutritional supplements and elemental diets in burned patients. To determine the hormonal and dietary factors which influence nitrogen balance in thermally injured troops.

24. (U) Use of defined diets or specific nutrient loading administered by the oral or intravenous route and determine nitrogen balance, amino acid patterns, or serum protein response.

25. (U) 75 07 - 76 09 Nitrogen balance in burned man plateaus when the metabolic rate--carbohydrate ratio approaches unity. Addition of exogenous insulin can effect further ritrogen sparing. Phenylalanine flow increases in the postburn period as an index of muscle catabolism.

# TERMINATION REPORT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# **REPORT TITLE: EVALUATION OF GASTROINTESTINAL ABSORPTION AND NUTRITIONAL EFFICACY OF STANDARD HIGH PROTEIN DIETS IN BURNED SOLDIERS**

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Douglas W. Wilmore, M.D. John P. Peterson, Captain, AMSC

Reports Control Symbol MEDDH-288(R1)

Unclassified

# ABSTRACT

#### **PROJECT NO.** 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

**REPORT TITLE: EVALUATION OF GASTROINTESTINAL ABSORPTION AND NUTRITIONAL** EFFICACY OF STANDARD HIGH PROTEIN DIETS IN BURNED SOLDIERS

US Army: Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Douglas W. Wilmore, M.D. John P. Peterson, Captain, AMSC

Reports Control Symbol MEDDH-288(R1)

Improved and sustained dietary intake has continued as a major priority in supporting thermally injured patients. Additional dietary supplements and tube feedings have been developed in the metabolic kitchen by modifying commercially available liquid diet formulas. These enteral feedings are supplemented by hypercaloric glucose and amino acid infusions with or without fat emulsions. The specific role of dietary tyrosine and gastrointestinal absorption of other nutrients in injured patients have been studied and are presented in separate reports.

Gastrointestinal absorption High protein diets Trace elements Humans

				1. ABENC	Y ACCESS	ow.	2. DATE OF SUN	NARY	REPORT C	WTROL SYNBOL	
RESEARCH	AND TECHNOLOGY	WORK UNIT S	UMMART	DA	OE 69	82	76 10	01	DD-DR	LE(AR)636	
& BATE PREV SUMAY	4. KIND OF SUMMARY	S. SUMMARY SCTY	. WORK SECURITY	7. REGRA	DING	86 DR	ST'H INSTR'N	BA SPECIFIC	DATA-	LEVAL OF SUM	
75 07 01	K. COMP	U	U	NA		1	NL	A YES	10	A VORK UNIT	
10. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK AREA NUMBER WORK UNIT NUMBER							
. PRIMARY	61102A	3A161102	3A161102B71R 01 119								
L. CONTRIBUTING						1. 1. 1					
C. CONTRIBUTING								<u></u>			
11. TITLE (Proceds with	Beautity Closel Heatlan Code)	• (U) A The	erapeutic T	riai	of An	tac	id in Pr	e o itio	n of t	he	
Clinical Co	mplications As	ssociated I	With Gastri	c Muc	osal	Dis	<u>ease in</u>	Burned	Soldie	rs (44)	
12. SCIENTIFIC AND TE	CHHOLOGICAL AREAS										
003500 Clin	ical Medicine										
15. START DATE		14. ESTIMATED CON	PLETION DATE	IL FUND	ING ASENC	;Y		14. PERFORM	ANCE METH	OD	
74 01		76	06	DA				C. In	-House		
17. CONTRACT/GRANT	Not Applica	able		-	URCES ES	TIMAT	E A PROFESS	OHAL MAN YR	S & PUN	(In thousands)	
A DATES PRECTIVE:		EXPISATION:			PRECEDIN	•		-			
N HUMBER:*				FISCAL 76			.6		2	20	
S. TYPE:		& ANOUNT:		YEAR	CUNNENT						
& KIND OF AWARD:		f. CUM. AMT.									
10. AESPONSIBLE 200	DREANIZATION			30. PERF	0111111601	RGANII	TION				
"AME!" US Army	Institute of	Surgical	Research	NAME:*	US Ar	my	Institut	e of Su	rgical	Research	
-		-		Burn Study Branch							
Appendit"Fort Sam Houston, Texas 78234			4	ADDRESS!" Fort Sam Houston, Texas 78234							
	······································					-					
				PRINCIP	AL INVEST	IGATO	R (Puminh LEAN	il U S. Academi	e jastikuk <b>un</b> g		
NESPONS OLE INDIVIOUAL			MAME® Albert J. Czaja, MD								
HAME Basil A. Pruitt, Jr, COL, MC			TELEPHONE: 512-221-5712								
TELEPHONE: 512-221-2720			SOCIAL SECURITY ACCOUNT NUMBER								
EI, OEHEPAL USE ASSOCIATE INVESTIGATORS											
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	-	Davi	d K	. Teegar	den, MA	J, MC		
				HANE:	Basi	.1 A	. Pruitt	, JI, C	OL, M	C DA	
IL REVIONDE (Presede	BACH with Security Cheels	sation Code)									

(U) Curling's Ulcer; (U) Burned soldiers; (U) Gastritis; (U) Antacid

ES. TECHNICAL OBJECTIVE, * 34. APPROACH, 24. PROGRESS (Purnich Individual paragraphs Identified by munitor Procedu text of each with Sonathy Classification Code.)

23. (U) To determine if neutralization of hydrogen ions by antacid administration to burned soldiers who manifest disruption of the gastric mucosal barrier will prevent progressive gastric mucosal damage, gastric hemorrhage, or perforation.

24. (U) All patients admitted to the Institute of Surgical Research Within 72 hours after sustaining greater than 35% total body surface injury will be considered for this study. A lithium flux test will be performed as has been previously described within the 72-hour postburn period. The patients will then be randomly assigned to receive a standard antacid preparation or no neutralization of gastric acid. The patients with disruption of the gastric mucosal barrier who receive antacid therapy will be statistically compared as regards hemorrhage and perforation to those with disruption of the gastric mucosal barrier who receive no antacid therapy.

25. (U) 75 07 - 76 09 This project is now completed. Conclusions from the data show that antacids significantly reduce gastroduodenal mucosal complications following thermal injury when compared to a nontreatment group. Of 27 patients treated with antacids, only one complication developed. However, of the 27 patients in the non-treatment group, seven major complications, including perforation of the gastrointestinal tract and significant hemorrhage occurred.

#### FINAL REPORT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

# REPORT TITLE: A THERAPEUTIC TRIAL OF ANTACID IN PREVENTION OF THE CLINICAL COMPLICATIONS ASSOCIATED WITH GASTRIC MUCOSAL DISEASE IN BURNED SOLDIERS

#### US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

David K. Teegarden, M.D., Major, MC Joseph C. McAlhany, Jr., M.D., Major, MC Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

Unclassified

#### ABSTRACT

#### PROJECT NO. 3A161102B71R-01, RESEARCH IN BLOMEDICAL SCIENCE-SURGERY

REPORT TITLE: A THERAPEUTIC TRIAL OF ANTACID IN PREVENTION OF THE CLINICAL COMPLICATIONS ASSOCIATED WITH GASTRIC MUCOSAL DISEASE IN BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: David K. Teegarden, M.D., Major, MC Joseph C. McAlhany, Jr., M.D., Major, MC Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

This study prospectively investigated the effectiveness of antacids in the prevention of stress ulcerations following thermal injury.

The integrity of the gastric mucosal barrier (GMB) was assessed, using the lithium flux technique. Twenty one of the 54 study patients had abnormal GMB's. Twenty-seven patients received 60 cc of an antacid preparation containing a combination of aluminum hydroxide and magnesium hydroxide each hour; the remaining 27 received no antacid therapy. Clinical complications were compared in the two groups (Table 1).

1	HD	ւո	1

	Antacid Group	No Antacids
Total patients	27	27
Complications	7	1
Gastrointestinal hemorrhage	6	1
Perforation	1	0
۰. «۲۰۰۰» «۲۰۰۰» «۲۰۰۰» و ۲۰۰۰» و ۲۰۰۰		والا ومعتمله الكرياس والأالا المترسل والمحت فالمواقده

#### Conclustons

As can be noted in Table 1, the antacid group demonstrated a markedly decreased incidence of gastrointestinal complications when compared to the nontreatment group. Based on this data, this unit currently administers antacids for prophylaxis in the prevention of erosive gastritis and ulceration following thermal injury.

# PRESENTATIONS AND/OR PUBLICATIONS

McAlhany, J. C., Jr.: Antacid Control of Complications from Acute Listroduodenal disease after burns. The American Association for the Surgery of Trauma, Scottsdale, Arizona, 11 September 1975.

# PUBLICATIONS

McAlhany, J. C., Czaja, A. J., and Pruitt, B. A., Jr.: Antacid control of complications from acute gastroduodenal disease after burns. J Trauma 16:645-649, 1976.

Curling's ulcer Burned soldiers Antacid Gastritis

#### ANNUAL PROGRESS REPORT

PROJECT NO. 3A16!102B71R-01, RESEARCH IN BLOMEDICAL SCIENCE-SURGERY

REPORT TITLE: PULMONARY COMPLICATIONS IN THERMAL INJURY

ł

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Victor Lam, MD, Major, MC James K. Sims, MD, Major, MC Gary W. Welch, MD, PhD, Lieutenant Colonel, MC Peter A. Petroff, Jr, MD, Major, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BLOMEDICAL SCLENCE-SURGERY

REPORT TITLE: PULMONARY COMPLICATIONS IN THERMAL INJURY

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

investigators: Victor Lam, MD, Major, MC
James K. Sims, MD, Major, MC
Gary W. Welch, MD, PhD, Lieucenant Colonel, MC
Peter A. Petroff, Jr, MD, Major, MC

Reports Control Symbol MEDDH-288(R1)

A review of 39 completed patient records from July to September 1976 indicates that pulmonary complications continue to be a common problem in the burned patient. Of major interest to the clinician are the pathophysiology, early detection, and therapy of pulmonary injury associated with burns. Previous pulmonary studies in the burned patient utilized tests developed to quantify chronic obstructive lung and restrictive lung disease. These conventional measurements are inadequate for early detection of pulmonary edema, inhalation injury, and upper airway obstruction. In the next year, more specific methods will be required to evaluate alternate therapies in the pulmonary complications of the burned patient.

Inhalation injury Thermal Cojury Pulmonary complications Therapy

#### PULMONARY COMPLICATIONS IN THERMAL INJURY

A review of 39 completed patient records from July to Septemper 1976 indicates that pulmonary complications continue to be a common problem in the burned patient. Of major interest to the clinician are the pathophysiology, early detection, and therapy of pulmonary injury associated with burns.

#### Incidence of Inhalation Injury

Previous reports of the incidence of inhalation injury by standard clinical criteria is about 3%. With the advent of fiberoptic bronchoscopy and 133Xenon lung scans, the diagnosis of inhalation injury due to toxic fumes and/or products of incomplete combustion can be made earlier in the hospital course. Of the 39 patients reviewed, 15 had the diagnosis of inhalation injury for a 38% incidence. Fiberoptic bronchoscopy was performed at bedside on 20 patients and xenon scans, as a diagnostic procedure, on 12.

For inhalation injury, the clinical problems are accurate early diagnosis, possible acute upper airway obstruction, and therapeutic options. For inhalation injury with smoke inhalation, upper respiratory tract findings may not indicate degree of pulmonary parenchymal involvement. Serial determination of pulmonary interstitial fluid volume may be of prognostic usefulness.

Our current therapeutic regimen for inhalation injury includes intermittent positive pressure breathing, mucomyst, chest physiotherapy, occasional staroids, bronchodilators, and humidified oxygen. Aerosolized Gentamycin has been used according to protocol study. Among other possible therapeutic options available for evaluation are prophylactic CPAP, incentive spirometry, and synthetic surfactant. The current therapy is nonspecific and has not been evaluated in a prospective double-blind clinical study. At this time, there is little scientific evidence for the efficacy of IPPB in pulmonary disease.

#### Acute Upper Airway Obstruction

Acute upper airway obstruction has necessitated emergency airway management in six cases in the past 15 months and in two cases in the 39 cases reviewed.

The use of steroids has been advocated. However, quantitative evaluation of upper airway obstruction has been difficult. Proposed methods for evaluation of upper airway obstruction include the use of ultrasound probe to measure traches dimensions and oscillometric resistance measurements to determine nasal and trachesistance with gases of different viscosity.

#### Acute Respiratory Failure

Of 39 patients reviewed, 17 required the ventilatory support of a volume-cycled respirator during their hospitalization. All except three patients with tracheostomies were managed via an endotracheal tube. Many patients required high alrway pressures and post-expiratory pressure to maintain adequate ventilation and arterial oxygenation. The low pulmonary compliance was due in part to increased interstitial pulmonary fluid or pulmonary edema.

Postmortem findings on eight patients showed pulmonary edema and congestion, indicating fluid overload (with respect to pulmonary parenchyma) is a major factor in the pathogenesis of acute respiratory distress syndrome in these patients. It is suspected but not proven that increased capillary permeability plays a major role; however, other possibilities include the role of vasoactive substances and inhomogenous distribution of arteriolar constriction because of hypoxemia. Early measurement of pulmonary interstitial fluid volume would be useful in fluid management in patients prone to the acute respiratory distress syndrome.

#### Sulfamylon Topical Therapy

Three patients were suspected of having an adverse reaction to Sulfamylon topical therapy with metabolic alkalósis and hyperventilation. Despite the intermittent application and rotational use of Sulfamylon, the pulmonary complications continued to occur, perhaps in predisposed patients. Earlier detection of Sulfamylon hypersensitivity would be clinically valuable. 'Carbon dioxide respiratory center testing may allow differentiation of susceptible patients.

#### Other Complications

There were no reports of pleural effusion or pulmonary emboli in the period reviewed. Perhaps the absence of pulmunary emboli in this seemingly susceptible class of almost immobile patients is a protective result of a hyperkinetic metabolic effect on circulation. Pneumothorax occurred in three patients with two cases of latrogenic nature, secondary to central intravenous line insertion. Twelve patients had the clinical and laboratory diagnosis of bacterial pneumonia, including two cases of hospital-acquired aspiration pneumonia.

#### Inhalation Injury and Surfactant Therapy-

inhalation injury, associated with pulmonary failure, adds significantly to the morbidity and mortality of thermal injury. Even in the absence of inhalation injury, previous research has shown a significant decrease in pulmonary surfactant.

In this study, tracheostomized goats were subjected to smoke inhalation. Dynamic compliance prior to injury was compared to dynamic compliance following treatment with nebulized dipalmityllecithin or with the DPL solvent.

Histopathology showed inhalation injury in all animals; no post-treatment difference between the two groups was observed. The dynamic compliance, however, in the surfactant group increased 67.8% over control values while it decreased 9.58% in the solventtreated group. Although the number of animals was too small to achieve statistical significance, this experiment raises interesting possibilities as to the use of surfactant in the treatment of inhalation injury.

#### PUBLICATIONS AND/OR PRESENTATIONS

None
DECEMPON							1. DATE OF BURSARY REPORT CONTINUE STUDIE			
REJEARCH	AND TECHNOLOG	T WURK UNIT S	UMMARY	DA	OF 6390	5	76 10 01		DD-DR&E(AR)636	
S. BATE PREV SUMRY	4. KIND OF SUMMARY	S. SUMMARY SCTY	A. WORK SECURITY	7. REGI	ADING	14 DH	IP'H WSTR'N	A SPECIFIC	OATA-	D. LEVEL OF SUM
75 07 01	H. TERM	U	U U	N	A	N	r.	CONTRACTOR	ACCESS	A WORK WHET
18. NO./CODES:8	PROGRAM ELEMENT	PROJECT	NUMBER	TASK	AREA NUME			WORK UNIT	NUMBER	
a, PRIMARY	61102A	3A161102	B71R		01		316			
L CONTRIBUTING						-	80.80° 78	S 6.33000	872878	
C. CONTRIBUTING									0.38	
II. TITLE (Procedo with ;	penantly Classification Code	' (U) Evalu	uation of D	opam	ine (3	,4-7	Cehydrox	vphevle	thyla	mine) For
Treatment of	f Septic Shoc	k i ) Burned	l Troops (4	4)						
12. SCIENTIFIC AND TEC	CHNOLOGICAL AREAS									
003500 Clin	ical Medicine									
13. START DATE		14. ESTIMATED COMP	LETION DATE	IL FUN	DING ABENCI	1		IL PERFORM	HCE MET	NOD
75 02		76 09		D	A		1	C. I	n-Hou	se
TT. CONTRACT/GRANT	Not Applic	able		16. RES	OUNCES EST	MATE	A PROFESSI		L PUN	BE (In Annually
A DATES/EFFECTIVE:		EXPIRATION:			1 76				2	1
h Hungtai"				PISCAL 7T 2				<u>i</u>	6	
		E AMOUN*:		YEAR	COMMENT					
L KIND OF AVANDI		S. CUM. AMT.								
				30. PER	'ORINA CR	ANIZA	TION			
CAME!" US Army	Institute of	Surgical F	Research	HANE:*	JS Arm	, ïn	stitute	of Sur	rical	Research
		-		Clinical Division						
Fort Se	m Houston, To	e <b>xa</b> s 78234		Fort Sam Houston, Texas 78234						
					TOLE Sam HOUSLON, TEXAS /8234					
				PRINCIPAL INVESTIGATOR (Pumish BAR II U.S. Associate politicia)						
HAND BREIT A	Donaidet Ton			HAME:	Gary	:1-2	221 571			
TANK DODIL A	-221-2720	COL, MC		TELEP	HONE: U	12-	-221-3/1.	6		
TELEPHONE: JIA				SOCIAL	SECURITY A	ccou	NT NUMBER:			
BORTON THE	TLIGENCE NOT	CONSTREEM		ABOCIA	DOUG I		10 101 101-1			
		CONSIDEREL	,	HARE	Doug	.as	w. wilin	ore, MD		
IL REVERIES (Procedo B	ACH with Bonatily Classific	eifan (ode)		HARE:						DA
(II) Dopamine	. (II) Sentia	shock (II)	Burn into	****	(11) Car		-	/77\ 7	Jum e	
IS TECHNICAL OBJECTI	VE. # 14 APPROACH, 14 1	ROGRESS (Funnish Int	Avidual paragraphs (day	- <u>Y</u> j illied hy	number proce	du fezi	I of seek or the see		iumani	5
23. (U) To	evaluate the	use of dop	amine in t	reati	ment of	se	ptic sho	ock in h	ourne	a l
soldiers.										

24. (U) Patients meeting the criteria of reduced urine output, hypotension, and evidence of reduced central perfusion will have a Swan-Ganz thermal dilution cardiac output catheter inserted for determination of cardiac output. After baseline studies have been obtained, an infusion of dopamine will be administered intravenously at a rate to raise mean arterial blood pressure to 60 torr. Cardiac outputs will then be determined again.

25. (U) 75 07 - 76 09 Eleven patients received intravenous dopamine. Of the 11, only one survived the septic episode. Dopamine produced an increased cardiac index and reduced peripheral resistance at low doses. Higher doses also increased cardiac index and raised the systemic resistance when compared to low dose administration.

refights to accurations when articulation approved

# FINAL REPORT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

REPORT TITLE: EVALUATION OF DOPAMINE (3,4-DEHYDROXYPHENYL-ETHYLAMINE) FOR TREATMENT OF SEPTIC SHOCK IN BURNED TROOPS

# US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEX 15 78234

1 July 1975 - 30 September 1976

Investigators:

Gary W. Welch, MD, PhD, Lieutenant Colonel, MC Charles Drueck III, MD Robert W.J. Baird, MD, Major, MC Douglas W. Wilmore, MD

Reports Control Symbol MEDDH-283(R1)

UNCLASSIFIED

# ABSTRACT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

# REPORT TITLE: EVALUATION OF DOPAMINE (3, 4-DEHYDROXYPHENYL-ETHYLAMINE) FOR TREATMENT OF SEPTIC SHOCK IN BURNED TROOPS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Gary W. Welch, MD, PhD, Lieutenant Colonel, MC Charles Drueck III, MD Robert W.J. Baird, MD, Major, MC Douglas W. Wilmore, MD

Eleven extensively burned patients with an average age of 30.2 years and an average burn of 62.5% of the total body surface who developed septic shock have been treated with dopamine in dosage ranging from 4.6 - 46.1 mcg/kg/min. This agent further elevated an already increased cardiac index in these patients with little effect on mean arterial blood pressure. Pulmonary vascular resistance was slightly elevated while systemic vascular resistance was decreased with low doses of dopamine and essentially unchanged with higher doses. Both left ventricular and right ventricular stroke work were significantly increased following administration of dopamine. Only one of the 11 patients survived despite the improvement in cardiac performance brought about by dopamine. The high mortality in this group of patients reflects an inability to eliminate the source of sepsis in the patients studied.

Dopamine Septic shock Burn injury Cardiac output Humans

# EVALUATION OF DOPAMINE (3,4-DEHYDROXYPHENYLETHYLAMINE) FOR TREATMENT OF SEPTIC SHOCK IN BURNED TROOPS

Sepsis remains the most frequent cause of death in extensively burned patients. Septic shock, which occurs frequently in patients with gram-negative infections, is often refractory to treatment by fluid loading, which may only precipitate pulmonary edema or the administration of sympathomimetic agents, such as catecholamines, which may cause oliguria and even acute tubular necrosis. The use of a new vasoactive agent, dopamine, has been evaluated in a group of burn patients by assessing its ability to improve cardiac function while maintaining adequate peripheral and renal blood flow.

# Methods

Eleven patients with burns of more than 40% of the total body surface who developed hypotension and oliguria, unresponsive to volume loading, in the presence of a positive blood culture, have been studied (Table 1). Prior to the administration of any pressor agents, a Swan-Ganz 7f flow-directed thermal dilution cardiac output catheter was inserted either percutaneously or via a cutdown incision. If the patient's condition permitted a preinfusion, 2-hour creatining clearance was done. All pressures were transduced with a Trantec pressure transducer and displayed on a Tektronix 412 physiologic monitor. Cardiac outputs were done using an Instrumentation Laboratory thermal dilution cardiac cutput computer. Arterial pressures were determined by either a direct intra-arterial catheter or the infrasonde blood pressure monitor. Simultaneous systolic time intervals were recorded on an 8-channel Electronics for Medicine recorder, using standard limb leads for the EKG, a Statham strain gauge for the carotid pulse tracing, and a Hewlett Packard microphone for the phonocardiogram. Following baseline studies, dopamine infusion was begun at a dose of 4,6 mcg/kg/min. The studies noted above were then repeated. The dopamine infusion was then increased in 400 mcg/ min increments to a maximum of 46.1 mcg/kg/min until the desired response was obtained, another agent was used, or the patient expired.

# Results

Two hundred thirty-six serial determinations of cardiac output, pulmonary artery pressure, widge pressure, central venous pressure, blood pressure and pulse rate were performed before (Table 2) and during dopamine administration. The dopamine dosage employed ranged from 4.6 to 46.1 mcg/ kg with an average low dose of 7.25 mcg/kg/min and an average high dose of 24.2 mcg/kg/min. When given in low doses, dopamine increased the cardiac index, pulmonary vascular resistance, and left ventricular stroke work. These changes were accompanied by a decrease in systemic vascular

# TABLE 1

# DOPAMINE IN SEPTIC SHOCK

# PATIENTS

	TOTAL	LOW DOSE	HIGH DOSE
NUMBER	11	8	و
SURV I VORS	*	O	
MALE	7	۲	νı
FEMALE	4	Ś	4
AVERAGE AGE	30.18 + 3.70	30.38 <u>+</u> 5.18	28.33 + 3.02
AVERAGE % BURN	62.45 + 3.49	58.81 <u>+</u> 3.65	64.06 <u>+</u> 3.73
AVERAGE % 3°	34.64 + 5.74	31.12 <u>+</u> 7.27	32.22 + 6.80

# TABLE 2

# DOPAMINE IN SEPTIC SHOCK

<b>2</b> CHANGE	+ 55\$	- 42\$	<b>1</b> 89 -	- 42\$	- 46	+ 24
PRE-PRESSOR	4.95 +.77	51.75 +5.34	707.67 <u>+</u> 93.71	156 <u>+</u> 30.68	30.30 ±5.39	8.98 1.3
NORMAL	3.2	د <mark>1+</mark>	2180 <u>+</u> 210	270 145	ጜ፟፟፟፟፟፟፟	8.76 <u>+</u> 2
	С.І. (L/min/M ² )	MABP (mm Hg)	SVR Dynes-sec cm ⁵ .H ²	PVR Dynes-sec cm ⁵ . 42	LVSV Gram Meter M2	RVSW Gram Meter N2

resistance to an average of 17% below prepressor levels, with the lowest value recorded being 66 dynes sec/cm². M². The right ventricular stroke work increased significantly, reflecting the increase in pulmonary vascular resistance. In high doses, the cardiac index, systemic vascular resistance and the pulmonary vascular resistance increased, although none of the increases were statistically significant. The left and right ventricular stroke work increased significantly, reflecting the elevated cardiac index and increased vascular resistances. The mean arterial blood pressure increased by 23%.

Comparison of the response to low dose dopamine with high dose dopamine revealed that the higher dose increased systemic vascular resistance and left ventricular stroke work (Table 3). The 27% increase above low dose levels observed with the higher doses represented only a 5% increase of vascular resistance above prepressor levels. A 21% increase of mean arterial blood pressure was associated with administration of the higher doses of dopamine.

Low dose dopamine caused a 17% decrease in systemic vascular resistance in association with a 35% increase in pulmonary vascular resistance, with no effect on mean arterial blood pressure. In high doses, dopamine increased both systemic vascular resistance and cardiac index, resulting in an increase in average mean arterial blood pressure to 64 mm Hg, a 23% increase over prepressor levels.

# Discussion

In these 11 study patients with extensive burns and documented sepsis, hypotension, indicated by a mean arterial blood pressure 42% less than predicted, appears directly related to the altered vascular resistance, which was 68% below predicted. An incomplete compensation has been achieved in such patients by elevation of the cardiac index to levels averaging 55% above predicted values. This compensation, if untreated, will lead to progressive cardiovascular failure. When volume loading failed to ameliorate these alterations, dopamine was started to preserve renal blood flow while increasing vascular resistance and supporting cardiac function to reverse the shock state. In both high and low dosage, dopamine increased cardiac index, mean arterial blood pressure and pulmonary vascular resistance, although the elevations did not attain statistical significance. Low dosage dopamine appeared to decrease systemic vascular resistance.

The effect of dopamine infusion on the pulmonary resistance may be an important consideration in patients suffering from respiratory insufficiency and hypotension. An increase in resistance by redistributing pulmonary blood flow may actually increase ventilation-perfusion abnormalities. Dopamine produced a rise in pulmonary vascular resistance, even though pulmonary resistance was below normal prior to dopamine infusion, and this effect may limit

# TABLE 3

# DOPAMINE IN SEPTIC SHOCK

	PRE PRESSOR	LOW DOSE DOPANINE (7.25 ± 0.92 mcg/kg/min)	high dose dopanine (24.2 + 3.1 mcg/kg/min)
C.I. (L/min/M ² )	4.95 <u>+</u> .77	5.44+ <u>+</u> .43	5.48 <u>+</u> .39*
MABP (mm Hg)	51.75 <u>+</u> 5.34	52.88 <u>+</u> 5.74☆	64.08 +5.912
SVR Dynes-sec cm5.H2	707.62 <u>+</u> 93.71	584.88 <u>+</u> 68.01*	/45.38 <u>+</u> 65.31*
PVR Dynes-sec cm5.H2	156 <u>+</u> 30.68	210.62 +57.07*	189 <u>+</u> 23.45*
LVSV Gram Heter H ²	30.30 <u>+</u> 5.39	36.68 <u>+</u> 4.65*	40.80 <u>+</u> 6.01 ²
RVSV Gram Heter H ²	8.98 <u>+</u> 1.90	14.25 <u>+</u> 2.82 ²	13.63 <u>+</u> 1.82 ²

No significance
1. p .01
2 p .05

the clinical usefulness of the agent. In both low doses and high doses, dopamine significantly increased both left ventricular and right ventricular stroke work with this inotropic effect predominating and bringing about a transient improvement in the hemodynamic status of these patients.

This study has demonstrated that dopamine is capable of improving cardiac function in septic burn patients. Only at the higher dose levels, however, did it increase systemic vascular resistance and even then the elevation was not to significant levels. The decreased systemic vascular resistance characteristic of septic shock was, in fact, exaggerated by infusion of low doses of dopamine, which would, therefore, appear to offer no unique benefit to the septic burn patient. The mortality of 91% in this group of patients (one of 11 survived) reflects not the ineffectiveness of dopamine but an inability to eliminate the source of sepsis and the persistance of the deleterious effects of the septic process.

Fublications and/or Presentations

None

RESEARCH	AND TECHNOLOG	WORK UNIT S	UMMARY	AGEN	OF GI	101	2. DATE OF SUN	0.1	REPORT C	ONTROL SYNBOL
			In wood sections	DA	Ur ou	1 26	10 TO			
		SUBBART SCIT	WONK SECONIT	) REUN	NDING .			CONTRACTOR	ACCESS	A WORK INNT
75 07 01	H. TEKM	U	<u> </u>	NA NA		N.		U YES U		
-	61102A	331611028	710	100		BEN	305	WORK ON		
A CONTRIBUTING	011026	JALULLULL	/10	<u>+</u>			<u> </u>	••••••		
C CONTRIBUTING	łi			+						· .
11 TITLE (Procodo WICh	Security Cleaselication Code	(U) Prolo	ongation of	Skir	h Allo	gra	ft Survi	val by		
Immunosuppr	essive Therap	v (Upjohn )	ATG) in Sol	diers	s With	Ma	ssive Th	ermal I	njury	(44)
12 SCIENTIFIC (HO TE	CHNOLOGICAL AREAS	<u>1</u>	······ • ·····························							
003500 Clin	ical Medicine			_						
13 START DATE		14 ESTINATED COM	PLETION DATE	TIS. PUNC	HIG AGEN	CY		IS PERFORM	ANCE METH	100
75 04		76 09		DA				C. In-House		
17 CONTRACT/GRANT	Not Applic	able		18 NES	OURCES ES	-	& PROFESSIONAL MAN YRS		S & FUNC	% (In theysends)
A DATES/EFFECTIVE.		EXPIRATION:			76		.5		18	
S. HUNDER *				FISCAL	71			.2		7
C TYPE		& ANOUNT		YEAN						
. KIND OF AWARD		f. CUM. AMT.								
19 RESPONSIBLE COUR	DRGARIZATION	L		120 7 6 11		RUARI -	7~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~		raica	Pesearch
HAME US Army	Institute or	Surgical	Research	NAME "	05 A	rmy inia	INSLILUU Al Divie	ie or ou	II YICA.	L Vesentor
Dent C	Se standon f	7073	A		· Foi	rure	an Houst	JON TON	rae 7	8234
ADDRESS FORT S	Sam Houston, 1	'exas 1023	4	AUUNE	1. 1.01		din nouse	.011, 101		525.
				BINCIP	AL INVEST		- /Fumish BEAN I	i II S. Acadami	Institutions	
				NAME	Hug	h D.	Peterse	n, COL,	MC	
HANE Basil	A. Pruitt, Jr	COL. MC		TELET	-	512-	221-3301	-		
TELEPHONE 5]	12-221-2720	,, .		SOCIAI	SECURIT	* ACCO	UNT NUMBER			•
21 GENERAL USE		<u></u>		-	TE INVEST	IGATOR	<b>1</b> 5			
FOREIGN INT	FELLIGENCE NOT	CONSIDERE	D	NAME:	Gary	Wel	ch, LTC,	, MC		
				NAME	Doug	las	W. Wilmo	ore, MD		DA
21. KEYWORDS (Procede	BACH with somelly Classifi	collon Codo) (U)	Autograft;	(U)	Mesh	graf	t; (U) I	xcisior	OFT	ascia;
(U) Antithy	ymocyte globul	in; (U) Is	olation; (	U) La	minar	flo	w room;	(U) Homog	<pre>graft;</pre>	(U)Humans
23 TECHNICAL OBJECT	TIVE - 24 APPROACH, 25	PROGRESS (Pumish in	ndividual paragraphs ide	entilied by	number pr	ocodo la	st of pack with 3	curity Classifi	tellan Cede j	,
23. (U) T	The objective	of this pr	otocol is t	to se	e if ]	burn	ed soldi	ers wit	h ext	ensive
injuries ca	an be effectiv	ely treate	d with ant	ithym	ocyte	glo	bulin ca	using s	suppre	SSION OI
cellular im	mmuni <i>.</i> y and al	llowing pro	longation of	of al	logra	ft t	ake in a	ireas of	exc1	sion.
				• • •			•			avestor
24. (U) Pá	atients in the	15 to 40	age group v	with	ful1-	thic	kness ti	iermai .	injury	greater
than 60% of	f their body s	surface and	without s:	ignır	icant	pul	monary :	injury w	111 D	e illuuno-
suppressed	with antithym	nocyte glob	ulin and so	erial	ly ex	cise	ana yi	arteu v	Vitn a ⊦o rom	variable
autograft a	and then with	fresh homo	graft. The	e hom	ograi	twi	lli De a.	Llowed	becom	alli li So availe
place. The	e patient's in	nmunosuppre	ssion will	cont	inue	ana	as aonoi	f Sites	Decon	e avaii-

autograft and then with fresh homograft. The homograft will be allocated with autograft will be surgically excised and replaced with autograft. Immunosuppression will continue until all but 15% of the grafted surface is covered with autograft.

25. (U) 75 07 - 76 09 To date two patients have been studied; a 25 year-old female with an 85% thermal injury and an 18 year-old female with a 90% thermal injury. Both underwent excision and immunosuppression; however, both had a rather rapid septic course. It has been elected to defer any more trials with the immunosuppression until a patient isolator in the form of a vertical laminar flow unit has been obtained for the unit.

# TERMINATION

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

REPORT TITLE: PROLONGATION OF SKIN ALLOGRAFT SURVIVAL BY IMMUNOSUPPRES-SIVE THERAPY (UPJOHN ATG) IN SOLDIERS WITH MASSIVE THERMAL INJURY

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 1 July 1975 - 30 September 1976

Investigators:

Hugh D. Peterson, DDS, MD, Colonel, MC Douglas W. Wilmore, MD Basil A. Pruitt, Jr., MD, Colonel, MC

3

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

## ABSTRACT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

REPORT TITLE: PROLONGATION OF SKIN ALLOGRAFT SURVIVAL BY IMMUNOSUPPRES-SIVE THERAPY (UPJOHN ATG) IN SOLDIERS WITH MASSIVE THERMAL INJURY

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78:34

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Hugh D. Peterson DDS, MC, Colonel, MC Douglas W. Wilmore, MD John Whelchel, MD, Lieutenant Colonel, MC*

Reports Control Symbol MEDDH-288(R1)

The object of this study is to evaluate the prolongation of survival of allograft which has not been tissue typed. Massive burn wounds fatal by historic comparison, are to be excised in stages and covered with nontyped allograft from cadaver source. The patients are then immunosuppressed with antithymocyte globulin and the grafts allowed to take and remain in place. As donor sites become available the allograft is excised and replaced with isograft. This has been a feasible treatment in children with extensive burns. To date at the Institute of Surgical Research two patients have been managed in this fashion, neither has been a survivor and both have died of sepsis.

At present we are awaiting installation of a laminar flow controlled environment unit in order to better isolate the patient prior to further attempts. However it must be noted that we see very few patients with greater than 65 to 70% third degree burn without significant inhalation injury. The study will be resumed when the laminar flow capability has been acquired.

Autograft Mesh graft Excision of fascia Antithymocyte globulin

Isolation Laminar flow room Homograft Humans

*Wilford Hall Air Force Hospital, Lackland AFB, Texas

DECEMBEL AND TECHNOLOGY WORK LINIT CLIMMARY								NTROL STMBOL			
REPEARCH		- 40KK URII 3		DA C	XG 6955	L	76 10	UI	DD DR	E(AR)636	
76 10 01	4 KIND OF SUMMARY H. TERM	s sunnary scty" U	6 WORK SECURITY"	7 REGR. N	ADING" A	NI	I'N INSTR'N	D YES	ACCESS	LEVEL OF SUM	
10 NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK /	REA NUMB	ER		WORK UNIT	NUMBER		
. PRIMARY	61102 <b>a</b>	3A161102	371R	01	·	_	122				
S. CONTRIBUTING						_					
C. CONTRIBUTING		ļ		L							
(U) The Eff Balance Fol	ect of Dextro lowing Therma	se and Ami 1 Injury i	no Acid So n Burned So	lutio	ns on $1$ rs $(44)$	litr	ogen Ex	cretion	and N	litrogen	
003500 Clin	CHHOLOGICAL AMEAS [®] Lical Medicine	2									
13 START DATE		14. ESTIMATED CON	LETION DATE	TIL FUN	NING AGENCY			IS PERFORMA	HCE NETHO	0	
75 11		76 09		DA	1		1	C. In-	House		
17 CONTRACT/GRANT	Not Applica	ble		18 RES	DURCES ESTI	-	A PROFESSIO	HAL MAN YRS	. FUND	(In these anda)	
A DATES/EFFECTIVE		E> PIRATION:			PRECEDING		.7		17		
NUMBER.*							10				
4 TYPE:	TYPE: 4 ANOUNT- YEAR CURRENY										
& KIND OF AWARD		f. CUM. AMT.									
IS RESPONSIBLE JOD C	REANIZATION			20 PERI	ORMING ORG	ANIZA	TION				
HANE!" US ATMY	Institute of	Surgical R	esearch	wwww.US Army Institute of Surgical Research							
		2		Burn Study Branch							
ADDRESS!" Fort	Sam Houston,	Texas 782	34	ADDRES	" Fort	San	n Housto	n, Texa	s 782	34	
	-										
				PRINCIP	AL INVESTIG	ATOR	Fumioh BEAN II	US Anndemic	(nelifiction)		
RESPONSIBLE INDIVIDU	AL			NAME	Will:	Lam	S. McDo	ugal, M	AJ, MC	:	
NANG Basil A	. Pruitt, Jr,	COL, MC		TELEPHONE 512-221-2943							
TELEPHONE: 51	2-221-2720			90C(AI	SECURITY A	ccou	NT NUMBER				
21 GENERAL USE				A550CIA	TE INVESTIG	ATORE					
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	NAME:	Cle	nent	t L. Sla	de, CPT	, MC		
				NAME.	Dou	las	s W. Wi	lmore,	MD	DA	
ZL KEYWONDS (Frecade	SACH with Security Classifi	collon Code) (U)	Nitrogen b	alanc	e; (U)	The	ermal in	jury; (	U) De	trose;	
(U) Amino a	cid solutions	s; (U) Huma	ns								
23 TECHNICAL OBJECT	IVE, 24 APPROACH, 25	PROGRESS (Furnish I	ndtvidual paragraphs id	millied by	number Preci	de (*2)	i ei each wi <b>d</b> Se	curity Classifics	illan Code )	<b>.</b>	
23. (U) To	determine th	ne efficacy	of dextro	se ad	minist:	rati	ion as o	pposed	to pro	tein	
administrat	ion in the fo	orm of amin	o acid sol	ution	s. Th	e ei	ffect th	at one	or bot	n ot	
these solut	ions have on	nitrogen h	alance wil	l be	determ	ine	d and ev	aluatio	n will	be made	
of the solu	tion or combi	ination of	solutions	which	will	be i	most app	ropriat	e tor		
preservatio	on of nitrogen	n balance i	n the ther	mally	injur	ed s	soldier.				
24. (U) Th	ne patients su	istaining g	reater that	n 30 <del>1</del>	total	pod	dy surfa	ce area	burn	who are	
admitted to	the Institue	te of Surgi	.cal Resear	ch wi	ll be	deer	med suit	able fo	r the	study	
providing t	hey have no p	prior histo	ry of live	r dis	ease,	inte	estinal	disorde	rs or	other	
chronic deb	bilitative dis	seases. Th	ese patien	ts wi	.11 be	rand	domized	and t	y will	receive	
either dext	rose alone an	nd will hav	ve daily 24	-hour	urine	cl	earances	for u	ermin	nation of	
nitrogen lo	oss or they wa	ill receive	e dextrose	plus	amino	acio	d soluti	ons or	amino	acid	
solution al	lone. The row	utine blood	ls which ar	e nor	mally	drai	wn will	provide	the i	necessary	
serum infor	mation. The	routine un	ine collec	tions	which	ar	e obtain	ed will	be a	nalyzed	
for total r	nitrogen and a	amino acid	excretion	levcl	s in a	ddr	tion to	those s	ubsta	1065	
which are r	normally obta:	ined.									
25. (U) 75	5 11 - 76 09 '	Iwenty-two	patients w	ith h	ody su	rfa	ce therm	ual inju	ries o	of 30% or	
greater hav	ve thus far b	een studied	l. The ini	tial	data 1	ndı	cates th	at calo	rie po	r calori	
low dose gl	Lucose and low	w dose amir	no acid sol	utior	ns achi	eve	sımılar	affect	s as i	iitrogen	
sparing. N	No support has	s been gair	ned for the	hype	othesis	th	at amino	nitroy	en alo	one	
provides a with respec	better nitro	gen balance Sparing:	e. Similar however, c	find ardia	lings w ne outp	ut -	appurer and live	it in se er funct	ptic j ion a	patients 5 measure	
by the Inde patients re	cyanine y ceiving only	excretion mino acid	test were solutions	. mar} . Ti	edly a iese de	lte fec	red in d ts could	pathol be cor	ogic i rected	ay in Fy	
adding gluc	cose to the in mixture.	nfusate. (	urrently,	studi	es are	be	ing cond	lucted t	o dete	imine	
"A rollable to contracto	we upon arteneter's epury	· * # }									
DD, #AR 4149	8 PREVIOUS AND 1498 1	EFITIONS OF TH 1 MAR 66 FOR	ILS FORM ARE OF ARMY SE ARE	218	1 U / FO 67E		-4V8 A ' N()	* 00			

.

•

# TERMENATION

# **PROJECT NO.** 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

REPORT TITLE: THE EFFECT OF DEXTROSE AND AMINO ACID SOLUTIONS CH NITROGEN EXCRETION AND NITROGEN BALANCE FOLLOWING THERMAL INJURY IN BURNED SOLDIERS

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 1 July 1975 - 30 September 1976

Investigators:

W. Scott McDougal, M.D., Major, MC Douglas W. Wilmore, M.D. With the technical assistance: Calvin P. Kennedy

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

# ABSTRACT

# PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCES

# REPORT TITLE: THE EFFECT OF DEXTROSE AND AMINO ACID SOLUTIONS ON NITROGEN EXCRETION AND NITROGEN BALANCE FOLLOWING THERMAL INJURY IN BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: W. Scott McDougal, M.D., Major, MC Douglas W. Wilmore, M.D.

Report Control Symbol MEDDH-288 (R1)

Near isotonic hypocaloric diets containing glucose and amino acids significantly diminish nitrogen loss in severely burned patients. The impact of both substrates on nitrogen balance was similar and additive. Indices of organ function improved with glucose addition to the diet. Glucose, together with amino acids, is the most efficacious hypocaloric diet for short-term alimentation in the critically ill patient.

Nitrogen balance Thermal injury Dextrose Amino acid solutions Humans

# THE EFFECT OF DEXTROSE AND AMINO ACID SOLUTIONS ON NITROGEN EXCRETION AND NITROGEN BALANCE FOLLOWING THERMAL INJURY IN BURNED SOLDIERS

Thermal injury, like trauma of other etiologies, results in a catabolic state which must be limited and ultimately reversed if recovery is to occur. A positive nitrogen balance may be obtained by the provision of adequate protein and calories. Unfortunately, in the immediate post traumatic period, gut motility is impaired and if nutrients are to be given, they must be administered intravenously.

Isosmotic solutions which provide calories are useful in limiting nitrogen loss. Such solutions may be administered by peripheral vein and are particularly advantageous in situations where lack of oral intake is of limited duration, difficult fluid and electrolyte problems require frequent alterations in infusion rate or electrolyte composition or frequent administration of intravenous medications is required. The purpose of this study is to define the composition of a near isosmotic solution with respect to amino acid and/or glucose content which will provide maximal effect per calorie in reducing nicrogen loss and preserving optimal organ and metabolic functions in traumatized patients with and without systemic infection.

#### MATERIALS AND METHODS

Thirty-two thermally injured patients with an average burn size of  $54.7 \pm 18.3\%$  and a mean age of  $35.2 \pm 7.2$  years were resuscitated with 3 to 4 cc lactated Ringer's per per cent burn per body weight in kilograms. Immediately following the 24 hour resuscitative period, the patients were divided into three groups. Nutrients were given isosmotically by vein and caloric intake was not permitted by mouth during the three to nine day study period. The first group (Group 1) received glucose free infusates either devoid of calories or containing amino acids (FreAmine 11, McGaw). Each patient was maintained on a constant dose for a 24 hour period; however, the quantity of amino acids administered on a day to day basis was randomly varied.

The second group (Group II) was given either 60 or 120 grams glucose per  $m^2$  per 24 hours or 60 or 120 grams of glucose per  $m^2$  per 24 hours plus amino acids (Fre Amine II, McGaw). The quantity of aminonitrogen administered was maintained at a constant infusion rate for 24 hour periods; however, the amount of amino acids given from day to day was varied in a periodic random manner.

The third group (Group III) received 60 grams glucose plus approximately 11 grams amino-nitrogen (Fre Amine 11, McGaw) per m² per 24 hours continuously for three to nine days.

When a positive blood culture was obtained, the patient was permanently excluded from the above groups. During the course of therapy, 4 patients from Group I, 10 patients from Group II, and 1 patient from Group III had positive blood cultures at the beginning and termination of a 24 hour infusion period. These bacteremic patients provided the data for analyzing the effect of various infusions in the presence of systemic infection.

Daily blood cultures were obtained in all patients throughout the study. Plasma was analyzed every 24 hours for chloride, bicarbonate, blood urea mitrogen, creatinine, glucose (by the glucose oxidase method), calcium and inorganic phosphate utilizing the Auto-analyzer (Technicon); for sodium and potassium by flame photometry (International Laboratories 343); and for osmolality by freezing point depression (Advance Osmometer). Arterial blood was analyzed at least once a day for pH, pCO₂, pO₂ and O₂ saturation (Corning 165).

Blood samples drawn at the midpoint and end of each 24 hour period were immediately separated and the plasma stored at -15 to -20 degrees centigrade until assayed. The samples were analyzed for insulin by a modification of the technique of Hales and Randle! and for free fatty acids by the method of Trout, Estes and Friedberg.² Plasma preserved with Trasyiol and sodium EDTA drawn at the same times and stored in the same manner was analyzed for glucagon by the immunoassay method of Unger utilizing the 30 K antibody.³

Twenty-four hour urine collections were obtained throughout the study period and analyzed for sodium and potassium by flame photometry (International Laporatories 343), for chloride, creatinine, glucose (glucose oxidase method), calcium, inorganic phosphate, and urea nitrogen with the Auto-analyzer (Technicon) and for total nitrogen utilizing the sulfuric perchloric acid digested method of Kjeldahl.

Intake and output as well as body weight were recorded on a daily basis. Patients who were hypotensive or required a cardiotonic drug to support blood pressure, had a prior history of diabetes, or had significant renal insufficiency were excluded from the study.

Calculations: Nitrogen balance was corrected for alterations in the blood urea nitrogen for each 24 hour period and expressed per square

1. Hales CN and Randle PJ: Immunoassay of insulin with insulinantibody precipitate. Biochem J 88:137, 1963.

2. Trout DL, Estes EH, Jr., Friedberg SJ: Titration of free fatty acids of plasma: a study of current methods and a new modification. J Lip Res 1:199, 1960.

3. Faluona GR and Unger RH: Glucagon in "Methods of Hormone Radio" immunoassay", ed. Juffee BM and Behrman HR. Academic Press, New York and London, 1974, pp 317-330.

meter of body surface area. The difference between the BUN at the inception and the termination of the study period was multiplied by 0.60 by the body weight in kilograms. This provided an estimate of nitrogen retained by the patient not reflecting protein synthesis and assumes that urea distributes throughout the body water in a concentration equal to that of plasma. The quantity was either added or subtracted according to whether accumulation or loss occurred from the total nitrogen excreted in the urine. This quantity was subtracted from the nitrogen intake and the result divided by the body surface area in square meters.

The non-urea nitrogen excreted in the urine is composed of uric acid, ammonia, creatinine, amino acids and peptides. If urinary creatinine, ammonia and uric acid are relatively constant between groups as they were in this study, an estimate of the amino acids lost in the urine expressed as a function of those infused may be obtained by subtracting the urine urea nitrogen from the total nitrogen and dividing

the quantity by the amount of infused nitrogen  $(\underline{TUN-UUN})$ . An expression  $N_{IN}$ 

of the nitrogen composition of the urine is provided by dividing the urine urea nitrogen by the total nitrogen excreted in the urine (UUN/TUN). Glomerular filtration rate, fractional sodium and fractional potassium excretions, sodium, potassium and water balance were calculated in the standard manner. Caloric load was determined by multiplying the grams of glucose and/or amino acids in the infusates by 4 Kcalories.

Linear regression lines expressing nitrogen balance as a function of nitrogen intake were determined by the method of least squares. Significant differences for the slopes of the lines as well as the population of points described by the regressions between the different groups was determined by the analysis of covariance. The student t-test was used to determine significant differences between groups of data.

Two matrices were formed, one for non-bacteremic and the other for bacteremic patients. Carbohydrate (glucose) doses of 0, 60 and 120 grams per  $m^2$  formed one side of the matrix while amino-nitrogen doses of 0, 10 and 15 grams per  $m^2$  formed the other. The nitrogen balance as a function of a combination of these doses was determined by calculation u: izing the appropriate regression equation which had been obtained from the data of Groups I and II.

### RESULTS

Ten patients with an average total body surface area burn of  $48.7 \pm 19.7\%$  and a mean age of  $36.4 \pm 19.5$  years were infused with amino acid or electrolyte solutions devoid of glucose. Daily blood cultures were regative during each of the 24 hour balance periods depicted by the triangle's in Figure 1. The linear regression line illustrated in Figure 1 was fitted to the values by the method of least squares ( $r^2 =$ 

.7407) and is described by the equation: y = .7234x - 17.6798.

Ten patients with a mean injury of  $53.5 \pm 19.1\%$  of the body surface area and an average age of  $33.6 \pm 16.6$  years were given 60 grams glucose per m² as the sole caloric load per 24 hours or 60 grams glucose plus 5.59 to 20.15 grams amino-nitrogen per m² per 24 hour period. Blood cultures were negative for each of the daily balance periods depicted by the X's in Figure 1. The linear regression line determined by the method of least squares illustrated in Figure 1 has an r² = .8450 and is described by the equation: y = .6131x - 10.7070.

Six patients who had sustained a 61.7 ± 15.5% total body surface area burn and had an average age of  $47.3 \pm 14.7$  years were infused with 120 grams glucose per m² per 24 hours or 120 grams glucose plus 7.02 to 20.00 grams amino-nitrogen per m² per 24 hours. All blood cultures during each 24 hour balance period depicted by the circles in Figure 1 were negative. The linear regression line illustrated for the data in Figure 1 (r² = .8769) has the form of: y = .5820x -7.5343.

The slopes of the linear regression lines for the three groups are not significantly different; however, each line represents a distinct population and is significantly different from each of the other two regressions by the analysis of covariance. (60 grams glucose + AA versus 120 grams glucose + AA, p < .001; AA versus 60 grams glucose + AA, p < .001; and AA versus 120 grams glucose + AA, p < .001).

Six patients with an average total body surface area burn of 59.6  $\pm$  16.6% and a mean age of 24.7  $\pm$  9.0 years were infused by peripheral vein with 60 grams glucose plus 11.32  $\pm$  1.49 grams amino-nitrogen per m² per 24 hours continuously for three to nine days. All blood cultures up to and including the study period were negative. Twenty four hour nitrogen balances calculated on the final day of the study revealed an average balance of -3.80  $\pm$  1.05 grams per m². The value calculated from the linear regression equation of the 60 gram glucose plus amino-acid group for a nitrogen intake of 11.32 grams is -3.77 grams per m². The similar values support the validity of 24 hour dose schedules in Groups 1 and 11.

Four patients with a mean burn size of  $43.5 \pm 7.6$ % and an average age of  $24.0 \pm 8.1$  years had positive blood cultures (one for Serratia marcescens, 3 for Klebsiella pneumoniae) at the beginning and termination of the 24 hour infusion period. These patients received glucose free solutions, containing either no amino-nitrogen or 5.69 to 18.99 grams amino-nitrogen per m² per 24 hours. Each 24 hour balance period is depicted by triangles in Figure 2. The linear regression line for the data ( $r^2 = .6239$  illustrated in Figure 2 is described Ly  $\gamma = .4550X$ -24.0010.

Six patients who had sustained an average total body surface area burn of  $50.3 \pm 16.5\%$  and had a mean age of  $30.0 \pm 16.3$  years developed

Figure 1. Nitrogen balance per square meter body surface corrected for alterations in blood urea nitrogen as a function of nitrogen intake in nonbacteremic patients infused with either amino acids, 60 grams glucose per  $m^2$  plus amino acids or 120 grams glucose per  $m^2$  plus amino acids.



Figure 2. Nitrogen balance per square meter body surface corrected for alterations in blood urea nitrogen as a function of nitrogen intake in bacteremic patients infused with either amino acids, 60 grams glucose/  $m^2$  plus amino acids or 120 grams glucose/m² plus amino acids.



positive blood cultures for Klebsiella pneumoniae at the inception and termination of a 24 hour study period in which they had received 60 grams glucose plus 6.03 to 20.15 grams amino-nitrogen per m² per 24 hours. The individual balance data for each 24 hour period is depicted by the X's in Figure 1. The linear regression line  $(r^2 = .2524)$  for the data is described by y = .2817X - 13.3368.

Five patients with an average burn of  $66.6 \pm 15.4\%$  and an average age of  $40.4 \pm 16.8$  years had positive blood cultures (one for Staphylococcus aureus, 4 for Klebsiella pneumoniae) at the beginning and termination of a 24 hour period in which they had received 120 grams glucose per m² per 24 hours as the sole caloric source or 120 grams glucose plus 1.17 to 14.82 grams amino-nitrogen per m² per 24 hours. Each 24 hour balance period is demonstrated in Figure 2 by the circles. The linear regression line for these data has an r² of .6384 and may be described by y = .4047X - 15.4545. There was no significant difference in age or burn size among any of the groups.

The slopes of the three regression lines in Figure 2 are not significantly different nor is there any significant difference between the nitrogen balance data for 60 gram glucose plus amino acids and 120 gram glucose plus amino acids. An analysis of covariance reveals that the regression line of the population receiving amino-acids is significantly different from both the regression line of the 60 gram glucose plus amino acid (p < .001) or 120 gram glucose plus amino acid (p < .001) groups.

The slopes of the regression lines for non-bacteremic and bacteremic patients receiving the same infusate are not significantly different. The populations represented by the regression lines which received the same infusates for bacteremic and non-bacteremic patients were significantly different: amino acid group,  $p \le .001$ ; 60 gram glucose plus amino acid group,  $p \le .001$  and 120 gram glucose plus amino acids, p < .001.

Figure 3 illustrates nitrogen balance as a function of caloric load irrespective of source for both non-bacteremic and bacteremic patients. The linear regression line for the non-bacteremic group has an  $r^2$  of .7808 and may be defined by y = .0221X - 16.0954. The linear regression line for the bacteremic group  $(r^2 = .6405)$  is represented by y = .0196X - 23.4426. The slopes between the two lines are not significantly different but each line differs significantly from the other and represents a distinct population as determined by the analysis of covariance (p < .021).

Table 1 is a comparison of the 24 hour urinary data for 60 or 120 gram glucose plus amino acids versus glucose free amino acid infusates in bacteremic and non-bacteremic patients. Significant differences are seen in the non-bacteremic patients for the ratio of urine urea nitrogen to total urinary nitrogen, p < .005, the ratio of the quantity total

Figure 3. Nitrogen balance per square meter body surface corrected for alterations in blood urea nitrogen as a function of caloric intake irrespective of source for nonbacteremic and bacteremic patients.



urinary nitrogen minus urine urea nitrogen to nitrogen intake, p < .001, for per cent fractional sodium excretion, p < .001, and for elevation in blood urea nitrogen, p < .01. In the bacteremic patients, only the more pronounced elevation in blood urea nitrogen in the amino acid group differed significantly from the glucose plus amino acid patients (p < .01).

There were no significant differences in the 24 hour urinary data illustrated in Table I for the amino acid infusates between bacteremic and nonbacteremic patients. A comparison in the glucose plus amino acid groups between bacteremic and non-bacteremic patients demonstrates a significant difference for the fraction of excreted amino acids as a function of those infused p < .005, UUN:TUN, p < .05, GFR, p < .05, percent fractional sodium excretion, p < .05 and change in plasma blood urea nitrogen, p < .01. Water balance and sodium and potassium loads were similar for the various infusates in both non-bacteremic as well as bacteremic patients.

Table II depicts the plasma values obtained for the various infusates for insulin, free fatty acids and glucagon in bacteremic and non-bacteremic patients. Significant differences are observed in nonbacteremic patients between those receiving amino acid infusates alone and those administered glucose plus amino acids for insulin, p < .001, free fatty acids, p < .001, glucagon, p < .001 and insulin to glucagon molar ratio, p < .001. Bacteremic patients demonstrated significant differences between the two types of infusions for insulin, p < .001, and free fatty acids, p < .001. A comparison between bactermic and non-bacteremic patients receiving the same infusion reveals a significant difference for glucagon in patients receiving amino acid infusates free of glucose (p < .001) and insulin in patients receiving glucose plus amino acids, (p < .05).

The nitrogen balance may be expressed as a function of both the carbohydrate and amino-nitrogen input by forming a matrix. The nitrogen balance for a particular dose or combination of doses is calculated from the linear regression equations describing the data in Figures 1 and 2. Matrices for non-bacteremic and bacteremic patients are illustrated in Figures 4 and 5, respectively.

## DISCUSSION

A hypermetabolic response occurs in both the uninfected and infected traumatized patient. Negative nitrogen balance, energy expenditure and weight loss are of far greater magnitude in the critically ill than in the uninjured chronically starved individual. Indeed, more than 200 grams protein may be catabolyzed per day to satisfy energy demands in the traumatized patient. Left unabated, the catabolic response results in excessive erosion of body mass, impaired organ function, an irreversible disruption of homeostasis and death. The provision of sufficient nitrogen and calories can reverse the process and thereby

	AMING ACIDS	GLUCOSE & AMINO ACIDS	۲
TUN-UUN Nin		<u></u>	,
Non-bacteremic	0.419 <u>+</u> .181	0.125 <u>+</u> .099	< .001
Sacteronic	0.777 <u>+</u> .268	0.862 <u>+</u> .357	N.S.
UUN TUN			
Non-bacteremic	0.705 <u>+</u> .060	0.811 <u>+</u> .090	< .005
Sactoremic	0.647 <u>+</u> .132	0 0.704 <u>+</u> .125	N.S.
GFR ml/min/m ²			
Non-bacteremic	66.7 <u>+</u> 26.0	74.4 <u>+</u> 31.5	N.S. '
Becteronic	56.2 <u>+</u> 28.6	53.9 <u>+</u> 21.2	N.S.
NaFxEx %			
Non-bacteromic	0.670 <u>+</u> .194	0.142 + .012	< .001
Bacterenic	0.383 ± .211	0.299 <u>+</u> .182	N.S.
KFXEX X			
Non-bacteremic	22.01 <u>+</u> 8.42	19.34 <u>+</u> 7.07	N.S.
Bacteremic	29.37 <u>+</u> 19.90	25.11 <u>+</u> 8.81	N.S.
A BUN mgt/m ²			
Non-bacteremic	6.81 <u>+</u>	0.29 ±	
Bacteremic	9.16 <u>+</u>	1.42 <u>+</u>	< .01

# TABLE I. RENAL AND PLASMA DATA FOR NON-BACTERENIC AND BACTERENIC PATIENTS INFUSED WITH EITHER AMINO ACIDS OR GLUCOSE PLUS AMINO ACIDS.

	AMINO ACIDS	GLUCOSE & AMINO ACIDS	Ρ
FREE FATTY ACIDS (mEq/1)	<u> </u>		
Non-bacteremic	0.443 <u>+</u> 0.143	0.165 <u>+</u> 0.058	.001
Bacteremic	0.402 <u>+</u> 0.115	0.175 <u>+</u> 0.051	.001
(NSULIN (uU/m1)	****	, μα μα προστά του ματικού του ματικού του	
Non-bacteremic	5.24 <u>+</u> 0.98	68.00 <u>+</u> 10.53	.001
Bacteremic *	2.97 <u>+</u> 0.90	23.04 <u>+</u> 4.77	.001
GLUCAGON (pg/m1)			
Non-bacteremic	125.49 <u>+</u> 33.08	634.55 <u>+</u> 114.83	.001
Bacteremic	476.65 <u>+</u> 80.21	539.43 <u>+</u> 217.84	N.S.
INSULIN: GLUCAGON			
Non-bacteremic	.0267 <u>+</u> .0275	.2303 <u>+</u> .1925	.001
Bacteremic	.0087 <u>+</u> .0056	.0759 <u>+</u> .0815	N.S.

Ĵ

TABLE II. PLASMA FREE FATTY ACIDS, INSULIN, GLUCAGON AND INSULIN TO GLUCAGON RATIO FOR NON-BACTEREMIC AND BACTEREMIC PATIENTS INFUSED WITH EITHER AMINO ACIDS OR GLUCOSE PLUS AMINO ACIDS.

## FIGURE 4. NITROGEN BALANCE AS A FUNCTION OF DOSE OF INTRAVENOUS INFUSION OF CARBOHYDRATE AND PROTEIN ADMINISTERED ISOSMOTICALLY IN NON-BACTEREMIC PATIENTS.

		GLUCO	SE (grams)	
( sue		0	60	120
EN (gra	0	-17.68	-10.71	-7.53
41 TROG	10	-10.45	-4.58	-1.71
AMINO	15	-6.83	-1.51	-1.20

FIGURE 5. NITROGEN BALANCE AS A FUNCTION OF DOSE OF INTRAVENOUS INFUSION OF CARBOHYDRATE AND PROTEIN ADMINISTERED ISOSMOTICALLY

IN BACTEREMIC PATIENTS.

GLUCOSE	(grams)
---------	---------

( swe		0	60	120
EN (gr	0	-24.00	-13.34	-15.45
NITROG	10	-19.45	-10.52	-11.41
AMINO	15	-17.18	-9.11	-9.38

achieve an anabolic state which is necessary for full functional recovery. Critically ill patients are often incapable of eating; however, positive nitrogen balance and satisfaction of energy demands have been achieved by intravenous administration of hyperoncotic glucose and protein infusates.⁴ These solutions which require administration through a central line, can serve as a source of systemic infection, can result in life threatening metabolic derangements⁸ and are impractical in the immediate post injury period when fluid balance, electrolyte composition and blood volume are being restored. Near isosmotic solutions which can be administered by peripheral vein obviate many of the hazards of the hyperoncotic infusates and are more practical when acute fluid, electrolyte and volume abnormalities require correction. With their use, the severe catabolic response is limited but often not corrected since satisfaction of energy requirements which may exceed 3000 calories per day, require unacceptably large quantities of fluid when nutrients are given in near isosmotic concentrations.

Glucose infusions in doses up to 100 grams per day provide a protein sparing effect which is proportional to the amount administered. Greater quantities result in very little additional reduction in nitrogen losses.⁹ Amino acid infusions free of glucose have also been shown to reduce proteolysis. Indeed, several reports indicate that a more substantial nitrogen sparing effect is achieved when amino acids are administered alone as compared to infusions of dextrose or dextrose in combination with amino acids.^{10,11} "The proposed mechanism for this effect involves mobilization of endogenous fat stores as an alternative energy source. Infused dextrose causes an increase in circulating insulin which inhibits the release of free fatty acids from fat stores. The reasoning is that since sufficient quantities of free fatty acids are unavailable, energy demands are met by increased proteolysis. If dex-

4. Dudrick SJ, Wilmore DW, Vars HM and Rhoass JE: Long-term parenteral nutrition with growth, development, and positive nitrogen balance. Surgery 64:134, 1968.

5. Freeman JB, Lemire A and MacLean LD: Intravenous alimentation and septicemia. Surg Gynec Obstet 135:708, 1972.

6. Sillon JD Jr., Schaffner W, Van Way CW Ill, and Meng HC: Septicemia and total parenteral nutrition. JAMA 223:1341, 1973.

7. Curry CR and Quie PG: Fungal septicemia in patients receiving parenteral hyperalimentation. NEJM 285:1221, 1971.

8. Dudrick SJ; MacFayden BV, VanBuren, CT, Ruberg RL and Maynard AT: Parenteral hyperalimentation: metabolic problems and solutions. Am Surg 176:259, 1972.

9. Gamble JL: Physiological information gained from studies on the life raft ration. Harvey Lect 42:247, 1947.

10. Hoover HC Jr., Grant JP, Gorschboth C, and Ketcham AS: Nitrogensparing intravenous fluids in postoperative patients. NEJM 293:172, 1975.

11. Blackburn GL, Flatt JP, Clowes, GHA, and O'Donnell TE: Peripheral intravenous feeding with isotonic amino acid solutions. AM J Surg 125:447, 1972.

trose is omitted from the infusate and only amino acids are given, insulin levels remain low and large quantities of free fatty acids are then released which provide the necessary energy substrate. Therefore, lesser amounts of endogenous protein need be catabolyzed to supply necessary energy demands.¹² Confirmation of these observations requires dose response date in a single population for both glucose and amino acid infusates. Unfortunately, these data have heretofore been unavailable. It should also be apparent that the most appropriate near isosmotic infusate must not only provide superior nitrogen sparing when compared to other solutions but must also provide the necessary form of substrate for optimal metabolic activity and organ function.

Figure I illustrates dose response date for amino acid and amino acid plus glucose infusates. The validity of 24 hour dose schedules demonstrated in this figure was confirmed by obtaining almost identical values from patients who had received constant dose infusions spanning 3 to 9 days (Group III). These data indicate that calorie for calorie in a dose range appropriate for peripheral vein administration with acceptable fluid loads amino acids are no more effective than 60 grams glucose per m² plus amino acids in achieving nitrogen balance. Indeed, 60 grams protein per m² (9.6 grams nitrogen) gives the same nitrogen balance as 60 grams glucose per m² free of amino acids. Although 120 grams glucose per m² plus amino acids improves nitrogen balance over that achieved with 60 grams glucose per m² plus a comparable amino acid dose, it does so disproportionately less. Therefore, 60 grams glucose per m² plus amino acids achieves the greatest effect per gram.

Free fatty acid levels were significantly elevated and insulin concentrations significantly reduced in the amino acid infused patients when compared to those receiving amino acids plus glucose (Table II). The insulin and free fatty acid levels in the amino acid infused are comparable to those reported for patients who had received protein alone and were reported to demonstrate a superior nitrogen sparing effect.

In the face of elevated free fatty acids and decreased insulin levels in the amino acid group, we were unable to demonstrate any superiority in terms of nitrogen balance of amino aicd infusates over glucose + amino acids. The exclusion of glucose from 2 dietary regimen on grounds of nitrogen sparing, therefore seems unwarranted.

The observation that there was no superiority of amino acid over glucose plus amino acid infusions may be explained by the fact that our nitrogen balances were corrected for alterations in the blood urea nitrogen. Indeed, the BUN rose 6 to 9 mg % per m² in the amino acid group compared to approximately 1 mg% in the glucose plus amino acid patients.

^{12.} Blackburn GL, Flatt JP, Clowes GHA, O'Donnell TF and Hensle TE: Protein sparing therapy during periods of starvation with sepsis or trauma. Ann Surg 177:588, 1973.

(p <.01). If not taken into account, what appears as improved nitrogen balance, may in actuality be retained urea. It may be most significant in patients who are administered glucose after having been maintained on protein. Under these circumstances, large amounts of urea will be excreted in the urine merely a reflection of more effective clearing of retained urea.

The balance studies were not corrected for the increased nitrogen loss which occurs from the burn wound. Wound nitrogen losses have been measured during the period of eschar separation and exposure of granulation tissue;¹³ however, as long as the eschar remains intact, as it did in all study periods, losses are minimal. Since patients from each of the study groups had comparable burn sizes and ages, the losses cancel themselves when the three groups are compared; i.e., there is no quantitative difference between the groups; however, all regression lines must be shifted the same distance in a negative direction if actual nitrogen balance for an individual patient is to be determined from the graphs.

Catabolism is more pronounced in bacteremic patients as illustrated in Figure 2. The regression line for the same infusate in bacteremic patients is significantly lower (p < .001) than in non-bacteremics. The response to amino acids alone is much poorer than when glucose plus amino-acids are given. Indeed, 60 grams glucose per m² alone is comparable to 125 grams protein (20 gm N) in terms of affecting similar nitrogen balances. Sixty grams glucose per m² plus amino acids provided the maximal nitrogen sparing effect since no significant difference was noted between 60 or 120 grams glucose/m² plus amino acids. The matrixes provide a measure whereby the optimal utilization with the lowest dose of carbohydrate and amino acids may be determined. Both non-bacteremics and bacteremics achieve optimal effect per gram nutrient at a level of 60 grams glucose + 10 grams amino nitrogen/m²/ 24 hours.

Although nitrogen balance is an important indicator of which type of infusate is more desirable, crucial to the success of any nutrient program is adequate provision of proper substrate for support of the metabolic demands of the organism. Chronically starved patients who derive their energy needs from free fatty acids improve their hepatic function significantly when dextrose is administered.¹⁴ Liver function as determined by indocyanine green clearance is impaired during infusion of amino acid solutions alone and may be restored to normal by adding glucese to the regimen. The effects are similar to those observed in patients who have gram negative septicemia; however, unlike the ab-

13. Coroff HS, Pearson E and Artz CP: An estimation of the nitrogen requirements for equilibrium in burn patients. Surg Gynec Obstet 112:159, 1961.

14. Demole M and Grosqurin J: Influence du glucose sur l'hyperbilirubinemie provoguee. Compt rend Soc de biol 116:12, 1934. normalities observed when amino acids are administered alone, hepatocyte function in ceptic patients cannot be corrected simply by administering glucose.¹⁵

Renal active transport processes for sodium and amino acid resorption are also impaired when amino acids serve as the sole exogenous energy source. In the face of similar sodium loads and sodium balances, fractional sodium excretion is diminished when glucose is added to the regimen. Thus addition of glucose results in more effective transport of sodium out of the filtrate. The kidney is better able to retain filtered amino acids as illustrated by the reduced fraction of infused amino acids excreted in the urine and the improved fraction of urea excreted to total nitrogen upon the addition of glucose to the infusate.

Total urinary nitrogen is comprised of ammonia, uric acid. creatinine, amino acids, peptides and urea nitrogen. Creatinine excretion, urinary pH and serum uric acic 'evels were similar in both the amino acid infused patients and those given amino acids plus glucose. Therefore, changes in TUN minus urea most likely reflect changes in the excretion of amino acids and peptides. Thus, TUN minus UUN divided by infused amino acids provides an estimate of the amount of amino acids lost expressed as a function of the amount infused. Also the ratio of UUN to TUN is a relative approximation of amino acids metabolized (Table I). Both ratios indicate more effective retention of amino acids and suggest greater incorporation into protein with lesser amounts degraded for gluconeogeneris when glucose is infused with the amino acids. Glucagon was elevated when compared to normal levels (75pg/ml) in the amino acid group in nonbacteremics but in bacteremics was markedly elevated and did not differ from values observed for patients infused with glucose plus amino acids. Glucagon stimulates hepatic glycogenolysis and gluconeogenesis, and therefore has been viewed as : "hormone of energy release."16

Insulin facilitates transport of amino acids into cells and hence their incorporation into protein. Glucose which accompanies the amino acids into the cells may well be necessary if the process of protein synthesis is to occur.¹⁷

In experimental animals glucagon is elevated after amino acid infusions and in normal subjects diminished following glucose

15. McDougal WS, Wilmore DW, Pruitt BA Jr.: Glucose dependent hepatic membrane transport in non-septic and septic thermally injured patients. Abstracts Society for Academic Surgery, 1976.

16. Unger RH: Glucagon physiology and pathophysiology. NEJM 285: 443, 1971.

17. Wool LG and Krahl ME; Incorporation of C14 - histidine into protein of isolated diaphragms: interaction of fasting, glucose and insulin. Am J Physiol 197:367, 1959.

administration.¹⁶,¹⁸ Thus, glucagon modulates the amount of energy available according to the specific needs of the organism. The elevated levels observed in the bacteremic and nonbacteremic glucose plus amino acid groups suggest that the stimulatory effect of amino acids in a catabolic patient overrides the suppressive effect of glucose and that glucagon may well be required for optimal amino acid utilization.

The molar ratios of insulin to glucagon in all groups is less than 1 and indicates that the glucose flow across the liver involves gluconeogenesis primarily from amino acid precursors. The larger ratios observed in the glucose plus amino acid groups is supportive of quantitatively lesser amino acid breakdown in this group when compared to patients receiving glucose free amino acid infusates.

The stress of bacteremia in the severely traumatized patient appears to be a powerful stimulus for glucagon production, since bacteremic patients receiving amino acids demonstrated exceedingly high levels. Since this is obviously an extreme energy demanding situation, the glucagon response is to be expected. The relatively low level, although elevated above normal, observed in non bacteremic patients receiving only amino acids would therefore appear to be less than that needed for optimum surgery utilization.

The efficacy of amino acid solutions administered alone is unsubstantiated in these studies since their infusion when compared to 60 grams glucose/m² plus amino acids given in equal caloric loads provides similar nitrogen balances in nonbacteremic patients and markedly inferior balances in the critically ill bacteremic patient. These observations were made in the face of exceedingly low insulin levels and markedly elevated free fatty acid concentration in the amino acid group. The administered amino acids are not utilized as well'as an energy source when glucose is absent from the infusate as demonstrated by the reduced UUN:TUN ratio and the increased fraction of infused amino acids excreted in the urine. Indeed, the efficacy of low insulin levels is questionable since it is known that this hormone enhances transport of amino acids into cells and hence their incorporation into protein.¹⁶ The absence of glucose also impairs hepatic and renal transport processes.

# PRESENTATIONS AND/OR PUBLICATIONS

None

^{18.} Unger RH, Ohneda A, Aguilar-Parada E, and Eisentrout AM: The rate of aminogenic glucagon secretion in blood glucose homeostasis. JJ Clin Invest 48:810, 1969.

				1. AGENCY ACCESSION" 2. DATE OF				P BUIMARYS REPORT CONTROL STWROL		
RESEARCH	AND TECHNOLOG	r work unit s	UMMARY	DA C	DE 6954	1	76 10	01	DD-D	R&E(AR)636
S. DATE PREV SUN'RY	4. KIND OF SUMMARY	S. SUMMARY SCTY	. WORK SECURITY	7. REGR	DEFO	84 D'S		SA SPECIFIC	DATA	LEVEL OF SUN
75 07 01	H. TERM	U	U	1	IA		NL	A YES		A. WORK UNIT
10. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK A	REA NUMB	ER		WORK UNI		ŧ
	61102A	3A161102B	71R		01		302			
b. CONTRIBUTING	61101A	3A16110A9	lc	(	00		076	• • .•	·	
C. CONTRIBUTING								· · · ·	<u>.</u>	
11 TITLE (Procodo with	Security Classification Code	<u>ب</u>								
(U) Ex	cision of Esc	har in Burn	ned Soldier	s (44	4) •			-		
12. SCIENTIFIC AND TE	CHHOLOGICAL AREAS									
003500 Clin	ical Medicine									
13. START DATE		14. ESTINATED CON	PLETION DATE	IS FUN	NNG AGENC	¥		16 PERFORM	ANCE MET	CHOD
71 07		76 (	09	DA				C. In	-Hous	e
W. CONTRACT/GRANT Not Applicable				18. RES	18. RESOURCES ESTIMATE & PROFESSIONAL NAN YRS					IDE (In those ands)
& DATES/EFFECTIVE:		EXPIRATION			76		.2			3
N. NUMBER:*				FISCAL	7T		.1			1
G TYPE:		4 ANOUNT		YEAR	CUNNERT		1			
& KIND OF AWARD:		f. CUM. AMT.								
19. RESPONSIBLE DOD	DEGANIZATION			20. PER	ORMING OR	GARIZ	ATION			
HANE! IS ATT	Tretitute of	Surgical R	esearch	-	IS Arm	υ T1	nstitute	of Sur	rgical	Research
00 ALMY	THEFTCALC OF	ourgrour in	cocur on	Burn Study Branch						
ADDRESS!	am Nouston M	OV20 7973	4	ADDRENS [®] Fort Com Houston Moura 70224						
FOLCS	all houscon, 1	exas /025	-1	1	POLC	Sa	" nousco	MI, ICAL	13 /0	234
				PRINCIP	AL INVESTI	GATOR	(Fumish SEAN )	ll U S. Anednai	c (netitutia	•
RESPONSIBLE INDIVIDU	AL			NAME	Hugh	D.	Peterso	n, COL,	, MC	
NAME Basil	A. Pruitt, Jr	, COL, MC		TELEPHONE 512-221-3301						
TELEPHONE: 5	12-221-2720	• • •		SOCIAL	SECURITY	ACCO	INT NUMBER.			
21. GENERAL USE				ASSOCIA	TE INVESTI	GATOR	15			
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	NAME:	Barr	v A	. Levine	MAJ.	MC	
			-	HAME:						DA
ZI. KEVBORDS (Procedo	BACH -IS LOOSILY CLASSIN	callon Code) (11)	Laser: (U)	Bovi	e caut	erv	; (U) In	vesting	a faso	ia;

(U) Eschar; (U) Autograft; (U) Humans 13. TECHNICAL OBJECTIVE,⁶ 24. APPROACH, 28. PROGRESS (Pumlish Individual paragraphic identifiled by number Photodo leat of each with geountly Classification Cade.)

23. (U) To evaluate the carbon dioxide laser for use in excision of full-thickness burns with attention towards time involved blood loss, and graft take involved in such excisions on burned soldiers.

24. (U) Symmetrical excisions were performed on patients who were candidates for excision to the investing fascia. Laser excisions were compared to both scalpel and electrocautery excisions from the standpoint of blood loss and electrocautery excisions, time for excision, graft take and pre-and postop excision quantitative microbiology.

25. (U) 75 07 - 76 09 The laser was found to be superior to scalpel excisions as far as blood loss, but was just as time consuming. The laser produced slightly less blood loss when compared to the electrocautery but took significantly longer time for the surgical procedure. The graft take in all three modalities was excellent. Because of the expense of the laser the surgical unwieldingness of the instrument and failure to demonstrate any significant difference from electrocautery, the laser in its present state was abandoned as an instrument for excision. If the machine can be made less cumbersome and less expensive, it should find a role in surgical excision. Sequential excision has been used to good effect in treatment of burns of the dorsum of the hand, especially those which are deep second-degree burns. Use of "meshed" graft to cover the excised wound bed results in maximum graft take and satisfactory long-term coverage and functional results.

Available to contractore upon originator's approval

DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 1488A I NOV 68 AND 1498 I I MAR 68 (FOR ARMY USEI ARE OBSOLETE

# TERMINATION

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY REPORT TITLE: EXCISION OF ESCHAR IN BURNED SOLDIERS

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 5 July 1975 - 30 September 1976

Investigators:

Hugh D. Peterson, D.D.S., M.D., Colonel, MC Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

PROJECT NO. 3A161102B71R-01, RESEARCH IN BIOMEDICAL SCIENCE-SURGERY

REPORT TITLE: EXCISION OF ESCHAR IN BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Hugh D. Peterson, D.D.S., M.D., Colonel, MC Basil A. Pruitt, Jr., M.D., Colonel, MC

Report Control Symbol MEDDH-288 (R1)

Studies prior to this reporting period demonstrated that the 100 watt carbon dioxide laser was not a practical instrument for excision to the investing fascia. The main disadvantages being its bulk and unwieldiness, plus cost.- It was in no way equivalent to the Bovie cautery except in excising large planar surfaces. With abandonment of the laser our attention has been focused more on the various modalities of excision and the refinement of patient selection criteria.

During this reporting period, 78 of 254 patients or 30% underwent excision procedure. These included excision to the investing fascia with immediate grafting, early sequential excision to viable tissue with delay in grafting, tangential excision with immediate grafting and delayed escharectomy.

Laser Bovie cautery Investing fascia Sequential excision Tangential excision Eschar

### EXCISION OF ESCHAR IN BURNED SOLDIERS

With abandonment of evaluation of the  $CO_2$  laser, attention has been focused upon patient selection for excision and evaluation of the various techniques of excision. A total of 78 patients have undergone some form of excision during this reporting period.

The criteria for excision at the US Army Institute of Surgical **Research are fairly well established.** There are three categories of patients, the first category is small burns either full thickness or deep second degree injuries that can be excised in their entirety and grafted immediately. This results in a closed wound that heals within two weeks, and greatly decreased morbidity of small nonlife threatening burns. The second category is burns in the 30 to 70% size that have a significant area or areas of full thickness or very deep partial thickness that lend themselves to excision. This does not decrease hospital time fair removes areas prone to burn wound problems. This is especially true if the legs or perineum are involved. A subcategory is burns of this size that have special areas that will benefit from excision by way of earlier return to function and a better final skin coverage. At our unit this is largely limited to the hands. The final category is burns greater than 70% with a large portion of full thickness injury. Historically these have done very poorly and conservative management with a topical agent offers no improvement in mortality, no matter what agent is used. In this final category some form of excision procedure seems at this time to be the only rational approach to attempting to increase survival.

The various types of excision being evaluated are, excision to the investing fascia with either immediate isografting or coverage with a biologic dressing, tangential excision and grafting, sequential excision to viable tissue and grafting, and lastly escharectomy either early or late.

Twenty-five patients were treated with excision to the investing fascia. There was a total of 35 procedures accomplished in these 25 patients. Seventeen patients were in the 40 to 70% burn size and there were six deaths in this group. This single figure means very little since it must be borne in mind that the patients subjected to an excision are those that have the largest areas of full thickness injury, There were a total of eight patients with burns greater than 70% that underwent formal excision with no survivers. In general the plan is to excise the patient as soon as they are hemodynamically stable from their resuscitation and when necessary to stage the excision so that never more than two extremities are done at one time. The excisions that do the best are those that can be isografted on fascia at the time of excision. It is of interest to note that all of the deaths in 40 to 70% burn size were at the upper end of that category and did not have sufficient donor sites to allow initial isografting. All the nonsurvivers in that group were subjected to the problems of repeated homograft changes. In the
burns greater than 70% none of them had available donor areas so that more than a small portion of the initial excision could be isografted. Both of these demonstrate the great difficulty of wound coverage in large excisions.

Tangential hand excisions have been carried out at the US Army Institute of Surgical Research for the past two and a half years. They follow the classic principles of tangential excision and immediate grafting and are limited to deep partial thickness injuries and full thickness injuries where there are no obvious deep structures involved. The hands are isografted immediately in all instances and when the criteria for patient selection are rigidly adhered to functional return is complete. The hands are excised as soon as the patient is hemodynamically stable post resuscitation and the procedure is not done in patients with burns in excess of 60% unless it is done in concurrence with some other form of excision requiring a general anesthetic. During this reporting period twenty three patients underwent tangential excision of the hands with a total of 34 hands being excised. There were five deaths in that age group, all in large burns that underwent hand excision concomitantly with other excision procedures. In the surviving patients those that had strict adherence to selection criteria demonstrated a complete return of function while those where the criteria were extended had various extensor defects.

Sequential excision to viable tissue with delayed grafting, more properly termed escharectomies, were carried out in 27 patients with a total of 31 procedures. The escharectomies applied in two instances. Early in the postburn course to rid the patient of eschar and start an earlier preparation of the wound for definitive grafting. Eleven patients were treated in this fashion early in the postburn course and there were seven deaths. These were all in burns greater than 60%. The early escharectomy was employed either because the burn wound did not lend itself to formal excision to the investing fascia. i.e. perineums, buttocks or other difficult areas to excise to the fascia. The higher incidence of mortality in like size burns would suggest that early escharectomy is not tolerated as well as excision to the investing fascia with coverage with a biologic dressing. The second instance in which escharectomy is performed is later in the postburn course to remove tenacious eschar. This is done in the third to fourth week postburn, 16 patients were subjected to this treatment during the last re- * porting period with no mortality. This is an excellent technique to remove tenacious eschar and hasten the time to grafting. We have been successful with the delayed escharectomy using either biologic dressings in the form of allograft or xenograft or simply coarse mesh gauze as sulfa soaks.

Excision in the truly large burns, those exceeding 65 to 70% is certainly a time consuming and difficult procedure, but with the

complete failure of topical agents or conservative measures to improve mortality in burns that size or larger it appears evident that some excision modality is at this time the only rational approach to improving survival.

# PUBLICATIONS AND/OR PRESENTATIONS:

\$

Levine NS, Salisbury RE, Peterson HD, Pruitt BA Jr. Clinical Evaluation of the Carbon Dioxide Laser for Burn Wound Excision: A Comparison of the Luser, Scalpel, and Electrocautery. J Trauma 15: 800-807, 1975.

Levine NS, Peterson HD, Salisbury RE, Pruitt BA Jr: Laser, Scalpel, Electrosurgical, and Tangential Excisions of the Third Degree Burns: A Preliminary Report. J Plast Reconstr Surg 56:286-296, 1975.

DA OD 6978       76 10 01       DDDMmanNA         DA OD 6978       76 10 01       DDDMmanNA         TS OT 01       D. CHANGE       U       U       NA       NL       DEVENTION       LUMEAR	RESEARCH	AND TECHNOLOGY	Y WORK LINIT S	UNHARY	1. AOEI	CY ACCERNIC	ľ	L DATE OF SU	MARY	REPORT	REPORT CONTROL STUBOL		
1 And Party Normal Finder December 1       1 During Party Normal Finder December	NEJEANCI				DA	OD 6978		76 10	01	DD-D	R&B(AR)636		
Disk Coords         Product Number         Fask Alla Number         Fask Alla Number         Fook UNIT Number           - Pressor         61102A         33161102BS05         00         081           - Conventione         61102A         33161102BS05         01         184           - Conventione         61102A         33161102BS05         01         184           - Conventione         61102A         33161102BS05         01         184           - Conventione         Based Status         Status         184         184           - Conventione         Based Status         Status         184         184           - Conventione         Based Status         Status         184         Status         184           - Status         Not Applicable         Based Status         Status         184         Status         184         184           - Status         Not Applicable         Based Status         Status         184         184         184         184           - Status         Not Applicable         Based Status         Status         184         184         184         184           - Status         Isset Status         Status         184         184         184         184	75 07 01	D. CHANGE	U U	U U	P. REGR	Adine" pi		NL	CONTRACTO		A WORK WHT		
a reason         61102A         351611102BS05         00         081           covernmentee         01         5184         01         5184           U. Unit (reast are term) Classified Card, (U) Evaluation of Synthetic Sheeting as Operating Room Dreimate Card, (U) Evaluation of Synthetic Sheeting as Operating Room Dreimate Card, (U) Evaluation of Synthetic Sheeting as Operating Room Dreimate Card, (U) Evaluation Card, (U) Evaluation Card, (U) Evaluation Card, (U) Military Burn Unit (U)         11 reason         11 reason         12 reason <td>10. NO./CODES:⁹</td> <td>PROGRAM ELEMENT</td> <td>PROJECT</td> <td>NUMBER</td> <td>TASK</td> <td>AREA NUMBE</td> <td>R</td> <td></td> <td>WORK UN</td> <td></td> <th>R</th>	10. NO./CODES: ⁹	PROGRAM ELEMENT	PROJECT	NUMBER	TASK	AREA NUMBE	R		WORK UN		R		
Lowerservise         61102A         3A161102B71R         01         184           Conversions         61102A         3A161102B71R         01         184           Conversions         61102A         3A161102B71R         01         184           Material For Use in a Military Burn Unit (44)         In Summary Conversions         In Summary Conversions         In Summary Conversions           Material For Use in a Military Burn Unit (44)         In Summary Conversions         In Summary Conversions         In Summary Conversions           Material For Use in a Military Burn Unit (44)         In Summary Conversions         In Summary Conversions         In Summary Conversions           Material For Use in a Military Burn Unit (44)         In Summary Conversions         In Summary Conversions         In Summary Conversions           Summary Conversions         In Summary Conversions         In Summary Conversions         In Summary Conversions           Summary Conversions         In Conversions         In Conversions         In Summary Conversions           Summary Institute of Surgical Research         In Summary Conversions         In Summary Conversions           Summary Institute of Surgical Research         In Summary Conversions         In Summary Conversions           Summary Institute of Surgical Research         In Summary Conversing Conversing Summary Conversions		61102A	3\$1611102	2BS05	0	)		.081					
<pre>c contrasting construction of synthetic Sheeting as Operating Room Dra Material For Use in a Military Burn Unit (44) Use Scientific and Technological Ansatz USESS (Construction of the State Sta</pre>	. CONTRIBUTING	61102A	3A161102	371R	0	L		194		an a bh			
11. Unit (Present all series (Case/Marine Case/Units)       (1) Evaluation of Synthetic Sheeting as Operating Room Dri Material For Use in a Military Burn Unit (44)         11. Unit (Present as reconcedent. astrong 003500 Clinical Medicine it first Marter Case/Units astrong m. Contract/Mart Not Applicable astrong trace astrong m. Contract/Mart Not Applicable astrong trace astrong m. Contract/Mart Not Applicable astrong trace astrong method astrong trace astrong trace astrong trace astrong method astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astrong trace astron	C. CONTRIBUTING					فالدينية الندار التدنية الكمال	ļ						
Material For Use in a Military Burn Unit (44)         003500 Clinical Medicine         11 MARKET STATUT         12 MARKET STATUT         13 MARKET STATUT         14 MARKET STATUT	11. TITLE (Procode with	Security ClocelRection Code	) [*] (U) Evalu	ation of Sy	ynthe	tic She	et	ing as (	Operati	.ng Ro	om Drape		
14 EXEMPTION ON THE TRADE CONCLETION DATE         15 FORM DATE         16 FORM DATE         16 FORM DATE         16 FORM DATE         16 FORM DATE         17 CONTRACT/SAMMT         16 FORM DATE         16 FORM DATE         16 FORM DATE         17 FORM DATE          18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         18 FORM DATE         <	Material Fo	or Use in a Mi	litary Bur	n Unit (44)	)				,,				
003500 Clinical Medicine       It Fundance Medicine         10 MARK MARK       Not Applicable       DAY       C. In-House         10 Confective       Exemation       It Fundance Methods       It Fundance Methods         10 Confective       Exemation       It Fundance Methods       It Fundance Methods         10 Confective       Exemation       It Fundance Methods       It Fundance Methods         10 Confective       Exemation       It Fundance Methods       It Fundance Methods         10 Confective       Exemation       It Fundance Methods       It Fundance Methods         10 Confective       Exemation       It Fundance Methods       It Fundance Methods         10 Confective       Amount       It Fundance Methods       It Fundance Methods         10 Confective       Amount       It Fundance Methods       It Fundance Methods         10 Confective       Amount       It Fundance Methods       It Fundance Methods         10 Confective       It Fundance Methods       It Fundance Methods       It Fundance Methods         10 Confective       It Fundance Methods       It Fundance Methods       It Fundance Methods         11 Confective       It Fundance Methods       It Fundance Methods       It Fundance Methods         11 Confective       It Fundance Methods	12. SCIENTIFIC AND TE	CHHOLOGICAL AREAS											
70 07       76 09       DW       C. In-House         * OCONTACT/SAMPT       Not Applicable       ************************************	003500 Clir	<u>ical Medicine</u>	-		TIS PUN	NNG AGENCY			IN. PERFOR	MANCE NE			
70 07/2000       1/2 07/2000       1/2 07/2000       1/2 07/2000       1/2 07/2000         A MODIFICATIONE       EXPRESSION       EXPRESSION<				-		. 1		1					
ADDERAFT THE ADDRESS OF A DEPARTMENT AND A DEPARTMENT OF A DEP	70 07		76_0	9		<u> </u>		1	<u> </u>	n-Hou	se		
A NUMBER. ² a YUEL a YUEL		Not Applic	Cable		16. HES	SUNCES ESTIN	ATE	A PROFESS	ONAL MAN Y		NCE (In Moundab)		
a type:       4 ANDUNT:       1       4         a type:       4 ANDUNT:       77       .2       7         a type:       1       77       .2       7         a type:       1       77       .2       7         a type:       2       7       .2       7         a type:       2       2       7       .2       7         a type:       2       2       2       7       .2       7         a type:       2       2       2       2       7       .2       7         a type:       2       2       2       2       2       2       2	h. uimera.*				PIECAL	76		.2					
a KND OF AWARD:       1. CUM. AMT.       77       .2       7         W. REFORMED FOOD CREAMEATION	G TYPE:		A ANOUNT:		YEAR	CUNNERY		<u>+ · · · </u> <u>4</u>					
<ul> <li>REMPONSIBLE DOD ORGANIZATION</li> <li>REMPONSIBLE DOD ORGANIZATION</li> <li>REMPONSIBLE DOD ORGANIZATION</li> <li>REMEATION DOLAL</li> <li>REMPONSIBLE NOTIONAL</li> <li>REMEATING ORGANIZATION</li> <li>REMEATIN</li></ul>	. KIND OF AWARD		& CUM. AMT.		1	77		.2		7			
<ul> <li>MARE⁶ US Army Institute of Surgical Research</li> <li>ADDREM⁶ Fort Sam Houston, Texas 78234</li> <li>ADDREM⁶ Fort Sam Houston</li> <li>ADDREM⁶ Fort Sam Houston, Texas</li></ul>		RGANIZATION											
<ul> <li>ADDATENS' Fort Sam Houston, Texas 78234</li> <li>ADDATENS' Fort Sam</li></ul>		Institute of	Surgigal B	ogoarch	MAME [®] US Army Institute of Surgical Research								
ADDRESS ⁶ Fort Sam Houston, Texas 78234 ADDRESS ⁶ Fort Sam Houston, T	Clinical Division												
<pre>FORE Sain Houseon, Texas 70234 REFORMENT Sain Houseon, Texas 70234 REFORMENT MULTION FORE Sain Houseon, Texas 70234 REFORE THE INVESTIGATION (Formula Base 102 - 221-2720 REFORE 512-221-2720 REFORE 512-221-2720 REFORE S12-221-2720 REFORE S12 REFORE S12-21-2720 REFORE S12 REFORE S12 REFORE S12 REFORE S12-21-2720 REFORE S12-21-2720 REFORE S12 REFORE S12 REFORE S12-21-2720 REFORE S12 R</pre>	ADDRESSI® Fort Sam Houston, Texas 78234												
<ul> <li>PRINCIPAL INVESTIGATOR (Foundat BEAR II U &amp; Actional positional)</li> <li>PRESPONDED E INDIVIDUAL</li> <li>MANNE BASIL A. Pruitt, Jr, COL, MC</li> <li>TELEPHONE: 512-221-2720</li> <li>POREIGN INTELLIGENCE NOT CONSIDERED</li> <li>PREVENE: Robert B. Lindberg, PhD</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE:</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE:</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE:</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE: Robert B. Lindberg, PhD</li> <li>MANNE:</li> <li>MANNE: Robert B. Lindberg, PhD</li> <l< td=""><td colspan="9">Fort Sam Houston, Texas 18234 Fort Sam Houston, Texas 18234</td></l<></ul>	Fort Sam Houston, Texas 18234 Fort Sam Houston, Texas 18234												
<ul> <li>RESPONSIBLE (NOUVDOAL MAKE BASIL A. Pruitt, Jr, COL, MC TELEPHONE: 512-221-2720</li> <li>IN GENERAL USE FOREIGN INTELLIGENCE NOT CONSIDERED</li> <li>REVEAL USE FOREIGN INTELLIGENCE NOT CONSIDERED</li> <li>REVEAL USE FOREIGN INTELLIGENCE NOT CONSIDERED</li> <li>REVEAL USE Infections; (U) Surgical drapes; (U) Military burn unit; (U) Operating room based infections; (U) Surgical drapes; (U) Surgical gowns</li> <li>U TECHNICAL OUTCUVE, IL APPROACH, IL PROPER (Control of the properties of a Spunbonded Olefin-cellulosic Lamin sheeting as surgical drapes and yowns. A decrease in bacterial seeding of operative wounds via drapes will minimize postoperative wound infections decreasing subsequer morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinica use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	PRINCIPAL INVESTIGATOR (Pumleh BRAN II U.S. Academic prolibulion)												
<ul> <li>NAME BASIL A. Pruitt, Jr, COL, MC</li> <li>TELEPHONE: 512-221-2720</li> <li>NO CAL SECURITY ACCOUNT NUMBER:</li> <li>ASSOCIATE INVESTIGATORS</li> <li>ASSOCIATE INVESTI</li></ul>		AL			NAME:	• Basil	A.	Pruitt	, Jr, C	OL, M	С		
TELEPHONE:       512-221-2720         BI. GENERAL USE       ANDOLASSE CONTRACTORS         FOREIGN INTELLIGENCE NOT CONSIDERED       NAME: Robert B. Lindberg, PhD         IN EXPRESSION (Notice LaCE of a Second Constitution Code)       (U) Military burn unit; (U) Operating room based         Infections; (U) Surgical drapes; (U) Surgical gowns       NAME:         IN TECHNICAL Code of a APPROACH. IN PROPERTY (Invitation Code)       (U) Military burn unit; (U) Operating room based         IN TECHNICAL Code of a APPROACH. IN PROPERTY (Invitation Code)       (U) Surgical drapes; (U) Surgical gowns         23. (U) Evaluation in terms of draping characteristics, absorbency, physician acceptance, and bacterial barrier qualities of a Spunbonded Olefin-cellulosic Lamin sheeting as surgical drapes and yowns. A decrease in bacterial seeding of operative wound infections decreasing subsequer morbidity and mortality in injured troops.         24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.         25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration	NAME Basil A.	Pruitt, Jr,	COL, MC		TELEP	HONE 51	2-	221-272	0				
FOREIGN INTELLIGENCE NOT CONSIDERED HARE Robert B. Lindberg, PhD HARE Robert B. Lindberg, PhD	TELEPHONE: 512	2-221-2720			BOCIA	SECURITY A	cou	INT NUMBER:					
FOREIGN INTELLIGENCE NOT CONSIDERED NAME: Robert B. Lindberg, PhD NAME: REVERSE (Provide EACH THE Search Closel Reveals (CU) Military burn unit; (U) Operating room based infections; (U) Surgical drapes; (U) Surgical gowns IN TECHNICAL OBJECTIVE. ⁹ 14 APPROACH, IN PROPERTY (Control and Property Closel Provide Closel Provide Closel Provide Closel	I. GENERAL USE				ASSOCIA	TE INVESTIGA	TOR	•					
<ul> <li>NAME:</li> <li>If KIVEOAGE //www.com/com/com/com/com/com/com/com/com/com/</li></ul>	FOREIGN INT	ELLIGENCE NO	CONSIDERE	D	NAME	Robert	В	. Lindb	erg, Ph	D			
<ul> <li>infections; (U) Surgical drapes; (U) Surgical gowns</li> <li>infections; (U) Surgical drapes; (U) Surgical gowns</li> <li>is reconstruction of terms of draping characteristics, absorbency, physician acceptance, and bacterial barrier qualities of a Spunbonded Olefin-cellulosic Lamin sheeting as surgical drapes and yowns. A decrease in bacterial seeding of operative wounds via drapes will minimize postoperative wound infections decreasing subsequer morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>					NAME:						DA		
<ul> <li>infections; (U) Surgical drapes; (U) Surgical gowns</li> <li>23. (U) Evaluation in terms of draping characteristics, absorbency, physician acceptance, and bacterial barrier qualities of a Spunbonded Olefin-cellulosic Lamin sheeting as surgical drapes and jowns. A decrease in bacterial seeding of operative wounds via drapes will minimize postoperative wound infections decreasing subsequer morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	II. KEYWORDE (Process)	EACH with Sometry Closer	(U)	Military b	urn u	nit; (U	)	Operatio	ng roon	base	đ		
<ul> <li>23. (U) Evaluation in terms of draping characteristics, absorbency, physician acceptance, and bacterial barrier qualities of a Spunbonded Olefin-cellulosic Lamin sheeting as surgical drapes and jowns. A decrease in bacterial seeding of operative wounds via drapes will minimize postoperative wound infections decreasing subsequer morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetrative</li> </ul>	infections;	(U) Surgical	drapes; (	U) Surgica.	1 gow	ns							
<ul> <li>23. (U) Evaluation in terms of draping characteristics, absorbency, physician acceptance, and bacterial barrier qualities of a Spunbonded Olefin-cellulosic Lamin sheeting as surgical drapes and jowns. A decrease in bacterial seeding of operative wounds via drapes will minimize postoperative wound infections decreasing subsequer morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	IS TECHNICAL OBJECT	IVE, 24 APPROACH, 21	PROGRESS (Fuminh II	velviduai paragrapha idi	ntilled by	number protec	No 703	ii ol oosh with S	courity Classi	Realism Code	• )		
<ul> <li>acceptance, and bacterial barrier qualities of a Spunbonded Olerin-cellulosic Samin sheeting as surgical drapes and jowns. A decrease in bacterial seeding of operative wounds via drapes will minimize postoperative wound infections decreasing subseques morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	23. (U) EV	valuation in t	erms of dr	aping char	acter	istics,	_a	bsorben	cy, pny	sicial	n Tamdu akar		
<ul> <li>sheeting as surgical drapes and jowns. A decrease in bacterial seeding of operative wounds via drapes will minimize postoperative wound infections decreasing subsequer morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	acceptance,	and bacteria	11 barrier	qualities	or a	spundon	ae	a Olerii	n-cerru	110810	aminated		
<ul> <li>wounds via drapes will minimize postoperative wound infections decreasing subsequer morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	sheeting as	s surgical dra	apes and jo	wns. A de	creas	e in ba	CT	eriai s	eeaing	or op	erative		
<ul> <li>morbidity and mortality in injured troops.</li> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	wounds via	drapes will n	ainimize po	stoperative	e wou	na inte	Cτ	ions de	creasin	ig sub	sequent		
<ul> <li>24. (U) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	morbidity a	and mortality	in injured	troops.									
<ul> <li>24. (0) Laboratory assessment of bacterial barrier of synthetic sheeting. Clinical use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	<b>24</b> (11) 7-		assess of	hastorial 1	hawwi	ar of a		thatia	abootir	~ ~	linical		
<ul> <li>use of drapes on burn patients to determine surgeon acceptability. Photographic documentation of draping characteristics, absorbency, and "run-off". Pre- and postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics.</li> <li>25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration</li> </ul>	24. (U) La	boratory asse	essment or	Dacterial I	Darri	er or s	yn ~-	chetic :	Bueech	19. C	Linicai		
postoperative cultures at margin of operative field. Temperature monitoring to, determine heat transmission characteristics. 25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration	use of drag	es on burn pa	actence to	determine :	surye	nou acce	a a	"municy	. FIIOU	ograp.	A		
determine heat transmission characteristics. 25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration	documentati	ion of draping	y character	f onersting		ancy, an	mn	araturo	monito	vring	4 to:		
25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration	postoperati	lve cultures a	it margin o	toristics	6 116	ita. le	тъ	eracure	MONTEC	TTIG			
25. (U) 75 07 - 76 09 Bacterial barrier properties of drape materials have been assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration	uetermine r	leat transmiss	Sion charac	CELISCICS.									
assessed and the results analyzed. Differences in penetration exist between organisms and between drapes but interaction is also significant. Overall penetration	25 /11 75	07 - 76 00	Bactorial	harrior pr	oner+	ios of	đ٣	ano mat	ariale	have	heen		
organisms and between drapes but interaction is also significant. Overall penetration	25. (U) /5	$\frac{1}{2} \frac{1}{2} \frac{1}$	baccertal	Difform	opert	n nenet	~~	tion ev	iet het				
- Oldanitama and hermeen arabea har inceraction to also alduratemic. Actail benefic	assessed di	nd between dr	anaryzeu.	nteraction	ie a	len ein	n i	ficant	Overs	ncen 11 ne	netration		
was 12-30% of test sites and differences between drames are not great suggesting as	ULYALLISUS C	of test sites	apes but I	rances hat	13 0 Wear	dranee	ar	e not a	reat er	adest	ing an		
was 12-500 of lest sites and differences between drapes are not great suggesting a	inadomiato	harrier funct	ion of all	samples	Redu	ction o	f	sheetin	g pore	size	 or		
"tighter" weave of material appear necessary to prevent passage of bacteria.	"tighter" "	warrier runce	rial annear	necessary	tor	revent	- pa	ssage o	f bacte	ria.			

AveilaNe to confrectors upon originators approval DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 1498A 10V 68 AND 1495 1 1 MAR 68 (FOR ARMY USE) ARE OBSOLETE 244

# ANNUAL PROGRESS REPORT

#### PROJECT NO. 3S1611102BS05, MILITARY BURN RESEARCH

REPORT TITLE: EVALUATION OF SYNTHETIC SHEETING AS OPERATING ROOM DRAPE MATERIAL FOR USE IN A MILITARY BURN UNIT

# US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Basil A. Pruitt, Jr., MD, FACS, Colonel, MC Robert B. Lindberg, PhD Arthur D. Mason, Jr., MD

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

#### PROJECT NO. 3S16111023S05, MILITARY BURN RESEARCH

## REPORT TITLE: EVALUATION OF SYNTHETIC SHEETING AS OPERATING ROOM DRAPE MATERIAL FOR USE IN A MILITARY BURN UNIT

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Basil A. Pruitt, Jr., MD, FACS, Colonel, MC Robert B. Lindberg, PhD Arthur D. Mason, Jr., MD

Reports Control Symbol MEDDH-288(R1)

The bacterial barrier properties of six coded samples of synthetic surgical drape material have been assessed. The penetration at 42 sites of each drape material by inocula of five organisms, Pseudomonas, Serratia, Klebsiella, E. coli and Staphylococcus, has been assessed using a test system developed at this Institute.

Penetration rates between drapes were significant only for <u>Klebsiella, E.</u> coli and <u>Staphylococcus</u>, but the variation within drapes was also significant, indicating that the variation may be either in the drape samples or the evaluation technique. No consistently "best" drape was identified. Mean penetration of drape samples by organisms ranged from 18 to 37%, with the penetration by <u>Pseudomonas</u> significantly higher than that for other organisms. Penetration of drape samples by the three organisms showing significant differences in penetration rates varied from 12 to 30%, indicating that all of the tested samples were imperfect microbial barriers and were penetrated too readily to afford significant protection. The findings suggest that two drape samples showed slightly better barrier properties and that any physical or processing characteristics common to those two samples should be the subject of further study and evaluation.

Military burn unit Operating room-based infections Surgical drapes Surgical gowns

## EVALUATION OF SYNTHETIC SHEETING AS OPERATING ROOM DRAPE MATERIAL FOR USE IN A MILITARY BURN UNIT

Commonly used surgical drapes are imperfect microbial barriers and as such may inadequately protect the operative wound of combat-injured soldiers from contamination by organisms from adjacent skin areas or the environment. Drapes made from synthetic fibers generally have poor draping characteristics, do not transmit water vapor and permit quantitative run-off of liquids. Earlier evaluations indicated that sheeting made from Spun-bonded Olefin possessed excellent microbial barrier properties and transmitted water vapor but not liquid water, but had marginal draping characteristics and still permitted liquid run-off. In an attempt to both minimize liquid run-off and soften the material, a single layer of Spunbonded Olefin was then sandwiched between two layers of absorbent cellulosic fibers. Such sheeting was better accepted clinically but the laminating procedure compromised the barrier properties of the sheeting and permitted the irregular passage of test bacteria when microbial transmissivity was assessed using laboratory techniques developed at this Institute. Spunbonded Glefin sheeting of varied thickness, produced by various fabrication procedures, is being evaluated to define those characteristics of a clinically acceptable surgical drape material which will provide a reliable bacterial barrier.

#### Methods

Two and one-half inch discs were cut from six coded samples of synthetic sheeting and sterilized in glass Petri dishes using ethylene oxide. Test strains of <u>Staphylococcus aureus</u>, <u>Pseudomonas aeruginosa</u>, <u>Klebsiella</u> <u>pneumoniae</u>, <u>Escherichia coli</u> and <u>Serratia marcescens</u> collected from clinical microbiology specimens were used as the inocula. Cultures of these organisms stored in suspensions of sterile milk at -70°, thawed and grown for 18 hours in trypticase soy broth, constituted the inoculum.

The sterile discs of sheeting were placed aseptically on fresh blood agar plates. Six drops of each test culture were placed on each disc, spaced equidistant at a distance of one cm. from the edge of the disc of sheeting material. The drops were left in place for four hours at room temperature. If the drops of inoculum did not spread and were still evident at the end of the four-hour exposure time, the remaining fluid was meticulously removed by aspiration with a sterile capillary pipette. The sheeting discs were then removed from the culture plates and the plates incubated at 37°C for 24 hours, following which the plates were examined for evidence of growth at each site of inoculum drop placement. Seven replicates were run for each test organism so that penetration of 42 individual inoculum drops could be assayed.

#### Results

The fraction and percentage of penetration of inoculum sites of each sheeting sample by each of the test organisms is shown in the table. Statistical assay of these results revealed no significant difference in penetration by <u>Pseudomonas and Serratia</u> organisms between drapes, but did reveal significant differences of penetration of the drape materials by the other three test organisms, <u>Klebsielia</u>, <u>F. coli</u>, and <u>Staphylococcus</u>. For the latter three organisms, the difference in drape penetration was significant at the one per cent level, but it should be noted that the interaction was also significant at the one per cent level.

#### Discussion

Chi square analysis of the test organism penetration of the six coded surgical sheeting samples revealed no systematic difference among samples based on sample code. The differences among drapes and among organisms were found to be significant but the interaction was also significant, necessitating analysis of drape penetration by individual organisms. Significant differences existed between drape penetration by Klebsiella, E. coli and <u>Staphylococcus</u>, but not by <u>Pseudomonas</u> or <u>Serratia</u>. The chi square within drapes was significant or nearly significant for all organisms, suggesting some real variation in either the drape samples or the evaluation technique.

Counting of positive penetration has been conservative, i.e., if a single area of bacterial colonies touches two sample sites, both were considered to have shown penetration, but it was possible that penetration had occurred at only one site with spreading of the colonies. Such a counting procedure may increase the variation in penetration observed beyond an expected value.

If the results of the <u>Pseudomonas</u> and <u>Serratia</u> testing are excluded, the interaction term remained significant and there was no difference between penetration by the remaining three organisms, i.e., <u>Klebsiella</u>, <u>E</u>. <u>coli</u> and <u>Staphylococcus</u>. The penetrations of each of these three individual organisms showed significant differences among drapes but the sampling technique itself, as previously noted, showed wide variation. Among those organisms where the difference in penetration between drapes was significant, Friedman two-way nonparametric analysis suggested there to be no consistently "best" drape. That analysis did suggest that if samples # 298-132-1 and #298-132-4, possess a common property imparted by either thickness or fabrication processing, further studies of that property should be made.

Mean penetration of drape samples by organisms range from 17.5% for Serratia to 36.5% by <u>Pseudomonas</u> organisms, with only the penetration rate by <u>Pseudomonas</u> being different from the other organisms. Determination of mean penetration by each drape material revealed lower rates of penetration of the two drape materials previously noted, both in terms of penetration by all organisms and penetration by Klebsiella, E. coli and Staphylococcus.

Organism:	Klebsie	dia sp	ц Ш	coli	Ps aeru	ginosa	Serr	atia	S aur	eus	Tot	als
Sample Code:	Pos	сно	Pos	а¢	Pos	đP	Pos	скр	Pos	d٩	Pos	aro
298-132-1	7/42	16.7	5/ 42	11.9	10/42	23.8	6 _{/42}	14.3	3/42	7.1	³¹ / ₂₁₀	14.7
298-132-2	17/42	40.5	6/ 42	14.3	15/42	35.7	6 _{/42}	14.3	\$/#2	14.3	50/210	23.8
298-132-3	8/42	19.1	12/ ₄₂	28.6	18 _{/42}	42.9	6/ 42	14.3	6/ 42	14.3	50/ ₂₁₀	23.8
298-132-4	1/42	2.4	3/42	1.7	11/42	26.2	5 _{/ 42}	11.9	14/42	33.3	^{34,} /210	16.2
298-132-5	9/ 42	21.4	15/42	35.7	19/42	45.2	10/42	23.8	11/42	26.2	64 / 210	30.5
298-132-6	10/42	23.8	19/42	45.2	19 _{/42}	45.2	11/42	26.2	8 _{/42}	19.1	⁶⁷ / ₂₁₀	31.9

,

Over-all penetration occurred in 12-30 % of test sites and the differences between drapes were not great, indicating that all the tested samples were penetrated too readily by the test organisms and are, in essence, inadequate microbial barriers. The findings suggest that a "tighter" sheeting is required, especially if organisms like <u>Pseudomonas</u> are to be excluded from passage in the presently employed test system.

T AGENCY ACCESSION ⁸							2 DATE OF SUBBARY						
RESEARCH	AND TECHNOLOGY	r work unlt s	SUMMARY	DA C	D 6	976	76 10 (	01	DD-D	R&E(AR)636			
1. DATE PREV SUWRY	4 KIND OF SUMMARY	S. SUMMARY SCTY	S WORK SECURITY	7 REGR	ADING	an D	SB'N INSTR'N	SA SPECIFIC I	DATA-	S LEVEL OF SUM			
75 07 01	D. CHANGE	U	U	NA			IL	Dres [	3 10	A. WORK UNIT			
10 NO /CODES *		PROJECT	NUMBER	TASK	AREA	NUMBER		WORK UNIT	NUMBE	3			
	61101A	3A161101	A91C	1	00		083						
b. CONTRIBUTING													
C. CONTRIBUTING													
11 TLE (Procodo with ;	Socurity Classification Code	) [•] (U) Studi	es of the l	Effec	t o	f Vari	ations (	of Tempe	ratur	e and			
Humidity on	Energy Deman	ids of the	Burned Solo	dier	ın	a Cont	rolled	Metabolı	c Roc	om (44)			
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS												
003500 Clin	ical Medicine												
13 START DATE		14 ESTINATED CON	PLETION DATE	IS FUN		GENCY		16 PERFORM	NCE WET	HOD			
71_07		Cont		DA				C. In	-Hous	se			
17 CONTRACT'GRANT	Not Applicat	Applicable IN RESOURCES ESTIMATE & PROFESSIONAL MAN YRS & FUNDS (In PROFESSIONAL MAN YRS & FUNDS (In						DE (In thousands)					
A DATES/EFFECTIVE EXPINATION 76 1.2 4						44							
NUMBER PISCAL 7T 0.3 5													
C TYPE		AL AMOUNT				-		-		2.4			
E KIND OF AVARD I. CUM AMT 77						7		<u>.0</u>		36			
THE RESPONSIBLE DOD ONGANIZATION													
"AME ''''''''''''''''''''''''''''''''''''					INAME US Army Institute of Surgical Research								
ACUMPSS *				Burn Study Branch									
Fort Sam Houston, Texas 78234				Fort Sam Houston, Texas 78234									
						PRINCIPAL INVESTIGATOR (Pumion SEAN II U.S. Academic Institution)							
					NAME Dougals W. Wilmore, MD								
RESPONSIBLE INDIVIDUAL					PHONE	512-3	21-5712						
NAME BASII A. Pruitt, Jr., COL, MC 512-221-2720					L SECL	UNITY ACCO	UNT NUMBER						
TELEPHONE         512-221-2720           21 GENERAL USE					TE IN	VESTIGATO	RS						
FORFICN INTELLIGENCE NOT CONSIDERED					NAME Arthur L. Mason, Jr, MD								
FOREIGN INTELLIGENCE NOT CONSIDERED					HAME BASIL A. Pruitt, Jr, COL, MC DA								
12 KEYWORDS (Freede Back with Seculty CloselReation Code) (U) Metabolism; (U) Heat loss; (U) Evaporative water loss;													
(U) Controlled environment; (U) Humans; (U) Critical temperature; (U) Temperature													
23 TECHNIC IL OBJECTIVE * 24 APPROACH, 25 PROGRESS (Furnish Individual paragrapho Identified E iumbar Procedo test of each with Security Classification Code )													
23, (U) To define the relationship between surface cooling and hypermetabolism in a													
controlled ambient environment, to determine the mediator of the profound hyper-													
catabolic r	cesponse follo	owing therm	mal injury,	and	the	mecha	anisms o	f stress	-indi	uced			
heat produc	ction in burne	ed soldiers	5.										
24. (U) 1	The use of a c	controlled	encironmen	tal s	stud	ly root	n to mea	sure met	abol:	ic rate			
at various	temperatures	; concomita	antly, meas	ureme	ents	ofwa	ater los	s, heat	prod	action,			
core temper	cature, mean s	skin tempei	rature, and	cald	ula	ition (	or heat	transfer	coe	rricients			
and router	of heat loss	Simultar	noously mo	asure	າມອາມ	its of	blood s	ubstrate	· ur	ine and			

25. (0) 75 07 - 76 09 Prostaglandin levels in the blood of burn patients were normal and the burn wound did not appear to be a site of synthesis which contributed to system levels. Serum from burn patients resulted in a febrile response when injected into the thermoregulatory area of rabbits, and the mediator substance is thought to be endogenous pyrogen.  $T_2$ , the active thyroid hormone, is low following injury, and  $T_2$ degradation appears to be toward alternate pathways, resulting in reversed T, synthesis, The significance of this enemical hypothyroidism is unknown. Glucose flow unimpaired by conury per secus dominished by supervening sepsise. Infusion of glucagon increases core temperature and metabolic rate in sectic patients and increases hepatic gluconeogene 15.

plasma catecholamines, blood hormone levels, and correlation of total bod- metabolism

L S E UM ARE AS CE E SH ANGE SE ARE 185 F I F RMS 498A 1 NOV 65 DD. **** 1498 ) 1

with energy demands are performed.

#### ANNUAL PROGRESS REPORT

PROJECT NO. 3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH

### REPORT TITLE: STUDIES OF THE EPFECT OF VARIATIONS OF TEMPERATURE AND HUMIDITY ON ENERGY DEMANDS OF THE BURNED SOLDIER IN A CONTROLLED METABOLIC ROOM

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Douglas W. Wilmore, MD Arthur D. Mason, Jr., MD Louis H. Aulick, PhD, Major, MSC Basil A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

Unclassified

# ABSTRACT

PROJECT NO. 3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH

REPORT TITLE: STUDIES OF THE EPFECT OF VARIATIONS OF TEMPERATURE AND HUMIDITY ON ENERGY DEMANDS OF THE BURNED SOLDIER IN A CONTROLLED METABOLIC ROOM

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Douglas W. Wilmore, MD Arthur D. Mason, JR., MD Louis H. Aulick, PhD, Major, MSC Basil A. Pruitt, Jr., Colonel, MD

Reports Control Symbol MEDDH-288(R1)

To determine the afferent mediator of the hypermetabolic response to injury, blood samples from thermally injured patients were analyzed for prostaglandin A, E, and F. Serum concentrations were unchanged from normals and femoral vein concentrations were similar in patients with and without leg injuries. Injection of serum from burn patients into the preoptic area of rabbits elicited a febrile response in 9 of 13 samples tested while serum from 6 normals evoked no response. The factor in the serum from burn patients was heat labile and limulus-lysate negative, suggesting the pyrogenic material is endogenous pyrogen.

Studies in 5 burn patients demonstrate the presence of clinical hypothyroidism following injury; there is a fall in serum  $T_3$  and a rise in the attenuated component, reversed- $T_3$ . The significance of this hypothyroidism is yet to be determined.

Endogenous pyrogen Prostaglandin Thyroid hormones

## STUDIES OF THE EFFECT OF VARIATIONS OF TEMPERATURE AND HUMIDITY ON ENERGY DEMANDS OF THE BURNED SOLDIER IN A CONTROLLED METABOLIC ROOM

Burn patients are hypermetabolic and hyperthermic. At any ambient temperature studied, core and mean skin temperatures in thermmally injured patients are elevated 1-2° above temperatures observed in normal individuals (1). When allowed to adjust ambient temperature, burn patients select a warmer environment than normals to achieve comfort, in spite of the fact that the patients have elevated core and mean skin temperatures (2). The ambient temperature selected is gen-erally related to burn size. Thermally injured patients may shiver in thermal neutral or warm environments (29-32°C), and patients with extensive thermal injuries rarely sweat, even in hot environments (>35°C). This evidence has been interpreted to suggest that thermal injury results in an elevation of the central reference temperature, and this apparent upward shift in setpoint activates the sympathetic nervous system which in turn raises body temperature. Fever secondary to infection is the result of two distinct physiologic mechanisms: vasoconstriction which prevents heat loss from the body and hypermetabolism which increases heat production.

The burn wound is highly vascularized and surface blood flow is increased; core-skin heat transfer increases with burn size (3), and leg blood flow is largely dependent on the extent of surface injury to the extremity (4). Because mechanisms for vasoconstriction and core insulation are not normal, the burn patient must rely on hypermetabolism as the primary mechanism for increasing body temperature.

The cause of the upward reset in the reference temperature in burn patients is unknown. Blocking afferent nervous input to the brain by topical anesthesia or applying topical anesthesia to the

and printed for and the function of the second s

1. Wilmore DW, Long JA, Skreen R, Mason AD Jr, Pruitt BA Jr: Catecholamines: mediator of the hypermetabolic response to thermal injury. Ann Surg 180:653-668, 1974.

2. Wilmore DW, Orcutt TW, Mason AD Jr, Pruitt BA Jr: Alterations in hypothalamic function following thermal injury. J Trauma 15:697-703, 1975.

3. Wilmore DW, Mason AD Jr, Johnson DW, Pruitt BA Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physiol 38:593-597, 1975.

4. Aulick LH, Wilmore DW, Mason AD Jr, Pruitt BA Jr: Influence of the burn wound on peripheral circulation in thermally injured patients. In press, J Appl Physiol. burn wound has not reduced the hypermetabolic response (5). A-single patient studied with spinal cord transection was markedly hypermetabolic in spite of almost complete denervation of the burn wound. However, the brain appears essential for the post-traumatic metabolic response, and hypermetabolism was observed in a burn patient with brain death.

Post-traumatic hypermetabolism is also attenuated during norphine anesthesia (6). This evidence suggests that afferent nervous input to the brain is not the cause of the hypermetabolic response following injury but that the brain (hypothalamus) is necessary to receive the signal and to integrate the final neurohormonal response. This report evaluates factors which are transmitted in the blood and could initiate the upward shift in central reference temperature, and evaluates new evidence which demonstrates further interaction between the sympathetic nervous system and metabolism of thyroid hormone.

#### Methods

· And the second second second second second second

Arterial and venous blood samples were obtained between the third and 31st postburn day from six burn patients (mean age 30 years, mean burn size 43 per cent), the plasma immediately separated, indomethacin added and the aspirates frozen. Prostaglandin A, E, and F were analyzed, using specific antibody techniques, and the results compared with blood obtained from normal controls. In addition, arterial venous samples were drawn simultaneously from 15 patients (mean age 30 years, mean burn size 45 per cent total body surface), with burned and unburned lower extremities to determine the contribution of the injury to the prostaglandin levels.

Heparinized venous blood was collected from burn patients and normals in pyrogenic-free syringes, the serum separated and transferred to pyrogen-free vials which were heat sealed. The samples were frozen until analysis, when they were thawed, the vial opened using sterile technique, and a micro aliquot injected through indwelling chronic cannula in rabbits, placed by standard sterotatic technique so the

^{5.} Wilmore DW: Hormonal Responses and their Effects on Metabolism. Surgical Clinics of North America, edited by G. A. Clowes, Jr. (In press).

^{6.} Taylor JW, Hander EW, Skreen R, Wilmore DW: Effect of CNS narcosis on the sympathetic response to stress. J Surg Research 20: 313, 1976.

distal tip lies in the preoptic area of the hypothalamus. Rectal temperature was monitored over the next two hours. Serum that yielded a febrile response was heat-treated and re-evaluated, and also tested by limulus lysate assay for endotoxin.

Serial plasma samples were obtained from five male thermally injured patients (mean age 29 years, mean burn size 66.5 per cent TBS) and evaluated by radioimmunoassay techniques for T₄, T₃, reverse T₃ and TSH.

#### RESULTS

Concentrations of prostaglandin A, E, and F were similar to those observed in normals (Table 1), and did not attain levels similar to those reported in pregnant women. All arterial concentrations appeared slightly lower than venous levels, presumably because the lung activity degradates these components. The burn wound did not contribute to the production of PGE or F; blood obtained from patients with and without leg burns showed similar concentrations in the femoral vein. Absolute prostaglandin levels were unrelated to burn size and did not change with time postburn.

Injection of normal serum into the preoptic area of the rabbit resulted in a 0.1°C rise in rectal temperature in 6/6 control subjects. Serum from 9/13 burn patients elicited a febrile response (0.63-0.93°C) following hypothalamic injection. Limulus lysate assay was negative in all these samples, and, after heat treatment of the serum, the response was attenuated, suggesting endogenous pyrogen mediated the thermogenic response.

T₄ levels were below normal levels in the burn patient, but the free thyroid index was normal, suggesting that T₄ concentrations were within low physiologic ranges and the low concentrations resulted from the low serum proteins which are observed following thermal injury. In contrast, triiodothyronine (T₃), the active hormone, was abnormally low throughout the first two weeks of injury (Table 2), and returned to normal with recovery. In contrast, there was an early rise in reverse T₃ which returned to normal levels during the second week postinjury. Further studies during septic episodes in burn patients two to three weeks postinjury demonstrate that the marked rise in reverse T₃ (>125 ng^C) is associated with this complication; T₃ is further suppressed. Despite the depressed T₃ and low T₄ values, basal TSH values are <10  $\mu$  IV/ml, suggesting alterations in the hypothalamus-pituitary-thyroid axis.

			(Mean ± S.E.	-	,	
Subjects	No.	No. of Samples	Sample Site	PGE	PGF	PGA
3urn patients (3-31 days)	Q	25	Arterial	06 <del>-</del> 1066	360 ± 70	1600 ± 90
3urn p <b>a</b> tients (6-21 days)	Q	14	Arterial	950 ± 120	410 ± 120	1590 ± 120
Vorma 1 s	4	4	Venous	1020 ± 290	01 7 COL	1520 ± 80
Pregnant women*	9	ę	Venous	2680 ± 480	490 ± 130	3220 ± 390
²regnanî women*	9	Q	Uterine vein	2530 ± 580	440 ± 130	2670 ± 450
3urn patients	15	15	Arterial	850 ± 134	281 ± 114	
			Femoral vein	974 ± 140	375 ± 234	
Burn patients (no leg burn)	4	4	Arterial	713 ± 160	251 ± 157	8
			<pre>*Femoral vein</pre>	889 ± 230	356 ± 167	
Burn patients (leg burns)	11	11	Arterial	838 ± 177	201 <b>± 101</b>	-
			Femoral vein	1005 ± 177	382 ± 261	

*From the data of Secker, R. A. and Sperbff, L.

Table 1. Prostaglandin Concentrations (PG/ml Plasma)

たいまたないでいた

۳ ۲

257

Postburn Day	T4 (μg%)	FTI	T ₃ (ng%)	RT ₃ (ng%)	TSH (µ IU/ml)
3	2.5	3.8	52	94	4.1
6	2.5	3.0	26	104	2.2
8	3.7	6.1	36	76	4.2
10	5.1	6.2	31	85	3.2
13	3.8	5.0	16	56	3.7
15	5.0	6.3	44	66	4.7
Norma1 Range	4.5-11.5	3.8-13.4	80-180	36-84	10

Table 2. Thyroid Studies in Burn Patients

## DISCUSSION

a for the second state of 
Prostaglandin synthesis in tissues is stimulated by hypoxia, resulting in the elaboration of PGE, a vasodilator which antagonizes the constrictive effect of catecholamines to increase blood flow to the anoxic tissue. PGE (but none of the other prostaglandins) injected into the hypothalamus causes a febrile response (7). It is appealing to speculate that prostaglandins serve as mediators for local dilatation of the wound and also stimulate the alteration in central reference temperature. This hypothesis was not confirmed by by the data. Because prostaglandins are rapidly degradated at the tissue level, these present analytical techniques may not be sensitive enough to detect increased prostaglandin activity and further analysis of PGE metabolites is now being planned. However, these prostaglandins are increased in pregnant women, and these patients do not exhibit alterations in the thermoregulatory mechanisms.

A febrile response was observed when serum from burn patients was injected into the temperature regulation center of the rabbit. This substance is heat labile and thought to be endogenous pyrogen, the biochemical product of lymphocytes and other phagocytic cells. Previous attempts to detect endogenous pyrogen in man during bacterial or viral infection have not yielded such a high percentage of febrile responses, suggesting that the inflammatory responses in the burn wound may continually liberate this mediator.

Whether endogenous pyrogen is in fact the mediator of the metabolic response to thermal injury, or just one of a number of factors which mediate post-traumatic events, is not known. Animal studies are in progress to evaluate the effect of the white cell in eliciting the metabolic and hyperpyrexic response to injury.

Burn patients appear chemically hypothyroid, although metabolic rate is elevated because of increasing discharge of the sympathetic nervous system. A fall in T₃ has been associated with increased gluconeogenesis in man, which apparently alters T₄ conversion to T₃ in the liver. Reversed T₃, an iodinated compound with low biological activity, appears to be generated, although the exact site of synthesis of this substance is not known. The effect of this hypothyroidism on cell and organ function in injured man is now being evaluated in an animal mode'.

7. Stitt JT, Hardy JD, Stolwijk JAJ: PGE, fever: its effect on thermoregulation at different low ambient temperatures. Amer J Physiol 227:622, 1974.

## REFERENCES

1. Nilmore DN, Long JA, Skreen R, Mason AD Jr, Pruitt BA Jr: Catecholamines: mediator of the hypermetabolic response to thermal injury. Ann Surg 180:653-668, 1974.

2. Wilmore DW, Orcutt TW, Mason AD Jr, Pruitt BA Jr: Alterations in hypothalamic function following thermal injury. J Trauma 15:697-703, 1975.

3. Wilmore DW, Mason AD Jr, Johnson DW, Pruitt BA Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physiol 38:593-597, 1975.

4. Aulick LH, Wilmore DW, Mason AD Jr, Pruitt BA Jr: Influence of the burn wound on peripheral circulation in thermally injured patients. In press, J Appl Physiol.

5. Wilmore DW: Hormonal Responses and their Effects on Metabolism. Surgical Clinics of N America, edited by G. A. Clowes, Jr. (In press).

6. Taylor JW, Hander EW, Skreen R, Wilmore DW: Effect of CNS narcosis on the sympathetic response to stress. J Surg Research 20: 313, 1976.

7. Stitt JT, Hardy JD, Stolwijk JAJ: PGE, fever: its effect on thermoregulation at different low ambient temperatures. Amer J Physici 227:622, 1974.

#### Acknowledgements

The prostaglandin assay was performed by Doctor Richard Becker, Department of Endocrinology, University of Texas Health Science Center at San Antonio, and Doctor Leon Speroff, Department of Obstetrics and Gynecology, Yale University School of Medicine. The microinjection studies were performed by Doctor Keith Cooper, Head, Division of Medical Physiology, University of Calgary, Calgary, Canada; r-T₃ assay was performed by Doctor Kenneth Berman at the Walter Reed Army Hospital.

# PRESE'ITATIONS

Becker RA: Depressed serum trilodothyronine (T₃) levels following thermal injury. 60th Annual Meeting of the Federation of America Society for Experimental Biology, Anaheim, California, 10 April 1976.

# PUBLICATIONS

•

Becker RA, Johnson DW, Woeber KA, Wilmore DW: Depressed serum tritodothyronine  $(T_3)$  levels following thermal injury. Fed Proc 35: 216, 1976.

RESE 4 AND TECHNOLOGY WORK UNIT SUMMARY										ONTROL STREOL
KE3P 4	AND TECHNOLOG	WUKK UNIT S	UMMARY	DA	OG 695	51	76 10	01	DD-DI	AE(AR)636
1. DATE PREV SUMRY	4 KIND OF SUMLARY	S. SUMMARY SCTY	. WORK SECURITY	-	ADING	-	SO'N INSTR'N	IN SPECIFIC	DATA-	LEVEL OF SUM
76 10 01	D. CHANGE	<u> </u>	U	NA		NL		VES I	] но	A. WORK UNIT
19. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK	AREA NUM	BER		WORK UNI		
& PRIMARY	61101A	3A161101	A91C	ļ	00		079			
D. CONTRIBUTING	ļ	ļ	·····	<b> </b>			·	. •	•	
C. CONTRIBUTING				1		ليب				
Thermal Ini	117V Hypermeta	(U) Deve.	Lopment and	i Ana	LYSIS	OI i	an Anima	1 Model	tor (	the Post
12. SCIENTIFIC AND TE	CHNOLOGICAL AREAS	borre kesp		111 C.	le bui	neu	Soluter	(44)		
003500 Clin	ical Medicine									
13 START DATE		14. ESTIMATED COM	LETION DATE	IS PUN	DING AGEN	CY		IS. PERPORM	ANCE MET	100
76 01		Cont	t	DA	1		1	с. 1	n-Hous	se
17 CONTRACT/GRANT	Not Applicab	le		18 RES	DURCES ES	TIMATE	A PROFESSI	ONAL MAN YR	L FUN	DE (In these ande)
A DATES/EFFECTIVE		EXPIRATION			76	,	6			
N HUNDER *				FISCAL	<u>7</u> <u>7</u>		.3		3	
G TYPE.		4 AMOUNT		YEAR	CURRENT 77		1.			
& KIND OF AWARD		f. CUM. AMT.			11		1.0		16	
TR. RESPONSIBLE DOD C				44. PER  	URTING OI	-GANIZ	A TION	L		
HANE US Army	-	US Ar	my :	Institut	e of Su	rgical	Research			
			Burn	Stu	udy Bran	ch				
Fort	94		rort	sam	Houston	, Texas	/823	14		
		PRINCIP	AL INVEST	GATOR	(Fumioh SEAN I	IUS Academic	[nelikulian)			
RESPONSIBLE INDIVIDU		-	Davi	d N	Herndo	n, CPT	MC			
NAME Basil A. Pruitt, Jr, COL, MC					HONE 51	2-22	21-4440	.,,		
TELEPHONE: 51		SOCIAL	SECURITY	ACCO	UNT NUNBER					
21 GENERAL USE ASSOCIATE INVESTIGATORS										
FOREIGN INTELLIGENCE NOT CONSIDERED NAME:										
		Allin Andel		HAME						DA
31. KEYWORG (Priced EACH with Secondly Clearly constitution Code) (U) Oxygen consumption; (U) Hypermetabolism;										
(U) Thermal injury; (U) Thyroid; (U) Catecholamines										
23. (U) To develop a reliable and easily reproducible small animal model of the								e		
postburn hypermetabolic response of humans; to better delineate the endocrine medicato								edicators		
of this resp	oonse; to inve	estigate th	e afferent	mech	anism	s ti	hat init	iate th:	is res	ponse;
to investiga	te the respon	ise of isol	ated tissu	es to	dete	rmir	he which	organs	are h	yper-
metabolic ar	ha which are i	ormal or h	ypometabol	ic po	stpur	n; t	otabol:	r aeline	eate t	ne
24 /II) And	manisms of the	lermoyenesi	s in the p	o die	th ny	hetu toy	ac foll	. Stdte. γwα•(∍)	enam	anec-
thetized it	(b) hurned or	60% THS L	uth standa	e uis rd se	ald	in 4	ach of	the fol	lowing	treatmen
groups: (1)	thyroidectomi	Led one mo	nth prior	to st	udv.	(2)	adrenal	ectomize	ed one	month
prior to stu	dv. (3) irrad	lrated 2 wk	s prior to	stud	y to	prod	luce leul	copenia	(4)	treated
with B.I.D.	injections of	1 mam/kam	indometha	cin s	tarti	ng 2	days n	rior to	start	of study
drug to cont	inue through	study peri	od. (5) tr	eated	with	5 d	lifferen	t topica	al ant	i-
microbials.	(6) treated w	vith varyin	g diet, (7	) 1nf	ected	wit	h topica	al pseud	lomona	s.
These groups	will be prep	ared in ra	ts mature 1	but c	rowin	g (]	.80 gm ) 1	rats at	their	plateau
growth phase	e (approx. 540	) gm) guine	a pigs mat	ure b	ut gr	owir	ng (appro	<b>54</b> 0	gmn)a	nd
guinea pigs	at plateau gi	owth phase	(i.e. 100	Ogm).	Who	le b	ody oxy	jen cons	sumpti	on will
be determine	ed in modified	l Warburg a	pparati on	eact	day	from	n seven d	lays pri	lor to	start
of study to	the time of h	ourned anim	als comple	te wo	und h	ealı	ng. In	vitro t	issue	oxygen
lactate gluc	ose and glyco	ogen consum	ption or p	roduc	tion	of 1	liver sli	ices, di	aphra	ст
sections and	white fat ce	ell suspens	ions will	be ma	de fr	om a	ill treat	ment gi	coups	atvarying
periods in t	he postburn h	ypermetabo	lic respon	se ar	d com	pare	ed with r	natched	contr	o <b>l.</b>
25. (U) 76	01 - 76 09 A	reproducib	le rat and	guir	ea pi	g mc	del demo	onstrati	ing 40	to 31%
increases in	i metabolic ra	ite post 60	and 40% b	urns	respe	ctiv	vely, rel	lative t	o mat	ched
controls. A	11 studies at	thermal n	eutral (32	•() }	ave b	een	establig	sned p≮.	.0001,	ror
Aveilable to contractor	LUGIEA. e upon originator's approv	•i	ند ب <u>ن زار در من این می</u> در در م							
DD. FORM 1498	B PREVIOUS E	DITIONS OF THE	S FORM ARE OB	SOLET	E DUF	RMS	1498A 1 NO	¥ 55		
			2	6.2						

.

## ANNUAL PROGRESS REPORT

### PROJECT NO: 3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH

## **REPORT TITLE:** DEVELOPMENT AND ANALYSIS OF AN ANIMAL MODEL FOR THE POST THERMAL INJURY HYPERMETABOLIC RESPONSE FOUND IN BURNED SOLDIERS

## US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MED CAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

David N. Herndon, MD, Cpt, MC Douglas W. Wilmore, MD Arthur D. Mason, Jr., MD Ronald V. Condon, Jr., SP4 Edwin W. Hander Basil A. Pruitt, Jr, MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

### PROJECT NO. 3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH

## REPORT TITLE: DEVELOPMENT AND ANALYSIS OF AN ANIMAL MODEL FOR THE POST THERMAL INJURY HYPERMETABOLIC RESPONSE FOUND IN BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1976-30 September 1976

Investigators: David N. Herndon, MD, Cpt, MC Douglas W. Wilmore, MD Arthur D. Mason, Jr., MD Ronald V. Concon, Jr., SP4 Edwin W. Hander Basil A. Pruitt, Jr, MD, Colonel, MC

Reports Control Symbol MEDDH-288 (R1)

A reproducible small animal model simulating the post thermal injury hypermetabolic response of humans was developed. The oxygen consumption of one hundred 60% burned rats was approximately 1.33 times that of a group of 100 weight-paired controls from the 5th through the 45th post burn day when the animals were studied at 27°C and 32°C (p < 0.0001 at both temperative tures). The metabolic rates of normal and burned rats were increased at 27°C relative to their metabolic rates at 32°C. These increases were proportional in each group. A group of 20 guinea pigs with 40% burns, 70 guinea pigs with 60% burns and 110 weight-paired controls gave similar results. The ratios of their observed to predicted normal metabolic rates were 1.33, 1.39, and 1.00 respectively at  $32^{\circ}C$  (p<0.0001). The metabolic rates of turned mature rates were also elevated over those of weight-paired controls. The oxygen consumption of a group of forty 60% burned, 540 gm rats was 1.35 times that of weightpaired controls. The mortality of a 60% burn in these mature rats was 90% by post burn day 30, whereas it was only 15% in 180 gm rats burned during their linear growth chase (180-500 gm). When compared with 60 intact normal rats from the 7th to the 25th post burn day, the ratio of observed/predicted oxygen consumption of thyroidectomized rats was 0.61. that of 20 thyroidectomized burned rats was 1.00, and that of 60 normal burned rats was increased to 1.41 (p<0.01).

Oxygen consumption Hypermetabolism Thermal injury Thyroiu Catacholomines

## DEVELOPMENT AND ANALYSIS OF AN ANIMAL MODEL FOR THE POST THERMAL INJURY HYPERMETABOLIC RESPONSE FOUND IN BURNED SOLDIERS

Hypermetabolism is a salient feature of the human response to thermal injury. Recent studies by Wilmore, et al have shown that the resting metabolic rate increases with burn size in a curvilinear fashion (approaching 70 to 75 Kca//M⁻/hr as burn size exceeds 50% of total body surface). This is approximetabolism; peritonitis can cause a 25% increase in energy consumption, a multiple traum: patient on a respirator can have up to a 75% increase, but even this is only equivalent to the rate generated by a 30% burn. The catabolic demands of this response are huge. Prior to the rate of continuous hyperalimentation at the Institute of Surgical Research, patients with greater than 40% total body surface burns lost approximately 25% of their preburn weight in the first 6 weeks post injury. Recent work by Wilmore, et al has shown that the ability of a thermally injured patient to maintain hypermetabolism in the face of a cold stress can be exhausted and that this exhaustion is correlated with mortality.

The stimulus, mediators, and mechanisms of end organ response that produce the hypermetabolic reaction to thermal injury are not fully understood. Wilmore, et al. demonstrated a correlation between metabolic rate and urinary catecholamine excretion. They were further able to decrease the hypermetabolism of burned patients via a combination of alpha and beta adrenergic blockade or beta adrenergic blockade alone indicating that catecholamine is one of the more significant efferent mediators of this response. Glucagon, which is also increased in the pos' burned state, has been shown to be capable of increasing metabolic rate of normal as well as burned individuals. Thyroid hormone, mediator of hypermetabolism in other conditions, has been shown not to be a primary mediator. "" Burn victims have, in fact, in some situations been shown to have

1. Wilmore DW, Long JM, Mason AD, Jr, Skreen FW, 2010 It BA, Jr: Catecholamines: Mediator of the hypermetabolic respondence to permakinjury. Ann Surg 180: 653 (Oct) 1974.

2. Wilmore DW, Mason AD, Jr, Johnson DW, Pruitt BA, Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appi Physiol 38: 593 (Apr) 1975.

3. Ibid 2.

4. Aulick LH, Wilmore DW, Mason AD, Jr: Mechanism of glucagon calorigenesis Fed Proc 35: 401 (Abst # 1077) (Mar) 1976.

5. Cope O, Nardi GL, Quijano M, Rovit RL, Stanbury JB, Wight A: Metabolic rate and thyroid function following acute thermal injury in man.Ann Surg 137:165, 1953.

6. Caldwell FTR, Osterholm JL, Sower ND, Moyer CA: Metabolic response to thermal trauma of normal and thyroprivic rats in 3 environmental temperatures. Jun Surg 150: 976 (Dec) 1956. diminished thyroid hormone levels.⁷ The precise character of the mediating hormonal milieu has not been delineated. The afferent stimuli that induce this milieu are unknown. The character of end organ and cellular response are unknown. Further definition of these mechanisms might lead to an ability to better support and modulate the process.

Various groups have developed animal models to investigate the hypermetabolic response to thermal injury. Morgan, et al ^o found insensible water loss to be significantly elevated in burned rats. Lieberman and Lansche measuring metabolic rate and insensible water loss of thermally injured rats found both to be elevated but "the rate of increase in oxygen consumption was less than the increase of caloric loss incident to the increased rate of insensibly lost water in these rats". From this and the evidence of one burned rat whose metabolic rate was returned to normal by placing a water impermeable covering over the burn they concluded that the hypermetabolic reaction to thermal injury was attributable to heat loss from the evaporation of water. Caldwell in a series of studies using a Benedict calorimeter came to similar conclusions.^{6,10} He showed a significant increase in metabolic rate in 20% burned rats relative to unburned controls studied at 24°C from the 2nd through the 65th post injury day. However, when these same animals were studied at 32°C, in the thermoneutral range for rats, there was no difference in the metabolic rate of the 2 groups. His interpretation, like that of Lieberman and Lansche, was that post burn hypermetabolism in rats is providing energy 2 needed to offset evaporative heat losses. Human studies by Wilmore, et al and by Barr, et al' clearly demonstrate that the human hypermetabolic response post thermal injury is not abolished by raising ambient temperature to thermoneutral, 32°C. Further, Zawacki, et al showed that covering the burn wound with a water impermeable membrane did not decrease caloric utilization. In man evapoarative water loss is not the primary stimulus of the hypermetabolic response but is elevated in parallel with it and may be the method by which the increased heat generated in response to other signals

7. Becker RA, Johnson DW, Woeber KA, Wilmore DW: Depressed serum triiodothyronine (T₂) levels following thermal injury. Fed Proc 35: 216, 1976.

8. Morgan HL, Andrews RP, Jurkiewicz MJ: The effect of thermal injury on insensible weight loss in the rat. Surg. Forum 6: 78, 1955.

9. Lieberman ZH, Lansche JM: Effects of thermal injury on metabolic rate and insensible water loss in the rat. Surg Forum 7: 83, 1956.

10. Caldwell, FT, Hammel H, Dolan F: A calorimeter for simultaneous determination of heat production and heat loss in the rat.J Appl Physiol 21:1665, 1966.

11. Barr PO, Birke G, Liljedhal SO, Plantin O: Oxygen consumption and water loss during treatment of burns with warm dry air, Nature 1: 104, 1963.

12. Zawacki BE, Spitzer KW, Mason AD, Jr, Johns LA: Does increased evaporative water loss cause hypermetabolism in burned patients? Ann Surg 171: 236, 1970. is dissipated. In recent animal studies ^{13,14} Farkas, et al showed small increases in oxygen consumption in 59 rats with 27% burns over those of paired controls. This increase was significant in animals studied at 32°C in an atmosphere with a humidity of greater than 60%.Oxygen consumption was less in both burned and control animals at 32°C than at 24°C. ¹ Aikawa, et al studying the hypermetabolic response of guinea pigs with 11% total body surface burn at 24.8°C reported increases in post burn metabolic rates which were correlated with increases in uninary catecholamine excretion. In this study elevation in total energy production could not be explained by comensurate elevations in measured evaporative heat loss. They concluded that increases in evaporative heat loss post thermal injury were secondary and "could at most account for a small portion of the post burn hypermetabolism".¹⁴

The objective of this investigation was to develop a reliable and easily reproducible animal model c the post burn hypermetabolic response of humans that would allow investigations of the stimulus, mediators, and mechanisms of end organ response to that reaction.

## MATERIALS AND METHODS

and the second secon

Various groups of animals were burned over 20, 40, and 60% of their total body surface by immersion in boiling water as previously reported by Walker and Mason. The animals were anesthetized with intraperitoneal pentobarbital. (with the exception of guinea pigs which were anesthetized with inhalation penthrane). The animals were clipped of fur prior to burning. Weight-paired control animals were also anesthetized and clipped of fur, they were then dipped in tepid water. All animals, except as noted below, were resuscitated with 1 cc of Ringer's lactate/kg/% burn injected intraperitoneally at the time of burning and with another equal injection 8 hours post burn. Whole body oxygen consumption was determined over 1 to 2 hour periods prior to burn (or sham manipulation) and at intervals of 1 to 4 days thereafter to approximately 45 days post burn. **Control animals were reclipped of fur twice each week.** Oxygen consumption was studied by placing animals in air-tight desiccator jars over a layer of sodalime, (a CO, absorbent). The jar was connected in series to an Erlenmever flask equilibrated with 100% oxygen. The flask was connected in series to a water bath in which all apparatus was submerged. The bath temperature and the ambient temperature in the metabolic chambers were kept constant at the predetermined level + 1.5°C. As the animals breathe, CO, is trapped by the sodalime, and a vacuum is created which, after overcoming the surface tension of the system, pulls oxygen from the flask to the jar and water from the bath to

^{13.} Farkas LG, McCain WG, Birch JA, James J: The effects of four different chamber climates on O consumption and healing of severely burned rats. J Trauma 13(10):911,1973.

^{14.} Aikawa N, Caulfield JB, Thomas RJS, Burke JF Postburn hypermetabolism: Relation to evaporative heat loss and catecholamine level. Surg Forum 26: 74, 1975.

^{15.} Walker HL, Mason AD, Jr: A standard animal burn. J Trauma 8: 1049, 1968.

the flask in a volume proportional to the amount of oxygen consumed. Sixteen jars are run simultaneously, 8 experimental animals and 8 weight-paired controls. The length of the experiment in minutes, the volume of water pulled into the Erlenmeyer flask in ccs, the weight of the animal in grams, the ambient temperature in degrees centigrade in the metabolic chambers and the barometric pressure at the time of the experiment were recorded. From these measurements, oxygen consumption/unit weight/unit time could be calculated after correcting for standard temperature and pressure. All animals were studied on different days at ambient jar temperatures of 27°C and at ambient temperatures of 32°C, thermoneutral. Oxygen consumption/unit time of experimental animals were compared with oxygen consumption of weight-paired controls over periods of time post injury by the Students unpaired "t" test, and the Scheffe modification of the "t" test when required.

#### RESULTS

and the constraint of the state of the set of the state o

The Effect of a 60% Thermal Injury on the Growth Rate of Rats Burned at 180 gms of Weight: The growth rate of 50 rats given a 60% burn when they weighed 180 gms was compared with that of 50 sham burned rats weighing 180 gms. As seen in Figure 1, the regression line defining post burn day versus weight for burned animals from the date of burn to the 80th postburn day is higher than and has a steeper slope than the regression line defining the control animals. The line defining the rate of growth of the burned animals from the 7th to 80th post burn days (Figure 2) is, however, seen to be parallel to but significantly elevated over that of the controls. This indicates that when rats are burned at 180 gms, a point early on their linear growth phase, they undergo a growth arrest of approximately 7 days after which they regain at a normal rate. At any given time post burn, however, the burned rats are lighter than controls weightpaired at the time of burning. Therefore, any comparison affected by weight of the animals must be corrected for this discrepancy.

Comparison of the Metabolic Rate of 60% Thermally Injured Growing Rats with that of Predicted Weight-Paired Controls from the 1st to 60th Post Burn Day with all Animals being Studied at an Ambient Temperature of 27°C. The amount of oxygen consumed/hour is plotted against weight for 40 control rats sham burned at 180 gm weight and studied every 3 days post sham at an ambient temperature of 27°C (Figure 3). The regression line defined by the equation Y = 160.70 + 0.7709X is a fair predictor of this relationship,  $(r^2 \text{ value} = 0.07)$ . If one divides the observed oxygen consumption/hour of control rats of a given weight by the oxygen consumption/hour that would be predicted for given weights by the above regression line and compares these ratios to post burn day, a relationship with slope confidence limits of -0.0019to 0.0005 and a Y intercept of 1 is obtained (Figure 4). This confirms the validity of the above regression line as a predictor of normal oxygen consumption/unit weight. In Figure 5, the oxygen consumed/hour by a group of 40. 60% burned rats studied at an ambient temporature of 27°C, divided by the oxygen consumption that would be predicted for a normal rat of the same weight, und r the same conditions, is plotted against post burn day. A curve is defined that is very similar to that obtained for thermally injured patients. Metabolic rate increases from a preburn level which is equivalent to the controls to an



OONH BOKZ OAY

FIGURE 1





LD ZYCB HNOD



N. 34. 18

territerin an Albert

in the second

271



OBSERVED VO2/ PREDICTED VO2

sty is a star share to the the starts

のないであるというない

Ş

る、このないないないないから



22.4.1

elevated plateau which is relatively constant from the 5th to the 45th post burn day, at which time the metabolic rate gradually decreases to control levels by the 60th post burn day. If the ratios of observed burn over predicted normal oxygen consumption are plotted against post burn day from the 5th to the 45th post burn day (Figure 6), they are found to define a horizontal line with a Y intercept of 1.36. If these values are compared with the observed/predicted ratios of the controls, this post burn hypermetabolic response is found to be highly significant ("t" value 18.7 with 202 degrees of freedom p40.0001 – Figure 7). At 27°C growing rats have an increase in metabolic rate from the 5th through the 45th post burn day of approximately 34% above controls of the same weight.

Comparison of the Metabolic Rate of 60% Thermally Injured Growing Rats with that of Predicted Weight-Paired Controls Studied at an Ambient Temperature of 32°C. The amount of oxygen consumed/hour is plotted against weight for 40 control rats sham burned at 180 gm weights and studied every 3 days post sham at a 32°C ambient temperature, thermoneutral, (Figure 8)., A regression line is defined by the equation Y = 74.6441 + 0.0072X with an r⁴ value of 0.86. Dividing the observed oxygen consumption/hour of control rats of given weights by the amount of oxygen that would be predicted from the above equation to be consumed by a rat of that weight, one obtains a line with slope confidence limits of -0.00117 and + 0.0011 with a Y intercept of 1 (Figure 9). The ratio of oxygen consumed/hour by a group of 40,60% thermally injured rats studied at the ambient temperature 32°C, divided by the predicted amount of oxygen that would be consumed by control rats of similar weight under similar conditions, again defines a curve very similar to that of thermally injured humans (Figure 10). The metabolic rates rise over the first 5 days post burn to an elevated plateau 32% higher than that of controls. The elevation in metabolic rate is relatively constant from the 5th to the 45th post burn day after which it gradually decreases to normal by the 60th post burn day. Comparison of the metabolic rates of the burned rats studied at 32°C with normals under similar conditions (Figure 11), shows the 32% increase in the burned animals' metabolic rate over controls to be highly statistically significant (unpaired "t" value 15.5 with 189 degrees of freedom p<0.0001). This indicates that growing rats, like humans, are hypermetabolic post thermal injury when studied at an ambient temperature of 32°C. Increasing temperature does decrease the amount of oxygen consumed by both normal and burned animals of any given weight as can be seen by the compatison of oxygen consumption/hour against weight for control animals at 32°C and at 27°C (Figure 12). The regression line defining the animals studied at 32°C is significantly lower than that defining those studied at 27°C (p<0.01). The comparison of the burned rats studied at the 2 temperatures (Figure 13) also shows that the oxygen consumption is lower at the warmer temperatures; but the decrease in metabolic rate of burned animals at the higher ambient temperature is comparable to the reduction observed in the control rats.

<u>The Effect of Animal Age on Metabolic Rate.</u> Mature rats, that had reached the plateau of their growth phase (540 gm), were submitted to 60% thormal injuries and their metabolic rates were found to be significantly elevated over weight-paired controls from the 5th through the 26th post burn day,  $p \ge 0.001$ (Figure 14). The increase in metabolic rate of these mature 60% burned rats was



to some weather .

275







FIGURE 8


FIGURE 9

NUM OF STREET

いたれた

}



AND AND AND AND AND AND AND

Sa Shi ka ka maraka



OBSERVED/PREDICTED 02 CONSUMPTION





· · --

FIGURE 13



h .. Y

1.41 times that of their weight-paired controls. The mortality of these mature animals following a 60% thermal injury was markedly greater than that of growing rats after a similar injury. 85% of the large rats succumbed within the first 28 post burn days. Only 10 to 15% of the growing rats succumbed to a similar injury. The cause of death in the adult rats, confirmed by autopsy appearance of gram negative wound invasion and miliary hematogenous bacterial deposits, as well as several pre-death blood cultures, was gram negative wound invasion and sepsis. The frequency of death in these mature rats after a 60% burn more closely approximates the human response to thermal injury than does that of the growing rat. It is presumed that there is something protective in the anabolic milieu of the growing rat to thermal injury. The adult rat may be a better model of the human response.

Comparison of the Metabolic Rates of 40% and 60% Thermally Injured Guinea Pigs with the Predicted Metabolic Rates of Weight-Paired Controls Studied at 32°C Ambient Temperature, Thermoneutral. A group of 20 guinea pigs was given a 40% thermal injury at 280 gm body weight (an early point on the linear growth phase). A second group of 70 guinea pigs was given a 60% burn at 540 gm body weight (a point midway on the linear growth phase for Both groups were compared with weight-paired controls and quinea pigs) with each other. Figure 15 shows that the oxygen consumed/hour/unit weight. of control guinea pigs studied in an ambient temperature of 32°C can be described by the equation Y = 145.7246 + 0.5817X with an r² value of 0.66. When observed oxygen consumption/hour for animals of given weight is divided by the oxygen consumption/hour that would be predicted for that weight from the above equation and the results plotted against post sham days, a horizontal line with a Y intercept of 1 is defined as would be predicted (Figure 16). When the observed oxygen consumption of 40% burned guinea pigs is divided by that predicted for that weight and plotted against post burn day, a curve similar to that of rats and humans is defined, increasing over the first 5 days post burn to a plateau that is constant from the 5th to the 45th post burn day, gradually returning to normal by the 60th post burn day (Figure 17). If the rates of oxygen consumption for the burned animals from the 5th to the 45th post burn days are compared with the predicted normals of the animals of the same weight, the observed increase is highly significant with a p < 0.0001, the burns being elevated 1.31 times the controls. The 20, 60% burned guinea pigs, by the same method of analysis, were found to have an elevation of 1.39 times that of controls, again highly significant with p < 0.0001 (Figure 18).

The Effect of Resuscitation on Metabolic Rate Post Thermal Injury. Groups of 10 controls and 10,60% burned growing rats were respectively resuscitated with 1 cc/kg/% burn of Ringer's lactate at the time of burn and 2 cc/kg/ % burn Ringer's lactate. No differences in the degree of hypermetabolicm post thermal injury could be ascertained among these four groups.

The Effect of Thyroidectomy on Metabolic Rate. Ten thyroidectomized control rats, 10 thyroidectomized 60% burned rats, 10 non-thyroidectomized control rats, and 10 non-thyroidectomized 60% burned rats we e compared. All rats were 540 gm at the time of burn or sham. Metabolic rates were compared from the 5th to the 25th post burn days at which times the metabolic rates of all



<u> 285</u>



いたいないないないないないないないないないないないです。

OBSERVED/PREDICTED 02 CONSUMPTION

. ...



and the second state of the second state of the second state of the

24. L. 24





September 1

\$12. The

----

OBSERVED/PREDICTED O2 CONSUMPTION

groups were found to be constant. Their ratios were: thyroidectomized unburned rats,  $0.61, \leq$  thyroidectomized burned rats 1.01 = to non-thyroidectomized controls  $1.00, \leq$  non-thyroidectomized burns 1.41. These results were significant with a p value of  $\leq 0.05$  by the Scheffe test. This indicates that thyroid hormone levels are not a primary mediator of the post burn hypermetabolic response but do condition it.

## DISCUSSION

and the second state of th

Previous small animal models that have been developed to simulate the human post burn hypermetabolic response have shown significant increases in oxygen consumptions with various burn sizes, when the animals were studied at relatively cold environments (24°C). consistent increases in metabolic rate when the animals were studied at thermoneutral 32°C, ³ and this increase was quite small. The human post burn hypermetabolic response is clearly not temperature dependent.^{1,2} The increases in metabolic rate in humans can not be explained by increased energy requirements generated by evaporative heat loss. as has been the explanation ments generated by evaporative heat loss, for the increased metabolic rates in many of the above animal studies. The present study establishes an animal model using rats and guinea pigs of varying stages of maturity, all of which showed large (on the order of 130-140of controls) and significant (p<0.0001 for all groups) post burn hypermetabolic reactions at thermoneutral (32°C). The metabolic responses of these animals was temperature sensitive as both normal and burned animals showed lower metabolic rates at 32°C than at relatively cold 27°C. But the decreases in metabolic rate were proportional for the burned and unburned animals. The characteristics of the metabolic rate versus post burn day of the animals, studied in this project closely simulate those previously reported for humans, increasing to a plateau elevation from the 5th to the 45th post burn day and then gradually returning to normal as the burn wound heals.

The reason this study shows increases in metabolic rates of burned animals is probably burn size related. The largest injury inflicted in earlier studies was the 27% burn of Farkas, ¹³ our burns were 40 and 60%. All previously studied animals were on their linear growth phase, burns being applied to rats when their weight was 180 to 220 gm. A 20-30% burn is fatal to only 5-15% of animals in this anabolic milieu and probably does not represent a sufficient stress to institute a hypermetabolic response.

## PRESENTATIONS and/or PUBLICATIONS None

ОАТЕ РЯСУ SUM'AY 75 07 01 D NO /CODES * PRIMARY CONTRIBUTING CONTRIBUTING CONTRIBUTING U) EVALUAT (U) EVALUAT (U) EVALUAT CONTRICTORANT CONTRACT/ORANT CONTRACT/ORANT CONTRACT/ORANT CONTRACT/ORANT CONTRACT/ORANT	H. TERM H. TERM PROGRAM ELEMENT 61102A Soundy Classification Code ion of Calcin NHOLOGICAL AREAF ical Medicing	b SUMMARY SCTYP U PROJECT 3A161102H J um Metabol:	e work security U NUMBER 371P	7 AEGRA N TASK A O	A NREA NUM	DA DISO NI DER	и INSTR'N 066	A SPECIFIC D. ON TRACTOR A S YES	ATA- CCESS MQ NUMBER	LEVEL OF SUN
75 07 01 75 07 01 75 07 01 75 07 01 75 07 01 76 07 01 76 07 01 76 01 77	H. TERM PROGRAM ELEMENT 61102A Soundy Clease/Heating Code ion of Calcin NHOLOGICAL AREAF ical Medicing	U PROJECT 3A161102F um Metabol:	U HUMBER 371P	N TASK A O	A Rea NUM 2		066	ONTRACTOR A	NO NUMBER	A FORK UNIT
D NO /CODES * PRIMARY CONTRIBUTING CONTRIBUTING CONTRIBUTING TITLE (Precede with 1 (U) EVALUAT CONTRIBUTING AND TEC 003500 Clin START DATE 74 01 CONTRACT/ORANT DATES/EFFECTIVE: NUMBER *	PROGRAM ELEMENT 61102A soundy classifiestion Code ion of Calcin nHOLOGICAL AREAS [®] ical Medicine	PROJECT 3A161102H um Metabol:	NUMBER 371P	TASK A	2	DER	066	NORK UNIT	NUMBER	
PRIMARY CONTRIBUTING CONTRIBUTING TITLE (Proced with 1 (U) EValuat SCIENTIFIC AND TEC 003500 Clin START DATE 74 01 CONTRACT/ORANT ADATES/EFFECTIVE: NUMBER *	61102A	3A161102F	371P	0	2		066			
CONTRIBUTING CONTRIBUTING TITLE (Proced with a (U) EValuat scientific and tec 003500 Clin stant Date 74 Ol CONTRACT/ORANT DATES/EFFECTIVE: NUMBER *	esenty cleaselfleasilen Code ion of Calcin unological Ament [®] ical Medicine	,• um Metabol:	ism in Burn		· <u>····</u> ·····			*****		
CONTRIBUTING 1 TITLE (Proced with 1 (U) EValuat 2 SCIENTIFIC AND TEC 003500 Clin 3 START DATE 74 01 CONTRACT/ORANT NOATES/EFFECTIVE: NUMBER *	Saunty Clease Heating Code ION OF Calcin HNOLOGICAL AREAS [®] ICal Medicing	um Metabol:	ism in Burn	1						
U) Evaluat (U) Evaluat Scientific and tec 003500 Clin Stant Date 74 01 Contract/orant A Dates/effective:	amily Closeffeetion Code ion of Calcin MNOLOGICAL AREAS [®] ical Medicine	of Metabol:	ism in Burn							
2 SCIENTIFIC AND TEC 003500 Clin 3 START DATE 74 01 7 CONTRACT/ORANT A DATES/EFFECTIVE: humber * 1 TYPE	HNOLOGICAL AREAS [®] ical Medicine			ed Tr	oops	(44)				
3 START DATE 74 01 7 CONTRACT/ORANT N DATES/EFFECTIVE: N NUMBER *	·····	e		<u></u>						
74 01 CONTRACT/ORANT DATES/EFFECTIVE:		14. ESTIMATED COM			DING AGEN	CY	T	IS PERPORAA	NCE METH	100
7 CONTRACT/GRANT N DATES/EFFECTIVE: N NUMBER ⁴ 1 TYPE		76 09		DA	1		1	C. In-	House	l -
N DATES/EFFECTIVE: N NUMBER ⁴ 1 Type	Not Appl	icable		18 RES		TIMATE	-		L PUN	36 (jn Shoucasdo)
NUNDER *		EXPIRATION:			76		.3		1.0	
LITYPE				FISCAL	<u>7</u> 1		<u> </u>		2	
		A MOUNT:		YEAR	CONNERT					
KIND OF AWARD		I. CUM. AMT					L			
RESPONSIBLE DOD O	RGANIZATION	L				N#4812A				1
AME." US Army	Institute of	Surgical	Research	HANE:*	USA	rmy	Institut	e of Su	rgica	i Researc
	• · · · • • · · · · · · · · · · · · · ·		24	-		ecab	Nonepe-	инсп Точес	. 707	27
Fort S	am Houston,	Texas /82	34		- rort	. san	noustor	i, rexas	102	. J 72
				PRINCIP	AL INVEST	SATOR	(Purstan 38 API II	U.S. Atodomic	(ne i frestan)	,
	AL			HAME	• Rich	ard	H. Merri	11, LTC	, MC	
Basil /	A. Pruitt, Jr	, COL, MC		TELE		12-2	21-5416			
TELEPHONE. 512	2-221-2720			10CIA	L BECURIT	* ACCOU	NT NUMBER			
I GENERAL USE				-			•			
FOREIGN IN	TELLIGENCE NO	T CONSIDER	ED	-	Will	iam	D. Myers	s, LTC,	MC	
				HAME	Harı	ty R.	Jacobs	on, MAJ,	MC	DA
T KEYBOADS (Procede	RACH all south Classifi n motaboliem.	(11) Rurne	: ([]) Renal	fai	lure:	(U)	Hypoca	cemia:	(U) HI	umans
(U) Calciu	IVE. 24 APPROACH. 34	PROGRESS (Fumilia	Individual paragraphs in	tentified to	number P	Nesade Ini	il of each with St	ourity Classifts	atten Cada	,
23. (II) T	ne objective	of this st	udy is to d	leter	mine t	:he π	echanis	us of hy	poca:	lcemia
and hypoca	lciuria in th	ermally in	jured patie	ents.	In a	addit	ion the	role of	Ecald	cium in
the produc	tion of renal	impairmen	t in therma	ally	injure	ed tr	oops is	to be a	asses	sed.
		-		-						
24. (U) C	alcium balanc	ce studies	will be per	form	ed on	burn	patien	ts, with	n all	calcium
intake and	output measu	ured. If,	as we exped	ct, t	he pat	tient	is in j	positive	e cal	cim
balance, w	e will invest	tigate the	skin as the	e sit	eof	calci	um depo	sition.	Skì	n
biopsies a	nd eschar bio	opsies will	be perform	ned a	nd and	alyze	d for c	alcium.	In	addition,
parathyroi	d hormone lev	vels will b	e determin	ed, a	nd th	e res	ponse t	c the d	evelo	pment
of hypocal	cemia will be	e documente	d. This w	ill b	e cor	relat	ed with	any fu	nctio	nal change
in renal m	etabolism.									
25. (U) 7	5 07 - 76 09	This prot	ocol has b	een c	omple	ted v	vith the	entry (	of 20	patients
into the s	tudy. To dat	te the resu	ults of the	seru	m and	urin	ne studi	es have	been	tabulated
but the re	sults from th	he dietitia	an concerni	ng th	e int	ake (	of calci	um and	phosp	horus are
not yet av	ailable. In	addition,	the studie	s of	parat	horm	one to b	e perfo	rmed	by
Dr. Goldsm	ith at the A	udie Murphy	/ Veteran A	dmini	strat	ion 1	lospital	in San	Anto	nio have
not yet he	en completed	due to the	e temporary	inte	rrupt	10n 3	in his a	ssay pr	ocedu	re. It i
expected t	hat the resu	lts will be	e tabulated	and	concl	us 10	ns drawn	within	the	next
several mc	nths.									
1										
1										
A suileble to contract	iora ign ertamaterta av	NUTEI							ni succe Thigh	

Sec. 1

Contraction of the second

ないない

and the provide the second second second second

and a second 
#### TERMINATION

# PROJECT NO. 3/161102B71P-02 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the s

REPORT TITLE: EVALUATION OF CALCIUM METABOLISM IN BURNED TROOPS

## US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON; TEXAS 78234

1 July 1975 - 30 September 1976

#### Investigators

Richard H. Merrill, MD, Lieutenant Colonel, MC William D. Myers, Lieutenant Colonel, MC Harry R. Jacobson, Major, MC

Reports Control Symbol MEDDH-288(RI)

UNCLASSIFIED

## ABSTRACT

## PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

REPORT TITLE: EVALUATION OF CALCIUM METABOLISM IN BURNED TROOPS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Richard H. Merrill, M.D., Lieutenant Colonel, MC William D. Myers, M.D., Lieutenant Colonel, MC Harry R. Jacobson, MD, Major, MC

Reports Control Symbol MEDDH-288 (RI)

This study has been completed with the entry of 20 patients into the study. Tc date the results of the serum and urine studies have been tabulated but the results from the dietitian concerning the intake of calcium and phosphorus are not yet available. The observed early hypocalcemia has been a consistent observation. Skin calcium levels have not been elevated so the calcium is either being excreted in the urine and/or stool, or being sequestered. In view of the observations with acute renal failure where it tends to be elevated, it is unlikely that the bone is acting as a sump. The studies of parathormone (P^{*}H) to be performed by Dr. Goldsmith at the Audie Murphy VA Hospital, San Antonio, have not been completed due to the temporary interruption in his assay procedure. It is expected that the results will be tabulated and the conclusions drawn within the next several months. It is hoped that a method for measuring urinary CAMP will be developed at USAISR shortly, and these studies will suppert the PTH data.

Calcium Burns Renal failure Hypocalcemia Humans

RESEARCH	AND TECHNOLOG	WORK UNIT SU	MARY	I AGEN	CY ACC	6057	2 DATE OF SU	01	REPORT CONTROL SYMBOL DD-DRAE(AR)616		
A DATE PREV SUMPRY	4 KIND OF SUMMARY	S. SUMMARY SCTY	WORK SECURITY	DA BEAR	Ding	0952	70 10	ISA SPECIFIC	DATA TO LEVEL OF SUM		
75 07 01	D. CHANGE	U	U	NA			NL	CONTRACTOR	TOR ACCESS A WORK W		
10 NO /CODES:*	PROGRAM ELEMENT	PROJECT N	UMBER	TASK A	REAI	NUMBER	WORK UNIT NUME				
	611018	3016110109		00 082							
b. CONTRIBUTING											
C. CONTRIBUTING											
11 TITLE (Procode with	Security Classification Code	(U) Laborat	ory Inves	tigat	ion	of t	he Mecha	nisms of	E scqu	ired	
Leukocyte Dysfunction Following Thermal Injury (44)											
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS										
003500 Clin	nical Medicin	2									
13 START DATE		14. ESTIMATED COMPL	ETION DATE	15. FUNT	HNG A	GENCY		IS PERFORM	ANCE NETH	00	
75 12		Cont		DA	1		1	C. In	-House		
17 CONTRACT/GRANT	Not Applic	able		-	DURCE	S ESTIMAT	-	IONAL MAN YR		\$ (In these ands)	
& DATES/EFFECTIVE		EXPIRATION			PRECI	76	1.0		1	4	
N NUMBER				FISCAL		7т	.3		4		
G 1998		A ANOUNT		YEAR	CONN	ENT					
& KIND OF AWARD:		I. CUM. AMT.		77 1.0 20					0		
IS RESPONSIBLE DOD	BLE DOD ORGANIZATION 30 PERFORMING ORGANIZATION								1.		
NAME." US Army	Institute of	Surgical Re	search	-	US	Army	Institut	e of Su	rgical	Research	
				i	М	licrob	iology B	ranch			
ADDRESS:" Fort	Sam Houston, '	<b>Texas 78234</b>	ł		• F	ort S	am Houst	on, Tex	as 78	234	
				1							
				PRINCIP	AL INV	ESTIGATO	R (Furnish SEAN	II U S Acad wh	Institution)		
RESPONSIBLE INDIVIDU	JAL			NAME.	• Al	bert	T. McMan	us, CPT	, MSC		
NAME: Basil	A. Pruitt, Jr	., COL, MC		TELEPHONE: 512-221-3411							
TELEPHONE:	512-221-2720			SOCIAI	SECU	RITY ACCO					
21. GENERAL USE	,			ASSOCIA	TE INV	ESTIGALO	A1				
FOREIGN IN	TELLIGENCE NO	T CONSIDERED	)	HAME:	Ar	thur	D. Mason	, Jr.,	MD		
				HAME						DA	
IL KEYWORDS (Procedo	BACK with somethy classifi	callen Code)									
(U) Burns;	(U) Leukocyt	es; (U) Gluc	cose oxida	tion	; (U	)) Lat	ex phago	cytosis			
23 TECHNICAL OBJECT	11VE, * 24. APPROACH, 25.	PROGRESS (Fumish ind	ividual peragraphie id	entitled by	nunbe	r procode i	est of each with a	jeculty Cleesift	cation Code )		
23. (U) E	fforts will b	e made to es	stablish o	ne or	c mo	ore me	tabolic	basis f	or acq	uired	
leukocyte	dysfunction f	ollowing the	ermal inju	ry.	Est	ablis	hment of	specif	ic nut	ritional	
or environ	mental effect	s may allow	for corre	ctive	e ma	inagem	ent.				
				_				•			
24. (U)	Initial effor	ts will be t	to measure	gluo	cose	meta	bolism i	n norma	1 and	burned	
patients	leukocytes.	Purified and	d washed g	ranu	locy	vte po	pulation	s will	be exa	mined	
for oxidiz	for oxidization of Carbon 14 labeled glucose. Hexose monophosphate shunt and										

glycolysis activity will be estimated by release of  $14_{CC}^{2}$  from respectively 1-14C glucose. Measurements will be made on resting and latex particle (=0.8 u) stimulated cells.

25. (U) 75 12 - 76 09 A standarized latex concentration response assay has been developed. Normal values and confidence limits have been established. Examination of patients' cells has begun with eight patients studied to date. In both stimulated and unstimulated leukocytes from burn patients both pathways of glucose metabolism were suppressed.

Available to contractors upon originator's approval PREVIOUS COLITIONS OF THIS FORM ARE DESOLETE DD FORMS 14984 I NOV 85 AND 1498 I I MAR 68 (FOR ARMY USE) ARE DESOLETE

A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PERSON AND A PER

# PROGRESS REPORT

## PROJECT NO. 3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH

# REPORT TITLE: LABORATORY INVESTIGATION OF THE MECHANISMS OF ACQUIRED LEUKOCYTE DYSFUNCTION FOLLOWING THERMAL INJURY

## US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

a de calata

a 10 e 10 d

t.

1 July 1975 - 30 September 1976

Investigators:

Albert T. McManus, Jr,Cpatain,MSC Thomas J. Lescher, MD,Major,MC Arthur D. Mason, Jr,MD Basil A. Pruitt,Jr, MD,Colonel, MC

Report Control Symbol MEDDH-288 (R1)

UNCLASSIFIED

#### ABSTRACT

## PROJECT NO.3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH

## REPORT TITLE: LABORATORY INVESTIGATION OF THE MECHANISMS OF ACQUIRED LEUKOCYTE DYSFUNCTION FOLLOWING THERMAL INJURY

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Albert T. McManus, Jr, Captain, MSC Thomas J. Lescher, MD, Major, MC Arthur D. Mason, Jr, MD Basil A. Pruitt, Jr, MD, Colonel, MC

Report Control Symbol MEDDH-288 (R1)

The metabolic alterations underlying postburn anomalies of granulocyte function have been studied by examining "C-glucose exidative metabolism in granulocytes from burned patients under serum free, chemically defined conditions. Granulocytes from burned patients and controls were prepared from heparinized (10 units/ml) venous blood by dextran sedimentation, NH₂Cl treatment and hypaque-ficoll density gradient centrifugation. Cells were examined for "CO₂ production from either "C-1-glucose or "C-6-glucose during a 60 minute incubation at 37°C with or without an optimal stimulatory dose of 0.79 u latex particles. Reaction mixtures contained 2-4.5 x 10° granulocytes/ml and a specific activity of 1 u Ci/5.5 uM glucose/ml. Phagocytosis of latex particles in the absence of serum was confirmed by scanning electron microscopic observation.

Examination of 33 normal male volunteers, ages 20-43, indicated ¹⁴CO, production values from C-1 or C-6 glucose to be distributed according to a log normal distribution. Eight thermally injured male patients, ages 18-37, with a range of total body surface burn of 60-90%, mean 70.2%, were examined 72-96 hours postburn. Patient mortality was 75%. Data are presented as In mean CPM/10° cells + 95% confidence limits.

	1*-C-C	lucose	6*-C-Glucose						
	w/o Latex	w/Latex	w/o Latex	w/Latex					
Control	5.5853 <u>+</u> 0.1306	7.3800 <u>+</u> 0.0851	2.4784 <u>+</u> 0.1292	4.4244 <u>+</u> 0.1544					
Burn	4.6115 ± 0.3678	6.6893 <u>+</u> 3355	1.3668 <u>+</u> 0.3167	2.5557 <u>+</u> Q.4795					

Significance testing showed that cells from patients had reduced  ${}^{14}CO_2$  production under all tested conditions (p < 0.001). This constant decrease

in oxidative metabolism of glucose by granulocytes following thermal injury may be a basis for the leukocyte dysfunctions previously demonstrated in like patient populations.

Burns Leukocytes Glucose oxidation Latex phagocytosis

•

RESEARCH AND TECHNOLOGY WORK UNIT SUMMARY			DA	OG 6953	3 DATE OF 5 76 10	01	REPORT CONTROL SYNGOL DD-DR&E(AR)636				
75 J7 01	A. KIND OF SUMMARY	S. SUMMARY SCTY	A. WORK SECURITY	7 REGR	ADING ^{II} BA D	NL	AL SPECIFIC		A. WORK WAT		
10. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK	AREA HUNDER	T	R				
. PRIMARY	61101A	3A161101A	.91C	0	0	084					
S. CONTRIBUTING				1							
C. CONTRIBUTING						1	<u></u>				
11. TITLE (Procode mith .	Security CloselHeation Code	(J) Echoc	ardiograph	ic Ev	aluation	of Left	Ventric	nilar			
Performance	in the Sever	ely Burned	Military	Popul	ation (4	4)					
12. SCIENTIFIC AND TE	CHNOLOGICAL AREAS								·······		
003500 Clin	ical Medicine	•									
IS START DATE		14 ESTIMATED COM	LETION DATE	IS FUN	DING AGENCY		16 PERFORM	ANCE MET	HOD		
76 06		Con	t	DA	1		C. Ir	-Hous	Se .		
17. CONTRACT/BRANT	Not Applicat	le		10. NES	DURCES ESTIMAT		BIONAL MAN YR		till the themendet		
& DATES/EFFECTIVE:		EXPIRATION:			PRECEDING	-			10		
\$ NUMBER:*				FISCAL	70	.9		1 4	18		
& TYPE:		4 AMOUNT:		YEAR	CUNNER		······				
. KIND OF AWARD		LCUM, AMT.			77	.5			25		
H. RESPONSIBLE DOD C	MEANIZATION				ORMING URGANI	ZATION					
ADDRESSI® Fort Sam HOUSTON, Texas 78234 RESPONSIBLE INDIVIDUAL Surgical Study Branch Fort Sam Houston, Texas 78234 PRINCIPAL INVESTIGATOR (Pumlet Man II & S Assessed prelimition) NAME® James F. Dorethy, MAJ, MC							78234 v				
TELEPHONE: 51	2-221-2720	, соц, мс		TELEPHONE 512-221-5712 SOCIAL SECURITY ACCOUNT NUMBER							
I. SEMERAL USE				ASSOCIA	TE INVESTIGATO	RS					
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	MANE							
				NAME.					DA		
IL REVIERCE (Procedo I	LACH with Somelly Classific	ailan Code) (U) P	ostburn she	ock;	(U) Echoo	cardiogr	aphy;				
(U) Resusci	tation burned	soldiers;	(U) Left	ventr	icular fu	inction					
B TECHNIC & OBJECT	VE. 4 24 APPROACH, 28.	PROGRESS (Furnish in	Nviduai paragrapha idi	millind by	number Precede s	ezi el each with	Security Classific	ellan Code			
23. (U) I especially depression	o evaluate le in postburn s is a direct c	ft ventric hock phase onsequence	ular funct. . To eval of severe	ion i uate ther	n the the the hypot mal inju	ermally thesis t cy.	injured hat myoc	patie ardia	ent, al		
24. (U) S graphically	erial left ve and correlat	ntricular ed to the	performance clinical st	e pro tate	files wi of the pa	ll be de atient.	erived ec	chocar	dio-		
25. (U) 7 serial pati assess the endocarditi patients wi these patie	6 06 - 76 09 ent studies h configuration s. No valvul th blood cult nts who died	Due to de ave just b of cardia ar vegetat ures posit autopsy ex	layed equip een initia c values in ions were : ive for Sta amination :	pment ted. n pat ident aphyl revea	delivery The inst lents sus ified by ococcus a led norma	y and su trument spected echocar aureus a a'. cardi	bsequent has beer at havir diograph and in th ac value	: cali used ng það ny in ne fou es.	ibration l to eterial 15 ar of		

-

- ----

i....

HAMELOFE SPRIVEL PREVIOUS EDITIONS OF THIS FORM ARE DESOLETE OD FORMS 1438A I NOV 85 AND 1498 1 I MAR 18 FOR ARMY USE ARE OBSOLETE 297 DD FORM 1498

-

ş

\$

#### ANNUAL PROGRESS REPORT

the statistical of the statistic best and the statistics

## PROJECT NO. 3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH

REPORT TITLE: ECHOCARDIOGRAPHIC EVALUATION OF LEFT VENTRICULAR PER-FORMANCE IN THE SEVERELY BURNED MILITARY POPULATION

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUATON, TEXAS 78234

> > 1 July 1975 - 30 September 1976

Investigators:

James F. Dorethy, M.D., Major, MC

Reports Curtrol Symbol MEDDH-288(R1)

Unclassified

#### ABSTRACT

**PROJECT NO.** 3A161101A91C-00, IN-HOUSE LABORATORY INDEPENDENT RESEARCH **REPORT TITLE:** ECHOCARDIOGRAPHIC EVALUATION OF LEFT VENTHICULAR PER-

FORMANCE IN THE SEVERELY BURNED MILITARY POPULATION

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: James F. Dorethy, M.D., Major, MC

**Reports** Control Symbol MEDDH-288(R1)

Complex cardiovascular interactions following acute thermal injury have been studied by indirect methods of myocardial performance. Myocardial depression has been implicated as an etiology for decreased cardiac output during the early postburn period. No direct measurements of left ventricular performance have been reported.

This study proposes to utilize sensitive echocardiographic measurements of LV internal dimensions to evaluate postburn myocardial contractility. Due to delayed equipment delivery and subsequent calibration, only two serial studies have been carried out in the early resuscitative phase. Fourteen other patients have been studied serially for possible endocarditis. Two patients have been evaluated during septic shock.

No specific conclusions can be drawn from this small group of patients. However, this noninvasive technique appears feasible for serial studies of LV performance post-thermal injury.

Resuscitation Burned soldiers Echocardiography Left ventricular function Postburn shock

RESEARCH A	RESEARCH AND TECHNOLOGY WORK UNIT SUMMARY				CY ACCE	389	76 10 0	1	REPORT CONTROL STREDG DD-DR&E(AR)636		
76 10 01 H	RIND OF SUMMARY I. TERM	S. SUMMARY SCTY	6. WORK SECURITY	7 ALGR	DING	NL	ig'n insta'n	THE SPECIFIC	ACCESS	A WERK UNIT	
10 NO / CODES:* PI	ROGRAN ELEMENT	PROJECT	NUMBER	TASK /	REA NU	MBER					
B PRIMARY	61102A	3A161102	2B71P	(	)2		070				
S. CONTRIBUTING											
C. CONTRIBUTING								di carifuan andu			
11 TITLE (Procode with Some	with Classification Code)	•	• _								
(U) Renal Fun	(U) Renal Function in the Burned Soldier. I. Histology (44)										
12 SCIENTIFIC AND TECHN	OLOGICAL AREAS										
003500 Clinic	al Medicine										
15 START DATE		14 ESTIMATED COMP	LETION DATE	IS PUN	NHS AGE	NCY		16. PERPORM	ANCE MET	HOD	
75 O ₁		76 09	-	DA				C. In-House			
17 CONTRACT/GRANT	CONTRACT/GRANT NOt Applicable					ESTIMATE	-	NORMAL MAN YAS & FUNDS (In these and a)			
& DATES/EFFECTIVE:		EXPIRATION			76		.5			22	
NUNDER:*				FISCAL 7T			.2			3	
G TYPE:		4 ANDUNTI		YEAR	CURREN	•	T				
		I. CUM. AMT.									
IT RESPONSIBLE DOD ORE	ANIZATION			30. PERI	0744116	ORGANIZ	ATION				
MAPE:" US Army I	institute of	Surgical 1	Research	NANE:"	JS Ar Re	my I enal	nstitute Section,	of Sur Metabo	gical lic B	Research ranch	
ADDRESS: Fort Sam	Houston, T	exas 78234	1	A DOME IN	" For	t Sa	m Housto	n, Texa	s 78.	234	
				PRINCIPAL INVERTIGATOR (Fundob SEAR II U.S. Associate publication)							
RESPONSIBLE INDIVIDUAL				NAME:	Will	liam	D. Myers	, LTC,	MC		
NAME: Basil A.	Pruitt, Jr	, COL, MC		TELEP	MONE:	512-	221-6532				
TELEPHONE. 512-2	21-2720			10CHA	SECURIT	TY ACCO	INT NUMBER:				
21 GENERAL USE				A360CI A		-	6				
FOREIGN INTEL	LIGENCE NOT	CONSIDERE	כ	NAME:	The	mas	Rizzo Jr	, MAJ,	MC		
				NAME:	Ric	hard	H. Merr	ill, LT	С, МС	DA	
II. KEYWORDS (Freesde EAC	H all socally Classific	atlan Code)									
(U) Glomerulu	is; (U) Scan	ning EM; (	J) Transmis	sion	EM;	(U) B	urn; (U)	Azotem	ia; (	U)Humans	
23 TECHNICAL OBJECTIVE.	TECHNICAL OBJECTIVE, * 24. APPROACH, 25. PROGRESS (Punnish Individual paragraphic Idaniiliad by number Prezide real of each with security Classification Code.)										

Service and the service and the service of the serv

23. (U) To identify electronmicroscopic glomerular pathology postburn and relate it to renal function changes in burned soldiers.

24. (U) All patients in the burn unit with immediate postmortem authorization will undergo within one hour postmortem percutaneous needle biopsy of kidney. Tissue will be examined by light, electron, and immunofluorescent microscopy. Premortem renab function will be monitored by urinalysis, BUN, serum creatinine, urine creatinine, urine sodium, and urine potassium. Patients will be grouped by serum creatinine and pattern of function (6p.1:Cr less than 1.2; 6pII:Cr more than 1.2; A:prerenal; B:renal). Clinical course noting type of burn, crush injury, resuscitation, coagulopathy, shock, sepsis; hypoxia, and medications will be recorded for future correlation.

25. (U) 75 07 - 76 09 'Twenty-five patients have been studied. Ten specimens were proressed for immunoglobulin deposition and were interpreted as having no immunoglobulins present. Seventeen had interpretable transmission EM findings. The transmission EM findings included subendothelial rarefaction, segmental foot process fusion, increased mesangial matrix, and subendotnelial dense deposits. No correlation could be found between renal function and EM alterations. Presently an effort is being made to correlate renal function with surface alterations demonstrated by scanning EM.

## TERMINATION

## PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

REPORT TITLE: RENAL FUNCTION IN THE BURNED SOLD:ER. I HISTOLOGY

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators

NO.

William D. Myers, MD, Lieutenant Colonel, MC Thomas Rizzo, Jr., MD, Major, MC Richard H. Merrill, MD, Lieutenant Colonel, MC John McPhaul, MD, Colonel, MC

Reports Control Symbol MEDDH-288 (RI)

UNCLASSIFIED

## ABSTRACT

# PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

## REPORT TITLE: RENAL FUNCTION IN THE BURNED SOLDIER. I. HISTOLOGY

U. S. Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: William D. Myers, MD, Lieutenant Colonel, MC Paulette Langlinais, MS Thomas Rizzo, Jr., MD, Major, MC Richard H. Merrill, MD, Lieutenant Colonel, MC John McPhaul, MD, Colonel, MC

Reports Control Symbol MEDDH-288(R)

and the later of the second states of the second st

The purpose of this study was to determine if fibrin disposition immune complex disease, or toxic basement membrane alteration could explain the azotemia seen in the thermally injured patient. To answer this question patients undersent needle biopsy of the kidney within one hour of death and the tissue was submitted for light, electron, and immunofluorescent microscopy. Of the 25 samples 17 were adequate for electronmicroscopy, and 9 were adequate for immunofluorescent microscopy. Electronmicroscopy revealed spreading of lamina rara interna in 15, atypical subendothelial dense deposits in 5, increased mesangial matrix in 9, and focal foot process fusion in 7. Immunofluorescent microscopy of IgG, IgM, IgA, fibrinogen, and complement was negative in 8. One patient exhibited 1+ linear staining of uneven intersity from glomerulus to glomerulus of IgC, and trace to 1+ mesangial streaking of Complement (B,C). Histopathologic findings could not be correlated with serum creatinine levels in these patients.

Most electonmicroscopic alterations were considered to be due to early autolysis. There was no evidence that fibrin or immune complex deposition or basement membrane alteration was responsible for the altered renal function.

Glomerulus Scanning EM Transmission EM Burn Azotemia Humans

## RENAL FUNCTION IN THE BURN PATIENT: I. HISTOLOGY

The incidence of renal insufficiency following thermal injury has decreased with improved initial fluid resuscitation. However, even with adequate resuscitation, azotemia, which progresses to marked renal insufficiency, is occasionally seen. The typical pattern is that of a "prerenal" insufficiency manifested by a BUN/creatinine ratio greater than 10, urine urea nitrogen/serum urea nitrogen or urine creatinine/ serum creatinine ratio greater than 20, urine specific gravity greater than 1.015, and a low excretion of sodium with a high excretion of potassium. The tubules continue to perform maximally as glomerular filtration progressively declines, revealing a glomerulotubular imbalance.

Graber and Sevitt² observed occasional lipid deposits in glomeruli and areas of tubular necrosis in the thermally injured patient. However, in spite of microscopic pathology, the tubules were able to reabsorb sodium and excrete potassium normally. It may be that the reduction in urine flow allows the tubules to reclaim most sodium from the slowly moving filtrate. A block in plasma flow from the glomerulus to the tubule could explain the clinical findings. Graber and Sevitt described multiple fine droplets of fat within the glomerular tufts and possibly within the endothelium and epithelium. However, no electron microscopy was performed. Tissue submitted from ISR autopsy specimens in the past for electron microscopy and immunofluorescent microscopy have been unsatisfactory for detailed study. Tissue obtained within one hour of death will hopefully provide interpretable, study material.

This study will attempt to identify any electronmicroscopic glomerular pathology associated with normal and abnormal renal function in the thermally injured patient. All patients with immediate postmortom authorization will undergo within one hour postmortem percutaneous needle biopsy of the kidney. Tissue will be examined by light and electron microscopy at the ISR Lab. Renal tissue will also be rapidly frozen and sent to Dr. McPhaul at Wilford Hall Air Force Hospital for immunofluorescent microscopy to determine the presence of IgG, IgA, fibrin, and complement. Renal function will be monitored by ur.nalysis, BUN, serum creatinine, urine creatinine, urine sodium, and urine potassium. Patients will be grouped by serum creatinine and pattern of function (Group 1. Cr less than 1.2; Group II: Cr greater than 1.2; A: prerenal; B: renal). Clinical Course, noting time of burn,

1. Moncrief JA: Burns. New Eng J Med 288: 444, 1973.

2. Graber IG, Sevitt S: Renal function in burned patients and its relationship to morphological changes. J Clin Path 12: 25, 1959.

crush injury, resuscitation, coagulopathy, sheck, sepsis, hypoxia, and medications will be recorded for future correlation.

#### RESULTS

Twenty-five patients were biopsied up to two hours after death. Of the twenty-five patients, only seventeen had interpretable electronmicroscopic findings. The electronmicroscopic glomerular alterations observed were subendothelial rarefaction in sixteen patients, segmental foot process fusion in eleven patients, increased mesangial matrix in ten patients, and subendothelial dense deposits in five patients. Of the eight uninterpretable specimens, four contained no glomeruli, two were mixed, and two were lost.

Of the twenty-five patients biopsied only ten had adequate samples for immunofluorescence microsocpy. Nine specimens failed to demonstrate immunofluorescence to IgG, IgM, IgA, complement, or fibrinogen. One patient exhibited 1+ linear staining of uneven intensity from glomerulus to glomerulus of IgG, and trace to 1+ mesangial streaking complement (B₁C).

Table I summarizes relationships between electron microscopy, immunofluorescent microscopy, urine fibrin split products, maximum serum creatinine, burn size, and recognized nephrotoxins.

#### DISCUSSION

Several years ago Graber and Sevitt² addressed the relationship between renal histopathology and function. Tubular necrosis was observed in patients with normal renal function, and normal tubules were observed in patients with clinical renal failure. They concluded that there was little correlation between renal tubule histology and function, but suggested that a pretubular anatomic defect might explain the discrepancy. This study examined the relationship of renal function to glomerular histology.

Our clinical observations suggested that the renal insufficiency observed in the burn population was of the "prerenal" type. This was suggested by a BUN rising out of proportion to creatinine, very low urinary sodium excretion, very high urinary potassium excretion, UUN/BL N and Ucr/Scr ratio greater than 20. Such a pattern is compatible with glomerular tubular imbalance i.e. decreasing filtration in the presence of good tubular function. Mechanical blocking of flow from the blood side to the urine side of the capillary loop as in acute glomerulitis could give this functional pattern.

This study found no significant immune deposits in the glomeruli thus ruling out antigen-antibody type damage. Fibrin was also absent on electron and light microscopy. However, all most all glomerular capillary loops demonstrated some abnormality on electron microscopy, but none of these abnormalities could be correlated with level of serum creatinine. These results suggest that there is no gross mechanical blockage of flow from the blood side to the Bowman's space side of the capillary loop by

(EX P MORT)	<u>EN</u>		0 PSP	<u>.</u>		TOXING				
PK (60)	Subon. rare w/ DD	Neg	*1	1.)	-	Colis, V				
VP (60)	Suban, rare, Sag PP Fus.	Neg	16: 1	5.0	73	Colis -				
TD (44)	Suben. rare, Seg 🏴 Pus	Neg	4: 1	386	66	Cells				
WE (60)	Suben, rare, Seg PP Fus, M	U	1:1	2.186		G				
MG (40)	Subon. rare, Seg PP Fus, M	Neg	<b>8</b> :1	4.9 ⁸⁶	•	Cells , G				
<b>TF (60)</b>	Subon. rare, Seg PP Fus, M	U	<b>8:1</b>	1.6	. <b>5</b> 8	Colis				
DK (60)	Suben.rare Seg PP Fus, M	U		2.3	76					
JC (60)	Subon. rare, Sog FP Fue, M	U		. 2.3 ⁸⁶	82	Colis				
WR (60)	Suben. rare, Seg PP , Kus, M		4: 1	<b></b>	82	Colis				
ICEN (45)		٠	٠	2.6		Celle G.				
RM (90)	Suben, rare, Seg PP Fus, M	Neg	4:1	6.4 ⁸⁶	57	Cells'G				
06 (120)	Bil, DM, Seg FP Fus	Neg	1:1	2.186		Cells				
CL (90)	Mined	U	12:1	BUN30						
LAD (30)	Minad	U	1:1	0.8	67	Cety, G				
LJ (130)	NG	U	8: 1	3	75					
C\$ (34)	NG	U	4: 1	6.2 ⁸⁶	67	Cely V				
RD (99)	Suben. rare, M	U	4:1	<b>0.6</b> ·	50					
DC (130)	NG	U	1:1	3.686	51	G				
RS (124)	NG	U	1:1	3.5 86	78	Cells				
68	Suben.rare w/DD		#1	4 ⁸⁶	50	Celis, G				
WC	Suten rarew/DD, SegFPFus, M		4:1	1.2	33	Cells, V				
AV				6.386	<b>\$7</b>	Cells,G				
DB (60)	Suben.rare,M	Neg		0.7	79					
DB (60)	Subon . rore (Minimal)	Neg		1.3	70					
W5 (60)	Suben, vara w/DD	Neg		4.4 ATN	ж	Celis				

----

-----

ر با جو

and the weather and the second se

ł

# KEY TO TABLE I

Suben, rare - Subandoth: Hat rerifaction	DD - Dense deposit
Seg. PP Fus - Segmental Jeet Process Fusion	M ~ Mesongial hypertrophy
BM - Thickened Basement Membrane	DM - Diabetes Mellitus
Mixed - Specimens kviermixed	NG - No giomeruli
U - Unsatisfactory	85 - Burn Syndrame (Prerenel)
ATN - Acute Tubular Necrosis	Calis - Colymicin
V - Vancemycin	G - Gentamycin
PSP - Fibrin Split Products	

immunoglobulins or fibrin. Although electronmicroscopic alterations could account for reduced filtration, it is unlikely since serum creatinine varied from 0.7 to 8.0 regardless of electronmicroscopic alterations. The histologic findings in this study may represent only postmortem autolysis.

Recently scanning electronmicroscopy has revealed glomerular surface alterations in acute renal failure models², again raising the question of relationship between histologic and functional alteration. These surface changes and their relationship to renal function need to be further evaluated. Such an investigation is presently underway.

## PRESENTATIONS AND/OR PUBLICATION

None

^{3.} Cox JW, et al: Studies on the mechanism of oliguria in a model of unilateral acute renal failure. J. Clin. Invest. 53: 6, 1974.

DECENDEN						sion	2. DATE OF SUMMARY ARE			PORT CO	TROL STHEOL
RESEARCH	AND TECHNJLUGT	WORK UNIT 5	UMBART	DA	OG 69	61	76 10	01		DD-DR	E(AR)636
I. DATE PREV SUM'RY	4 KIND OF SUMMARY	S SUMMARY SCTY	S WORK SECURITY	T REGR	ADING	- 0		AL SPEC	IFIC DA	TA- 0	LEVEL OF SUM
76 10 01	H. TERM	U	U	N	A	1	NL	1 ves			A VORK UNIT
10 NO /CODES *	PROGRAM ELEMENT	PROJECT	NUMBER	TASK AREA NUMBER WORK UNIT NUMBER						UMBER	
L PRIMARY	61102A	3A161102	B71P	01 021							
& CONTRIBUTING											
C CONTRIBUTING							<u></u>		•	•	
11 TITLE (Procode with	Security Classification Code;	•									
(U) Localiz	ation of Infe	ection With	in the Uri	nary	Tract	of	Burned	Soldi	ers	(44)	
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS										
003500 Clir	nical Medicine	2									
13 START DATZ		14 ESTIMATED CON	LETION DATE	IS FUNDING AGENCY				14. PER	16. PERFORMANCE METHOD		
75 11		76 09		DA				C. In-House			•
17 CONTRACT/GRANT NOT Applicable				IS RESOURCES ESTIMATE & PRO			E A PROFES	PROFESSIONAL MAN YRS & FUNDS (In these			(In these ands)
A DATES/EFFECTIVE		EXPIRATION			76		A			15	
NUMBER *				FISCAL 7T		.2			Ž		
-		A ANOUNT.		YEAR	CONNEN		**************************************				
. KIND OF AWARD		f. CUM. AMT.		Ι.							
IS RESPONSIBLE DOD	DRGANIZATION		1	SO PER	PORMING	AGANI	LATION			· · · · ·	T
NAME TIS AVMU	Institute of	Surgical B	ecoarch	-	-		t		Curre		Decemple
oo namy	Institute of	burgicar r	esearch	US Army institute of Surgical Research							
ADDRESS." Fort	Com Nouston P	Downe 7037		ADDRES	Ner Rer		Section				
roit.	sall nouscon,	lexris /023			FOI	τ 30	am Houst	con, 1	exas	/84	234
				PRINCIP	-	TISATO	R (Fumish SEAN		danic ja	e i i tro Manj	
RESPONSIBLE INDIVIDU	IAL				• Will	iam	D. Myei	cs, LI	C, M	IC	
HAME Basil A. Pruitt, Jr, COL, MC				TELEPHONE 512-221-6532							
TELEPHONE. 512	2-221-2720			SOCIAL	L SECURIT	* ****					
21 GENERAL USE ASSOCIATE INVESTIGATORS											
FOREIGN IN	TELLIGENCE NOT	CONSIDERE	D	-	Dr.	Ge	rsh				
				NAME Thomas Rizzo, MAJ, MC							DA
THE REAL PROPERTY OF	RACH WIR RAWARD Closell	anti-m dadat								_	

(U) Yeast; (U) Bacteria; (U) Immunofluorescence; (U) Urimary tract; (U) Humans

23 TECHNICAL OBJECTIVE." IL APPROACH, IL PROGRESS (Pumile) individual programe identified by number procede lost of each side pocurity Classification code) 23. (U) To determine the degree of spread of infection from the lower to the upper urinary tract in burned soldiers.

24. (U) Patients with indwelling bladder catheters will have urine examined daily for bacteria and/or yeast. Those patients found to have organisms in the urine will have their urine processed to determine if the organisms are coated with immunoglobulin. Patients found to have organisms which are coated with immunoglobulin will have urine sent to bacteriology for culture and sensitivity testing so that appropriate antibiotic therapy can be instituted. Those patients coming to post mortem examination will have detailed examination of the urinery tract for organism invasion.

25. (U) 75 11 - 76 09 Twenty patients have been evaluated to date. Of the five with bacteruria none were demonstrated to have immunoglobulin coating, suggesting only bladder infection. Of the fifteen with candiduria, only those with hyphae were found to have immunofluorescence. No correlation has been made between anatomic site of infection by postmortem examination and yeast immunoglobu in coating. No conclusions can be drawn until more patients are studied.

silable to contractore upon originator's approve

and the star strength

which is a start of the start o

PREVIOUS EDITIONS OF THIS FORM ARE /BSOLETE DD FORMS 1498A 1 NOV 66 AND 14981 1 MAR 68 (FOR ARMY USE) 4 48 OBSOLETE 307

#### TERMINATION

# PROJECT NO. 3A161102B71P-01, EASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

# REPORT TITLE: LOCALIZATION OF INFECTION WITHIN THE URINARY TRACT

## US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators

William D. Myers, M.D., Lieutenant Colonel, MC Harvey A. Gersh, M.D., Major, MC Thomas Rizzo, Jr., M.D., Major, MC Marvin Forland, M.D. V. Thomas, PhD

Reports Control Symbol MEDDH-288(RI)

UNCLASSIFIED

#### ABSTRACT

## PROJECT NO. 3A161102B71P-01 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

## REPORT TITLE: LOCALIZATION OF INFECTION WITHIN THE URINARY TRACT

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: William D. Myers, M.D., Lieutenant Colonel, MC Harvey A. Gersh, M.D., Major, MC Thomas Rizzo, Jr., M.D., Major, MC Marvin Forland, M.D. V. Thomas, PhD

Report Control Symbol MEDDH-288 (RI)

are strike here and the strike and the strike is the strike is the strike strike is the strike of the strike and the

Budding yeast and hypae are frequently observed in the urine of thermally injured patients requiring prolonged indwelling urinary blacker catheterization and systemic antibiotics. This investigation was designed to determine if yeast infections of the urinary tract can be localized by staining for immuno oglobulin on the yeast, to determine what percent of catheterized patients develop upper tract disease, and to determine if appropriate therapy can reduce the incidencof sepsis by identifying upper tract disease early.

Twenty patients have been evaluated to date. Of the 5 with bacteriuma, none were demonstrated to have immunoglobulin coating suggesting no extrabladder infection. Of the 25 with candiduria, only those with hyphae were found to have immunofluorescence. No correlation has been made between anatomic site of infection by postmortem examination and yeast immunoglobulin coating. No conclusions can be drawn until more patients are studied.

Bacteriuria Immunoglobulin Infection Soldier Yeast Immunofluorescence

### LOCALIZATION OF INFECTION WITHIN THE URINARY TRACT

Budding yeast and hyphae are frequently observed in the urine of thermally injure patients requiring prolonged indwelling urinary bladder catheterization and systemic antibiotics. Hyphae in the urine have been considered a sign of tissue invasion, rather than just a sign of contaminated urine in the bladder; however, if the infection is limited to the bladder, it can be managed by local irrigation of the bladder with 5 flucytosine or amphotericin, thus reducing the toxicity of systemic therapy². If the kidney was infected, local bladder irrigation would not be sufficient. Recently, it has been reported that bacteriuria can be localized as coming from the kidney or bladder by staining the urine for bacteria coated with immunoglobulins^{3,4}. Those bacteria originating in the kidney would be immunoglobulin coated and those coming from the bladder should not be coated. However, one must be careful when the patient is a male because prostatitis can a'so coat bacteria; therefore, "everything that coats" is not always of kidney origin. If yeast in the urine, originating from areas other than the bladder, were also coated with an immunoglobulin, then this could be used to determine if the patient should be managed with systemic antifungal therapy in addition to local therapy.

Previous investigators at this institute have reported that once the urinary catheter is removed, bacteriuria frequently resolves without therapy. Those patients who did not have spontaneous resolution of bacteriuria may have had upper tract infections requiring further management. By staining the bacteria in the urine for immunoglobulin, one can identify this group and manage them appropriately.

This investigation was designed to answer the following questions:

1. Can yeast infections of the uninary tract be localized by staining for immunoglobulin on the yeast?

2. What percentage of catheterized patients develop upper tract disease?

3. Can appropriate therapy reduce the incidence of sepsis of identifying upper tract disease early?

^{1.} Kosinn PJ, Taschdjian CL: Enteric Candidiasis: diagnosis and clinical considerations. Pediatrics 30: 71, 1962.

^{2.} Wise GJ et al: Candidal cystitis. JAMA 224: 1636, 1973.

^{3.} Thomas V, et al: Antibody-coated bacteria in the urine and the sites of urinery tract infection. New Eng J Med 290: 558, 1974.

^{4.} Thomas V, et al: Immunoglobulin levels and antibody - coated bacteria in urines from patients with urinary tract infections. Pros Soc Exper Biol Med 148: 1198, 1975.

^{5.} Reckler JM: Urinary tract infections in burned patients. USAISR Annual Research Progress Report, June 1971, pg 11.

4. After appropriate therapy, does coated or uncoated bacteriuria recur?

#### METHODS

All patients found to have yeast, by microscopic examination of the urine, will have urine set aside for immunofluorescence. Those patients coming to postmortem examination will be specifically observed for yeast infection of the kidney, ureter, bladder wall, and prostate, as well as other areas of tissue invasion. The pathologic findings will be correlated with the immunofluorescence of the organisms in the urine.

All patients with indwelling catheters will have urine screened for bacteria which, in turn, will be subjected to immunofluorescence. Clean catch-midstream urine will also be screened after the urinary bladder catheter is removed. Those patients found to have "coated" bacteriuria, as well as those with "noncoated" bacteriuria found 48 hours after removal of the urinary catheter, will be treated with the appropriate antibiotic, as determined by culture and sensitivity. Urine will be constantly screened after any therapy for urinary tract infection.

Staining of bacteria and yeast in the urine for immunoglobulin will be done by the technique described by Thomas, et al

#### RESULTS

Twenty patients have been evaluated to date. Of the 5 with bacteriuria, none were demonstrated to have immunoglobulin coating suggesting no extrabladder infection. Of the 15 with candiduria, only those with hyphae were found to have positive immunofluorescence tests. No correlation has been made between anatomic site of infection by postmortem examination and yeast immur oglobulin coating. No conclusions can be drawn until more patients are studied.

## PUBLICATIONS AND/OR PRESENTATIONS

None

RESEARCH	AND TECHNOLOGY	WORK UNIT S	UMMARY			200	- 76 10 C		ABPORT CONTROL SYMMOL			
		-	-	DA	Ur 6	388	10 10 0	1				
S DATE PREV SUM RY	4 KIND OF SUMMARY	S SUMMARY SCTY	WORK SECURITY	P REGRA	OIN G"	84 DI	sa'n instr'n	ON TRACTO	DATA-	LEVEL OF SUM		
75 07 01	H. TERM	U	Ŭ		NA	h	IL K YRS			A USER UNIT		
10 NO CODES *	PROGRAM ELEMENT	PROJECT	NUMBER	TASK AREA NUMBER WORK UNIT NUME								
& PRIMARY	61102A	3A161102	B71P		)2		069					
S. CONTRIBUTING												
C CONTRIBUTING				·····								
11 TITLE (Procode with	Security Classification Code	(U) To De	termine the	e Rol	e of	the	Prostag]	andins	(PG)	in the		
Response to	o Volume Expan	nsion in th	e DogA M	odel	of C	hange	s in In	iured T	roops	(44)		
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS									<u></u>		
003500 Clir	nical Medicine	9										
13 START DATE		14 ESTIMATED COM	PLETION DATE	IL FUNC	NNG AGE	ENCY				ANCE METHOD		
75 02		76 09		DA	DA   C.				n-House			
" CONTRACT/GRANT NOT Applicable			IS RESOURCES ESTIMATE			L & PROFESSIONAL MAN YRS & PUT			51 (jn thevearda)			
& DATES/EFFECTIVE:		EXPIRATION			PREEE	SING	-					
N. NUMBER *				PISCAL 76		.4		1 13				
		4 ANDUNT		YEAR	CORNEL		+		-			
& KIND OF AWARD		f. CUM. AMT.										
IS RESPONSIBLE DOD	DRGANIZATION			3 PERI	ORMAN	ORGANI	ATION			1		
HAME .* ITS Arms	. Institute of	E Surgical	Pegearch		110		Tuchikut			l Dogo zvel		
00 ALM	Y INSTITUTE O	Jurgicar	Research	1	03	ALMY			urgica	1 ISSEALC		
ADDRESS * Down	Com Houston			ADDREN	ме ^{н•} —	Cabol	LIC Brand	cn, Nep	nrotog	y svc		
Fort	sam Houston,	Texas /82	:34	1	FO	ort Sa	m Housto	on, Tex	as /8	234		
				PRINCIP		STIGATO	n / Pursiak MAAN I		e inclination			
				HALL & DATE & DIT MAT MC								
Basil 2	A. Pruitt. Tr	COT. MC				512	-221-4600	2007 MC				
AND DUGTT ?	5102001_070A			TELEPHONE 012-221-4098								
TELEPHONE.	512-221-2720				. BECUR	TTT ACCO	NAL HA <b>RE</b> .					
TO GENERAL USE	ASSOCIA	TE INVE	STIGATO	R5								
FOREIGN INTELLIGENCE NOT CONSIDERED				-	Ric	hard	H. Merr	111, LT	C, MC			
				NAME						DA		
ZZ KEYWORDS (Procedo	BACH WID Somethy Classifi	(U)	Animal; (U	) Ren	al;	(U) 1	Autoregul	Lation;	(U) D	iogs;		
(II) Prostar	Tanding: (II)	Volume evr	angion				-					

des a

. All and a second

Academ

and a complete the second structure of the second 
and the second second second

これのないないのであるとない いいないがい

Charles Bally word

23 TECHNICAL OBJECTIVE,* 24. APPROACH, 25. PROGRESS (Furnish Individual paragraphs Idmilliod by number Protodo last of ash midi powelly Classification Cade.)

23. (U) In the established dog model of volume expansion, responses in adaptation and autoregulation occur. Renal prostaglandins are thought to play a role in this autoregulation. The objective in this study is to observe response to volume expansion in the absence of prostaglandins as a model of changes in injured troops.

24. (U) Volume expansion in the dog model will be accomplished with 7.5% body weight expansion with Ringer's Lactate (RL). For comparison, 6.5ml/kg body weight of hyperoncotic albumin (HA) will be used to study the difference between vascular and interstitial expansion (RL) and vascular expansion alone (HA). The role of prostaglandin (PG) in a response to volume expansion will be tested by inhibiting PG synthesis with IV Indomethacin 2mg/kg body weight.

25. (U) 75 07 - 76 09 The study was designed to observe animals response to RL and HA before and after PG inhibition. The data show no effect of PG inhibition on baseline studies and in no way alter the response to volume expansion with either RL or HA.

vallable to contractore upon originator's

approve

#### TERMINATION

## PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

# REPORT TITLE: TO DETERMINE THE ROLE OF THE PROSTAGLANDINS (PG) IN THE RESPONSE TO VOLUME EXPANSION IN THE DOG -A MODEL OF CHANGES IN INJURED TROOPS

## U S ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators

David B. Olin, MD, Major, MC Richard H. Merrill, MD, Lieutenant Colonel, MC

Reports Control Symbol MEDDH-288 (RI)

UNCLASSIFIED
#### PROJECT NO. 3A161102B71R-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE- SURGICAL PATHOLOGY

#### **REPORT TITLE:** TO DETERMINE THE ROLE OF THE PROSTAGLANDINS IN THE RESPONSE TO VOLUME EXPANSION IN THE DOG - A MODEL OF CHANGES IN INJURED TROOPS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

**Period covered in this report:** 1 July 1975 - 30 September 1976

Investigators: David B. Olin, MD, Major, MC Richard H. Merrill, MD, Lieutenant Colonel, MC

**Reports Control Symbol MEDDH-288 (RI)** 

Isotonic volume expansion in man and animals results in slight changes in glomerular filtration rate (GFR) and renal blood flow, but dramatically increases urinary sodium excretion and fractional excretion of sodium (FE $_{N_2}$ ). The factors leading to this response remain unclear. Physical factors (Starling forces) may account for the response by increasing the proximal tubular backleak of sodium. The concept of a hormonal factor which may be natriuretic (III factor) arose from work of De Wardner in which cross perfused dogs receiving blood from volume expanded animals underwent natriures is despite the fact that they themselves were not volume expanded. The character and mechanisms of release of this factor remained to be defined. Work from this laboratory and others has shown prostaglandins E (PGE) to have natriuretic properties when infused into the renal artery. This effect seems to be mediated by vasodilation, as filtration decreases and proximal tubular reabsorption of sodium, glucose, and bicarbonate are decreased. The physiologic singificance of intra-arterial infusions of PGE have been challenged due to the low levels of PGE in the circulation and its site of synthesis has resulted in increased urinary sodium. To further test the role of PG in renal auto-regulation, PG synthesis was inhibited by Indomethacin. To observe the role of PGE in volume expansion Rir arts Lactate (RL) was used. Further studies were performed with hyperoncotic albumin (HA) to observe any variation in PG's role in the presence of interstitial volume contraction which results with hyperoncotic albumin.

Animal Renal Autoregulation Dogs Prostaglandins Volume expansion

#### TO DETERMINE THE ROLE OF THE PROSTAGLANDINS IN THE RESPONSE TO VOLUME EXPANSION IN THE DOG - A MODEL OF CHANGES IN INJURED TROOPS

Isotonic volume expansion in man and animals results in slight changes in glomerular filtration rate (CFR) and renal blood flow, but a dramatic increase in urinary sodium excretion and fractional excretion of sodium (FE_{NIR}). The factors leading to this response remain unclear. Physical factors (Starling Forces) may account for the response by increasing the proximal tubular backleak of sodium. The concept of a hormonal factor which may be natriuretic (III Factor) arose from work of De Wardner in which cross perfused dogs receiving blood from volume expanded animals underwent natriuresis despite the fact that they themselves were not volume expanded. The character and mechanisms of release of this factor remained to be defined. Work from this laboratory and others has shown prostaglandins E (PGE) to have natriuretic properties when infused into the renal artery (1,2,3). This effect seems to be mediated by vasodilation, as filtration fraction decreases and proximal tubular reabsorption of sodium glucose, and bicarbonate are decreased. The physiologic significance of intra-arterial infusions of PGE have been challenged due to the low levels of PG in the circulation and its site of synthesis being in the renal medulla. Infusion of arachidonic acid (immediate precursor of PGE in the kidney) into the renal artery, indirectly increasing renal PG synthesis have resulted in increased urinary sodium. To further test the role of PG in renal auto-regulation, PG synthesis was inhibited by indomethacin. To observe the role of PGE in volume expansion Ringer's lactate (RL) was used. Further studies were performed with hyperoncotic albumin (HA) to observe any variation in PG's role in the presence of interstitial volume contraction which results with hyperoncotic albumin.

#### METHODS

the second state of the second state of the second

2

Studies were performed in the Metabolic Branch, USAISR. Mongrel dogs weighing between 8 and 20 kilograms were utilized. The animals were given nothing by mouth after 6 pm of the day prior to surgery. The animals were anesthetized with sodium pentothal and respirations were maintained by a Byrd respirator. Supplemental doses of sodium pentothal were given as needed. Femoral vessels were isolated for monitoring arterial pressure, blood sampling, and intravenous infusions. Both ureters were isolated through a small suprapubic incision. In a few animals, a left flank incision

3. Gross J, Barlter F: Effects of prostaglandins  $E_1$ ,  $A_1$ , and  $F_2$  x on renal handling of salt and water. Am J Physio 225: 218, 1973.

^{1.} Martinez-Maldanado M, Tsparas N, Eknoyan G, Suki W: Renal actions of prostaglandins: Comparison with acetylcholine and volume expansion. Am J Physio 222: 1147, 1972.

**^{2.}** Johnston H, Herzog J, Lauler D: Effect of prostaglandin E, on renal hemodynamics sodium and water excretion. Am J Physio 213: 939, 1967.

was utilized to isolate the renal vein for determinations of para-aminohippurate (PAH) extraction.

all animals were given 20 to 40 ml/kg 5% Dextrose in water intravenously in order to establish a water diuresis during control periods. All urine output was replaced with 5% Dextrose in water. Loading doses of Inulin 15 mg/Kg and PAH 8 mg/Kg were given and infusion of Inulin and PAH was started. The animal was allowed 30 minutes to recover from surgery and then baseline control periods of 10 to 15 minutes each were performed.

After the control periods, the animals were treated in the following manner:

Group I - Ringer's lactate alone - 7.5% body weight volume expansion Group II - Indomethacin 2 mg/Kg body weight Group III - Hyperoncotic albumin - 6.5 ml/Kg body weight

The response was then observed, measuring clearance of Inulin, clearance of PAH, urinary sodium excretion, fractional excretion of sodium, osmolar clearance, clearance of free water, blood pressure and hematocrit. In addition to the above data, serum and urinary calcium, phosphorus, potassium, and chloride were obtained. After the response was observed, the following was done to each animal group:

Group I - Indomethacin ? mg/Kg IV Group II - Sub group A - Ringer's lactate 7.5% body weight Sub group B - Hyperoncotic albumin 6.5 ml/ Kg Group III - Indomethacin 2 mg/kg IV

Repeat data was collected for comparison with the previous data for clearance of Inulin, clearance  $C_{PAH}$ , urinary sodium excretion( $U_{Na}V$ )

fractional excretion of sodium,  $(FE_{Na})$  clearance of Osmoles,  $(C_{OSM})$  free

water clearance,  $(C_{H_2O})$ , blood pressure and hematocrit. In addition, 2

to three animals were used as controls for both Ringer's lactate volume expansion and hyperoncotic albumin expansion, and received no Indomethacin.

#### LABORATORY METHODS

Laboratory tests were performed in the Biochemistry Laboratory of the Institute of Surgical Research. All tests were performed on both serum and urine utilizing the following techniques. Sodium and potassium were done under routine techniques on the flamephotometer. Chloride determinations were performed on the Buchler-Cotlove chloridimeter utilizing standard techniques. Serum and urine values for Inulin, PAH, calcium, phosphorus, creatinine, and blood urea nitrogen were all performed on the autoanalyzer using adaptation of standard methods. These techniques were modified by the personnel of the institute of Surgical Research Laboratory for utilization in this study. Calculations of urinary sodium excretion using urine sodium concentration and volume as well as fractional excretion of sodium utilizing urinary sodium excretion and glomerular filtration rate as estimated by the clearance of Inulin were all performed utilizing standard physiologic formulas. Clearance data for osmolality and the calculations of free water clearance were made using the formula  $V = C_{osm} + C_{H_nO}$ .

All studies utilizing Indomethacin were performed 30 minutes after the intravenous dosage was administered (4).

The animals were sacrificed at the end of the study,

The data represents single observations in each animal in each condition studied. Paired T analysis was performed on these data. Mean and standard deviations of each group were determined for the purposes of comparison.

#### RESULTS

Results from animilas in Group I receiving 7.5% body weight volume expansion with RL are in Table I. Following volume expansion there were significant increases in urine flow,  $FE_{Na}$ ,  $C_{osm}$ , and  $C_{H_2O}$ . After Indomethacin

was given there were no charges in any of these parameters.

Group II animals were given 2 mg/kg Indomethacin. There was no change in any parameter (Table I). Following the Indomethacin, Group IIA received 7.5% body weight volume expansion with RL. There was significant change in urine flow, FE_{Na}, and C_{H₂O} with slight but not significant increases in C_{osm} (Table II)

The changes were similar to those of Group I. Group IIB received 65ml/kg LA volume expansion following the Indomethacin (Table III). These animals also showed no change in any parameter following Indomethacin. After HA there were significant increase in urine flow, FE_{Na}, and C_{H₂O}. These changes were similar

to Group III animals. Group III animals received 6.5ml/lg HA (Table IV). The HA resulted in significant increase in urine flow and  $C_{H_2O}$  yet no significant

increase in  $FE_{Na}$ . Indomethacin resulted in no change in any parameter other than slight decrease in  $FE_{Na}$  (p = 05).

^{4.} Vane JR: Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. Nature (New Biology) 231: 232, 1971.

TABLE I

# RINGERS LACTATE (4)

	CONTROL	EXPERIME	NTAL	EXP + IND	10
FLOW	1.1±.99	6.2±2.26	P=0.025	6.57±1.46	ŃS
GFR	52.77±15	74±31.4	SN	78.8±31.4	NS
FENa	0.7±.85	3.32±1.82	P=0.01	4.37±3.0	NS
Cosm	1.73±.73	3.38±.65	P<.025	3.95±1.22	NS
CH30	<b>-</b> .64±.73	2.78±2.0	PC.025	2.59±7.49	SN
BP ,	129±12	114±8	SN	139±26	SN
нст	(1)	32(1)			

and the dependence of the second s

1990 C. C. C.

がないないです

A STATE OF A STATE OF A STATE OF A STATE OF A STATE OF A STATE OF A STATE OF A STATE OF A STATE OF A STATE OF A

のないないないないのであるというないないないというからいという

9.82±4.21<.025 3.75±2.68 NS 55.18±25 NS 4.8±1.36<.01 5.7±2.93<.05 136±11 NS R CONTROL + INDO 2.49±1.8 1.48±.89 .92±.23 .8±1.28 49±25 135±12 FLOW FENa Cosm с_{Н2}0 GFR ВР

.

「「「「「「「」」」をいたいできたいできたというできた」というできたので、

a state and a state of the stat

## TABLE III

	CONTROL	CONTROL + INDO	HA
FLOW	1.94	2.23 NS	11.9<.001
GFR	60.77	60.51 NS	61.65 NS
FENa	.47	.99	5.58<.01
Cosm	1.8	2.0 NS	4.23 NS
0 <b>6H</b> 3	69	1.0	7.45<.05
48) 1	174	174 NS	174 NS

.

TABLE IV

# SALT POOR ALBUMIN (7)

	CONTROL	EXPERIMENTAL	EXP + IND	8
Flow	2.75±3.3	9.36±4.85 P<0.001	9.14±5.6	NS
GFR	62±11.7	70.6±21.2 NS	72.42±19	NS
FE _{Na}	1.39±1.76	2.6±2.2 NS	2.14±1.93<	:.05
Cosm	2.71±1.6	3.95±2.9 NS	3.63±3, 19	NS
с _{Н2} о	.74±2.8	6.68±3.12 №0.005 ·	7.2±2.74	NS
ВР	154±27	152±23.9 NS	161.5±23 I	NS
ĤCĨ	32.9±5.6	25.6±3.6 P<0.001		

#### COMMENTS

Volume expansion with Ringer's lactate results in both interstitial as well as intravascular volume expansion. Our data suggest that prostaglandin inhibition results in no charge from control period data and does no affect response to 7.5% body weight volume expansion with Ringer's lactate. This lack of response was observed both before volume expansion and 30 to 45 minutes after volume expansion. This observation has also been made by Kirschenbaum et al (5).

Further studies in unanesthetized animals suggest that prostaglandin inhibition leads to a natriuresis (6). This observation is contrary to previous observations in anesthetized animals and may be important. Sonnenburg and Stein have suggested that since prostaglandins are synthetized in the medulla of the kidney, they may act locally (6,7). This eliminates the need to postulate a circulating hormone like effect of the prostaglandins. The theory can be stated that interstitial volume contraction leads to an increase in prostaglandin synthesis thus, increasing collecting duct sodium reabsorption. In interstitial expansion, the opposite would occur.

During volume expansion with hyperoncotic albumin, similar degrees of intravascular volume expansion (judged by plasma volume measurements and changes in hematocrit) do not result in comparable natriuresis as with Ringer's lactate. (8,9) The lesser natriuresis is seen despite an increase in volume in urine and an increase in free water clearance suggesting similar distal delivery of sodium. Hyperoncotic albumin results in interstitial volume contraction. Thus, prostaglandin synthesis could be stimulated and distal tubule or collecting duct sodium reabsorption increased to explain the observation of a lesser degree of natriuresis. The present data show no change in fractional excretion of sodium, urinary sodium excretion, free water clearance, or osmolar clearance when prostaglandin inhibition was performed both prior to volume expansion as well as after volume expansion. These studies were performed in anesthetized animals to confirm these observations. The role of renal prostaglandin in renal autoregulation is still unclear. Further testing needs to be performed utilizing anatagonists and blockers of prostaglandin synthesis other than Indomethacin.

6. Stein J, Kirschenbaum M, Bay W, Osgood R, Fenis T: Role of collecting duct in the regulation of sodium balance. Cir Res Supp 36 & 37, 1-119, 1975.

7. Sumenburg H: Renal response to blood volume expansion: Distal tubular function and urinary excretion. Am J Physio 223: 916, 1972.

8. Levy M, Levinsky N: Proximal reabsorption and intrarenal pressure during colloid infusions in the dog. Am J Physiol 220: 415, 1971.

9. Howards SS, Davis BB, Knox FG, Wright FS, Berliner RW. Depression of fractional sodium reabsorption by the proximal tubule of the dog without sodium diuresis. J Clin Invest 47: 156

^{5.} Kirschenbaum M, Stein J: The effect of inhibition of prostaglandin synthesis on urinary sodium excretion in the conscious dog. JCI 57: 517, 1976.

#### PUBLICATIONS AND/OR PRESENTATIONS

None

¥

#### REFERENCES

1. Martinez-Maldinado M, Tsparas N, Eknoyan G, Suki W: Renal actions of prostaglandins: Comparison with acetylcholine and volume expansion. Am J Physiol 222: 1147, 1972.

2. Johnston H, Herzog J, Lauler D: Effect of prostaglandin E₁ on renal hemodynamics sodium and water excretion. Am J Phsyiol 213: 939, 1967.

3. Gross J, Barlter F: Effects of prostaglandins E, A₁, and  $F_2x$  on renal handling of salt and water. Am J Physiol 225: 218, 1973.

4. Vane J: Inhibition of prostaglandin synthesis as a mechanism of action for asprin-like drugs. Nature (New Biology) 231: 232, 1971.

5. Kirschenbaum M, Stein J: The effect of inhibition of prostaglandin synthesis on urinary sodium excretion in the conscious dog. J. Clin Invest 57: 571, 1976.

6. Stein J, Kirschenbaum M, Bay W, Osgood R, Fenis T: Role of collecting duct in the regulation of sodium balance. Cir Res supp 36 & 37, 1-119, 1975.

7. Sonnenburg H: Renal response to blood volume expansion: Distal tubular function and urinary excretion. Am J Physiol 223: 916, 1972.

8. Levy M, Levinsky N: Proximal reabsorption and intrarenal pressure during colloid infusion in the dog. Am J Physiol 220: 415, 1971.

9. Howards S, Davis B, Knox F, Wright F and Berliner R: Reabsorption by the proximal tubule of the dog without sodium diuresis. J Clin Invest 47: 1561, 1968.

				-						والمتباد والمراقع الشاعات
RESEARCH	AND TECHNOLOG	Y WORK UNIT S	JUMMARY	DA (	DF 638	7	76 10 0	1	DD-D	CONTROL STHEOL R&E(AR)636
5 DATE PREV SUMPRY 75 07 01	H. TERM	S SUMMARY SCTT	6. WORK SECURITY	7 AEGR	ADING	NL	PN 14578"N	CONTRACTOR	ACCESS	S. LEVEL OF SUM
10 NO /CODES *	PROGRAM ELEMENT	PROJECT	T NUMBER	TASK	AREA NUM	IDER		WORK UNIT	NUMBER	A
a. PRIMARY	61102A	3A16110;	2B71P	02			068			
b. CONTRIBUTING				<u> </u>						·····
C CONTRIBUTING				<u> </u>			-			
II TITLE (Procode with	Security Classification Code	"(U) To De	termine Whe	ther	the P	rost	aglandi	ns are 1	Impor	tant in
Development	t of Acute Re	nal Failur	eA Rabbit	. Mode	<u>el of</u>	Rena	<u>l Failu</u>	re in Sc	oldie	rs (44)
12 SCIENTIFIC AND TEC	CHNOLOGICAL AREAS	-								
13 START DATE	Alcal Medicin	C TIA. ESTIMATED CON	PLETION DATE	TIE PUN	DING AGEN	<del>CY</del>		16 PERFORM		THOD
75 02	I	76	09		1		1	C. In-	-Hous	e
17 CONTRACT/GRANT	Not Appl	icable	55	10 RET	OURCES ES	TIMATE	A PROPESS	IONAL MAN YRS	Th PU	H DS (In draw ands)
& DATES/EFFECTIVE	**** •*EE	EXPIRATION.		<b> </b>	PRECEDIN	18	1		+	
A NUMBER *				FISCAL	76		.4		1	0
G TYPE.		4 ANOUNT:		YEAR	CURRENT		1		1	
& KIND OF AWARD		f. CUM. AMT	•		1					
IS RESPONSIBLE DOD (	DRGANIZATION	Į		30 PER	FORMIPG OF	REANIZA	TION	T	-	
HANE." US ATTWY	Institute of	Surgical	Research	NAME *	US Arm	av In	stitute	of Sur	rical	Research
····.		• <b>,</b> ·		1	Met	abol	ic Bran	ch		
ADDRESS . Fort	Sam Houston,	Texas 782	34	ADDRES	** For	ct Sa	m Houst	on, Texa	as 7	8234
				1		-		•		
]				PRINCIP	AL INVEST	ISATOR/	(Fumioh BEAN (	il V S. Anndomic	<u>Inetineties</u>	•
RESPONSIBLE INDIVIDU	JAL			NAME	• Dav	id B	. Olin,	MAJ, MO	2	
NAME Basil	A. Pruitt, Jr	, COL, MC		TELEF	PHONE 51	.2-22	1-4698			
NAME SIL LIL LOSS   TELEPHONE 512-221-2720 BOCIAL SECURITY ACCOUNT HUMBER										
TELEPHONE. JIZ ZZI Z/ZO 21 GENERAL USE ABSOCIATE INVESTIGATORS DI DI D										
				NAME:	Ricn	ara	H. Merr	ill, Lu	С, МС	;
FOREIGN IN	TELLIGENCE NO	T CONSIDER	ED	NAME:	:					Uh
22 REYTORUS (Pressee)	FACH WIN POWER COM			Bro	-+		~~ (II)	Pahkite		
(U) ACU.e	renal lallure	-; (U) Gent		ELU:	SLAYIG		S; (U)	ROLUGI CO	-Hen Cad	
2) TECHNICAL ( BJCC)	WE, as aremunum, in	model of	Contamicin	nenh	rotori	i ci tv	eimula	ting a (	rause	» of
23. (U) 1	O estabilsh a	. MODEL OL .	Gentamituri	пери	LOLONI	Cr cr	DTHATA	lung a		; 01
renal fait	ure in injuie	a soluters	•							
24 (II) N	ion Zoeland wh	site rabbit	e will he i	+111	wid ar	in he	ven d <b>a</b> j	lv IM i	niect	ions of
24. (U) in Contamicin	ew searand with	for two was	be whother	nech	rotoxi	icity	ven de	establ:	ished	1.
Gentamitum	CO ODSELVE I	UL LWO WEE		مديط شروو	100000	.0103		,	*****	••
25 (11) 7	76 09	New Zeala	and white r	■hhit	e mair	tair	ed on r	ecular (	diets	and
25. (U) /	10 05 - 10 05	Mew Deura	lo while i	^ ຣນໂ	fered	no j	11 effe	ets and	mair	tained
receiving	40 mg/ky and 0	during	two wook st	i aur. Fudv	nerior	4. Z	in addit	ional s	+indv	comparing
Stable Ser	um electrolyc	es auring	two weer at	or NA	Cl and	ι Α agg	in foll	owed for	r two	weeks
effect or	changing arin	King water	to NHACL C	JI WA	ti and	l ayu chon:	+ Nc	.Uweu Io.	nd wh	ite
showed no	effects other	than milu	. decrease :	jer um	DICOL		te. ne	w beara		1100
rabbits to	lerate massiv	'e doses or	Gentamicii	1 neb	NEOLOY	uc.	/. 110	19, stury	y or	doute
renal fail	ure in this a	inimal is n	or practice	¥1.						
1										
1										
1										
1										
I										

and the states

olikov, nr sokoči

形は長

ALL AND ALL AND

DD FORM 1498 PREVIOUS ED TIONS OF THIS FORM THE OBSOLETE DD FORMS 1488A I NOV 88 ND 1488 - 1 MAR 68 FOR ARMY USED ARE OBSOLETE 374

#### TERMINATION

#### PROJECT NO. 3A161102B71P-02 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

#### **REPORT TITLE:** TO DETERMINE WHETHER THE PROSTAGLANDINS ARE IMPORTANT IN DEVELOPMENT OF ACUTE RENAL FAILURE - A RABBIT MODEL OF RENAL FAILURE IN SOLDIERS

#### US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

#### Investigators

David B. Olin, MD, Major, MC Richard H. Merrill, MD, Lieutenant Colonel, MC

Reports Control Symbol MEDDH-288 (RI)

UNCLASSIFIED

#### PROJECT NO. 3A16112B71R-02 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

#### REPORT TITLE: TO DETERMINE WHETHER THE PROSTAGLANDINS ARE IMPORTANT IN DEVELOPMENT OF ACUTE RENAL FAILURE -A RABBIT MODEL OF RENAL FAILURE IN SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: David B. Olin, MD, Major; MC Richard H. Merrill, MD, LTC, MC

Reports Control Symbol MEDDH-288 (RI)

The clinical observation of Gentamicin nephrotoxicity is well established. Patients receiving Gentamicin may experience transient renal insufficiency or frank renal failure. In the burn soldier, with a high risk of systemic infection, Gentamicin is a commonly used antibiotic. In an effort to further study the pathogenesis of Gentamicin nephrotoxic renal failure, an attempt was made to create an animal model. Previous studies utilizing the rat have been reported. The New Zealand white rabbit was chosen for the present investigation because of the larger size animal and the observation of high levels of circulating prostaglandins E and A (personal observation).

The observation of renal failure associated with aminoglycoside antibiotics is common, yet the mechanisms are not clear. The drug is filtered and reabsorbed in proximal and distal tubules. It is thought that renal failure results from the concentration of drugs in the tubular cell since volume depletion seems to increase reabsorption and thus predispose to renal failure. The rabbit model was proposed to evaluate the pathogenesis of renal failure associated with Gentamicin. Massive doses were utilized (10 to 20 times therapeutic dosage) to insure high dose exposure. Blood levels were measured at 4 to 8 hours and found to be in the 20 to 40 ug mg/ml range showing adequate absorption of the intramuscular injections. It is concluded that the rabbit is not a practical animal model to investigate Gentamicin induced renal failure, since it has not been possible to induce even modest renal imp⁻.rment in this species. The mechanisms of handling of Gentamicin in the rabbit kidney were not investigated, but may be of interest in future investigations.

Renal Failure Gentamicin Animal Prostaglandins Rabbits

						(minger					
RESEARCH	I AND TECHNOLOG	Y WORK UNIT S	SUMMARY	DA (	DE 6980	он ^а С	2 DATE OF SUE 76 10 0	1	REPORT O	CONTROL SYMBOL	
L DATE PREV SUMRY	4 KIND OF SUMMARY	S SUMMARY SCTY	. WORK SECURITY	7 REGR	ADING	84 DF	BB'N INSTR'N	Sh SPECIFIC	DATA -	LEVEL OF SUM	
75 07 01	H. TERM	U	U	N7	Α	N	L	I ves		A WORK UNIT	
ID NO /CODES.*	PROGRAM ELEMENT	PROJECT	TNUMBER	TASK (	ARE! NUMB	IER		WORK UNIT	T NUNDER	l	
L PRIMARY	61102A	3A16110	2B71P		02		067				
. CONTRIBUTING											
C. CONTRIBUTING	L	[					L			•	
11, TITLE (Procode wim)	Security Classification Code;	* (U) The F	Effects of	Calci	ium on	th	e Renin-	Angote	nsin S	ystem-	
Use of an A	Animal Model	of Hyperter	nsion in Mi	.litar	cy Pers	soni	nel (44)				
SCIENTIFIC AND IL	CHHOLOGICAL ANEAD										
UUSSUU CLLI	nical Medicin	C	IN STICH DATE	Tes FUN							
74.03	!	76.00	PLE HOR ONLS		i i	<b>′</b>	1	14 PERFORM	ANCE METH	HOD	
74 UL		10 09		DA			- <b> </b>	[C. In-1	House		
A DATEL/EFFECTIVE:	NOT APPIL	Cable		16. RES	DURCES ESTIN	INATE	L & PROFESSI	ONAL MAN YRS	E PUNI	Di (In thoseando)	
		EAFING HAR			76		4 .4	E.	14		
4 TYPE.		A AMOUNT		YEAR	EUNINERY-				2		
A KIND OF AWARD		f. CUM. AMT.					ļ				
IS RESPONSIBLE DOD O	DRGANIZATION	[		20 PERI	FORMING ORE	SANIZ	ATION				
TIS ATT	Thetitute of	Surgical	Percent	1	11C Arm		*				
	INSCICUCE OF	Surgicar .	Research		US ALM Motat	- 19 1 -	INSTILUL	e or su L	rgical	Research	
ADDRESS." Fort Sam Houston, Texas 78234 ADDRESS Fort Sam Houston, Texas 78234											
FOLL .	ADDARESS." Fort Sam Houston, Texas 78234 ADDARESS * Fort Sam Houston, Texas 78234										
			•	PRINCIP	AL INVESTIG	ATOR	- / Pumish SEAN II	I II E. Academic	*****		
RESPONSIBLE INDIVIDU	AL		ľ	NAME. ⁴	' Richa	ird	H. Merr	ill, LT(	C. MC		
NAME Basil A.	. Pruitt, Jr,	COL, MC	,	TELEP	HONE 51	2-:	221-5416	~,	<b>~</b> ]		
TELEPHONE. 512-	-221-2720		,	SOCIAL	. SECURITY A	ACCO	UNT NUMBER				
21 GENERAL USE				ASSOCIA		ATOR	18				
FOREIGN INT	TELLIGENCE NOT	I CONSIDERF	ED I	HAME:							
	·			NAME		_				DA	
IL KEYWORDS (Procede 1	EACH with Somethy Closelfic	ellen Code)									
(U) Calcium	m; (U) Renin;	(U) Angiot	censin; (U)	Нурє	rtensi	.on;	; (U) Do	gs			
23 TECHNICAL OBJECTI	IVE, 24 APPROACH, 28 P	PROGRESS (Pumish in	idividual paragrapha ida	millied by	number Prece		at al each with go	curity Classifics	tion Code.)		
23. (U) 10	⊃ assess we . ∽alated to hu	rortension	in militar	ຍ ອບ. 	Inurac.	.011 i	OI UNE A	reniin ai	JATOre	nsin	
System as .	related to my	per cension	In mitticur	y per +pe	soneri	 mai	-+-1 mod	-1 Vai	-lone	]	
24. (U) A	Serres or us	JS WIII NC	infueed di	ront1	expert.	,1005. (+ ~	ho ronal		510us hre	+ha	
concentration	of renin and	intensin w	ill he meas	ured.	y inco	i u roci	t cannul	ar cery	, anu f +hp	ronal	
production.	ontralatora	1 bidnov w	ill be mean	urea unit if	Ly une Fuend a	eu. Ind	will set	auron e. Tue ag i	r che	-mh1	
25 (II) 7F	$5 \wedge 7 = 76 \wedge 9$	The expert	ionco with	+ho ]	iantlv	יווט זים י	will co. nestheti:	red anir	1 CONC mal mo	del in	
investigati	ing the effect	+ of calci	n on the r	onin	-rync+1	 inoni	in evet	an does	not a		
fire conclu	ing the error	Arown Ber	cause the a	mount	of pr	an;	aratory (	Sin uous	100 u 101 u	in animal	
model is co	neiderahle.	it has been	n impossibl	a to	etabil	er. ize	- +he an	imal wit	+n u ⊧hin g	12 CILLING	
moner is of	period of tir	ne for the	evneriment	ອີເບ ຈີ່ກາ	rotocol	، مله هو بر ا	‴h≏ ຕໄດ	tillar Moriilar	filty Filty	ation	
rate and re	enal blood fle	ow have var	ried consid	erabl	v with	vin	the con	trol per	riod a	ind the	

ha this interview was a statistic to the second 
alarderser a franks

and the second second second second

ellebie to contractore upon originator's approval PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DUFFORMS 1498A AND 14981 THAR 68 FUN ARMY USEL ARE OBSOLETE NOV 68 

results obtained could easily be explained by the instability of the animal during the immediate post operative period rather than by the effects of calcium infusion. The trend of renal research is now directed toward the study of the awake intact animal. We have returned to the difficult problem of creating a suitable intact animal model.

utilizing custom designed silastic catheters and fittings, some early success has been achieved in the creation of the awake intact dog model. It is to be emphasized that the time spent on the creation of the dog model is necessary and important since it will represent an important advance in future renal studies. It is anticipated that this model will form the basis of future experimentation from this laboratory.

Using newer techniques for protecting the kidneys during prolonged surgery and

#### TERMINATION

× · ·

#### PROJECT NO. 3A161102B71P-02 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

#### REPORT TITLE: THE EFFECTS OF CALCIUM ON THE RENIN-ANGIOTENSIN SYSTEM - USE OF AN ANIMAL MODEL OF HYPERTENSION

#### US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators

Richard H. Merrill, MD, Lieutenant Colonel, MC William D. Myers, MD, Lieutenant Colonel, MC Thomas J. Lescher, MD, Major, MC

Reports Control Symbol MEDDH-288 (RI)

UNCLASSIFIED

#### PRCJECT NO. 3A161102B71R-02 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

#### **REPORT TITLE:** THE EFFECTS OF CALCIUM ON THE RENIN-ANGIOTENSIN SYSTEM - USE OF AN ANIMAL MODEL OF HYPERTENSION

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Richard H. Merrill, M.D., Lieutenant Colonel, MC William D. Myers, MD, Lieutenant Colonel, MC Thomas J. Lescher, MD, Major, MC

Reports Control Symbol MEDDH-288 (RI)

The experience with the lightly anesthetized animal model in investigating the effect of calcium on the renin angiotensin system does not allow firm conclusions to be drawn. Because the amount of preparatory surgery in this animal model is considerable, it has been impossible to stablize the animal whithin a reasonable period of time for the experimental protocol. The glomerular filtration rate and renal blood flow have varied considerable within the control period and the results obtained could easily be explained by the instability of the animal during the immediate postoperative period rather than by the effects of calcium infusion. In addition it is clear from the recent Nephrology literature that many of the assumptions classically made concerning experiments in the anesthetized dog bear little physiologic significance. The trend of renal research is now directed toward the stucy of the awake intact animal. We have returned to the difficult problem of creating a suitable intact animal model which will be suitable for sophisticated experiments involved in this protocol and others to follow. Because of the requirement of multiple vessel cannulation as well as exteriorizing ureters, other extent models will not suffice. Using newer techniques for protecting the kidneys during prolonged surgery and interruption of blood flow based on successes in the field of transplantation and utilizing custom designed silastic catheters and fittings, some early success has been achieved in the creation of an awake intact dog model. It is anticipated that a successful model will be developed within the next few months which will enable the completion of this protocol. It is to be emphasized that the time spent on the creation of the dog model is necessary and important since it will represent an important advance in future renal studies. It is anticipated that this model will form the basis of future experimentation from this laboratory.

Calcium Renin Angiotensin Hypertension Dogs

				-		OH [®] 1		MARY [®]	REPORT	CONTROL STREET
RESEARCH	AND TECHNOLOGI	WORK UNIT S	UMMARY	DA	OG 695	59	76 10 0	1	DD-D	R&E(AR)636
S DATE PREV SUMRY	4 KIND OF SUMMARY	S SUMMARY SCTY	. WORK SECURITY	7 REGR	DING	84. DH		SPECIFIC	DATA-	. LEVEL OF OW
76 10 01	H. TERM	U	U	N	A	N	L.	Ves		A. VORK UNIT
10 NO /CODES	PROGRAM ELEMENT	PROJECT	NUMBER	TASK A	REA NUME	IER		WORK UN	TNUMBE	
A PRIMARY	61102A	3A161102	2B71P		01		020			
S. CONTRIBUTING									•	
C CONTRIBUTING								<u> </u>	<u></u>	
11 TITLE (Procodo with	Security Classification Code;	(U) The	Use of Lyon	phili	zed Ve	nou	s Homogr	afts f	or Per	ripheral
AV Fistula	in Eurned Sol	diers (44)								_
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS									
003500 Clin	ical Medicine	2								
13 START DATE		14 EFTINATED COM.	LETION DATE	IS FUNC	NHG AGENC	¥		IS. PERFOR	ANCE ME	MGD
75 12		76 09		DA				C. I	n-Hou	5e
17 CONTRACT/GRANT	Not Applica	uble		18. RES	DURCES EST	INATE	A PROFESS	DHAL MAN Y		106 (In these such)
& DATES/EPPECTIVE		EXPIRATION:			PRECEDINI		1			
S. NUMBER *				PIBCAL 76 .5 14						
G TYPE		4 ANOUNT:		VEAR CURRERY						
& KIND OF AWARD.		f. CUM. AMT.								
IS RESPONSIBLE DOD	DRGANIZATION			SO. PERI	-	GANIZ	TION			
NAME." TIS Arms	Institute of	Surgical P	esearch	NAME.*	TIC L	· m 1 z	Inctitut		urgio	al Pogoard
00 ALMY	Institute of	Surgical N	escarch	i	Mot	.my .aho	lia Prov	e ur s	urgree	ai Nesearci
ADDRESS.* Fort	Sam Houston	Towar 792	34	ADDREN	e Het		am Bouet	icn fon The	Va/	79224
FOIL	ram Houscon,	Texas /02		1	FOI			.01, 19	AdS	10234
				PRINCIP	-	SATOR	(Funish SEAN ·	W S. Anndhan	ic jaolikukka	•
RESPONSIBLE INDIVIDU	AL			NAME.	Ric	har	d H. Mer	rill,	LTC,	MC
HANE Basil A.	Pruitt, Jr,	COL, MC		TELEP	HONE	512	-221-541	.6		
TELEPHONE: 5]	12-221-2720			BOCIAL	SECURITY	-				
21 GENERAL USE		·····		-	-	SATOR	•			
FOREIGN INT	TELLIGENCE NOT	CONSIDERE	D	-	Willi	Lam	D. Myers	LTC.	MC	
			-	NAME:	Thoma	as J	. Lesche	r. MAJ	MC	DA
22. KEYWORDS (Process	BACH all Security Clevell	callen Code) (U) H	omografts;	(U)	AV Fis	stul	a; (U) \	/ein gr	aft;	
(U) Hyperal	limentation;	(U) Sepsis;	(U) Therm	al in	jury;	(U)	Humans			
23. (U) Cent	TVE. 24 APPROACH, 25 tral venous ca	atheters ha	ve long be	en in	plicat	ted	in conti	ibutin	g to	systemic

sepsis in hospitalized patients. This problem is especially critical in the thermally injured soldiers since infection is the principal cause of morbidity and mortality in this patient population. As the trauma patient is usually unstable and/or requires prolonged intravenous therapy for parenteral hyperalimentation and drug therapy, and any maneuver that would decrease the duration of intravenous catheter placement might well reduce the morbidity and mortality in these patients. Usually a central catheter is not required except for pressure monitoring and therefore a convenient peripheral access site would be desirable; however, in patients with massive burns peripheral sites are scarce and often short-lived. Creation of an artificial arterial-venous shunt in an extremity might well provide a more durable access which could be used not only for drug therapy and parenteral alimentation but also for recurrent diagnostic blood sampling. 24. (U) Patients with burns ranging from 30-60% will be entered into this study. The patients will be assigned alternately to the study or control group. Those in the experimental group will have placed a peripheral lyophilized venous homograft within 48 hours of injury. A careful tabulation will be made in both groups of all complications of intravenous therapy. In the event of death all peripheral sites that have been cannulated will be inspected for evidence of infection or thrombosis, and evidence for systemic infection as a principal cause of death will be accessed by the pathologist. 25. (U) 75 12 - 76 09 To date, two burn patients requiring prolong intravenous alimentation have had lyophilized cadavar allograft veins inserted as AV fistulae. In The other graft one patient graft sepsis necessitated removal two weeks post insertion. remained patent and free of sepsis and was ligated at the time of the patients discharge Appropriate candidates are deemed to be rare even though the technic has been shown to be feasible.

Available to contractors upon originations of this form are obsolets DD forms 1488A 1 NOV 68 DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 1488A 1 NOV 68 AND 1488 1 1 MAR 68 (FOR ARMY USE) ARE OBSOLETE 330

#### TERMINATION

### PROJECT NO. 3A161102B71P-01 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

#### REPORT TITLE: THE USE OF LYOPHILIZED VENOUS HOMOGRAFTS FOR PERIPHERAL AV FISTULA IN BURNED SOLDIERS

10.14

#### US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

#### Investigators

Richard H. Merrill, MD, Lieutenant Colonel, MC Hugh D. Peterson, DDS, MD, Colonel, MC William D. Myers, MD, Lieutenant Colonel, MC Thomas J. Lescher, MD, Major, MC

Reports Control Symbol MEDDH-288(RI)

UNCLASSIFIED

#### PROJECT NO. 3161102B71P-01 BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

#### REPORT TITLE: THE USE OF LYOPHILIZED VENOUS HOMOGRAFTS FOR PERIPHERAL AV FISTULA IN BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Peciod covered in this report: 1 July 1975 - 30 September 1976

Investigators: Richard H. Merrill, MD, Lieutenant Colonel, MC Hugh D. Peterson, D.D.S., MD, Colonel, MC William D. Myers, MD, Lieutenant Colonel, MC Thomas J. Lescher, MD, Major, MC

#### **Reports Control Symbol MEDDH-288 (RI)**

Central vences catheters have long been implicated in contributing to systemic sepsis in hospitalized patients. This problem is especially critical in the thermally injured patient since infection is the principal cause of morbidity and mortality in this patient population. It was for this reason that this investigation was initiated.

To date two burn patients requiring prolonged intravenous alimentation have had lyophilized cadaver allograft veins inserted as A-V fistula. In one patient graft sepsis necessitated removal 2 weeks post insertion. The other graft remained patent and free of sepsis and was ligated at the time of the patients discharge. Appropriate candidates are deemed to be rare even though the technic has been shown to be feasible.

In the past eight months that the protocol has been in effect no further patients have been entered. It is the judgement of the primary physicians caring for the patients that suitable patients for this protocol will be few and therefore the study cannot be successfully concluded. As the input of patients into this study has been negligible, it is the feeling of the principal investigator that this study cannot be concluded within any reasonable time span utilizing this patient population, so the protocol is therefore terminated with no patients studied except for the two mentioned above.

Homografts AV fistula Vein graft Hyperalimentation Sepsis Thermal Injury Humans

							D. 78 0.0 000			
RESEARCH	AND TECHNOLOGY	WORK UNIT S	UMMARY	ה אין	CT ACCESSION		76 10 0	анант  } ] 	REPORT DD-DI	CONTROL SYMBOL RAE(AR)636
L DATE PREV SUMAY	A. KIND OF SUMAARY	S SUMMARY SCTV	A VORK STOUNTY	DA C		045.0	TH INSTA'N	AD SEFCIFIC D	ATA - 1	LEVEL OF NUM
76 10 01	H. TERM	U	U	NA		NL		YES	ACCESS	A. WORK UNIT
10 NO /CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TASK A	REA NUMBE	•		WORK UNIT	NUMBER	2
a PRIMARY	61102A	3A161102	B71P	0	2	_	070			
S. CONTRIBUTING				<b> </b>		_				
C. CONTRIBUTING				ĺ						
(U) Determi	ination of Ar	erial and	Tissue Oxy	gen I	ension	in	Normal	Man to	Deve	lop a
Technic to	Measure Tiss	ie Oxygen 1	ension in	Burne	d Soldi	ers	3 (44)			
003500 C111	nical Medicine	A ESTIMATED COM	LETION DATE		NNG AGENCY			IS. PERFORMA	HCE MET	HOD
76 01		76 09	•	DA	1	1	l	c. 1	n-Ho	use
17 CONTRACT/BRANT	Not Applica	ole		10. REA	DURCES ESTIM	ATE	& PROFESS	ONAL MAN YES	5 FUI	(D6 (In these as de)
& DATES/EFFECTIVE:	**	EXPIRATION:					1		1	
				FISCAL	76		{ .	5	1	9
G TYPE.		& ANOUNT:		YEAR	CUMBERY					
& KIND OF AWARD		f. CUM. AMT.					L			
TO RESPONSIBLE DOD	DREANIEATION			30. PERI	ORMING OREA	NIZA	TION			
HANE!" US Army	Institute of	Surgical I	Research	HANE!T	IS Army	Ins	stitute	of Surg	ical	Research
Burn Study Branch ADDRESS:" Fort Sam Houston, Texas 78234 ADDRESS:" Fort Sam Houston, Texas 78234										
ADDRESS." Fort Sam Houston, Texas 78234										
Fort Sam Houston, Texas 78234										
				PRINCIP	AL INVELTIGA	10R (	T C T	I U S. Anodomic .	ine likeliee M.C	U
RESPONSIBLE INDIVIDU	JAL Domuški Ta	007 MG		HANE:	CLAME 51	:nt   )_	221~444	ae, cri,	MC	
HANE BASIL	A. (Pruitt, Jr 12-221-2720	, COL, MC		BOCIAL	SWCURITY AC	. Z-	221-444 	0		
TELEPHONE: D	12-221-2720			AMOCIA		TORS				
FORETCH TH	TELLICENCE NO	T CONSTDER	FD	NANES	Garv	w.	Welch.	LTC. MC	2	
FORETON TH	TELETOPHCE NO			NAME:	Peter	с. А	. Petro	ff, MD	-	DA
EL REVIORDS (Procedo	BACH with Longity Class IN	cailan Cada)	· · · · · · · · · · · · · · · · · · ·					<u>-</u>		
(U) Oxygen	; (U) Tension	; (U) Arte:	rial tissue	; U	Elect	cod	e; (U)	Normal n	han	
23 TECHNICAL OBJECT	IVE, . 24 APPROACH, 25.	PROGRESS (Furnish )	ndividual paragraphs id	untilled by	number prese	la te ui	i of each with a	ecurity Classifics	tilen Cedi	
23. (U) T	o determine i	f a predic	table relat	ionsl	nip exis	sts	betwee	n arteri	lai p	02 and
tissue oxy	gen tension a	s mersured	by an indw	elli	nd oxAde	en .	electro	de. It	such	a
relationsh	ip exists in	normal man	then tissu	le oxy	ygen tei	nsı	on may	be a use	erut	measure
of periphe	ral perfusion	in injure	d soldiers.							
			a	momb	wa af	-1.0	nhucia	ian eta	FF AF	the TSP
24. (U) P	articipants 1	n this stu	obtained	anento An	ST2 OT 1	ale	ctrode	will he		ed in the
Detailed 1	niormed conse	the muscle	ticene of	070	arm and	an	indwel	ling art	teria	l line
subcuraneo	aced in the h	rachial or	radial art	erv	of the	oth	er arm	Usina	the	Ohio
will be pi	aceu III ule L annaratus Fl	02  of  30 +	o 100% will	be	adminis	ter	ed. Ai	terial	502 a	nd tissue
no2 will h	apparatus fi	isly determ	ined at eac	h le	vel of	F10	2.		•	
POZ WIII D	e shiurcaleot	orl accerm					_ /			
1										

25. (U) 76 01 - 76 09 Five normal adults have been studied. In each subject there is a linear relationship between arterial p02 and tissue oxygen tension. The slope of this line varies between subjects. There thus appears to be a range of normal. The shape of the curve suggests a relationship to age, physical conditioning and smoking history.

Available to contractors upon originator's approval

the second second second second

ž

DD HAR 61498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 1488A 1 HOV 65 AND 1488 1 1 MAR 68 (FOR ARMY USE) ARE OBSOLETE 333

#### TERMINATION REPORT

PROJECT NO. 3A161102B71P-Q2, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE-SURGICAL PATHOLOGY

REPORT TITLE: DETERMINATION OF ARTERIAL AND TISSUE OXYGEN TENSION IN NORMAL MAN TO DEVELOP A TECHNIC TO MEASURE TISSUE OXYGEN TENSION IN BURNED SOLDIERS

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 1 July 1975 - 30 September 1976

Investigators:

Clement L. Slade, MD, Captain, MC Gary W. Welch, MD, Lieutenant Colonel, MC Peter A. Petroff, MD, Major, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### PROJECT NO. 3A161102B71P+02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE-SURGICAL PATHOLOGY

#### REPORT TITLE: DETERMINATION OF ARTERIAL AND TISSUE OXYGEN TENSION IN NORMAL MAN TO DEVELOP A TECHNIC TO MEASURE TISSUE OXYGEN TENSION IN BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Clement L. Slade, MD, Captain, MC Gary W. Welch, MD, Lieutenant Colonel, MC Peter A. Petroff, MD, Major, MC

Reports Control Symbol MEDDH-288(R1)

Banadista the midden and see a subscription to be a disput to remain set

It is now possible to measure tissue oxygen levels with an indwelling oxygen electrode which gives an instantaneous readout of tissue oxygen tension. It is likely that tissue oxygen tension is a function of blood flow in the area that is being evaluated. If such a relationship exists, tissue oxygen tension can be used as an index of tissue perfusion during resuscitation of thermally injured soldiers.

Tissue oxygen levels vary with blood oxygen levels  $(p0_2)$ . Since  $p0_2$  varies from patient to patient within a predictable range, it is important to know if a predictable, linear relationship exists between  $p0_2$  and tissue oxygen. The useful data point in any study of peripheral perfusion would thus be the difference between  $p0_2$ and tissue oxygen levels with a larger disparity expected when the area under investigation is receiving less blood flow.

The purpose of this study is to determine if a predictable relationship exists between  $pO_2$  d tissue oxygen as measured by an indwelling oxygen electrode. Six normal human subjects were studied with a narrow gauge oxygen electrode (1.2 French, 0.16 inch) placed percutaneously in the rectus femoris muscle. The electrode is attached to a light-weight meter (International Bio-Physics Cooperation). Simultaneous tissue oxygen and blood  $pO_2$ measurements were made while inspired oxygen was varied between portion to a s

room air and 100%. There was a wide range of  $pO_2$  and muscle  $O_2$  at high levels of inspired  $O_2$ . At  $pO_2$  of 100 to 200, a plot of  $pO_2$  against muscle  $O_2$  was linear and there was less variation from subject to subject. This is the range of  $pO_2$  at which most trauma subjects will be studied. This system may thus prove to be a useful tool in the study of resuscitation of thermally injured soldiers.

Normal man Electrode Oxygen Tension Arterial tension

				-	NET ACCE					
RESEARCH	AND TECHNOLOG	T WORK UNIT S	UNHARY	DA	OG 690	52	76 10	01	DD-D	R& B(AR)436
& BATE PREV SUMPRY	. KIND OF SUMMARY	S. SUMMARY SCTY	. WYOR SECURITY		A	7			ATA	
76 10 01	H. TERM	U	U		NA	N	L			A NORE UNIT
NO./CODES:"	PROGRAM ELEMENT	PROJECT	HUMBER	TANK					-	
-	61102A	3A161102	B71P		01		022			
-										20 Sec. 19
C. CONTRIBUTING									ais s	
TIL TITLE (Proceds with	Security Classification Code	" (U) Serum	Pepsinoge	n Le	vels i	in th	e Therm	ally Inj	ured	Soldier:
A Possible	Predictor of	Gastroduod	lenal Disea	<b>BE</b> (	44)					
		_								
003500 C11	nical Medicin			110 700						
26.00		76.00			1	•••	•			
T CONTRACT/SRANT	Not Inslig	10 09.		DA				<u>C. I</u>	-HOU	ie
& DATEMEFFECTIVE	NOT APPITC				The state of			MAL HAN YAS	1	ti (to theready
A HUMBER:					1 76		.4		1 1	B
-		L ANDUNT:		VEAR	-				÷	\$
. KIND OF AVAND:		I. CIMA ANAY			}		}		1	
	MGANIEATION			A. PER	CILLOS A	115 ) r			<u> </u>	
		Const 1 *								
US AIMY	TUSTICUTE OI	Surgical I	vəsçal CD	la ant	US AT	my I	nstitut	e oi Sui	gica	nesearch
				l	Burn	i Stu	dy Bran	ch		
rort	sam Houston, "	rexas 7823	94		- Fort	: San	Housto	n, Texas	78	234
					AL MAREN	168768 (   17	Puilles <b>1949</b> H	W.L. Anothenis J A.R. M.R. T	NACT	
HESPONSIOLE HIDIVISO	n. A. Praitt. Jr	COL. MC			DEVIC	E17	1889aru	sn, mau,	ПÇ	
512	-221-2720			VILLO		27.4-	<u> </u>	3		
TELEPHONEL JLC	-a. 6/6V			THEIAL		ACCOM				
FORRICH THE	PTITCENCE NO		'n	ABOCIA	78 HIVEST	647668	**			
FOREIGN IN.	TELLIGENCE NO.	I CONSIDERE	L L		Dr. 3	amito	LI			
	ACT with Security Closelles	ellen Code) (***) **	(11)	HADE:			/** \ ****			DA.
(III) Dombio		(U) H	umans; (U)	Stre	ess ul	ceri	(U) Hei	norrhægi	c gai	tritis;
IL TECHNICAL OBJECTI	ULCEI GISCAS		Burn Inju	IY	augular Die					
22 (m) m	. Johowału o di						4 a b 3 a			
23. (U) It	o decermine i	serum pep	sinogen is	Vetz		IGT	repre b	edictor	ot	
. gastroquode	enal disease	corrowing p	urn injury	in a	201016					
0.4 /m\ m						• • • •				
24. (U) TI	ne method invo	DIV90 18 58	mpling ser		c regu	lar	interva.	LS, and	meas.	iring
the pepsind	gen levels to	o determine	trends an	a coi	LIGIGI	e th	18 W1CD	CIJUICS	T GAN	dence or
gastroduode	enal disease	in the burn	patient.							
25. (U) 7	76 03 - 76 09	Eleven pa	tients hav	e bei	en stu	died	in pile	ot proje	ct.	
Although no	o definitive of	conclusion	can be rea	ched	with	such	small :	numbers,	twg	patients
clearly sho	wed elevation	n of the se	rum pepsin	ogen	as an	alyz	ed by ra	ngiozman	noasi	may prior
to gastroir	ntestinal hem	orrhage. I	he second	phase	e of t	his	study w	ill be t	0 001	relate
the serum p	pepsinogen lev	vels with e	ndoscopic	findi	ings.					
										1
										1
										ſ
										[
										1
										1
										1
										1
										I
Available to contractore	une etimeter's envot	a i		i nga mga ma						
DD, <b>****</b> 1498	PREVIOUS E	DITIONS OF THIS	RMY USEI ARE	BOLETI Desoli	K. DD 70 Kte		496 A. 1 NOV			
			33	37						

#### TERMINATION REPORT

PROJECT NO. 3A161102B71P-01, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

**REPORT TITLE:** SERUM PEPS (NOGEN LEVELS IN THE THERMALLY INJURED SOLDIER: A POSSIBLE PREDICTOR OF GASTRODUODENAL DISEASE

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

David K. Teegarden, M.D., Major, MC l. M. Samloff, M.D. (Professor of Medicine, UCLA)

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

PROJECT NO. 3A161102B71P-01, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

REPORT TITLE: SERUM PEPSINOGEN LEVELS IN THE THERMALLY INJURED SOLDIER: A POSSIBLE PREDICTOR OF GASTRODUODENAL DISEASE

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: David K. Teegarden, M.D., Major, MC I. M. Samloff, M.D. (Professor of Medicine, UCLA)

Reports Control Symbol MEDDH-288(R1)

Little attention has been directed to pepsin as an etiologic factor in ulcerative disease of the stomach and duodenum following thermal injury. Pepsin is a proteolytic enzyme that is formed from the cleavage of its precursor, pepsinogen. Its cells of origin are the chief cell and mucous neck cell in the stomach. Pepsinogen is a proteolytic enzyme which attacks soluble native proteins and is therefore an endopeptidase. Previous studies in duodenal ulcer disease and other diseases of the intestine have shown pepsinogen to increase prior to a gastrointestinal hemorrhage.

This study proposes to analyze serum pepsinogen levels and evaluate as a predictor of gastrointestinal erosion and ulceration following thermal injury.

Thermal injury Burns Stress ulceration Hemorrhagic gastritis Pepsinogen

#### SERUM PEPSINOGEN LEVELS IN THE THERMALLY INJURED PATIENT: A POSSIBLE PREDICTION OF GASTRODUODENAL DISEASE

Little attention has been given pepsin as an etiologic factor in ulcerative disease of the stomach and duodenum. Pepsin, a proteolytic enzyme, attacks soluble native proteins within the molecule and is therefore an endopeptidase. It is present in its cells of origin, the chief cell and mucous neck cells, as a zymogen precursor called pepsinogen. Other unidentified cells also contain pepsinogens. A leakage of digestive enzymes into the blood stream is a common occurrence and can be used, for instance, in detecting diseases of the pancreas, utilizing the serum lipase and amylase. How pepsinogens gain access to the circulation is unknown; serum levels have been shown to reflect the secretory activity and morphologic status of mucosa and have been helpful in the diagnosis of pernicious anemia, duodenal ulcer, and gastric cancer.

Pepsinogens are separated by immunochemical methods into two groups: PGI and PGII. The two groups differ in mucosal distribution and cellular origins. Both groups are found in the fundic mucosa.

The purpose of this study is to analyze the predictive value of the stomach enzyme precursor, pepsinogen, as an indicator of gastrointestinal hemorrhage resulting from stress erosions and ulceration following thermal injury.

#### Results

Eleven patients with a greater than 30% total body surface burn were included in a pilot study. The serum pepsinogen levels were measured daily for seven days, then every other day for the next seven days, and, finally, at weekly intervals for the next six weeks. The samples were submitted to Dr. Samloff's laboratory for analysis, utilizing a competitive binding double antibody radioimmunoassay,

Of the 11 patients in the pilot study, two patients demonstrated abnormally elevated serum pepsinogen levels, one of which was associated with a gastrointestinal hemorrhage. The remaining patients demonstrated essentially normal levels.

No specific conclusions can be drawn as to the reliability of the serum pepsinogen in reflecting morphology of the gastric mucosa; however studies correlating the serum pepsinogen level and endoscopic appearance of the stomach and duodenum will be performed in the future.

#### PRESENTATIONS AND PUBLICATIONS

None.

*****			-					*******		
RESEARCH	I AND TECHNOLOG	Y WORK UNIT S	JUMMARY	DA	ACT ACCESSION 6957	٦	# DATE OF 000	01	DD-D	PAE/AR1636
S DATE PREV SUMAY	T4. KIND OF SUMMARY	TA. SUMMARY SCTY	. WORK SECURITY	D. 8661				THE SPECIFIC	DATA-	LEWEL OF SUM
76 10 01	H. TERM	υ	U	N	A	NI		CONTRACTOR		A VORK UNIT
10 NO./CODES:*	PROGRAM ELEMENT	PROJECT	T NUMBER	TABK	AREA NUMB	an J	·	WORK UNIT	T NUMBER	1
- PRIMARY	61102A	3A161102	2B71P		02	J	067			
b. CONTRIBUTING						J	Report to the second se			
C. CONTRIBUTING								<u>ilin, 245</u>	<u></u>	
TILLE (Protos	Becauty Classification (	/ (U) Alter	ations in	the r	Sacter1		Flora of	f the St	omach	and
	1 INCOSCINE :	following a	hermal inj	ury 1	in a ML	lit	ary Pop	ulation	(44)	
003500 CL	inical Medici									
13 START DATE	IIIGI NULL	ILE THATED COM	PLETION DATE	TIS PUP		<del>,                                     </del>		Tie PERPORN		
76 03	,	76 09		DA	1		i	C In-	Units	-
17. CONTRACT/GRANT	Not Appli	cable		10. RE/	HOURCES ESTY			MAL MAN YR	The	:
& DATES/EFFECTIVE:		EXPIRATION:	1	<b> </b>	76		+		+	
-			1	FISCAL	<u>. <u>7</u>°</u>		: <u>i</u>			7 1
C TYPE:		4 ANDUKT:	1	YEAR	COMMENT.		1		1	
& KIND OF AWARD:		f. CUM. AMT.	, 		<u> </u>		1			
IS RESPONSIBLE DUD	REAN EATION			30. PER	FORMINE ORE	ANITA	ATION			
HANG!" US ATTRY	/ Institute of	f Surgical	Research	a ven E 18	US Army	3' I'	nstitute	a of Sur	gica]	Research
 			,	1	Burn	St	udy Brar,	n <b>ch</b>		
ADDRESS:" Fort Sam Houston, Texas 78234 Fort Sam Houston, Texas 78234										
1			,						-	
	•••		,	-	AL HIVESTIC.	хтен, К.	(Pumin 2017	188. Anomena ; Aan. MAJ	MC	1
HANNE Basil F	A. Pruitt, Jr	COL. MC	,	TELE	- Davies	12-	-221-294	2 2	1 110	
TALEPHONE: 5]	2-221-2720	/ ••••, ···	,	BOCIA	L SECURITY A			,		
21. OENERAL USE	·····	<del></del>			ATE INVESTIGE	ATON	A			
FOREIGN IN	ITELLIGENCE NO	OT CONSIDER	ED /	HAME	Willir	am	S. McDo	ugal, MA	J, MC	1
1			······································	-	Trumar	n M	. Sasak	I, MAJ,	MC	DA
21. X EV 45 A 65 (Pressee 1	JACH all Jonathy Classifie	collan Codo)		in the second se				. <u></u>		
(U) Bacter	ial flora; (l'	J) Intestin	al flora;	(U) E	small be	<u>we</u>	<u>1; (U) F</u>	lumans		
23. TECHNICAL OBJECTI	VE, SA APPROACH, 28.	PROGRESS (Purnish In	idividual paragrapho iou	milliod by	maker preser	<b>de</b> 1621	I of each with po	ourly Classifics	illen Code (	>
23. (U) T	o determine t	che nature	of the bact	teria	al flora	<b>a</b> o'	f the st	comach a	.nd sm	all
intestine	following bur	in injury 1	.n soldiers.	•						
	- 1 intact	tenha v	-11 ha ing		* +++*AU(	~~ <b>b</b> _		ha		-" ~~~~~lo
24. (0) m	. long incest.	inal twe w	lli De ince - and emall	STLEG	/ TRIUUy	Jn ⊣ ∼nh	the nuse	3 and Day for cult	Cterr	al sampre
WIII	Calling Lim	ine summer.	, ally consta	DOWE	/L GIN -	. بيناد	Milley -	OI GULL	ure -	eculizane.
25. (U) 7	⁷⁶ 03 - 76 09	Hypomoti]	ity of the	inte	estinal	tr	act fol	lowing t	·herm?	iniury
has limite	ad the technic	que of gast	rointestin	al ir	atubatic	on.	Until	this te	chnic	nie has
been furth	her improved,	cultures c	of the smal	1 bov	wel cont	ten	t cannot	be obt	ained	with
reliabilit	.y.			-	•-	/~ .	• • • • • •	, <b>-</b>	<b>RL</b>	
l i	-									
1										

Sector and the sector of the s

ないためのないないない

and the second rest of the second 
Sale of the second

#### TERMINATION REPORT

#### PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

#### REPORT TITLE: ALTERATIONS IN THE BACTERIAL FLORA OF THE STOMACH AND UPPER SMALL INTESTINE FOLLOWING THERMAL INJURY IN A MILITARY POPULATION

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Maria and Andrew States (Mary States)

a hite the Kerst

David K. Teegarden, M.D., Major, MC Wiiliam S. McDougal, M.D., Major, MC Truman M. Sasaki, M.D., Major, MC Barry L. Davison, M.S., Captain, MSC Robert B. Lindberg, Ph.D.

Reports Control Symbol MEDDH-288(R1)

Unclassified

#### PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE, SURGICAL PATHOLOGY

REPORT TITLE: ALTERATIONS IN THE BACTERIAL FLORA OF THE STOMACH AND UPPER SMALL INTESTINE FOLLOWING THERMAL INJURY IN A MILITARY POPULATION

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: David K. Teegarden, M.D., Major, MC William S. McDougal, M.D., Major, MC Truman M. Sasaki, M.D., Major, MC Barry L. Davison, M.S., Captain, MSC Robert B. Lindberg, Ph.D.

Reports Control Symbol MEDDH-288(R1)

The role of the bacterial flora of the small intestine in infection of the thermally injured soldier has previously been undefined. Bacteria normally coexist in a relationship which is mutually beneficial to both the bacteria and the human. Under various conditions, these bacteria may proliferate with metabolic and infectious sequelae. Experimental studies have also shown that bacteria in the stomach may predispose to the development of stress ulceration.

This study proposes to investigate the bacterial flora of the stomach and small intestine in the thermally injured soldier.

Thermal injury Infection Stress ulceration Bacterial flora

#### ALTERATIONS IN THE BACTERIAL FLORA OF THE STOMACH AND UPPER SMALL (NTESTINE FOLLOWING THERMAL INJURY IN A MILITARY POPULATION

Bacteria are normal inhabitants in certain segments of the human intestine. They normally coexist in a relationship mutually beneficial to both organisms. Under normal situations, the stomach and upper small bowel are relatively sterile compared to the distal small bowel and colon, where luxuriant growth of bacteria is typical. Alterations in this balance between intestinal bacteria and man may result in adverse metabolic, blochemical, and infectious sequelae. It is well known that luxuriant bacterial growth in the small bowel may result in malabsorption, presumably by a deconjugation of bile acids. rendering intestinal juice with a bile acid concentration below the critical micellar concentration. This occurs primarily under conditions of impaired motility and stagnation of intraluminal contents from various causes. Intestinal organisms may also produce substances that either favor or inhibit the growth of other organisms. Transfer of antibiotic resistance from one organism to another is promoted by proliferation of various microbial strains. Bacteria also occupy a pivotal role in absorption of folate and Vitamin B-12.

The role of intestinal bacterial overgrowth in the pathogenesis of sepsis has not been established. Endotoxin of intestinal origin has been shown to be etiologically important in early deaths from large burns in animal studies. It can be postulated that hypoperfusion of splanchnic organs presumably occurs in the immediate postburn period. This might allow proliferation of organisms on an ischemic mucosa and possibly a port of entry into the circulation.

Following injury, patients with a total body surface burn of greater than 30% are being included in the study. A polyvinyl intestinal tube with sampling ports is employed to obtain samples from the stomach and upper small bowel near the ligament of Treitz. Samples of intestinal juice are collected within 72 hours of thermal injury and submitted for anaerobic and aerobic cultures.

#### Results

To date, luxuriant bacterial growth following thermal injury has not been established. A predominant organism has not appeared. A variety of both gram negative and gram positive organisms, as well as yeast, have been recovered. Although the colony concentrations are small, <u>Klebslella</u> has been the predominant gram negative organism recovered. Insufficient data has been collected to reach conclusions concerning the role of the bacterial flore of the stomach and small intestine following thermal injury:

PUBLICATIONS AND/OR PRESENTATIONS

None.

ANY MAN

\$

				1 JENCI	ACCESSION	2. DA	TE OF SUM	MARY	REPORT	CONTROL STREOL
RESEARCH	AND TECHNOLOGI	WORK UNIT S	JUMMARY	DA O	6956		76 10	01	DD-D	R&E(AR)636
. DATE PREV SUMRY	4 KIND OF SUMMARY	S SUMMARY SCTY	. WORK SECURITY	7. REGRAD	HO ^B 04. C	D16 8" N	INSTR'N	SA SPECIFIC	DATA	P. LEVEL OF SUM
76 10 01	H. TERM	U	<u> </u>	NA		NL		Ves	0.0	A. WORK UNIT
10 NO /CODES.	PROGRAM ELEMENT	PROJECT	NUMBER	TASK AR	EA NUMBER			WORK UNI	T NUMBE	N
& PRIMARY	61102A	3A16110	02B71P	0	2	T.	068			
S. CONTRIBUTING		ļ				1		N 49 A.		
C. CONTRIBUTING		<u> </u>		l		<u>سبا</u>		····		- 18 <u>1</u> 19 1
11 TITLE (Procode with	Security Classification Code;	) [*] (U) Evalu	uation of A.	ltered	Metabo	lis	m of I	Phenyla	lanin	e and
Tyrosine 1	n Buined Sold:	iers (44),								
	CHNOLOGICAL AREAS									
003590 CT11	nical Medicine	3 114. ESTINATED CON	PLETION DATE	TIL FUNDI	A AGENCY			14. PERFOR	ANCE ME	
75.1		76.00			1	I	ĺ			
13 LZ	Not Applia	10 09		DA		-+-		<u> </u>	In-Ho	use
A DATES/CPPECTIVE:	MOL WHATTO	ADLE			HECEDINS	** +*		DNAL MAN TE		106 (74 (74 (74 (74 (74 (74 (74 (74 (74 (74
h sumara;*				FISCAL	76 7m		.9			32
G TYPE:		4 ABOUNTS		VEAR T	UNNEWY	-+-	<u></u>			
& KIND OF AWARD:		L.CUM. AMT		1 1					}	
IS RESPONSIBLE DOD	ORGANIZATION	1	·	IN PERFO	MING OREAN	IZATIO	<b>300</b>			-r
	- Tratituta a	f Currai anl						C. 0		
The Contraction	A TURCICUCE O	I SULGICAL	Research	1	Army I	Inst	ltute	OI SUI	gicar	Research
ADDRESS:* Fort	Cam Houston	<b>Maxaa</b> 70	<b>~~</b> <i>A</i>	ADDRESS	Burn a	Stua	ly Bran	ncn		~~~ 4
FULL	Sam Houston,	Texas /o	235		ΓΟΤΤ 2	sam	HOUSTO	on, Tex	as /	8234
PRINCIPAL INVESTIGATOR (PURIOR BAR II & S. Andreic Invitation) RESERVICIONAL HARES DAVID N. HEYDOOD, CPT. MC										
ARSPONSIBLE INDIVIDUAL NAME: David N. Herndon, CPT, MC										
RESPONSIBLE INDIVIDUAL NAME: David N. Herndon, CPT, MC NAME: Basil A. P. LIT. COL. MC TELEPHONE: 512-221-4440										
TELEPHONE. 5	12-221 2720	,,		SOCIAL	ECURITY ACC	COUNT	NUMBER:			
21. SENERAL USE				AMOCIAT	E INVESTIGAT	085				
FOREIGN IN	TELLIGENCE NO	T CONSIDER	ED	NAMEI	Willian	n S.	McDou	ugal, M	iaj, m	C
				HANE	Douglas	в W.	Wilma	ore, MD	)	DA
TT. KEVBORDS (Presede	BACH with Security Classif	tellen Code) (U)	Thyroid; (	U) Phe	enylalar	nine	e; (U)	Tyrosi	.ne;	
(U) Hyperm	etabolism; (U	) Burns; (	U) Catechol	amines	;; (U) 1	Huma	ans			
TECHNICAL OBJECT	TIVE, " IL APPROACH, IL	PROGRESS (Furnish	Individual paragraphs id	lantified by n	unber procede	iont of	aanh with Si	ourly Classi	teation Cod	v .
23. (U) I	t is proposed	to compar	e the kinet	ics of	disapp	pear	ance o	of oral	. phen	ylalanine
and tyrosi	ne loads in t	hermally i	njured sept	ic and	l non-se	epti	.c sold	diers,	versu	s normal.
controls.	· • • .			-	·		- •			• . •
24. (U) N	ormal volunte	ers and se	ptic and no	n-sept	ic the	rmal	ly in	jurea p	atien	ts with
burns >403	of their tot	al body su	rface (who	have	stabils	shed	l norma	al GI I	uncti	on), will
be asked t	o give inform	ed consent	and on suc	cessi	/e days	art	er 6-1	hour Ia	ists p	ю 
administer	ed phenylalan	ine (.1g/k	g/body wt)	and ty	rosine	.05	s g∕kg	boay w	/eignt	. Venous
blood samp	les will Le a	rawn at ti	me U for Da	sal co	ncentra	atic	ons or	serum	pheny	lalanine,
tyrosine,	glucose, and	liver runc	tion tests.	AIte	er the c	orai	LOad	or ami	no ac	ids the
D1000 W111	be drawn at .	hourly int	ervais for	6 now	s and a	anaı	Lyzea :	tor typ	OSine	, pneny-
lalanine a	nd glucose.	The amino	acid levers	. WIII	De meas	sure	ed in o	dupiica	te by	specific
flurometri	c tecnnics an	d on the a	mino acid a	nalyse	er. Rai	tec		appeara	ince c	or the 3
measured s	ubstances wil	T be compa	red between	norma	al, buri	nea	septio	c and r	burneo	i
non-septic	patients.	The secol all a							1	1 4 1
25. (U) /	5 12 - 70 Uy	Pnenylaia	nine and ty	LOSTU	admin.	1Ste 7 ma	er u a:	s apove		.1 Thermail
Injured pa	ionts fasting	urn size J	on range 40	-03% A	markod'		roato	mean c r than	norma	

in strate and the state of the

injured patients mean burn size 56% range 40-85% and to 7 normals mean age 28. In burned patients fasting phenylalanine levels were markedly greater than normals 1.77=B 1.00=Norm, but rates of disappearance of phenylalanine and appearance of tyrosine arter phenylalanine load were the same in normals as in burns. One hour glucose levels were however increased greater in burns. Aglucose burns =  $22\Delta$  glucose norm = 4. These data indicate that in burns peripheral breakdown of phenylalanine and its flow to glucose is greater than normal. Burns tyrosine tolerance tests were markedly lower than normals. They were comparable to tyrosine tolerance tests of myxedematous patients.

DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORME 1456A I NOY 68 AND 1406 I I MAR 65 (FOR ARMY USE) ARE OBSOLETE 346

#### TERMINATION

#### PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

#### REPORT TITLE: EVALUATION OF ALTERED METABOLISM OF PHENYLALANINE AND TYROSINE IN BURNED PATIENTS

#### US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 December 1975 - 30 September 1976

Investigators:

David N. Herndon, MD, Cpt, MC Douglas W. Wilmore, MD Arthur D. Mason, Jr., MD John Dubois, B.S. Basil A.Pruitt, Jr, MD, Col, MC

Reports Control Symbol MEDDH-288(R1)

Unclassified

### PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE

REPORT TITLE: EVALUATION OF ALTERED METABOLISM OF PHENYLALANINE AND TYROSINE IN BURNED PATIENTS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 December 1975 - 30 September 1976.

Investigators: David N. Herndon, MD, Cpt, MC Douglas W. Wilmore, MD Arthur D. Mason, Jr, MD John Dubois, B.S. Basil A. Pruitt, Jr, MD, Col, MC

Reports Control Symbol MEDDH-288(R1)

Previous studies at this institution have shown all serum amino acid levels with the sole exception of phenylalanine to be decreased in thermally injured patients. This study investigated abnormalities in phenylalanine and tyrosine metabolism in burned individuals from the 7th to the 14th post burn day. Seven controls were compared with 7 non-septic patients (total body surface burn 40 - 65%, mean 49%) and with 5 septic patients (total body surface burn 40-60%, mean 50%). Oral loads of 100 mg/kg of phenylalanine and 50 mg/kg tyrosine were administered to all subjects on separate days. Phenylalanine and tyrosine tolerance curves were determined from hourly post ingestion blood samples. The phenylal nine flow was at least equal to that in controls in the non-septic patients and was significantly (p < 0.05) increased in the septic patients indicating that an increase in phenylalanine level in thermally injured patients is an index of increased peripheral production and not due to decreased metabolism. Increases in serum tyrosine levels in response to an oral load were markedly lower in patients than in controls (p<0.01) indicating a malabsorption of tyrosine similar to that observed previously in myxedematous patients. The rate of tyrosine disappearance following a phenylalanine load was increased in the non-septic patients relative to controls (p < 0.05).

Thyroid Phenylalanine Tyrosine Hypermetabolism Burns Catecholamines

#### EVALUATION OF ALTERED METABOLISM OF PHENYLALANINE AND TYROSINE IN BURNED PATIENTS

Increased basal metabolic rate, protein catabolism, and weight loss Recent evi-2 characterize the metabolic response to major thermal injury. dence indicates that catecholamines mediate this hypermetabolic response. Goodall and associates first demonstrated that increased, urinary excretion of adremaline and noradrenaline following large burns.³ The urinary excretion of catecholamines varies with the degree of injury and is related to the oxygen consumption of the patient. Catecholamine infusion in normal man increases metabolic rate,⁵ and combined alpha and beta adrenergic blockade or beta adrenergic blockade alone in a group of thermally injured patients significantly reduced metabolic rate.² Goodall and Haynes demonstrated that some patients with extensive thermal injuries have depleted adrenal medullary catecholamine stores.⁶ In additional studies, Goodall demonstrated an increase in conversion of dopamine to norepinephrine and suggested that dietary supplements of catechol procursors be added to the patients' diet to insure adequate catecholamine stores.  $\frac{1}{8}$  A recent evaluation of serum amino acid levels in burn patients at this Institute⁸ demonstrated a reduction in the concentrations of all essential amino acids with the sole exception of phenylalanine which was elevated. Phenylalanine is a precursor in catecholamine synthesis and the phenylalanine to tyrosine ratio in normal man is approximately 1.

1. Wilmore DW: Nutrition and metabolism following thermal injury.Clin Plast Surg 1: 603, 1975.

2. Wilmore DW, Long JM, Mason AD, Jr, Skreen RW, Pruitt BA, Jr: Catecholamines: mediator of the hypermetabolic response to thermal injury. Ann Surg 180: 653, 1974.

3. Goodall, McC, Stone C, Haynes BW, Jr: Urinary output of adrenaline and noradrenaline in severe thermal burns. Ann Surg 145: 479, 1957.

4. Harrison TS, Seaton JF, Feller I: Relationship of increased O₂ consumption to catecholamine excretion in thermal burns. Ann Surg 165: 169, 1967.

5. Stone DJ, Katz H, Sarkor TK, Singzon J: Ventilatory response to alpha-adrenergic stimulation and inhibition. J Appl Physiol 34: 619, 1973.

6. Goodall McC, Haynes BW, Jr: Adrenal medullary insufficiency in severe thermal burn. J Clin Invest 39: 1927, 1960.

7. Goodall McC, Alton H: Dopamine (3-hydroxytyramine) replacement and metabolism in sympathetic nerve and adrenal medullary depletions after prolonged thermal injury. J Clin Invest 48: 1761, 1969.

8. Coleman RA, Wilmore DW: Unpublished data, 1975.

9. Perry HL, Trachler B, Hansen S, MacDougal I. A simple test for heterozygosity for phenylketonuria. Clin Chim Acta 15: 47-55, 1962.
This ratio in burned patients is elevated to approximately 1.6.⁸ Similar elevations in intracellular phenylalanine levels have been observed in surgically stressed patients 1, 12 and in the serum levels of children with kwashiorkor and malnutrition. 11, 12 It is unknown if this elevation of phenylalanine is due to a block of its conversion in the liver to tyrosine or glucose, or whether it is secondary to increases in catabolsim of protein in the periphery. This study was designed to determine whether flow of phenylalanine and tyrosine is abnormal in the thermally injured patient. This information may determine whether dietary supplementation with these catechol precursors or co-factors of their enzymes would increase the catecholamine stores thereby increasing reserves in stressed, thermally injured patients.

# BACKGROUND

Phenylalanine and tyrosine metabolism have been extensively studied because in phenylketonuria there is a congenital absence of the enzyme phenylalanine hydroxylase which converts phenylalanine to tyrosine. Evaluation of parents of homozygotes for this recessive disease has demonstrated that conversion of phenylalanine to tyrosine is decreased in heterozygotes as well as homozygotes. Other studies have demonstrated that serum concentrations of phenylalanine following an oral load of phenyl-

10. Vinnars E, Bergstrom Z, Furst P. Influence of the postoperative state on the intracellular free amino acids in human muscle tissue. Ann Surg 182: 665, 1975.

12. Holt LE, Jr, Snyderman SE, Norton PM, Roitman E. The plasma aminog: am as affected by protein intake.p. 32, Protein Nutrition and Free Amino Acid Patters (L.H. Leathem, ed), New Brunswick, NJ, Rutgers Univ Press, 1968.

13. Knox WE: Metabolism of phenylalanme and tyrosme. In Symposium on Amino Acid Metabolism. W.D. McElroy, H.B. Glass, eds. John Hopkins Press, Baltimore, 1955, p. 836.

^{11.} Arroyave G, Wilson D, deFunes C, Behar M. The free amino acids in blood plasma of children with Kwashiorkor and Marasmus. Am J Clin Nutr 11: 517, 1962.

^{14.} Hsia DY, Driscoll KW, Troll WF, Knox WE Detection of phenylalanine tolerance tests of heterozygous carriers of phenylketonuria Nature 178: 1239 (Dec 1) 1956.

^{15.} Hsia DYY, Paine RS: Phenylketonuria detection of the heterozygous carrier. J Ment Defic Res 1:53,1957.

^{16.} Hsia DYY: Phenylketonuria the phenylalanine-tyrosine ratio in the detection of the heterozygous carrier. J Ment Defic Res 2, 8, 1958.

^{17.} Hsia DY, Driscoll KW Detection of the heterozygous carriers of phenylketonuria. Lancet 271: 1337 (29 Dec) 1956.

^{18.} Jarvis GA Detection of heterozygotes for phenylketonuria.Clin Chim Acta 5: 471, 1960.

^{19.} Berry H, Sutherland B, Guest GM Phenylalanine tolerance tests on relatives of phenylketonuria children. Am J Human Genet 9 310, 1957.

alanine ranging from 0.1 to 0.33 gm/kg/body weight are increased in heterozygotes when compared to the curves generated in normal man. Several investigators^{16,18,20,21} have followed serum tyrosine levels in heterozygotes after phenylalanine loading and found it to be significantly lower than the tyrosine response generated from the same phenylalanine load in normal man. Evaluation of these individuals using intravenous loading technics showed no discriminative improvement with intravenous loading compared with oral loading tests². Phenylalanine conversion to tyrosine and tyrosine tolerance tests are also abnormal in certain liver diseases.²² In all of these studies large oral loads of phenylalanine and tyrosine were given to a wide variety of patients. This study analyzes phenylalanine and tyrosine metabolism by similar technics in thermally injured patients.

# MATERIALS AND METHODS

all the restored a sublimer as a brief of the second second second second second second second second second s

Seven control subjects, 4 males, 3 females, ages 23-34, mean age 28, all with no past history of liver disease, no family hisotry of phenylketonuria and with normal liver function tests at the time of the study were compared with 5 septic burn patients (defined by having positive blood cultures on the day of the study, 2 patients had Klebsiella sepsis and 3 had Pseudomonas sepsis). Their burn sizes ranged from 40-65% of total body surface, mean 50% total body surface. Their ages varied from 19-60, with a mean age of 49. Controls and septic burns were compared with 7 uncomplicated, thermally injured patients, burn size 40-65% of total body surface, mean of 31. The burned patients were studied on the 7th to 14th post burn days.

After a 12-hour fast, patients and controls were given (1) an oral load of phenylalanine of 100 mg/kg at time zero. The phenylalanine was made palatable by mixing with 100 gm of dietetic applesauce Blood was drawn at hourly intervals starting prior to ingestion and running to 5 hours post-ingestion. Serum was analyzed by colorimetric methods (Sigma kit) for phenylalanine and tyrosine levels, and for glucose levels. (2) At a time 3 to 5 days after the phenylalanine loading tests, patients and controls were given 50 mg/ kg of tyrosine orally, made palatable by mixing with applesauce. Blood was drawn at hourly intervals starting prior to ingestion and running to 5 hours post-ingestion. Serum was analyzed for phenylalanine , tyrosine and glucose levels as above. Values at fasting and each hourly interval post-oral load were compared between the three groups by analysis of variance using Scheffe technic for multiple comparisons. When discrete disappearance curves could be defined by linear regressions, these were compared by analysis of variance and their rates of disappearance determined from the regression slope constants.

20 Wang HL, Morton NE, Waisman HA: Increased reliability for the determination of the carrier state in phenylketonuria. Am J Human Genet 13: 255, 1961.

^{21.} Anderson JA, GravemH, Ertel R, Fisch R: Identification of heterozygoteswith phenylketonuria on basis of blood tyrosine responses. J Pediat 61: 603, 1962.

^{22.} Levine RJ, Conn N: Tyrosine metabolism in patients with liver disease. J Clin Invest 46: 2012, 1962.

# RESULTS

A. Phenylalanine Tolerance Tests, 100 mg/kg Administered Orally at Time Zero. When the appearance-disappearance curves of phenylalanine in mg/100 cc of serum versus time in response to an oral load of 100 mg/kg of phenylalanine of the 7 controls are compared with all 12 burn patients (Fig. 1), the fasting levels are not statistically separate nor are the 4 and 5 hour post-ingestion values. The 1, 2, and 3 hour post-ingestion values for the patients are significantly decreased below those of the controls (p<0.025, p<0.01, and p<0.001 respectively). A regression line fit from the 1 to 4 hour, mean values for the controls has an equation Y = 14.7921 - 2.3893X with an r² of 0.9808, a similar line for the patients has an equation Y = 10.0729 -1.6304X with an r² of 0.9075. By analysis of variance, the difference in the slopes of these lines is not statistically significant but the control line is relatively elevated over that of the patients, p<0.01.

When the appearance-disappearance curves for tyrosine in mg/100 cc of serum versus time in hours in response to a 100 mg/kg phenylalanine oral load were plotted, (Fig 2) the responses of the patients could not be statistically separated from the responses of the controls.

The glucose levels in response to the above load were significantly higher in all of the patients relative to controls at all times, p < 0.01 (Fig.3); but there was no change in response to the phenylalanine load from fasting levels in either group (Fig.4).

When the 5 septic patients are compared to the 7 controls, the phenylalanine tolerance curve in mg/100 cc/hour shows the septic patients fasting levels to be significantly greater than that of the controls. (Septic patients mean mg phenylalanine/100 cc serum at time Zero = 2.330 with a standard deviation of 0.74; controls mean mg phenylalanine/100 cc serum at time Zero = 1.1514 with a standard deviation of 0.38; p < 0.01 (Fig.5). Since the fasting levels are different in these 2 groups, disappearance rates must be calculated from changes from fasting levels (Fig. 6). The equation defining the disappearance of the controls is Y = 2.39X + 13.64,  $r^2 = 0.98$ . That for the septic patients is Y = 2.03X + 10.3938,  $r^2$  = 0.92. The slopes of these lines, or the rates of disappearance, are statistically equal. The increases from fasting are statistically equal at 1 and 2 hours post-ingestion but at 3, 4, and 5 hours postingestion the increases are significantly greater in the controls than in the patients (p < 0.01). The tyrosine appearance-disappearance curves to the phenylalanine loads were the same in the controls and in the septic burn patients (Fig. 7).

When the 7 non-septic or uncomplicated burns are compared to the 7 controls, the fasting phenylalanine levels and rates of appearance and disappearance of phenylalanine are identical, but the patient curve is significantly lower at all times beyond fasting, p<0.001 (Fig.8). The tyrosine appearance-disappearance curves are notable in that the fasting tyrosine level is greater in the uncomplicated burn patients than in the controls (Fig. 9). Therefore, appearance-disappearance rates in this instance must be



ì

いいないないないであるというないためのないがないためであり







442.00

Figure 3

w



Salt and



PHENYLALANINE TOLERANCE TEST

and the second second

AT A SALE AND A SALE AND A



CALL STATE AND IN CALL

ALC: NO

いたのであるというで、「「たいたい」というになった。

12 Hel 17





Figure 8

;



è

examined by comparing changes from fasting as in Figure 10. When this is dene, there is no difference in the 2, 3 and 5 hour changes from fasting between the 2 groups, but the 4 hour values are significantly lower in the uncomplicated burn group than in the control group (p < 0.05). The regression line describing control disappearance from 2 to 4 hours post-ingestion of phenylalanine is described by the equation  $Y = 0.1150X + 1.6628 r^2 \overline{2} 0.96$ ; that for the uncomplicated patients is Y = 0.2427 + 1.4299 with an  $r^2$  of 0.98. The slope, or rate of disappearance for the burns is significantly greater in the non-septic burned group than in the control group, p < 0.05.

If the changes from fasting phenylalanine levels in response to 100 mg/ kg oral dose of phenylalanine of the septic and non-septic patients are compared as in Figure 11, the regression line describing the uncomplicated controls is Y = 6.387 - 1.2899% with an  $r_2^2$  of 0.94 and the line for the septic burns is Y = 10.3938 - 2.0318% with an  $r_2^2$  of 0.95. The septic patients have a greater elevation at one hour p  $\lt$  0.05, and a greater slope from 1 - 5 hours, p $\lt$ 0.05.

B. Serum Tyrosine Response in mg/100 cc of Serum to 50 mg/kg Tyrosine Oral Load at Time Zero. When changes in levels of phenylalanine and glucose from fasting are compared after a 50 mg/kg tyrosine oral load no statistical difference at any period could be discerned between any group of patients and controls (Fig. 12,13). When the tyrosine appearancedisappearance curves are compared for this tyrosine load, the rates of disappearance could not be differentiated but the degree of elevation from fasting was only a fraction in the patients of that seen in the controls at any given time post-ingestion, p < 0.01 (Fig. 14). The curves of septic and nonseptic patients could not be distinguished.

# DISCUSSION

A recent review of the serum concentrations of all essential amino acids to the thermally injured patients treated at this Institution showed all essential amino acids to be decreased with the sole exception of phenylalanine. Examination of the major pathway for phenylalanine metabolism (Fig. 15) would suggest that increased phenylalanine levels could be from either increased production of phenylalanine, decreased conversion of phenylalanine to tyrosine, or shunting of phenylalanine to other metabolic pathways such as gluconeogenesis. Our data show that the rate of disappearance of phenylalanine, and its conversion to tyrosine are equal in uncomplicated and complicated burns and in normals. In septic burns, since their fasting phenylalanine levels are greater than controls, the phenylalanine flow which is equal to the fasting concentration times the rate of disappearance, is increased. It is probable that the phenylalanine flow is even greater than suggested by these data as the extracellular water of thermally injured patients from the 7th to the 14th post burn day is greater than that of controls. This would cause the space occupied by phenylalanine to be larger in patients than in controls, Phenylalanine flow would be increased by the proportion by which this space is enlarged. Elevated phenylalanine levels in burned patients are then secondary to increased peripheral production and not due to decreased conversion to tyrosine. Phenylalanine levels can be taken to represent a



ě.

Figure 10



Figure 11

Sa Mashainin a Linno



Contraction of the second second

÷,







Charles and the second

____



# MAJOR PATHWAY FOR CATECHOLAMINE BIOSYNTHESIS:

Figure 15

and the production of the prod

متحقيد أاللايا

further catabolic marker in the thermally injured.

The tyrosine tolerance curves of the patients showed much smaller increases from fasting than the controls. This indicates either a much greater rate of tyrosine utilization in the burns or, more likely, a failure of absorption of tyrosine. It is interesting that the abnormalities in oral tyrosine tolerance curves observed in burned patients here, closely mimic the previously described abnormalities in tyrosine tolerance curves of myxedematous patients (Fig. 16). The tyrosine appearance-disappearance curves from 100 mg/kg phenylalanine load of uncomplicated burns compared with controls showed a more rapid rate of disappearance of tyrosine in the burns than in the controls. This would indicate that in burns at least this abnormal tyrosine tolerance test is a result of malabsorption.

Decreased absorption of tyrosine and increased flow of phenylalanine in the thermally injured patient would indicate a need for dietary supplementation of these important precursors of catecholamines and thyroid hormone.

# PRESENTATIONS AND/OR PUBLICATIONS

None

^{23.} Rivilin RS, Melmon KL, Sjoerdsma A: An oral tyrosine tolerance test in thyrotoxicosis and myxedema. N Engl J Med 272: 1143 (June 3), 1965.





the the state of the state of the second second

				1. AGENCY ACCESSION			2. DATE OF SUM	NARY	REPORT CONTROL SYMBOL				
RESEARCH AND TECHNOLOGY WORK UNIT SUMMARY			DA OG 6960		76 10 01		DD-DR&E(AR)636						
S DATE PREV SUM'RY	4 KIND OF SUMMARY	S SUMMARY SCTY	. WORK SECURITY	T REOR	DING	84. DI	BO'N INSTR'N	OL SPECIFIC	DATA-	. LEVEL OF SUM			
76 10 01 ·	K. COMP	U	υ	NA		NL		X YES		A. WORK UNIT			
10 NO /CODES *	PROGRAM ELEMENT	PROJECT NUMBER			REA NUME	PER		WORK UNIT NUMBER					
& PRIMARY	61102A	3A161102B71P			)2		069						
S. CONTRIBUTING									•				
C CONTRIBUTING								•					
11 TITLE (Procode with	security Closelfication Code	" (U) Evalu	ation, Don	or Si	te Hea	aliı	ng and Me	esh Gra	ft Sp	read With			
Hyperbaric	Oxygen Treat	ments in Bu	rned Soldi	ers (	44)								
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS												
003500 Cli	nical Medicin	e											
13 START DATE		14. ESTIMATED COM	PLETION DATE	IS FUNDING AGENCY				16 PERFOR	MANCE METHOD				
76 01		76 09	)	DA				C. In-House					
17 CONTRACT/ORANY	Not Appli	cable		18. RESOURCES ESTIMATE			-	ONAL MAN YR		L FUNDE (In Housends)			
A DATES/EFFECTIVE		EXPIRATION:			PRECEDING				10				
NUMBER *				FISCAL	/0	<u></u>				12			
G TYPE.		4 AMOUNT:		YEAR	CUMMENT								
& KIND OF AWARD		LCUM, AMT.											
IS RESPONSIBLE DOD	ORGANIZATION	_		30. PERI	PORMINGOR	GANI	ZATION						
HAME!" US Army	Institute of	Surgical	Research		US Ar	my	Institut	e of Su	rgica	1 Research			
				Clinical Division									
ADDRESS." Fort	Sam Houston.	Texas 782	34	ADDAEss!"Fort Sam Houston, Texas 78234									
1010		ICAUD /OD		Ì									
				PRINCIPAL INVESTIGATOR (Punnish SEAN II U.S. Association (politonian,									
RESPONSIBLE INDIVIDUAL				HAME' Hugh D. Peterson, COL, MC									
HANE Basil A. Pruitt, Jr, COL, MC				TELEPHLIE 512-221-3301									
TELEPHONE: 512-221-2720				SOCIAL SECURITY ACCOUNT NUMBER									
21. GENERAL USE				AMOCIATE INVESTIGATORS									
l				NAME.									
FOREIGN INTELLIGENCE NOT CONSIDERED NAME:							DA						
TE. REV BORDS (Procedo	EACH of B. Jonally Classi	faatlan (ada)											

(U) Hyperbaric oxygen; (U) Mesh graft; (U) Donor site; (U) Human3

13 TECHNICAL OBJECTIVE.* 34 APPROACH, 35 PROGRESS (Puriod Individual paragraphs Hamilting by number Protects test of each with Security Classification code.)
23. (U) The objective of this study is to attempt to demonstrate that hyperbaric oxygen shortens the healing time of donor sites and increases the rate of spread of mesh graft applied to suitable recipient areas of burned soldiers.

24. (U) The patients were divided into two groups, those receiving hyperbaric oxygen treatments b.i.d. at 2.4 atmospheres of 100% pure oxygen and those not receiving this treatment. In both groups the uniform depth donor site was placed with a Davol dermatome and the time required for healing of this area recorded. A portion of mesh graft was placed on a healthy granulating bed and photografted every two days to observe the filling of the interstices of the mesh.

25. (U) 76 01 - 76 09 The patients treated with hyperbaric oxygen healed their documented depth donor sites at a rate equal to or slower than the patients not subjected to hyperbaric treatments. The mesh graft spread in patients not receiving treatments. A separate part of this study showed that documented depth donor sites in volunteers heal at the same rate with hyperbaric oxygen as when untreated. The unescapable conclusion from our work is that hyperbaric oxygen does not accelerate the boaling of partial thickness donor sites nor does it speed the healing of mesh graft. Because of the total lack of demonstrable response to hyperbaric oxygen this study was discontinued and the chamber removed from the unit.

FINAL REPORT

PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

berther his data and the street street in the second street with the

REPORT TITLE: EVALUATION, DONOR SITE HEALING AND MESH GRAFT SPREAD WITH HYPERBARIC OXYGEN

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 1 July 1975 - 30 September 1976

Investigators:

Hugh D. Peterson, DDS, MD, Colonel, MC Basil A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

### ABSTRACT

# PROJECT NO. 3A161102B71P-02, BASIC RESEARCH IN SUPPORT OF MILITARY MEDICINE - SURGICAL PATHOLOGY

**REPORT TITLE:** EVALUATION, DONOR SITE HEALING AND MESH GRAFT SPREAD WITH HYPERBARIC OXYGEN

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, 1X 78234

Period covered in this report: 1 January 1976 - 30 September 1976

Investigators: Hugh D. Peterson, DDS, MD, Colonel, MC Basil A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

Among the many benefits claimed for hyperbaric oxygen in the treatment of burned patients, are decrease in time of donor site healing and increased rapidity of mesh graft spread. These were evaluated on documented depth split thickness skin graft donor sites on convalescent patients requiring grafting and on "mesh" grafts with documented size interstices. The hyperbaric oxygen group were treated with 2.4 atmospheres of 100% oxygen for 60 minutes b.i.d. from the time of graft harvest and placement of graft until healing was complete. No acceleration could be noted when compared with a control group treated in the same fashion but without hyperbaric oxygen treatments. Because of failure to demonstrate any improvement in healing time or mesh graft spread and the difficulty of administering hyperbaric oxygen to patients with large burns, the study was abandoned and the hyperbaric chamber removed from the unit.

Hyperbaric oxygen Mesh graft Donor site

# EVALUATION, DONOR SITE HEALING AND MESH GRAFT SPREAD WITH HYPERBARIC OXYGEN

There are four purported advantages of administration of hyperbaric oxygen to burn patients; namely, 1) decrease in amount of infection. 2) prevention of conversion of deep second degree burn to third degree burn, i e., less areas need grafting, 3) decrease in amount of resuscitation fluid needed in the first 48 hours, henceforth less edema and 4) decreased healing time c^c donor sites and increased rapidity of mesh graft spread. A single person hyperbaric oxygen chamber was obtained and it was elected to first evaluate mesh graft healing and healing of documented depth donor sites. For the initial evaluation small donor sites were placed on either the upper thigh or the forearm of patients requiring grafting procedures. These were done with the Davol dermatome which takes a set thickness graft. The small specimens were pinned on a tongue blade and submitted to pathology for direct measure of the thickness. The donor sites were treated open with fine mesh gauze. The subjects were treated to one hour of 100% oxygen at 2.4 atmospheres twice a day until healing occurred. Healing being defined as separation of the coagulum with total epithelialization. Once the donor site treated with hyperbaric oxygen had healed and separated, a like donor site was placed on the other extremity. Its depth was documented, and it was treated in the same fashion with the exception of the hyperbaric oxygen treatments. Time to healing in the like depth donor sites was 13 to 14 days with and without hyperbaric oxygen. There was no demonstrable decrease in the time of healing.

Donor site healing and mesh graft spread were evaluated in convalescing patients in the 4th to 5th week postburn with wounds of healthy granulation tissue ready for grafting. At the time of grafting a small portion of uniform depth graft was taken with the Davol dermatome and a portion of the skin taken for definitive grafting was also submitted to pathology for measurement of depth. The donor sites were treated in a usual open fashion after placing a single layer of fine mesh upon them. A three inch square area of healthy granulation tissue was picked and a representative portion of the 3:1 meshed graft was placed thereupon with maximum expansion and photographed with a ruler in the field. The meshed graft was the dressed with fine mesh gauze and coarse mesh gauze scaked in 5% Sulf. ylon solution. The patients to be subjected to hyperbaric oxygen were treated in this fashion and then treated with 2.4 atmospheres of 100% oxygen for one hour b.i.d. to include one treatment the day of surgery. Only two patients were in the hyperbaric oxygen group and four in the control group. This is obviously a very small group but since there was absolutely no decrease in healing time of the documented depth donor sites nor any increase in the rapidity of mesh graft spread, the investigaters were reluctant to increase the size of the series. If there is significant benefit from treatment with hyperbaric oxygen in increasing healing, it should be demonstrated in a very small series of patients. A difference that would require 50 patients in either group to document statistical significance would be of little clinical value.

Because of our failure to document any improved healing, it was elected to discontinue the hyperbaric oxygen study and return the hyperbaric chamber to the manufacturers.

PUBLICATIONS AND/OR PRESENTATIONS

None.

RESEARCH AND TECHNOLOGY WORK UNIT SUMMARY			DA OF 6393			2. DATE OF BU 76 10	01	REPORT CONTROL SYMBOL DD-DR& C(AR)636					
& DATE PREV SUM'RY	A DATE PREV SUM'RY & KIND OF SUMMARY S. SUMMARY SCTY			7. REGRADING"		EB'H INSTR'H	Sh SPECIFIC	DATA-	. LEVEL OF SUN				
75 07 01	H. TERM	U	U	NA N		NI	<b>.</b>	U YES		A. WORK UNIT			
IO NO /CODES.	PROGRAM ELEMENT	PROJECT	TASK AREA NUMBER			WORK UNIT NUMBER							
& PRIMARY	62110A	3A1621102	821	00			110						
b. CONTRIBUTING													
C CONTRIBUTING		L											
II TITLE (Procode with	Soburity Classification Code	" (U)Low Fl	ow Vein Ana	stom	osis i	n I	ogs Ant:	icoagula	ated 1	With			
Ancrod: A M	odel for Bloc	d Vessel R	<u>epair in Ir</u>	iture	d Troo	)ps	(44)	·····					
002500 014	ical Madiaina												
13 START DATE	ical Medicine	IL. ESTIMATED COM	PLETION DATE	IL PUNI		Y		14. PERFOR		FHOD			
75 01		75 06		DA	1		1	C. I	-Hou	se			
17 CONTRACT/GRANT	Not Applica	ble		10. RES		TIMAT		IONAL MAN YE	8   h. 70	NOS (In these and it)			
A DATES/EFFECTIVE		EXPIRATION:		<b></b>	PRECEDIN				-				
NUMBER.*				FISCAL	76		.5			18			
G TYPE:		4 ANDUNT:		YEAR	CONNERT	-							
& KIND OF AWARD:		f. CUM. AMT											
IS RESPONSIBLE DOD C	RGANIZATION			<b>38.</b> PERI		GANI	LATION						
HANE."US Army	Institute of	Surgical R	lesearch	HANEI [#]	US Art	ny 1	Institute	e of Su	gica	l Research			
_				Metabolic Branch									
ADDRESS!" Fort S	am Houston, 7	Cexas 7823	4	ADDARses Fort Sam Houston, Texas 78234									
				PRINCIPAL INVESTIGATOR (Pumiek SSAN If U.S. Asadesis prefinition)									
RESPONSIBLE INDIVIDU Racil	AL A Druitt .Tr	COT. MC		TELEPHONE: 512-221-2411									
NAME DUSII	-221-2720	, con, nc		TELEP		2-2	221-3411						
TELEPHONE, JLL													
FOREIGN INT	ELLIGENCE NOT	CONSIDERE	D	HAME	W- 11-		N Nrde						
			-	NAME:	John	W.	Sagartz	, VD		DA			
22. KEYWORDS (Presede	BACH will Somethy Closell	cation Code)		4									
(U) Coagula	tion; (U) And	rod; (U) V	ascular sur	gery	; (U)	Dog	js						
23 TECHNICAL OBJECT	IVE.* 24 APPROACH. 25	PROGRESS (Pumish )	individual paragrapho ide	milled by	number pri	tada k	est of each with s	penally Classif	eation Cud	<b>b.</b> j			
23. (U) A	ncrod is ath	rombin-like	e enzyme der	rived	from	the	e venon (	of the I	lalaya	an pit			
viper. It	is a very pot	ent antico	agulant wit	h fe	w unde	esir	able sid	de effe	cts.	Vein			
reconstruct	ions followir	ng the pene	trating or	crus	h inju	irie	es which	occur	frequ	ently			
in modern w	arfare have r	not been po	ssible with	n con	venti	onal	antico	agulant	5. W	ith Ancrod			
it may be p	ossible to pe	erform vasc	ular recons	struc	tion :	in d	cases wh	ich wer	e pre	viously			
regarded an	inoperable.	The purpo	se of this	stud	y is '	to	levelop	a vascu	lar s	urgical			
model which	is highly p	redisposed	to thrombos	sis a	nd to	ut:	llize th	15 mode	i in	cne			
study of Ar	icrod as a sui	rgical anti	coagulant :	tor u	se in	in	jurea so	laters.					
	· · · · · · · · · · · · · · · · · · ·				hilst		llu in a	aroup	of si	tube v			
24. (U) Tr	ie remoral ve:	in was sure	Jically exp		and a	51 a.	astomos	od Ve	noara	ms were			
Mongrel dog	s. The Vein	was then c	cut transve	rsery	ang	. eai	145 (01105	eu, ve	liogra				
performed a	it two and for	ur weeks po	St Operation	.crì.									
25 /11) 74	07 - 75 OP	Most of	the vein a	naeto	moses	re	mained n	atent a	t two	weeks			
25. (0) /:	$\frac{1}{100} = \frac{1000}{100}$	which wer	the vern a	nastu 1 at	two w	ee k	s had re	canaliz	ed an	d were			
postoperati	Lvery. verns	which were	slv The d	na ie	thus	no	t an ade	guate m	odel	for these			
patent by a	nause of its	active fin	cinolytic s	vster	and	ano	ther mod	el will	be n	ecessary.			
studies dec	cause of its	accive rap	THOTICTO D	1									
1													

- --

• Available to contractors upon originatore approve. DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 1488A I NOV 68 AND 1498 I I MAR 68 (FOR ARMY USE) ARE OBSOLETE 376

# TERMINATION

. . ....

PROJECT NO. 3A162110A821-00, COMBAT SURGERY

and the design of the second 
40° 77 7

REPORT TITLE: LOW FLOW VEIN ANASTOMOSIS IN DOGS ANTICOAGULATED WITH ANCROD--A MODEL FOR BLOOD VESSEL REPAIR IN INJURED TROOPS

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

> > 1 July 1975 - 30 September 1976

Investigators

Clement L. Slade, M.D., Captain, MC Willard A. Andes, M.D., Major, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

### ABSTRACT

PROJECT NO. 3A162110A821-00, COMBAT SURGERY

REPORT TITLE: LOW FLOW VEIN ANASTCMOSIS IN DOGS ANTICOAGULATED WITH ANCROD--A MODEL FOR BLOOD VESSEL REPAIR IN INJURED TROOPS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Clement L. Slade, M.D., Captain, MC Willard A. Andes, M.D., Major, MC

**Reports** Control Symbol MEDDH-288(R1)

Ancrod is a thrombin like enzyme derived from the venom of the Malayan pit vipor. This agent is a very potent anticoagulant with few undesirable side effects. It is possible this anticoagulant could be used as an adjunct in vascular surgical procedures which are predisposed to thrombosis. The purpose of this study was to develop a model vascular surgical procedure which was highly predisposed to thrombosis and to use this model in the study of Ancrod as a surgical anticoagulant.

Femoral vein anastomoses were performed bilaterally on six adult mongrel dogs who were not anticoagulated. Follow up venograms at two and four weeks postoperatively, revealed a high rate of patency of the anastomosis. Veins which had clotted at two weeks postoperatively, were found to have recanalized by four weeks postoperatively, therefore the project was terminated. The dog is thus not an adequate model for these studies. It may be necessary to use primates to continue this work.

Coagulation Ancrod Vascular surgery Dogs

PET APCH AND TECHNOLOGY MORE INNE STANKAR					ICY ACCES	SION	2. DATE OF BU	MMARY®	REPORT CO	TROL STREDL		
REL ARUN	AND TECHNOLOGI	WORK UNIT S	UMMARY	DA	OE 69	984	76 10	01	DD-DR	E(AR)636		
S. DATE PREV SUN'RY	4. KINP GF STIMMARY	S. SUMMARY SCTY	. WORK SECURITY	7. REOR	ADING	-	BR'N INSTR'N	IS SPECIFIC	DATA- 9.	LEVEL OF SUM		
75 07 01	K. COMP	U	U	N	A	N	L	CONTRACTOR	ACC ESS	A. WORK UNIT		
ID. NO./CODES:		PROJECT	NUMBER	TAS	AREA NUN	-	TORK UNIT HUNBER					
	62110A	3A162110	A821	1	00		114					
. CONTRIBUTING												
C. CONTRIBUTING								•• • •				
11. TITLE (Proceds with )	spective Co	mpar	ison a	Stud	y of Su	famylon	and S	ilver				
Sulfadiazin	e in the Trea	tment of B	urned Troop	os (4	4)		_					
18 SCIENTIFIC AND TEC	CHNOLOGICAL AREAS											
003500 Clin	ical Medicine	1										
13 START DATE		14 ESTIMATED COMP	LATION DATE	TE FUN	DING AGEN	CY		16 PERPORMA		0		
73 12		76 09		DA			L	C. In-House				
17. CONTRACT/ GRAN T	Not Applicab	le		10 RES	N RESOURCES ESTIMATE			IONAL MAN YRS	L FUND	(In those and a)		
& DATES/EFFECTIVE		EXPIRATION	I		PHECEDINE		,5	,5		18		
A NUMBER *				PISCAL	PISCAL 7T		.2		4	4		
C TYPE:		& ANDUNT:	1	YEAP	COMMENT				1			
& KIND OF AWARD:		f. CUM. AMT.										
IS RESPONSIBLE DOD O	REANIZATION			30. PERF		RGANEL	ATION		-	1		
HANE!" US ARTY	Institute of	Surgical A	Research	HAME:*	US AI	mv	Institut	e of Su	rgical	Research		
				Surgical Study Branch								
Fort S	am Houston, T	exas 78234	4	ADDRESS!" Fort Sam Houston, Texas 78234								
		<b>Undu</b>	•		• • •				uə , -	237		
				PRINCIPAL INVESTIGATOR (Pumion BRAN II U.S. Academic Inclination)								
RESPONSIBLE INDIVIDUA	AL.		1	NAME* Hugh D. Peterson, COL, MC								
HANE Basil A.	Pruitt, Jr,	COL, MC		TELEPHONE 512-221-3301								
TELEPHONE: 51	2-221-2720			LOCIAL SECURITY ACCOUNT NUMBER								
21. GENERAL USE				ASSOCIATE INVESTIGATORS								
FOREIGN INT	ELLIGENCE NOT	CONSIDERED	ן כ	*AME: Arthur D. Mason, Jr, MD								
	خورداد برواندين المترافعات متاكر			NAME.						DA		
SI, KEYBORDS (Procedo S	ACH will Be outly Classific	ellen Code) (U) 7	fopical the	erapy	; (U)	Bur	n injury	(; (U) S	ulfamy	lon		
(mafenide);	(U) Silvaden	e (silver s	sulfadiazin	ne);	(U) Hu	ımanı	S					
A TECHNICAL OBJECTI	VE. 4 24 APPROACH. 25.	PROGRES" (Furnish int	divinuel serentestates	atiliad be	sumber Ba	C a da 14.						

23. (U) The objective of this study is to compare silver sulfadiazine with Sulfamylon in six groups of surned soldiers, two groups by age 16 to 40, and greater than 40 and three by burn size, 30 to 50 percent, up to 70 percent and greater than 70 percent. The indices to be evaluated are survival, acid base balance, pain on application, bacteriology of the burn wound, pulmonary function, and clinical status of the injured troops.

entified by number Precede test of each with Security Classification Code )

24. (U) Burns greater than 30 percent, seen within the first 72 hours, were randomized by pairs in the aforementioned groups and placed in one of the two agents. The pulmonary functions, acid base balance, wound bacteriology were then followed meticulously until the eschar was entirely separated or until the patient expired.

25. (U) 75 07 - 76 09 The study demonstrated that Sulfamylon offered far superior control of the gram-negative population of the burn wound, but had deleterious effects when applied in a regular fashion to burns over 30 percent. The deleterious effects were manifested by respiratory problems, decreased G1 function, cerebral changes and earlier death not characterized by sepsis. The study demonstrated that Sulfamylca was most likely not a good agent for initial burn therapy in burns larger than 30 percent but should be reserved for wound problems where it has been very efficacious at this unit. Since the termination of this study silver sulfadiazine has been used as the initial agent we have had increasing problems with the gram-negative flora of the wound and attention is now being turned to an alternating therapy of silver sulfadiazine and Sulfamylon in the early postburn course with careful moderation systemic toxicity to the Sul amylon.

DD. FORM 1498

A service and a service to the service in the service of the servi

A MARKAN CALL TANKS

# FINAL REPORT

. . . . . . . . .

# PROJECT NO. 3A162110A821-00, COMBAT SURGERY

# **REPORT TITLE: A PROSPECTIVE COMPARISON STUDY OF SULFAMYLON AND SILVER SULFADIAZINE IN THE TREATMENT OF BURNED TROOPS**

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Hugh D. Peterson, DDS, MD, Colonel, MC Arthur D. Mason, Jr., MD Basil A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

### ABSTRACT

## PROJECT NO. 3A162110A821-00, COMBAT SURGERY

# REPORT TITLE: A PROSPECTIVE COMPARISON STUDY OF SULFAMYLON AND SILVER SULFADIAZINE IN THE TREATMENT OF BURNED TROOPS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Hugh D. Peterson, DDS, MD, Colonel, MC Arthur D. Mason, Jr., M.D. Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

During all of the reporting period only silver sulfadiazine was used as the initial topical agent. It was noted to have decreasing efficacy against gram negative organisms mainly Klebsiella. In addition Pseudomonas organisms have been more commonly recovered from the burn wounds, in terminal patients. This decreasing efficacy was highlighted by earlier positive blood cultures and earlier time to death in those patients dying with silver sulfadiazine treatment. Sulfamylon burn cream was again used for wound complications and several patients had wound degeneration which was recouped by subeschar infusion and intermittent application of Sulfamylon burn cream. Sulfamylon burn cream when used early in the burn course was again noted to have a deleterious effect on some patients. Probably the most outstanding thing noted in survivals during this period was that there were only two deaths in the 15 to 40 age group of burns less than 50%. One of these was a patient that entered the hospital with a cerebral death secondary to inhalation injury, the second was a 50%burn that died at 126 days with a nonspecific myocarditis. This low mortality in the 15 to 40 age group in burns of 50% or less contrasts to the 18 deaths in that age group in 1970 and the 14 deaths in that age group in 21973, the last year in which Sulfamylon was used as a primary topical agent. Figures such as this further document the benign quality of silver sulfadiazine in relation to systemic effect on the patient. However this fact is at least in part counter balanced by the poor control of Klebsiella sp.

This study has been terminated because we have continually demonstrated our earlier findings of better bacterial control by Sulfamylon, which must be balanced against its systemic effects in patients with burns of more than 30% of the total body surface. At present we are evaluating a treatment regimen in the early post burn period of 12 hours application of silver sulfadiazine followed by 12 hours application of Sulfamylon alternating in this fashion until the eschar has separated. With this maneuver we have been able to maintain the bacterial density in the eschar at a level of  $10^{11}$  to  $10^{21}$  less than if only silver sulfadiazine is used. However an increase in the survival of patients with extensive burns has not been confirmed and we have noted some patients to show the systemic effects of Sulfamylon application. At present we are continuing to pursue this course in patients with burns larger than 50%. The findings of this study suggest that there is probably no ideal topical agent and that emphasis should be placed on surgical removal.

Topical therapy Sulfamylon (mafenide) Silvadene (silver sulfadiazine)

Burn injury Humans

DESCARCH AND TECHNOLOCY MORY INIT SHIMLARY				-	CY ACCESSI	-	DATE OF SU	UMMARYS REPORT CONTROL STREOL					
RESEARCH	AND TECHNOLOGI	WURK UNIT S	UMMART	DA	OF 639	1	76 10 01			DD-DR&E(AR)436			
S. DATE PREV SUM'RY	4 KIND OF SUMMARY	S SUMMARY SCTY	S WORK SECURITY	PRESA	DING	- 016	B'H INSTA'H	CONTRACTOR		TA - P.	LEVEL OF DUM		
75 07 01	к. ~	U	U	NA			NL	2 vz	. 🛛	10	A TORK UNIT		
10 NO /CODES.*	PROC LEMENT	PRUJECT	NUMBER	TASK AREA NUMBER			WORK UNIT NUNBER						
	62110A	3A162110A	821	<u> </u>	00		105						
S. CONTRIBUTING		_					ty		n ji ji				
C. CONTRIBUTING													
11 TITLE (Procode with	Security reselfication Code	*(U) Evalua	tion of th	e Eff	ect of	Fr	esh-Fro	zen	Plasm	na on			
Leukocyte (	Chemotaxis in	Burned Sol	diers (44).										
12 SCIENTIFIC AND TE	CHNOLOGICAL AREAS												
003500 Cli	nical Medicine	3											
13 START DATE		14. ESTIMATED CONF	PLETION DATE	IL PURC	IS. FUNDING AGENCY		14. PERFORMANCE MET			CE METHO	0		
74 09		75 10	)	DA	DA		<b>_</b>	<u> </u>		-House			
17 CONTRACT/GRANT	Not Applical	ble		18	OURCES EET	MATE	A PROFEM	BRIONAL MAN YAS L FUNDS (In these and ag					
N DATES/EFFECTIVE: CXP		EXPIRATION.	CZPIRATION.		PRECEDINE	-	i	-		18			
L NUMBER *	S. NIIMBER *				76	<u> </u>		4					
G TY <b>PE.</b>		4 ANOUNT:		YEAR	CORRENT		1						
S. F OF AWARD		I. CUM. AMT.											
19. RESPONSIBLE DOD	DRGANIZATION				PORMINGOR	GANIZ.	A TION	L					
NAME:" US Army	Institute of	Surgical H	Research	HANE:"	JS Army	/ Ir	stitute	e of	Surgi	ical	Research		
				Burn Study Branch									
ADDRESS:* Fort	Sam Houston, '	<b>Texas 782</b> 3	34	ADDAESS Fort Sam Houston, Texas 78234									
				PRINCIPAL INVESTIGATOR (Pumion BEAN II U.S. Academic Inclinution)									
RESPONSIBLE INDIVIDU	IAL			NAME. James W. Taylor, MAJ, MC									
HAME Basil A. Pruitt, Jr, COL, MC			TELEPHONE: 512-221-2943										
TELEPHONE: 512-221-2720					SOCIAL SECURITY ACCOUNT NUMBER								
21 GENERAL USE					ASSOCIATE INVESTIGATORS								
					****James M. Long, III, MD								
FOREIGN IN	I *****Arthur D. Mason, Jr, MD DA												
(II) Burne	(11) Leukocyt	es: (II) Ch	emotaxis.	(II) H	mans:	(11)	Plasma	•					

and a state of the second s

23 TECHNICAL OBJECTIVE,⁴ 24 APPROACH, 24 PROGRESS (Pumilsh and programs idmilled to make Protect itsi of each with provide Classification Control 
24. (U) Patients with burns over 55% of the total body surface are divided on admission into two groups. One receives the routine resuscitation, the other receives fresh-frozen plasma on colloid during their resuscitation. The functional chemotactic index is determined serially in these patients according to the Warden modification of the Boyden technique namely the ability of leukocytes to migrate through nucleopore filter toward a complement dependent chemotactic, CASEN-SERUM. Other patients who have a low chemotactic index but who had received no fresh-frozen plasma prior to determination, are treated with three units of fresh-frozen plasma per day and the functional chemotactic index determined during the period of treatment with freshfrozen plasma.

25. (U) 75 07 - 75 10 Serial studies of functional chemotactic index (FCI) in 24 burn patients between 3 and 21 days postburn revealed a significantly higher mean FCI (61.4) in those who survived than in those who died (mean FCI 44.7). Twelve patients with low FCI who received fresh-frozen plasma daily showed an increase in mean FCI from 37.7 to 56.0. The changes in patients receiving fresh frozen plasma was not merely an effect of time.

a coro va l

ntractora upon ortginato

# FINAL REPORT

# PROJECT NO. 3A162110A821-00, COMBAT SURGERY

# REPORT TITLE: EVALUATION OF THE EFFECT OF FRESH FROZEN PLASMA ON LEUKOCYTE CHEMOTAXIS IN BURNED SOLDIERS

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

James W. Taylor, M.D., Major, MC James M. Long, III, M.D., Lieutenant Colonel, MC Arthur D. Mason, Jr. M.D. Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

## ABSTRACT

PROJECT NO. 3A162110A821-00, COMBAT SURGERY

REPORT TITLE: EVALUATION OF THE EFFECT OF FRESH-FROZEN PLASMA ON LEUKOCYTE CHEMOTAXIS IN BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 73234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: James W. Taylor, M.D., Major, MC James M. Long [1, M.D., LTC, MC Arthur D. Mascn, Jr., M.D. Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288 (R1)

Warden and his associates reported that leukocytes from burned patients showed decreased chemotaxis and that after 72 hours post burn the impairment of leukocyte chemotaxis is directly correlated with the clinical status of the patient and is highly predictive for ultimate mortality, and that in vitro normal serum can restore chemotactic function of the granulocytes from burned patients to normal.

In the present study, chemotaxis of leukocytes from burned patients was studied using a modified Boyden chamber following the methods described by Warden. All studies were performed in triplicate with the person counting the cells blinded as to the source of the specimen. Eightyfour determinations of the Functional Chemotactic Index (FCI) were made on 24 burned patients between the third and twenty-first post burn day. Those burned patients who ultimately survived showed a mean FCI of 61.4 (range 19.7 to 118) compared to 44.7 (range 11 to 100.7) for the nonsurvivors (p 0.01).

Subsequently, 12 burned patients who were first demonstrated to have a low FCI were treated with three units of fresh frozen plasma per day, and the FCI was determined serially. The mean FCI following treatment was 56.0 (range 36.2 to 93.9) compared to 37.7 (range 15.4 to 68.4) before treatment (p 0.01). Comparison of this group with similar patients from our earlier series who received no fresh frozen plasma showed that this was not merely an effect of time.

This study reconfirmed the findings that ultimate mortality in burned patients is associated with significant depression of the FCI and demonstrated that leukocyte chemotaxis in burned patients can be enhanced in vivo by some constitutent of fresh frozen plasma.

Burns Leukocytes

Chemotaxis Humons Plasma
RESEARCH	AND TECHNOLOGY	WORK UNIT SI	UNILARY	I. AND	CV ACCEN		2. BATE OF SHE		DD-DR	WITHOL STREEL
		-				<u></u>			DATA- B	-
1 MATE PREV 300								CONTRACTO	ACC 238	
75 07 01	H. TERM		<u> </u>		<u> </u>	N		La ves		
IS. NO./CODES:*	PROGRAM ELEMENT	PROJECT	NUMBER	TABK A	REA HUN			HORK UNI	THUNGER	
a, PREMARY	62110A	3A162110	<u>N821</u>	<u> </u>	00		10/			
. CONTINUTING		L		<u> </u>						·
C. CONTROUTING		L								
II TITLE (Proceds and )	Soturity Classification Code	(U) Evalua	ition of Ga	istric	Phys:	iol	ogic Dis	curbanc	ces As	sociated
With Therma	<u>l Injury in a</u>	Military I	Population	(44)						
L SCIENTIFIC AND TES	CHIOLOGICAL AREAD									
003500 Clin	ical Medicine									
11 STAR" DATE		14. ESTRATES CON	PLETION DATE	TIL FUNC		¥		H. PERFOR	NANCE MET	400
72 01		76 09		DA				C. II	n-Hous	е
T CONTRACT BRANT	Not Applic	able		14. MEN	OURCES EST		-		RS & FUN	DE (In Bronada)
& DATES/EFFECTIVE.		EXPIRATION:			76		1	-5		16
L				PHICAL	7T		1	.2		6
C TYPE:		4 ANDUNT:		YEAR	COMMENT		1		1	
& KIND OF AVARD:		f. CUM. AMT.	·		I					
W, RESPONSIBLE DOD ORGANIZATION										
ANNE? US Army Institute of Surgical Research "ANNE? US Army Institute of Surgical Research										
	Y ALLOUZOULE -	· · · · · · · · · · · · · · · · · · ·		i	Bu	rn	Study Br	anch	-	
ADDRESS." BOTH C	am Houston . T	Novas 7873	4		Fort	Sa	m Housto	n. Tex	as 78	234
FOLC 5	And Houseen's .	CALD / ULU	3							
i i				PRINCIP		64 10	n (Punulah SBAR I		de jaothuilan	•
	VAL			HAME!	• Dav	id	K. Teega	rden,	MAJ, M	IC .
HANE Basil A	A. Pruitt, Jr,	, COL, MC		TELES	Pron E	512	-221-653	2		1
	2-221-2720	-		SOCIA	L SECURITY	-	UNT NUMBER-			ļ
21. OENERAL USE				-		GATO	**			
FORETCH TNT	FELLIGENCE NOT	CONSIDERE	Ð	HANE	Basil	A.	Pruitt,	Jr, C	OL, MC	:
FUREIGN INI	Linit Grater		2		Albe	rt .	J. Czaja	, M.D.	•	DA
IL KEYBORDS (Procedo	BACH will Security Closell	fanden (ada) /11)	Thormal in	inru.	(11) 5	hirr	natient	e · (11)	Evalu	ation:
		(V)	THETHER TH	յա չ,	(0) 1	NIT I	l pacient	3, (0)		u czossy

(U) Gastric physiologic Disturbances

the second of the second states and the second states of the second second second second second second second s

23. (U) To study the gastric pathophysiology of the thermally injured soldier, so as to better define etiologic factors responsible for Curling's ulcer.

24. (U) Evaluation to be carried out on thermally injured soldiers with greater than 30% TBS area injury admitted to the USAISR. Study will be stratified so that a group of patients in the 30 to 50% TBS area injury and the second group in the 50 to 70% 185 area injury will be included. Investigative procedures will be performed within 24 hours if possible and at 72 hours postburn. Burns of greater than 50% body surface area will also be studied at 5 to 7 days post injury and all patients will then be studied between the 9th and 12th day post injury and at 30 days or discharge. Studies will encompass: (1) gastric endoscopy with photography and biopsy for semiquantitative mucous determination, (2) ion flux across the gastric mucosa, (3) coagulation studies, (4) measurements of gastric clearance of radioactive isotopes, and (5) evaluation of the role of bacteremia.

25. (U) 75 07 - 76 09 Early gastritic and duodenitic changes are present in 7-80% of patients with burns of 40% or more of the body surface. Gastric acid production is increased markedly in those patients who develop clinically significant complications of Curling's ulcer. Gastrin level assay suggests an intact gastrin control mechanism. Prophylactic antacid therapy reduces bleeding from this variant of stress ulcer disease.

DD FORM 1498 PAR VIOUS EL ANO 1488 1

### TERMINATION REPORT

# PROJECT NO. 3A162110A821-00, COMBAT SURGERY

### REPORT TITLE: EVALUATION OF GASTRIC PHYSIOLOGIC DISTURBANCES ASSOCIATED WITH THERMAL INJURY IN A MILITARY POPULATION

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

David K. Teegarden, M.D., Major, MC Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

### PROJECT NO. 3A162110A821-00, COMBAT SURGERY

## **REPORT TITLE: EVALUATION OF GASTRIC PHYSIOLOGIC DISTURBANCES ASSOCIATED** WITH THERMAL INJURY IN A MILITARY POPULATION

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: David K. Teegarden, M.D., Major, MC Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

Endoscopic abnormalities of the gastroduodenal mucosa develop in 78% of thermally injured patients with burns involving 25% or greater of the total body surface.

Physiologic studies revealed that production of gastric mucosubstances was normal and not etiologically significant. Increased permeability of the gastric mucosa to hydrogen ions was not consistently noted in those with endogenous abnormalities but correlated with clinical progression of disease.

Current studies include evaluation of serum pepsinogen as a predictor of gastroduodenal mucosal disease, duod@nopyloric reflux of bile salts as an etiologic factor in stress ulcers, and evaluation of cimetidine as a therapeutic agent in prevention of stress ulcers following thermal injury.

Evaluation Gastric physiologic disturbances Thermal injury Burn patients

### EVALUATION OF GASTRIC PHYSIOLOGIC DISTURBANCES ASSOCIATED WITH THERMAL INJURY IN A MILITARY POPULATION

Over the past several years, gastrophysiologic studies have been performed in the thermally injured soldier to determine those factors that may be important in the etiology of erosive and ulcerative disease of the stomach and duodenum. Patients with a greater than 25% total body surface burn were at most risk to develop gastroduoderal complications and were included in the physiologic studies.

#### Gastroduodenal Endoscopy.

Endoscopy of the esophagus, stomach, and duodenum was performed in a serial fashion. Seventy seven thermally injured patients were included in the study and ranged in ages from 16 to 74. Superficial gastric mucosal injury was noted in approximately 87% of the patients. The gastric lesions were most often noted in the fundus and body of the stomach, and the antrum was less often involved but was abnormal in one third of the patients. Discrete ulcerative disease was found in 26.5% of the patients, and, of these, six had concomitant duodenal ulcers. Duodenal inflammation was found in 81% of those patients, and was noted to be present only in those patients with a greater than 35% total body surface burn. Duodenitis was the most frequently encountered diagnosis, and was defined by the appearance of erythema, edema, and increased mucosal friability. In two patients, ulcerative disease was seen to evolve from superficial erosive disease. Discrete duodenal ulceration was encountered in 26% of the patients, and all were noted to have accompanying duodenitis.

It can be concluded from this endoscopic evaluation that superficial gastroduodenal mucosal injury is a frequent accompaniment of severe thermal injury. This study emphasizes that this disease is a diffuse process and involves many portions of the gastroduodenal mucosa. Discrete ulceration in either the stomach or duodenum was not present unless accompanying superficial inflammatory changes were noted around the ulceration.

#### Physiclogic Studies

Histochemical techniques were employed to study the mucous protective barrier of the stomach. It has been postulated in previous literature that a decrease in mucus production, or a change in its characteristics, might render the gastric mucosal membrane more susceptible to damage. Of the nine patients studied, all were noted to have a normal production of gastric mucus. It was therefore concluded that mucus substances produced in the stomach are not an important etiologic factor in the development of acute gastroduodenal stress ulceration foilowing thermal injury. A second study utilized the lithium flux technique to evaluate back-diffusion of the hydrogen ion in the gastric mucosa and the integrity of the gastric mucosal barrier. Our data documented that increased back diffusion of hydrogen ions was not an etiologic factor in the early development of gastric lesions following thermal injury. However, disruption of the gastric mucosal barrier seemed to correlate with endoscopic and clinical progression of mucosal disease, suggesting that hydrogen ion is an important contributing factor in the progression of disease and perhaps is useful as a prognostic index.

#### Current Physiologic Studies in Progress

Little attention has previously been given to the role of pepsin digestion of the gastric mucosa following thermal injury. Currently, we are utilizing a radioimmunoassay of serum pepsinogen as a reflector of gastroduodenal morphology. A pilot study has been performed, and indicates that serum pepsinogen levels rise markedly prior to gastrointestinal bleeding in two of 11 patients studied. Although the numbers are small, we are currently evaluating the endoscopic appearance of the gastroduodenal mucosa and determining if correlations exist between the serum pepsinogen and the endoscopic appearance of the stomach.

In the experimental animal, disruption of the gastric mucosal membrane can be accomplished easily by topical application to the gastric mucosa of substances that have detergent properties, such as bile, aspirin, urea, etc. Reflux of bile through the pylorus into the stomach has been postulated as an etiologic factor in the development of erosive and ulcerative disease of the stomach. We currently are using a steroid dehydrogenase enzymatic assay to quantitate bile reflux early in the postburn period. Bile reflux will be compared between thermally injured and normal controls and will be correlated with endoscopic findings.

Since the presence of hydrochloric acid is essential in the experimental production of stress ulceration, control of hydrogen ion secretion by the gastric mucosa would seem a logical method of treatment. Histamine is felt to be the final common pathway in gastric acid secretion regardless of the stimulus employed. A new group of drugs has been developed which antagonizes the action of hist mine on the parietal cells of the gastric mucosa. Cimetidine is a third generation histamine antagonist, and is being given in a double-blind fashion to determine its effectiveness in prevention of gastroduodenal disease. This study is currently in progress.

#### Summary

Previous gastrophysiologic studies have been extensive in this unit, yet fail to define the etiologic factors solely responsible for Curling's ulcer. Future studies, as outlined above, are underway to further define the pathophysiologic events important in this disease process. In addition, various therapeutic modalities are being studied to determine the most effective in the treatment of this disease process.

#### PRESENTATIONS

None

#### PUBLICATIONS

1. Czaja AJ, McAlhany JC, Jr., Pruitt BA Jr: Acute gastroduodenal disease following thermal injury: an endoscopic evaluation of incidence and natural history. New Eng J Med 291:925-929, 1974.

2. McAlhany JC Jr, Czaja AJ, Cathcart RS III, Spicer SS, Foley FD, Pruitt BA Jr: Histochemical study of gastric mucosubstances after thermal injury: correlation with endoscopic evidence of acute gastroduodanal disease. J Trauma 15:609-612, 1976.

3. McAlhany JC Jr, Czaja AJ, Villareal Y, Mason AD Jr, Pruitt BA Jr: The gastric mucosal barrier in thermally injured patients: correlation with gastroduodenal endoscopy. Surg Forum 25:414-416, 1974.

4. Czaja AJ, McAlhany JC Jr, Andes WA, Pruitt BA Jr: Pathogenesis of acute duodenal disease after burns: role of acid secretion. Gastroenterology 68:1023, 1975.

5. Czaja AJ, McAlhany JC, Pruitt BA Jr: Acute duodenitis and duodenal ulceration after burns. Clinical and Pathological characteristics. JAMA 232:621-624, 1975.

6. Czaja AJ, McAlhany JC, Pruitt BA Jr: Gastric acid secretion and acute gastroduodenal disease after burns. Arch Surg 111:243-245, 1976.

		I AGEN	CY ACCESS	ION	DATE OF SU	MMARY	REPORT	CONTROL SYMBOL	
RESEARCH AND TECHNOLOGI	WORK UNIT SUMMARY	DA (	DE 698	1	76 10	01	ם-עם	R&E(AR)636	
A DATE PREV SUM'RY 4 KIND OF SUMMARY	S. SUMMARY SCTY	7 REGR	DING	-	W'N INSTR'N	SA SPECIFIC	DATA	. LEVEL OF SUM	
75 07 01 K. COMP	υυ	NZ	1		IL	O ves		A. VORK UNIT	
10 NO /CODES:* PROGRAM ELEMENT	PROJECT NUMBER	TASK /	REA NUM	DER		WORK UNI		R	
- PRMARY 62110A	3A162110A821	0	)0		109			·····	
S. CONTRIBUTING								S	
C. CONTRIBUTING	· · ·					• •			
11 TITLE (Procode with Soburity Classification Code)	* (U) The Zinc Requir	ement	s of	the	BUrned	Rat and	the	Influence	
of Zinc on LDH Activity	, Growth Rate and Wou	nd He	aling	:A M	lodel of	Burned	Sold	liers (44)	
12 SCIENTIFIC AND TECHNOLOGICAL AREAS	<u> </u>								
003500 Clinical Medicine	e								
13 START DATE	TA ESTIMATED COMPLETION DATE	IS FUN	NHG AGENC	Y		IS PERFORM	ANCE ME	CONTRACTOR	
73 11	76 09	DA				C. In	-Hous	e	
" CONTRACT/GRANT Not Applical	ble	18. RES	DURCES ES	-	A PROFES	HONAL MAN YR	-	HDE (In these ands)	
A DATES/EFFECTIVE:	EXPIRATION.		PRECEDIN		}				
N. NUMBER.*		FISCAL	76			.3		8	
4 TYPE	4 ANOUNT:	YEAR	CURNERY.						
& KIND OF AWARD:	I. CUM. AMT.						L		
IS RESPONSIBLE DOD ORGANIZATION		B. PERI	ORNING OF	GANIZ	TION				
HAME!" IIS Army Institute of	Surgical Research	NAME:**	IS Arm	v Tr	stitute	of Sur	gical	Research	
ob Aimy institute of	Surgical Research	i	т. Т.	y 11 ahor	atory 7	nimal B	ranch	Nesedicii	
ADDRESS:" Fort Sam Houston Th		ADDRES	" For	+ 5=	m House	on Tev	ac 7	8234	
FOIL Sam nouscon, it	EXAS /0234		FOL	6 30	un nous	JUIT, IEA	as i	0234	
		PRINCIP	AL INVESTI	GATOR	(Furnish SEAN	II V S. Academic	: Inclination	•	
RESPONSIBLE INDIVIDUAL		-	Dona	1d 3	Johns	son, MAJ	, VC		
HAME Basil A. Pruitt, Jr,	COL, MC	TELEP	HONE	512	-221-49	951			
TELEPHONE: 512-221-2720		SOCIAL	SECURITY	ACCOL					
21. C ,NERAL USE		ASSOCIA	-	GATOR				1	
FOREIGN INTELLIGENCE NO	CONSIDERED	NAME:	Ysid	ro V	/illarre	al, BS			
		NAME:						DA	
DA UNAME: DA DA UNAME: DA									
(U) Lactate dehydrogena:	se; (U) Rats			,	•	•			
TECHNICAL OBJECTIVE. 24 APPROACH, 38	PROGRESS (Furnish Individual paragraphs Ide	ntilled by	number pre	tode les	I of each with	security Classifi	tellen Cod	•.)	
23. (U) To determine th	he effect of zinc dos	age (	on tla	sma	zinc le	evels an	d the	arowth	
rate of burned rats as	an animal model for t	he z	inc re	ດນ່າ	ements	of burn	ed so	ldiers.	
				1		•• ••••			
24 (II) Groups of scale	t burned rate (60% to	tal }	odv e	urf:	ice) wi	l be fe	d a z	inc	
free dist and administer	red daily doses of zi	nc = a	otato	int	ranori	oneally	ասել տե	e animale	
will be weighed and obs	arved for signs of zi	no de	ficio	nov	Dlag	na zinc	• 14 will	bo	
will be weighed and obs	erved for sight of 21		sticie sticie	ncy. +iaa	. 1105i		WILL	ne	
decermined at 10 days po	schurn and the anima	тэег	a chailt	CT 26					
25 (11) 75 07 76 00	No signs of sine def	ini		r	oon i-	the how	- 6 a	nimala	
25. (U) $15.07 - 16.09$	NO SIGNS OF ZINC der	TCTEL	icy we	re s	een in		uea 'g	uiimdis.	
No significant differen	ces were found in the	plas	sma zi	nco	or weig	nt g <b>ain</b>	OI TI	ie animals	
receiving zinc over the	se receiving no zinc.	We:	ight g	ain	was not	t imporv	ed wi	th zinc	
administration.									
1									
5									
l									
1									
1									
l									
1									
Available to contractors upon originator's series	v61		_						

'

-

- --

. -

- --

~ •

DD FORM 1498 PREVIOUS EDITIONS OF THIS FORM ARE OBSOLETE DD FORMS 1488A I NOV 68 AND 1499 1 1 MAR 68 (FOR ARMY USE) ARE OBSOLETE 397

#### FINAL REPORT

# PROJECT NO. 3A162110A821-00, COMBAT SURGERY

# REPORT TITLE: THE ZINC REQUIREMENTS OF THE BURNED RAT AND THE INFLUENCE OF ZINC ON LDH ACTIVITY, GROWTH RATE AND WOUND HEALING: A MODEL OF BURNED SOLDIERS

## US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Donald J. Johnson, DVM, Major, VC Ysidro Villarreal, BS Harrel L. Walker, MS Arthur D. Mason, Jr., MD

Report Control Symbol MEDDH-288(R1)

UNCLASSIFIED

### ABSTRACT

# PROJECT NO. 3A152110A821-00, COMBAT SURGERY

# **REPORT TITLE:** THE ZINC REQUIREMENTS OF THE BURNED RAT AND THE INFLUENCE OF ZINC ON LDH ACTIVITY, GROWTH RATE AND WOUND HEALING: A MODEL OF BURNED SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Donald J. Johnson, DVM, Major, MC Ysidro Villarreal, BS Harrel L. Walker, MS Arthur D. Mason, Jr, MD

Report Control Symbol MEDDH-288(R1)

The objective of this study is to determine if rats with extensive scald burns have an increased requirement for the trace metal zinc. Rats were scald burned over 60% of their total body surface and administered parenteral zinc at 0, 50, and 100 micrograms per day. The animals were fed a zinc-free diet to minimize the intake of exogenous zinc. The animals were weighed preburn and at 7 and 10 days postburn. Plasma zinc levels were determined at 10 days postburn.

No significant differences were found in the postburn weights or the plasma zinc levels of animals treated with zinc and those with no zinc source. No lesions of zinc deficiency were seen.

Zinc requirements Burns Wound healing Lactate dehydrogenase Rats

# THE ZINC REQUIREMENTS OF THE BURNED RAT AND THE INFLUENCE OF ZINC LDH ACTIVITY, GROWTH RATE AND WOUND HEALING: A MODEL OF BURNED SOLDIERS

The zinc requirement of man is not fully known and the reported cases of zinc deficiency are rare." A zinc deficiency syndrome in man was reported in 1963 by Prasad in Egyptian males with a dwarfing and hypogonadal syndrome (1). Zinc has recently been reported to be essential in vitamin A metabolism. Depressed vitamin A levels in plasma were found in zinc deficient rats and in children with low plasma zinc (2). The low vitamin A levels may account for the epithelial lesions seen in zinc deficient rats and other animals.

Altered zinc levels have been reported in burn patients (3). Previously we have found low zinc levels in the plasma of burned rats with 60% total body surface burns (TBS). The tissue levels of zinc did not correlate with zinc dosage or time post burn. Lactic dehydrogenase activity was only altered during the first 48 hours post burn.

### METHODS

Male, Sprague-Dawley rats weighing between 172-200 grams were used in this study. All animals were anestheteized with pentobarbital sodium intraperitoneally (0.6 mg/100 gm body weight) and the hair clipped on the dorsum of the back and the abdomen. The animals in all groups were 3^o scald burned over 60% TBS by the method of Walker and Mason (4). All animals were fed a zinc-free diet (Nutritional Biochemical Co.) ad libitum, housed in stainless steel wire bottomed cages, and given water ad libitum. Fifty or 100 ug of zinc, as zinc acetate in saline, was administered daily intraperitoneally. Normal saline was administered to the zinc deficient animal group intraperitoneally once daily. The animals were weighed on the day of burning and at 7 and 10 days post burn. The animals were then anesthetized with chloroform to obtain a blood sample by intracardiac puncture. The blood was heparinized and plastic tubes used to store and centrifuge the samples. Plasma zinc was determined by atomic absorbtion spectrophotometry using a model AA-5 Varian Techtron.

^{1.} Prasad AS, Miale A, Jr, et al: Biochemical studies in dwarfism, hypogonadism and anemia. Arch Int Med 111: 407, 1963.

^{2.} Smith JC, et al: Zinc: A trace element essential in Vitamin A metabolism. Science 181: 954-955, 1973.

^{3.} Larson DL, Maxwekk R, Abston S, Dobrkovsky M: Zinc deficiency in burned children. Plas Reconstr Surg 46: 13-21, 1970.

^{4.} Walker HL, Mason AD, Jr: A standard animal burn, J Trauma 8: 1049-1051, 1968.

### RESULTS

The plasma zinc levels are shown in Table 1. Blood samples from 8 animals are in each group. The mean plasma zinc levels did not differ significantly between groups using analysis of variance (ANOV). Four of 8 samples in the zinc- free group and 3 of 8 samples in the 50 ug/day zinc had plasma zinc levels below 100 ug/100 ml plasma. All animals receiving 100 ug zinc daily had plasma zinc levels in excess of 100 ug/100 ml plasma. 100 ug/ 100 ml is considered the low normal value for the rat and man.

0	109.9 <u>+</u> 12*	60 - 165
50	111.3 <u>+</u> 11	70 - 160
100	142.5 <u>+</u> 28	110 - 185
	0 50 100	$\begin{array}{cccccccccccccccccccccccccccccccccccc$

#### Table 1. Mean Plasma Zinc Levels at 10 Days Postburun

All values are ug/100 ml plasma

The body weights are shown in Table 2. The animals lost weight between burning and the 7th postburn day. The administration of zinc did not result in any diminution of the weight loss which occurs early in the post burn period. The loss of weight did not differ significantly between groups. Between the 7th and 10th post burn day, the animals began to gain weight at a rate normal for unburned animals of from 3-6 gm daily. At 10 days post burn, the weights between groups did not differ significantly.

### Table 2. Mean Weight (grams + S.E.)

Group	Treatment ug/Zinc/Day	0	Days Postburn 7	10
1	0	183.5 <u>+</u> 1.04	167.3 <u>+</u> 2.4	187.0 <u>+</u> 2.4
11	50	184.3 <u>+</u> 1.94	171.6 <u>+</u> 2.4	186.5 <u>+</u> 3.8
111	100	182.6 <u>+</u> 1.44	164.6 <u>+</u> 2.2	181.8 + 2.6

#### DISCUSSION AND SUMMARY

The absence of lesions of zinc deficiency in this study and earlier studies in which animals were observed for 30 days postburn on zinc-free diets and without exogenous zinc indicates that zinc is conserved by the animals and that large zinc losses are not occurring through the burn wound and urine. The rats eat their eschar as it separates which serves as a zinc source for the animal which the burn patient would not have.

The mean plasma zinc levels in 47 unburned animals from our rat colony is 150.6 + 12.8 ug/100 ml. Individual values range from 100 - 231 ug/100 ml plasma. In a test of the diet, a group of 5 unburned animals fed the zinc-free diet for 4 months had plasma zinc levels of 37.5 - 87.5 ug/100 ml. Three of the 5 animals had lesions of zinc deficiency and in the one animal which had nc lesions, the plasma zinc level was 87.5 ug/100 ml. In plasma, one-third of the zinc is firmly bound to globulin and the remaining two-thirds is loosely bound to albumen (5). The unburned animals fed the test diet for 4 months and with lesions of zinc deficiency had plasma zinc levels compatible with a complete depletion of the loosely bound zinc. In the present study, the plasma zinc determinations were all above values which would indicate a complete depletion of the lossely bound zinc.

The age and weight of the rat when deprived of zinc may be important to produce lesions of zinc deficiency. We have not seen an immediate rapid decline in weight gain or plasma zinc in unbured animals weighing 180-200 gm. Most investigators using rats for nutritional studies use weanling rats which weight approximately 45 gm. Animals of this age and weight do not have a large zinc pool to maintain plasma and tissue zinc levels above the minimum required for normal growth and the enzymatic processes for which zinc is essential.

The negative results in producing a zinc deficiency syndrome in burned rats indicates that 180-200 gm rats have sufficient reserves to prevent complete depletion of zinc in the critical early post burn period. The administration of exogenous zinc in excess of daily requirements did not significantly alter changes in the plasma zinc levels or the weight loss and gain in the early post burn period.

PUBLICATIONS AND/OR PRESENTATIONS

None

^{5.} Prasad AS (ed): Zince Metabolism. Charles C. Thomas, Springfield; III., 1966.

C				1 AGEN		104	DATE OF SUN	MARY [®]	EPORT	CONTROL STREOL
RESEARCH AN	D TECHNOLOGY	WORK UNIT S	UMMARY	DA O	G 696'	7	76 10 0	1	DD-DI	R&E(AR)636
L DATE PREV SUM'RY 4. KI	IND OF SUMMARY	S. SUMMARY SCTY	. WORK SECURITY	7 REGR	DING	-	P'H INSTR'N	SPECIFIC D	ATA ·	. LEVEL OF SUM
76 10 01 H	. TERM	U	ប	NA		N	L		110	A WORK UNIT
18. NO./CODES:* PR	OGRAM ELEMENT	PROJECT	NUMBER	TASK A	REA HUM	BER		WORK UNIT	NUMBER	R
• PR76ARY 6	2110A	3A162110	A821	0	0		105			
S. CONTRIBUTING				ļ			· • • • • • • • •		• •	
CONTRIBUTING		<b>-</b>				╾				, de la companya de l
11. TITLE (Proceds with Securi	ity Closellisetten Codej	(U) The F	elationshi	р вет	ween	LIMD	BTOOD I	10W and	Met	apolism
FOLLOWING The	Print Injury	/ in solaie	ers (44)		~~~~~					
003500 Clinic	al Modicin	•								
THE STARY BAYE	al Medicine	14. ESTIMATED COMP	LETION DATE	SE PUNC		Y			CE MET	HOD
76 02		76 09		DA			J .	C. In-	Hous	e
TT CONTRACT/BRANT	Not Appl	cable		-	UNCES EST	-		DNAL MAN YRS	-	195 (In those ands)
A BATESPEFFECTIVE	nee oppe	EXPIRATION:			76		1.0	<u> </u>	3	5
N. HUNDER:*				FISCAL	7 <b>T</b>			3	1	i
G TYPE:		4 ANOUNT:		YEAR	CONNENT					
& KIND OF AWARD:		f. CUM, AMT.								
19 RESPONSIBLE DOD ORAA	NIZATION			30. PERI		16AHIZ/	TION			
"ANE." US Army In	stitute of	Surgical H	esearch	HALE.*	IS Arm	y In	stitute	of Surg	ical	Research
· · · · · · · · · · · · · · · · · · ·				1	Sur	- gica	1 Study	Branch		
Fort Sam	Houston, '	Texas 7823	34	ADDREN	" For	t Sa	m Houst	on, Texa	s 7	8234
										_
				HAME.		ie H	(Aulic)	ν	MSC	•
HANEL Dagil A	Druitt Tr	COL MC		TELEP	HONE 5	12-2	21-5712		100	
TELEBRONEL 512-	221-2720	, cob, me		SOCIAL	BECURITY	ACCON	NT NUMBER:			
EI. OENERAL USE	221-2120			ABOCIA	-		1 · · ·			
FORETON INTEL	LIGENCE NO	r CONSIDER	ED	-	Do	uqla	s W. Wi	lmore, M	D	
				-	Ar	thur	D. Mas	on, Jr,	MD	DA
I WANKE: Artnur D. Mason, Jr, MD DA I KEVVERKE // Second Each with Schulty Closel Scotlan Code) (U) Cardiac output; (U) Glucose; (U) Pyruvate; (U) Humans										
(U) Plethymog	maphy; (U)	Limb blood	flow; (U)	Mear	<u>skin</u>	and	body t	emperatu	res	
11. TECHNICAL OBJECTIVE.	24 APPROACH, 25	PROGRESS (Pumish in	dividual persographe ide	niilied by and n	number pro	iede (ea liem	isloom was p and re	turity classifies 1 ato tho	Van Çede m t∩	) (a)extent
23. (U) 10 de	stermine in	rdiac outru	1000 110# 11. (c)ther	mal e	anili	briu	m. and (	d)genera	1 le	vel of
or thermal my	abolism of	burned sol	diers (8-1	()) ar	nd nor	mal	control	s.		
24 (II) Limb	Blood Flow	via venous	occlusion	plet	hysmo	grap	hy usua	lly of t	otal	leg but
arms may also	be used wi	th selected	l patients.	-	-		-	-		
Cardiac Output	- Indicat	or (cardio	green) dilu	tion						
Total Body Met	abolism -	Indirect ca	alorimetry	via d	collec	tior	ofexp	ired air	in	Douglas
bags and analy	vsis on mas	s spectropl	nometer.							
Arteriovenous	sampling a	cross limb	- Analysis	for	oxyge	n co	onsumpti	on and c	arbo	n dioxide
production as	well as up	take of glu	cose and F	'FA ar	nd rel	ease	e of pyr	uvate an	d la	ctate.
Body Temperatu	ires - Rect	al and seve	eral skin t	empe:	rature	S WI	ll be m	onitored	БУ	copper-
constantan the	ermocouples	•						hwium in	ani	ot warm
General Protoc	<u>col</u> - After	post-abso	rptive subj	ect a	it the	rmal	equili	be dote	r qui	et, warm
(29°C) room, h	his resting	metabolic	rate and b	n nie	temper	acui	h and $8$	-10 limb		od flow
limb to be stu	idied Will	then be pro	followed by	care	liac c	~9±0±	st deter	mination	is (3	and
measurements t	blood asmu	3 WIII De . Jing Whenes	ver feasihl	e, J	imb bl	.000d	flow me	asuremer	its w	ill be
arteriovenous	bioon samp	ental time	estimated	at t	vo to	thre	e hours	•		-
25 (n) 76 02	$2 \rightarrow 76 \ 09 \ B$	lood flow	is increase	d on	ly to	the	burned	legs and	l the	close
relationahin h	petween leg	blood flow	w and the s	size (	of leg	bui	n sugge	st. that	. mos	st of the
increased peri	ipheral blo	od flow is	directed t	the the	e woun	nd wł	ule the	extent	of 1	local
injury is the	major dete	rminant of	limb flow.	Ter	nperat	urc	changes	of the	woun	nd will
modify leg blo	ood flow in	a predicta	able manner	- 11	ncreas	ing	tempera	ture, Ca	use Lhia	d flow
and vice versa	a. Leg oxy	gen consum	ption and s byway and r	iubst: Not el	rate t nough	data	a is avi	lable fo	br cc	omment.
Available to contractore up	un erigeneterte appro					_			_	

• •

the Harden Street Street Street Street

terry parter and press of the second

and his statistic to the second second

hated carly analysis of a

Children Athens

DD 10-P 00 1498 PREVIOUS EDITIONS OF THIS "ORM ARE OBSOLETE DD FORMS 14984 I NOV CE AND 1498 1 1 MAR 68 (FOR ARMY USE) ARE OBSOLETE

### TERMINATION REPORT

## PROJECT NO, 3A162110A821-00, COMBAT SURGERY

# REPORT TITLE: THE RELATIONSHIP BETWEEN LIMB BLOOD FLOW AND METABOLISM POLLOWING THERMAL INJURY IN SOLDIERS

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Louis H. Aulick, Ph.D., Major, MSC Douglas W. Wilmoré, M.D. Arihur D. Mason, Jr., M.D. Basil A. Pruitt, Jr., M.D., Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

### ABSTRACT

### PROJECT NO. 3A162110A821-00, COMBAT SURGERY

### REPORT TITLE: THE RELATIONSHIP BETWEEN LIMB BLOOD FLOW AND METABOLISM FOLLOWING THERMAL INJURY IN SOLDIERS

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: Louis H. Aulick, Ph.D., Major, MSC Douglas W. Wilmore, M.D. Arthur D. Mason, Jr., M.D. Basil A. Pruitt, Jr., M.D., Colonel, MC

Simultaneous measurements of resting leg blood flow and surface and rectal temperatures were performed in 45 studies of nine normals and 28 burn patients. The patients had burns from 3-86%of the total body surface with leg injury ranging from 0-87.5% of the leg surface.

In the patient group, blood flow was essentially normal in the uninjured legs, increased in a curvilinear manner with the size of the leg burn, and approached a plateau of 8.0 ml/100 ml·min as the per cent of leg burn exceeded 60%. Increasing leg surface temperature by  $5^{\circ}$ C increased blood flow to the same extent in patients with burned and unburned extremities as it did in normals. Peripheral blood flow is increased following thermal injury.

Peripheral circulation wound blood flow Thermal injury

# THE RELATIONSHIP BETWEEN LIMB BLOOD FLOW AND METABOLISM FOLLOWING THERMAL INJURY

Following successful resuscitation of burn shock, cardiac output rises above normal resting levels. The increase in total circulation is closely related to the per cent of the total body surface burned, accompanies an increase in resting metabolic rate, and may reach levels 2-3 times normal (1,2).

Indirect evidence suggests that much of the extra blood flow in the burned patient is directed to peripheral tissues. Gump, et al (3), found that splanchnic blood flow represented a smaller portion of the cardiac output in three severely burned patients than it did in normal individuals or patients with postoperative infection. Wilmore and collaborators (4) suggested that much of the elevated peripheral blood flow is directed to the skin. They found that burned patients maintained above normal surface temperatures despite increased evaporative cooling of the wound, and concluded that this elevated skin temperature could be the result of increased superficial blood flow. In addition, the coefficient of core-to-skin heat conductance, an index of skin blood flow, was twice normal in burn patients in a variety of thermal environments.

While these heat balance studies tend to support the concept of increased peripheral (surface) blood flow in the burned patient, the influence of the burn wound on the distribution of this elevated peripheral flow is unknown. Through the use of venous occlusion plethysmography, this study provides the first measurements of peripheral blood flow in thermally injured patients.

1. Baxter CR: Fluid volume and electrolyte changes of the early postburn period. In: <u>Clinics in Plastic Surgery</u> (Vol. 1, No. 4), edited by J A. Moncrief, Philadelphia, Pa.: W. B. Saunders, 1974, p. 653-709.

2. Unger A, 'aynes BW Jr: Hemodynamic studies in severely burned patients. Jurg Forum 10:356-361, 1959.

3. Gump FE, Price JB Jr, Kinney JM: Blood flow and oxygen consumption in patients with severe burns. Surg Gynec Obstet 120: 23-28, 1970.

4. Wilmore DW, Mason AD Jr, Johnson DW, Pruitt BA Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physiol 38:593-597, 1975.

### MATERIALS AND METHODS

#### Subjects

Twenty eight thermally injured patients, with a mean burn size of 39.5% of total body surface, range 3-86%, and nine normal individuals were studied. The subjects ages ranged from 15-55 years. Patients selected for this study were 1) free of any pre-existing disease prior to injury; 2) hemodynamically stable after an un-eventful resuscitation; 3) in a normal state of hydration with a hematocrit greater than 33 and without abnormalities in serum electrolyte concentrations, osmolality, or pH; 4) free of systemic infection as determined by clinical symptoms and signs and daily blood cultures; 5) 6-21 days postinjury, before significant healing had occurred; and 6) able to participate in the study. Patients represented a wide range of thermal injury, in relation to both the size of the total body surface burn (%TBS) and the extent of leg injury (%LB). The wounds were treated by a variety of techniques. The majority of patients were treated by the exposure method and received topical applications of a silver sulfadiazine cream (Silvadene^R cream) to the injured surface, but some were treated with 11% mafenide acetate topical burn cream (Sulfamylon^K cream), saline, or 5% mafenide-saturated dressings, 0.5% silver nitrate soaks, or cutaneous allograft biological dressings. The last treatment of the wound was at least 8 hours prior to study. In the nine patients treated by closed techniques, the dressings were removed only from the left leg.

### Study Design

All experiments took place in the environmental chamber described previously (4). Room temperature was maintained at 30°C, and relative humidity ranged between 40-50%. Control subjects were studied in shorts or shorts and halter, and patients were similarly draped with light cotton towels. The postabsorptive subject was moved to the study room in the early morning and placed in bed. Water was given on request and, in some patients, intravenous infusion of 0.04 molar sodium chloride solution was maintained to insure normal hydration. Nine copper-constantan thermocouples were attached to the skin at the same sites for all

^{4.} Wilmore DW, Mason AD Jr, Johnson DW, Pruitt BA Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physiol 38:593-597, 1975.

subjects (dorsum of foot, lateral and posterior calf, posterior and anterior thigh, dorsum of hand, forearm, abdomen and low back). Leg skin temperatures were monitored from both legs, using five additional thermocouples, in patients with asymmetrical leg burns. In those patients treated in dressings, the thermocouples were placed on the wound, under the dressing. A rectal probe was inserted to a distance of 10 cm from the external anal sphincter. All temperatures were recorded at 5-minute intervals throughout the study. After the subject had rested quietly for at least one hour, the left leg was inserted into a soft, pliable, water-impermeable boot, and then placed in a full-length plethysmograph. Water was added to the plethysmograph at a temperature equal to the mean leg skin temperature, and maintained at this temperature through a 30-minute equilibration period and the following 8-10 blood flow measurements. A 2-3 minute interval separated the blood flow determinations.

In selected studies, water temperature was then raised 5°C above neutral bath temperature and, following 30 minutes of equilibration, a second set of leg blood flow measurements was performed. At the end of each study, the plethysmograph was drained and the water volume determined. Total time for each study was less than three hours, and the subjects usually slept throughout the study.

### Leg Blood Flow Measurements

The validity (5,6,7), simplicity, and noninvasive character of venous occlusion plethysmography make this the optimal approach to the study of peripheral blood flow. The plethysmograph used in this study is basically a rectangular box of 6.4 mm plexiglass, slightly tapered toward the foot, with an overall length of 86 cm and a volume of 51.9 lites (Fig. 1). It consists of three sections: the thigh plate assembly with supporting rings, a full length top, and a "trough" section made up of the floor, two sides, and distal wall. A rubber gasket covered with high vacuum stopcock grease is used to obtain a watertight seal when these sections are bolted together. The top is raised at the distal end to accommodate the foot. On this elevated portion of the top section is

5. Formel PF, Doyle JT: Rationale of venous occlusion plethysmography. Circ Research 5:354-357, 1957.

6. Greenfield ADM, Whitney RJ, Mowbray JF: Methods for the investigation of peripheral blood flow. Brit Med Bull 19:101-109, 1963.

7. Landowne M, Katz EN: A critique of the plethysmographic method of measuring blood flow in the extremities of man. Am Heart J 23:644-675, 1942



. .

Ftgure 1. Leg plethysmograph.

a "chimney" (7.5 cm I.D.) which permits water displacement from 'he plethysmograph with limb swelling..

A loose-fitting, full length, boot of thin (0.05 mm) clear, polyvinyl sheeting was used to form a watertight seal around the leg when in the plethysmograph, preventing fluid exchange across the injured limb and minimizing contamination of the wound. A narrow tube of this same material is attached to the toe of this boot. The tube is passed through a small port in the top of the plethysmograph to permit air, caught between the skin and the boot, to escape as water is added. As soon as the boot is fully collapsed, the exit port is plugged with the connecting tube in place.

A circular cuff of polyvinyl sheeting at the proximal end of the boot serves to attach the boot to the thigh plate of the plethysmograph. Before the leg is placed into this boot, the proximal cuff is pulled through the opening of the thigh plate, stretched tight over a raised, grooved lip of this opening, and held in place by an O-ring. To minimize bulging of this flexible dam, the cuff is backed by three supporting rings (one 12.8 mm felt ring sandwiched between two rings of 6.4 mm plexiglass). A set of rings of various dimensions was prepared to accommodate limbs of different sizes. The supporting rings are held in place by a 6.4 mm aluminum plate bolted to the thigh plate.

The leg is slipped into the boot with the thigh plate and supporting rings attached. The thigh plate is advanced up the leg until the felt supporting ring forms a close, nonconstricting cuff around the upper thigh (approximately 15-20 cm distal to the symphysis pubis). The leg is then placed into a nylon mesh sling, and the three sections of the plethysmograph are locked into place. Throughout the rest of the study, the subject rests supine with the leg abducted about 30° at the hip and at the level of or slightly above the heart.

Water is added through a port in the bottom of the plethysmograph until it reaches a set level in the chimney. In order to standardize hydrostatic pressure across the limb, this same level is re-established prior to each blood flow determination. Water temperature is maintained at the mean leg skin temperature by a Forma refrigerated and heated bath and circulator, which circulates ethylene glyrol at a set temperature through copper tubing 'ocated in the floor of the plethysmograph. The water is continuously stirred by two small centrifugal pumps.

Venous occlusion is accomplished by rapidly inflating a 10 cm wide tourniquet cuff placed on the exposed thigh. Inflation pressure is created by filling a large air reservoir located beneath the plethysmograph. In some patients with burns on the thigh, a topical anesthetic (2% viscous lidocaine) was applied to the area under the tourniquet cuff to reduce pain resulting from inflation. The distance between the cuff and the plethysmograph is minimized but may be adjusted in order to distinguish the inflation artifact from actual limb volume changes. Venous occlusion pressures were varied in each experiment to maximize the rate of limb swelling. The usual occlusive pressure ranged from 80-140 mmHg.

Change in limb volume with venous occlusion causes water to rise in the chimney. The associated increase in column hydrostatic pressure is converted to an electrical signal (Statham Model P23BB pressure transducer), amplified (Sanborn 964 recorder), and "ecorded on a Leeds and Northrup Model 620 recorder.

Using this recording system, the inflation artifact is readily identifiable and lasts 2-3 seconds following limb occlusion. Leg volume changes between 4 and 10 seconds were used to determine leg blood flow. In the vast majority of measurements, this portion of the curve was linear; but, when this segment was curvilinear, the slope between 4-6 seconds of occlusion was read.

The plethysmograph is calibrated with the leg in place prior to the experiment and after each two or three blood flow measurements. This is accomplished by injecting known volumes of water through a three-way stopcock located on one side of the chamber.

At the end of each study, the volume of water drained from the plethysmograph is measured. Limb volume is calculated by subtracting the measured volume plus the volume of the boot from the known capacity of the plethysmograph and leg blood flow is expressed per 100 ml of leg volume. The 8-10 flow measurements are averaged to determine mean resting leg blood flow.

This plethysmograph was designed and constructed specifically for use in burn patients. The long leg model was chosen for several reasons. First, it provided for comfortable, full-length support of burned limbs in order to achieve resting blood flow measurements in this particular patient population. Second, by incorporating a major portion of the limb, the effect of variations in the extent of limb injury could be evaluated. Finally, a plethysmographic cuff was not applicable in burn limbs because variations in the wound location could affect limb expansion, thereby precluding the use of circumferential changes of one segment as being representative of volume alterations of the entire limb.

#### Body Temperature Measurement

Total body mean skin temperature  $(\overline{T}_{sk})$  was calculated by appropriately weighting nine skin temperatures according to the estimated surface area being represented (8):

 $T_{sk} = 0.143$  T anterior thigh + 0.081 T posterior thigh +

**0.168** T lateral calf + 0.105 T low back + 0.140 T forearm + 0.050 T dorsumm of hand + 0.315 T abdomen.

Mean leg skin temperature  $(T_{skl})$  was calculated in a similar manner from five leg skin temperatures:

Tsk1 = 0.365 T anterior thigh + 0.122 T posterior thigh + 0.250 T lateral calf + 0.083 T posterior calf + 0.179 T foot.

### RESULTS

Forty-five separate studies of resting leg blood flow and simultaneous measurements of rectal and surface temperatures were performed in the nine control subjects and 28 burn patients (Table 1). With an increase in the size of total body surface injury, there was a general increase in peripheral blood flow, central and surface temperatures and per cent leg burn (Table 2). The increase in leg blood flow was poorly related to total body surface burn ( $r^2 = 0.33$ ) and rectal temperature ( $r^2 = 0.27$ ), suggesting that the extra leg blood flow was not a manifestation of a generalized systemic response to injury.

Because some patients with burns as great as 50% of the total body surface had uninjured legs, the local influence of the wound on peripheral circulation was assessed by comparing a group of patients with burned legs with another group having comparable total body surface injuries but no burn on the leg under study (Table 3). The systemic response to injury was essentially the same in both groups of patients, as reflected by comparable rectal and total body mean skin temperatures. Leg blood flow and mean leg skin temperatures, however, were significantly increased by the local presence of the burn wound. Conversely, leg blood flow in the uninjured limbs of burned patients was similar to that of the normal controls.

8. Mitchell D, Wyndham CH: Comparison of weighting formulas for calculating mean skin temperature. J Appl Physiol 26:616-622, 1969.

TABLE 1. Physical characteristics of subjects, extent of burn injury, body temperatures, and leg blood flow

					Ő	um Size		Temperature	s (°C)	
Subject	Age (years)	Sex	Weight* (kg)	PBD Studied	X TBS	X Burn on Leg Studied	Rectal (T _{re} )	Mean Skin (Ī _{sk} )	Mean Leg Skin (T _{sk L} )	Leg Blood Flow** (mi/100 ml•min)
Controls										
4	24	Σ	75.0		0	0	37.2	34.2	34.1	3.29 ± 0.09
q					0	0	36.9	34.4	33.9	3.12 ± 0.06
lc					0	0	37.0	34.7	33.7	2.97 ± 0.17
8	8	X	82.7		0	0	36.5	34.0	33.7	3.45 ± 0.10
e	35	Σ	104.5		0	0	37.1	34.7	33.9	2.30 ± 0.07
4	33	Σ	79.5		0	0	37.1	34.7	33.8	2.41 ± 0.06
2	35	Σ	2.67		0	0	37.1	34.3	32.6	2.02 ± 0.05
9	28	Σ	63.6		0	0	36.9	34.2	34.0	3.06 ± 0.24
7	23	Σ	ו.וע		0	0	36.9	34.0	33.6	1.73 ± 0.16
8	23	LL.	51.4		0	0	37.1	35.1	34.8	2.83 ± 0.20
σ	22	Li.	65.9		0	0	37.3	35.0	34.5	2.91 ± 0.18

-

~

----

ŝ	
R	1
5	
Ē	
and	
s.	
L'	
rat	
ape	
<b>ب</b>	
<b>b</b> d	1
<u>م</u>	
Ę	
În ji	
F	
hu	
of	
ňt	ł
xte	
<u>ت</u> ه	
cts	
þje	
sul	
of	
ics	
Sti	ŀ
er	
act	
har	
0	
ica	ļ
hys	
ā	{
<u> </u>	
μ	
1A	ł

.

N. A.

No. Bachard Links (1997)

	Leg Blood Flou** (ml/:00 ml·min)		2.87 ± 0.10	3.24 ± 0.17	6.52 ± 0.13	4.91 ± 0.11	2.61 ± 0.18	2.53 ± 0.10	2.56 ± 0.08	5.27 ± 0.29	1.77 ± 0.07	8.39 ± 0.27	7.05 ± 0.07
s (°C)	Mean_Leg Skin (Tsk_)		34.4	35.0	35.7	35.5	33.8	34.2	34.6	35.1	33.2	36.1	35.2
Temperature	Mean Skin (T _{sk} )		35.0	35.2	35.3	35.6	34.9	35.2	34.7	36.1	34.5	36.6	35.8
	wectal (Tre)		37.1	37.4	37.4	37.9	37.9	37.6	37.8	38.0	37.6	38.7	37.7
m Size	Burn on Leg Studied		0	0	32.5	12	0	0	2.5	45	o	20	50
Bu	r TBS		m	6.5	6,5	01	12	17.5	81	19.5	23	24.5	24.5
	PBD Studied		12	7	7	9	01	12	01	18	7	Π	18
	Weight* (kg)		77.4	54.5		84.0	80.8	93.1	65.7	52.3	83.0	83.5	
	Sex		Σ	Σ		Σ	Σ	Σ	X	Σ	Σ	ŗ	
	Age (years)		ଛ	15		61	36	36	18	22	40	29	
	Subject	Burn Patients		28***	\$P	m	4	ŝ	9	7	ø	9a	<b>q</b> 6

-

409

TABLE 1. Physical characteristics of subjects, extent of burn injury, body temperatures, and leg blood flow

					78	m Size	•	Temperature	s (°C)	
Subject	Age (years)	Sex	Weight" (kg)	PBD Studied	Set a	Burn on Leg Studied	Rectal (T _{re} )	Mean Skin (T _{sk} )	Mean Leg Skin (T _{skl} )	Leg Blood Flow ⁴⁴⁶ (ml/100 ml-min)
01	16	×	62.0	2	25.5	0	38.2	37.0	35.8	3.07 ± 0.14
11a	ଷ	X	83.7	2	8	20	38.6	36.2	35.9	7.48 ± 0.19
<b>d[</b> [				21	28	20	39.1	36.8	36.9	11.73 ± 0.82
12	18	Σ	69.6	=	35	10	37.8	35.7	34.2	3.50 ± 0.13
13	33		71.8	13	8	0	38.7	35.4	34.5	3.43 ± 0.07
14a	23	X	61.8	~	41.5	45	38.8	35.9	0.7.	·5.88 ± 0.20
146				*	41.5	45	37.5	35.8	35.2	6.24 ± 0.16
15	S	X	63.4	6	43	45	38.2	36.3	36.4	8.60 ± 0.45
16 <b>a</b>	55	X	62.6	15	45	0	37.0	34.6	3.9	3.95 ± 0.11
16b				21	45	0	38.2	34.7	34.1	4.20 ± 0.18
17	25	LL.	69.5	6	45	63	37.8	36.2	35.4	7.59 ± 0.25
18	<b>\$</b>	Σ	٥٥	12	45	17.5	38.1	35.7	33.9	4.04 ± 0.06
ől	61	X	72.8	12	84	12.5	39.6	37.2	36.7	4.48.± 0.11

	eg Skin Leg Blood Flow ⁴⁺ دل (ml/100 ml-min)	.7 8.54 ± 0.11	.1 2.99 ± 0.21	.8 3.17 ± 0.41	.7 6.24 ± 0.24	.9 6.36 ± 0.09	.2 9.24 ± 0.42	.8 7.75 ± 0.15	.9 8.33 ± 0.17	.2 7.51 ± 0.13	
()•) sa	i Mean Lí (T _{si}	36	36	7	Ŕ	35	35	8	35	33	1
Temperatur	Mean Skin (T _{sk} )	36.4	36.7	36.0	36.3	36.3	36.5	37.2	36.5	33.2	
_	Rectal (Tre)	38.1	38.9	38.3	38.3	38.7	39.6	38.9	38.7	36.3	
ım Size	Burn on Leg Studied	63.0	0	o	52.5	37.5	3	77.5	8	87.5	
B	× TBS	89	23	25	23	50.5	61.5	74	78	82	
	PBD Studied	6	10	17	21	10	21	7	n	8	
	Weight* (kg)	73.1	73.2		77.2	59.1	68.2	81.6	61.1	58.4	
	Sex	16	Σ		X	X	X	Σ	Σ	Σ	
	Age (years)	19	26		28	8	51	22	24	28	
	Subject	20	21a	21b	22	23	24	25	26	27	

TABLE 1. Physical characteristics of subjects, extent of burn injury, body temperatures, and leg blood flow

*Body weights on initial study. **Mean ± S.E. ***Patient had a burned (2b) and unburned (2a) leg.

TABLE 2	?. (	General	effect	<u>s of</u>	total	body	burn	size	on	body	temperatures	and	leg
blood f	Flow	(mean :	<u>t</u> S.E.	or r	ange)								

έ.

Burn Size	Controls	<25%	25-49%	>49%
No. of Subjects	9	9	11	8
No. of Studies	11	11	14	9
Age (Years)	28 (22-35)	26 (15-40)	27 (16-55)	24 (19-28)
Weight (kg)	74.8 (51.4-104.5)	74.9 (52.3-93.1)	70.8 (61.8-88)	69.8 (58.4-81.6)
% Total Body Surface Burn	0	15.0 (3-24.5)	40,5 (25,5-48)	66.5 (50-86)
% Leg Surface Burn	0	12.0 (0-45)	29.5 (0-70)	59.5 (0-87.5)
Postburn Day Studied	_	11 (618)	12 (7-21)	14 (7-21)
Temperatures (°C)				
Rectal	37.0 ± 0.1	37.7 ± 0.1	38.3 ± 0.2	38.4 ± 0.3
Mean Skin	34.5 ± 0.1	35.4 ± 0.2	36.0 ± 0.2	35.8 ± 0.5
Mean Leg Skin	33.9 ± 0.3	34.8 ± 0.3	35.3 ± 0.3	35.0 ± 0.6
Leg Blood Flow (ml/100 ml·min)	2.74 ± 0.16	4.34 ± 0.67	5.91 ± 0.69	6.39 ± 0.72

	Controls	No Leg Burn*	Leg Burn**		
No. of Subjects	9	8	8		
No. of Studies	11	10	9		
Age (Years)	28 (22-35)	30 (15-40)	23 (15-33)		
Weight (kg)	74.8 (51.0-104.5)	70.7 (54.5-83)	68.8 (57.3-84)		
1 Total Body Surface Burn	0	25.5 (3-50)	26.0 (6.5-50)		
% Leg Surface Burn	0	0	27.5 (2.5-52.5)		
Postburn Day Studied	-	12 (7-21)	12 (6-21)		
Temperatures (°C)					
Rectal	37.0 ± 0.1	37.9 ± 0.2	38.0 ± 0.1		
Mean Skin	34.5 ± 0.1	35.4 ± 0.3	35,8±0.2		
Mean Leg Skin	33.9 ± 0.2	34.6 ± 0.3	35.4 ± 0.2+		
Leg Blood Flow (ml/100 ml•min)	2.74 ± 0.16	3.13 ± 0.21	5.89 ± 0.68 ⁺		

And the second state of the second state

 TABLE 3.
 Local effects of leg burn wound on limb blood flow and

 surface temperatures (mean ± S.E. or range)

*Patients No. 1, 2a, 4, 3, 10, 13, 16a, 16b, 21a, 21b

**Patients No. 2b, 3, 6, 7, 9a, 9b, 12, 15, 22

+Unpaired t-test with patients without leg burn, p < .05

In all subjects studied, leg blood flow increased with the extent of the leg burn (Fig. 2). The relationship appeared curvilinear: limb blood flow approaching 8.0 ml/100 ml·min as the size of the leg burn exceeded 60%.

Increasing leg surface temperature by raising bath temperature 5°C increased limb blood flow to the same extent in patients with burned and unburned extremities as it did in the normal controls (Fig. 3). This circulatory response in the burned limbs was not solely determined by vasodilatation of the uninjured skin since comparable changes in leg blood flow occurred in patients with leg burns as great as 85% of the leg surface.

#### DISCUSSION

Leg blood flow measurements in these patients clearly demonstrate that peripheral blood flow is increased during the hypermetabolic-hyperdynamic phase of thermal injury. Since this extra blood flow was closely related to the extent of local injury, and not found in the uninjured legs, the increased peripheral blood flows appear to be directed primarily to the burn wound. As observed in earlier total body measurements and confirmed by this study, elevated blood flow accounts for a higher mean leg skin temperature on burned limbs. This is most evident in the two groups of patients with and without leg burns but comparable core and total body mean skin temperatures (Table 3). Since trese two groups maintained similar levels of body heat content, the higher skin temperatures on the burned legs are the result of higher superficial blood flows. Considering the increased evaporative cooling which occurs from the burn wound (9,4), the impact of increased superficial blood flow on maintaining an elevated surface temperature is even more impressive.

Additional observations in our laboratory support the concept of wound-directed peripheral blood flow. In third degree wounds, which are associated with superficial vascular thrombosis (10), leg blood flow is near control levels shortly after injury. Flow

9. Harrison HN, Moncrief JA, Duckett JW Jr, Mason AD Jr: The relationship between energy metabolism and water loss from evaporation in severely burned patients. Surgery 56:203-211, 1964.

4. Wilmore DW, Mason AD Jr, Johnson DW, Pruitt BA Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physiol 38:593-597, 1975.

10. Order SE, Moncrief JA: <u>The Burn Wound</u>. Springfield, Ill: Charles C. Thomas, 1965.







Figure 3. The effects of a 5°C increase in water temperature on leg blood flow in normal controls (subjects 1a, 2, 3, 4), patients without leg burns (patients 1, 10, 13, 16a, 16b), and patients with burned legs (patients 3, 6, 7, 14a, 14b, 20, 25, 27). While the flow to the burned legs is elevated, the response to leg heating was the same in all subjects. Each point represents the mean surface temperature and the mean and S.E. of the leg blood flow for each group. increases to reach levels predictable from the size of leg burn by the end of the first week, associated with formation of a richly vascularized wound bed. In contrast, partial-thickness injury does not ablate the superficial vascular bed and blood flow is elevated in these legs as soon as circulatory volume is restored. In addition, leg blood flow measurements were performed in patient no. 28 prior to and five days after excision to fascia of the 32.5 leq surface wound. Leg blood flow dropped from 5.27 ml/100 ml·min to 3.33 ml/100 ml·min with the removal of the burn tissue.

the conduction is realized an interaction of the second state and the second state of the second states are second states and

Other studies suggested that increased peripheral blood flow occurs in burned patients (3) and related the increase in total body core-to-skin heat conductance to burn size (4). Patients with large burns were characterized by diminished core-to-skin insulation. This study demonstrates that the increased peripheral blood flow and heat conductance previously reported are confined to the burn wound. Patients with small burns have less of an insulative deficit, and, consequently, can store heat more effectively than extensively injured patients. Severely burned individuals usually do not increase body remperature in proportion to burn size and often maintain cooler skin and wound surface temperatures than patients with smaller burns. .Since the insulative deficit increases with burn size, but the rate of metabolic heat production reaches maximal levels (approximately twice normal) in patients with burns greater than 60% TBS injuries (11), the more extensively injured patients are extremely vulnerable to hypothermia when exposed to cool environments.

The plateau or upper limit of leg blood flow to the more extensively burned limbs does not reflect maximal vasodilation of the wound, since blood flow to these legs increased when the temperature of the surface was raised 5°C. Vascular smooth muscle tone in the granulating wound is therefore, at least, sensitive to changes in local temperature, but the additional influences of other local metabolic factors and/or central reflex activity on wound perfusion remain to be identified. Since peripheral or wound blood flow is affected by variations in surface temperature, an elevation in ambient temperature should permit the burned patient to increase wound blood flow and presumably improve healing.

3. Gump FE, Price JB Jr, Kinney JM: Blood flow and oxygen consumption in patients with severe burns. Surg Gynec Obstet 120: 23-28, 1970.

4. Wilmore DW, Mason AD Jr, Johnson DW, Pruitt BA Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physiol 38:593-597, 1975.

11. Wilmore DW, Long JM, Mason AD Jr, Skreen RW, Pruitt BA Jr: Catecholamines: Mediator of the hypermetabolic response to thermal injury. Ann Surg 180:653-669, 1974. In summary, peripheral blood flow measurements in 28 burn patients revealed that leg blood flow was increased only to the injured limbs. Leg blood flow was closely related to the extent or size of leg burn, indicating that most of the increased peripheral blood flow was directed to the burn wound. Variations in surface temperatures modified this peripheral vascular response to injury. The dominance of local wound effects on the distribution of peripheral blood flow in patients with marked systemic alterations in metabolism and thermoregulation demonstrates the priority of the wound in the body's homeostatic response to injury.

#### ACKNOWLEDGEMENT

The authors acknowledge the valuable assistance provided by Hartmut Arnhold, who designed and constructed the full-leg plethysmograph and the technical support provided by Specialists Frank Henderson and David Dahlquist.

#### REFERENCES

1. Baxter CR: Fluid volume and electrolyte changes of the early postburn period. In: <u>Clinics in Plastic Surgery</u> (Vol. 1, No. 4), edited by J. A. Moncrief. Philadelphia, Pa.: W. B. Saunders, 1974, p. 693-709.

2. Unger A, Haynes BW Jr: Hemodynamic studies in severely burned patients. Surg Forum 10:356-361, 1959.

3. Gump FE, Price JB Jr, Kinney JM: Blood flow and oxygen consumption in patients with severe burns. Surg Gynec Obstet 120: 23-28, 1970.

4. Wilmore DW, Mason AD Jr, Johnson DW, Pruitt BA Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physic: 38:593-597, 1975.

5. Formel PF, Doyle JT: Rationale of venous occlusion plethysmography. Circ Research 5:354-357, 1957.

6. Greenfield ADM, Whitney RJ, Mowbray JF: Methods for the investigation of peripheral blood flow. Brit Med Bull 19:101-109, 1963.

7. Landowne M, Katz LN: A critique of the plethysmographic method of measuring blood flow in the extremities of man. Am Heart J 23:644-675, 1942.

8. Mitchell D, Nyndham CH: Comparison of weighting formulas for calculating mean skin temperature. J Appl Physiol 26:616-622, 1969.

9. Harrison HN, Moncrief JA, Duckett JW Jr, Mason AD Jr: The relationship between energy metabolism and water loss from evaporation in severely burned patients. Surgery 56:203-211, 1964.

10. Order SE, Moncrief JA: <u>The Burn Wound</u>. Springfield, Ill: Charles C. Thomas, 1965.

11. Wilmore DW, Long JM, Mason AD Jr, Skreen RW, Pruitt BA Jr: Catecholamines: Mediator of the hypermetabolic response to thermal injury. Ann Surg 180:653-669, 1974.

PRESENTATIONS AND/OR PUBLICATIONS

None.

RESEARCH AND TECHNOLOGY WORK UNIT SUMMARY			UMMARY	DA OG 6966			76 10 01		DD-DR&E(AR)636		
76 10 01	4 KIND OF SUMMARY H. TERM	S. SUMMARY SCTY	R. WORK SECUNITY	7 REGR	ADING" A	94. 0161 NJ	rn Insta'n J	CONTRACTO		A WORK UNIT	
NO NO /CODES."	PROGRAM ELEMENT	PROJECT NUMBER		TASK /	TASK AREA NUMBER		WORK UNT NUMBER			R	
PRIMARY	62110A	3A16211	0	0	I	107					
. CONTRIBUTING						ļ.	S	<u>.</u>	<u> </u>		
CONTRIBUTING						ŀ		<u> saitai</u>	in a chuir a c	<u></u>	
(U) Intrare Intrarenal 2 SCIENTIFIC AND TEC	Distribution CHARLES AND	tribution: of Blood F	A Non-inva low in a Ma	asive odel	Tec Sim	chnique ilating	For D Burne	etermin d Soldi	ing Ti ers (4	ne 14)	
START DATE	SOUU CLINICAL MEGICINE		PLETION DATE	18. FUN		ENCY					
75 11		76 00			1		1				
CONTRACT/GRANT	Not Appliq	10 09						<u>1 C.</u>	In-Ho	18e	
	NOT APPIICA	DIC		10. HES	TAKE	DING .	A PHOPEN	HOHAL MAN Y	N 1.70	HOS (In Coursesta)	
				I BIRCAL		6 m		.5 .1		19	
G TYPE.		4 ANOUNT:		YEAR	EUNIN		<b>∤:</b>				
		L CUMA ANAT			ł		ļ				
RESPONSIBLE DOD	DRGANIZATION	1.004. 241		D. PER	PO RINKI N	6 ORGANIZA	TION				
•				{					*- <b></b> :	<u> </u>	
ADDRESS:" Fort Sam Houston, Texas 78234 PRINCIPAL INVESTIGATOR / DURY INSTITUTE OF Surgical Research Burn Study Branch Fort Sam Houston, Texas 78234 PRINCIPAL INVESTIGATOR / DURY INSTITUTE											
RESPONSIBLE INDIVIDUAL				NAME:" William S. McDougal, MAJ, MC							
NAME Basil A. Pruitt, Jr. COL. MC			TELEPHONE 512-221-4440								
TELEPHONE, 512-221-2720			SOCIAL SECURITY ACCOUNT NUMBER:								
I GENERAL USE				ASSOCIATE INVESTIGATORS							
FOREIGM INTELLIGENCE NOT CONSIDERED			D	NAME: William D. Myers, LTC, MC							
			-	HAME: Harry R. Jacobson, MAJ, MC DA							
I. KEYWORDS (Procede	RACH + In Security Classifi	cotion Code)		•					4		
(U) Intrare	nal blood dis	tribution;	(U) DMSA;	(U)	Rena	al scar	1; (U)	Dogs			
23. (U) To of blood fl	o determine a ow in a labor	non-invasi atory anim	Ne techniquial model s	ue 'fo imula	r ev ting	valuati y the h	ing int ourned	rarenal soldier	dist.	ribution	
24. (U) An as a jugula kidneys wil microsphere kidneys har counting th invasive so microsphere	imals will be ir cannula. I l be scanned. s will be inj vested and th he distribution can technique es will be sou	anestheti imercapto Followin ected intra e exact re n of injec and actual ght.	zed. A car succinic a og the term aarterially enal distrij ted micros renal blo	rotid cid w inati Th butio phere od fl	art ill on c e ar n of s. ow c	tery ca be ind of the himal w f blood A corn listril	unnula fused scanni vill th flow celatio pution	will be intrave ng labe en be s will be n betwe as det	place nously lled acrif: dete: en the ermine	ed as well y and the iced, the cmined by e non- ed by the	
25. (U) 75 2 days to 3 was used as of Dimercap to cortical analysis of but too lit altering bl	5 11 - 76 09 8 months in an 5 a control. 5 tosuccinic ac 6 medullary bl 7 microsphere 5 tle data is a 6 cod flow are	Fourteen r attempt t Preliminar id uptake ood flow i distributi s yet avai also being	ats have have have have have have have have	ad th nal b indic mal k . A ucted a mea conf	e ri lood ate idne rati kid ning irm	ight un flow, that u ey a ra io of f lneys h gful ar this r	reter 1 The sing a atio of 1.75 wa nave al nalysis method.	igated contral graphi 1.73 f s obtai so been . Othe	for per atera cal an or con ned by stud: r mean	eriods of L kidney halysis rtical Y ied ns of	

Aveileble to contractore upon originator's approval DD 1 wan as 1498 PREVIOUS EDITIONS OF "HIS PORM ARE OBSOLETE DD FORMS 1488A 1 NOV 68 AND 1498 1 1 MAR 68 (FOR ARMY USE) ARE OBSOLETE 420

### TERMINATION

PROJECT NO. 3A162110A821-00, COMBAT SURGERY

REPORT TITLE: INTRARENAL BLOOD DISTRIBUTION: A NONINVASIVE TECHNIQUE FOR DETERMINING THE INTRARENAL DISTRIBUTION OF BLOOD FLOW

> US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSION, TEXAS 78234 1 July 1975 - 30 September 1976

> > Investigators:

V. Scott McDougal, M.D., Major, MC William D. Myers, M.D., Lieutenant Colonel, MC Robert J. McAuley, M.S., Captain, MSC Robert Lull, M.D., Lieutenant Colonel, MC

Reports Control Symbol MEDDH-288 (R1)

UNCLASSIFIED
#### ABSTRACT

#### PROJECT NO. 3A162110A821-00, COMBAT SURGERY

REPORT TITLE: INTRARENAL BLOOD DISTRIBUTION: A NONINVASIVE TECHNIQUE FOR DETERMINING THE INTRARENAL DISTRIBUTION OF BLOOD FLOW

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 September 1976

Investigators: W. Scott McDougal, M.D., Major, MC William D. Myers, M.D., Lieutenant Colonel, MC Robert J. McAuley, M.S., Captain, MSC Robert Lull, M.D., Lieutenant Colonel, MC

**Reports** Control Symbol MEDDH-288(R1)

The intrarenal distribution of blood flow was determined by a new noninvasive technique. Technetium labeled dimercapto-succinic acid was injected intravenously and timed retention by the kidney recorded. Graphical analysis of the uptake curve provided values for two zones of blood flow and compared favorably to values for the cortex and juxtamedullary region as determined by microsphere injection. When the distribution of blood flow was altered by ureteral ligation, the graphical values compared favorably with those obtained by microsphere analysis.

Renal Blood Flow Dimercaptosuccinic Acid Differential Renal Blood Flow

# A NONINVASIVE TECHNIQUE FOR DETERMINING THE INTRARENAL DISTRIBUTION OF BLOOD FLOW

A curious form of renal insufficiency is periodically observed in the thermally injured patient. Glomerular filtration rate is diminished, but unlike classic acute renal failure oliguria does not occur and urinary sodium excretion is exceedingly small. The syndrome is heralded by laboratory evidence of prerenal failure in the absence of recognizable clinical causes. The avidity of the nephrons for sodium, their continued ability to secrete potassium, and the early prerenal character of the syndrome might be explained by a shift in the intrarenal blood flow from the cortex to the medulla. Thus, the nephrons with the long loops of Henle which are known to have an increased capacity for sodium reabsorption as well as urea retention are maximally perfused. The cortical nephrons would contribute little to the formation of urine, thus explaining the observed abnormalities. Supportive evidence for this hypothesis has been gained from the experimental animal in which a redistribution of renal blood flow has been demonstrated in the early postburn period.¹

Two methods are currently available for determining the distribution of renal blood flow. The first involves an intra-arterial injection of labeled microspheres followed by section of the kidney, isolation of the various zones and counting the microspheres in the tissue specimens. This technique obviously cannot be applied to patients. The second method utilizes a radioactive nuclide of an inert gas, either xenon or krypton, and involves injection of the nuclide directly into the renal artery followed by recording the disappearance of radioactivity from the kidney. The disappearance curve is stripped into four components, the first three representing specific zones of blood flow.2,3,4 The method is cumbersome, requires manipulation of an arterial cannula and does not lend itself to studying acutely ill patients who demand continuous intensive care monitoring. Therefore, if the proposed mechanism of acute renal failure in the thermally injured patient is to be substantiated, a more practical approach must be designed. The distribution of blood flow determined by a nuclide injected intravenously followed by

3. Ladefoged J, Pedersen F; Renal blood flow, circulation times and vascular volume in normal man measured by the intra-arterial injection - external counting technique. Acta Physiol. Scand. 69:220, 1967.

4. Rosen SM, Hollenberg NK, Dealy JB, Merril JP: Measurement of the distribution of blood flow in the human kidney using the intraarterial injection of 133Xe. Relationship to function in the normal and transplanted kidney. Clin. Sci. 34:287, 1968.

^{1.} Carter JG, Wells CH: Intrarenal redistribution of blood flow in the early postburn period. J Trauma 15:877, 1975.

^{2.} Ladefoged J: Renal cortical blood flow and split function test in patients with hypertension and renal artery stenosis. Acta Media Scandinavia 179:6, p 641, 1966.

a renal scan and graphical analysis would obviate many of the difficulties and risks of the washout method and would lend itself quite well to the study of critically ill patients. The purpose of this study is to determine if the nuclide technetium labeled dimercapto-succinic acid (DMSA) given intravenously followed by renal scanning and graphical analysis is an accurate method of determining the intrarenal distribution of blood flow.

#### MATERIALS AND METHODS

Preparation of the experimental animals. Eight Sprague Dawley rats were anesthetized with sodium pentobarbital (0.025 mg per gm body weight), the abdomen entered through a lower midline incision and the right ureter ligated. The animals were allowed food and water ad lib. They were reanesthetized from two to sixty days postligation with sodium pentobarbital, a jugular and superior aortic arch cannula placed, and the animals positioned under a Searle Hp Pho-Gamma camera. One millicurie of DMSA (Medi-physics) was injected intrajugularly and both kidneys scanned at 30 second intervals over a 45-minute period. At the termination of the experiment, approximately 40,000 Strontium 85 (1.0 uc) labeled microspheres (3 M Company) were injected directly into the aorta and the animal sacrificed. The kidneys were removed, weighed and placed in glutaraldehyde. Twenty-four hours later each kidney was sagittally sectioned and the cortex, juxtamedullary and medullary zones dissected Each tissue zone was weighed, the technetium allowed to decay and the strontium counted in a standard well scintillation counter (Chicage Nuclear).

<u>Graphical analysis</u>. The uptake curve conforms to the general equation  $q = C + (A - C) e^{-t/k}$  where q is the value at time t, A, the value at time 0 and C the value of the asymptote.⁵ The curve obtained for the uptake of DMSA by the kidney is of the second order and therefore the general form of the equation may be expanded to q = C + (A - C) $e^{-t/ka} + (B-C)e^{-t/kb}$  where q is the amount of nuclide contained in the kidney at time t, A the quantity of nuclide present as a function of the first component at t = 0, B the quantity of nuclide as a function of the second component at t = 0 and C the asymptote.

Several requisites must be satisfied if the uptake equation is to be related to blood flow: 1) DMSA must achieve rapid equilibrium between the blood and the renal tissue, 2) Once taken up by the tissue, significant back diffusion into the blood must not occur, 3) there should be little excretion into the tubular lumen, and 4) the partition coefficient must be the same for all renal zones. Given these assumptions, as t approaches 0, the renal vein concentration of DMSA also approaches 0 and the arterial concentration is proportional to the blood flow. If the

^{5.} Defares JG, Sneddon IN: An introduction to the mathematics of medicine and biology. Year Book Publishers, Chicago, 1960, pp. 554-552.

blood flow for specific areas differs from that in other regions, the equation will be greater than the first order and the concentration of nuclide represented by each component will be proportional to the blood flow of a particular area. Therefore, from the expanded equation when t = 0, A and B represent the components of total isotope which would be retained if all the isotope injected were presented to the kidney simultaneously and instant equilibrium were achieved. Since the components are proportional to blood flow, the percentage distribution follows directly.

If the same assumptions regarding the rate constants for inert gas washout curves are made for the uptake curves, then flow per gram tissue for the zone represented by the rate constant may be calculated by the equation  $K_n = F/W$ , where  $K_n$  is the rate constant for component n, F the flow, W the weight of the tissue and the partition coefficient between tissue and blood for the indicator.⁶

# RESULTS

The ratio of component A to component B determined by graphical analysis for seven normal kidneys was  $1.73 \pm 0.45$ . The ratio of cortical to juxtamedullary blood flow determined by microsphere injection was  $1.76 \pm 0.59$  for four normals and did not differe significantly from the ratio obtained by graphical analysis. Three ligated kidneys demonstrated a graphical ratio of  $1.47 \pm 0.23$  which did not significantly differ from the two ligated kidneys analyzed by microspheres (1.58 + 0.32). The values obtained for ligated kidneys did not significantly differ from those for normal kidneys in either the microsphere or graphical groups.

Rapid counting of three normal kidneys and bolus injection of the nuclide demonstrated a sharp increase in counts over the kidney to a maximum followed by a slight fall in the total count. As the bolus circulated and raturned to the kidney, the shar uptake followed by a moderate dip was repeated. Simultaneous measurement over the heart confirmed the bolus effect without dissipation (counts over the heart during the period when the bolus was elsewhere in the vascular system revealed values close to background). Since the peak kidney counts represent total isotope presented to the tissue and the troughs that which is retained, the ratio of the two gives the partition coefficient for DMSA for the entire kidney. Seven curves in three rats were analyzed giving a ratio of  $0.76 \pm 0.05$ .

Timed bladder scans were obtained in 8 animals, each with at least one normal kidney, and demonstrated insignificant accumulation of activity during the first 10 to 15 minutes following injection. The ratio of remal activity to bladder activity at 15 minutes was 17:1. Counts of the vascular system, bladder, background, liver and kidney revealed almost

6. Ladefoged J: Measurements of the renal blood flow in man with the ¹33Xenon washout technique. Scandinav. J. Clin. & Lab. Investigation 118:299, 1966.

### exclusive accumulation in the latter.

### DISCUSSION

Dimercapto-succinic acid labeled with technetium is rapidly taken up by the renal tissue as demonstrated by the analysis of a bolus of the agent passing through the kidney with the determination of the quantity retained and the partition coefficient. Retention by the kidney is demonstrated by the rapid fall off in radioactivity from the heart and lack of accumulation in the liver and muscular tissue. The bulk of nuclide remained in the kidney and initially, the bladder was free but after 30 minutes, a moderate amount was present. Thus, the agent satisfies the criteria of retention and insignificant dissipation during the first 10 to 15 minutes following injection. The uniformity of the partition coefficient throughout the kidney cannot be accepted with certainty since no experimental maneuvers were performed which would shed light on the question. Therefore, the criteria necessary of DMSA for determining regional renal blood flow from the uptake curve have for the most part been met.

Microsphere determination of regional blood flow is generally accepted as the most accurate method and the very close correlation of the microspheres to graphical analysis of DMSA uptake in the normal kidneys is encouraging. One method of altering renal blood flow is by obstructing the kidney.⁷ Although the correlation between injected microsheres and DMSA analysis for the obstructed kidneys was quite good, very little alteration in blood flow was obtained with ligation.

Although promising, confirmation of the usefulness of DMSA analysis for regional renal blood flow determination must come from correlations with microspheres in animals who have significant alterations in regional perfusion. Such studies are currently in progress.

#### PRESENTATIONS AND/OR PUBLICATIONS

None

^{7.} Bay WH, Stein JH, Rector JB, Osgood RW, Ferris TF; Redistribution of renal cortical blood flow during elevated ureteral pressure. Am J Physiol. 222:33, 1972.

RESEARCH	AND TECHNOLOG	WORK UNIT S	UMMARY	1. AGEN	G 696	10H [#] 2	DATE OF SUN	MARY [®]	DD-D	CONTROL SYNBOL R&E(AR)636
S. BATE PREV SUWRY	4. KIND OF SUMMARY	S. SUNMARY SCTY	. WORK SECURITY	7 REGR	DING	-	"N INSTR'N	BA SPECIFIC D	ATA-	. LEVEL OF DUE
76 10 01	H. TERM	U	U	NA	<b>L</b>	NL			H0	A. WORK WIT
18. NO /CODES:*		PROJECT	NUMBER	TASK A	REA NUM	DER		WORK UNIT	NUMBE	R
	62110A	3A16211	0A821	C	0		115			
b. CONTRIBUTING										
C. CONTRIBUTING							•			
11. TITLE (Presede with	Security Classification Co A	• (U) Gluca	agon and He	at Pr	oduct	ion	ir Norma	al Man a	nd S	eptic
Burn Patie	nts in a Mili	tary Popula	ation (44)							
12. SCIENTIFIC AND TE	CHNOLOGICAL AREAS									
003500 Cli	nical Medicin	e								
IS. START DATE		14. ESTIMATED COM	PLETION DATE	IL PUNI	NNG AGENI	CY.		116. PERFORMA Í	NCE MEI	/#00
75 07		76 09		DA			ļ	C. In-	Hous	e
17. CONTRACT/GRANT	Not Applic	able		18 RES	OURCES ES	TINATE	A PROFESSI	ONAL MAN YRS	L FU	H D& (In those ands)
A DATES/EFFECTIVE:		EXPIRATION			76	•	.7			25
b. NUMBER:*				FISCAL	<u>7</u> T				<b>_</b>	9
G TYPE:		4 ANOUNT:		TEAR			1			
. KIND OF AWARD:		LCUM. AMT		ļ	L		<u> </u>			
10. RESPONSIBLE DOD	DREANIZATION			20 PER	PORMINGO	RGANIZA	TION			
HAME!" US Army	Institute of	Surgical	Research	HAME.*	US Ar	my I	nstitut	e of Sur	gica	1 Researc
					Su	rgic	al Stud	y Branch	L	
ADDREM:* Fort	Sam Houston,	Texas 782	34	ADDRES	• Fort	Sam	Housto	n, Texas	; 78	234
				1						
1				PRINCIP	AL INVEST	IGATOR	(Funish SEAN :	IU.S. Anademic	ine i i fetie	n)
RESPONSIBLE INDIVID	JAL			HAME	• Dou	gias	W. W11	more, M.	υ.	
HANE: Basil	A. Pruitt, Jr	, COL, MC		TELEI	HONE 5	12-2	21-4/33			
TELEPHONE: 51	2-221-2720			- SOCIA	L SECURIT	ACCOU	NT NUMBER			
21. GENERAL USE				ABOCIA	TE INVEST	IGATORI				
FOREIGN IN	TELLIGENCE NO	T CONSIDER	ED		Loui	.s H.	Aulick	, MAJ, N		
				<u> </u>	Arth	ur L	. Mason	, Jr., P	1.D.	DA
AZ. KEYBORDE (Provide	RACH WIN Southly Class			-		/ <b>••</b> •	<b>T</b> 1			
(U) Calori	genesis; (U)	Gluconeoge	nesis; (U)	GIUC	ago:	(0)	Humans			
ES. TECHNICAL OBJEC	TIVE," 24 АРРИОАСН, 20	PROGRESS (Fumilar								., hast
23. (U) 1	o evaluate the	e contribu	tion of her	Datic	gruce	neoç	Jenesis	to tota.		iy neat
production	in burned ar	id burned i	nfected so	laier	s.					
										llaving
24. (U) F	ive normal me	n and four	burn patie	ents	WILL I	)e 31	uuleu L	core t		Siture
the infusi	on of 2 mg g	ucagon for	two nours.	. Ux	ygen d	const		COLE C	ember	
serum gluc	cose, insulin,	and gluca	igon whill be	e mea	surea	seri	taily tr	irouynou	C C 1	s study.
							in habl			d hurn
25. (U) 7	75 07 - 76 09	Glucagon	increased I	netab	0110 1	rate	in both	I NOTMAL		ant of
patients.	The effect a	appears dos	e related	ana o	ccurs	in v	ctwe wi	an the m	i ast	ent or a thig
glucose fi	com the liver	to the per	ciphery. G	Iucos	e ini	1510	n does i	ιοτ αυρι	ICat	e unis
effect.										
1										
1										
1										
1										
ļ										
ł										
1										

history with the second second second second second between the second second second second second second secon

in the second

ĝ

# TERMINATION REPORT

# project no. 3A162110A821-00, COHBAT SURGERY

# **NEPORT TITLE: GLUCAGON AND HEAT PRODUCTION IN NORMAL MAN AND SEPTIC** BURN PATIENTS IN A MILITARY POPULATION

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Louis H. Aulick, PhD, Major, MSC Douglas W. Wilmore, MD Arthur D. Mason, Jr., MD

Reports Control Symbol MEDDH-288(R1)

Unclassified

### ABSTRACT

PROJECT NO. 3A162110A821-00, COMBAT SURGERY

REPORT TITLE: GLUCAGON AND HEAT PRODUCTION IN NORMAL MAN AND SEPTIC BURN PATIENTS IN A HILLTARY POPULATION

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Investigators: Howard L. Aulick, PhD, Major, MSC Douglas W. Wilmore, MD Arthur D. Mason, Jr., MD

Reports Control Symbol MEDDH-288(R1)

Metabolic rate, urinary catecholamines, serum glucose, and insulin were determined serially in 23 studies in six normal men during two hours of rest followed by two hours of intravenous glucagon infusion. Glucagon doses ranged from 0-2 mg/hr. Metabolic rate increased 20% above resting levels. This relationship was: metabolic rate,  $kca1/m^2/hr = 42.2$ , dose mg/hr 0.055,  $r^2 = 0.758$ , p<0.001. Glucagon calorigenesis was unrelated to uninary catecholamine excretion rate and unaffected by beta adrenergic blockade in three subjects. Glucose, insulin, and respiratory quotient increased with glucagon administration, indicating augmented mass flow of glucose from the liver to the periphery. Glucagon calorigenesis was not related to increased cellular uptake of glucose: glucose infusion in six studies produced similar hyperglycemia and insulin responses but failed to produce comparable hypermetabolism. Glucagon infusion increased heat production, core temperature, and blood glucose in four septic burn patients. Glucagon calorigenesis is associated with hepatic glucose production and these effects are not mediated by the beta adrenergic nervous system.

Glucagon Hypermetabolism Glucose flow

### GLUCAGON AND HEAT PRODUCTION IN NORMAL MAN AND SEPTIC BURN PATIENTS

Injury or infection in man increases the mass flow of glucose from the liver to peripheral tissues, and the rate of hepatic glucose production is related to the extent of injury and degree of hypermetabolism.¹ Gram negative bacteremia in burn patients decreases both mass flow of glucose and oxygen consumption.^{2,3} This study, describing the calorigenic response to glucagon, relates this response to other glucagon-mediated metabolic events and evaluates mechanisms which could mediate the increased heat production.

### METHODS AND MATERIALS

### Subjects

Six healthy normal males participated in this study, which was performed over a four-month period (Table 1). All subjects were active, without previous disease processes, and were well acquainted with metabolic and respiratory testing. The subjects maintained their customary dietary intake and activity status throughout the fourmonth period of investigation and were rarely studied more than once a week. The night before the study, the subjects did not eat after 7:00 P.M., but were allowed to drink water until midnight.

# Procedure

The studies were performed in an environmental chamber, previously described, with ambient temperature maintained at 28°C and a relative humidity of 50 per cent. Subjects entered the room between 6:00 and 7:00 A.M. on the day of the study, changed to light cotton shorts, and rested in the recumbent position in bed. A probe of copper constantan was placed 8 cm into the rectum, and temperature was monitored at 10-minute intervals. Intravenous infusion needles were placed in large bore veins of one or both forearms for infusion and blood sampling, and patency was maintained by infusion pumps with 0.04 molar saline infusion at 60 ml/hr. After 30 to 60 minutes

1. Wilmore DW, Mason AD Jr, Pruitt BA Jr: Alterations in glucose kinetics following thermal injury. Surg Forum 25:81-83, 1975.

2. Wilmore DW, Mason AD Jr, Pruitt BA Jr: Altered glucose flow in septic burn patients. In press, Surg Gynec Obstet.

3. Wilmore DW, Long JA, Skreen R, Mason AD Jr, Pruitt BA Jr: Catecholamines: mediator of the hypermetabolic response to thermal injury. Ann Surg 180:653-668, 1974. CHARACTERISTICS OF SUBJECTS STUDIED

TABLE 1

ł.

の日本になっていたが

When a straight a with the solution with the second state of the

A CONTRACTORY

2

•

	Respiratory Quotient	0.88 ± 0.01	0.84 ± 0.01	0.84 \$ 0.01	0.82 ± 0.02	0.79 ± 0.01	0.80 ± 0.02	0.83
n	V02 (L/Min)	0.197 ± 0.003	0.236 ± 0.002	0.246 ± 0.002	0.274 ± 0.001	0.257 ± 0.004	0.240 ± 0.003	0.242
sal Measurement (Mean ± S.E.)	Metabolic Rate (kcal/m²/hr)	36.1 ± 0.6	34.3 ± 0.3	35.1 ± 0.2	36.5 ± 0.3	36.8 ± 0.5	35.8 ± 0.3	35.8
2	lnsulin (וש/שע)	10.6 ± 1.3	7.2 ± 1.4	7.6 ± 0.5	7.0 ± 2.7	8.2 ± 0.7	8.9 ± 0.3	8.2
	Serum Glucose (mg/100 ml)	86 ± 3	1 = 16	88 ± 1	83 ± 3	87 ± 1	86 _{, ±} 1	87
	BSA (m ² )	1.61	2.00	2.05	2.18	2.00	1.93	1.96
	Weight (kg)	51	74.5	78.0	88.6	76 <b>.</b> 8	1.62	76.0
	Age (Years)	21	37	34	26	24	ʻ 25	28
	Subject	-	7	s	4	Ś	9	re er

of recovery following venopuncture, the subjects voided, the time was recorded and designated as the start of the experiment. In the first **infusion study, the metabolic** rate of the subject was determined hourly for four hours. The four-hour study period was divided into a two-hour control period, followed by a two-hour experimental period. At the end of the two-hour control period, one intravenous infusion was replaced with a freshly prepared solution of 0.04 molar saline containing glucagon, with doses administered ranging from.0.05 mg to 2 mg per hour, with the infusion rate maintained at 60 ml per hour. Analysis of the solution by specific glucagon antibody confirmed the glucagon concen**trations in the infusate were within 10 per cent of the specified dose. Control studies were** performed in all subjects with the saline infusion maintained throughout the four hours and no glucagon given. The order of the dose of glucagon infused and control studies were randomized in all subjects. In addition, three subjects received a glucose infusion during the experimental period to evaluate the effect of glucose administration on metabolic activity. Venous blood samples were drawn for serum glucose and insulin concentration at the end of each hour of the control period and at 30-minute intervals throughout the experimental period. All subjects voided at the end of the twohour control and two-hour experimental periods, and the volume was **measured and an aliquot** acidified and frozen for catecholamine analysis.

A similar protocol was used to evaluate the effect of beta blockade on glucagon calorigenesis in three subjects. At the end of the two-hour control period, 5 mg of propanolol were administered intravenously as a loading dose, followed by a constant infusion of 0.08 mg propanolol per minute, which was given along with a steady infusion of 0.5 mg glucagon per hour. The efficacy of beta blockade was determined at the end of the experiment by the inability of subjects to respond to beta stimulation (2 ug isoproterenol IV) by increasing pulse rate, or demonstrating a rise in plasma free fatty acids.

#### Analytical Techniques

「ためたい」とない。「「「「たい」」となって、「「」」となっている。

ないために、いていたので、いたいからない

Gas exchange of cxygen and carbon dioxide was determined hourly by the open-circuit technique previously described.³ Two Douglas bags were collected over 10 minutes and the results of the two measurements averaged. Serum glucose, insulin and glucagon, and urinary catecholamines were determined by methods previously described.³

3. Wilmore DW, Long JA, Skreen R, Mason AD Jr, Pruitt BA Jr: Catecholamines: mediator of the hypermetabolic response to thermal injury. Ann Surg 180:653-668, 1974.

# Patients Studied

Four individuals with gram negative bacteremia and hypothermia were infused with two mg'glucagon per hour for a two-hour period. Core temperature was monitored in all patients by a rectal thermistor and metabolic rate and serum glucose were measured in three of the four individuals before and at the end of the study period.

# RESULTS

#### Metabolic Rate

Basal metabolic rate was within 3 per cent of predicted metabolic rate for trained subjects.⁴ Variations in a Douglas bag measurement during the control period or during the saline infusion did not exceed more than 6 per cent for any single subject studied over the two-hour basal or four-hour control periods. Maximum study-to-study variation in basal metabolic rate in the same subject did not exceed 10 per cent for any single individual studied over the four-month period.

Basal metabolic rate increased in a curvilinear manner with increasing doses of glucagon (Fig. 1). This metabolic effect was first evident at the dose of 0.25 mg glucagon per hour. Metabolic rate plateaued at approximately 20 per cent above basal levels of heat production. No significant difference in glucagon calorigenesis was noted between the first and second hour following glucagon infusion. Oxygen consumption for the first hour during all glucagon infusions was  $262 \pm 8$  ml/min and  $263 \pm 8$  ml/min for the second hour. Mean respiratory exchange ratios (RQ) moved from  $0.83 \pm 0.01$  during the basaí period to  $0.90 \pm 0.01$  during the first hour postinfusion, and  $0.90 \pm 0.01$  during the second hour. No significant alterations were observed in rectal temperature at any level of glucagon infusion.

# Altered Substrate Flow

Accompanying the alterations in metabolic rate were elevations in respiratory_exchange ratio and serum insulin and glucose concentrations. Hyperglycemia and insulin response were highly variable. In general, the early elevation of serum glucose correlated positively with increasing glucagon doses, but the hyperglycemic response was masked by the stimulation of insulin with glucagon doses >0.5 mg/hr. Glucose infusion in an intermediate (12 g/hr) and large dose (36 g/hr)

4. Fleisch A: Le metabolisme basal standard et sa determination au moyen du "Metabocalculator," Helv Med Acta 18:23, 1951.



Ftgure 1. Metabolic rate increases in a curvilinear manner but does not exceed a 20% increase above basal (left). The dose response curve is observed when expressing per cent change in basal metabolism as a function of the log of the glucagon dose (right).

resulted in hyperglycemia and increased insulin elaboration, and was associated with an increase in respiratory exchange ratio. However, glucose infusion failed to produce comparable levels of heat production attained with glucagon doses which resulted in similar alterations in glucose and insulin concentration (Table 2).

#### Role of Catecholamines

and the state on a distribution of the state 
Sympathetic activity, as quantitated by urinary excretion rate of catecholamines, was not related to glucagon calorigenesis. The mean excretion rate of urinary catecholamines was  $5.6 \pm 0.6 \,\mu\text{g}$  per two hours during the control period,  $6.7 \pm 0.8 \,\mu\text{g}$  per two hours during all periods of glucagon infusion, and  $6.2 \pm 0.8 \,\mu\text{g}$  per two hours during the infusion of glucagon doses of  $0.25 \,\mu\text{mg}$  per hour or greater (doses associated with increased heat production). These differences were not significant. Propanolol administration in doses known to adequately block metabolic responses to beta adrenergic stimulation did not significantly alter increased calorigenesis, which occurred following the 0.5 mg per hour glucagon infusion. Similarly, alterations in glucose or insulin response were not observed by beta adrenergic blockade (Fig. 2).

#### Patient Studies

Core temperature increased with glucagon infusion in all patients (Table 3). This was associated with an increase in metabolic rate and a rise in blood glucose, similar to the alterations that were observed in normal individuals.

#### DISCUSSION

Glucagon calorigenesis has been described in both animals and man, although the exact mechanisms of the increased heat production were not known. In small animals, glucagon administration stimulates epinephrine elaboration, and it was suggested that the increased calorigenesis was mediated through the sympathetic nervous system.⁵ However, in this study, urinary catecholamines failed to show any elevation in this neurohormonal transmitter following glucagon administration, and heat production was unaffected with beta blockade. Tests of thyroid function have been performed following glucagon administration in man, and thyroid hormone is not increased following infusion

5. Davidson IWF, Salter JM, Best CH: The effect of glucagon on the metabolic rate of rats. Am J Clin Nutr 3:560, 1960.

1
6
F
N

METABOLIC EFFECTS OF GLUCAGON ADMINISTRATION AND GLUCOSE INFUSION

	CONTROL	EXPERIM	ENTAL
Gludose Infusion (36 g/hr), N = 2			
Serum glucose	86 ± 2	220 ± 44	285 ± 7
Insulin	6.3 ± 0.9	15.8 ± 1.8	35.0 ± 2.2
Metabolic Rate	35,4 ± 0.0	37.2 ± 1.0	37.5 ± 1.7
Respiratory Quotient	0.76 ± 0.0	0.84 ± 0.04	0.86 ± 0.04
Clucagon infusion $(0.25 \text{ mg/hr})$ , N = 4			
Serun 61 ucose	93 ± 2	155 ± 16	118 ± 12
Insu) in	8.4 ± 1.1	77.8 ± 27.9	40.4 ± 19.4
Metabolic Rate	35.9 ± 0.7	39.2 ± 1.5	40.3 ± 0.2
Respiratory Quotient	0.82 ± 0.02	0.90 ± 0.04	0.90 ± 0.02





. EFF	ECT 0	F TWO-H	OURS OF G	LUCAGON	INFUSION	GIVEN AT	A DOSE OF	2 mg/HOU	77
				Hetabol (kcal/	ic R <b>ate</b> m²/hr)	Rectal Ter	mperature C)	Blood 6 (mg/10)	lucose 0 mì)
Patients	Age	ະ ສູ່ກອງ Burn	P'åD Studied	Before	After	Before	After	Before	After
	20	63	4	45.1	54.2	36.5	37.0	110	220
2	17	69	28	55.9	68.2	35.8	36.5	150	246
w	20	65.5	15	36.2	55.1	36.2	36.6	312	246
4	33	46.5	39	1	I	36.7	37.0	124	249

-
0
Ť.
2
Ő.
÷
ō
Ē
2
<b>.</b>
2
ß
ň
>
୍ର 🏾
ž
-
=
ç.
2
0
z
a
-
$\sum_{i=1}^{n}$
z.
2
_
>
0
ð
ŝ
1
0
-11
N
_
E.
イ
Ξ
2
7

TABLE 3

of this hormone.⁶ The possibility that the calorigenic effect was related to the mass flow of glucose into the peripheral tissues was also evaluated in this study. Although a slight increase in metabolic rate was noted during glucose infusion, the specific dynamic effect of glucose is small, and does not account for the total elevation of oxygen consumption which was observed in the normal subjects.

Glucagon acts specifically on the liver to enhance gluconeogenesis. Previous studies in severely injured patients reveal that the mass flow of glucose from liver to peripheral tissue is correlated closely with heat production, thus reflecting energy costs of gluconeogenesis. That heat production increases with the translocation of glucose from liver to peripheral tissue is observed by the increase in heat production which is accompanied by a rise in serum insulin and respiratory quotient. Diminished mass flcw of glucose occurs in septic injured man, but heat production, core temperature, and mass glucose flow are increased and return toward preseptic levels with the infusion of glucagon, thereby supporting the concept that the liver is the primary target organ which fails during gram negative endotoxemia. Alterations in circulation may affect liver function in septic man and glucagon may improve cardiac output, peripheral perfusion, and organ function in the patients studied. However, it is unlikely that increased support of splanchnic circulation alone accounted for the calorigenic effect of this hormone because a consistent rise in metabolic rate was observed in the normals following glucagon administration.

It should be noted that the maximum rates of heat production following a pharmacologic dose of glucagon infusion were relatively small. With infusion doses which mimicked portal glucagon levels observed in fasting man (0.05 mg/hr), no significant change in metabolic rate was observed. Burn patients have glucagon levels which are much higher than those observed in normal or fasting man, and this hormone may augment the heat production which occurs following activation of the adrenergic nervous system in injured man. The physiologic significance of this calorigenic effect of glucagon is yet to be defined.

6. Trisotto A, Busnardo B, Frezzato S, Girelli ML, Federspil G: Effect of glucagon on oxygen consumption in man. In <u>Current Topics</u> on <u>Glucagon</u>, edited by M. Anstoni, et al, Padoya, 1971, p. 193.

7. Wilmore DW, Lindsey CA, Moylan JA, Faloona GR, Pruitt BA Jr, Unger RH: Hyperglucagonemia after burns. Lancet 1:73-75, 1974. REFERENCES

1. Wilmore DW, Mason AD Jr, Pruitt BA Jr: Alterations in glucose kinetics following thermal injury. Surg Porum 26:81-83, 1975.

2. Wilmore DW, Ma.on AD Jr, Pruitt BA Jr: Altered glucose flow in septic burn patients. In press, Surg Gynec Obstet.

3. Wilmore DW, Long JA, Skreen R, Mason AD Jr, Pruitt BA Jr: Catecholamines: mediator of the hypermetabolic response to thermal injury. Ann Surg 180:653-668, 1974.

4. Fleisch A: Le metabolisme basal standard et sa determination au moyen du "Metabocalculator," Helv Med Acta 18:23, 1951.

5. Davidson IWF, Salter JM, Best CH: The effect of glucagon on the metabolic rate of rats. Am J Clin Nutr 3:560, 1960.

6. Trisotto A, Busnardo B, Frezzato S, Girelli ML, Federspil G: Effect of glucagon on oxygen consumption in man. In <u>Current Topics</u> on Glucagon, edited by M. Anstoni, et al, Padova, 1971, p. 193.

7. Wilmore DW, Lindsey CA, Moylan JA, Faloona GR, Pruitt BA Jr, Unger RH: Hyperglucagonemia after burns. Lancet 1:73-75, 1974.

# PRESENTATIONS

Aulick LH. "Mechanisms of Glucagon Calorigenesis," presented at the American Physiological Society meeting (Federation of American Societies for Experimental Biology), Anaheim, California, 10 April 1976.

#### PUBLICATIONS

Aulick LH, Wilmore DW, Mason AD Jr. Mechanisms of glucagon calorigenesis. Fed Proc 35:401, 1976.

RESEARCH	I AND TECHNOLOG	Y WORK UNIT S	UMMARY	1 AGEN	OF (	6399	2. DATE OF N 76 10	76 10 C1 DD-DR&E		
1 DATE PREV SUMPRY	4. KIND OF SUMMARY	5. SUMMARY SCTY	A. WORK SECURITY	7. REOR	ADING#	94. D	GO'N INSTR'N	BA SPECIFIC	ACCESS	D LEVEL OF SUN
10 NO (CODE:								IN YES	U #0	~
A BRIMARY	621103	221621107		TASK			100	WORK UN	IT NUMBER	R
	0211UA	JAI62IIUA	4021	<u> </u>	0		106		<u>,</u>	
C. CONTRIBUTING				<u> </u>				•••••	• •	
1 TITLE (Protode with	Security Classification Code	1 [*] (11) Domos	alined Con	1		in the	E ificio de la			
Tubalation	Tudowe in a l	(U) Aeros	ollzed Gen		in :	in in	e propny	lactic	Treat	ment or
12. SCIENTIFIC AND TE	THILLY THE A	MILICALY PC	pulation (	44)						
003500 Cli	nical Medicin	<u> </u>								
13. START DATE		14. ESTINATED CON		IS. PUN	DING AC	ENCY		14. PERFOR	MANCE MET	HOD
74 07		76	6 0 9	A	1		1	C. Tr	Hone	A
T CONTRACT/GRANT	Not Appli	rable					-	1 0	T.	
A DATES/EFFECTIVE:	not ippit	EXPIRATION			INNEE!	I ESTIMATI	E & PROFES	SIONAL MAN Y	RS & PUI	IDE (In those and a)
b. summer."				FIRCAL	1 4	6 T		.4		15
G TYPE		4 AMOUNT.		YEAR	COMM			·		
S KIND OF AWARD		f. cupa anav		1						
IS RESPONSIBLE DOD	ORGANIZATION	1. COM. AMT.		20. PERI	FORMIN	GORGANI				
	Tu	ـــــــــــــــــــــــــــــــــــــ		1	••• •			L		
Burn Study Branch										
ADDRESS:" Fort Sam Houston, Texas 78234										
Fort	Sam Houston,	Texas 782	234	AUURES	• F	ort S	am Houst	ton, Tex	kas 7	8234
				PRINCIP	AL INV	ESTIGATO	R (Fumish SEAN	IIUS, Asadem	is institution	U
RESPONSIBLE INDIVID	JAL A Druitt Tr	COT NO		NAME	• Ba	rry A	. Levine	e, MAJ,	MC	
NAME DOSIL	A. PLUICC, JI	, COL, MC		TELEP	HONE	512	-221-460	04		
TELEPHONE: DL	2-221-2720			50CIAI	SECUI	RITY ACCO	UNT-NUMBER			
DODUTINE TH			-	ASSOCIA	TE INV	ESTIGATO	RS .			
FOREIGN IN	TELLIGENCE NO	r considere	sD	NAME:	Hug.	h D.	Peterson	h, COL,	MC	
				NAME.	Cle	ment	L. Slad	e, CPT,	MC	DA
			Humans; (U	) inr	nata	tion	injury;	(0) Pre	eumonı	a;
(U) Prophy	laxis; (U) Ge	ntamicin;	(U) Aerosol	1 zed		. <u> </u>				
23 TECHNICAL OBJECT	TIVE, 24 APPROACH, 28	PROGRESS (Pumlah li	ndividual paragraphs id	entitled by	number	Prezede li	ezt of each with	Security Classi	Restian Cade	
23. (U) T	he purpose of	this expendence	ciment is t	o det	cerm	ine i	f aeros	olized g	gentam	icin is
an effecti	ve agent in p	reventing p	pneumonia i	n bui	rned	sold	iers who	o have s	sustai	ned
inhalation	injury.									
24. (U) T	his will be a	double-bli	ind study.	The	pat	ients	will b	e divid	ed int	o treat-
ment and c	ontrol groups	. The grou	ups will be	subo	livi	ded b	y age a	nd burn	size.	To ente
the study	the patient m	ust have e	ither a xen	on so	can	or fi	beropti	c broncl	noscop	ic
evidence o	f inhalation	injury. Al	ll patients	will	l re	ceive	an aer	osolized	d agen	t, either
gentamicin	3.0 mg/kg/24	hrs divided	l into 3 do	ses o	or n	ormal	saline	on the	same	schedule.
Therapy co	ntinues for 1	0 days fol	lowing iniu	rv oi	r ນກ	til a	ກບຸກນາກ	onary i	nfiltr	ates have
cleared	Teukens tube	miltures w	ill be take	n on	al+	ornat	o dave	in an a	ttomnt	+0
identify +	he emergence	of recietar	nt etrain.	of h	40±0	ria	c uuyo	un a	ccempt	
THENCITLY C	The emergence	or restardi	it otratilo	<b>2</b> 10	~~					
25 /11) 7	6 01 - 76 00	20 mandan		l nati	i ont		0 0011 0	ntored	the et	udu No
25. (0) /	0 UI - 70 UY	20 random.	ry serected	pat:	rent F Fr	s nav		ncerea	cne st	.uuy. NO
conclusion	s can be draw	n rrom this	s small gro	up at	u un	IS TI	me. Th	e code :	remain	15
unbroken a	na it is not	possible to	o identity	τωο (	aist	inct	groups	on the l	Dasis	or
clinical c	ourse.									

delandin ayahan katalah nangari di kata sa kat

#### **TERMINATION**

# PROJECT NO. 3A162110A821-00, COMBAT SURGERY

# REPORT TITLE: AEROSOLIZED GENTAMICIN IN THE PROPHYLACTIC TREATMENT OF INHALATION INJURY

US ARMY INSTITUTE OF SURGICAL RESEARCH BROOKE ARMY MEDICAL CENTER FORT SAM HOUSTON, TEXAS 78234

1 July 1975 - 30 September 1976

Investigators:

Barry A. Levine, MD, Major, MC Hugh D. Peterson, DDS, MD, Colonel, MC Robert B. Lindberg, PhD Basil A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

UNCLASSIFIED

#### ABSTRACT

PROJECT NO. 3A162110A821-00, COMBAT SURGERY

**REPORT TITLE:** AEROSOLIZED GENTAMICIN IN THE PROPHYLACTIC TREATMENT OF INHALATION INJURY

US Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Period covered in this report: 1 July 1975 - 30 5eptember 1976

Investigators: Barry A. Levine, MD, Major, MC Hugh D. Peterson, DDS, MD, Colonel, MC Robert B. Lindberg, PhD Basil A. Pruitt, Jr., MD, Colonel, MC

Reports Control Symbol MEDDH-288(R1)

Inhalation injury associated with cutaneous thermal injury, and the airborne pne_monia which frequently occurs as a complication, remain clinical problems for which there is no satisfactory treatment. Despite the use of various mechanical devices for pulmonary toilet and respiratory support, the prognosis for an inhalation injury associated with a large cutaneous thermal injury remains dismal. Although the early problems, e.g. laryngeal edema, bronchospasm, and pulmonary edema, can be treated, airborne pneumonia occurs with regularity and often with a fatal outcome.

It has been suggested that aerosolized antibiotics have a role in the treatment of airborne pneumonias. In various studies, aerosolized polymixin and gentamicin used prophylactically have been shown to reduce significantly the incidence of airborne pneumonia in certain groups of high risk patients. All observers have described the emergence of resistant strains of bacteria during such treatment.

This study's purpose is to investigate the effect of gentamicin, used prophylactically, on the course of inhalation injury. The questions asked are:

1) is overall survival affected?

2) is the incidence of airborne pneumonia affected?

3) Does the bacterial flora of the respiratory tract change and do resistant bacterial strains emerge?

This is a prospective, double blind study. The patients are divided into treatment and control groups. These groups are further subdivided by age and burn size. To enter the study, a patient must have evidence of inhalation injury proven by xenon scan and/or bronchoscopy. All patients receive an aerosolized agent, either gentamicin (3.0mg/kg/24 hours divided into 3 doses) or normal saline on the same schedule. Therapy continues for 10 days following injury or until pulmonary infiltrates have cleared. Luken's tube cultures of bronchial secretions are taken on alternate days in an attempt to identify the emergence of resistant strains of bacteria.

There have now been 20 patients randomized into the study. No conclusions can be drawn from so small a group and more patients will be studied before the code is broken and results analyzed.

Inhalation injury Pneumonia Prophylaxis Gentamicin Aerosolized

### PUBLICATIONS

huishedt. Die seuschwiegen werde sternen in

the street of the the restriction and the street street with the street street the street s

and a start of the second and the start of the second and the second second sectors of the second second second

### 1 July 1975 - 30 September 1976,

Czaja ÅJ, McAlhany JC, Pruitt BA, Jr: Acute duodenitis and duodenal ulceration after burns. Clinical and pathological characteristics. JAMA 232: 621-624, (May 12) 1975. (Not received until July)

Czaja AJ, McAlhany JC, Jr, Andes WA, Pruitt BA, Jr: Acute gastric disease after cutaneous thermal injury. Arch Surg 110: 600-605 (May) 1975. (Not received until July).

Czaja AJ, Rizzo TA, Jr, Smith WR, Jr, Pruitt BA, Jr: Acute liver disease after cutaneous thermal injury. J Trauma 15: 887-894 (Oct) 1975.

Merrill RH: Acute venous hemodialysis using the unipuncture apparatus. Am J Surg 132: 410-414 (Sep) 1976.

Wilmore DW: Alimentation in injured and septic patients. Heart & Lung 5: 791-792 (Sep-Oct) 1976.

Wilmoro DW, Mason AD, Jr, Pruitt BA, Jr: Alterations in glucose kinetics following thermal injury. Surg Forum 26:81-83, 1975.

Wilmore DW, Orcutt TW, Mason AD, Jr, Pruitt BA, Jr: Alterations in hypothalamic function following thermal injury. J Trauma 15:697-703 (Aug) 1975.

McAlhany JC, Jr, Czaja AJ, Pruitt BA, Jr: Antacid control of complications from acute gastroduodenal disease after burns. J Trauma 16: 645-649 (Aug) 1976.

Pruitt BA, Jr, Mason AD, Jr, Hunt JL: Burn injury in the aged or high-risk patient, In The Aged and High Risk Surgical Patient (Ed, JH Siegel, PD Chodoff). Grune & Stratton, Inc. 1976, pp -523-546.

Pruitt BA, Jr, Lindberg RB, McManus WO: Bacteriology, antibiotics, and chemotherapy, In Hand Surgery, 2nd ed. (Ed, E Flynn), The Williams & Wilkins Co. Baltimore, 1975, pp 451-480.

Salisbury RE, Pruitt BA, Jr: Burns of the Upper Extremity. Vol. XIX in series, Major Problems in Clinical Surgery (Consulting Editors, J E Dunphy, Paul A. Ebert), W B Saunders Co., Philadelphia, 1976.

Long JM III, Wilmore DW, Mason AD, Jr, Pruitt BA, Jr: Comparison of carbohydate and fat as caloric sources. Surg Forum 26:108-110, 1975.

Levine NS, Lindberg RB, Salisbury RE, Mason AD, Jr, Pruitt BA, Jr: Comparison of coarse mesh gauze with biologic dressings on granulating wounds. Am J Surg 131: 727-729 (June) 1976.

Levine NS, Salisbury RE, Peterson HD, Pruitt BA, Jr: Clinical evaluation of the

carbon dioxide laser for burn wound excisions: A comparison of the laser, scalpel, and electrocautery. J Trauma 15:800-807 (Sep) 1975.

Wilmore DW, Mason AD, Jr, Johnson DW, Pruitt BA, Jr: Effect of ambient temperature on heat production and heat loss in burn patients. J Appl Physiol 38: 593-597 (Apr) 1975. (Reprints not received until July).

Levine NS, Salisbury RE, Mason AD, Jr: Effect of early surgical excision and homografting on survival of burned rats and of intraperitoneally-infected burned rats. Plast Reconstr Surg 56: 423-429 (Oct) 1975.

Levine NS, Salisbury RE, Seifter E, Walker HL, Mason AD, Jr: Effect of Vitamin A on tumor development in burned, unburned, and glucocorticoid-treated mice inoculated with oncogenic virus. Experientia 32: 1309-1312 (Nov 15) 1975.

Taylor JW, Hander EW, Skreen R, Wilmore DW: The effect of central nervous system nacrosis on the sympathetic response to stress. J Surg Res 20: 313-320, (Apr) 1976.

Hunt JL, Agee RN, Pruitt BA, Jr: Fiberoptic bronchoscopy in acute inhalation injury. J Trauma 15: 641-649 (Aug) 1975.

Czaja AJ, McAlhany JC, Pruitt BA, Jr: Gastric acid secretion and acute gastroduodenal disease after burns. Arch Surg 111: 243-245 (Mar) 1976.

McAlhany JC, Jr, Czaja AJ, Cathcart RS, III, Spicer SS, Foley FD, Pruitt BA, Jr: Histochemical study of gastric mucosubstances after thermal injy: Correlation with endoscopic evidence of acute gastroduodenal disease. J Trauma 15: 609-612 (Jul) 1975.

**Goodwin MN, Jr, Raulston GL, Pruitt BA, Jr:** Hypertrophic scar in the experimental animal. In The Ultrastructure of Collagen (ed. JJ Longacre). Charles C. Thomas, Publishers, Springfield, IL, 1976.

Wilmore DW, Mason AD, Jr, Pruitt BA, Jr; Insulin response to glucose in hypermetabolic burn patients. Ann Surg 183: 314-320 (Mar) 1976.

Merrill RH, Tasch R: Iron absorption in hemodialyzed patients. Transactions, ASAIO, XXII: 69-72, 1976.

Nash DA, Jr, Merrill RH: Kidney transplanation in the active duty soldier: Case report. Mil Med 151: 545-546, 1976.

Levine NS, Peterson HD, Salisbury RE, Pruitt BA, Jr: Laser, scalpel, electrosurgical, and tangential excisions of third degree burns. A preliminary report. Plast Reconstr Surg 56: 286-296 (Sep) 1975.

Pruitt BA, Jr: Management of fluid and electrolyte disturbances in burn patients. In Monograph published by Searle Library of Electrolytes.Searle & Co. San Juan PR, 1976, pp 1-25. Hunt JL, Pruitt BA, Jr: Management of thermal injury. In Traumatic Surgery, Medical Ou'line Series (Ed, NE McSwain, Jr). Medical Examination Publishing Co, Inc, Flushing, NY, 1976, pp 131-153.

Fruitt BA, Jr, Long JM, Agee RN: Management of burn injuries. In Current Management of Trauma in Surgery and General Practice (Ed. T. Matsumoto). Charles C. Thomas, Springfield, IL, 1975 pp 166-183.

Manual on Control of Infection in Surgical Patients. Committee on Control of Surgical Infections of the Committee on Pre-and Postoperative Care (Subcommittee, WA Altemeier, JF Burke, BA Pruitt, Jr, WR Sandusky). Am Coll Surg, Chicago, 1976.

Andes WA: Normoblastemia after thermal injury. Am J Surg 131:725-726 (June) 1976.

Pruitt BA, Jr: Nutrition for the hypermetabolic burn patient. In Proc of International Symp on the Treatment of Burns, Beverwijk, Netherlands, 1975, pp 360-400.

Pruitt BA, Jr: Postburn hypermetabolism and nutrition of the burn patient. In Manu. I on Nutrition in Surgical Patients, Fhiladelphia, WB Saunders Co, 1975, pp 295-412.

--

Wilmore DW, Pruitt BA, Jr: Parenteral nutrition in burn patients, InTotal Parenteral Nutrition (ed., J Fisher). Little, Brown & Co., Boston, 1976, pp 231-252.

Hunt JL, Mason AD, Jr, Masterson TS, Pruitt BA, Jr: The pathophysiology of acute electric injuries. J Trauma 16: 335-340 (May) 1976.

Pruitt BA, Jr, Erickson DR, Morris A: Progressive pulmonary insufficiency and other pulmonary complications of thermal injury. J Trauma 15:369-379.

Brown WL, Bowler EG, Mason AD, Jr. Pruitt BA, Jr: Protein metabolism in burned rats. Am J Physiol 231: 476-482 (Aug) 1976.

Levine NS, Lincberg RB, Mason AD, Jr, Pruitt BA, Jr: The quantitative swab culture and smear: A quick, simple method of determining the number of viable aerobic bacteria on open wounds. J Trauma 16: 89-94 (Feb) 1976.

Levine NS, Atk ns A, McKeel DW, Peck SD, Pruitt BA, Jr: Spinal cord injury following electrical accidents: Case reports, J Trauma 15:459-463 (May) 1975. (received after last reporting).

Wilmore DW, Long JM, Mason AD, Jr, Pruitt BA, Jr: Stress in surgical patients a neurophysiologic reflex response. Surg Gynec Obstet 142: r257-269 (Feb) 1976.

Taylor JW, Plunkett GD, McManus WF, Pruitt BA, Jr: Thermal injury during pregnancy. Obst Sync 47: 434-450 (Apr) 1976.

Andes WA, Baron JP, Bowman RP, Rickles FR: Thermal injury in Von Willebrand disease.Managment of burn wound bleeding. Arch Surg 111: 280-283 (Mar) 1976.

Salisbury RE, Levine NS, McKeel DW, Jr, Wade CR: Tendon sheath reconstruction with artificial implants: A study of ultrastructure. In Am Acad of Orthopedic Surgeons Symp on Tendon Surgery in the Hand (Ed. JM Hunter, LH Schneider) St Louis CV Mosby Co, 1975, pp 59-65.

Lindberg RB, Pruitt BA, Jr, Mason AD, Jr: Topical chemotherapy and prophylaxis in thermal injury. In Chemotherapy, William JD & Geddes AM (eds). Plenum Pub. Corp., NY, Vol. 3:351-359, 1976.

Agee RN, Long JM, III, Hunt JL, Petroff PA, Lull RJ, Mason AD, Jr, Pruitt BA, Jr: Use of ¹³³Xenon in early diagnosis of inhalation injury. J Trauma 16:218-224 (Mar) 1976.

Petroff PA, Hander EW, Mason AD, Jr: Ventilatory patterns following burn injury and effect of Sulfamylon. J Trauma 15: 650-655 (Aug) 1975.

# DISTRIBUTION LIST

5 copies to: US Army Medical Research and **Development Command** Washington, DC 20314 12 opies to: **Defense Documentation Center** ATTN: DDC-TCA Alexandria, Virginia 22314 1 copy\(o; * Superintendent Addemy of Health Sciences, US Army ATT: AHS-COM For Sam Houston, Texas 78234 1 copy to: Dir of aiol & Med Sciences Div Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 1 copy to: CO, Nava Medical R&D Command National Neval Medical Center Bethesda, Haryland 20014 1 copy to: Dir of Prof Sycs Office of the Surgeon General Department of the Air Force Washington, DC 20314 1 copy to:

Director of Defense Research and Engineering ATTN: Assistant Director (Environmental & Life Sciences) Washington, DC 20301

TUS GOVERNMENT PRINTING OFFICE 1977-774-44 1513