

8

3



ECTOPARASITES OF AFRICAN MAMMALS

FINAL SCIENTIFIC REPORT

C. Selby Herrin

and

Vernon J. Tipton

November 1976

Supported by

U. S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND Washington, D. C. 20314

Contract No. DADA17-73-C-3042

Center for Health and Environmental Studies Brigham Young University Provo, Utah 84602

> Approved for public release; distribution unlimited

FILE COPY

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.



BO November 2076	TA. TOTAL NO. OF PAGES	Th. NO. OF REFS				
DADA17-73-C-3042 299	CORIGINATOR'S REPORT NUMBERISI					
	SA. OTHER REPORT HOUS (An	y other numbers that may be coolgoed				
Approved for public release; distr	ribution unlimited	Heesetignatid Mices Horocco				
11. SUPPLEMENTARY NOTES	John F. Reinert, Chief, Entomology U.S. Army Medical Wachington, D.C.	y Research Branch 1 Research & Development				
The study of ectoparasites of At on "potential vectors and reservoirs consisted of ectoparasites from appro- probably more than 500 species of whit taken because (1) epidemiological stu- taxonomic base., (2) a study of ectop actions among animal reservoirs of di- meters for ectoparasites and their ho- patterns. Of the four major groups of mesostigmatid mites and the fleas the fleas were mounted and many identified in the Journal of the Egyptian Public from northwest Africa were mounted and half of those collected in southern is mite genus <u>Laelaps</u> in Africa has been paper is currently being prepared for p <u>Haemolaelaps</u> in northwest Africa south mites have been mounted on slides, but is been done with the ticks. The tick colle who will assume responsibility for the ick terminated with considerable work yet to	of diseases in overseas are oximately 100,000 African sm ich many are yet undescribed udies of arthropod-borne dis parasites may provide inform isease, and (3) an understand dealt with, considerably mor an with the ticks and the tr ed. TA paper on the fleas of c Health Association. Almost nd identified at least to get Africa were mounted. A major n published in the Great Bass publication by a graduate stude of the Sahara. Approximately almost none have been identifi- action is being prepared for sh dentification work of this group	eas." The collection mall mammals, and represent d. The study was under- seases require a sound mation concerning inter- nding of ecological para- ing of epidemiological re work was done with the rombiculid mites. All the f Morocco was published st all mesostigmatid mite enus, however, only about or paper dealing with the sin Naturalist. Another ent dealing with the genus half of the trombiculid ded. Very little work has hipment to another scientist p. Thus, this project was				

Security Classification	LINK				LINKC	
AREY WORDS AND AND A REY WORDS	ROLE	-	OLE	WT	ROLE	WT
African	(sineance	viva:	9125 1	NE BAR	y for	(e)(b)
Arthropod Parasites		As it	eneve N	NU 203	OY MAY	目的
Arthropod-borne Diseases	ana ang ang ang ang ang ang ang ang ang	an en ander		the locate	Rate.	
Chiggers	6 779 1 179 1 10 7					
	- C.Inchi		(nage)	10 633	1 CAMAA	
Ecological Distribution	Canage of several	al and an	gen ve v	of 5 1.270	and the second	1.511.017
Ectoparasites .	la suame.		· ward		alan -	
Epidemiological Studies	alt. t. the	TYNY A	OF STREET	arianati arianati	00 291	1553
Ethiopian Region	to farming	ern rorniga	*******	er the same	and the second second	a second and
Fleas	and the second second	· · · · ·	i		mas a	
Geographical Distribution				N B A	अर्थित्वर विश्व हो जनसङ्ख्या	6 QQ
Haemolaelaps	17	. The	1		-ES-SA	DADA
Host-parasite Relationships	1 0	1 mile		anders en te and a	Con 7 D	には強め
Ixodoidea	1	1				
Laelaps		are i				
Mammals		-				
Mesostigmatid Mites	And the line of a	sara fina	7.6	22/37 673	14 (3 (1 (1 (1 (1 (1 (1 (1 (1 (1 (1 (1 (1 (1	A 1 21 G
Morocco	a) para p	1000				r g q ð
Numerical Taxonomy		gennesenmente este			and the second	
Reservoirs						
Selgem Database Management System					(A	
DOSCON And A STREAM			e engine in		TEL	
Ticks	in at Arms	12518	9940	y/ d.f.	11.72 13	1
Trombiculid Mites	rkgrade eo	15 7 1 1 1 14 7 2 1	625Y	00109	10 09	्र १२ र देव
Vectors a shipper casesalb shipd-bogendine to as	so ida ho a	etjere	500	nadit	er co v	dedo
se or arthrough a diverse a seasch a divers a concerning initer-		series og	an rei		80603 80 No	1 DER
mase, and (3) an understanding of seclogical para	rs of dise	10.0412	en li	mrns i	noma	tions
ACC 35: 17 inc		S and	est ce	16002	e tot	iene.
nor a section L	groups del Res chen	107,10 1 901	our bns	i ang Iatin (tel	t d'acto	1200
	dent fines	it was	bris.	betou	KE BYLS	258
identified at least to delug house only about	i tridutin bried inte	inter an a	B and	17 (A. 17) 1910 - 17	stration of	983
tica were mounted. I regor pager deera a with th	ucharn Ar	02 0	0.000	1105	szönd "	10.11
DISTRIBUTION/AVAILABILITY CODES	a deved add	65447	$A^{*}_{i}(t)$	20312	1 200	42 GR
D: J/CT SPLCIAL	Clean NOT DO	rederad Market	gain	a yura	1100 2	1946
nost some bave been to stiffed. Voru an unrik has	a sourcer es, but sin	bile on	eewst tool n	nom ni	2Gelas Ad avr	4 1.5
on is being prepared he ship on to the selentist	inspillos stal	t aci Th	. 8,45	o di n	Flw sra	No mai
	the the lifer	b they i	12mgs	ion in	0838	in th
a dane with this enormous collection of ectoportables	dict they sha	chief of the	No or B	the corta	The Dest	miner

I. Introduction and Background

The ectoparasites collected from the approximately 100,000 African mammals constitute the largest single collection of ectoparasites made in any geographical or political subdivision of the world. There are probably more than 500 species of ectoparasites in this collection, many of which are yet undescribed. The African continent is noted for the number and variety of endemic arthropod-borne diseases. There were several practical reasons for studying the ectoparasites of African mammals. Epidemiological studies of arthropod-borne diseases require a sound taxonomic base, and a study of ectoparasites and their hosts may enhance understanding of epidemiological patterns.

The major objectives of this project were to (1) determine parameters related to the geographical and ecological distribution of ectoparasites of Africa, (2) provide a reference collection of identified ectoparasites with well documented associated data available to workers with interest in epidemiology and the ectoparasite fauna of Africa, (3) enter the data into a computer data base system (SELGEM system for information management) so that it would be readily available to those working with a specific disease biocenose involving African ectoparasites, and (4) provide data, illustrations and keys through publication of scientific papers which may be used by epidemiologists and entomologists to identify ectoparasites found on mammals of Africa.

The project on ectoparasites of African mammals was restricted to four major groups of arthropod parasites: (1) mesostigmatid mites, (2) trombiculid mites, (3) ticks and (4) fleas. All other arthropod parasites (lice and other miscellaneous insects) in the collections were returned to the Smithsonian Institution for distribution to other specialists for study. The largest group of specimens was the mesostigmatid mites, thus a greater percentage of effort was devoted to this group. Each of these four major groups are treated separately in the report.

Very few of the ectoparasite collections were prepared for museum storage. Some slides of fleas and mesostigmatid mites were labeled with identification labels, but none were labeled with collections data labels. Thus preparation of specimens for museum storage could not be completed with host identifications because the necessary collection data which was only partially available. Since the beginning of the project the Smithsonian Institution Mammal Division Personnel have been entering the collection data into the SELGEM data base management system. We were periodically provided with computer listings of collection records for our own use as they became available. In this way the extoparasites were associated with the hosts from which they were collected. During the nearly six years duration of the project, the major emphasis was in the preparation of the ectoparasites for study; that is, sorting them into major taxonomic groups, shipment of Anoplura, Mallophaga, Streblidae and Nycteribiidae back to the Smithsonian Institution for distribution to appropriate entomologists, mounting of specimens on micriscope slides, labeling and general identification. A letter has been sent to the U.S. Army Research and Development Command requesting a modest financial support to assist in further sorting and shipment of the ectoparasite specimens.

II. Accomplishments

A. Mesostigmatid Mites

The objective of this phase of the study was to investigate the systematics of mesostigmatid mites parasitic on small mammals of the Ethiopian region, including taxonomic revisions of certain genera, and host-parasite relationships of all groups. The specific aims in accomplishing this objective were to (1) make a systematic revision of the genus <u>Laelaps</u> of the Ethiopian region, based on numerical taxonomic analysis and host parasite associations; (2) make a numerical taxonomic analysis of the genus <u>Haemolaelaps</u> of the Ethiopian region; (3) prepare identification keys to all mesostigmatid mite groups occurring on African small mammals; (4) prepare complete descriptions of all new species of African mesostigmatid mites for publication; and (5) investigate the host-parasite and ecological relationships of all parasitic mesostigmatid mites of Africa.

Almost all mesostigmatid mites from northwest Africa have been mounted and identified at least to genus, however, only about half of those collected in southern Africa have even been mounted. A major paper dealing with the mite genus <u>Laelaps</u> in Africa has been published in the Great Basin Naturalist (Herrin and Tipton, 1976, ¹). Another paper is currently being prepared for publication by a graduate student dealing with the genus Haemolaelaps in northwest Africa south of the Sahara. This paper should be submitted for publication within the next few months. Within these two papers 4 new species of <u>Laelaps</u> and 10 new species of Haemolaelaps are described.

Preliminary identification keys to the families and genera of mesostigmatid mites of African small mammals were prepared and are available but are not ready for publication. These keys to families and genera will have to be further refined, and keys to species of all genera prepared before being submitted for publication. Also it is inappropriate to publish these keys outside the context of a systematic review of a particular taxon. Descriptions of new species of African Laelaps and Haemolaelaps mites have been made, but no other descriptions have been prepared for publication. Approximately 12 new species in addition to the Laelaps and Haemolaelaps noted above have been found among the African ectoparasite material. There are no plans at present for the publication of descriptions for these new species.

Herrin, C. Selby and Vernon J. Tipton. 1976. A systematic Revision of the Genus Laelaps s. str. (Acari: Mesostigmata) of the Ethiopian Region. Great Basin Naturalist 36(2):113-205, June 1976.

Studies dealing with host-parasite and ecological relationships of all parasitic mesostigmatid mites of Africa could not be undertaken until after the identifications had been completed and the collection data (host, locality, ecological, etc.) assembled.

All mesostigmatid mite specimens, described as well as undescribed material, will be temporarily deposited in the Brigham Young University Life Science Museum until competent specialists and adequate funding can be obtained to complete the unfinished work.

B. Trombiculid Mites

The objectives were to conduct a systematic study of Ethiopian trombiculid mites, construct keys and make illustrations to facilitate their identification, and provide data relative to host-parasite relationships and other ecological parameters. The specific aims for accomplishing this objective were to (1) complete identification of the African chiggers; (2) construct keys and illustrate characters of taxonomic importance for major taxa; (3) describe new taxa and redescribe other taxa requiring such treatment; and, (4) compile and evaluate host-parasite data in terms of epidemiological significance.

Approximately 5,000 chiggers were mounted, however, all but about 300 specimens were from Morocco. Generic and specific determinations were made for about 1,300 of those mounted. Specific identifications were limited because of the lack of a reference collection of type material. Type specimens for about 80 species were located, but only 20 specimens were received. The material which was identified represents 13 genera or subgenera and some 22 species, including several undescribed species and at least three new genera or subgenera. Thus, evidence indicates considerable undescribed material in the Moroccan collection alone, and the same would be expected in other areas of Africa. There are still approximately 3,250 vials of chiggers yet to be mounted and most of the entire collection is left unidentified.

Progress in the preparation of identification keys, descriptions and illustration of characters was limited until more work could be done in assimilating a reference collection and in the identification of additional material. Thus, very little was accomplished in the description of new species or the preparation of keys and illustrations. Compilation and evaluation of host-parasite data could not be undertaken because of the lack of collection data and complete identifications of the material.

C. Ticks

The objectives of this phase of the project were to identify ticks and make data on host-parasite relationships available to other workers engaged in studies of African Ixodoidea. The specific aims for accomplishing these objectives were to (1) identify all ticks in the Smithsonian-African collection, (2) attempt to make correlations between immature forms and adults, and (3) investigate the host-parasite and ecological relationships of African ticks.

Some progress was made in the preliminary identification of the ticks to genus. The tick collections, sorted by country, collector and collectors number, have been shipped to Dr. Carlton M. Clifford, National Institute of Allergy and Infectious Diseases, Rocky Mountain Laboratory, Hamilton, Montana 59840, who will assume responsibility for the identification of this collection, as well as any other studies which may be undertaken.

D. Fleas

The objectives of this part of the project were to conduct a systematic study of African fleas, compare the flea fauna of each political subdivision with the flea fauna of other political subdivisions and attempt to explain the differences on the basis of physical and biological parameters, and provide publications containing data on host-parasite relationships, distribution, and taxonomic keys and illustrations. The specific aims for accomplishing these objectives were to: (1) complete identification of the approximately 50,000 fleas in the Smithsonian-African collection; (2) publication of the paper on the fleas of Morocco; and (3) prepare publications containing descriptions of new taxa, taxonomic keys, illustrations, host-parasite and ecological relationships, and geographical distribution of all other political subdivisions of Africa.

All fleas have been processed and prepared for identification. Considerable work on the identification of this material was done in collaboration with Lt. Michael Hastriter and Dr. Robert Traub. A paper dealing with the fleas of Morocco was published in the Journal of the Egyptian Public Health Association (Hastriter and Tipton, 1975,²). Reprints of this publication are ready for distribution.

²Hastriter, Michael W. and Vernon J. Tipton. 1975. Fleas (Siphonaptera) Associated With Small Mammals of Morocco. J. Egypt. Publ. Hith Assoc. 50 (2): 79-169, Since a portion of the flea collection was sent to Dr. Robert Traub early in the history of the project, it is essential that this phase of the project be a cooperative venture between Dr. Traub and Lt. Michael Hastriter, or other workers, to avoid duplication of effort. Plans for joint authorship of papers on flea taxa to be described were formulated and Lt. Hastriter had identified about 30% of the fleas of Africa. The flea collection will be temporarily placed in the Brigham Young University Life Science Museum until definite arrangements are made for the completion of the work by Dr. Robert Traub and Lt. Michael Hastriter or other competent specialists.

1

BIBLIOGRAPHY

Hastriter, Michael W. and Vernon J. Tipton. 1975. Fleas (Siphonaptera) Associated with Small Mammals of Morocco. J. Egypt. Public Hith. Assoc. 50(2):79-169.

Herrin, C. Selby and Vernon J. Tipton. 1976. A Systematic Revision of the Genus Laelaps s. str. (Acrai: Mesostigmata) of the Ethiopian Region. Great Basin Nat. 36(2):113-205.

Sakamoto, Calvin J. and C. Selby Herrin. The Genus <u>Haemolaelaps</u> (Acarina: Laelapidae) in the Northwest Ethiopian Biogeographical Region. (manuscript in preparation).

DISTRIBUTION

Contract No. DADA17-73-C-3042

Name of Contractor:

Principal Investigator:

November 1976

C. Selby Herrin

Date of Report:

Final Scientific Report

4 copies

12 copies

1 copy

1 copy

HQDA (SCRDRP) Washington, D. C. 20314

Brigham Young University

Defense Documentation Center Attn: DDC-TCA Cameron Station Alexandria, Virginia 22314

Superintendent Academy of Health Sciences, U. S. Army Attn: AHS-COM Fort Sam Houston, Texas 78234

Center for Health and Environmental Studies

Dean School of Medicine Uniformed Services Univ. of the Health Sciences Office of the Secretary of Defense 6917 Arlington Road Bethesda, MD 20014

Reprints of Publications

10 copies of each

HQDA (SGRDRD) Washington, D. C. 20314