

AD-A046 103

NAVAL INTELLIGENCE SUPPORT CENTER WASHINGTON D C TRA--ETC F/6 8/3
OSCILLATIONS OF THE ICE COVER EXCITED BY INTERNAL WAVES OF THE --ETC(U)
SEP 77 V N SMIRNOV

UNCLASSIFIED

NISC TRANS-3964

NL

1 OF 1
AD
A046103



END
DATE
FILMED

12-77

DDC



DEPARTMENT OF THE NAVY
NAVAL INTELLIGENCE SUPPORT CENTER
TRANSLATION DIVISION
4301 SUITLAND ROAD
WASHINGTON, D.C. 20390

3

AD A 046103

CLASSIFICATION: UNCLASSIFIED

APPROVED FOR PUBLIC RELEASE, DISTRIBUTION UNLIMITED

TITLE:

6 Oscillations of the Ice Cover Excited by Internal Waves of the Arctic Ocean

(Kolebaniya ledyanogo pokrova, obuslovlennyye vnutrennimi volnami ledovitogo Okeana),

10 AUTHOR(S): V. N. / Smirnov, V. N.


PAGES:

5

21 Trans. of SOURCE:

AN SSSR, Doklady Vol. 206, No. 5, 1972

(USSR) v206 n5 p1106-1108
1972.



ORIGINAL LANGUAGE: Russian

TRANSLATOR: C

14 NISC TRANS - 3964

12 5P.
APPROVED P.T.K.

DATE 6 Sep 1977

11

DDC FILE COPY:
NO. 407 682

OSCILLATIONS OF THE ICE COVER EXCITED BY INTERNAL WAVES OF THE ARCTIC OCEAN

[Smirnov, V.N., Kolebaniya ledyanogo pokrova, obuslovlennyye vnutrennimi volnami Ledovitogo Okeana, AN SSSR, Doklady, Vol. 206, No. 5, 1972, pp. 1106-1108, Russian]

→ In 1970 tiltmeters installed on Drift Station North Pole 20 recorded oscillations of the ice cover with periods of up to 25 minutes. Preliminary results of the investigation indicate that these oscillations are excited by internal waves generated in the highly stratified waters of the Arctic ocean basin. ↑

[1106

The pack ice was approximately 3.0 by 5.0 km across and was about 5 m thick. Vertical oscillations of the ice pack were recorded by tiltmeters, horizontal pendulums recording tilting of the ice surface. The pendulum displacement was converted into electric current and recorded on a moving strip of photographic paper. Four identical tiltmeters were installed on the ice cover. Two of the instruments were installed at the same point, at right angles to each other. The remaining tiltmeters were located along a straight line at 25 m intervals. Such arrangement of the instruments made it possible to visually correlate the oscillation phases and to determine the phase velocity and the direction of propagation of the wave. Oscillations from all four sensors were recorded simultaneously by the N-700 light-beam oscillograph. The frequency response of the recording channel is flat in the range of periods observed and the sensitivity to tilts is 0.030 s arc/mm. The double amplitude $2A$ of vertical displacements of the ice cover was determined from the formula $2A = \lambda \phi / \pi$, where λ is the wave length and ϕ is the maximum tilt of the ice floe during propagation of the wave. Fig. 1 shows typical recordings of oscillations of the ice cover. Relatively short-period wave ($\tau = 20$ to 25 s), such as are caused by swell, are superposed on long-period oscillations. Fig. 2 shows vector diagrams of polarization of long-period waves in the plane of the ice floe. A histogram of the periods (Fig. 3) was constructed for a half-year cycle of continuous observations. Of the $n = 1120$ periods determined, approximately 70 percent fall in the range 6 to 13 minutes.

[1107

Table 1 gives the values of wave parameters during specified periods of observation: period T , phase velocity c , wave length λ , double amplitude $2A$, and propagation direction α . All calculations were made on the assumption that the drift velocity is zero, i.e. the Doppler effect was not taken into account.

Numbers in right margin indicate pagination in original text.

DISTRIBUTION FOR	White Section	☒	☐	DISTRIBUTION/AVAILABILITY CODES	BY	D. 1000	SP. CIAL
	Buff Section	☐	☐				
UNCLASSIFIED				A			




Fig. 1 Typical record of oscillation of the ice cover obtained with the aid of special azimuthal placement of tiltmeters




Fig. 2 Wave polarization in the horizontal plane for the following observation periods:
1 - July 11; 2 - August 27;
3 - August 28




Fig. 3. Histogram of oscillation periods of the ice cover

These are the first recordings of surface gravity waves in ice-covered sea, having phase velocities between 0.5 and 2.0 m/s. The results of theoretical investigations and processing of observational data on flexural gravity waves in the period range between 0.1 and 1000 s are published in references [1 to 4]. It was established that natural oscillations of the ice cover are caused primarily by the breakup of ice, effects of wind, and transmission of gravity waves incident on the ice from the free water side.

Date	T, s	c m/s	λ , m	α , °	2A, mm
11 VII	840	1.6	1340	57	4.00
27 VIII	660	1.1	730	26	4.36
28 VIII	880	1.1	980	41	4.90
28 VIII	800	1.2	960	36	4.33
28 VIII	900	1.0	900	180	3.12
28 VIII	1080	0.7	760	180	2.66
14 IX	700	2.0	1400	50	4.20
14 IX	670	0.5	330	303	0.84
14 IX	670	0.6	400	300	1.00

The results of our observations appear to indicate that the source responsible for generation of such slow waves is found in the ocean depths. This assumption appears to be substantiated by the fact that the propagation parameters of such types of waves along the ocean surface (Table 1) are in good agreement with the existing concepts of short-period internal waves. It can be assumed that internal waves, which are proper oscillations of the layers of stably-stratified sea, reach the surface causing oscillations of the ice cover with the same period, phase velocity and direction of propagation. The polarization of internal waves (Fig. 2) indicates a sufficiently stable progressive wave with a specific direction of propagation. [1108] This important factor may serve as a definite criterion making it possible to establish the characteristics of wave motion. Thus, oscillations of the ice cover with very low propagation velocities may provide data on the dynamic state of deep water masses.

In conclusion the authors would like to express their gratitude to V.T. Spitsyn for the assistance provided on the drifting station and also to V.P. Gavrilov, V.G. Savchenko and B.Ye. Kheysin for a useful discussion.

Arctic and Antarctic

Received on July 2, 1975

Scientific Research Institute,
Leningrad

REFERENCES

1. Hunkins, K. Journal of Geophysical Research "Waves in the Arctic Ocean", Vol. 67, No. 6, 1962, pp. 2477-2489.
2. LeSchack, L.A. and R.A. Haubrich, Journal of Geophysical Research, "Observations of Waves on an Ice-Covered Ocean," Vol. 69, No. 18, 1964, p. 3815-3821.
3. Smirnov, V.N. Problems of the Arctic and Antarctic (Problemy Arktiki i Antarktiki), Vol. 29, 1968, p. 70.
4. Smirnov, V.N., Ye.M. Lin'kov, Fizika Zemli, No. 8, 1967, p. 88.