

AD-A046 379

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER F/G 12/1  
BOUNDS FOR THE SET OF SOLUTIONS OF FUNCTIONAL-DIFFERENTIAL EQUA--ETC(U)  
AUG 77 K L NICKEL DAAG29-75-C-0024  
MRC-TSR-1782 NL

UNCLASSIFIED

| OF |  
ADA  
046 379



END  
DATE  
FILMED  
12-77  
DDC

AD A 0 46379

⑨ MRC Technical Summary Report, #1782

⑥ BOUNDS FOR THE SET OF SOLUTIONS OF FUNCTIONAL-DIFFERENTIAL EQUATIONS.

⑩ Karl L. Nickel

⑪ B.S.

Mathematics Research Center  
 University of Wisconsin-Madison  
 610 Walnut Street  
 Madison, Wisconsin 53706

DDC  
 RECEIVED  
 NOV 15 1977  
 F

⑪ August 1977

⑫ 25p.

(Received July 27, 1977)

⑭ MRC-TSR-1782

⑮ DAAG29-75-C-φφ24

DDC FILE COPY

Approved for public release  
 Distribution unlimited

Sponsored by  
 U. S. Army Research Office  
 P. O. Box 12211  
 Research Triangle Park  
 North Carolina 27709

1473  
 221 200

1B

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

BOUNDS FOR THE SET OF SOLUTIONS OF FUNCTIONAL-DIFFERENTIAL EQUATIONS

Karl L. Nickel \*

Technical Summary Report #1782

August 1977

ABSTRACT

Sets of systems of ordinary functional-differential equations with Volterra type functionals under sets of initial values are considered. Upper and lower bounds are constructed for the sets of all solutions. Classes of such problems are given where these bounds are optimal. The main tool is a Lemma of Max Müller on inequalities. Also ideas from interval mathematics are used.

AMS (MOS) Subject Classification: 34J05

Key Words: Functional-differential equations and inequalities, Volterra type, Solution set, Optimal bounds, Maximal interval solution

Work Unit Number 1 (Applied Analysis)

SIGNIFICANCE AND EXPLANATION

If differential equations

$$u'(t) = f(t, u(t)), \quad u(0) = \alpha$$

appear in Applied Mathematics there is normally not just one right hand side  $f(t, u)$ . Instead of this a whole set  $\{f\}$  of right hand sides must be considered. This is due to many facts such as: data errors, data intervals obtained from measurements, approximation of  $f$  by a more suitable function, poor knowledge of the laws involved etc. The same is true for the initial

---

\* Visiting from the University of Freiburg/Germany.

"value"  $\alpha$  which is usually a set  $\{\alpha\}$ . Hence the above initial value problem has to be replaced by the inclusion problem

$$u'(t) \in \{f(t, u(t))\}, \quad u(0) \in \{\alpha\} .$$

It is normally completely impossible to solve all the real problems which are combined in this set of problems. The goal of the following paper is therefore to find at least lower and upper bounds to the set of all such solutions. This can always be done. Since these bounds are sometimes very pessimistic, classes of such problems are given where the bounds obtained are optimal.

The main ideas of this paper are also valid in the more general case where  $f$  does depend as a functional upon the unknown solution  $u$ . This is written in the form  $f(t, u(t), u(\cdot))$ . Therefore the theory of this paper also includes integro-differential equations and difference-differential equations. Sets of such problems do occur for example in Economics and in Biology.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	APR 80/01 SPECIAL
A	

BOUNDS FOR THE SET OF SOLUTIONS OF FUNCTIONAL-DIFFERENTIAL EQUATIONS

Karl L. Nickel \*

1. Introduction

In the following paper systems of functional-differential equations

$$(1) \quad u'(t) = f(t, u(t), u(\cdot)) \quad \text{for } 0 < t \leq T$$

are considered under the initial conditions

$$(2) \quad u(0) = \alpha .$$

Herein  $u = (u_1, u_2, \dots, u_n)$ ,  $f = (f_1, f_2, \dots, f_n)$  and  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$  are  $n$ -vectors.

As usual  $u$  or  $u(t)$  means the value of the function  $u$  at the point  $t$ ; moreover  $u'(t) = du/dt$ . Opposite to this the notation  $u(\cdot)$  means that  $u$  is regarded as an element of the class of admissible functions. Hence  $f(\cdot, \cdot, u(\cdot))$  is a functional on  $u$ ; in what follows only special "Volterra" functionals will be regarded.

If  $f$  is continuous then the system (1), (2) is equivalent to the system of functional-integral equations

$$(3) \quad u(t) = \alpha + \int_0^t f(s, u(s), u(\cdot)) ds \quad \text{for } 0 \leq t \leq T .$$

It is the subject of the following paper to find bounding functions  $v(t)$ ,  $w(t)$  such that for every solution  $\hat{u}(t)$  of (1), (2) or (3)

$$(4) \quad v(t) \leq \hat{u}(t) \leq w(t) \quad \text{for } 0 \leq t \leq T .$$

Hence the classical theory of maximal and minimal solutions for differential equations appears as a special case of these results.

If a solution  $\hat{u}$  of (1), (2) is uniquely determined then it is trivial that (4) is satisfied for  $w := v := \hat{u}$ . It is therefore interesting to switch to a more general problem: Let  $\{\alpha\}$  be a set of initial values and let  $\{f\}$  be a set of right hand sides to (1). Then the more general initial value problem

---

\* Visiting from the University of Freiburg/Germany.

$$(5) \quad u'(t) \in \{f(t, u(t), u(\cdot))\} \text{ for } 0 < t \leq T,$$

$$(6) \quad u(0) \in \{a\}$$

is considered. Let  $\{\hat{u}\} = \{\hat{u}(t)\}$  be the set of all solutions of (5), (6). Again two functions  $v(t)$ ,  $w(t)$  are looked for such that (4) is true for any solution  $\hat{u} \in \{\hat{u}\}$ . If one writes  $[v, w]$  for the function interval from the two bound functions  $v$  and  $w$  then this can be written as

$$(7) \quad \{\hat{u}\} \in [v, w].$$

It is in general quite simple to find rough bounds  $v, w$ . In what follows special emphasis is therefore given to the look for "optimal" bounds. Here "optimality" means the following: let there exist the infimum and the supremum of the set  $\{\hat{u}\}$  such that

$$(8) \quad v = \inf\{\hat{u}\}, \quad w = \sup\{\hat{u}\}.$$

In that case one can call  $[v, w]$  the "interval hull" of  $\{\hat{u}\}$ . It is the goal of this paper to find classes of sets  $\{f\}$  and  $\{a\}$  such that (7) and (8) are true.

In order to get such results a lemma of Max Müller (1927) on differential inequalities is essential. This lemma has been published exactly 50 years ago. For decades however, it remained widely unnoticed. In what follows this lemma will be extended to the case of functional-differential inequalities. This will be done by extending an old paper of the author (Nickel (1961)).

It should finally be remarked that the problem of this paper and some of the formulations have been strongly influenced by the ideas of interval mathematics.

## 2. Notations and assumptions

Let  $n \in \mathbb{N}$ ,  $0 < T \in \mathbb{R}$ ,  $I := [0, T]$ ,  $I_0 := (0, T)$ ,  $\alpha \in \mathbb{R}^n$ . The  $n$ -vectors  $\alpha$ ,  $u(t)$ ,  $f(t, u, u(\cdot))$  are written as  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ ,  $u = (u_1, u_2, \dots, u_n)$ ,  $f = (f_1, f_2, \dots, f_n)$ . Together with the  $k$ -th component  $u_k$  of the vector  $u$  also the  $n-1$ -vector  $u^k = (u_1, u_2, \dots, u_{k-1}, u_{k+1}, \dots, u_n)$  is used.

Let the class  $Z$  of the admissible functions be defined as the set of all function vectors  $u : I \rightarrow \mathbb{R}^n$ , continuous on  $I$  such that the derivative  $u' = du/dt$  exists in  $I_0$ .

The notation  $u(\cdot)$  means that  $u$  is an element of the class  $Z$ . Opposite to this  $u$ ,  $u'$  or  $u(t)$ ,  $u'(t)$  mean the values of these functions at the point  $t$ .

Inequalities  $v(t) \leq w(t)$  or  $v \leq w$  are always meant componentwise as  $v_k(t) \leq w_k(t)$  for  $k = 1(1)n$ . Inequalities of the kind  $v(\cdot) \leq w(\cdot)$  are meant both componentwise and pointwise for all points in the definition set.

For  $v \leq w$  the interval  $[v, w]$  is defined as the set  $[v, w] = [v(t), w(t)] := \{z \in \mathbb{R}^n \mid v(t) \leq z \leq w(t)\}$ . Similarly  $[v(\cdot), w(\cdot)] := \{z \in Z \mid v(\cdot) \leq z(\cdot) \leq w(\cdot)\}$ .

Let the dependence of  $f_i(t, u, u(\cdot))$  of any component  $u_k(\cdot)$  for  $k = 1(1)n$  be that of a "Volterra" functional. Here a functional  $g(t, z(\cdot))$  is called "Volterra" if it represents a mapping in  $\mathbb{R}$  such that the value of  $g$  at the point  $t$  depends only on the value  $z(s)$  for  $0 \leq s \leq t$  ( $g$  depends only upon "the past" of the function  $z(\cdot)$ ).

Examples of Volterra functionals are

$$g(t, z(\cdot)) := \int_0^t K(t, s, z(s)) ds,$$

$$(9) \quad g(t, z(\cdot)) := z(\tau \cdot t) \quad \text{with} \quad 0 \leq \tau \leq 1,$$

$$(10) \quad g(t, z(\cdot)) := \begin{cases} z(t - s) & \text{for } 0 \leq s \leq t, \\ \alpha(t - s) & \text{for } 0 \leq t < s \end{cases}$$

with some given function  $\alpha(t)$  for  $-s \leq t < 0$ .

These examples show that the theory given in this paper can be applied to: differential equations, (Volterra) integro-differential equations, difference-differential equations

with retarded argument and naturally also to combinations of these equations.

In order to simplify the results the following notation will be used for the components  $f_k$  of  $f(t, u, u(\cdot))$ : the second argument  $u \in \mathbb{R}^n$  of  $f$  is broken up in the component  $u_k$  with the same index  $k$  as  $f_k$  and in the rest vector  ${}_k u$ :

$$f_k = f_k(t, u_k, {}_k u, u(\cdot)) .$$

Furthermore it is suitable to have a special notation for the set of all functions  $f_k$  if the arguments lie in certain intervals. This will be denoted by

$$\{f_k(t, u_k, [{}_k v, {}_k w], [v(\cdot), w(\cdot)])\} := \{f(t, u_k, {}_k u, u(\cdot)) \mid u_k \in [{}_k v, {}_k w], u(\cdot) \in [v(\cdot), w(\cdot)]\} .$$

From section 7 to the rest of this paper only functions  $f$  will be regarded which are partially monotone. The corresponding definitions will be given in section 7.



### 3. Existence

The following theorem is the extension of the well known Peano existence theorem for systems of differential equations:

Theorem:

Let  $f$  be defined and continuous on  $I \times \mathbb{R}^n \times Z$ . Then (1), (2) and (3) are equivalent. If  $f$  is bounded there exists (at least) one solution  $\hat{u} \in Z$  of (1), (2).

If  $f$  is not bounded then there exists a solution of (1), (2) at least in a largest interval  $0 \leq t < T_1 \leq T$ .

Proof: The equivalence is trivial. For the existence the fixed point theorem of Schauder is applied to equation (3). The main ideas are exactly the same as in the case of differential equations. They are described in the book of Walter (1970), p. 23-25.

#### 4. The Lemma of Max Müller

Lemma:

Let the functions  $v, w \in Z$  with  $v \leq w$  satisfy the following inequalities

$$(11) \quad v(0) < \alpha < w(0) ,$$

$$(12) \quad v'_k < \{f_k(t, v_k, [{}_k v, {}_k w], [v(\cdot), w(\cdot)])\} ,$$

$$(13) \quad w'_k > \{f_k(t, w_k, [{}_k v, {}_k w], [v(\cdot), w(\cdot)])\}$$

for  $t \in I_0$  and  $k = 1(1)n$  .

Then any solution  $\hat{u} \in Z$  of (1), (2) is bounded by

$$(14) \quad v(t) < \hat{u}(t) < w(t) \quad \text{for } t \in I .$$

Corollaries:

1) If  $u \in Z$  is a solution of the inequalities  $u(0) \leq \alpha$ ,  $u' \leq f(t, u, u(\cdot))$  in  
 $I_0$  then

$$u(t) < w(t) \quad \text{for } t \in I .$$

2) Similarly

$$\bar{u}(t) > v(t) \quad \text{for } t \in I$$

for any solution  $\bar{u} \in Z$  of the inequalities  $u(0) \geq \alpha$ ,  $u' \geq f(t, u, u(\cdot))$  in  $I_0$ .

3) If all  $f_k$  are strictly monotone (increasing or decreasing) with respect to  
(at least) one of the components of  $u(\cdot)$ , it then suffices to have the  $\geq$  - and  
 $\leq$ -signs in (12) and (13) instead of the  $>$ - and  $<$ -signs.

Remarks:

1) This Lemma has been formulated and proven by Max Müller (1927) as Theorem 5 on the pages 13 to 15 for the special case where  $f$  does not depend upon  $u(\cdot)$ . See W. Walter (1970), p. 93-94.

2) The original notation of M. Müller was very inconvenient. It has here been replaced by the interval notation.

3) Kindly note that there are no assumptions to be made with respect to the function  $f$ , such as continuity, monotonicity etc.

Proof: The original proof of M. Müller carries right over to this case.

5. Example

In the following example  $n = 2$ . The functional (9) is used for  $\tau = 1/2$  together with the other functional

$$\int_0^t (u_1^2(s) + u_2^2(s)) ds .$$

Let

$$f_1(t, u, u(\cdot)) := -2u_1^2 + u_2/(1 + t^2) + (u_1(t/2) + u_2(t/2))/2 ,$$

$$f_2(t, u, u(\cdot)) := 1 - \sin(\pi u_1/2) + 2u_2(1 - 2u_2) + \frac{1}{2t} \int_0^t (u_1^2(s) + u_2^2(s)) ds ,$$

$$0 < \alpha_1, \alpha_2 < 1 .$$

Define  $v_1 = v_2 := 0, w_1 = w_2 := 1$  for  $t \geq 0$ . Then one verifies easily for  $t > 0$

$$f_1(t, v_1, [v_2, w_2], [v_1(\cdot), w_1(\cdot)], [v_2(\cdot), w_2(\cdot)]) = 0 + [0, 1]/(1 + t^2) + [0, 1] \geq 0 ,$$

$$f_1(t, w_1, [v_2, w_2], [v_1(\cdot), w_1(\cdot)], [v_2(\cdot), w_2(\cdot)]) = -2 + [0, 1]/(1 + t^2) + [0, 1] < 0 ,$$

$$f_2(t, v_2, \dots) = [0, 1] + 0 + [0, 1] \geq 0 ,$$

$$f_2(t, w_2, \dots) = [0, 1] - 2 + [0, 1] \leq 0 .$$

Since  $f_1$  and  $f_2$  are strictly monotone increasing with respect to both components  $u_1(\cdot)$  and  $u_2(\cdot)$  the third corollary to the lemma can be used. This gives the a priori estimate  $0 < \hat{u}_1(t), \hat{u}_2(t) < 1$  for  $t \geq 0$  for any solution  $\hat{u}$ .

## 6. Uniqueness and error bounds

For ordinary differential equations all uniqueness theorems can be derived from the Lemma of M. Müller. In his original paper he did use the lemma exactly for that purpose. Similarly probably all known a posteriori bounds can be proven with its help. Therefore a very large number of such theorems can immediately be proven also for the case of functional-differential equations. It is possible to translate all the results from the book of W. Walter to these extended systems of equations. Some first results have already been given in the old paper by K. Nickel (1961). It is however, not the purpose of this paper to publish such theorems.

7. Monotonicity conditions

Definition (unconditionally partially isotone/antitone/monotone):

Let  $g(x_1, x_2, \dots, x_m)$  be a mapping  $g : D \rightarrow \mathbb{R}$  with  $D \subseteq \mathbb{R}^m$ . Again the notation  $g(x_1, \dots, x_m) = g(x_k, x)$  is used. The function  $g$  is called unconditionally partially isotone or antitone on  $D$  with respect to the variable  $x_k$  if

$$g(y_k, x) \geq g(x_k, x) \text{ for } y_k \geq x_k \text{ or } y_k \leq x_k$$

and for all  $(x_k, x), (y_k, x) \in D$ . A function which is either unconditionally partially isotone or unc. part. antitone is called unconditionally partially monotone.

Kindly note that the function  $g(x_1, x_2) := x_1 \cdot x_2$  is unconditionally partially monotone with respect to  $x_1$  and  $x_2$  on  $D := [0, \infty) \times [0, \infty)$ , but not on  $D := \mathbb{R}^2$ .

Definition (monotonicity class  $M$ ):

Let the class  $M$  consist on all functions  $f(t, u, v(\cdot))$  for which the following is true: Each function  $f_k(t, u_k, u, v(\cdot))$  is unconditionally partially monotone on  $I_0 \times \mathbb{R} \times \mathbb{R}^{n-1} \times Z$  with respect to any component of  $u$  and  $v(\cdot)$ , but not necessarily with respect to  $u_k$ .

If  $f \in M$  then it is convenient to write

$$f_k(t, u_k, u, v(\cdot)) = f_k(t, u_k, u^+, u^+, v(\cdot)^+, v(\cdot)^+)$$

This clearly means that the vectors  $u_k$  and  $u(\cdot)$  are divided in the two sets of components for which  $f_k$  is isotone (+) and antitone (-).

Let  $f \in M$ . Then in the Lemma of M. Müller the (rather inconvenient) inequalities of sets can be replaced by real inequalities (which are much simpler to handle). In this case the Lemma reduces to the

Special case of the Lemma:

Let  $f \in M$ . Assume that (11) is true for functions  $v, w \in Z$  with  $v \leq w$  and that furthermore

$$v_k^+ < f_k(t, v_k, v_k^+, w_k^+, v(\cdot)^+, w(\cdot)^+),$$

$$w_k^+ > f_k(t, w_k, w_k^+, v_k^+, w(\cdot)^+, v(\cdot)^+)$$

for  $t \in I_0$  and  $k = 1(1)n$ .

Then any solution  $\hat{u} \in Z$  of (1), (2) is bounded by (14); furthermore the corollaries 1) to 3) hold.

The proof comes immediately from the Lemma together with the definition of the class  $M$ .

### 8. Construction of bounds

In what follows the special case of the Lemma will be used to construct bound functions  $v$  and  $w$ . The above problem (1), (2), (4) will be replaced by a more general since no additional difficulties are generated by that extension:

The initial data  $\alpha$  in the initial conditions (2) are to be replaced by a set  $\{\alpha\}$ . For simplicity assume  $\{\alpha\} \subseteq [\underline{\alpha}, \bar{\alpha}]$  with two vectors  $\underline{\alpha}, \bar{\alpha} \in \mathbb{R}^n$  and  $\underline{\alpha} \leq \bar{\alpha}$  such that

$$(15) \quad \underline{\alpha}, \bar{\alpha} \in \{\alpha\} \subseteq [\underline{\alpha}, \bar{\alpha}].$$

Hence the initial conditions (1) are to be replaced by (6). Let  $\{\hat{u}\}$  be the set of all solutions of (1) under the set of all initial conditions (6). Wanted are bounds  $v, w$  such that (7) is true.

Assume that  $f$  is continuous and bounded on  $I \times \mathbb{R}^n \times Z$ . The following sequence of problems for  $v \in \mathbb{N}$  is considered:

$$(16) \quad \begin{cases} v_k = f_k(t, v_k, v_k^{v+1}, w_k, v_k^{(\cdot)+}, w_k^{(\cdot)+}) - \frac{1}{v}, \\ w_k = f_k(t, w_k, w_k^{v+1}, v_k, w_k^{(\cdot)+}, v_k^{(\cdot)+}) + \frac{1}{v} \end{cases}$$

for  $t \in I_0$  and  $k = 1(1)n$ ,

$$(17) \quad v(0) = \underline{\alpha} - 1/v, \quad w(0) = \bar{\alpha} + 1/v.$$

By the existence theorem of section 3 there exists for every  $v \in \mathbb{N}$  at least one solution  $v, w \in Z$  of the coupled system (16), (17). An arbitrary solution is picked up and called  $(v^v, w^v)$ . Then

$$v^v < v^{v+1} < \hat{u} < w^{v+1} < w^v$$

for all  $v \in \mathbb{N}$  and for any solution  $\hat{u} \in Z$  of (1), (2). This follows by the special case of the Lemma and by the definition of the right hand sides in (16) and in (17).

Since the sequences  $\{v^v\}$  and  $\{w^v\}$  are monotone and bounded. As sequences of measurable functions they have measurable limit functions  $\underline{v}(t) := \sup\{v^v(t)\}$  and  $\bar{w}(t) := \inf\{w^v(t)\}$ . Furthermore

$$(18) \quad \underline{v} \leq \hat{u} \leq \bar{w}.$$

One can show as usual that these sequences are uniformly convergent on  $I$  and that  $\underline{v}, \bar{w} \in Z$  (see W. Walter (1970), p. 68). Furthermore the pair  $(\underline{v}, \bar{w})$  satisfies the

following functional-differential system in  $I_0$  consisting of  $2n$  equations

$$(19) \quad \begin{cases} \underline{v}_k = f_k(t, \underline{v}_k, \underline{v}_k^+, \bar{w}_k^+, \underline{v}(\cdot)^+, \bar{w}(\cdot)^+) , \\ \bar{w}_k = f_k(t, \bar{w}_k, \bar{w}_k^+, \underline{v}_k^+, \bar{w}(\cdot)^+, \underline{v}(\cdot)^+) \end{cases}$$

under the  $2n$  initial conditions

$$(20) \quad \underline{v}(0) = \underline{\alpha}, \quad \bar{w}(0) = \bar{\alpha} .$$

If  $\hat{u} \in Z$  is a solution of (1), (2), then the pair  $(\hat{u}, \hat{u})$  is also a solution of (19),

(2). In general the reverse is however, not true, i.e. the functions  $\underline{v}, \bar{w}$  are in general no solutions of (1), (6). Hence they are not minimal or maximal solutions to (1), (6) in the usual sense (see however, section 11). In what follows the interval  $[\underline{v}, \bar{w}]$  will be called maximal interval solution of (1), (6).

The reason for this notation comes from the following: Define the interval operator  $F[v, w] = (F_1, F_2, \dots, F_n)$  by its  $k$ -th component as follows:

$$F_k[v, w](t) := [\underline{\alpha}_k + \int_0^t f_k(s, \underline{v}_k, \underline{v}_k^+, \bar{w}_k^+, \underline{v}(\cdot)^+, \bar{w}(\cdot)^+) ds , \\ \bar{\alpha}_k + \int_0^t f_k(s, \bar{w}_k, \bar{w}_k^+, \underline{v}_k^+, \bar{w}(\cdot)^+, \underline{v}(\cdot)^+) ds] .$$

Any solution  $\hat{u}$  of (1), (6) satisfies

$$\hat{u} \in F[\hat{u}, \hat{u}] .$$

Moreover the interval  $[\underline{v}, \bar{w}]$  is a fixed interval of the operator  $F$  by (19) and (20).

By construction  $[\underline{v}, \bar{w}]$  is the smallest fixed interval of  $F$  for which (18) is true.

If  $\underline{\alpha} = \bar{\alpha} = \alpha$  and if maximal and minimal solutions of (1), (2) exist then they are equal to  $\underline{v}$  and  $\bar{w}$ .

This idea consists therefore in replacing the usual ordering relation  $\leq$  (componentwise with respect to  $k$  and pointwise with respect to  $t$ ) by the inclusion  $\subseteq$  as a new ordering relation (also componentwise and pointwise).



9. Bounds for the solutions of sets of functional-differential equations

The equations (1), (2) are now being replaced by inclusions (5), (6).

Theorem:

Let (15) be satisfied by the set of initial values  $\{a\}$ . Assume that there exist two right hand sides  $\underline{f}, \bar{f} \in \{f\}$  with  $\underline{f} \leq \bar{f}$  such that for all solutions  $\hat{u} \in Z$  of (5), (6)

$$\underline{f}(t, u, u(\cdot)), \bar{f}(t, u, u(\cdot)) \in \{f(t, u, u(\cdot))\} \subseteq [\underline{f}(t, u, u(\cdot)), \bar{f}(t, u, u(\cdot))] \text{ in } I_0.$$

Assume furthermore that the two functions  $\underline{f}, \bar{f} \in Z$  and are continuous and bounded on  $I \times \mathbb{R}^n \times Z$ . Construct the maximal interval solutions to  $\underline{a}, \underline{f}$  and  $\bar{a}, \bar{f}$  to the problem (1), (2) and call them  $[\underline{v}, \underline{w}]$  and  $[\bar{v}, \bar{w}]$ . Then  $\underline{v} \leq \bar{w}$  and for the set  $\{\hat{u}\}$  of all solutions  $\hat{u} \in Z$  of (5), (6) the inclusion

$$(21) \quad \{\hat{u}\} \subseteq [\underline{v}, \bar{w}]$$

is true.

Remarks: 1) There is nothing assumed for one of the right hand sides  $f \in \{f\}$  if  $f \neq \underline{f}, \bar{f}$ , only the existence of that function. If  $f$  for example is not continuous, no solution  $\hat{u}$  of (1), (2) may exist. Kindly note that the theorem deals only with existing solutions.

2) In order to find the two bounds  $\underline{v}, \bar{w}$  for all (in general  $\infty$  many) solutions of (5), (6) one has to determine the maximal interval solutions of two coupled systems with  $2n$  equations each. The functions  $\bar{v}$  and  $\underline{w}$  are a "side effect" of this procedure, they are not needed for the inclusion (21). They do have however, a meaning as "inner" bounds to  $\{\hat{u}\}$  in the sense of interval mathematics.

Proof: Let  $\hat{u}$  be a solution of (5), (6). Then  $\hat{u}' = f(t, \hat{u}, \hat{u}(\cdot)) \leq \bar{f}(t, \hat{u}, \hat{u}(\cdot))$  in  $I_0$ . Then  $\hat{u} \leq \bar{w}$  in  $I$  by the corollary 1 of the lemma and by the construction of  $(\bar{v}, \bar{w})$ .

The inequality  $\hat{u} \geq \underline{v}$  is shown similarly which finishes the proof.

10. Two examples

1) Let  $n = 1, \alpha = 0,$

$$(22) \quad f(t, u, u(\cdot)) := 2(\sqrt{|u(t)|} + u(\tau \cdot t))$$

and

$$\{f\} := \{f | 0 \leq \tau \leq 1\} .$$

The functional used is  $u(\tau \cdot t)$  by (9). I do not know if the problem (1), (2) with  $f$  by (22) can explicitly be solved for  $\tau \neq 0, 1.$

Since  $f$  is isotone in  $u(\cdot)$  the two inequalities (12) and (13) of the Lemma are decoupled. By putting  $v(t) := -\epsilon, 0 < \epsilon < 1$  one gets  $v(0) = -\epsilon < 0 = \alpha$  and

$$0 = v'(t) < 2(\sqrt{|v(t)|} + v(\tau \cdot t)) = 2\sqrt{\epsilon} (1 - \sqrt{\epsilon}) .$$

Hence by the Lemma  $\hat{u}(t) > -\epsilon$  for any solution  $\hat{u} \in Z$  of (1), (2). For  $\epsilon \rightarrow 0$  one gets  $\hat{u}(t) \geq 0$  and therefore by (22) there is also  $\hat{u}'(t) \geq 0$  for any solution  $u.$

Hence  $0 \leq \hat{u}(\tau \cdot t) \leq \hat{u}(t)$  for any solution, therefore one can define

$$\underline{f}(t, u, u(\cdot)) := 0, \bar{f}(t, u, u(\cdot)) := 2(\sqrt{|u(t)|} + u(t)) .$$

The maximal interval solution is found easily as  $\underline{v} := 0, \bar{w} := (\exp t - 1)^2$  hence

$$(23) \quad \hat{u}(t) \in [0, (\exp t - 1)^2] \text{ for } t \geq 0 ,$$

for all solutions  $\hat{u}$  of (1), (2) with (22).

If one now changes the functional (9) in (22) to (10) one gets the function

$$(24) \quad f(t, u, u(\cdot)) := \begin{cases} 2(\sqrt{|u(t)|} + u(t-s)) & \text{for } 0 \leq s \leq t , \\ 2\sqrt{|u(t)|} & \text{for } 0 \leq t \leq s , \end{cases}$$

where  $0 \leq s < \infty.$  With this right hand side (24) one gets a whole set of difference-differential equations with retarded argument. The same ideas as above give exactly the same bound functions  $\underline{v}$  and  $\bar{w}.$  Hence also in this case (23) is true.

The same can be said for the third different right hand side

$$(25) \quad f(t, u, u(\cdot)) := 2(\sqrt{|u(t)|} + (\int_0^t |u(s)|^p ds)^{1/p})$$

with  $1 \leq p \leq \infty.$  The functional in this case is the Volterra  $p$ -norm with the sup norm for  $p = \infty.$

Since in all three cases the bounds  $\underline{v}, \bar{w}$  are solutions itself to  $u' = \underline{f}, u' = \bar{f}$  one gets in addition the optimality condition

$$(26) \quad \underline{v}, \bar{w} \in \{\hat{u}\} \subseteq [\underline{v}, \bar{w}] .$$

This result is highly surprising. The three problems with the different right hand sides (23), (24) and (25) most certainly have completely different solutions and therefore also different solution sets. In spite of this fact all three sets have the same bounds and furthermore these bounds are optimal.

This example shows also, that it is very often simpler to look for bounds  $\underline{v}, \bar{w}$  such that (26) is true than to try to solve the equations.

2) Let  $n = 2$ , the given system is

$$\begin{cases} u_1' = u_2, & u_1(0) = \alpha_1, \\ u_2' = -u_1, & u_2(0) = \alpha_2. \end{cases}$$

The uniquely determined solution is

$$\hat{u}_1(t) = \alpha_1 \cos t + \alpha_2 \sin t, \quad \hat{u}_2(t) = -\alpha_1 \sin t + \alpha_2 \cos t .$$

Let  $\alpha_1 \in [0,1], \alpha_2 \in [0,1]$ . Then by the rules of interval arithmetic (see R. E. Moore (1966))

$$\begin{aligned} \hat{u}_1(t) &\in [0,1] \cos t + [0,1] \sin t, \\ u_2(t) &\in [-1,0] \sin t + [0,1] \cos t . \end{aligned}$$

The set of solutions is hatched in Figure 1. In the picture also the "main" solution for  $\alpha_1 = \alpha_2 = 1/2$  is shown. The extended system (19) reads here as

$$\begin{aligned} v_1' &= v_2, & v_1(0) &= 0, \\ v_2' &= -v_1, & v_2(0) &= 0, \\ v_3' &= v_2, & v_3(0) &= 1, \\ v_4' &= -v_1, & v_4(0) &= 1. \end{aligned}$$

The solution to this system is unique, hence the maximal interval solution




FIGURE 1

$$2v_{-1}(t) = -e^t + \sin t + \cos t ,$$

$$2v_{-2}(t) = -e^t + \cos t - \sin t ,$$

$$2\bar{w}_1(t) = e^t + \sin t + \cos t ,$$

$$2\bar{w}_2(t) = e^t + \cos t - \sin t .$$

Hence the functions  $\underline{v}, \bar{w}$  "back away" from the "main" solution (see Figure 1) as fast as  $e^t/2$  to below and to above.

For  $t = 2\pi$  one sees  $\{\hat{u}_1(2\pi)\}, \{\hat{u}_2(2\pi)\} \subseteq [0,1]$ . But  $\bar{w}_k(2\pi) - v_k(2\pi) = e^{2\pi} = 535.4 \dots$  for  $k = 1,2$ . The real set of solutions  $\{\hat{u}\}$  is therefore surpassed at  $t = 2\pi$  by  $[\underline{v}, \bar{w}]$  by a factor of more than 500 and this grows rapidly worse for larger values of  $t!!!$

This example was first discussed by R. E. Moore (1966) by using geometrical reasoning.

11. Optimal bounds for the set of solutions: maximal and minimal solutions

By section 9 one sees that it is enough to restrict the survey to the solution of two real functional differential equations if one wishes to bound the solutions of sets of such equations. Hence the following optimality considerations are given only for systems of the type (1). Since it is no aggravation the initial inclusion (6) has however, been used instead of the initial condition (2). It is always assumed that (15) is true.

As is shown by the second example of section 10 the maximal interval solution  $[\underline{v}, \bar{w}]$  gives in general not optimal bounds to the set of solutions  $\{\hat{u}\}$ . In this and in the next section classes of problems will be given such that there is optimality either in the sense of (26) or at least of (8).

Let  $f \in M$ . Assume furthermore that all functions  $f_k(t, u_k, u, v(\cdot))$  are unconditionally partially isotone with respect to any of the components of  $u$  and  $v(\cdot)$ . This is called "quasimonotone increasing" by W. Walter (1970) in the case of differential equations. With this condition the equations for  $v$  and  $w$  in (16) and (19) are decoupled from each other. Therefore the functions  $\underline{v}$  and  $\bar{w}$  are even solutions of (1). In the case of differential equations these are the well known minimal and maximal solutions (see W. Walter 1970, p. 95). Hence (26) is true which implies (8), therefore the bounds  $\underline{v}$  and  $\bar{w}$  are optimal.

A simple example for this case was given in section (0.1). Another example for  $n = 2$  is the system of differential equations

$$(27) \quad \begin{cases} u_1' = 2\sqrt{|u_1|}, & u_1(0) = 0, \\ u_2' = +2\sqrt{|u_1|}, & u_2(0) = 0. \end{cases}$$

One finds easily  $\underline{v}_1(t) = v_2(t) = 0, \bar{w}_1(t) = \bar{w}_2(t) = t^2$ .

12. Further cases with optimal bounds

If one changes in the system (26) the sign in the second equation one gets

$$\begin{cases} u_1' = 2\sqrt{|u_1|}, & u_1(0) = 0, \\ u_2' = -2\sqrt{|u_1|}, & u_2(0) = 0. \end{cases}$$

Now the right hand sides are not anymore "quasimonotone increasing". The set of all solutions can be described quite easily, one finds  $\{\hat{u}_1(t)\} \subseteq [0, t^2]$ ,  $\{\hat{u}_2(t)\} \subseteq [0, t^2]$ .

The maximal interval solution of (19), (20) produces the bounds

$$\underline{v}_1(t) := 0, \underline{v}_2(t) := -t^2, \bar{w}_1(t) := t^2, \bar{w}_2 := 0.$$

They are again optimal bounds. Opposite to the results of section 11 the functions  $\underline{v}$  and  $\bar{w}$  are not anymore solutions of (1) (but certainly of (19)).

By inspection one sees however, that the "crossed" couples  $(\underline{v}_1, \bar{w}_2)$  and  $(\bar{w}_1, \underline{v}_2)$  each are a solution to (1). This is responsible for the optimality. Because if  $\underline{v}$  and  $\bar{w}$  are at least componentwise solutions of (1) then there exist no smaller intervals  $[v, w] \subseteq [\underline{v}, \bar{w}]$  such that (4) is valid. Hence in this case (8) is true which means optimality.

The classes of problems (1), (6) considered in this section are extensions of this example.

Definition (monotonicity matrices): Let  $f \in M$ .

Define for  $i = 1(1)n$

$$a_{ii} := 1,$$

$$a_{ik} := \begin{cases} 0 & \text{if } f_i \text{ does not depend upon } u_k, \\ +1 & \text{if } f_i \text{ depends isotone upon } u_k, \\ -1 & \text{if } f_i \text{ depends antitone upon } u_k \end{cases}$$

for  $i \neq k = 1(1)n$ ,

$$b_{ik} := \begin{cases} 0 & \text{if } f_i \text{ does not depend upon } u_k(\cdot), \\ +1 & \text{if } f_i \text{ depends isotone upon } u_k(\cdot), \\ -1 & \text{if } f_i \text{ depends antitone upon } u_k(\cdot) \end{cases}$$

for  $k = 1(1)n$ .

The matrices  $A = (a_{ik})$  and  $B = (b_{ik})$  are called the monotonic matrices to  $f$ .

Definition (monotonicity condition (M)): Let  $f \in M$ .

Assume the existence of associate<sup>\*</sup> matrices  $A' = (a'_{ik})$ ,  $B' = (b'_{ik})$  to the monotonicity matrices  $A$  and  $B$  such that for all  $i, k = 1(1)n$

$$(28) \quad \left\{ \begin{array}{l} a'_{ik} \in \{+1, -1\}, \quad b'_{ik} \in \{+1, -1\}, \\ a'_{ik} = a_{ik} \quad \text{for } a_{ik} \neq 0, \\ b'_{ik} = b_{ik} \quad \text{for } b_{ik} \neq 0, \\ a'_{ik} = b'_{ik}, \\ a'_{ik} = a'_{li} \cdot a'_{lk}. \end{array} \right.$$

Then  $f$  is said to satisfy the condition (M).

Remark: By (28) one sees immediately

$$a'_{ik} = a'_{li} \cdot a'_{lk} \quad \text{for all } l = 1(1)n.$$

Theorem:

Assume (15). Let  $f \in M$  be continuous and bounded on  $I \times \mathbb{R}^n \times Z$ . Assume that  $f$  satisfies the condition (M). Let  $[v, \bar{w}]$  be the maximal interval solution to (1), (6).

Then  $[v, \bar{w}]$  is even the interval hull of the set of all solutions  $\{\hat{u}\}$  of (1), (6), i.e. (8) is true.

Proof: By construction of  $[v, \bar{w}]$  the inclusion (18) is true. Define the two function vectors  $p = (p_1, p_2, \dots, p_n) = p(t)$ ,  $q = (q_1, q_2, \dots, q_n) = q(t)$  by

$$p_k := \begin{cases} v_k & \text{for } a'_{lk} = +1, \\ \bar{w}_k & \text{for } a'_{lk} = -1, \end{cases}$$

$$q_k := \begin{cases} \bar{w}_k & \text{for } a'_{lk} = +1, \\ v_k & \text{for } a'_{lk} = -1. \end{cases}$$

Certainly  $p_k(0) \in \{a\}$  and  $q_k(0) \in \{a\}$ . Moreover the vectors  $p$  and  $q$  are by construction and by (28) both solutions of (1). Hence  $v$  and  $\bar{w}$  are componentwise composed of solutions of the problem (1), (20). Hence (8) is true.

<sup>\*</sup> These need not be uniquely determined.



REFERENCES

- Moore, R. E., Interval Analysis. Prentice Hall (1966).
- Müller, Max, "Über die Eindeutigkeit der Integrale eines Systems gewöhnlicher Differentialgleichungen und die Konvergenz einer Gattung von Verfahren zur Approximation dieser Integrale. Sitz.-ber. Heidelberg. Akad. Wiss., Math.-Naturw. Kl. (1927), 9.Abh.
- Nickel, K., Fehlerabschätzungs - und Eindeutigkeitsätze für Integro-Differentialgleichungen. Arch. Rat. Mech. Anal. 8 (1961), 159-180.
- Walter, W., Differential and Integral Inequalities. Springer, 1970.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1782 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) BOUNDS FOR THE SET OF SOLUTIONS OF FUNCTIONAL- DIFFERENTIAL EQUATIONS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Karl L. Nickel		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1 (Applied Analysis)
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE August 1977
		13. NUMBER OF PAGES 21
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Functional-differential equations and inequalities Volterra type Solution set Optimal bounds Maximal interval solution		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Sets of systems of ordinary functional-differential equations with Volterra type functionals under sets of initial values are considered. Upper and lower bounds are constructed for the sets of all solutions. Classes of such problems are given where these bounds are optimal. The main tool is a Lemma of Max Müller on inequalities. Also ideas from interval mathematics are used. ue		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)