



DB.S

OFFICE OF NAVAL RESEARCH Contract N00014-76-C-0229 Project No. NR 207-040

TECHNICAL REPORT NO. 122

AD A 0 46445

FILE COPY

IN VITRO EFFECTS OF METHYLPREDNISOLONE SODIUM SUCCINATE AND <u>E. COLI</u> ORGANISMS ON NEUTROPHILS IN BABOON BLOOD

> L. B. Hinshaw, B. K. Beller, J. A. Majde, L. T. Archer, and G. L. White

> > Prepared for Publication

Circulatory Shock



University of Oklahoma Health Sciences Center Departments of Pathology & Physiology & Biophysics Parys Oklahoma City, Oklahoma and

Loyola University Stritch School of Medicine Department of Microbiology Maywood, Illinois

11 October 1977

Reproduction in whole or in part is permitted for any purpose of the United States Government

Distribution of this report is unlimited

OFFICE OF NAVAL RESEARCH Contract N00014-76-C-0229 Project No. NR 207-040

# TECHNICAL REPORT NO. 122

# IN VITRO EFFECTS OF METHYLPREDNISOLONE SODIUM SUCCINATE AND

E. COLI ORGANISMS ON NEUTROPHILS IN BABOON BLOOD

L. B. Hinshaw, B. K. Beller, J. A. Majde,

L. T. Archer, and G. L. White

Prepared for Publication

in

Circulatory Shock

University of Oklahoma Health Sciences Center Departments of Pathology & Physiology & Biophysics Oklahoma City, Oklahoma

and

Loyola University Stritch School of Medicine Department of Microbiology Maywood, Illinois

11 October 1977

Reproduction in whole or in part is permitted for any purpose of the United States Government

Distribution of this report is unlimited

#### ABSTRACT

The corticosteroid, methylprednisolone sodium succinate (MP), has been observed to prevent hypoglycemia in experimental septic shock; however, detrimental actions of various corticosteroids on polymorphonuclear leukocyte function have been reported. The present study was designed to determine if MP depresses glucose metabolism of leukocytes or adversely affects neutrophil survival, or whether it modifies the mortality rate of live <u>E</u>. <u>coli</u> in baboon blood <u>in vitro</u>. Results show that therapeutically effective concentrations (13 µg/ml blood) and high doses (130 µg/ml blood) of MP exert no detrimental influences on glucose utilization or survival of neutrophils in the absence or presence of <u>E</u>. <u>coli</u> organisms in concentrations of  $4.2x10^7$  and  $2.3x10^8$ organisms/ml blood. <u>E</u>. <u>coli</u> organisms, however, increase neutrophil mortality rate and glucose uptake of the blood. These findings support the view that MP does not adversely influence leukocyte metabolism and survival, nor does it modify the mortality rate of live <u>E</u>. <u>coli</u>.



KEY WORDS: septic shock, neutrophils, polymorphonuclear leukocytes, glucose uptake, baboon, <u>E</u>. <u>coli</u> organisms, corticosteroids in shock, methylprednisolone, <u>in vitro</u> blood study

### INTRODUCTION

The incidence of septic shock has increased almost twenty-fold in the United States during the past twenty years (1), and its critical nature has been underscored in a recent review in which over 130,000 deaths per year are estimated to occur from a total number of 330,000 cases (1).

Experimental septic shock has been studied extensively in animals during the past several decades in order to understand the mechanism of septic shock in man. Pathophysiological manifestations in the nonhuman primate administered varying doses of live <u>E</u>. <u>coli</u> organisms include progressively developing systemic hypotension and hypoglycemia in all nonsurviving animals (2). Hypoglycemia has been reported in human septic shock in adults and children (3-5) and in the canine species administered lethal doses of <u>E</u>. <u>coli</u> organisms (6,7). Current findings strongly suggest that hypoglycemia results from the adverse effects of <u>E</u>. <u>coli</u> on hepatic gluconeogenesis (6) and from increased glucose utilization (8,9), including accelerated uptake by the blood in the canine species (10,11) and increased utilization in man (12,13).

Our recent studies were conducted on canine blood to which varying doses of live <u>E</u>. <u>coli</u> were administered <u>in vitro</u>. Results demonstrated consistent accelerated glucose disappearance at all doses of <u>E</u>. <u>coli</u> or endotoxin with significantly greater disappearance rates of glucose at higher dosages (10). Increased utilization of glucose was traced to the white blood cell population (10) with inference that the neutrophil was the cell responsible for the accelerated uptake.

Circulating phagocytic cells, particularly neutrophils, have been shown to exert a potent antibacterial defense in septic shock (14-16). Defects in neutrophil function are considered to have a significant negative impact on the ability to recover from bacterial infections, including septic shock (17,18).

The current consideration of the use of corticosteroids in the treatment of clinical and experimental septic shock is a confusing issue because of the conflict of findings in regard to therapeutic effects and a possible depressant action of the steroids on leukocyte function. Beneficial actions of gluco-corticoids have been reported for the treatment of septic shock in animals (19) and man (20). Several studies, however, have described adverse effects of steroids on polymorphonuclear leukocyte function, including inhibition of chemotaxis (21,22), depression of mobilization (23) and diminished bactericidal capability (24-28). Other reports have shown that certain corticosteroids exert no adverse effects on the bactericidal activities of leukocytes (27,29), including engulfment and killing of enteric bacteria (30), or may enhance their function (31).

The purpose of the present study was to answer the questions, "Does the steroid methylprednisolone sodium succinate (MP) depress glucose metabolism of leukocytes?"; "Does the addition of MP adversely affect neutrophil survival?"; and "Does MP modify the survival rate of live <u>E</u>. <u>coli</u> added to blood <u>in vitro</u>?". Results from these experiments provide evidence for the absence of adverse effects of MP on neutrophil metabolism and survival and indicate that its presence neither accelerates nor depresses the survival rate of live <u>E</u>. <u>coli</u> organisms.

## METHODS

In vitro studies were carried out on the blood from six adult baboons of random sex weighing between 12.6 and 16.2 kg. Each animal was fasted overnight, restrained by means of a squeeze cage device the next morning and administered ketamine hydrochloride (Ketaset; Bristol-Myers Co., Syracuse, N.Y.), 5 mg/kg, intramuscularly. Within 5-10 minutes the baboon was gently lifted

from the cage and blood was drawn from a femoral vein with a needle and plastic syringe wet with heparin (1000 U/ml). The blood was immediately divided into 5 ml volumes in six separate plastic tubes each containing 0.05 ml of heparin (20,000 U/ml), capped, gently mixed and placed on ice. Appropriate volumes of saline or saline plus low or high concentrations of methylprednisolone sodium succinate were then added to three of the tubes of blood, while the remaining three tubes received live E. coli organisms alone or together with low or high doses of the steroid. The six tubes of blood from the same baboon were then simultaneously placed in a water bath at 37°-38°C and incubated. In vitro blood glucose concentrations were determined hourly with a Beckman glucose analyzer (Beckman Instruments; Fullerton, Ca.) possessing an accuracy of ±3 mg%, and each experiment was terminated when blood glucose concentrations fell to an average concentration of 14 mg% (SE, ±1). Initial and final white blood cell (WBC) counts were measured with an automatic particle counter (Coulter  $Z_F$ ; Hialeah, Fla.), and the WBC differentials were estimated by microscopic examination of blood smears stained with Wright's stain.

Preparation of the live organisms for use in these studies was as follows: <u>E. coli</u>, Type B, isolated from a stool specimen at Childrens Memorial Hospital, Oklahoma City, Okla., was maintained in the lyophilized state at 4°C after growth on tryptic soy agar (TSA). The <u>E. coli</u> was reconstituted as needed with 2-3 ml tryptic soy broth and incubated at 37°C for 4-6 hours. Fresh TSA slants were inoculated from the broth suspension using a sterile cotton swab and incubated at 37°C for approximately 18 hours. The cells were then washed from the slants with 2-3 ml of physiological sterile saline (PSS). The washing was centrifuged, the supernatant was discarded and the cells were resuspended in PSS. The cell suspension was then adjusted with PSS to a predetermined

density monitored by reading percentage of transmittance on a Coleman spectrophotometer. <u>E</u>. <u>coli</u> viability counts were done at the beginning and end of all <u>in vitro</u> experiments, using standard serial dilution and pour-plate techniques.

These experiments were designed to evaluate the separate effects of low and high doses of live E. coli organisms and the corticosteroid, methylprednisolone sodium succinate, on glucose uptake and survival of polymorphonuclear leukocytes, specifically neutrophils, in baboon blood in vitro. Accelerated uptake of glucose by the blood in response to E. coli addition was ascribed to increased metabolic activity of the white blood cells in earlier reports (10,32). E. coli concentrations were calculated from data on intact baboon experiments conducted in this laboratory (2). The  $LD_{100}$ ,  $1x10^{10}$  organisms/kg, determined in the earlier in vivo studies, was calculated for use in the present in vitro studies by assuming an in vivo blood volume of 75 ml/kg and instantaneous mixing of organisms in vivo, and the in vitro "high concentration" thus selected was 1.3x10<sup>8</sup> organisms/ml blood. The "low concentration", 1/5 LD<sub>100</sub>, was utilized in order not to overwhelm the phagocytic capacity of neutrophils (70(±12) organisms per neutrophil--LD<sub>100</sub>--vs. 19(±3) organisms per neutrophil--1/5 LD<sub>100</sub>). Methylprednisolone sodium succinate (Solu-Medrol; The Upjohn Company, Kalamazoo, Mich.) was added to blood samples with and without E. coli organisms in concentrations equaling the in vivo half-life concentration in dogs,  $13 \mu$ g/ml serum (unpublished data from The Upjohn Company), and ten times this dosage, 130  $\mu$ g/ml, the latter chosen to determine if high levels of the steroid would be detrimental to the neutrophil. Six samples of blood from the same baboon were simultaneously incubated during a 3-6 hour period, three samples containing saline alone or with low or high doses of steroid, and three receiving E. coli alone or E. coli together with low or high doses of steroid. Blood from the same baboon was alternately used in separate experiments with

high or low doses of <u>E</u>. <u>coli</u> separately added to the blood. Results from all experiments were analyzed using the <u>t</u> test for paired or unpaired data.

## RESULTS

Figures 1A and 1B depict in vitro effects of varying methylprednisolone sodium succinate (MP) and live E. coli organism concentrations on mean cumulative glucose uptake in the blood from six baboons. Results from Figure 1A indicate that neither low nor high concentrations of MP depress glucose uptake of the blood in the absence of E. coli organisms (lower curve); however, during the first two hours glucose utilization increases in contrast to the saline controls (p<0.005). The presence of live organisms (LD<sub>100</sub>; upper curves) results in a markedly elevated glucose uptake during all time periods (p<0.05). High concentrations of MP increase cumulative glucose uptake by the end of the second and third hours (p<0.05). Figure 1B illustrates the in vitro effects of MP and 1/5 LD<sub>100</sub> live <u>E</u>. <u>coli</u> organisms on mean cumulative glucose utilization in the blood from six baboons. Results in the lower three sets of curves in which MP alone was added to blood are indistinguishable from those in Figure 1A (lower curves), the studies having been carried out on the original six baboons, but on different days. In distinction from the results shown in Figure 1A, the addition of a lower dose of E. coli elicits a lesser, though significant, effect on accelerating glucose utilization (p<0.05) with the exception of the first hour. Also, as was the case in Figure 1A, the presence of MP in higher concentrations results in increased glucose utilization in the presence of live organisms (p<0.05). Findings from these data show that neither low nor high concentrations of methylprednisolone depress the cumulative uptake of glucose by the blood during a 5-hour period in the absence or presence of less than lethal to  $LD_{100}$  concentrations of live E. <u>coli</u>

organisms. The addition of organisms, however, elicits notable increases in glucose utilization, quantities approximately doubling in the presence of high concentrations (2.3x10<sup>8</sup> organisms/m1).

The question as to possible effects of MP on neutrophil survival was addressed by data shown in Tables IA and IB. Table IA presents findings pertaining to the in vitro effects of MP and LD100 live E. coli organisms on the blood neutrophil concentration in six baboons. Results show that low and high concentrations of MP do not alter the survival rate of mature and immature neutrophils during a mean in vitro exposure time of 2.6 hours. Addition of E. coli organisms in an LD<sub>100</sub> concentration of 2.3x10<sup>8</sup> organisms/ml blood markedly decreases the numbers of mature neutrophils (p<0.02), and the numbers of degenerated neutrophils are significantly elevated within 2.6 hours (p<0.05). The addition of low and high concentrations of MP to blood containing lethal concentrations of E. coli did not alter the numbers of mature, immature and degenerated neutrophils in comparison to blood containing E. coli alone (p>0.05). Table IB provides data for the in vitro effects of MP and 1/5 LD100 live E. coli organisms on neutrophil mortality in baboon blood. Results show that the addition of low and high concentrations of MP does not alter the survival rate of mature and immature neutrophils during a mean exposure time of 4.3 hours. The duration of experiments was based on the time required for blood glucose to fall below 20 mg%, and times were consistently observed to be directly related to the concentration of E. coli (see Table IA for comparison). Addition of live organisms at a  $1/5 \text{ LD}_{100}$  concentration (4.2x10<sup>7</sup>/ml blood) decreased the numbers of mature neutrophils (p<0.01) and increased the quantities of degenerated neutrophils (p<0.02). Addition of low and high concentrations of MP to blood in the presence of E. coli did not change the numbers of mature, immature and degenerated neutrophils in comparison to blood containing E. coli

alone (p>0.05). These findings indicate that although varying doses of <u>E</u>. <u>coli</u> markedly depress the concentrations of mature neutrophils, incubation with either low or high concentrations of methylprednisolone has no effect on the survival rate of neutrophils. The steroid added to blood in the absence of <u>E</u>. <u>coli</u> does not accelerate the loss of mature and immature neutrophils.

<u>Table II</u> illustrates the effect of MP on <u>E</u>. <u>coli</u> mortality in the blood from six baboons observed <u>in vitro</u>. Results show that high or low concentrations of MP added to blood in the presence of  $LD_{100}$  and 1/5  $LD_{100}$  quantities of live <u>E</u>. <u>coli</u> do not influence the mortality rate of <u>E</u>. <u>coli</u> during incubation times of 2.6 and 4.3 hours, respectively. Mean concentrations of organisms decreased between 40 and 50% during a 2.6-hour period of incubation with  $LD_{100}$  concentrations of <u>E</u>. <u>coli</u> with and without MP and decreased between 16 and 46% with lower concentrations of live organisms during a 4-hour period. These findings indicate that neither low nor high blood concentrations of methylprednisolone exert positive or negative influences on the <u>E</u>. <u>coli</u> mortality rate <u>in vitro</u>.

It was thought to be of interest to determine the effects of MP and <u>E</u>. <u>coli</u> organisms on lymphocyte mortality. Initial concentrations of lymphocytes were compared to final concentrations in each of 12 types of experiments (N=6 in each of the 12 experiments). <u>Table III</u> demonstrates that the numbers of lymphocytes per mm<sup>3</sup> of blood remain relatively constant in all conditions of the experiments during mean incubation periods of 2.6 and 4.3 hours. The only change was a significant increase in mean lymphocyte concentration in the study utilizing the combination of high doses of <u>E</u>. <u>coli</u> and low concentrations of MP (p<0.05), which presumably was due to limitations of accuracy in the counting method.

DISCUSSION

The increasing incidence of septic shock (1) must reflect in part the lack of understanding of its underlying mechanisms. Recent work has emphasized metabolic defects in both its experimental and clinical forms, and in particular

the role of glucose has been emphasized (33). Hypoglycemia has been reported in dogs and nonhuman primates administered live <u>E. coli</u> organisms (2,6,7) and in children and adults subjected to septic shock (3-5). The hypoglycemia is not necessarily a terminal event but may occur during the early course of shock in nonsurviving animals (2,7). Available data suggest that the primary cause of hypoglycemia is failure of hepatic gluconeogenesis (6), and although increased glucose utilization is suspected in clinical septic shock (12,13), published reports from canine studies have documented a significant disappearance of glucose in the blood following live E. coli administration (10,11). In the latter studies (10,11), the white blood cell population was responsible for the accelerated uptake of glucose, and observations implicated the neutrophil as the primary cell accountable. The neutrophil has been shown to exert a potent antibacterial defense in septic shock (14-16,34,35), and defects in its function have been reported to adversely influence the course of bacterial infections and septic shock (17,18). It appears that the circulating phagocyte provides an important defense for the host (14,36,37) but exacts a high metabolic cost in the performance of its role (10,11,38).

Corticosteroids have been studied extensively as a mode of treatment in experimental or clinical septic shock, and although findings are controversial, recent reports with more refined criteria for administration of the agent and its evaluation have shown it to exert a beneficial effect on survival (19,20). However, the actions of steroids on neutrophil function are uncertain since varying adverse effects, including defective mobilization, inhibition of chemotaxis and depression of bactericidal capacity, have been reported (21,22,24-28).

It was thought to be important to characterize the effects of a prominently utilized corticosteroid, methylprednisolone sodium succinate, on neutrophil function and survival and on the mortality rate of live <u>E</u>. <u>coli</u> organisms added

to baboon blood in an in vitro system. Experiments were designed to answer the following questions, "Does methylprednisolone (MP) depress glucose metabolism of leukocytes and survival of neutrophils, and is it able to modify the survival rate of live E. coli?" Differing concentrations of steroid and organisms were employed in order to include most experimental or clinical eventualities. Blood from the same baboons was used in all procedural maneuvers in order to reduce experimental variables to a minimum. The first question to be answered by the present study was, "Does MP depress glucose metabolism of leukocytes?" Findings show that the presence of both low and high concentrations of MP does not depress the uptake of glucose in the blood during a 3-5 hour period in the absence or presence of lethal to less than lethal ( $<LD_{100}$ ) concentrations of live E. coli organisms. The presence of MP with or without E. coli organisms increases cumulative uptake of glucose by the blood during significant portions of the incubation period. The second question addressed in the present study was, "Does the addition of MP adversely affect neutrophil survival?" Findings indicate that incubation of blood in the absence of E. coli organisms with low or high concentrations of MP does not affect the survival rate of mature and immature neutrophils. Adding low or high doses of live E. coli to blood markedly decreases the concentrations of mature neutrophils, but adding MP has no effect on the survival rate of neutrophils. The final question to be answered was, "Does MP modify the survival rate of live E. coli added to blood in vitro?" Findings from the present study showed that low or high concentrations of MP exert neither positive nor negative influences on E. coli mortality. Although E. coli concentrations decreased significantly during incubation with blood, MP had no influence on mortality rates of the organisms during a 3-5 hour period of observation.

Results from the present study suggest that the corticosteroid, methylprednisolone sodium succinate, employed in therapeutic concentrations or greatly exceeding these levels, exerts no demonstrable detrimental direct effects on , neutrophil glucose metabolism or neutrophil survival. The introduction of live <u>E</u>. <u>coli</u> into baboon blood results in a marked increase in glucose consumption, confirming earlier studies using <u>E</u>. <u>coli</u> or endotoxin alone in canine blood (10, 11), and the presence of organisms greatly elevates the mortality rate of neutrophils, which is unaltered by the presence of the steroid in low and high concentrations. In addition to observations on neutrophils, concentrations of lymphocytes remained relatively constant in all experiments. Since the data presented in the present study are confined to <u>in vitro</u> conditions and do not evaluate possible indirect effects of this steroid on blood leukocytes, their application to the <u>in vivo</u> state must await further investigation.

#### REFERENCES

- 1. McCabe WR: Gram-negative bacteremia. Adv Intern Med 19: 135, 1974.
- Hinshaw LB, Benjamin B, Coalson JJ, Elkins RC, Taylor FB, Jr, Priće JT, Smith CW, Greenfield LJ: Hypoglycemia in lethal septic shock in subhuman primates. Circ Shock 2: 197, 1975.
- Berk JL, Hagen JF, Beyer WH, Gerber MJ: Hypoglycemia of shock. Ann Surg 171: 400, 1970.
- Rackwitz R, Jahrmärker H, Prechtel K, Theisen K, Grohmann H: Hypoglykämie während kreislaufshock. Klin Wschr 52: 605, 1974.
- 5. Yeung CY: Hypoglycemia in neonatal sepsis. J Ped 77: 812, 1970.
- Groves AC, Woolf LI, O'Regan PJ, Beach C, Hasinoff C, Sutherland WH: Impaired gluconeogenesis in dogs with <u>E</u>. <u>coli</u> bacteremia. Surgery 76: 533, 1974.
- Archer LT: Hypoglycemia in conscious dogs in live <u>Escherichia</u> <u>coli</u> septicemia: A chronic study. Circ Shock 3: 93, 1976.
- 8. Archer LT, Benjamin B, Lane MM, Hinshaw LB: Renal gluconeogenesis and increased glucose utilization in shock. Am J Physiol 231: 872, 1976.
- Peyton MD, Hinshaw LB, Greenfield LJ, Elkins RC: Hypoglycemic effects of endotoxin in eviscerated dogs. Surg Gynec Obstet 141: 727, 1975.
- Hinshaw LB, Beller BK, Archer LT, Benjamin B: Hypoglycemic response of blood to live <u>Escherichia coli</u> organisms and endotoxin. J Surg Res 21: 141, 1976.
- Hinshaw LB, Beller BK, Archer LT, White GL: Associated leukocyte responses in the lethal aspects of <u>E</u>. <u>coli</u> shock. Proc Soc Exptl Biol Med 155: 179, 1977.

- Roe CF, Kinney JM: The caloric equivalent of fever. II. Influence of major trauma. Ann Surg 161: 140, 1965.
- 13. Kinney JM: Energy expenditure and tissue fuel in the septic patient. In Forscher BK, Lillehei RC, Stubbs SS (eds): "Shock in Low- and High-Flow States." Amsterdam: Excerpta Medica, 1972, p 145.
- Stossel TP: Phagocytosis: The department of defense. New Engl J Med 286: 776, 1972.
- Meakins JL: Pathophysiologic determinants and prediction of sepsis.
   Surg Clin N Am 56: 847, 1976.
- Postel J, Fortado D, Schloerb PR: Effect of prolonged bacteremia on leukocyte bactericidal function. Surgery 81: 180, 1977.
- Solberg CO, Hellum KB: Neutrophil granulocyte function in bacterial infections. Lancet 2: 727, 1972.
- Hellum, KB, Solberg CO: Granulocyte function in bacterial infections in man. Acta Path Microbiol Scand 85: 1, 1977.
- 19. Pitcairn M, Schuler J, Erve PR, Holtzman S, Schumer W: Glucocorticoid and antibiotic effect on experimental gram-negative bacteremic shock. Arch Surg 110: 1012, 1975.
- Schumer W: Steroids in the treatment of clinical septic shock. Ann Surg 184: 333, 1976.
- 21. Ward PA: The chemosuppression of chemotaxis. J Exptl Med 124: 209, 1966.
- 22. Rivkin I, Foschi CV, Rosen CH: Inhibition of <u>in vitro</u> neutrophil chemotaxis and spontaneous motility by anti-inflammatory agents. Proc Soc Exptl Biol Med 153: 236, 1976.
- 23. Peters WP, Holland JF, Senn H , Rhomberg W, Banerjee T: Corticosteroid administration and localized leukocyte mobilization in man. New Engl J Med 286: 342, 1972.

The Alder Second

- 24. Mandell GL, Rubin W, Hook EW: The effect of an NADH oxidase inhibitor (hydrocortisone) on polymorphonuclear leukocyte bactericidal activity. J Clin Invest 49: 1381, 1970.
- 25. Chretien JH, Garagusi VF: Corticosteroid effect on phagocytosis and NBT reduction by human polymorphonuclear neutrophils. J Reticuloendo Soc 11: 358, 1972.
- Olson GE, Polk HC: <u>In vitro</u> effect of ascorbic acid on corticosteroidcaused neutrophil dysfunction. J Surg Res 22: 109, 1977.
- Funfer MM, Olson GE, Polk HC, Jr: Effect of various corticosteroids upon the phagocytic bactericidal activity of neutrophils. Surgery 78: 27, 1975.
- 28. Cline MJ: Drugs and phagocytes. New Engl J Med 291: 1187, 1974.
- Olds JW, Reed WP, Eberle B, Kisch AL: Corticosteroids, serum, and phagocytosis: <u>In vitro</u> and <u>in vivo</u> studies. Infect Immunol 9: 524, 1974.
- 30. Hirsch JG, Church AB: Adrenal steroids and infection: The effect of cortisone administration on polymorphonuclear leukocytic functions and on serum opsonins and bactericidins. J Clin Invest 40: 794, 1961.
- 31. McConn R, Del Guercio LRM: Respiratory function of blood in the acutely ill patient and the effect of steroids. Ann Surg 174: 436, 1971.
- 32. Hinshaw LB, Archer LT, Beller BK, White GL, Schroeder TM, Holmes DD: Glucose utilization and the role of blood in endotoxin shock. Am J Physiol (in press).
- 33. Hinshaw LB: Concise review: Role of glucose in endotoxin shock. Circ Shock 3: 1, 1976.
- 34. Alexander JW, Dionigi R, Meakins JL: Periodic variation in the antibacterial function of human neutrophils and its relationship to sepsis. Ann Surg 173: 206, 1971.

- 35. Postel J, Schloerb PR, Furtado D: Pathophysiologic alterations during bacterial infusions for the study of bacteremic shock. Surg Gynec Obstet 141: 683, 1975.
- 36. Epstein RB, Waxman FJ, Bennett BT, Andersen BR: <u>Pseudomonas</u> septicemia in neutropenic dogs. I. Treatment with granulocyte transfusions. Transfusion 14: 51, 1974.
- 37. Graw RG, Jr, Herzig G, Perry S, Henderson ES: Normal granulocyte transfusion therapy. Treatment of septicemia due to gram-negative bacteria. New Engl J Med 287: 367, 1972.
- Heggers JP, Robson MC, Jennings PB, Fariss BL: Effects of glucose therapy on experimental <u>Escherichia coli</u> septicemia. J Surg Res 20: 33, 1976.

IABLE IA. <u>In Vitro</u> Effects of Neutrophil Con	Methylprednisolone Icentration in Baboo	Sodium Succinat on Blood (Mean±S	E; N=6, each g	100 <u>E</u> . <u>coli</u> Organ group)	I SmS ON
		Neutrophi	S		Total WRC
Experiment	Mature	Immature [	legenerated	Total†	(#/mm <sup>3</sup> )
Init	cial Concentrations	of Neutrophils	(#/mm <sup>3</sup> ; zero †	time; Mean±SE) —	
All experiments	3738(895)	225(100)	(0)0	3964(904)	7504(899)
Final	Concentrations of	Neutrophils (#/	'mm <sup>3</sup> ; +2.6±0.2	hours; Mean±SE) -	
Blood without <u>E</u> . <u>coli</u> a) Saline only b) MP (13 $\mu g/m1$ ) c) MP (130 $\mu g/m1$ ) (a) compared to (b) (a) compared to (c)	3960(776) 3838(690) 3830(597) NS NS	160( 64) 151( 68) 135( 63) NS NS	0( 0) 41( 41) 52( 52) NS NS	4394(802) 4030(644) 4034(666) NS NS	7817(747) 7933(744) 7817(639) p<0.05 NS
Blood with <u>E</u> . <u>coli</u> d) Saline + <u>E</u> . <u>coli</u> e) MP (13 $ug/m1$ ) + <u>E</u> . <u>coli</u> f) MP (130 $ug/m1$ ) + <u>E</u> . <u>coli</u> (d) compared to (e) (d) compared to (f)	1043(208) § 1295(217) § 1333(391) § NS NS	104(38) 117(45) 118(83) NS NS	1731(650) <sup>§</sup> 1364(542) 1829(368) <sup>§</sup> NS NS	2878(631)§ 2776(482) 3280(497) NS NS	7233(494) 7150(557) 7400(604) NS NS
<pre>(a) compared to (d) (b) compared to (e) (c) compared to (f)</pre>	Analysis of Effect p<0.02 p<0.005 p<0.005	t of <u>E</u> . <u>coli</u> on NS NS NS	Concentration p<0.05 NS p<0.005	of Neutrophils - p<0.05 p<0.05 p<0.05	NS p<0.05 p<0.05

\*2.3 X 10<sup>8</sup> E. coli/ml blood (LD100) <sup>†</sup>Total = mature + immature + degenerated neutrophils
<sup>\$p<0.05\$</sup>, initial compared to final concentrations

TABLE IB. In Vitro Effects of	Methylprednisolone	Sodium Succin	ate (MP) and Or	e-Fifth (1/5) LD <sub>1</sub>	00 E. coli
Organisms* on	Neutrophil Mortalit,	y in Baboon B	lood (Mean±SE; N	=6, each group)	
		Neutro	ohils		Total WBC
Experiment	Mature	Immature	Degenerated	Total <sup>†</sup>	(#/mm <sup>3</sup> )
	nitial Concentratio	ns of Neutropl	nils (#/mm <sup>3</sup> ; zer	o time; Mean±SE)	
All experiments	2329(401)	86(44)	20( 20)	2434(402)	5776(330)
Fina	l Concentrations of	Neutrophils	(#/mm <sup>3</sup> ; +4.3±0.3	hours; Mean±SE)	
Blood without <u>E</u> . <u>coli</u> a) Saline only b) MP (13 µg/ml) c) MP (130 µg/ml) (a) compared to (b) (a) compared to (c)	2530(467) 2550(394) 2274(395) NS NS	54(27) 72(18) 58(31) NS NS	39( 28) 31( 22) 116( 66) NS NS	2623(470) 2653(400) 2448(355) NS NS	6067(367)5 6233(352)5 5983(309)5 NS NS
Blood with <u>E</u> . <u>coli</u> d) Saline + <u>E</u> . <u>coli</u> e) MP (13 $\mu$ g/ml) + <u>E</u> . <u>coli</u> f) MP (130 $\mu$ g/ml) + <u>E</u> . <u>coli</u> (d) compared to (e) (d) compared to (f)	1436(426)\$ 1690(220) 1497(310)\$ NS NS	33(24) 141(73) 59(27) NS NS	821(210) § 479(209) 688(324) NS NS	2288(495) 2309(333) 2244(592) NS NS	5633(409) 5750(331) 5817(399) NS NS
Statisti	cal Analysis of Eff	ect of E. col	i on Concentrati	on of Neutrophils	
<ul><li>(a) compared to (d)</li><li>(b) compared to (e)</li><li>(c) compared to (f)</li></ul>	p<0.01 p<0.02 p<0.025	NS NS NS	p<0.02 NS NS	NSSS	NS p<0.001 NS
*4.2 X 10 <sup>7</sup> <u>E</u> . <u>coli/</u> ml blood (1, <sup>†</sup> Total = mature + immature + de	/5 LD <sub>100</sub> ) egenerated neutroph	ils			

<sup>9</sup>p<0.05, initial compared to final concentrations (limitation of method, ±600 cells/mm <sup>3</sup>; Coulter Diagnostics, Inc., Hialeah, Florida)

TABLE II. Effect of Methylprednisolone Sodium Succinate on E. coli Mortality in Baboon Blood In Vitro

		M)	lean±SE; N=6, ea	ch group)	•		
	High concentration	, <u>E</u> . <u>coli</u> —			Low concentration	, E. coli -	
<pre>Initial mean concentration (#E. coli/ml)</pre>	<pre>Final mean concentration (#E. coli/ml)</pre>	% change	Time of incubation (hrs)	<pre>Initial mean concentration (#E. coli/ml)</pre>	<pre>Final mean concentration (#E. coli/ml)</pre>	% change	Time of incubation (hrs)
		Cont	rol (no methylp	rednisolone)			
2.3×10 <sup>8a</sup>	1.3x10 <sup>8b</sup>	-42.3°	2.6	<b>4.</b> 2x107 <sup><i>a</i></sup>	4.0×10 <sup>7b</sup>	-16.4 <sup>d</sup>	4.3
(1.0×10 <sup>7</sup> )	(9.0×10 <sup>7</sup> )	(37.5)	(0.2)	(0.5×10 <sup>7</sup> )	(2.7×10 <sup>7</sup> )	(53.8)	(0.3)
		Methylpredn	isolone concent	ration = 13 g/ml			
2.3x10 <sup>84</sup>	1.2×10 <sup>8b</sup>	-50.3 <sup>c</sup>	2.6	4.2×10 <sup>7a</sup>	2.6×10 <sup>7b</sup>	-46.4 <i>d</i>	4.3
(1.0×10 <sup>7</sup> )	(7.4×10 <sup>7</sup> )	(30.8)	(0.2)	(0.5×10 <sup>7</sup> )	(1.8×10 <sup>7</sup> )	(35.2)	(0.3)
		Methylpredn	isolone concent	ration = 130 g/m			
2.3x10 <sup>84</sup>	1.4×10 <sup>8b</sup>	-39.6 <sup>c</sup>	2.6	4.2x10 <sup>7a</sup>	2.8×10 <sup>7b</sup>	-43.8 <sup>d</sup>	4.3
(1.0×10 <sup>7</sup> )	(7.1×10 <sup>7</sup> )	(29.6)	(0.2)	(0.5x10 <sup>7</sup> )	(2.0×10 <sup>7</sup> )	(37.6)	(0.3)
<sup>a</sup> Significant	difference between	initial value	is of low and hi	gh E. coli concen	trations (p<0.001)		
<sup>b</sup> No significa	nt difference betwee	en final valu	ies of low and h	igh <u>E. coli</u> conce	ntrations (p>0.05).		
<sup>c</sup> No significa	nt difference in per	cent changes	between contro	l and methylpredn	isolone-treated gro	. squo	
d <sub>No</sub> significa	nt difference in per	cent changes	between contro	l and methylpredn	isolone-treated gro	.sdnc	

TABLE III	. In Vitro Effects of	Methylprednisolone	: Sodium Succinate (MP) and	d <u>E. coli</u> Organisms	
	on Lymphocyte Concen	itration in Baboon	Blood (Mean±SE; N=6, each	group)	
	High Dose, E. coli		LOW	Dose, E. coli	
Mean initial concen	tration of Lymphocytes,	. 3410±467/mm <sup>3</sup>	Mean initial concentrat	ion of lymphocytes,	(459±634/mm <sup>3</sup>
	Final concentration of lymphocytes (termination of ex-		Fina of 1 (terr	l concentration ymphocytes mination of ex-	
Experiment	periment; #/mm <sup>3</sup> ±SE)	Significance*	Experiment perir	nent; #/mm 3_SE)	Significance
Saline (no E. coli)	3248(288)	NS	Saline (no <u>E</u> . <u>coli</u> )	3138(288)	NS
MP (13 µg/ml) (no <u>E</u> . <u>coli</u> )	3445(335).	NS	MP (13 µg/ml) (no <u>E</u> . <u>coli</u> )	3373(237)	NS
MP (130 µg/m1) (no E. coli)	3469(524)	SN	MP (130 µg/m1) (no <u>E</u> . <u>coli</u> )	3246(279)	NS
<u>E</u> . <u>coli</u> (2.3x10 <sup>8</sup> /m1 + saline	) 3930(426)	NS	E. coli (4.2x10 <sup>7</sup> /m1) + saline	3098(228)	NS
<u>E</u> . <u>coli</u> (2.3x10 <sup>8</sup> /m <sup>1</sup> + <u>MP</u> (13 µg/m <sup>1</sup> )	) 3851(439)	p<0.05†	<u>E. coli (4.2x10<sup>7</sup>/m1)</u> + MP (13 μg/m1)	3043(196)	NS
<u>E. coli (2.3x10 /ml</u> + MP (130 μg/ml)	) 3624(319)	NS	<u>E</u> . coli (4.2×10 <sup>7</sup> /m1) + MP (130 μg/m1)	3264(357)	NS
*Statistical signif of incubation: hig	icance between initial h dose <u>E. coli</u> group =	and final concentr 2.6±0.2 hr; low do	ations in each of the 12 s se <u>E</u> . <u>coli</u> group = 4.3±0.3	sets of experiments. 3 hr).	(Mean times

19ch

<sup>t</sup>Limitation of method,  $\pm 600$  cells/mm<sup>3</sup> (see Table 1B).

## LEGENDS FOR FIGURES

Figure 1A. In vitro effects of methylprednisolone soch succinate (MP) and LD<sub>100</sub> <u>E</u>. <u>coli</u> organisms on glucose utilization in baboon blood (six experiments, N=6 each experiment).

\* per ml blood

\*\* per ml blood - LD100

\*\*\* addition of <u>E</u>. <u>coli</u> results in increased glucose utilization, hours 1,2,3 (p<0.05)</pre>

Figure 1B. In vitro effects of methylprednisolone sodium succinate (MP) and  $1/5 \text{ LD}_{100} \stackrel{\text{E.}}{=} \frac{\text{coli}}{\text{organisms}}$  on glucose utilization in baboon blood (six experiments, N=6 each experiment).

\* per ml blood

\*\* per ml blood - <LD100

\*\*\* addition of E. coli results in increased glucose utilization, hours 2,3,4 (p<0.05)</pre>



DOCUME	NT CONTROL DATA -	R & D
Security classification of title, body of abstract a	nd indexing annotation must	be entered when the overall report is classified)
		UNCLASSIFIED
OKLAHOMA CITY, OKLAHOMA HEALTH SCIE	INCES CENTER	26. GROUP UNCLASSIFIED
PORT TITLE		
ON NEUTROPHILS IN BABOON BLOOD	DLONE SODIUM SUCCI	INATE AND E. COLI ORGANISMS
DESCRIPTIVE NOTES (Type of report and inclusive date	s)	
AUTHOR(S) (First name, middle in tial, last name)	an ng manana kana na ang mang mga ng mga	
L. B./Hinshaw, B. K./Beller, J. A.	/Majde, L. T./Arc	cher G. L./White
//11 Oct (100 1977)	78. TOTAL NO	224 P. 38
NONTRACT OR GRANT NO.	98. ORIGINAT	OR'S REPORT NUMBER(S)
PROJECT NO.	(1+)T'	R-122 ·)
NR 207-040	9b. OTHER RE	EPORT NO(S) (Any other numbers that may be assigned
d.		
0. DISTRIBUTION STATEMENT		
Distribution of this report is unl	imited.	
 I. SUPPLEMENTARY NOTES	12. SPONSORI	ING MILITARY ACTIVITY
1		Size of New 1 Descende
		FICE OF Naval Research
The corticosteroid, methylprednisolone sodium succinate (MP), has been observed to prevent hypoglycemia in experimental septic shock; however, detri- mental actions of various corticosteroids on polymorphonuclear leukocyte function have been reported. The present study was designed to determine if MP depresses glucose metabolism of leukocytes or adversely affects neutrophil survival, or whether it modifies the mortality rate of live <u>E. coli</u> in baboon blood in vitro. Results show that therapeutically effective concentrations $(13 \mu g/ml blood)$ and high doses $(130 \mu g/ml blood)$ of MP exert no detrimental influences on glucose utilization or survival of neutrophils in the absence or presence of <u>E. coli</u> organisms in concentrations of $4.2x10^7$ and $2.3x10^8$ organisms/ml blood. <u>E. coli</u> organisms, however, increase neutrophil mortality rate and glucose uptake of the blood. These findings support the view that MP does not adversely influence leukocyte metabolism and survival, nor does it modify the mortality rate of live <u>E. coli</u> .		
	1101	THEY Day
D FORM 1472 (PAGE 1)	701	

# OFFICE OF NAVAL RESEARCH BIOLOGICAL SCIENCES DIVISION BIOPHYSICS PROGRAM, Code 444 DISTRIBUTION LIST FOR TECHNICAL, ANNUAL AND FINAL REPORTS

Number of Copies	
(12)	Administrator, Defense Documentation Center Cameron Station Alexandria, Virginia 22314
(6)	Director, Naval Research Laboratory Attention: Technical Information Division Code 2627 Washington, D. C. 20375
(6)	Office of Naval Research Attention: Code 102IP (ONRL DOC) 800 N. Quincy Street Arlington, Virginia 22217
(3)	Office of Naval Research Biophysics Program Code 444 Arlington, Virginia 22217
(1)	Commanding Officer Naval Medical Research and Development Command National Naval Medical Center Bethesda, Maryland 20014
(1)	Chief, Bureau of Medicine and Surgery Department of the Navy Washington, D. C. 20375
(2)	Technical Reference Library Naval Medical Research Institute National Naval Medical Center Bethesda, Maryland 20014
(1)	Office of Naval Research Branch Office 495 Summer Street Boston, Massachusetts 02210
(1)	Office of Naval Research Branch Office 536 South Clark Street Chicago, Illinois 60605



Enclosure (3)

(1)	Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91106
(1)	Commanding Officer Naval Medical Research Unit No. 2 Box 14 APO San Francisco 96263
(1)	Commanding Officer Naval Medical Research Unit No. 3 FPO New York 09527
(1)	Officer in Charge Submarine Medical Research Laboratory Naval Submarine Base, New London Groton, Connecticut 06342
(1)	Scientific Library Naval Medical Field Research Laboratory Camp Lejeune, North Carolina 28542
(1)	Scientific Library Naval Aerospace Medical Research Institute Naval Aerospace Medical Center Pensacola, Florida 32512
(1)	Commanding Officer Naval Air Development Center Attn: Aerospace Medical Research Department Warminster, Pennsylvania 18974
(1)	DIRECTOR Naval Biosciences Laboratory Building 844 Naval Supply Center Oakland, California 94625
(1)	Commander, Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709
(1)	DIRECTOR OF LIFE SCIENCES Air Force Office of Scientific Research Bolling Air Force Base Washington, D. C. 20332

'

Commanding General Army Medical Research and Development Command Forrestal Building Washington, D. C. 20314

Y

(1)

(1)

Department of the Army U. S. Army Science and Technology Center - Far East APO San Francisco 96328

(1)

I

Assistant Chief for Technology Office of Naval Research, Code 200 800 N. Quincy Street Arlington, Virginia 22217

-3-