AD A 0 46582

KINETIC STUDIES FOR XeF AND KrF LASERS

Prepared for OFFICE OF NAVAL RESEARCH

September 1977

MATHEM



EST. INC.

DISTRIBUTION STATEMENT A Approved for public relear 1 Distribution Unlimited

SCIENCES NO

14 Sept 077 Program Code Number TE20 MSNW-1062 fandt STATUS REPORT, NO. 3 QUARTERLY R TITLE: KINETIC STUDIES FOR XEF AND KEF LASERS NOBO14-76-C-1066, DARPA Order-1806 Contract Number: 1 September 1976 Effective Date: Expiration Date: 30 November 1977 Amount: \$99,909 Sponsored by Defense Advanced Research Projects Agency DARPA Order No. 1806 A Principal Investigator: R. E. Center (206) 827-0460 C. H./Fisher/(206) 827-0460 Director, Physics Programs Physical Sciences Division Scientific Officer: Office of Naval Research Department of the Navy 800 North Quincy Street Arlington, Virginia 22117 MATHEMATICAL SCIENCES NORTHWEST, INC. P. 0.Box 1887 Bellevue, Washington 98009 The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U. S. Government. DISTRIBUTION STATEMENT A Approved for public release; Distribution Unlimited 40.4 \$58

TABLE OF CONTENTS

SECTION		PAGE
I	INTRODUCTION	1
II	EXPERIMENTAL RESULTS	2
	2.1. XeF(B^{1}_{2}) Radiative Lifetime	2
	2.2. $XeF(B^{1}/_{2})$ Collisional Quenching	4
	REFERENCES	9

LIST OF FIGURES

FIGURE

PAGE

1	The 353 nm Fluorescence from a Gas Mixture of He + 10 Torr Xe + 0.7 Torr UF6 at a Total Pressure of 125 Torr	3
2	Decay Frequency as a Function of Pressure for a He + 4% Xe + 0.3% UF $_6$ Gas Mixture	5
3	Variation of 353 nm Fluorescence Decay as a Function of Fluorine Partial Pressure	6

MOLOGIUM		-
NTIS GOC UNARROONC	White Service 2 Burn Seather 0 ED 0	-
JUSTIFICAT	tis on fil	
BY	TION /AVAILABILITY COOR	

LIST OF TABLES

1000

TABLE

1

Summary of $XeF(B\frac{1}{2})$ Radiative Lifetime and Quenching Data

PAGE

I. INTRODUCTION

The goal of this program is to determine the radiative lifetime and the collisional quenching rate coefficients for the XeF and KrF upper laser levels using a state selective, laser excited fluorescence technique. Details of the experimental procedure were described in our semi-annual report.⁺ During this reporting period, we have determined the XeF($B\frac{1}{2}$) radiative lifetime by monitoring the decay of the 353 nm XeF($B\frac{1}{2}$) radiative lifetime by monitoring the decay of the 353 nm XeF($B\frac{1}{2}$) fluorescence as a function of pressure. Preliminary collisional quenching rate constants have been determined for He, Xe, and F^{*}₂ from the variation of the fluorescence decay time constant as a function of partial pressure of the collision partner. Collisional quenching by Ar was investigated; however, no bimolecular rate constant was determined because of nonexponential decay of the XeF($B \oplus X$) emission in the presence of Ar. This behavior suggests collisional coupling to some other excited state, most likely the XeF($C\frac{3}{2}$) state.

Bimolecular collisional quenching rate coefficients by Ne and NF₃ remain to be determined during the next reporting period. In addition, termolecular quenching by He, Ne, and Ar will be investigated by increasing the partial pressures of these constituents to the 1 to 3 atm range. Collisional coupling between the B^{1}_{2} and C^{3}_{2} states will also be investigated by looking for fluorescence at 460 nm on the XeF(C + X) band.

II. EXPERIMENTAL RESULTS

2.1. XeF(B ½) Radiative Lifetime

Oscilloscope traces showing the 351 nm exciting pulse and the 353 nm fluorescence signal are presented in Figure 1. This signal was obtained with a He + 10 torr Xe +0.7 torr UF₆ gas mixture at 125 torr total pressure. We have switched to UF₆ as the F atom source because its UV absorption cross section (resulting in dissociation to produce an F atom) is two orders of magnitude larger than that of F_2 , which yields a substantial increase in the XeF fluorescence. However, the UF₅ produced during the flash has a low vapor pressure resulting in the formation of solid particles which increased the scattered light from the 351 nm exciting pulse on subsequent shots. Therefore, it was necessary to pump out the cell and refill it with fresh gas after every shot.

If the XeF(B $\frac{1}{2}$) excited state is lost only by radiative decay and bimolecular collisions, it will decay exponentially once the exciting laser pulse has terminated. A semilogarithmic plot of the 353 nm fluorescence decay read from the oscilloscope trace is presented in Figure 1. The slope of the straight line fit to the data yields the decay time constant τ^{-1} given by

 $\tau^{-1} = \tau_{B}^{-1} + k_{He} [He] + k_{Xe} [Xe] + k_{UF_{6}} [UF_{6}]$ $= \tau_{B}^{-1} + (k_{He}X_{He} + k_{Xe}X_{Xe} + k_{UF_{6}}X_{UF_{6}}) p$

where τ_{B} is the radiative lifetime of the XeF(B $\frac{1}{2}$) state, k_{i} and X_{i} are the quenching rate coefficient and the mole fraction of the ith component of the mixture, and p is the total pressure. Thus, a plot of the decay time constant τ^{-1} as a function of pressure for a





The 353 nm Fluorescence from a Gas Mixture of He+10 Torr Xe + 0.7 torr UF₆ at a Total Pressure of 125 torr. a) Oscillo-scope traces showing the 351 nm exciting laser pulse and the 353 nm fluorescence signal. b) Semi-logarithmic plot of the 353 nm fluorescence decay.

gas mixture with fixed composition should follow a straight line with the XeF(B $\frac{1}{2}$) radiative lifetime as the zero pressure intercept. A plot of the data obtained for a He + 4% Xe + 0.13% UF₆ gas mixture is shown in Figure 2. The zero pressure intercept of the straight line fit to the data gives a XeF(B $\frac{1}{2}$) radiative lifetime of 18.7 ± 1.4 nsec. This result is in good agreement with that obtained by Burnham and Harris from the dissociative excitation of XeF₂.²

2.2. XeF(B ½) Collisional Quenching

Bimolecular quenching rate constants for various molecules are obtained by measuring the XeF (B $\frac{1}{2}$) decay time constant as a function of the partial pressure of that constituent while holding the partial pressures of the other constituents fixed. The slope of the straight line fit to the data gives the two-body quenching rate constant for that molecule. To illustrate the procedure, a plot of the 353 nm decay time constant τ^{-1} as a function of F₂ partial pressure is shown in Figure 3. The partial pressures of the other constituents of the gas mixtures were held fixed at 250 torr He + 10 torr Xe + 0.3 torr UF_6 . The quenching rate constant for F_2 determined from the slope of the straight line fit to the data is $k_{F_{-}} = 1.2 \times 10^{-10} \text{ cm}^3$ molecule⁻¹ sec⁻¹. Using this same procedure, quenching rate constants for He and Xe have also been determined. These rate constants are listed in Table 1 along with the pressure range over which the quenching was studied. The measured rate constants are in fairly good agreement with those inferred by Brashears, Setser and Desmarteau from Stern-Volmer plots of the quenching of XeF(B $\frac{1}{2} \rightarrow X$) emission produced by photolysis of XeF2.3

No quenching rate constant is listed in the table for Ar even though it is a primary constituent of e-beam excited XeF laser



Figure 2. Decay Frequency as a Function of Pressure for a He + 4% Xe + 0.13% UF₆ Gas Mixture. The zero pressure intercept of the straight line fit to the data gives a XeF (B_2^1) radiative lifetime of 18.7 nsec.

Figure 3. Variation of 353 nm Fluorescence Decay as a Function of Fluorine Partial Pressure. The slope of the straight line fit to the data gives a quenching rate constant of 1.2 x 10^{-10} cm³ molecule⁻¹ sec⁻¹.

	XeF(B $\frac{1}{2}$) $\tau_r = 18.7 \pm 1.4$ nsec		
Molecule	kq(cm ³ Molecule ⁻¹ sec ⁻¹)	Pressure Range	
Не	2 x 10 ⁻¹²	75 - 750	
Xe	6 x 10 ⁻¹¹	5 - 80	
F ₂	1.2×10^{-10}	0.5 - 24	

Table 1

Summary of XeF (B 1/2) Radiative Lifetime and Quenching Data

mixtures. This is due to the fact that the 353 nm fluorescence in the presence of Ar exhibited nonexponential decay over much of the pressure range studied, making it difficult to extract a rate constant without a more complicated analysis. This behavior suggests collisional transfer from the $B^{1}/_{2}$ state during the decay period. The most likely candidate is the XeF($C^{3}/_{2}$) state which is nearly degenerate with the $B^{1}/_{2}$ state.⁴ During the next reporting period, the possibility of collisional coupling between the $B^{1}/_{2}$ and $C^{3}/_{2}$ states will be investigated by looking for broadband fluorescence at 460 nm on the XeF($C^{3}/_{2} \rightarrow X$) transition.

REFERENCES

- R.E. Center and C.H. Fisher, "Kinetic Studies for XeF and KrF Lasers," MSNW Semi-Annual Report, May 1977.
- 2. R. Burnham and N.W. Harris, J. Chem. Phys. <u>66</u>, 2742 (1977).
- 3. H.C. Brashears, Jr., D.W. Setser, and D. Desmarteau, Chem. Phys. Lett. <u>48</u>, 84 (1977).
- 4. T.H. Dunning and P.J. Hay, to be published.