AD-A046 865

FOREIGN TECHNOLOGY DIV WRIGHT-PATTERSON AFB OHIO
INTEGRAL EQUATION OF THE KINETIC THEORY OF THE COAGULATION OF C--ETC(U)
JUN 77 I M YENUKASHVILI
FTD-ID(RS)T-1055-77

NL

END
RUTTO

PD-A046865

FTD-ID(RS)T-1055-77

0

FOREIGN TECHNOLOGY DIVISION



INTEGRAL EQUATION OF THE KINETIC THEORY OF THE COAGULATION OF CLOUD FARTICLES

by

I. M. Yenukashvili



Approved for public release; distribution unlimited.

UNEDITED MACHINE TERANSLA

FTD-ID(RS)T-1055-77

28 June 1977

MICROFICHE NR: 740-77-C-000 762

INTEGRAL EQUATION OF THE KINETIC THEORY OF THE COAGULATION OF CLOUD FARTICLES

By: I. M. Yenukashvili

English pages: 12

Source:

Trudy VIII Vsesoyuznoy Konferentsii po Fizike Oblakov i Aktivnym Vozdeystviya, Leningrad, 1970, PP. 470-475

Country of origin: USSR

This document is a machine translation

Requester: FTD/PHE

Approved for public release; distribution unlimited

ACCRECION O	
erne	Witte Settles X
100	Buff Section (1)
MANNOWICE	0
JUSTIFICATIO	1
97	M/ATAILABILITY COOL
Blot	AVAIL and/or Special
	The second secon

THIS TRANSLATION IS A RENDITION OF THE ORIGI-MAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DI-

PREPARED BY:

TRANSLATION DIVISION FOREIGN TECHNOLOGY DIVISION WP-AFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Block	Italic	Transliteration	Block Italic	Transliteration
Aa	A a	A, a	Pp	R, r
5 6	5 6	B, b 800	C c . C .	S, s
Вв	B 4	V, v +k+	T T . T . M	T, t
Гг	r .	G, g don	уу у у	U, u
Дд	д в	D, d	ф 🏓 🍎	F, f
Еe	E .	Ye, ye; E, e*	X x X x	Kh, kh
жж	Ж ж	Zh, zh	Цц 4 4	Ts, ts
З э	3 1	Z, z	4 4 4 4	Ch, ch
Ии	и и	I, i	шш Ш ш	Sh, sh
Йй	яа	Y, y	Щщ Щ щ	Sheh, sheh
Нн	K *	K, k	ъъ 3 ъ	•
Лл	ЛА	L, 1	Ыы Ы и	Y, y
M M	M M	M, m	ьь Б .	
Нн	H ×	N, n	3 9 9 1	Е, е
0 0	0 .	0, 0	Ю ю ю	Yu, yu
Пп	Лж	P, p	Яя Яя	Ya, ya

*ye initially, after vowels, and after ъ, ъ; e elsewhere. When written as ë in Russian, transliterate as yë or ë. The use of diacritical marks is preferred, but such marks may be omitted when expediency dictates.

GREEK ALPHABET

Alpha	Α	α	•	Nu ota	N	ν	
Beta	В	β		d Xi Sha	Ξ	ξ	
Gamma	Γ.	Υ	4	Omicron	0	0	
Delta	Δ	δ		Pi	Π	π	
Epsilon	E	ε	C (Figure 17)	Rho	P	ρ	•
Zeta	Z	ζ		Sigma	Σ	σ	•
Eta	Н	'n		Tau	T	τ	
Theta	Θ	θ	•	Upsilon	T	υ	
Iota	I	TA	10510 30	Phi	Φ	φ	ф
Kappa	('K ⋅	×	400 and	0 Ch1	X	χ	(A
Lambda	٨	λ	TISTAGE?	Psi	¥	Ψ	9 % 4 %
Mu	M	μ		Omega	Ω	w	

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

. .

M. F.

Q A

1 1

*

C - (5

175

its water with the

resident med opti w Modeler von de

2384

T. TIPLOTERE T.

* N

7, 4 7, 7 XI, 10

01 . 57

da de

*

188

n

135

*

Russian	English
sin -	f sin
cos	cos
tg	tan
ctg	cot
sec	sec
cosec	csc
sh	sinh
ch	cosh
th	tanh
cth	coth
sch	sech
csch	csch
arc sin	sin ⁻¹
arc cos	cos ⁻¹
arc tg	tan ⁻¹
arc ctg	cot ⁻¹
arc sec	sec ⁻¹
arc cosec	csc ⁻¹
arc sh	sinh ⁻¹
arc ch	cosh-1
arc th	tanh-1
arc cth	* coth
arc sch	sech 1
arc esch	csch ⁻¹
14	10.00
504	
rot	curl
lg	log

GRAPHICS DISCLAIMER

All figures, graphics, tables, equations, etc. merged into this translation were extracted from the best quality copy available.

PARTICLES.

I. H. Yenukashvili.

Page 470.

Investigations in the theory of transfer of cloud particles in considerable degree are bonded with the development of the methods of the solution to kinetic equations. In this respect by very promising is presented writing of the kinetic equation of coagulation in the form of integral equation. In particular this is exhibited during the solution to the boundary-value problems of the kinetics of coagulation. This integral equation, including boundary conditions, in very compact form contains entire physics of the process of the transfer of cloud particles and can render/show more convenient than integrodifferential equation. Let us note that the integral kinetic equations successfully are utilized in the theory of transfer of neutrons [1], in aerodynamics of rarefield gases [2, 3].

The transport problem of cloud particles - this the problem of many particles and with study of the formation of the spectrum of

cloud particles in space and in time must be considered their absolute motions, mutual collisions and merging/coalescences.

Simultaneously with this the value of the distribution function of cloud particles in point with radius-vector r (x, y, z) they must be bonded not only with the values of distribution function in the points, close toward the end of vector r, but also with values in all points of cloud, including interfaces.

Let n (v, r, t) be the distribution function of cloud particles according to volumes v at the moment of time t about point in space with radius-vector r, so that value

dn == n(v, r, t) d2 dv

(1)

is for the moment of time t the mathematical expectation of the number of cloud particles, which have radius-vector and range r, r + dr and volume in the range v, v + dv; dQ = dx dy dz - elementary three-dimensional/space common/general/total, that adjoins toward the end of radius-vector r.

Distribution function n (v, r, t) in question makes sense of the density of the mathematical expectation of the number of aloud particles; however, together with this it simultaneously has also the sense of the density of distribution of the probability of the detection of one particle in the four-dimensional volume dvdQ.

Let us isolate in cloud the small volume element $d\Omega$ and determine in this volume about point with radius-vector r for the arbitrary moment of time t, the number of cloud particles with volume in the range v, v + dv, i.e., let us determine dn = n (v, r, t) $d\Omega$ dv.

Page 471.

Let \$\Pi\$ (v, r, t, v) be a probability of free motion in the time interval (v, t) of this cloud particle, which at the moment of time v is located in point with radius-vector r- (t-v) with and at the moment of time t proves to be at the point of space with radius vector r. Here c is speed of the motion of the cloud particle, which has volume v, which depends on v, and also on the velocity of the

sotion of air masses in cloud. The generalization of the physical derivation of the system of integral equations of the kinetics of the coagulation of cloud particles in the case of the accelerated motion of cloud particles presents no difficulties [4]. Special examination requires the case of the motion of cloud particles in the turbulent flow of air masses.

Let us examine certain cloud particle at certain moment of time to Let it is locate in point with radius-wector r and has volume v and speed c. It is possible to assert that at certain moment of time v < t it experienced the last/latter collision and that it as the cloud particle, possessing volume v and speed c, it was born in point with radius-vector r- (t-v) c at the moment of time v.

Further, let (v, r, t) be a function of generations, so that

 $dn_1 = \Phi(v, r, t) d2 dv dt \tag{2}$

gives the mathematical expectation of the number of cloud particles, which were being born in the elementary volume of space dR about point with radius vector r for a period of time dt and which have volume in the range v, v + dv.

Let us express function n (v, r, t) through the function of

generations and the probability of free aption. At the moment of time t > to (to - the initial soment) the mathematical expectation of the number of cloud particles with a volume of v, v + dv in the volume element of space 48 is composed of two pasts.

 $n(v, r, t) dv dQ = dn_0 + dn_2$

(3)

where dno is a mathematical expectation of the number of cloud particles with volume in the range v, v + dv, which reached in dQ from the initial state without collisions; dno is a mathematical expectation of the number of cloud particles with a volume of v, v + dv, which appear in cloud in time interval from to to t at the different moments of time v at the different points of space with radius-vector r- (t-v) c and which without collisions reach dQ up to the moment of time t.

It is obvious that

 $dn_0 = n [v, r - (t - t_0)c, t_0] \Pi(r, v, t, t_0) dv d2.$ (4)

Page 472.

Then, according to determination (2), value

will be equal to the mathematical expectation of the number of cloud particles, which are born within volume dΩ about point with radius-vector r- (t-v) c for a period of time dv and have a volume in the range v, v + dv. Prom this quantity of cloud particles to element of volume dΩ about the points with radius-vector r will r ach only the those, that for the extent/elongation of time (v, t) do not experience collisions. Therefore, if we expression (5) multiply by the probability of free motion Π (r, v, v, t) and to integrate over alternating/variable v from t₀ to t we will obtain the total quantity of cloud particles with a volume of v, v * dv, which apper in cloud in time interval (t₀, t) and the moment of time t enterelement of volume dΩ about point with radius-vector r. Thus,

$$dn_2 = dQ \, dv \int_L^t \Phi \left[v, \, r - (t - \tau) \, c, \, \tau \right] \cdot \Pi \left(r, \, v, \, \tau, \, t \right) d\tau. \tag{6}$$

Special examination requires the case, when occurs the emergence of cloud particles on interface, for example at the level of condensation. For those regions of the clouds, into which can enter cloud particles, which arcse on interface in time interval (t₀, t), in relationship/ratio (3) one should write the additional term, which considers a quantity of cloud particles with volume in the range v, v · dv, which appear on interface at the appropriate moment of time and which without collision reach dΩ about point with radius-vector r up to the moment of time t. In this case, it goes without saying, the boundary function of the generations is assumed known.

Substituting (4) and (6) in (3), we obtain

$$n(v, r, t) = n[v, r - (t - t_0)c, t_0] \Pi(r, v, t, t_0) +$$

$$+ \int_{t_0}^{t} \Phi[v, r - (t - \tau)c, \tau] \Pi(r, v, \tau, t) d\tau.$$
 (7)

Por obtaining the closed system of integral equations it is necessary to express the probability of free action and the function of generation by distribution function.

Let ΛQ_i be a probability of the random event, which consists of the fact that during time interval $\Lambda \tau_i$ examined cloud particle is adjacent to another cloud particle.

Page 473.

then we have

$$\Delta Q_1 = \Delta z_1 \int_0^{\infty} n |u, r + z_1 c, l + z_1| \sigma(v, u) du,$$

where

$$\sigma(v, u) = \pi E\left(\frac{3}{4\pi}\right)^{2/3} (v^{1/3} + u^{1/3})^2 |c(v) - c(u)|$$

- the effective volume of the collisions of the cloud particles, which have volumes v and u; E is a coefficient of capture

With small $\Delta \tau_i$ the probability of free motion for the cloud particle in question during time interval $\Delta \tau_i$ will be equal to

$$P_i = 1 - \Delta Q_i = e^{-\Delta Q_i}.$$

Then the probability of the free action of the cloud particle in question for a period of time $T = \sum_{i=1}^{n} \Delta \tau_i$ wild be

$$P = \prod_{i=1}^{n} P_{i} = \exp\left\{-\sum_{i=1}^{n} \Delta Q_{i}\right\} = \exp\left\{-\sum_{i=1}^{n} \left[\int_{0}^{\infty} n(u, \tau + \tau_{i}c, t + \tau_{i})\sigma(v, u) du\right] \Delta \tau_{i}\right\}.$$

Hence, replacing summing by integration, we obtain

$$\Pi(v, r, \tau, t) = \exp\left\{-\int_{0}^{t} \left[\int_{0}^{\infty} \sigma(v, u) n(u, r - (t - q)c, q) du\right] dq\right\}.$$
(8)

The last/latter relationship/ratio gives communication/connection

between the probability of free motion and the distribution function of cloud particles.

To the generation of cloud particles with a volume of v in the volume of space do for time dt give the collisions of cloud particles from volumes v-u, u, with this u it can take values from 0 to v.

Therefore the collision frequency, summed up in terms of all possible values of u, gives interesting us the mathematical expect tion of the number of generations of cloud particles with a volume of v, thus

$$\Phi(v, r, t) = \frac{1}{2} \int_{0}^{\pi} n(v - u, r, t) n(u, r, t) \sigma(v - u, u) du.$$
 (9)

Expression • (v, u) reflects mechanics and the statistics of the reaction between the pair of cloud particles.

equations (7), (8), (9) are the system of the integral equations which characterize the kinetics of the coagulation of cloud particles. The structure of these equations is such, that easily it is possible to produce the exception/elimination of functions !!, (1) and to obtain one integral equation for the distribution function of cloud particles.

Page 474.

BEST AVAILABLE COPY

Substituting (8), (9) in (7), we have

BEST AVAILABLE COPY

$$n(v, r, t) = n\{v, r - (t - t_0)e, t_0 | \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} + \int_t^{\infty} \exp\left\{-\int_t^{\infty} \left[\int_0^{\infty} \sigma(v, u)n(u, r - t_0)e, t_0\right] du\right\} dq\right\} dq$$

Let us use the operation

$$\frac{d}{dt} = \frac{\delta}{\delta t} + (\mathbf{c} \cdot \underline{\mathbf{A}})$$

to both parts of equation (7); taking into account the relationship/ratios:

$$\frac{d}{dt} \left\{ n \left[v, \dot{\tau} - (t - t_0 c, t_0) \right] = 0, \right.$$

$$\Pi \left(r, v, t \right) = \Delta,$$

$$\frac{d\Pi}{dt} = -\Pi \int_{-\pi}^{\pi} e(v, u) n(u, \tau, t) du,$$

ous aidean

BEST AVAILABLE COPY

we obtain

$$\frac{dn}{dt} = \Phi(v, r, t) - \int_{0}^{\infty} \sigma(v, u) n(u, r, t) du \left\{ \int_{0}^{\infty} \Phi[v, r - (t - v)c, v] \times \right.$$

$$\times \Pi(r, v, v, t) dv + n[v, r - (t - t_{0})c, t_{0}] \Pi(r, v, t, t_{0}) \right\}.$$

If, according to (7), expression in curly braces in the last/latter relationship/ratio is replaced through n (v, r, t), and function $^{\oplus}$ (v, r, t) - with expression (9), then we will obtain the known integrodiff equation of the kinetic theory of the coagulation of the cloud particles

$$\frac{dn}{dt} = -n(v, \tau, t) \int_{0}^{\infty} \sigma(v, u) n(u, \tau, t) du + \frac{1}{2} \int_{0}^{\infty} \sigma(v - u, u) n(v - u, \tau, t) n(u, \tau, t) du,$$

DOC = 77103601

BEST AVAILABLE COPY

As the asserical methods of the approximate solution of the system of integral equations (7), (8), (9) it is possible to utilize a method successive approximation, an iterative method, and also a method of torque/moments [5, 6]. Accepting as zero approximation either the imitial distribution or any analytical solution of the simplified problem of the kinetics of the coagelation of cloud particles, we compute the function of generaties on and the probability of free notion Π_0 . Substituting Φ_0 and Π_0 in equation (7), we obtain the first approximation of distribution function ng. Purther we compute Ot. II. 10. The fellowing approach/approximations are constructed analogously.

In conclusion let us note that the proposed above form of the notation of the physical derivation of the system of the integral egentiese of the himetics of the congulation of cloud particles can subconstally used also in the exemination of the timetics of the coagulation of dispersed particles and, is particular accessi particles.

LAYER TIME BIBLIOGRAPHY

ANTEPATYPA

^{1.} Дэвисо в В. Теория переноса нейтронов. Атомиздат, М., 1960. 2. Ввяландер С. В. Уравнения и постановка задач в аэродинамике разреженных газов. Аэродинамика разреженных газов. Информ. сб. ЛГУ, № 1.

Филиппов В. В. Вариант нестационарных кинетических уравнений. Аэродинамика разреженных газов. Информ. сб. ЛГУ, № 1, 1963.
 Велова А. В., Валавидер С. В. Интегральные кинетические уравнении

теории одноатомных газов при наличии внешнего поля массовых сил. Аэродинамика разреженных газов. Информ. сб. ЛГУ, 1963.

5. В а р а и ц е в Р. Г. Метод интегральных моментных кинетических уравнений. ДАН СССР, т. 151, № 5, 1963.

6. Е в у в а и в в а и М. М. О решения авменческого уравнения коагуляции. Изв. АН СССР, скр. гоофия. № 16, 1664.

REPORT DOCUMENTATION PAGE	BEFORE COMPLETING FORM ESSION NO. 3. RECIPIENT'S CATALOG NUMBER
FTD-ID(RS)T-1055-77	
. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED
INTEGRAL EQUATION OF THE KINETIC TH	EORY Translation
OF THE COAGULATION OF CLOUD FARTICL	ES 8. PERFORMING ORG. REPORT NUMBER
AUTHOR(e)	S. CONTRACT OR GRANT NUMBER(s)
	053 AF/INARA 017 AF/ MEXTRAG
	404 Ache
Foreign Technology Division	408 PROGRAM ELEMENT, PROJECT, TASK 410 ADYC
Air Force Systems Command U. S. Air Force	413 400
CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
A AFRANTON SYS CLARE 1	1970
MONITORING ASENCY NAME & ADDRESS(II dillerent from Controllis	12
CHARLES AND AND AND A ADDRESS OF MINORAL PAIN CONTROLL	g Office) 18. SECURITY CLASS. (or this report)
D (CAME (CRAMMER)	UNCLASSIFIED
C1A/C9C/ADD/SD 1	154. DECLASSIFICATION/DOWNGRADING SCHEDULE
DISTINGUTION STATEMENT (of this Ropert)	
Approved for public release; distri	bution unlimited
476D	
DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if	illferent from Report)
SUPPLEMENTARY NOTES	
KEY WORDS (Continue on reverse side if necessary and identify by bit	
REY WORLDS (Commiss on Over20 also it recorded y and recording by Sir	
ABSTRACT (Continue on reverse side if necessary and identify by blo	
ABSTRACT (Continue on reverse side if necessary and identify by blo	
20 04 10	
20, 04, 12	
20, 04, 12	
²⁰ ; 04; 12	

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

ORGANIZATION		MICROFICHE	ORGANIZATION		MICROFICHE	
A205	DMATC	1	E053	AF/INAKA		
A210	DNAAC	2	E017			
B344	DIA/RDS-3C	2 8		AEDC		
CO43	USAMITA	1		AFWL		
	BALLISTIC RES LABS	1	CERTIFICATION CONTRACTOR SECTION CONTRACTOR SECTION SECTION SE	ADTC		
	AIR MOBILITY RAD	1	E413	ESD	2	
	LAB/FIO			PTD		
C513	PICATINNY ARSENAL	1		CCN		
	AVIATION SYS COMD	ī		ETID	1	
C557	USATIC	1		NIA/PHS	3	
	FSTC	5		NICD	1 5	
	MIA REDSTONE	í		ALUD	3	
D008	NISC					
	USAICE (USAREUR)					
	ERDA	Ī				
	CIA/CRS/ADD/SD	7				
	DSTA (50L)					
	MSCEN (Code 121)	† •				
MASA/		•				
HADA	· .					
AFIT/	TÔ.	1				