

AD A 0 4 6 9 6 9

12
R

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0807

TECHNICAL REPORT NO. 13

Hydrolysis of Enkephalin by Cultured
Human Endothelial Cells and by Purified Peptidyl Dipeptidase

Prepared by

Ervin G. Erdős

Submitted for Publication

Departments of Pharmacology and Internal Medicine
University of Texas Health Science Center at Dallas
5323 Harry Hines Boulevard
Dallas, Texas 75235

04 November 1977

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Distribution of this report is unlimited

DDC
NOV 28 1977
F

AD No. _____
DDC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report 13 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) HYDROLYSIS OF ENKEPHALIN BY CULTURED HUMAN ENDOTHELIAL CELLS AND BY PURIFIED PEPTIDYL DIPEPTIDASE	5. TYPE OF REPORT & PERIOD COVERED Technical 6/1/77 - 12/15/78	
7. AUTHOR(s) Ervin G. Erdős, Alice R. Johnson, Nit T. Boyden	8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0807	
PERFORMING ORGANIZATION NAME AND ADDRESS Univ. TX Health Sci, Center at Dallas ✓ Dept. of Pharmacology 5323 Harry Hines Blvd., Dallas, TX 75235	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PHS-HL-14187	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Biological and Medical Sciences Division Medical and Dental Sciences Program, Code 444 Arlington, VA	12. REPORT DATE 11/04/77	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Technical rept., 1-2-78-15-2-88	13. NUMBER OF PAGES 10	
	15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of this Report) Distribution of this document is unlimited 12 12p. 14 TR-13	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)	11 4 Nov 77	
18. SUPPLEMENTARY NOTES Submitted for publication.	D D C RECEIVED NOV 28 1977 F	
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Amino peptidase, vascular endothelium, converting enzyme, kininase, opioid		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The experiments indicate that rapid inactivation of enkephalins may be due to at least two different enzymes present in tissues. A peptidyl dipeptidase which is a component of plasma membrane of various cells may degrade enkephalins by liberating a C-terminal dipeptide and an amino peptidase in endothelial cells may cleave peptides, that they are taken up as blood flows past the endothelial surface, at the N-terminal end.		

408 854

18

Hydrolysis of Enkephalin by Cultured Human Endothelial Cells
and by Purified Peptidyl Dipeptidase

Ervin G. Erdős, Alice R. Johnson, Nit T. Boyden

Departments of Pharmacology and Internal Medicine
University of Texas Health Science Center at Dallas
Dallas, Texas 75235

Send Galley Proofs to:

ACCESSION for	
NTIS	
DDC	
UNANNOUNCED	
JUSTIFIED	
BY	
DISTRICT	
A	

The biological activities of enkephalins are of great interest to numerous investigators since they have an affinity for opiate receptors and may mediate perception of pain¹. Studies of enkephalins have been hampered, however, because these endogenous pentapeptides are rapidly degraded by blood and tissues. For example, radiolabeled enkephalin is hydrolyzed in less than one minute when it is injected intraventricularly in rats².

Peptides transported in blood are continuously in contact with the vascular endothelium. These cells are more than just a diffusion barrier between blood and tissues since they contain enzymes that metabolize various peptides. Enzymes in vascular endothelial cells grown in culture cleave peptides such as bradykinin, angiotensin I and II³ and substance P⁴. Two of the enzymes in cultured human endothelial cells were identified as an aminopeptidase and a peptidyl dipeptidase³. These two enzymes have important functions in the control of both hypotensive and hypertensive peptides. The aminopeptidase may hydrolyze angiotensin I or II by cleaving the N-terminal aspartic acid. The peptidyl dipeptidase cleaves a dipeptide from the C-terminal end of bradykinin and angiotensin I. By this action it converts angiotensin I to II and inactivates bradykinin^{5,6}. In addition to endothelial cells, the brush border of the kidney⁶ and various parts of the central nervous system contain peptidyl dipeptidase activity^{7,8}. Because the substrate specificity of this enzyme (E.C. 3.4.15.1; kininase II or angiotensin I converting enzyme) indicated⁶ that enkephalins may also be cleaved, we investigated the action of purified enzyme on synthetic leucine- and methionine-enkephalins. We determined the peptide bond hydrolyzed and also characterized the mode of inactivation of enkephalins by cultured human endothelial cells.

Leu⁵-enkephalin (Tyr¹-Gly²-Gly³-Phe⁴-Leu⁵; Leu-Enk) and Met⁵-enkephalin (Met-

enk) were obtained from commercial sources. Endothelial cells from veins of human umbilical cords were grown in monolayer cell cultures as described ³. Peptidyl dipeptidase was purified to homogeneity by the method of Oshima et al. ⁹. Leucine aminopeptidase (E.C. 3.4.1.1) was purchased from Boehringer GmbH. Leu-enk or Met-enk (0.25 μmol) was incubated with 1.2 μg peptidyl dipeptidase or 0.12 μg aminopeptidase at 37° in 0.06 M NH_4HCO_3 buffer, pH 7.4. The molar ratio of peptidyl dipeptidase to substrate was 1 to 30,000. Cultured human endothelial cells were removed from monolayers by gentle scraping with a rubber spatula. These cells were then suspended and used in a concentration of one million cells per ml. The products of hydrolysis of enkephalins were determined by thin layer chromatography (t.l.c.).

The rate of cleavage of Met-enk by peptidyl dipeptidase was determined in a programmed amino acid analyzer by separating Tyr-Gly-Gly and Phe-Met on a 5.5 cm column in 0.2 N citrate buffer, pH 4.25. The column was calibrated with the synthetic tripeptide and dipeptide. Substrate (0.2 $\mu\text{mol/ml}$) and enzyme (2.4 $\mu\text{g/ml}$) were incubated at 37° and pH 7.4 in 0.1 M Tris containing 0.1 M NaCl. In control studies the cleavage of enkephalins was completely inhibited by one of the specific inhibitors of the enzyme: either by 10^{-4}M of the non-peptide SQ 20881⁶ or by 10^{-6}M of SQ 14225 (2-D-methyl-3-mercapto-L-proline) ¹⁰.

Both Leu-enk and Met-enk were cleaved by peptidyl dipeptidase at the Gly³-Phe⁴ bond to Tyr¹-Gly²-Gly³ and to Phe⁴-Leu⁵ or Phe⁴-Met⁵. Figure 1 shows the separation of the peptides in t.l.c. in two different systems. Fig. 1

The rate of hydrolysis of Met-enk was measured with an amino acid analyzer (Fig. 2). The enzyme and the substrate were incubated in a buffer more favor- Fig. 2

able for enzyme action than the one used for t.l.c. Instead of the NH_4HCO_3 buffer, the incubation medium contained 0.1 M NaCl in Tris buffer, since the enzyme cleaves most substrates faster in presence of Cl^- ⁵. Peptidyl dipeptidase cleaved Met-enk at a rate of 4.2 $\mu\text{mol}/\text{min}$ per mg protein and not at all in presence of SQ 20881. The same enzyme preparation cleaved the specific substrate Bz-Gly-Gly-Gly (10^{-3}M) at a rate of 7.4 $\mu\text{mol}/\text{min}$ when determined in the u.v. spectrophotometer ⁵.

Because cultured endothelial cells contain both aminopeptidase- and peptidyl dipeptidase-type enzymes we utilized t.l.c. to determine whether the N or C-terminal end of enkephalin is hydrolyzed first by enzymes of these cells.

Figure 3a shows the separation of products of hydrolysis of Leu-enk by endothelial cells. The $\text{Tyr}^1\text{-Gly}^2$ bond was cleaved first by the endothelial aminopeptidase. The enzyme was inhibited by the sequestering agent o-phenanthroline (10^{-3}M). In control studies a highly purified commercial leucine aminopeptidase (E.C. 3.4.11.1) was used (Fig. 3b) which also cleaved the pentapeptide to Tyr^1 and to tetrapeptide. We could detect the release of Phe-Leu by the action of endothelial peptidyl dipeptidase only after an incubation of 60 min or longer.

Fig. 3.

These experiments have shown that purified peptidyl dipeptidase cleaved enkephalins to tri- and dipeptide. The unprotected tripeptide liberated may be resistant to further degradation by the enzyme because of the free $\alpha\text{-NH}_2$ group.

Peptidyl dipeptidase is concentrated in various parts of the body, for example, in vascular endothelium of the lung and other organs, in kidney, in testicles ⁶, and pituitary gland ⁷. In the central nervous system it is present in structures such as choroid plexus¹¹, caudate nucleus ⁸ or striatum ⁷.

The rate of cleavage of Met-enk by peptidyl dipeptidase is quite substantial.

Under our conditions the rate was more than half of that of hydrolysis of Bz-Gly-Gly-Gly, the specific substrate of this enzyme,^{5,6} although the concentration of enkephalin was 1/5 of that of the optically active tripeptide. Since only peptides with free C-terminal end are cleaved by peptidyl dipeptidase⁶ resistance to the action of the enzyme may be one of the reasons for the markedly enhanced in vivo activity of the methionine amide analogue of enkephalin¹² over Met-enk. In tissues where peptidyl dipeptidase is concentrated, enkephalins may be readily degraded by this enzyme. Furthermore, by reacting with the enzyme enkephalins may interfere with the metabolism of other active peptides. Hypothetically they may inhibit the release of angiotensin II from angiotensin I or the inactivation of kinins by competing for the same enzyme.

Cultured human endothelial cells, however, inactivate enkephalins primarily by the action of an aminopeptidase, as demonstrated by the release of Tyr¹. The degradation of enkephalins by a similar aminopeptidase has been shown by using purified enzyme,¹³ plasma, brain homogenates,^{12,14} or a membrane fraction of rat brain² as sources of enzyme.

Our experiments indicate that rapid inactivation of enkephalins may be due to at least two different enzymes present in tissues. These enzymes presumably are localized in different structures of the cells. A peptidyl dipeptidase which is a component of plasma membrane of various cells⁶ may degrade enkephalins by liberating a C-terminal dipeptide and an aminopeptidase in endothelial cells may cleave peptides that they are taken up as blood flows past the endothelial surface at the N-terminal end,

Legends

Figure 1. Hydrolysis of Leu⁵- and Met⁵-enkephalin by purified hog kidney peptidyl dipeptidase (angiotensin I converting enzyme or kininase II) to Tyr-Gly-Gly and Phe-Leu or Phe-Met. The specific inhibitor SQ 14225 (10^{-6} M) inhibited the reaction. Peptides were incubated with the enzyme up to 60 min in presence and absence of inhibitor. a. Silica gel t.l.c. plate. Solvent: sec-butanol: acetic acid: water (4:1:5). Separation time 3 h. b. Aluminum oxide t.l.c. plate. Solvent: chloroform: methanol: NH_4OH (20:20:9). Separation time 1.5 h. The plates were sprayed with ninhydrin.

Figure 2. Hydrolysis of Met⁵-enkephalin by purified peptidyl dipeptidase determined in an automatic amino acid analyzer in absence (0—0) and in presence (0-----0) of the specific inhibitor SQ 20881 (10^{-4} M).

Figure 3. Cleavage of Tyr¹ from Leu⁵-enkephalin by human vascular endothelial cells (a) and by purified leucine aminopeptidase (b). Inhibitor: o-phenanthroline (10^{-3} M). Silica gel plates, conditions as in Fig. 1.

References

- ¹Snyder, S. H. & Simantov, R., *J. Neurochem.* 28: 13-20 (1977).
- ²Meek, J. L., Yang, H. -Y. T. & Costa, E., *Neuropharmacol.* 16: 151-154 (1977).
- ³Johnson, A. R. & Erdős, E. G., *J. Clin. Invest.* 59: 684-695 (1977).
- ⁴Johnson, A. R. & Erdős, E. G., in Substance P Nobel Symposium 37 (eds. Euler, U. S. von & Pernow, B.) 253-260 (Raven Press, New York 1977).
- ⁵Yang, H. -Y. T., Erdős, E. G. & Levin, Y., *Biochim. Biophys. Acta* 214: 374-376 (1970).
- ⁶Erdős, E. G., *Fed. Proc.* 36: 1760-1768 (1977).
- ⁷Yang, H. -Y. T. & Neff, N. H., *J. Neurochem.* 19: 2443-2450 (1972).
- ⁸Poth, M. M., Heath, R. G. & Ward, M., *J. Neurochem.* 25: 83-85 (1975).
- ⁹Oshima, G., Gecse, A. & Erdős, E. G., *Biochim. Biophys. Acta* 350: 26-37 (1974).
- ¹⁰Ondetti, M. A., Rubin, B. & Cushman, D. W., *Science* 196: 441-444 (1977).
- ¹¹Igić, R., Robinson, C. J. G. & Erdős, E. G., in Symposium on the Central Actions of Angiotensin and Related Hormones (eds. Buckley, J. P. & Ferrario, C. M.) 23-27 (Pergamon Press, Inc. New York 1977).
- ¹²Marks, N., Grynbaum, A. & Neidle, A., *Biochem. Biophys. Res. Comm.* 74: 1552-1559 (1977).
- ¹³Hughes, J., *Brain Res.* 88: 295-308 (1975).
- ¹⁴Hambrook, J. M., Morgan, B. A., Rance, M. J. & Smith, C. F. C., *Nature* 262: 782-783 (1976).

Acknowledgements

We wish to thank Mrs. C. J. G. Robinson for her cooperation in this project. These studies have been supported in part by N.I.H., U.S.P.H.S. PS0-HL14187, 1-RO1-HL-18826, RO1-HL16320, and by the ONR N00014-75-0807.



