

AD-A048 249

CLEMSON UNIV S C DEPT OF MATHEMATICAL SCIENCES
APPROXIMATION OF A COMPLETELY MONOTONE FUNCTION. (U)
OCT 77 K ALAM, W WALKER
N89

F/G 12/1


UNCLASSIFIED


N00014-75-C-0451
NL

| OF |
ADI
A048 249



END
DATE
FILMED
78
DDC





**DEPARTMENT
OF
MATHEMATICAL
SCIENCES**

**CLEMSON UNIVERSITY
Clemson, South Carolina**



6 APPROXIMATION OF A COMPLETELY
MONOTONE FUNCTION.

10 BY
KHURSHEED/ALAM & WALTER/WALKER

15 NAPP 14-75-C-0451

12 11p. 14 N89, TR-270

11 N89
9 OCTOBER 1977
TECHNICAL REPORT #270

DDC
RECEIVED
JAN 11 1978
B

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

1472
407 183 LB

APPROXIMATION OF A COMPLETELY MONOTONE FUNCTION

Khursheed Alam* & Walter Walker

Clemson University

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DOC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	and/or SPECIAL
A		-

ABSTRACT

∇ A function f on $(0, \infty)$ is completely monotone if it possesses derivatives of all orders, and the successive derivatives alternate in sign. It is shown that for each x the value of $f(x)$ lies between any two consecutive partial sums of the expansion of $f(x)$ in Taylor series. The given result can be applied to various functions such as the hypergeometric and confluent hypergeometric functions, which are widely used in applied mathematics. Some statistical applications are also given. ↑

Key words and phrases: Hypergeometric; Confluent Hypergeometric; Exponential function; Gamma distribution.

AMS Classification: Primary 26A86

Secondary 62E15

* The author's work was supported by the Office of Naval Research under Contract N00014-75-C-⁰⁴⁵¹Task NR 042-271.

1. Introduction and main results. A function f on $(0, \infty)$ is said to be completely monotone if it possesses derivatives $f^{(n)}$ of all orders and

$$(-1)^n f^{(n)}(x) \geq 0, \quad x > 0.$$

Typical examples of a completely monotone function are e^{-x} and $(1+x)^{-m}$, where m is a positive number. A wide class of functions which arise in applied mathematics have the completely monotone property. Consider, for example, the confluent hypergeometric and the hypergeometric functions, given by

$$\begin{aligned} \phi(a, b; x) &= \sum_{r=0}^{\infty} \frac{(a)_r}{(b)_r} \cdot \frac{x^r}{r!} \\ \psi(a, b; c; x) &= \sum_{r=0}^{\infty} \frac{(a)_r (b)_r}{(c)_r} \cdot \frac{x^r}{r!} \end{aligned}$$

Where $(a)_r = a(a+1)\dots(a+r-1)$. By the integral representation formulas (see e.g. Abramowitz and Stegun [1], 13.2.1, 15.3.1) we have that

$$\frac{\Gamma(b-a)\Gamma(a)}{\Gamma(b)} \phi(a, b; x) = \int_0^1 e^{xt} t^{a-1} (1-t)^{b-a-1} dt \quad (1.1)$$

$$b > a > 0$$

$$\frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \psi(a, b; c; x) = \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-tx)^{-a} dt \quad (1.2)$$

$$c > b > 0$$

From (1.1) and (1.2) it is seen that the confluent hypergeometric function $\phi(a, b; -x)$ is completely monotone for $b > a > 0$ and the hypergeometric function $\psi(a, b; c; -x)$ is completely monotone for $c > b > 0, a > 0$. If $a = b$ then $\phi(a, b; -x) = e^{-x}$.

The completely monotone property is important in the theory of probability and statistical analysis. It is known that a function f on $(0, \infty)$ is the Laplace transform of a probability distribution if and only if it is completely monotone and $f(0) = 1$. From Feller ([2], XI.1.4, Criterion 1 and 2) we have that if f and g are completely monotone then the product fg is also monotone and that if f is completely monotone and g is a positive function with completely monotone derivative then $f(g)$ is completely monotone (in particular, e^{-g} is completely monotone).

Let

$$S_n(x) = \sum_{r=0}^n f^{(r)}(0) \frac{x^r}{r!}$$

denote the partial sum of the Taylor series expansion of $f(x)$. The following theorem shows that if f is completely monotone then for each x the value of $f(x)$ lies between any two consecutive partial sums.

Theorem 1.1. Let n be a positive integer. If f is completely monotone then for each $x > 0$

$$S_{2n-1}(x) \leq f(x) \leq S_{2n-2}(x). \quad (1.3)$$

Proof: Let

$$h(x) = f(x) - S_{2n-1}(x).$$

We have $h^{(r)}(0) = 0$ for $r = 0, 1, \dots, 2n-1$. Since $h^{(2n-1)}(0) = 0$ and $h^{(2n)}(x) = f^{(2n)}(x) \geq 0$ therefore $h^{(2n-1)}(x) \geq 0$. Repeating

the argument we have that $h^{(2n-2)}(x) \geq 0$. Successive repetition of the argument yields $h(x) \geq 0$, establishing the first inequality in (1.3). The second inequality in (1.3) is proved similarly. \square

The result of Theorem 1.1 for $f(x) = e^{-x}$ is known. The following theorem and its corollary extend that result. Theorem 1.3 below gives a monotonicity property of the tail of the exponential series. The proof of the theorem is omitted. Let

$$M_t(x) = e^{-x} - \sum_{r=0}^t (-x)^r / r!, \quad x \geq 0.$$

Theorem 1.1. If t is even (odd) then for all $x \geq 0$ and $0 \leq v \leq 1$

$$M_t(x) + (-1)^t \frac{x^{v+t}}{\Gamma(v+t+1)} \geq (<) 0. \quad (1.4)$$

Proof: Let $x \geq 0$, $0 \leq v \leq 1$ and let $L(v)$ denote the quantity on the left side of the inequality (1.4). Since $M_t(x) \leq (>) 0$ if t is even (odd) by Theorem 1.1, we have that $L(v) \geq (<) 0$ if t is even (odd) for $v = 0$ and 1 .

Let $Z = x^{v+t} / \Gamma(v+t+1)$. We have

$$\begin{aligned} \partial^2 \log z / \partial v^2 &= -\partial \psi(v+t+1) / \partial v \\ &< 0 \end{aligned}$$

where ψ denotes the digamma function. Hence Z is either decreasing or increasing or first increasing then decreasing as v varies from 0 to 1. Suppose that t is even. Then $L(v)$ is either decreasing or increasing or first increasing then decreasing as v varies from 0 to 1. Since $L(v) \geq 0$ for $v = 0$

and 1, it follows that $L(v) \geq 0$ for $0 \leq v \leq 1$. Similarly, $L(v) \leq 0$ for $0 \leq v < 1$ if t is odd.

Corollary 1.1. If $0 \leq v \leq 1$ then for all $x \geq 0$

$$|M_t(x)| \leq x^{v+t}/\Gamma(v+t+1).$$

Let

$$m_{t+v}(x) = |M_t(x)|/x^{t+v}.$$

Theorem 1.3. As x varies from 0 to ∞ , $m_{t+1}(x)$ increases and $m_{t+v}(x)$ first increases then decreases for $0 < v < 1$.

Application: Let

$$g_v(x) = \frac{x^{v-1} e^{-x}}{\Gamma(v)}, \quad v, x > 0$$

denote the density function of the gamma distribution with v degrees of freedom. The distribution function is given by

$$\begin{aligned} G_v(x) &= \int_{-\infty}^x g_v(y) dy \\ &= \frac{x^v e^{-x}}{\Gamma(v+1)} \phi(1, v+1; x) \\ &= \frac{x^v}{\Gamma(v+1)} \phi(v, v+1; -x). \end{aligned} \tag{1.5}$$

Let X_i be a random variable distributed according to a gamma distribution with v_i degrees of freedom, $i=1, \dots, K$. Let X_1, \dots, X_K be jointly independent, and let $X^* = \max(X_1, \dots, X_K)$. The random variable X^* arises in various statistical problems, such as life testing. The m th moment of X^* is given by

$$\begin{aligned} \mu_m &= E(X^*)^m \\ &= \sum_{i=1}^K \int_0^\infty x^m \prod_{j \neq i}^m G_{\nu_j}(x) g_{\nu_i}(x) dx \\ &= \sum_{i=1}^K \int_0^\infty x^m \prod_{j \neq i}^m \frac{x^{\nu_j}}{\Gamma(\nu_j+1)} \phi(\nu_j, \nu_j+1; -x) g_{\nu_i}(x) dx \end{aligned}$$

Using Theorem 1.1 we obtain bounds on the value of μ_m , given by

$$\begin{aligned} \sum_{i=1}^K \int_0^\infty x^m \prod_{j \neq i}^m \frac{x^{\nu_j}}{\Gamma(\nu_j+1)} S_{2n-1}(\nu_j, \nu_j+1; -x) g_{\nu_i}(x) dx \leq \mu_m \leq \\ \sum_{i=1}^K \int_0^\infty x^m \prod_{j \neq i}^m \frac{x^{\nu_j}}{\Gamma(\nu_j+1)} S_{2n-2}(\nu_j, \nu_j+1; -x) g_{\nu_i}(x) dx \end{aligned} \quad (1.6)$$

where n is any positive integer and

$$S_n(a, b; x) = \sum_{r=0}^n \frac{(a)_r}{(b)_r} \cdot \frac{x^r}{r!}$$

denotes the partial sum of the confluent hypergeometric series $\phi(a, b; x)$. The left side of the inequality (1.6) reduces to

$$\left(\prod_{j=1}^K \Gamma(\nu_j) \right)^{-1} \sum^* (-1)^{\sum_{j \neq i} s_j} \Gamma(m + \sum_{j=1}^K \nu_j + \sum_{j \neq i} s_j) \prod_{j \neq i}^m ((\nu_j + s_j) s_j!)^{-1}$$

where \sum^* denotes summation over all non-negative integer values of $s_j \leq 2n-1$, $j \neq i$. The right hand side of (1.6) is reduced similarly. The relation (1.6) is useful in the computation of μ_m .

For another application consider a non-central gamma distribution with ν degrees of freedom and non-centrality parameter δ , given by the density function

$$g(x) = e^{-\delta} \sum_{r=0}^{\infty} \frac{\delta^r}{r!} g_{\nu+r}(x).$$

The m th moment of the distribution is given by

$$\begin{aligned} \mu_m(\delta) &= e^{-\delta} \sum_{r=0}^{\infty} \frac{\delta^r}{r!} \int_0^{\infty} x^m g_{\nu+r}(x) dx \\ &= e^{-\delta} \sum_{r=0}^{\infty} \frac{\delta^r \Gamma(m+\nu+r)}{\Gamma(\nu+r) r!} \\ &= \frac{\Gamma(m+\nu)}{\Gamma(\nu)} e^{-\delta} \phi(m+\nu, \nu; \delta) \\ &= \frac{\Gamma(m+\nu)}{\Gamma(\nu)} \phi(-m, \nu; -\delta) \end{aligned} \tag{1.7}$$

If $-\nu < m < 0$ the confluent hypergeometric function $\phi(-m, \nu; -\delta)$ is completely monotone in δ . Then the results of Theorem 1.1 can be used to derive bounds on the value of $\mu_m(\delta)$.

References

- [1] Abramowitz, M. and Stegun, I. A. (1970). Handbook of Mathematical Functions. Dover Publications.
- [2] Feller, W. (1965). An Introduction to Probability Theory and Its Applications, Vol. II. Wiley Publications.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER N89 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Approximation of a completely monotone function		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER TR #270 ✓
7. AUTHOR(s) Khursheed Alam Walter Walker		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0451 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS Clemson University ✓ Dept. of Mathematical Sciences Clemson, South Carolina 29631		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 042-271
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Code 436 Arlington, Va. 22217		12. REPORT DATE OCT. 1977
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Completely monotone function; Hypergeometric and Confluent hypergeometric functions; Gamma distribution.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A function f on $(0, \infty)$ is completely monotone if it possesses derivatives of all orders and the successive derivatives alternate in sign. The authors have shown that for each $x > 0$ the value of $f(x)$ lies between any two consecutive partial sums of the expansion of $f(x)$ in Taylor series.		

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

78

