AD-A048 486

VANDERBILT UNIV NASHVILLE TN DEPT OF GENERAL BIOLOGY F/G 6/13
IMMUNOCYTOCHEMICAL STUDIES OF 'SCHISTOSOMA MANSONI'.(U)
APR 77 B J BOGITSH

DADA17-73-C-3052
NL

UNCLASSIFIED

AD AO48486














END DATE FILMED 2- 78





IMMUNOCYTOCHEMICAL STUDIES ON SCHISTOSOMA MANSONI

FINAL SCIENTIFIC REPORT 1973-1976

BURTON J. BOGITSH

APRIL 7, 1977

SUPPORTED BY

US ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND WASHINGTON, DC 20314

CONTRACT NO. DADA 17-73-C-3052

DEPARTMENT OF BIOLOGY VANDERBILT UNIVERSITY NASHVILLE, TENNESSEE 37235

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

THE FINDINGS IN THIS REPORT ARE NOT TO BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION UNLESS SO DESIGNATED BY OTHER AUTHORIZED DOCUMENTS.

NO NO.

410520

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered) READ INSTRUCTIONS REPORT DOCUMENTATION PAGE BEFORE COMPLETING FORM 2. GOVT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER 1. REPORT NUMBER TYPE OF REPORT & PERIOD COVERED ITLE (and Subtitle) Final Report. Immunocytochemical Studies on Schistosoma 1973 - 1976 mansoni . 6. PERFORMING ORG. REPORT NUMBER 8. CONTRACT OR GRANT NUMBER(*) 7. AUTHOR(a) Burton J. Bogitsh DADA 17-73-C-3052 9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of General Biology 61102A Vanderbilt University 6 3A161102B71Q.00.007 Nashville, Tennessee 37235 12. REPORT DATE April 7, 19% US Army Medical Research and Development Command 13. NUMBER OF PAGES Washington, D. C. 20314 15. SECURITY CLASS. (of this report) 14. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office) Unclassified /2 15. DECLASSIFICATION DOWN GRADING 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 161102B71G 18. SUPPLEMENTARY NOTES 410 520 Hu 19. KEY WORDS (Continue on reverse side if necessary and identify by block number, Schistosoma mansoni; Miracidium; Cercaria; Trematode; Liver, mouse; Granuloma; Soluble Egg Antigen; Cercarial Antigen Protein; Ultrastructure; Freeze Dried Tissue; Antigen; Unlabeled Antibody; Schistosomules; Postacetabular gland; Preacetabular gland; Apical gland; Lateral penetration glands 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) AThe localization of soluble, cercarial antigen preparation (CAP) and soluble egg antigen (SFA) were localized in cercariae and miracidia, respectively at the light and electron microscope levels using the unlabeled antibody method In the cercaria, CAP was observed associated with the contents of the pre- and postacetabular glands and with the filamentous coat of mature cercariae. CAP components were also observed in the area surrounding the intrasporocyst cercar ae. In the eggshell enclosed miracidia, SFA was observed in the contents of the

The major objectives of the research were to localize soluble egg antigens (SEA) in eggshell-enclosed miracidium of <u>Schistosoma mansoni</u> and to evaluate several immunocytochemical techniques that will localize these antigens at the light and microscope levels of resolution. Of the procedures evaluated, the unlabeled antibody procedure proved to be the most sensitive and reproducible.

The localization of soluble egg antigen (SEA) in the eggshellenclosed miracidium of <u>Schistosoma mansoni</u> was performed at the light and
electron microscope levels using the unlabeled antibody method. Reaction
product was observed associated with the contents of the 3 major gland
cells described previously. Additionally, small vesicles were observed
that reacted positively for SEA. It was hypothesized that they may be
responsible for the secretion of the hatching fluid. SEA components were
also observed in the epidermis of the miracidium and in the area subjacent
to the eggshell.

The localization of the soluble, carcarial antigen preparation (CAP) in cercariae and schistosomules of Schistosoma mansoni was performed at the light and electron microscope levels using the unlabeled antibody method. Reaction produce was observed associated with the contents of the pre-and postacetabular glands and with the filamentous coat of mature cercariae. No reaction product was observed associated with the glycocalyx of schistosomules. However, several schistosomules did retain remnants of their filamentous coats and reaction product was observed associated with those remains. CAP components were also observed in the area surrounding the intrasporocyst cercariae.

PUBLICATIONS:

- Bogitsh, B.J. and C.E. Carter, 1975. Immunocytochemical studies of <u>Schistosoma</u>
 <u>mansoni</u>. I. Soluble egg antigen in eggshell-enclosed miracidium.

 J. Parasit. 61: 1031-1040.
- Bogitsh, B.J. and S.P. 1976. Immunocytochemical studies on <u>Schistosoma</u>
 <u>mansoni</u>. II. Soluble cercarial antigens in cercariae and schistosomule.
 J. Parasit. 62: 709-714.

These two reports were also presented at the annual meetings of the Southeastern Society of Parasitologists.

DISTRIBUTION LIST

4 copies

HQDA (SGRD-AJ) Washington DC 20314

12 copies

Defense Documentation Center (DDC) ATTN: DDC-TCA Cameron Station Alexandria, Virginia 22314

1 сору

Superintendent

Academy of Health Sciences, US Army

ATTN: AHS-COM

Fort Sam Houston, Texas 78234

1 сору

Dean School of Medicine Uniformed Services University of the Health Sciences Office of the Secretary of Defense 6917 Arlington Road Bethesda, Maryland 20014