

Document is available to the public through the National Technical Information Service, Springfield, Virginia 22161.

> Prepared for U.S. DEPARTMENT OF TRANSPORTATION FEDERAL AVIATION ADMINISTRATION Office of Aviation Medicine Washington, D.C. 20591

264 320



		1	Fechnical Report D	Documentation Page
1. Report No. 2	Government Acces	sion No. 3.	Recipient's Cotalog h	10.
FAA-A1- 8-5				
4. Title and Subrille		5.	Report Date	
		6.	Performing Organizati	ion Code
THREE REPORTS RELEVANT TO STR	ESS IN AVIAT	TONYPERSONNEL		
7. Author(s)			Performing Organizati	on Report No.
E. A. Higgins, M. T. Lategola	, and C. E.	Melton		
9. Performing Organization Name and Address		10	Work Unit No (TRAI	IS)
FAA Civil Aeromedical Institu	te	C C T	6	
P. O. Box 25062 Oklahowa City, Oklahowa 7313	5 5		. Contract of Grant No	».
		JPILL MA	Type of Report and F	Period Covered
12. Sponsoring Agency Name and Address	Citra	The Fills	$\lambda$	
Office of Aviation Medicine Redoral Aviation Administrati	i fil a	NAR 22 TEL	9	
800 Independence Avenue, S.W.		14	Sponsoring Agency (	lode
Washington, D.C. 20591	······			
15. Supplementary Notes		K.		
Work was performed under appr AM-A~76-PHY-86.	oved Tasks A	м-с-76-рну-82, Ам	-A-76-PHY-79,	and
16. Abstract SIn development of t	he aviation	stress protocol	simulation (A	SPS), the fol-
lowing conclusions were reach	ed: (1) In	experiments using	the ASPS, ca	rdiovascular
testing will be conducted in	parallel, bu	t separately; (2)	The time of	exposure to
altitude will be limited to 2 internal body temperature bl	h; and (3)	Measurements such	as visual ac	commodation,
will be included in the ASPS	only when ap	propriate.	conor rever,	and others
		• '		
Cardiovascular and pulmonary	parameters w	ere assessed unde	r simulated +	Gz and exer-
cise conditions in normal mai	es alter exp	osure to the ASPS	. Some param	eters were
ful physiological significance	e because of	the unavoidable	time lanse be	tween altitude
exposure and assessment. The	se prelimina	ry experiments se	rved to demon	strate that
meaningful physiological asse	ssments can	only be made duri	ng exposure t	o the alti-
tudes specified in the ASPS.		-		
Thirty-six controllor subject	· from provi	ous strong studio	e vore identi	fied who cub-
sequently suffered medical co	nditions sev	ere enough to req	uire waiver o	r retirement.
These subjects' stress indice	s were compa	red with those of	subjects who	had no known
pathology to see if any of th	e stress ind	icators were pred	ictive of pat	hological
conditions. The data showed	that control	lers who develope	d gastrointes	tinal pathol-
ogy had significantly ( $p < 0$ .	01) higher c	(st) <sup>than did their</sup>	normal count	erparts. At
Miemi ARTCC, che)was signific	antly elevat	ed (p < 0.05) in	the cardiovas	cular group.
17. Key Words Aviation Stress Pr	otocol	18. Distribution Statement		
imulation, Aeromodical Cortif	ication,			
eromedical Standards, Cardiov	ascular ving	Document is avail	able to the p	ublic through
inction, rhysiological Monitor	and Air	ne National lech	nicai informa inia 22161	tion Service,
raffic Controller.	, and All	bhrußtrein <sup>,</sup> Alf8	mia 22101	
19. Security Classif. (of this report)	20. Security Clus	sif, (of this page)	21. No. of Pages	22. Price
vcíassi fied	Unclassif	ied	34	
	L			L

1.1

. .

DEVELOPMENT OF THE AVIATION STRESS PROTOCOL--SINULATION AND PERFORMANCE, PHYSIOLOGICAL, AND BIOCHEMICAL MONITORING SYSTEMS: PHASE I

#### I. Introduction.

The Federal Air Surgeon, in his FY-75 Call for Estimates, said, "I have determined that the highest priority will be research in support of medical certification and medical standards development." In response to this priority statement, the Aviation Physiology Laboratory began to design a broad research program to provide clinical aviation medicine with information needed for medical certification decisions and for developing and revising medical standards.

The Aviation Stress Protocol--Simulation (ASPS) focuses on the fact that any assessment of fitness in pilots and air traffic controllers must take into account their potential for impaired performance. Not only must they perform their routine tasks at optimal levels for prolonged periods of time, but these persons must also retain the capacity to respond to task overloads in emergency situations. For pilots, there must be a normal tolerance to flight environmental stressors, particularly to the range of altitudes where general aviation pilots can fly without their requiring supplemental oxygen.

Because performance was an important consideration, performance measurements were made using the Civil Aeromedical Institute (CAMI) Multiple Task Performance Battery (MTPB) (1).

#### II. Experiment Design.

Healthy, male, paid volunteers (aged 21 to 28 years) served as test subjects. Interviews and physical examinations were conducted for all prospective subjects prior to selection. After selection, subjects were trained on the NTPB equipment and the vision testing equipment. Subjects were then tested individually in the CAMI research altitude chamber at each of four simulated altitudes. These (MSL) altitudes to which the subjects were exposed in random sequence were ground level, 8,000 ft, 10,500 ft, and 12,500 ft. Subjects reported to the laboratory without eating breakfast and were given a standard meal before beginning the experiment. After breakfast, subjects completed a sleep survey and a subjective rating scale

(Figure 1). Electrodes were attached to the subjects for heart rate recordings, and subjects were given a thermistor rectal probe for self-insertion for rectal temperature measurements. The subjects then reported to the altitude chamber for a 3-h test session. Heart rate was recorded continuously, rectal temperature and blood pressure were measured once during each 15-min period, and tests of visual accommodation (provided by Dr. K. W. Welsh of the Aviation Physiology Laboratory) were administered before the ses ion and at the end of each hour. Capillary blood samples from finger punctures were drawn before and after each test session for determinations of blood glucose and hematocrit. At the end of the 3-h test, subjects filled out a symptom checklist (Figure 2) and again executed a subjective rating scale (Figure 3). At the end of the 3-h test, urine was collected for measurements of epinephrine, norepinephrine, and 17-ketogenic steroids (2). After each chamber session, subjects were given a cardiovascular evaluation.

During each 3-h test session, the subjects performed four different combinations of tasks of the MTPB in 1-h cycles. (For a complete description of the tasks, see Jennings, Chiles and West(1).) Lights and meters monitoring tasks were performed throughout the test session. During the first 15 min of the hour, tracking arithmetic tasks were performed. During the second 15 min, arithmetic and problem solving were performed. During the third 15 min, pattern discrimination and problem were performed. And, during the final 15 min, pattern discrimination and tracking were performed. This sequence was then repeated during the second and third hours of the test session. The measure of performance used was an equal-variance composite of the various time and accuracy measures from the individual tasks.

Twelve subjects were selected and 10 completed the test series. Data collection was accomplished in three periods: March 15 to April 16, June 1 to June 24, and June 28 to August 2, 1976.

III, Results.

The sleep surveys indicated no difference in the quality or quantity of the subjects' sleep prior to the test days. According to the results of the subjective rating scales, attentiveness decreased significantly from the beginning to



5

Ħ

1





the second of a second

**此时这些自己的是一些不可** 

**建立的**。 一



Siteptons		DEGRE	st*		WHEN OCCURRED	
Difficult Concentration Difficult Coordination Drovsiness Arm/Hand Fatigue Headache throbbing-forchead temples ache -forchead temples sharp pains-forchead temples		of he point		e back of head back of head back of head		
Visual Symptoms blurring double vision eye fatigue or aching						
mulfling of sounds buzzing sound	0 0	0				
happiness depression irritability recurring thoughts detachment feeling dizziness			000000			

\* Mild - not of sufficient degree to impair performance Moderate - of sufficient degree to possibly impair performance Severe - of sufficient degree to <u>definitely</u> impair performance

Observer's Comments:

FIGURE 2. Symptom checklist.

\_\_\_\_\_Study \_\_\_\_ Think of how you felt near the end of the task period you just completed. Rate your feelings, attitudes, and emotions by circling the number which best describes them. It is important that you try to rate how you felt while still performing the task, and not now you may feel right now. \_ Condition \_ Period \_ Subject .o. Kane –

•

٠

; I

1

- 50

canada a contra desarrol de contra en estas anas

. 1.

	o	Very Inattentive	σ	Very Tired and Sleepy	5	Completely Relaxed	æ	Extremely Interested	ð	Indifferent
		,	ŝ		8		2		8	
		Inattentive	٢	Nore lired Than Usual	7	Relaxed	7	Moderately Interested	L	N£IdIy Annoyed
•	¢		¢.		4		9		4	
	ŝ	Åttentive	5	<pre>.bout My Usual Level .of thergy</pre>	2	lndifferent	5	Indifferent	~ ~	Sioderately Ánnoyed
•	4		-1		Ŀ		7		7	
	~	Quite Attentive	ſ	Xure Pep Than Usual	3	strathed	6	Moderately Lored	) 2013	Yuite Irritated
	5		-1				2		2	
	-	Very Áttent <b>iv</b> e		very Wide Avake and Lnergetic	1 V V	Strained	~	Extremely Bored	] : v r + + + + + + + + + + + + + + + + + +	Irritated

FIGURE 3. Subjective rating scale B.

- ---

5

•--

the end of the 3-h experiment at both 8,000 ft and 12,500 ft (Figure 4); a corresponding decrease in energy level was reported (Figure 5). Interest decreased with time of exposure at all altitudes, with statistical significance at the 0.05 evel for ground level and 8,000 ft and at the 0.01 level for 10,500 and 12,500 ft (Figure 6).

Concentration difficulty and drowsiness were the most frequently reported symptoms on the checklist; more subjects reported these two symptoms for 12,500 ft than for the other altitudes.

The visual accommodation tests yielded no statistically significant findings attributable to either altitude or duration of the test.

Hematocrit demonstrated a slight but statistically insignificant hemoconcentration through time at each of the four altitudes. The blood glucose was significantly less at the end of the experiments than at the beginning, but there were no differences attributable to altitude (Figure 7). There was a statistically significant increase in heart rate with increasing altitude (Figure 8). There were no statistically significant differences for systolic blood pressure, but diastolic pressure did demonstrate a statistically significant decrease ( $p \le .05$ ) at 12,500 ft (Figure 9).

The findings of the urine analyses are presented in Table 1.

Although the catecholamines, epinephrine, and norepinephrine demonstrated increasing values with each increase in altitude, the individual variances were large and the differences did not prove to be statistically significant.

The performance data from the MTPB exhibited great variability with no effects that could be attributed to altitude. In addition, the performance levels, when averaged over altitude conditions, showed a significant drop as a function of the number of test sessions experienced. Apparently, the subjects were losing interest in, or losing motivation to comply with the demands of the experiment.

IV. Discussion.

The series of experiments conducted during the development of the ASPS provided information indicating a need for



1

-



ない事業を使用する





مارغات الاغليات وللمأت ومقاولتهم أت

1

8

• -- •



ę

•

η.

•

.

hallen i

A COLORED F

FICURE 6.

والمحالية والمستحدث وأسأره الألهمية فتعاومهم مدامعا والألفان أترقنا فلسراء المستحد والمسارعة المسارعة المستعملين

9

**、**. .

•



۲. ۲.

.

÷,

,





Mean heart rate values through time as a function of altitude (N = 10). FIGURE 8.

ALC DE LA BA

日本市の記書



ų

「「「「「「「」」」」」



Altitude	Epinephrine	Norepinephrine	17-Ketogenic Steroids
(MSL)	(ng/h)	(ng/h)	(mg/h)
Ground	575,2	1725.1	0.6959
Level	±166.0	±386.3	±0.1361
8,000	651.3	1864.0	0.6014
Feet	±174.1	±526.5	±0.1446
10,500	745.5	2100.6	0.6695
Feet	±338.6	.,42.4	±0.1322
12,500	797.1	2213,6	0.6918
Feet	±386.1	,42.9	±0.2590

## TABLE 1. Mean Values\* and Standard Deviations\* for Urinary Hormones

\* N = 10

.

È.

-----

í

refinement of the protocol. Results of some of the subjective rating scales showed that increasing altitude accentuated changes that occurred primarily as a result of the long experimental procedure; i.e., interest decreased more at the two higher altitudes. Further, concentration difficulty and drowsiness were reported most frequently at 12,500 ft.

In mid-August 1976, conferences were held by Drs. Busby, Melton, Collins, Chiles, Lategola, and Higgins to evaluate the findings to that time. The general conclusions were:

A. The experiments were too long and too complicated.

B. The experiment protocol itself probably had greater effect on the subjects' responses than did the differences in altitude.

C. The data obtained from the MTPB and cardiovascular testing were not conclusive.

D. Further refinement of the protocol is indicated. In future experiments, the cardiovascular testing should be conducted separately, in parallel with other testing, and not at the end of an already long test procedure.

E. The time spent in the altitude chamber for a test session should be reduced to 2 h to preclude the boredom and fatigue associated with the longer testing procedure.

F. Additional tests should be conducted to determine the sensitivity of the MTPB for detecting differences due to altitude alone.

Between August 28 and November 18, 1976, shorter tests (2 h) were conducted at three altitudes; ground level, 12,500 ft, and 14,000 ft. Ten subjects were evaluated. The cardiovascular testing, visual accommodation tests, dietary control, and blood sampling were eliminated from the protocol. This series of tests was conducted primarily to evaluate the effects of altitude on complex performance. Unfortunately, subject motivation again appeared to fluctuate widely and again, altitude did not have a significant effect on performance. In the past, the problem was produced by rather frequent MTPB equipment malfunctions.

## V. Conclusion.

In future experiments using the ASPS, the cardiovascular testing will be conducted separately. The time of exposure to altitude and MTPB testing will be limited to 2 h. Measurements such as visual accommodation, internal body temperature, blood glucose, blood drug or alcohol level, and others will be included in the ASPS experiments only when appropriate. 

### REFERENCES

- Jennings, A. E., W. D. Chiles, and G. West: Methodology in the Measurement of Complex Human Performance: Two-Dimensional Compensatory Tracking, FAA Office of Aviation Medicine Report No. AM-72-21, 1972.
- Melton, C. E., J. M. McKenzie, B. D. Polis, S. M. Hoffmann, and J. T. Saldivar, Jr.: Physiological Responses in Air Traffic Control Personnel: Houston Intercontinental Tower, FAA Office of Aviation Medicine Report No. AM-73-21, 1973.

### ASSESSMENT OF CARDIOVASCULAR FUNCTION AFTER EXPOSURE TO THE AVIATION STRESS PROTOCOL--SIMULATION

#### I. Introduction.

The Aviation Stress Protocol--Simulation (ASPS) was designed to assess the potential for impaired performance by testing the capacity of pilots and air traffic controllers to: (i) maintain an optimum performance level for a prolonged period of time and (ii) respond to a task overload or emergency situation. For pilots, this capability includes the assessment of tolerance to flight environmental stresses in the range of altitudes to which general aviation pilots can legally fly without supplemental oxygen. The core of the ASPS (for pilots) is the exposure of the individual for single-day 3-h periods to a battery of simulated complex psychomotor tests (MTPB) at ground level (GL) and chamber altitudes at 8,000 (A), 10,500 (B), and 12,500 (C) feet (MSL). During the ASPS, task performance is monitored for deterioration of psychomotor performance and physiological and biochemical tests are administered to identify possible causes cf such deterioration.

As an adjunct procedure, cardiovascular tests were administered immediately following the ASPS to identify any functional deterioration due to the ASPS exposure <u>per se</u>. This report will deal only with the post-ASPS cardiovascular **testing**.

### II. Methods.

The cardiovascular test battery consisted of: (i) a test of orthostatic tolerance using a lower body negative-pressure (LBNP) box (supine) and (ii) a submaximum physical workload test using standardized, upright, seated, bicycle ergometry. The LBNP load and time sequences were a 3-min rest period, a 2-min exposure to -25 mm Hg LBNP ( $\approx$  +1.5 Gz), a 2-min recovery/rest period, and a 2-min exposure to -50 mm Hg LBNP ( $\approx$  +2.0 Gz). The ergometry load and time sequences were a 5-min rest period, a 3-min exposure to a 50 rpm/30 W workload, and a 3-min exposure to a 50 rpm/60 W workload. Parameters assessed during the LBNP test were pulmonary ventilation (using direct spirometry), heart rate (HR) and electrocardiogram (single-lead ECG), blood pressure (automatic sphygmomanometry), and temporal artery flow velocity (directional

Doppler equipment). During ergometry testing, the same parameters were assessed excepting the omission of the temporal artery flow velocity and the addition of oxygen uptake (analysis of quantitatively collected expired air).

Immediately following a given ASPS altitude exposure, the subject (seated in a wheelchair at all times) was "finger pricked" for microsampling of blood, wheeled to a private room for quantitative collection of urine, and then delivered to the cardiovascular testing room. The average elapsed time between the end of altitude exposure and start of the LBNP and ergometry tests were 44 and 63 min respectively.

A complete set of ASPS altitude exposures was achieved on a total of 10 normal male subjects (21-25 yr of age). The GL data served as baselines for comparing displacements due to altitudes A, B, and C.

#### III. Results.

<u>LBNP</u>. Both levels of +Gz exposure were well tolerated by all subjects without syncope or temporal artery flow reversal. Although syncope was absent, physiological displacements did occur in all subjects. In general, the displacements increased in rough proportion to the applied level of +Gz and the preceding altitude exposure. The quantitative altitude-related displacements of the systolic blood pressure (SBP), diastolic blood pressure (DBP), pulse pressure (PP), HR, temporal artery flow velocity (TAFV), and pulmonary ventilation ( $\Psi_{\rm p}$ ) are summarized in Tables 1-3. The SBP dropped during each +Gz exposure, but the drops and overall levels appeared to be approximately the same for GL as for preceding altitudes A, B, and C as compared to GL. The PP generally dropped during each +Gz exposure, and the overall levels were also lower following altitudes A, B, and C as compared to GL. The HR increased during each +Gz exposure, and the overall levels were also higher following altitudes A, B, and C as compared to GL. In each of the two +Gz exposures and the two preceding rest periods, most of the altitude-related increases in HR (A, B, and C compared to GL) were statistically significant ( $p \le 0.05$ ). In the altitude context of comparison, four of the DBP and two of the PP displacements were also significant.

The TAFV dropped during each +Gz exposure, but the overall levels were substantially higher following altitudes A, B, and

SBP and DBP Responses to +1.5 and +2.0 Gz at Ground Level, and at 8,000, 10,500, and 12,500 Ft Altitudes TABLE 1.

•

\*

i

					LB L	NP		1	
			SBP (	(mm Hg)			DBP (	(pH mm)	
		e:	+15 02	œ	+2002	œ	+15 62	æ	+20 Gz
Ground	Ε	1.711	115.7	118.1	8.111	65.1	65.5	68.1	64.5
Level (Control)	S	ä.	3.0	3°8	3.5	1.7	1,9	2.0	2.1
	ε	100.6	98.6	8-66	99.1	1.701	108.6	103.9	108.1
(% of Control)	SE	1.5	1.7	4 5	5°†	2.6	2.1	3.2	۵.4
10 500 ti	Ε	102.4	100.1	102.1	104.8	106.3	105.4	102.4	109.4
(% of Control)	SE	1.9	2.2	2.7	2.8	3.7	2.4	3.1	2.3
5 EAO fi	ε	6-66	100.2	100.6	100.5	104.1	102.9	104.5	108.4
(% of Control)	SE	1.3	3.4	2.8	3.3	2.2	3.0	4°0	9°7

T T

PP and HR Responses to +1.5 and +2.0 Gz at Ground Level, and at 8,000, 10,500, and 12,500 Ft Altitudes TABLE 2.

. . . .

•

۰ ۲

i

					LB L	NP		1	
			PP (n	nm Hg)			HR (I	(mdo	
		éx	+ 1.5 Gz	œ	+2062	æ	+15 02	œ	+20 Gz
Ground	Ε	52.1	50.2	50.0	47.3	55.8	56.7	58.2	65.0
Level (Control)	SE	2.4	3.1	3.£	5°7	2.3	2.2	2.8	2.6
8000 ft	E	92.7	• • • • • • • • • • • • • • • • • • •	98.2	95.7	104.8	108.3	105.1	105.8
(% of Control)	SE	2.5	3.3	5.6	4 <b>~</b> 8	1.9	3.2	3.6	0°†
10,500 ft	E	95.5	4.42	102.2	101-9	108.4	• 112.2	• 0.70I	• •
(% of Control)	SE	3.2	3.9	3.3	7.0	2.1	3.1	3.0	£"#
12 500 ft	٤	95.0	95.2	96.6	93.1	108.4	107.4	106.6	104.3
(% of Control)	SE	3.0	6 <b>.</b> 5	÷.5	6.9	3.4	3.4	F	e. 4
					•	•			

4

 $\tilde{r}_{E}$  and TAFV Responses to +1.5 and +2.0 Gz at Ground Level, and at 8,000, 10,500, and 12,500 Pt Altítudes TABLE 3.

- 50 B. J.

					LB	NP		ł	
			ўЕ (	lpm)			TAFV	(cm/s)	
		~	+15 Gz	ł,	+2061	æ	+15 02	R	+2.0 Gz
Ground	ε	7.6	7.2	g•1	8.8	9.7	8.9	8,1	1.5
Level (Centrol)	SE	0.5	0.6	<b>4</b> ~0	0.8	1.6	2.1	1.3	1.5
80004	E	101.6	8.211	105.9	0.62	108.4	132.7	143.8	\$°67[
(% of Control)	SE	6-5	12.0	10.6	3°5	14.6	6.12	28.5	31.2
10 500 ft	E	103.3	98.9	93.5	101.7	139.6	225.2	233.2	318.8
(% of Control)	SE	9.é	1.6	7.1	8.9	28.5	39.6	103.1	170.1
12 500 #	٤	t.62	1.121	107.2	94 <b>.</b> 1	137.8	188.0	223.6	258,8
(% of Control)	SE	8.0	27.0	6.7	9.1	19.2	59.3	61.3	3*11

C compared to GL. None of the observed blood flow velocity displacements was statistically significant. The  $V_{\rm p}$  response

to each +Gz exposure varied widely in direction and magnitude with no obvious consistent relationship to either +Gz or altitude.

<u>Ergometry</u>. Both the 30- and 60-W workloads were comfortably tolerated by all subjects. The SBP, DBP, PP, HR,  $\Psi_{\rm E}$ , and oxygen uptake  $(\Psi_{0})$  increased generally in direct

proportion to the applied ergometric load. With the exception of PP, the overall levels of these parameters were higher following altitudes A, B, and C as compared to GL. The overall levels of PP were lower following altitudes A, B, and C as compared to GL. The quantitative altitude-related displacements of the SBP, DBP, PP, HR,  $\nabla_{\rm E}$ , and  $\nabla_{\rm Q}$  are summarized in Tables

4-6. Statistically significant (P $\leq$  0.05) altitude-related changes occurred: (i) in DEP, PP, HR, and  $\nabla_0$  in the

preergometric rest period; (ii) in DBP, PP, and HR during the 30-W workload; and (iii) in HR and  $\Psi_{\rm p}$  during the 60-W workload.

#### IV. Discussion.

Some statistically significant altitude-related displacements in physiological parameters appeared in both the LBNP and ergometry segments of the post-ASPS cardiovascular assessment, even though these two testing segments nonoptionally followed the end of altitude exposure by an average of 44 and 63 min respectively. Because these displacements could possibly be accentuated by closer temporal proximity of the cardiovascular testing to the altitude exposures, it was decided that the cardiovascular testing would henceforth be run in parallel racher than in tandem with the core ASPS. The cardiovascular tests are to be applied at GL and during appropriate altitude exposures rather than after the altitude exposures. During the currently extended "shakedown" Phase I cesting of the core ASPS, the paralleling cardiovascular equipment and protocol will be shifted to one of the small altitude chambers. We should be ready to resume full-scale parallel testing when the core ASPS commences its Phase II operations. Running this test in parallel with the core ASPS tests further has the boneficial effect of relieving the tedium of the core tests by reducing the length of subject exposure.

S3P and D3P Responses to 30% and 60% Bicycle Ergcmetry Workloads at Ground Level, and at 8,000, 10,500, and 12,500 Ft Altitudes TABLE 4.

Ŋ

ų

٠,

• • •

Š.,

- 5.5

				ERGON	<b>IETRY</b>		
			SBP (mm Hg)			DBP (mmHg	(
		æ	30 W	<b>₽ 0 ₽</b>	¢	30 W	W 09
Ground	ε	116.4	128.6	138.6	6 <b>.</b> 43	<b>1.</b> 99	6,8
(Cantrol)	×	3.0	3.5	3.6	2.2	2.5	3.2
8.000 ft	E	100.5	99.2	102.4	1.905.7	• •	104.6
(% of Control)	SE	1.9	0.8	1.8	2.6	2.1	3.5
10.500 ft	ε	100.5	100.2	100.4	106.2	• • • • • •	106.0
(% of Control)	SE	1.3	1.9	1.8	2.3	3.2	2.7
12.500 ft	E	301.5	100.0	302.4	<b>105.</b> 6	106.0	106.9
(% of Curital)	8	2.7	2.7	2.1	5.9	5.2	5.8
	1	•					

7

ę

والمستعملية والشراطية ومعرفها والمستعمل والمستعمل والمستعمل والمتعالية والمستعمل والمستعم

à

PP and HR Responses to 30M and 60M Bicycle Ergometry Workloads at Ground Level, and at 8,000, 10,500, and 12,500 Ft Altitudes TABLE 5.

Î

				ERGON	ЕТНҮ		
			PP (mmHg			HR (bpm)	
		æ	N OE	60 W	¢	30 W	60 W
Ground	Ε	51.5	62.5	<b>6</b> 8.3	50.8	6-11	5*.2
Level (Control)	ß	3.2	3.7	3.9	2.1	3.1	4.5
8.000 ft	E	88.4	91.8	103.0	105.6	102.6	102.6
(% of Control)	SE	5.7	1.4	4.5	3-3	2.7	2.2
10 500 #	ε	89.5	£.09	6,59	107.7	105.1	104.2
(X of Control)	SE		۳°0	3.8	2.5	3.0	5.1
12-500 ft	ε	87.7	95.8	100.3	10.5	108.5	105.0
(% of Control	SE SE	50 1	3.8	ي. ال	3.0	3.3	3.1
	-			-	-	-	_

 $\tilde{v}_{E}$  and  $\tilde{v}_{0}^{}$  Responses to 30W and 60W Bicycle Ergometry Workloads at Ground Level, and at 8,000, 10,500, and 12,500 Ft Altitudes TABLE 6.

...

and the state of the second

ŝ,

•

				ERGON	IETRY		
			V <sub>E</sub> (Ipm)		.>	2 (ml/m/1	(g)
		~ ~	30 6	60 W	œ	30 W	60 W
	F						
Ground	Ε	8.6	16.1	22.6	3.2	8.5	12.7
Lever (Control)	Ж	0.4	0.7	0.8	0.2	٥.4	0.6
	T			•			
8 000 <del>1</del> 1	٤	104.3	97.2	109.1	105.3	100.3	103.3
(% of Control)	SE	6.4	4.5	2.9	3.2	4.5	2.7
	Ī						
	ε	104.2	94.3	103.8	117.9	8.66	101.7
(% of Control)	SE	9.2	3.1	2.2	6.9	5.0	2.8
	T			*.	•	201	g
12 500 11	E	107.4	201.8	106.8	110.7	0.001	0°001
(% of Control	3	3.9	6.0	2•3 5	5.9	5,2	<b>4</b> ,3
	-	-					

THE RELATIONSHIP BETWEEN STRESS-RELATED METABOLITES AND DISQUALIFYING PATHOLOGY IN AIR TRAFFIC CONTROL PERSONNEL

#### I. Introduction.

This laboratory's first study of stress in air traffic controllers was carried out in 1968 at O'Hare Tower. Since that time, 15 other studies have been carried out at towers, centers, TRACON's, and flight service stations (4-10). A total of 237 controllers representing trainees, journeymen, and supervisors have been studied; 189 of those are the subjects of this study--48 were either studied too recently for their data to be available or their urinary data were incomplete.

Sufficient time has elapsed since the first study at O'Hare that some controller subjects have suffered a variety of medical conditions severe enough to require either waiver or retirement. Notations regarding medical conditions requiring either of the above actions are in the controllers' files in the Aeromedical Certification Branch of the Civil Aeromedical Institute. These files were searched for controller subjects who developed medical conditions after they were studied. Thirty-six (19 percent of the total) such controllers were identified who developed pathology after the time of the stress study in which they served as subjects. These controllers showed pathology in one or more of three groups: gastrointestinal, neuropsychiatric, and cardiovascular (Table 1). A further breakdown of the three conditions by region and facility is shown in Table 2. 

#### II. Results.

An index developed in this laboratory (6) readily allows the comparison of excretion values for 17-ketogenic steroids, epinephrine, and norepinephrine. The average of the three individual indices ( $c_{st}$ ,  $c_e$ , and  $c_{ne}$ ) is the composite stress index (C<sub>s</sub>). The individual and composite indices for normal controllers (those without a pathology file) and those with pathological conditions are shown in Table 3. The table shows that for the whole population of controllers, those who developed gastrointestinal pathology had significantly ( $\underline{P} < 0.01$ ) higher c<sub>st</sub> than did their normal counterparts at the time they were studied. The index

### TABLE 1. Distribution of Diagnoses Among Three Major Disease Categories

•

-5

Disease		Percentage
Category	Diagnosis	Occurrence
Gastrointestinal		
	Gastric Ulcer	15.4
	Duodenal Ulcer	15.4
	Peptic Ulcer	15.4
	Gastritis	15.4
	Gall Bladder	7.7
	Other	30.7
Cardiovascular		
	Myocardia: Infarction	26.7
	Coronary Insufficiency	6.7
	Hypertension	33.3
	Angina	13.3
	Arteriosclerotic Heart Disease	13.3
	Arrhythmia	6.7
Neuropsychiatric		
	Anxiety Reaction	21.8
	Anxiety Depression	21.8
	"Neuroses"	8.7
	Cluster Headaches	4.3

の日本社会

4.3

4.3

8.7

4.3

21.8

2

Depression

Schizophrenia

Anxiety Neuroses

Personality Disorder

Psychosomatic Illness

Region	Facility	Gastro- intestinal	Cardio- vascular	Neuro- psychiatric
Southern				
	Opa Locka Tower	0	0	1
	Miami ARTCC	3	8	12
	Atlanta ARTCC	<u>0</u>	<u>1</u>	3
		3	9	16
Southwest				
	Houston Inter- continental Tower	1	1	0
	Fort Worth ARTCC	<u>1</u>	1	<u>3</u>
		2	2	3
Central				
	O'Hare Tower (1968)	4	1	2
Western				
	Los Angeles TRACON	0	0	0
	Oakland TRACON	<u>1</u>	<u>0</u>	0
		1	0	0

# TABLE 2. Distribution of the Three Major Disease Categories Among Regions and Facilities

			-	-			
	c st	<b>P</b> **	с е	<u></u> 2*	c <sub>ne</sub>	P*	
Normal	0.67	≤0.01	0.60		0.77		
Gastrointestinal	1.12	0.01	0.61	ns**	0,68	ns	
Cardiovascular	0.94	ns	0.74	ns	1.25	≤0.05	
Neuropsychiatric	0.64	ns	0.82	ns	0.85	ns	

## TABLE 3. Pathology and Grouped Stress Indices for the Entire Subject Population

\*p = Level of significance of difference between normal and pathological conditions \*\*ns = Not significant

Unpaired t-test

for norepinephrine (c<sub>ne</sub>) was also elevated significantly

 $(\underline{p} < 0.05)$  over the normals for the cardiovascular pathology group.

When the individual facilities are considered, c is

significantly ( $\underline{p} < 0.05$ ) elevated in the cardiovascular pathology group at Miami ARTCC (Table 4). There are no other points of significant difference.

### III. Discussion.

This study indicates that high levels of steroid excretion are related to the presence or later development of gastrointestinal disease, the most common "stress related" disease. As shown in Table 1, the principal subcategories of that disease relate to ulceration of the stomach and duodenum. Ulcers have long been known to be related to high levels of endogenous steroids (1,2,11).

At Miami Center, the occurrence of cardiovascular disease (25 percent hypertension, 75 percent coronary artery disease) is significantly related to high levels of norepinephrine excretion. Norepinephrine is the principal catecholamine liberated by the sympathetic nervous system and is, therefore, the main regulator of arteriolar resistance. Norepinephrine has also been shown to have pronounced effects in causing arrhythmias and fatty changes in the hearts of dogs (3).

This study suggests that excretion levels of 17-ketogenic steroids and norepinephrine may be predictive of heart disease, hypertension, and gastrointestinal ulceration. However, because of the small number of controllers in each diagnostic category, the statistics leave much to be desired; the findings reported here are, therefore, tentative, and this report should be viewed as preliminary. The pathology files will be periodically checked in the years to come to see if the statistical validity of these observations changes as more controllers join the list of those with pathology files.

ATC Facility	Disease Category	c <sub>st</sub>	<u>P</u> *	с <sub>е</sub>	<b>D</b> ,,	c <sub>ne</sub>	<b>Đ</b> <sup>%</sup>
O'Hare Tower							
	Normal	1 23		0 74		0 99	
	Gastrointestinal	1.73	nssk	0.71	ns	0.79	ns
	Cardiovascular	2.91	+	1.07	+ +	1.51	1. 1
	Neuropsychiatric	1.56	ns	0.71	ns	0.73	ns
Opa Locka Tower	r						
	Normal	0.67		0.74		1.21	
	Gastrointestinal	0.00	+	0.00	4	0.00	÷
	Cardiovascular	0.00	+	0.00	t.	0.00	't'
	Neuropsychiatric	0.17	†	0.63	†	0.42	t
Miami ARTCC							
	Normal	0.59		0.68		0.88	
	Gastrointestinal	0.73	ns	0.42	ns	0.78	ns
	Cardiovascular	0.74	ns	0.69	ns	1.43	0.05
	Neuropsychiatric	0.65	ns	0.84	ns	0.96	ns
Atlanta ARTCC							
	Normal	0.80		0.29		1.38	
	Gastrointestinal	0.00	+	0.00	†	0.00	†
	Cardiovascular	0.34	ተ	0.53	†	1.25	t
	Neuropsychiatric	0.59	ns	0.52	ns	1.31	ns
Fort Worth ART	сс						
	Normal	0.23		0.54		0.20	
	Gastrointestinal	0.09	Ŧ	0.88	+	0.36	÷
	Cardiovascular	0.60	+	1.04	+	0.16	†
	Neuropsychiatric	0.24	ns	1.21	ns	0.20	ns

# TABLE 4. Pathology and Individual Stress Indices for Various ATC Facilities

1.1

.

# TABLE 4 (Continued)

Facility	Category	c <sub>st_</sub>	£*	с <sub>е</sub>	<u>D</u> *	cne	<u>P</u> *
Houston							
Intercontine Tower	ntal						
	Normal	1.01		0.40		0.59	
	Gastrointestinal	1.77	t	0.96	+	0.57	4
	Cardiovascular	1.77	+	0,96	+	0,57	+
	Neuropsychiatric	0.00	ł	0.00	t	0.00	ł
Oakland TRAC	NC						
	Normal	0.37		0.85		0.48	
	Gastrointestinal	0.48	t	0.41	Ŧ	0.30	+
	Cardiovascular	0.00	÷	0.00	ł	0.00	
	Neuropsychiatric	0.00	†	0.00	+	0.00	÷
Los Angeles TRACON							
	Normal	0.63		0.38		0.79	
	Gastrointestinal	0.00	t	0.00	Ŧ	0.00	÷
	Cardiovascular	0,00	÷	0.00	+	0.00	
	Neuropsychiatric	0.00	+	0.00	ų. V	0.00	י לי

\*p = Level of significance of difference between normals and diseased

\*\*ns = Not significant

a se a constante de la constant La constante de la constante de

÷

1

+ = Number of cases insufficient for statistical treatment

7

- 1-+

#### REFERENCES

- Gray, S. J., J. A. Benson, H. M. Spiro, and R. W. Reifenstein: Effects of ACTH and Cortisone Upon Stomach, GASTROENTEROLOGY, 19:658-673, 1952.
- Kirsner, J. B., A. P. Klotz, and W. L. Palmer: Unfavorable Course of Gastric Ulcer During Administration of ACTH and Cortisone, GASTROENTEROLOGY, 20:27-29, 1952.

- Maling, H. M., and Benjamin Highman: Exaggerated Ventricular Arrhythmias and Myocardial Fatty Changes After Large Doses of Norepinephrine and Epinephrine in the Unanesthetized Dog, AM. J. PHYSIOL., 194:590-596, 1958.
- Melton, C. E., J. M. McKenzie, B. D. Polis, G. E. Funkhouser, and P. F. Iampietro: Physiological Responses in Air Traffic Control Personnel: O'Hare Tower, FAA Office of Aviation Medicine Report No. AM-71-2, 1971.
- Melton, C. E., J. M. McKenzie, R. C. Smith, B. D. Polis, E. A. Higgins, S. N. Hoffmann, G. E. Funkhouser, and J. T. Saldivar, Jr.: Physiological, Biochemical, and Psychological Responses in Air Traffic Control Personnel: Comparison of the 5-Day and 2-2-1 Shift Rotation Patterns, FAA Office of Aviation Medicine Report No. AM-73-22, 1973.
- Melton, C. E., J. M. McKenzie, J. T. Saldivar, Jr., and S. M. Hoffmann: Comparison of Opa Locka Tower With Other ATC Facilities by Means of a Biochemical Stress Index, FAA Office of Aviation Medicine Report No. AM-74-11, 1974.
- Melton, C. E., R. C. Smith, J. M. McKenzie, J. T. Saldivar, Jr., S. M. Hoffmann, and P. R. Fowler: Stress in Air Traffic Controllers: Comparison of Two Air Route Traffic Control Centers on Different Shift Rotation Patterns, FAA Office of Aviation Medicine Report No. AM-75-7, 1975.
- Melton, C. E., R. C. Smith, J. M. McKenzie, S. M. Hoffmann, and J. T. Saldivar, Jr.: Stress in Air Traffic Controllers: Effects of ARTS-III, AEROSP. MED., 47(9):925-930, 1976.
- 9. Smith, R. C.: The State-Trait Anxiety Inventory and the Assessment of Stress in Private Pilot Training. Proceedings

of the 80th Annual Convention of the American Psychological. Association, 7:64-65, 1972. ्रा त्यां स्वतिहेलकारी संक्रेफि र है में 2 होती. . विवस का ब्यावरी आधिक प्रैतिकारी संवति का ती कारत जन्म की की

يفيمنا العفاظيا بالمخرار

accurate avoid the risk of the final data

a contra Milling, Process Incollicities

 Smith, R. C., and C. E. Melton: Susceptibility to Anxiety and Shift Difficulty as Determinants of State Anxiety in Air Traffic Controllers, AEROSP. MED., 45:599-601, 1974.

and the second secon

ł

 Thorn, G. W., D. Jenkins, J. C. Laidlaw, F. C. Gcetz, J. F. Dingman, W. L. Arons, D. H. P. Streeten, and B. H. McCracken: Pharmacologic Aspects of Adrenocortical Steroids and ACTH in Man, N. ENGL. J. MED., 248:323-337, 1953.