



Annual technical rept. no. 1 AD A 051720 Annual Report of Research Sponsored by the Office of Naval Research (Power Program) 15 Contract: N00014-75-C-0470 Project NR 092-550 6 Chemical Kinetics of Excited States. Novel Application of Gas Phase EPR and Chemical Kinetics of Polyatomic Free Radicals. Principal Investigator an 18 David L. McFadden Department of Chemistry Boston College Chestnut Hill, MA 02167

This document has been approved for public release and sale; its distribution is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States Government.



493 740

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2. GOVT ACCESSION	NO. 3. RECIPIENT'S CATALOG NUMBER
Annual Technical Report I	
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED
CHEMICAL KINETICS OF EXITED STATES	Annual Technical Rep
NOVEL APPLICATION of ERR and	for 1977
CHEMICAL KINETICS OF POLYATOMIC FREE	6. PERFORMING ORG. REPORT NUMBER
RADICALS 7. AUTHOR(0)	8. CONTRACT OR GRANT NUMBER(.)
	/
David L. McFadden	, N00014-75-C-0470 New
PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Department of Chemistry	and be that that
Boston College	NR 092-550
Chestnut Hill, MA 02167	12 REPORT DATE
Office of Naval Research	January 25 1978
Power Program	13. NUMBER OF PAGES
Arlington VA 22217	17
14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office	e) 15. SECURITY CLASS. (of this report)
	Unclassified
	15. DECLASSIFICATION DOWN GRADING
	SCREDULE
Approyed for public release; distributi	on unlimited
Approyed for public release; distributi 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different	on unlimited
Approyed for public release; distributi 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES	on unlimited
Approyed for public release; distributi 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, If different 18. SUPPLEMENTARY NOTES	on unlimited
Approyed for public release; distributi 7. DISTRIBUTION STATEMENT (of the ebstrect entered in Block 20, if different 18. SUPPLEMENTARY NOTES	on unlimited
Approyed for public release; distributi 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block num	on unlimited
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block num Kinction Floring Coin Decement of the second statement of the second	ber)
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF, Radicals, Hydroca	ber) Spectrometry, Thons, Fluorocarbons
Approyed for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca	ber) Spectrometry, Trbons, Fluorocarbons
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca	on unlimited
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, If different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block number Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse side if necessary and identify by block number 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)	ber) Spectrometry, rbons, Fluorocarbons
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde 11 necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse elde 11 necessary and identify by block number Chemical kinetics research at Bost	ber) Spectrometry, Trbons, Fluorocarbons
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde 11 necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse elde 11 necessary and identify by block number Chemical kinetics research at Bost ceeding in two major areas. One area i	on unlimited trom Report) Spectrometry, trbons, Fluorocarbons er; con College is now pro- nvolves the direct study
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different 18. SUPPLEMENTARY NOTES 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse elde if necessary and identify by block number Chemical kinetics research at Bost ceeding in two major areas. One area i of the kinetics of gas phase chemical r	on unlimited trom Report) (Spectrometry, trbons, Fluorocarbons (no College is now pro- nvolves the direct study eactions of metastable
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse side II necessary and identify by block number Chemical kinetics research at Bost ceeding in two major areas. One area i of the kinetics of gas phase chemical r electronically excited atoms and molecu	en unlimited from Report) ber) Spectrometry, urbons, Fluorocarbons ber) con College is now pro- nvolves the direct study eactions of metastable les. In contract year
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse elde if necessary and identify by block number Chemical kinetics research at Bost ceeding in two major areas. One area i of the kinetics of gas phase chemical r electronically excited atoms and molecu 19. (Figure 1) denotes that the figure 1) of the successfully demonstrated that	An unlimited three Report) ber) Spectrometry, tbons, Fluorocarbons ber) con College is now pro- nvolves the direct study eactions of metastable les. In contract year Electron Paramagnetic
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the ebstrect entered in Block 20, if different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse elde if necessary and identify by block num Chemical kinetics research at Bost ceeding in two major areas. One area i of the kinetics of gas phase chemical r electronically excited atoms and molecu 1976 we successfully demonstrated that Resonance (EPR) detection of metastable	An unlimited them Report) ber) Spectrometry, trbons, Fluorocarbons ber) con College is now pro- nvolves the direct study reactions of metastable les. In contract year Electron Paramagnetic atoms in the gas phase
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessary and identify by block num Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse elde if necessary and identify by block numb Chemical kinetics research at Bost ceeding in two major areas. One area i of the kinetics of gas phase chemical r electronically excited atoms and molecu 1976 we successfully demonstrated that Resonance (EPR) detection of metastable can be combined with an electrical disc	An unlimited them Report) Spectrometry, trbons, Fluorocarbons ery con College is now pro- nvolves the direct study reactions of metastable les. In contract year Electron Paramagnetic atoms in the gas phase tharge-fast flow apparatus
Approved for public release; distributi 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different 18. SUPPLEMENTARY NOTES 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessary and identify by block must Kinetics, Electron Spin Resonance, Mass Metastable Atoms, CF <sub>3</sub> Radicals, Hydroca 20. ABSTRACT (Continue on reverse elde if necessary and identify by block must Chemical kinetics research at Bost ceeding in two major areas. One area i of the kinetics of gas phase chemical r electronically excited atoms and molecu 1976 we successfully demonstrated that Resonance (EPR) detection of metastable can be combined with an electrical disc D 1/AM 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE	An unlimited them Report) Spectrometry, trbons, Fluorocarbons (a) (a) (b) (c) (c) (c) (c) (c) (c) (c) (c

LECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)

to obtain absolute rate constants for reactions of metastable atoms. The major thrust of this work during contract year 1977 has been to utilize the experimental capability which was developed in the first two years of this program. We have measured absolute reaction rate constants for the deactivation of metastable nitrogen atoms  $(N^{D}_{3/2,5/2})$  by 17 different quenching partners, including  $O_2$ ,  $N_2O$ ,  $CO_2$ ,  $H_2$ ,  $CH_4$ ,  $C_2H_4$ ,  $C_2H_6$ ,  $C_3H_8$ ,  $C_4H_{10}$ ,  $C_5H_{12}$ ,  $CH_3CHCH_2$ ,  $C_2H_2$ ,  $CH_3F$ ,  $CF_4$ ,  $C_2F_6$ , and  $SF_6$ .

A second major project in our overall research effort was initiated during contract year 1977. An effort is being directed toward the development of new experimental techniques for the direct kinetics study of transient polyatomic free radical species such as  $CF_3$ . During contract year 1977 we have successfully detected  $CF_3$  radicals as products of an RF electrical discharge in mixtures of  $C_2F_6$  and He. Low energy electron bombardment ionization coupled with Time-of-Flight Mass Spectrometry was employed in the detection of these radicals.

1718		White Sect	- >
996		Bett Sectio	. 0
UNAHROUN	CED		D
JUSTIFICA	TION		
BY	9165 AN		
DV Distrino	7108/49.	AILADILITY	000 <b>ES</b>
ev ommanen Dist.	TION/AV	AILASILITY . and/or Si	CODES

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

#### Summary

Chemical kinetics research at Boston College is now proceeding in two major areas. One area involves the direct study of the kinetics of gas phase chemical reactions of metastable electronically excited atoms and molecules. In contract year 1976 we successfully demonstrated that Electron Paramagnetic Resonance (EPR) detection of metastable atoms in the gas phase can be combined with an electrical discharge-fast flow apparatus to obtain absolute rate constants for reactions of metastable atoms. The major thrust of this work during contract year 1977 has been to utilize the experimental capability which was developed in the first two years of this program. We have measured absolute reaction rate constants for the deactivation of metastable nitrogen atoms ( $N^2D_{3/2,5/2}$ ] by 17 different quenching partners, including O<sub>2</sub>, N<sub>2</sub>O, CO<sub>2</sub>, H<sub>2</sub>, CH<sub>4</sub>, C<sub>2</sub>H<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, C<sub>3</sub>H<sub>8</sub>, C<sub>4</sub>H<sub>10</sub>, C<sub>5</sub>H<sub>12</sub>, CH<sub>3</sub>CHCH<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>, CH<sub>3</sub>F, CF<sub>3</sub>H, CF<sub>4</sub>, C<sub>2</sub>F<sub>6</sub>, and SF<sub>6</sub>.

A second major project in our overall research effort was initiated during contract year 1977. An effort is being directed toward the development of new experimental techniques for the direct kinetics study of transient polyatomic free radical species such as  $CF_3$ . During contract year 1977 we have successfully detected  $CF_3$  radicals as products of an RF electrical discharge in mixtures of  $C_2F_6$  and He. Low energy electron bombardment ionization coupled with Time-of-Flight Mass Spectrometry was employed in the detection of these radicals.

## Progress in Research Program During 1977

# A. Electron Spin Resonance (ESR) - Flow Kinetics

One goal of our original proposal was to demonstrate that the ESR-discharge flow technique could be used to measure rate constants for metastable atom quenching reactions. During 1977 we have realized this goal by measuring absolute quenching rates for seventeen reaction systems involving metastable N atoms. These include both systems which have been previously studied by other methods (in order to check for possible systematic errors) and new systems for which comparative data is unavailable.

The flow tube apparatus is diagrammed in fig. 1. A known flow of N<sub>2</sub> and He is discharged to produce the meta-stable N atoms  $\binom{2D_{3/2,5/2}}{2}$ . Reactant gas and SF<sub>6</sub> (to



suppress electron noise) are introduced through the teflon reactant inlet which is a 1/8" dia. teflon tube perforated with approximately 20 small holes pointing upstream. Rate constants are obtained from measurements of  $N({}^{2}D_{5/2})$  ESR signal intensities versus the concentration of reactant gas in the flowtube. The details of these experiments will be described in a future publication (J. Chem. Phys.).

The results of our measurements are summarized in Tables I and II. Table I presents data on systems for which comparative values are available in the literature. The values in Table II represent first determinations. In choosing these systems for study we were motivated in part by the desire to obtain results for a series of similar reactions so that relationships between structure and reactivity might be investigated. For a series of hydrocarbons the length of the carbon chain has been varied. The effect of double and triple bonds in the hydrocarbon has been studied, and the effect of fluorine substitution for hydrogen in a series of fluorinated hydrocarbons has been investigated. These results will be compared with existing models and theories of quenching reactions.

During the final quarter of 1977 we initiated work on the detection of  $O({}^{1}D_{2})$  and  $N({}^{2}P_{J})$ . We have found in our studies of  $N({}^{2}D)$  that cooling the microwave discharge results in a significant increase in the  $N({}^{2}D)$  signal. We have constructed a flowtube which can be cooled to 77° K using liquid nitrogen.

	with various gase	25	
Reactant	Lin & Kaufman	Black et. al.	This work
°2	$4.5 \times 10^{-12}$	$7 \times 10^{-12}$	$4.5 \times 10^{-12}$
N20	$2.2 \times 10^{-12}$	$3 \times 10^{-12}$	$2.4 \times 10^{-12}$
co2	$2.6 \times 10^{-13}$	$6 \times 10^{-13}$	$5.2 \times 10^{-13}$
<sup>H</sup> 2		$5 \times 10^{-12}$	$2.7 \times 10^{-12}$
CH4		$3 \times 10^{-12}$	$3.5 \times 10^{-12}$
с <sub>2</sub> н <sub>4</sub>		$1.2 \times 10^{-10}$	0.60 x 10 <sup>-10</sup>

TABLE I. Rate Constants ( $cm^3/sec$ ) for reactions of N(<sup>2</sup>D) with various gases

References: C.-L. Lin and F. Kaufman, J. Chem. Phys. <u>55</u>, 3760 (1971).

G. Black, T.G. Slanger, G.A. St. John, and R.A. Young, J. Chem. Phys. <u>51</u>, 116 (1969).

	Reactant				
	CH4	C2H6	C3H8	C4H10	C5H12
Number	methane	ethane	n-propane	n-butane	neo-pentane
1	3.3x10 <sup>-12</sup>	2.3x10 <sup>-11</sup>	3.7x10 <sup>-11</sup>	4.5x10 <sup>-11</sup>	3.6x10 <sup>-11</sup>
2	3.7x10 <sup>-12</sup>	2.0x10 <sup>-11</sup>	3.1x10 <sup>-11</sup>	4.2x10 <sup>-11</sup>	3.5x10 <sup>-11</sup>
3	3.4x10 <sup>-12</sup>	2.0x10 <sup>-11</sup>	3.3x10 <sup>-11</sup>	4.7x10 <sup>-11</sup>	3.9x10 <sup>-11</sup>
4	3.2x10 <sup>-12</sup>	2.2x10 <sup>-11</sup>	3.3x10 <sup>-11</sup>	4.4x10 <sup>-11</sup>	3.6x10 <sup>-11</sup>
5	4.0x10 <sup>-12</sup>	2.3x10 <sup>-11</sup>	3.8x10 <sup>-11</sup>	5.6x10 <sup>-11</sup>	3.6x10 <sup>-11</sup>
6	3.0x10 <sup>-12</sup>	2.0x10 <sup>-11</sup>	3.7x10 <sup>-11</sup>	4.0x10 <sup>-11</sup>	3.3x10 <sup>-11</sup>
7	3.7x10 <sup>-12</sup>	2.2x10 <sup>-11</sup>		3.7x10 <sup>-11</sup>	2.9x10 <sup>-11</sup>
8		1.8x10 <sup>-11</sup>		2.3x10 <sup>-11</sup>	
9				2.6x10 <sup>-11</sup>	
Average	3.5×10 <sup>-12</sup>	2.1x10 <sup>-11</sup>	3.5x10 <sup>-11</sup>	4.0x10 <sup>-11</sup>	3.5x10 <sup>-11</sup>
	CH2=CH2	CH3CH=CH2	Сн≅сн	CH3F	CF3H
1	5.0x10 <sup>-11</sup>	11.6x10 <sup>-11</sup>	11.5×10 <sup>-11</sup>	1.2x10 <sup>-12</sup>	0.9x10 <sup>-13</sup>
2	7.3x10 <sup>-11</sup>	11.0x10 <sup>-11</sup>	9.8x10 <sup>-11</sup>	0.68x10 <sup>-12</sup>	1.2x10 <sup>-13</sup>
3	6.8x10 <sup>-11</sup>	5.6x10 <sup>-11</sup>		1.1x10 <sup>-12</sup>	1.3x10 <sup>-13</sup>
4	5.7x10 <sup>-11</sup>	4.3x10 <sup>-11</sup>		1.3x10 <sup>-12</sup>	1.0x10 <sup>-13</sup>
5	5.0x10 <sup>-11</sup>	9.6x10 <sup>-11</sup>		0.9x10 <sup>-12</sup>	1.2x10 <sup>-13</sup>
6		5.6x10 <sup>-11</sup>		1.8x10 <sup>-12</sup>	1.3x10 <sup>-13</sup>
7		8.9x10 <sup>-11</sup>		1.2x10 <sup>-12</sup>	1.3x10 <sup>-13</sup>
8		13.3x10 <sup>-11</sup>			1.5x10 <sup>-13</sup>
9		9.2x10 <sup>-11</sup>			1.4x10 <sup>-13</sup>
Average	6.0x10 <sup>-11</sup>	8.8x10 <sup>-11</sup>	10.6x10 <sup>-11</sup>	1.2x10 <sup>-12</sup>	1.2x10 <sup>-13</sup>

TABLE II. Rate Constants (cm<sup>3</sup>/sec) for reactions of N(<sup>2</sup>D) with various gases

# Table II con't.

There is no detectable quenching of  $N(^{2}D)$  by the following gases:  $CF_4$ ,  $C_2F_6$ , and  $SF_6$  the rate constant must be less than  $10^{-14}$  cc.sec<sup>-1</sup> Additional experiments on  $CH_2=CH_2$  and CH=CH are in progress at the time of this writing. There is evidence that  $O({}^{1}D_{2})$  and  $N({}^{2}P_{J})$  are produced in sufficiently large concentrations in flow discharges and that the problems of detection are associated in large part with wall deactivation. Cooling the walls to 77°K should increase the production of atomic species, reduce the rate of wall deactivation, and increase condensation of molecular impurities which can lower the metastable signals by means of unwanted quenching reactions. Fig. 2 shows a schematic drawing of this flowtube.



B. Polyatomic Free Radical Kinetics by Mass Spectrometry

Electric discharge-fast flow techniques coupled with detection by kinetic spectroscopy have been used extensively to study reactions of free atoms and to some extent of diatomic free radicals. We have initiated a program to extend the application of these techniques to the study of polyatomic free radicals. In order to accomplish this end two experimental problems must be solved. First, relatively pure free radical sources, uncomplicated by atom or radical impurities, are needed. Second, unambiguous methods of detection for polyatomic free radicals are necessary so that the steady state concentration of the radicals can be monitored.

Our work in this area began during this past contract year, 1977, and the initial studies have focused on the production and detection of trifluoromethyl radicals, CF3. Our approach is to develop a simple, clean, and relatively inexpensive source of CF3 radicals in the gas phase and to investigate at least three techniques for the direct measurement of the radical concentration. Selected reactions of CF3 will be studied subsequently. The first source of radicals to be investigated is an RF electrical discharge through a flowing mixture of  $C_2F_6$  and He gas. There is indirect evidence that this source will meet the requirements stated above (R.J. Lagow et.al. J. Amer. Chem. Soc. 97, 518 (1975)), but CF3 radicals have not previously been detected directly from such a source. During this past year we have accomplished the detection of CF, radicals from this source using electron bombardment mass spectrometry. The essential feature to note in these experiments is that the electron energy in the ion source of the mass spectrometer was only 13 eV. Although operating at low electron energy means a significant loss in sensitivity for CF<sub>3</sub> radical detection, it also eliminates spurious signals which result from fragmentation of stable species in the ion source. This point is expanded in a following paragraph.

A schematic diagram of the RF discharge source and the time-of-flight mass spectrometer ion source is shown in Fig. 3.



Three versions of the RF discharge source were designed, contructed, and tested during 1977. Typical operating conditions are shown in the figure. The mass spectrometer is a commercial unit, Bendix model 3012, which we have used without significant modification. Before presenting our results we will discuss the problems associated with electron bombardment ionization and how these problems can be avoided by limiting the electron bombardment energy. The essential problem encountered in the detection of CF<sub>3</sub> radicals by electron bomardment ionization mass spectrometry is that of <u>fragmentation</u>. The desired signal is a current of  $CF_3^+$  ions which is produced by ionization of CF<sub>3</sub> in the ion source as in equation (1).

$$CF_3 + e^- CF_3^+ + 2e^-$$
 (1)

The ionization potential of  $CF_3$  is 10.2 eV. If high energy electrons are used to ionize the radicals (70-100 eV) the efficiency of the ionization process (Eq. 1) is maximized but an additional spurious source of  $CF_3^+$  is also present due to the process of fragmentation represented in Eq. 2.

$$C_2F_6 + e^- CF_3 + CF_3 + 2e^-$$
 (2)

At 70-100 eV electron energy this latter process dominates the former since the concentration of undissociated  $C_2F_6$  is much larger than the CF<sub>3</sub> radical concentration. Therefore the detection of a CF<sub>3</sub><sup>+</sup> signal in the mass spectrum does not necessarily indicate the production of CF<sub>3</sub> radicals in the flow tube. Since the appearance potential of CF<sub>3</sub><sup>+</sup> from the fragmentation of  $C_2F_6$  as in Eq. 2 is approximately 14.5 eV, this source of CF<sub>3</sub><sup>+</sup> signal can be suppressed if the electron energy is lowered below 14.5 eV. For example, if the electron energy is adjusted to 13 eV, there is sufficient energy to ionize the CF<sub>3</sub> radicals but not to fragment C<sub>2</sub>F<sub>6</sub>. Therefore, the  $CF_3^+$  signal at this electron energy comes exclusively from the ionization of  $CF_3^-$  radicals. Unfortunately the ionization efficiency drops drastically as the electron energy is lowered so there is a reduced detection sensitivity at low electron bombardment energy. In our experiments we have made up for this loss of sensitivity by signal averaging the output of the mass spectrometer using a PAR Waveform Educator which accumulates the signal from multiple mass scans and averages out the noise.

The results of  $CF_3$  radical detection are shown in Fig. 4. The operating conditions are recorded in the figure. Each trace is the result of 60 mass scans accumulated over a 3 minute period. The upper trace shows a single mass peak at m/e = 69, the mass of  $CF_3$ . The lower trace shows the affect of turning off the RF discharge and scanning for 3 minutes through the same mass range. The signal also disappears when the discharge is 1 down on but the  $C_2F_6$  flow is turned off. These spectra were obtained at an electron energy of 13 eV. So far we have detected  $CF_3$  radicals with an electron energy as low as 11.5 eV, only 1.4 eV above the appearance potential for  $CF_3^+$  from  $CF_3$ .

In summary, we have successfully completed the initial phase of this research program by showing that direct measurement of  $CF_3$  radical production can be accomplished using low energy electron bombardment ionization mass spectrometry. Initial measurements indicate that an RF electrical discharge in  $C_2F_6$  and He is an efficient source of  $CF_3$  radicals which can be used for kinetic studies.

CF3t peak RF Discharge (23 watts) in A  $C_2F_6$  + He electron energy 13eV

same mass scan as A RF Discharge OFF

В

Fig 4

#### Recent Publications

\*Observation of the Opto-Acoustic Effect in the Microwave Region, G.J. Diebold and D.L. McFadden, Applied Phys. Letters, 29,

447 (Oct. 1, 1976).

The microwave analog of the optoacoustic effect was observed. Collisional relaxation of absorbed microwave energy between magnetic sublevels of gaseous molecular oxygen results in the production of an acoustical signal which is detected by a sensitive microphone.

\*Kinetics of the 0 +  $F_2$  Reaction: A Case of Low Reactivity of Elemental Fluorine, R.H. Krech, G.J. Diebold, and D.L. McFadden, J. Am. Chem. Soc. <u>99</u>, 4605 (1977). This gas phase reaction was studied by the EPR-Discharge Flow Technique. The rate constant obtained is given by the Arrhenius equation,  $k = 10^{12.99} \exp[-10.4/RT] \text{ cm}^3 \text{mor}^1 \text{sec}^1$ . The large activation energy for the reaction is attributed to the low polarizability of molecular fluorine.

\*An Empirical Correlation of Activation Energy with Molecular Polarizability for Atom Abstraction Reactions, R.H. Krech and D.L. McFadden, J. Am. Chem. Soc. <u>99</u>, 8402 (1977). Activation energies for a homologous series of excergic atom transfer reactions are shown to correlate inversely with molecular polarizability. Sixty-five reactions are presented. Predictions of activation energies for some reactions are included.

### Manuscripts in Preparation

\*Signal/Noise Enhancement in Gas Phase EPR Studies of Atomic Nitrogen and Rare Gas Metastables, G.J. Diebold and D.L. McFadden (to be submitted to J. Chem. Phys.).
\*Gas Phase Chemiluminescence in Alkali Halid Systems, D.L. McFadden, to appear in a book on the chemistry of alkali halide vapors, Academic Press, 1978.
\*Absolute Rate Coefficients for the Quenching of Metastable Nitrogen Atoms (N<sup>2</sup>D<sub>3/2,5/2</sub>) by Several Molecules, I.V. Rivas, B. Fell, and D.L. McFadden (J. Chem. Phys.).
\*Acknowledgment is made to ONR for partial support of this work.

D.L. McFadden received an Alfred P. Sloan Research Fellowship for 1977-1979.

# Additional Work

Chemiluminescent Reactions of Boron with Alkali Fluorides,

U.C. Sridharan, D.L. McFadden, and P. Davidovits,

J. Chem. Phys, 65, 5373 (1976).

Chemiluminescence from the Gas Phase Reaction of Atomic Boron with the Alkali Metal Fluorides, U.C. Sridharan, D.L. McFadden, and P. Davidovits, ACS Symposium Ser. 56, 136 (1977).

Chemiluminescence from the Gas Phase Reaction of Atomic Carbon with PbO, U.C. Sridharan, T. DiGiuseppi, D.L. McFadden and P. Davidovits (to be submitted to J. Chem. Phys.).

The Alkali Halide Vapors, P. Davidovits and D.L. McFadden, eds., Academic Press, New York, 1978.

Rate Coefficients for the Reactions,  $B + O_2 \rightarrow BO + O$  and Al +  $N_2 O \rightarrow AlO + N_2$  Using a Diffusion Technique, U.C. Sridharan, T.G. DiGiuseppi, D.L. McFadden, and P. Davidovits (To be submitted to J. Chem. Phys.).

#### DISTRIBUTION LIST

Office of Naval Research Code 473 Arlington, VA 22217 Attn: Dr. Miller

Office of Naval Research Branch Office 1030 East Green Street Pasadena, CA 91106 Attn: Dr. R. Marcus

Office of Naval Research Branch Office 536 S. Clark Street Chicago, IL 60605 Attn: Dr. J. Smith

Defense Documentation Center Bldg. 5 Cameron Station Alexandria, VA 22314

Office of Naval Research Branch Office 495 Summer Street Boston, MA 02210 Attn: Dr. L.H. Peebles

U.S. Naval Research Laboratory Code 2627 Washington, D.C. 20375

U.S. Naval Research Laboratory Code 2629 Washington, D.C. 20375

U.S. Army Research Office Chemistry Division P.O. Box 12211 Research Triangle Park, NC 27709

U.S. Army Research Office Physics Division P.O. Box 12211 Research Triangle Park, NC 27709

Professor S.N.B. Murthy Technical Director, Project SQUID Purdue University Lafayette, IN 47907 Naval Research Laboratory Code 6100 Washington, D.C. 20375

Naval Air Systems Command Code 440 Washington, D.C. 20360 Attn: Dr. H. Rosenwasser

Naval Sea Systems Command SEA-0331 Washington, D.C. 20362 Attn: Mr. J. Murrin

Naval Sea Systems Command SEA-0332 Washington, D.C. 20362 Attn: Dr. Amster

Naval Surface Weapons Center Research and Technology Dept-WR Silver Spring, MD 20910

Naval Weapons Center Code 608 China Lake, CA 93555 Attn: Ronald L. Derr

Air Force Office of Scientific Research Directorate of Aerospace Sciences Bolling Air Force Base Washington, D.C. 20332

Air Force Office of Scientific Research Directorate of Chemical Sciences Bolling Air Force Base Washington, D.C. 20332

Air Force Office of Scientific Research Directorate of Physics Bolling Air Force Base Washington, D.C. 20332