

UNITED STATES AIR FORCE AIR WEATHER SERVICE (MAC) USAF ENVIRONMENTAL TECHNICAL APPLICATIONS CENTER B-105198 SCOTT AIR FORCE BASE, ILLINOIS 62225 USAFETAC-PR-78-002 POINT WARNING CLIMATOLOGY OF MT. LAGUNA, CALIFORNIA by James R. Clark, Capt, USAF January 1978 Approved for public release: distribution unlimited DC POP ner MAR SO 1978

Review and Approval Statement

This report is approved for public release. There is no objection to unlimited distribution of this report to the public at large, or by DDC to the National Technical Information Service (NTIS).

This technical report has been reviewed and is approved for publication.

James R. Clark

JAMES R. CLARK Capt, USAF

Calicin C. Maegel

CALVIN C. NAEGELIN Capt, USAF Reviewing Officer

FOR THE COMMANDER

WALTER S. BURGMAN

Scientific & Technical Information Officer (STINFO)

REPORT DOCUMENTATION	READ INSTRUCTIONS BEFORE COMPLETING FORM				
USAFETAC-PR-78-002	2. GOVT ACCESSION NO.	3. RECIPIENT'S C	ATALOG NUMBER		
POINT WARNING CLIMATOLOGY O MT. LAGUNA, CALIFORNIA	F	5. TYPE OF REPO	ORT & PERIOD COVERED		
	6. PERFORMING ORG. REPORT NUMBER				
James R. Clark, Capt, USAF	8. CONTRACT OR GRANT NUMBER(*)				
PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Air Force Environmenta Applications Center Scott AFB, IL 62225	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS				
1. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE			
U.S. Air Force Environmental	l Technical	January	1978		
Applications Center Scott AFB, IL 62225	13. NUMBER OF PAGES				
4. MONITORING AGENCY NAME & ADDRESS(If different	nt from Controlling Office)	15. SECURITY CL	ASS. (of this report)		
		UNCLASSIFIED			
		15a. DECLASSIFI	CATION/DOWNGRADING		
7. DISTRIBUTION STATEMENT (of the abstract entered	i in Biock 20, 11 different fro	em Report)	BETTETENTION ST STETETENTION/AVAILABILITY C Bost. AVAIL and/or SPI		
8. SUPPLEMENTARY NOTES			A		
9. KEY WORDS (Continue on reverse aide if necessary a Meteorology Precipitation Temperature	nd identify by block number) Mt. Laguna Chubascos Tropical S	, CA torms			
 ABSTRACT (Continue on reverse side if necessary of A 17-year period of record v criteria. The four major st California are discussed. (precipitation, and wind are warning elements. 	was analyzed for torm types that Climatological presented mont	or weather t affect so values of thly for ir	warning outhern temperature, ndividual		

PREFACE

The USAF Environmental Technical Applications Center (USAFETAC) prepared this report to answer a request from Detachment 11, 12th Weather Squadron, Luke AFB, AZ. They requested climatological data to assist in issuing point weather warnings for a radar site at Mt. Laguna, California. Physical separation and geographical location make point warning forecasting for Mt. Laguna extremely difficult for the Luke AFB forecaster.

Climatological data were obtained from USAFETAC/OL-A, Asheville, NC and Mrs. Dorothy L. Haak, the National Weather Service Cooperative Observer at Mt. Laguna. As a result, a 17-year period of record was available for some point warning elements. Mrs. Haak's voluntary assistance is gratefully acknowledged.

This report answers a specific request and is not expected to have application beyond that request. Further questions on this problem or related problems should be referred to USAFETAC for consultation and study.

If the requestor or any other agency incorporates this report, in whole or in part, into another report, we request that USAFETAC be given proper credit and be furn-ished a copy of the new report if possible.

5 ...

POINT WARNING CLIMATOLOGY OF MT. LAGUNA, CALIFORNIA

Introduction

Mt. Laguna, California, is located approximately 45 miles east of San Diego and along the eastern edge of the Peninsula Mountain Range (see Figures 1a and 1b). In contrast to the Transverse Range, the eastern flank of the Peninsula Range plunges to the low desert floor. The Peninsula Range runs parallel to the coast and continues down into Mexico. Elevation of Mt. Laguna is approximately 6000 feet.

Storm Types Affecting Southern California

Blake (1933) and Reed (1932) categorized the weather patterns affecting southern California into four broad types. The North Pacific type includes all low-pressure systems that approach the mainland from the Pacific Ocean north of San Francisco. These northern storm tracks move southward with the approach of winter, affecting southern California from November to May. Of the four weather patterns, this type is the most prevalent and is responsible for the majority of winter precipitation. However, large excesses and deficiencies in seasonal precipitation totals are more likely due to an unusual number of storms of the other types than to variations in the North Pacific type.

The second weather pattern, designated as the South Pacific type, includes all disturbances that come from the Pacific Ocean south of San Francisco and north of the Tropic of Cancer. The majority of South Pacific storms form during November through March with a maximum in February. The storms may develop in several ways. They sometimes develop along the southern California coast on the tropical side of an eastern Pacific high pressure ridge, or they may move along the coast after passing around the ridge's eastern flank. The majority of the South Pacific type lows form in this second manner. Occasionally, secondary depressions may form in the lower end of a trough that lies near the southern California coast and move basically in the same direction as the other two major systems in this type. Precipitation amounts for the South Pacific storm type vary widely.

The third weather pattern, designated the interior type, consists of all active depressions that originate or develop over the plateau regions, the Colorado Valley, or the California interior. These complicated disturbances form primarily in the Great Basin, but may originate anywhere over the far western interior. Their growth and movement is erratic and hard to predict. The interior storms are most active from February to May, but they may form any month of the year.

The fourth weather pattern, designated the Mexican type, consists of two distinct phenomena: the tropical disturbances that occasionally move northward to southern California from the west coast of Mexico from August to November, and the few sporadic thunderstorms that develop in the mountains and deserts of lower and southern California from the middle of June to the middle of September. These sporadic thunderstorms are locally known as "Sonoras." The Mexican type is responsible for the smallest number of southern California storms, but may produce the most locally heavy precipitation. Eastern Pacific tropical cyclones, locally called "Chubascos," originate in the tropics south and west of Baja, California. They then move northward across western Mexico and southwestern California, again bringing locally heavy precipitation.

Climatology of Some Weather Warning Criteria

Precipitation. Precipitation in southern California is light except in the high mountains. According to the Climatic Summaries of the US, 1930, more than 75% of the total annual precipitation occurs during the winter months. In some areas of the mountains, excessively heavy rains may occur within short periods of time. The topography, elevation, and exposure have significant effect on the recorded precipitation amounts.

Mt. Laguna receives about 16.5 inches of precipitation annually. Approximately 20% of that precipitation occurs in December. Table 1 gives the average monthly precipitation at Mt. Laguna. The extreme values of precipitation illustrate the variability of precipitation in southern California.

Rainfall. Climatological records for Mt. Laguna from June 1943 to February 1950 indicate no rain days with rainfall greater than 2 inches per day. However, heavy

rainfall associated with strong thunderstorms or dissipating "Chubascos" may reach or exceed that amount (Fors, 1977). The "Chubascos" that approach or cross southern California produce the greatest single amounts of precipitation in the mountains. For example, thirteen inches were recorded at Mt. Wilson in September 1939 when a tropical cyclone moved inland near San Pedro. Mt. Laguna recorded 10.13 inches from Hurricane Kathleen in September 1976 (Fors, 1977) and over 4 inches from Doreen in August 1977 (Haak, 1977).

Snowfall. Snow accounts for about 50% of the precipitation received at Mt. Laguna during the winter months. Snow may occur from October through May, but the largest amounts are received in December. The annual average is about 61 inches. Days with snowfall greater than 2 inches in a 24-hour period occur most often from November through April.

Freezing Precipitation. Mt. Laguna receives freezing precipitation an average of 10 days a year. March has the largest number of days (3) with freezing precipitation. December, February, and April experience freezing precipitation an average of two days per month.

Temperature. Temperature extremes can vary greatly at Mt. Laguna (Table 1). Occasionally, January and May experience differences in the extreme maximum/minimum temperatures of over 64°F. Average maximum/minimum temperature differences vary from 16 to 23°F with the greatest deviations occurring in May.

<u>Tropical Cyclones</u>. In the past 60 years only five major tropical cyclones have affected California. Four of these storms caused significant precipitation in southern California. The most destructive was Kathleen in September of 1976 (Fors, 1977). Kathleen moved onshore about 190 nautical miles south of San Diego and moved northeast into the California desert region. During this period 10.13 inches of rain were recorded at Mt. Laguna, 10.78 inches at Mt. Wilson, and 14.78 inches on San Gorgonia Mountain northwest of Palm Springs. Thirteen inches of rain were recorded at Mt. Wilson in September of 1939 when a tropical cyclone moved inland near Los Angeles. That same year, two other cyclones passed less than 200 miles south of San Diego.

September is the month of the greatest frequency of Eastern North Pacific tropical cyclones and the month that most recurve into Mexico (Fors, 1977). Tropical cyclones, which follow paths that affect Mt. Laguna, usually develop in the first two weeks in September, with a maximum occurring around 10 September (Baum, 1975).

Thunderstorms. Blake (1933) associates two major synoptic patterns with most of the thunderstorms over the southern California mountains. One pattern consists of air approaching southern California from the south and east. The resulting "Sonoras" usually originate over the great deserts of lower and southern California and reach their maximum development over the mountains. The other pattern brings in moist, tropical air from dissipating tropical cyclones that move far enough north along the west coast of Mexico to affect southern California.

The Peninsula Range averages about 18 thunderstorm days per summer (Tubbs, 1972). At Mount Laguna the majority of thunderstorms occur from July through September with the maximum in August. Mt. Laguna averages ten thunderstorm days a year.

Hail is possible at Mt. Laguna during all months (Table 1); however, it is rare during the winter months. August and September have the most occurrences. Hail size varies with season but rarely exceeds one-half inch (Haak, 1977).

Tornadoes. Tornadoes are infrequent in southern California. One such occurrence near San Diego on Pebruary 23, 1971, was associated with a strong frontal system. Such rare occurrences are most often associated with thunderstorms or frontal systems. Tornadoes are not considered to be a weather warning problem at Mt Laguna.





Table 1. Climatological Data for Mt. Laguna, California, Station Number 23178, for the Period of Record from Jun 43 to Feb 50, and Jul 66 to Sep 77.

.

О.

0

Annual		42		16.48		60.6 - 0		001 40 #		•	
Dec		67 47 31		3.33 11.28 0.32		16.4 80.5 0				93	
Nov		76 54 37		1.95 6.02 0.22		2.1 9.8 0		0-1 #00		89	
Oct		77 626 44		0.84 3.04 0		0.6 0.0		0*0**		78	
Sep		90 32 52		1.17 10.69 0		000		0000#		75	one day
Aug		94 41 58		1.24 5.42 0		000		0000#		62	than c
Jul		6440 59440		0.59 3.12 0		000		0000#		64	+ Less
Jun		88 28 52		0.15 1.3 0		000		0*0**		114	
May		603 603 603 603 603 603 603 603 603 603		0.38 1.76 0		0.4 0.0		0*0 **		86]	onlv
Apr		78 56 34		0.99 3.68 0.09		7.6 45.8 0		0 1 * * *		95	Feb 50
Mar		72 51 32		2.44		13.3 61.5 0		o~***		100	1e 43 -
Feb		69 49 22		1.97 7.34 0		8.5 49.0		00*0*		100	IUC**
Jan		73 88 31 88 81		1.92 8.61 0		9.8 43.5	(sýe	•••••		125	vinc
	emperatures (^o F)	Extreme Max Extreme Min Average Max* Average Min*	recipitation (in)	Average Extreme Max Extreme Min	Snow (in)	Average Extreme Max Extreme Min	ccurrence (No. De	Precip >2 (in)* Freezing Precip Snow >2 (in)** Thunderstorm Hail	(ind (mph)	Extreme**	Jul 66 - Sep 77 6

5

REFERENCES

Baum, R. A., and Rasch, G. E.: "Digitized Eastern Pacific Cyclone Tracks," NOAA Technical Memorandum, NWS WR-101, 1975, 189 p.

Blake, Dean: "Sonora Storms," Monthly Weather Review, November 1923, pp 585-588.

- Blake, Dean: "Storm Types and Resultant Precipitation in the San Diego Area," Monthly Weather Review, Vol. 61, No. 8, August 1933, pp 223-225.
- Coffin, Harry: "Marine Influences on the Climate of Southern California in the Summer," PhD Dissertation, Department of Geography, University of California, Berkeley, 1961, 147 p.
- Court, Arnold: "Lightning Fire Incidence in Northeastern California," Technical Paper No. 47, U. S. Forest Service, Pacific Southwest Forest and Range Experimental Station, Berkeley, California, May 1960, 21 pp.
- Fors, James R.: "Tropical Cyclone Kathleen." National Weather Digest, Vol. 2, No. 3, August 1977, pp 6-20.

Haak, Dorothy L., personal communication, September 1977.

- Halvorson, D. A.: "Tornado and Funnel Clouds in San Diego County," Western Region Technical Attachment No. 71-33, July 21, 1971.
- Markham, Charles G.: "Baja California's Climate," Weatherwise, Vol. 25, No. 2, 1972.
- Reed, Thomas R.: "Weather Types of the Northeast Pacific Ocean as Related to the Weather of the North Pacific Coast," Monthly Weather Review, Vol. 60, December, 1932, pp 246-252.
- Rosendal, Hans E.: "Eastern North Pacific Tropical Cyclones, 1947-1961," Mariner's Weather Log, Vol. 6, No. 6, November 1962, pp 195-201.
- Serra C, Sergio: "Hurricanes and Tropical Storms of the West Coast of Mexico," Monthly Weather Review, Vol. 99, No. 4, April 1971, pp 302-308.
- Tubbs, Anthony M.: "Summer Thunderstorms Over Southern California," Monthly Weather Review, Vol. 100, No. 11, November 1972, pp 799-807.
- U.S. Department of Commerce, Weather Bureau, "Southern California and Owens Valley," <u>Climatic Summaries of the U.S. 1930</u>, Section 18, U.S. Government Printing Office, Washington, D.C.

6