

AD-A052 145

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/G 7/2
THE REACTION OF 3,3-DICHLOROALLYLTRIMETHYLSILANE WITH N-BUTYLLI--ETC(U)
MAR 78 D SEYFERTH, R E MAMMARELLA N00014-76-C-0837

NL

UNCLASSIFIED


1 OF 1
AD
A062145



END
DATE
FILMED

5-78

DDC



AD A 052145

AD No. _____
ODC FILE COPY

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-76-C-0837

Task No. NR 053-618

THE REACTION OF 3,3-DICHLOROALLYLTRIMETHYLSILANE
WITH n-BUTYLLITHIUM.

by

Dietmar Seyferth Robert E. Mammarella

Prepared for Publication

in the

Journal of Organometallic Chemistry

Massachusetts Institute of Technology
Department of Chemistry, 18-411
Cambridge, Massachusetts 02139

March 27, 1978

Reproduction in whole or in part is permitted for
any purpose of the United States Government

*Approved for Public Release; Distribution Unlimited

220 007 JOB

THE REACTION OF 3,3-DICHLOROALLYLTRIMETHYLSILANE WITH n-BUTYLLITHIUM.

Dietmar Seyferth* and Robert E. Mammarella#

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139 (USA)

SUMMARY


n-Butyllithium reacts with 3,3-dichloroallyltrimethylsilane to metalate the vinyl proton. Under the reaction conditions the $\text{Me}_3\text{SiCH}_2\text{C}(\text{Li})=\text{CCl}_2$ formed undergoes β -elimination of LiCl to give $\text{ClC}\equiv\text{CCH}_2\text{SiMe}_3$ whose subsequent reaction with n-butyllithium produces $\text{LiC}\equiv\text{CCH}_2\text{SiMe}_3$. Addition of trimethylchlorosilane gives $\text{Me}_3\text{SiC}\equiv\text{CCH}_2\text{SiMe}_3$. When two molar equivalents of n-butyllithium are used, further metalation of $\text{LiC}\equiv\text{CCH}_2\text{SiMe}_3$ gives $\text{LiC}\equiv\text{CCH}(\text{Li})\text{SiMe}_3$. The action of N-bromosuccinimide on $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ resulted in formation of $\text{Me}_3\text{SiCH}=\text{CHCCl}_2\text{Br}$.

SEARCHED	INDEXED	SERIALIZED	FILED
DIST. BY MAIL			
SP. CHAL			
A			


National Science Foundation Graduate Fellow, 1973-1976.

INTRODUCTION.

In a recent investigation,¹ ~~we~~ prepared the novel ambident allylic lithium reagent gem-chloro(trimethylsilyl)allyllithium, ~~was prepared~~ ^{WAS PREPARED} and ~~studied~~ its reactions with a variety of substrates ^{WAS STUDIED.}



Another trimethylsilyl-substituted allyllithium reagent which was of potential interest to us was II, and we report here con-




cerning our attempts to prepare this reagent. Although they were unsuccessful, the route examined was of interest since it involved the reaction of n-butyllithium with a molecule which had three different potential sites at which the lithium reagent could attack.

2 —


RESULTS AND DISCUSSION.

The organosilicon compound which was chosen as a potential precursor for II was 3,3-dichloroallyltrimethylsilane, $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$.^{*} This was readily prepared. The condensation of trichlorosilane with 1,1,3-trichloropropene by the method of Furuya and Sukawa² (eq. 1) gave $\text{Cl}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ whose methylation



with methylmagnesium bromide produced the desired silane.

The hoped-for reaction of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ with *n*-butyllithium is the one shown in eq. 2. Metalation α to a silyl group is a



favorable process since silicon stabilizes an adjacent negative charge.³ In fact, trimethylsilylallyllithium, $\text{Li}[\text{Me}_3\text{SiCHCHCH}_2]$, is readily prepared by reaction of allyltrimethylsilane with *n*-butyllithium.^{4,5} However, the two chlorine atoms in $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ also will have activating effects on the attack by *n*-butyllithium, and these will oppose the desired metalation process. On the one hand, they will enhance the acidity of the vinyl proton,

* The action of *n*-butyllithium on the isomeric $\text{Me}_3\text{SiCCl}_2\text{CH}=\text{CH}_2$ was reported in ref. 1.

making it more reactive toward a base; on the other hand, the two chlorine substituents will mutually enhance each other's reactivity toward lithium/chlorine exchange with an alkyllithium. Thus $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ offers three potential sites for attack by n-butyllithium.

In the initial experiment, a molar equivalent of n-butyllithium in hexane was added to a solution of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ in tetrahydrofuran (THF) which was cooled to -90° . After 30 min., an excess of trimethylchlorosilane was added to the reaction mixture. The products were an acetylenic silane, 2,2,6,6-tetramethyl-2,6-disila-3-heptyne, $\text{Me}_3\text{SiC}\equiv\text{CCH}_2\text{SiMe}_3$, in 26% yield, and 2,2,6,6-tetramethyl-2,6-disila-3-chloro-3-heptene, $\text{Me}_3\text{SiC}(\text{Cl})=\text{CH}-\text{CH}_2\text{SiMe}_3$, in 6% yield. A 68% recovery of the starting silane was realized.

Further experiments (Table 1) showed that longer reaction times or higher temperatures increase the yield of the acetylene. The last three experiments show roughly the same result, that just over one-half of the starting material is converted to the acetylene. Apparently, the conditions of the first experiment did not allow the reaction to go to completion.

The fact that about one-half of the starting material remains unreacted suggests that two molar equivalents of n-butyllithium are consumed per equivalent of substrate. However, doubling the amount of the lithium reagent caused more deprotonation of the substrate, and on reaction of the organolithium intermediates with trimethylchlorosilane, 1,3,3-tris(trimethylsilyl)propyne, $\text{Me}_3\text{SiC}\equiv\text{CCH}(\text{SiMe}_3)_2$, was formed in 19% yield,

TABLE 1. Reactions of 3,3-Dichloroallytrimethylsilane with One Molar Equivalent of n-Butyllithium. Trimethylchlorosilane Quench.


<u>Reaction Temperature, °C</u>	<u>Reaction Time, min.</u>	<u>Relative Yields*</u>	
		<u>Me₃SiC≡CCH₂SiMe₃</u>	<u>Me₃SiCH₂CH=CCl₂</u>
-90°	30	26**	68**
-90°	150	59	41
-46°	30	55	45
- 1°	30	53	47

* Relative ratio determined by GLC.

** Absolute yield determined by GLC (internal standard method).


in addition to $\text{Me}_3\text{SiC}\equiv\text{CCH}_2\text{SiMe}_3$ (37%). Such multiple substitution had been observed by West et al.⁶, who obtained the same tris-(trimethylsilyl)propyne in a reaction of an excess of *n*-butyllithium with propyne with subsequent addition of trimethylchlorosilane.

These observations are explicable in terms of initial metalation of the vinyl proton of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$. Previous work reported by Köbrich and his coworkers⁷⁻⁹ is pertinent. Low temperature reactions of *n*-butyllithium with equimolar quantities of mono-, di-(*cis*- and *trans*-) and trichloroethylene were found to result in metalation of the vinylic hydrogens to give α -chlorovinyl lithium compounds (eq. 3). The stabilities of



(R, R' = H and/or Cl)

these products depended on their configuration. The least stable were those with vicinal chlorine and lithium substitution in *trans* relationship. These decomposed by *trans*-elimination to give an acetylene (eq. 4). In those cases where a chloro-




(R, R' = H and/or Cl)


acetylene was produced, further reaction with *n*-butyllithium gave

alkynyllithium reagents. Such chemistry is involved in the $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2/\underline{n}\text{-C}_4\text{H}_9\text{Li}$ interaction.


The initial step must be metalation (eq. 5). The product



undergoes trans- β -elimination of lithium chloride (eq. 6).




The chloroacetylene produced reacts with n-butyllithium, giving Li/Cl exchange (eq. 7). The reaction sequence (5,6,7) utilizes



III


two molar equivalents of n-butyllithium per mol of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$. If excess n-butyllithium is present, the alkynyllithium reagent III undergoes further metalation to give the dilithium species IV (eq. 8). Addition of trimethylchlorosilane to such reaction



mixtures results in formation of the silylation products of III and IV, $\text{Me}_3\text{SiC}\equiv\text{CCH}_2\text{SiMe}_3$ and $\text{Me}_3\text{SiC}\equiv\text{CCH}(\text{SiMe}_3)_2$, respectively.

Since the initial lithium reagent produced (eq. 5) is only a transient species which undergoes rapid β -elimination of lithium chloride, it is clear why, as Table 1 shows, low reaction temperatures are not required. The first lithium reagent which persists, III, is stable at room temperature.

The small amount of 2,2,6,6-tetramethyl-2,6-disila-3-chloro-3-heptene formed in the $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2/\underline{n}\text{-C}_4\text{H}_9\text{Li}/\text{Me}_3\text{SiCl}$ reaction results from competitive Li/Cl exchange as shown in eq. 9.




Reagent V, an α -chlorovinyl lithium species, is sufficiently stable for reaction with trimethylchlorosilane to take place, giving $\text{Me}_3\text{SiC}(\text{Cl})=\text{CHCH}_2\text{SiMe}_3$. It is clear that the β -Cl effect on the acidity of a vinyl proton is greater than the α -silicon effect on the acidity of a methylene proton, and thus the desired chemistry (eq. 2) did not develop.


Our successful route to gem-chloro(trimethylsilyl)allyl-lithium involved the transmetalation reaction between $\text{Ph}_3\text{PbCH}_2\text{CH}=\text{C}(\text{Cl})\text{SiMe}_3$ and n-butyllithium. The required organolead precursor was prepared by reaction of Ph_3PbMgBr with $\text{Me}_3\text{SiC}(\text{Cl})=\text{CH}-\text{CH}_2\text{Cl}$.* A similar route should, in principle, be applicable to the synthesis of $\text{Ph}_3\text{PbCH}(\text{SiMe}_3)\text{CH}=\text{CCl}_2$, VI, and via reaction of

* Such a procedure also served well in the synthesis of gem-dichloroallyllithium¹⁰: $\text{CCl}_2=\text{CHCH}_2\text{Cl} \xrightarrow{\text{Ph}_3\text{PbMgBr}} \text{CCl}_2=\text{CH}-\text{CH}_2\text{PbPh}_3 \xrightarrow{n\text{-BuLi}} \text{Li}[\text{CCl}_2\text{CHCH}_2]$. It is apparent that such transmetalation processes occur more readily than vinyl proton metalation β to a vinylic chlorine atom.

the latter with n-butyllithium, of $\text{Li}[\text{Me}_3\text{SiCHCHCl}_2]$. This would require a suitable starting material for the preparation of the required lead compound. Allylic bromination of $\text{Me}_3\text{SiCH}_2\text{-CH}=\text{CCl}_2$ might provide a route $\text{Me}_3\text{SiCHBrCH}=\text{CCl}_2$, whose reaction with Ph_3PbMgBr then would be expected to give the desired VI. However, such allylic bromination of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ could well proceed with allylic rearrangement, giving $\text{Me}_3\text{SiCH}=\text{CHCCl}_2\text{Br}$ rather than the desired bromide. Indeed, Corriu et al.⁴ had reported that allylic bromination of allyltriphenylsilane with N-bromosuccinimide gave the rearranged bromide, $\text{Ph}_3\text{SiCH}=\text{CHCH}_2\text{Br}$. Nevertheless, we examined the allylic bromination of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ with N-bromosuccinimide in carbon tetrachloride in the presence of a catalytic amount of benzoyl peroxide. This reaction gave a 91% yield of a monobromination product. The NMR and IR spectra of this product, however, showed it to be $\text{Me}_3\text{SiCH}=\text{CHCCl}_2\text{Br}$, rather than the desired $\text{Me}_3\text{SiCHBrCH}=\text{CCl}_2$. Presumably, rapid rearrangement of the initially formed radical, VII, to the more stable VIII took place before bromine atom abstraction occurred.



VII



VIII

Such allyl radical rearrangements have been discussed in a review by Walling.¹¹

It would appear that there is no easy route to $\text{Li}[\text{Me}_3\text{SiCHCHCCl}_2]$.

EXPERIMENTAL.

General Comments. The "general comments" in the preceding paper¹ are applicable.

Preparation of 3,3-Dichloroallyltrichlorosilane and 3,3-Dichloroallyltrimethylsilane.

a) $\text{Cl}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$.

A two-liter, three-necked round-bottomed flask, equipped with a 500 ml pressure-equalizing addition funnel, a mechanical stirrer and a Claisen adapter fitted with a no-air stopper and a nitrogen inlet, was charged with 4.95 g (50 mmol) of CuCl , 139 ml (1.0 mol) of triethylamine (distilled from NaOH pellets) and 500 ml of diethyl ether. The addition funnel was topped with a no-air stopper and charged by syringe with 100.0 ml (1.0 mol) of 1,1,3-trichloropropene and 99.7 ml (1.0 mol) of trichlorosilane. The Claisen adapter was replaced with a reflux condenser topped with a nitrogen inlet tube and then the contents of the addition funnel were added dropwise during 80 min. while the reaction mixture was stirred and maintained at reflux. During the course of the addition the following color changes were observed: initially, olive green to colorless (after the first drops had been added) to yellow, then orange and, finally, brown. Subsequently, the addition funnel and the reflux condenser were

replaced by a glass stopper and a still head. About 400 ml of a trichlorosilane/diethyl ether mixture was distilled at 37°.

The residue was transferred, under nitrogen, to a distillation flask and was trap-to-trap distilled at 0.03 mm, temperature to ~150°, into a receiver cooled with liquid nitrogen. The distillate was fractionally distilled at reduced pressure to give 85.0 g (35%) of $\text{Cl}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$, bp 87-88.5° at 26 mm Hg.

(Found: C, 14.99; H, 1.32. $\text{C}_3\text{H}_3\text{Cl}_5\text{Si}$ calcd.: C, 14.74; H, 1.24). NMR (CCl_4): δ 2.49 (d, $J = 8$ Hz, 2H, CH_2) and 5.87 ppm (t, $J = 8$ Hz, 1H, $=\text{CH}$).

b) $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$.

The reaction of 85.0 g (0.35 mol) of $\text{Cl}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ with methylmagnesium bromide in diethyl ether from 1.40 mol of magnesium, using standard Grignard alkylation procedure (non-hydrolytic work-up), gave 54.35 g (85%) of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$, bp 73-74° at 31 mm Hg, n_D^{25} 1.4604. (Found: C, 39.47; H, 6.63; Cl, 38.77. $\text{C}_6\text{H}_{12}\text{Cl}_2\text{Si}$ calcd.: C, 39.35; H, 6.60; Cl, 38.71). NMR (CCl_4): δ 0.12 (s, 9H, Me_3Si), 1.65 (d, $J=9$ Hz, 2H, CH_2) and 5.85 (t, $J = 9$ Hz, 1H, $=\text{CH}$). IR(film): $\nu(\text{C}=\text{C})$ 1615 cm^{-1} .

Reaction of 3,3-Dichloroallyltrimethylsilane with n-Butyllithium.

A 500 ml, three-necked Morton flask, equipped with a mechanical stirrer, no-air stopper and a Claisen adapter fitted with a low-temperature thermometer and a nitrogen inlet tube, was charged with 2.874 g (15.7 mmol) of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ dissolved in 250 ml of THF. The solution was cooled to $-91^\circ \pm 2^\circ \text{C}^*$ and then

a 2.57 N solution of n-butyllithium (17.2 mmol) in hexane was

* Reaction temperatures are uncorrected. They were obtained using a pentane (total immersion type) thermometer which was immersed into the reaction mixture to a depth of ~3 cm. A stem correction of 8-10°C (to lower temperature) is appropriate.

added by syringe over a 15 min. period. The resulting clear, lavender solution was stirred at -90°C for 30 min. and then 2.60 ml (~20 mmol) of trimethylchlorosilane was added during a 2 min. period. The reaction mixture was stirred at -90°C for 30 min. The clear, colorless solution which resulted was allowed to warm slowly to room temperature and then was hydrolyzed to a dry end-point with saturated aqueous ammonium chloride. Filtration was followed by concentration of the filtrate at reduced pressure. GLC analysis of the residue (4 ft., 20% SE-30 at 110° and 6 ft. Carbowax at 110°) showed the presence of three components. These were isolated by GLC and were identified as starting material, $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$, 10.7 mmol, 68% recovery (by IR and NMR); $\text{Me}_3\text{SiC}\equiv\text{CCH}_2\text{SiMe}_3$, 4.08 mmol (26%), whose NMR spectrum [in CCl_4 : δ 0.17 (s, 18H, Me_3Si), and 1.52 ppm (s, 2H, CH_2)] matched that reported for this compound by West and Jones⁶; $\text{Me}_3\text{SiC}(\text{Cl})=\text{CHCH}_2\text{SiMe}_3$, 0.94 mmol (6%), n^{25}_D 1.4530 (lit.¹ n^{25}_D 1.4529), whose NMR and IR spectra matched those of an authentic sample (cf. preceding paper¹).

The results of other 1:1 $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2/\underline{n}\text{-BuLi}/\text{Me}_3\text{SiCl}$

reactions are given in Table 1.

A similar reaction was carried out in which 3.63 mmol of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ in 200 ml of THF at 1°C was treated with 7.97 mmol of *n*-butyllithium in hexane. The caramel-colored reaction mixture was treated with 10 mmol of trimethylchlorosilane. Work-up as above, followed by GLC analysis and product isolation by GLC, showed the following compounds to be present: $\text{Me}_3\text{SiC}\equiv\text{CCH}_2\text{-SiMe}_3$, 1.34 mmol (37%) and $\text{Me}_3\text{SiC}\equiv\text{CCH}(\text{SiMe}_3)_2$, 0.69 mmol (19%), whose IR and NMR spectra matched the published spectra of this compound^{6,12} [NMR (CCl_4): δ 0.15 (s, 27H, Me_3Si) and 1.05 ppm (s, 1H, CH)].

Reaction of 3,3-Dichloroallyltrimethylsilane with N-Bromo-succinimide.

A 200 ml, three-necked Morton flask, equipped with a reflux condenser and a nitrogen inlet tube was charged with 8.898 g (48.6 mmol) of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$, 9.225 g (46.2 mmol) of N-bromosuccinimide, a spatula tip-ful of benzoyl peroxide and 100 ml of carbon tetrachloride. The mixture was stirred and heated at reflux for 5 hr., at which time iodide-starch paper was negative. The succinimide was filtered and the filtrate was trap-to-trap distilled (room temperature at 0.03 mm Hg). The distillate was concentrated (to 75° , 33 mmHg) and the residue was trap-to-trap distilled at 0.07 mm (heat gun), to give 11.6 g (91%) of $\text{Me}_3\text{SiCH}=\text{CHCCl}_2\text{Br}$, n_D^{25} 1.5082 (Found: C, 27.95; H, 4.36; total halogen, 56.83; $\text{C}_6\text{H}_{11}\text{Cl}_2\text{BrSi}$ calcd.: C, 27.50; H, 4.23; total halogen, 57.56). NMR (CCl_4): δ 0.23 (s, 9H, Me_3Si), 4.03

(d, J = 12Hz , 1H, Si-CH=), 6.00 ppm (d, J = 12Hz, 1H, =CHCCl₂Br).

The NMR spectrum was almost superimposable with that of Me₃SiCH=CH-
CBr₃.¹³ IR(film): ν (C=C) 1605 sh, 1595 cm⁻¹.

Acknowledgments. This work was supported in part by the Office of Naval Research. Gifts of chemicals from Union Carbide Corporation are gratefully acknowledged.

REFERENCES

1. D. Seyferth and R. E. Mammarella, preceding paper: J. Organometal. Chem.
2. N. Furuya and T. Sukawa, J. Organometal. Chem., 96 (1975) C1.
3. D. J. Peterson, Organometal. Chem. Rev. A, 7 (1972) 295.
4. R. J. P. Corriu, J. Masse and D. Samate, J. Organometal. Chem., 93 (1975) 71.
5. D. Ayalon-Chass, E. Ehlinger and P. Magnus, J. Chem. Soc. Chem. Comm. (1977) 772.
6. R. West and P. C. Jones, J. Am. Chem. Soc., 91 (1969) 6156.
7. G. Kobrich and K. Flory, Chem. Ber., 99 (1966) 1773.
8. G. Kobrich and K. Flory, Tetrahedron Lett. (1964) 1137.
9. G. Kobrich, H. Trapp and I. Hornke, Tetrahedron Lett. (1964) 1131.
10. (a) D. Seyferth, G. J. Murphy and B. Mauzé, J. Am. Chem. Soc., 99 (1977) 5317.
(b) D. Seyferth, G. J. Murphy and R. A. Woodruff, J. Organometal. Chem., 141 (1977) 71.
11. C. Walling, in "Molecular Rearrangements", F. R. Mayo ed., Interscience, New York, 1963, p. 431.
12. R. West, P. A. Carney and I. C. Minco, J. Am. Chem. Soc., 87 (1965) 3788.
13. (a) D. Seyferth and M. Takamizawa, J. Org. Chem., 28 (1963) 1142.
(b) P. L. K. Hung, M. Sc. Thesis, Massachusetts Institute of Technology, 1973.

TECHNICAL REPORT DISTRIBUTION LIST

<u>No. Copies</u>		<u>No. Copies</u>	
Dr. R. M. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1	Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1
Dr. C. Quicksall Georgetown University Department of Chemistry 37th & O Streets Washington, D.C. 20007	1	Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540	1
Dr. M. F. Hawthorne University of California Department of Chemistry Los Angeles, California 90024	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1		

TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market Street, Rm. 447 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1		
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1		

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Reaction of 3,3-Dichloroallyltrimethylsilane with <u>n</u> -Butyllithium		5. TYPE OF REPORT & PERIOD COVERED Technical Report Interim
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Dietmar Seyferth and Robert E. Mammarella		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0837
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Massachusetts Institute of Technology Cambridge, Mass. 02139		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-618
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217		12. REPORT DATE March 29, 1978
		13. NUMBER OF PAGES 15
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approval for Public Release; Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in the Journal of Organometallic Chemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) organolithium compounds organosilicon compounds		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) n-Butyllithium reacts with 3,3-dichloroallyltrimethylsilane to metalate the vinyl proton. Under the reaction conditions the $\text{Me}_3\text{SiCH}_2\text{Li}=\text{CCl}_2$ formed undergoes β -elimination of LiCl to give $\text{ClC}\equiv\text{CCH}_2\text{SiMe}_3$ whose subsequent reaction with <u>n</u> -butyllithium produces $\text{LiC}\equiv\text{CCH}_2\text{SiMe}_3$. Addition of trimethylchlorosilane gives $\text{Me}_3\text{SiC}\equiv\text{CCH}_2\text{SiMe}_3$. When two molar equivalents of <u>n</u> -butyllithium are used, further metalation of $\text{LiC}\equiv\text{CCH}_2\text{SiMe}_3$ gives $\text{LiC}\equiv\text{CCH}(\text{Li})\text{SiMe}_3$. The action of N-bromosuccinimide on $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CCl}_2$ resulted in formation of $\text{Me}_3\text{SiCH}=\text{CHCCl}_2\text{Br}$.		

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)