

AD-A052 334

AEROSPACE CORP EL SEGUNDO CALIF IVAN A GETTING LABS

F/G 4/1

THE ALTITUDE DEPENDENCE OF THE LOCAL TIME VARIATION OF THERMOSP--ETC(U)

MAR 78 L R SHARP, D R HICKMAN, C J RICE

F04701-77-C-0078

UNCLASSIFIED

TR-0078(3960-04)-3

SANSO-TR-78-67

NL

| OF |

AD  
A052334



END  
DATE  
FILMED  
5-78  
DDC

AD A 052334

# The Altitude Dependence of the Local Time Variation of Thermospheric Density

②  
S

AD No.  
DDC FILE COPY

L. R. SHARP, D. R. HICKMAN, C. J. RICE, and J. M. STRAUS  
Space Sciences Laboratory  
The Ivan A. Getting Laboratories  
The Aerospace Corporation  
El Segundo, Calif. 90245

10 March 1978

Interim Report

APPROVED FOR PUBLIC RELEASE;  
DISTRIBUTION UNLIMITED

Prepared for  
SPACE AND MISSILE SYSTEMS ORGANIZATION  
AIR FORCE SYSTEMS COMMAND  
Los Angeles Air Force Station  
P.O. Box 92960, Worldway Postal Center  
Los Angeles, Calif. 90009

DDC  
RECEIVED  
ADD 3 1978  
F

This interim report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract No. F04701-77-C-0078 with the Space and Missile Systems Organization, Deputy for Advanced Space Programs, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by G. A. Paulikas, Director, Space Sciences Laboratory. Lieutenant Dara Batki, SAMSO/YCPT, was the project officer for Advanced Space Programs.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

Dara Batki

Dara Batki, Lt., USAF  
Project Officer

Robert W. Lindemuth

Robert W. Lindemuth, Lt Col, USAF  
Chief, Technology Plans Division

FOR THE COMMANDER

Leonard E. Baltzell

LEONARD E. BALTZELL, Col, USAF  
Asst. Deputy for Advanced  
Space Programs

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

<b>REPORT DOCUMENTATION PAGE</b>		<b>READ INSTRUCTIONS BEFORE COMPLETING FORM</b>
1. REPORT NUMBER <b>SAMSO TR-78-67<sup>v</sup></b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>THE ALTITUDE DEPENDENCE OF THE LOCAL TIME VARIATION OF THERMOSPHERIC DENSITY.</b>	5. TYPE OF REPORT & PERIOD COVERED <b>Interim rept.</b>	
	6. PERFORMING ORG. REPORT NUMBER <b>TR-0078(3960-04)-3</b>	
7. AUTHOR(S) <b>Lawrence R. Sharp, David R. Hickman, Carl J. Rice, Joe M. Straus</b>	8. CONTRACT OR GRANT NUMBER(S) <b>F04701-77-C-0078<sup>v</sup></b>	
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>The Aerospace Corporation El Segundo, Calif. 90245</b>	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Space and Missile Systems Organization Air Force Systems Command Los Angeles, Calif. 90009</b>	12. REPORT DATE <b>10 Mar 1978</b>	
	13. NUMBER OF PAGES <b>18</b> <span style="margin-left: 20px;"><b>(12) 20 p. 1</b></span>	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) <b>Unclassified</b>	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release; distribution unlimited.</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>Thermospheric Density Local Time Variation Semidiurnal Tides Thermospheric Models Thermospheric Energy Sources</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <b>Analysis of density measurements from the AE-E satellite have demonstrated that the local time variation in the thermosphere at low latitudes (&lt;20°) changes from predominantly diurnal at high altitudes (~300 km) to predominantly semi-diurnal at low altitudes (~140 km). This behavior is significant because it both affects the accuracy of density predictions at low satellite altitudes and confirms that the mesospheric semidiurnal tide is important to the dynamics of the lower thermosphere. Comparison of the results with the local time dependence found in the thermospheric model of Jacchia (1971) and the MSIS model of Hedin et al.</b>		

DD FORM 1473 (FACSIMILE)

409944

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Handwritten initials and scribbles in the bottom right corner.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

19. KEY WORDS (Continued)

20. ABSTRACT (Continued)

(1977 a and b) indicate that only the latter correctly models the transition to dominantly semidiurnal behavior at lower thermospheric altitudes near the equator.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

CONTENTS

INTRODUCTION . . . . . 3

DISCUSSION. . . . . 5

CONCLUSION. . . . . 15

REFERENCES . . . . . 17

FIGURES

1. Average Ratio of the Measured Density to the Jacchia (1971) Model Prediction as a Function of Local Solar Time. . . . . 6

2. Amplitude of the Diurnal and Semidiurnal Components of the Local-Time Variation for the AE-E Data Base Resulting From a Two Component Harmonic Decomposition. . . . . 8

3. Phase of the Diurnal and Semidiurnal Components of the Local-Time Variation Corresponding to the Amplitudes in Fig. 2. . . . . 9

ACCESSION FOR		
NTIS	Section	<input checked="" type="checkbox"/>
DDC	Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL	SPECIAL
A		

## Introduction

Data obtained from high altitude satellites have shown that the atmospheric density has a strong diurnal variation, maximizing near the subsolar point at around 14 LST. For the most part, atmospheric models which have been based principally on data gathered above about 250 km, where the diurnal character is very pronounced, have extrapolated this behavior to lower altitudes though with decreasing amplitude. Some indication that such a straightforward extrapolation may not be correct has been given by the San Marco 3 satellite data presented by Newton et al. (1975). These investigators found at 220 km two distinct peaks of equal amplitude at 11 LST and 18.5 LST. Newton et al. attributed this behavior to the different phase in the diurnal response of individual constituents O and N<sub>2</sub>. That the number density of each atmospheric constituent peaks at a different time of day has also been deduced by Mayr et al. (1974) from analysis of OGO-6 mass spectrometer data.

We have used density data from a cold cathode gauge on board the equatorial satellite Atmosphere Explorer E to investigate the diurnal behavior of the atmosphere from 300 km down to 140 km. The data are used to determine the phase and relative amplitude of the diurnal and semidiurnal components of the local time variation as a function of altitude. These parameters are valuable as an indication of the relative magnitude of energy sources for the lower thermosphere since a diurnal component may be expected to result from local heating while the semidiurnal component would be driven by tides propagating upwards from the stratosphere and mesosphere (Lindzen, 1976). Assignment of a

source would also depend, of course, on such details as phase and phase progression and the altitude-dependence of the wave amplitude.



## Discussion

The data used in this study consist of 5062 density measurements obtained using the cold cathode ion gauge known as Pressure Sensor A (PSA) on board the Atmosphere Explorer E (AE-E) satellite. The instrument has been described by Rice et al. (1973), and the data reduction procedure has been presented by Rice and Sharp (1977). AE-E is in a near-equatorial orbit, with an inclination of  $19.6^\circ$ . The data used in this study were gathered between December 1975 and October 1976 when AE-E was in an eccentric orbit (140 km  $\times$  2757 km). The data set was further limited by including only data obtained during periods of oriented (non-spinning) operation. Density and ephemeris data at fixed altitude levels at 5 km intervals (140-300 km) were tabulated for use in the present study. In order to minimize the complicating and incompletely modeled effects of geomagnetic activity, the data base excludes any data for which  $K_p \geq 5$  at any time during the period six hours prior to the time of measurement.

In order to isolate the local time variation in the lower thermosphere, all the other known contributions to the variability of the density were suppressed using the Jacchia (1971) model, evaluated using all appropriate input parameters, but with the local time set arbitrarily at midnight. The ratio R of the measured density to this model value was then averaged over each two-hour interval of local time. The results of this analysis are shown in Figure 1. Each curve represents the average of this ratio over the five altitude levels in a 20 km band centered at the designated altitude. (For example, the 290 km




Fig. 1. Average ratio of the measured density to the Jacchia (1971) model prediction as a function of local solar time. The local time parameter for the model is taken to be local midnight in all cases, but all parameters are handled in the manner prescribed in the model formulation. The data are binned at two-hour intervals. Each curve includes all altitude increments within the 20 km band centered on the designated altitude.

curve includes data from the 280-300 km range.) The curves thus represent the qualitative altitude dependence of the local time variation which occurs near the equator. There is a systematic transition from the clearly diurnal behavior at 290 km with a well-defined maximum near 14 hours local time and a broader minimum near midnight to the semidiurnal behavior at 140 km. In the transition region between these altitudes the daily variation is somewhat less well-defined due to the fact that neither the diurnal nor the semidiurnal mode is clearly dominant.

In order to study the altitude dependence of the daily density variation in greater detail, we have performed a two-component harmonic analysis of the ratio  $R$  after dividing by the mean value. The amplitude and phase of the diurnal and semidiurnal terms were calculated for each 5 km altitude step in the 140-300 km data base. Initially a three-component fit was made, but the terdiurnal component showed no phase continuity with altitude, and the amplitude at all points was too small to be statistically significant. Furthermore, neglect of the third harmonic in later analyses did not significantly affect either the amplitude or phase of the resultant diurnal and semidiurnal terms. A somewhat similar analysis using data from the San Marco 3 satellite at higher altitudes has been described by Broglia et al. (1976).

In Figures 2 and 3 we present the amplitude and phase, respectively, determined in each of these decompositions. The amplitudes and phases of the two modes are shown as functions of altitude, along with the corresponding predictions of two widely used empirical models of the thermosphere, the Jacchia (1971) model and the Mass Spectrometer-Incoherent Scatter (MSIS)




Fig. 2. Amplitude of the diurnal and semi-diurnal components of the local-time variation for the AE-E data base resulting from a two component harmonic decomposition. The error bars present a statistical measure of the goodness of fit at each altitude; for clarity they are only given every other altitude increment. Also plotted are the results of performing a similar decomposition on the density values calculated by means of the Jacchia (1971) and MSIS thermospheric models.




Fig. 3. Phase of the diurnal and semi-diurnal components of the local-time variation corresponding to the amplitudes in Fig. 2.

model of Hedin et al. (1977). In calculating the amplitude and phase predictions of the empirical models, model densities are determined for every point in our AE-E data set and the resultant values are then handled in the same manner as with the actual measured data.

Comparison of the amplitudes derived from the data with those of the empirical models leads to several conclusions. For both the data and the models, the amplitude of the diurnal mode decreases rapidly with decreasing altitude. However, at any given altitude the amplitude of the diurnal model predicted by the Jacchia (1971) model exceeds that actually measured by more than a factor of two. The altitude dependence of the amplitude of the MSIS diurnal mode closely resembles that of the AE-E data. Although the amplitude is uniformly less than the measured values, it is only slightly beyond the uncertainty limits for the fits. The amplitudes of the semidiurnal mode calculated from the data and both models are comparable at the higher altitudes treated here ( $\geq 240$  km). However, this overall agreement breaks down below  $\sim 200$  km, where the data and the MSIS model indicate an increase with decreasing altitude, whereas the semidiurnal mode in the Jacchia (1971) model continues to decrease monotonically. The measured semidiurnal mode surpasses the diurnal mode in amplitude at  $\sim 180$  km and is dominant at all lower altitudes. The essentially identical behavior of the semidiurnal mode in the MSIS model and in our AE-E data base extends to such details as the altitude at which minimum amplitude is seen ( $\sim 200$  km). The agreement of model and data is even more striking in the phase plot, Figure 3. The phase of the semidiurnal mode, which at high altitudes is almost constant between 2 and 3 hours

LST, makes near 200 km a nearly discontinuous transition to a phase angle between 8 and 10 hours LST. The Jacchia (1971) model extrapolates the behavior at higher altitudes downward to the lower thermosphere as well.

The measured phase of the diurnal mode shows good agreement with the models at high altitudes, but considerable disagreement at low altitudes. The phase in the measured data progresses from near 13 hours LST at 300 km toward earlier times, reaching 6 hours LST at 140 km. The phase of the MSIS model, on the other hand, though starting at a similar point at high altitude, progresses toward later times, reaching 2.5 hours LST at 140 km. The Jacchia (1971) model shows a diurnal phase which remains constant at 14 hours LST. This divergence can probably be attributed to the manner in which the data used in constructing the models was obtained. The data base for the Jacchia (1971) model contained no data below an altitude of  $\sim 260$  km; the constant phase results from extrapolating downwards the behavior of the upper thermosphere and, hence, good agreement in the lower thermosphere might not be expected. On the other hand, the low altitude data input to the MSIS model is almost exclusively from the Atmosphere Explorer C satellite. We believe that the behavior of the phase of the diurnal mode in the MSIS model may be an artifact arising from the strong coupling in the AE-C data base of local time with other parameters such as latitude and season. Because the amplitude of the diurnal mode is small at low thermospheric altitudes, such behavior has little effect on the density there. To test this hypothesis we performed the above analysis using our own ion gauge data from the AE-C

satellite and obtained results which were qualitatively similar to the MSIS predictions with respect to the diurnal phase progression at low altitude.

The MSIS model semidiurnal mode agrees quite well in phase with the data showing the sharp phase transition associated with the change from EUV to tidal domination below  $\sim 200$  km. The Jacchia (1971) model fails to reproduce this behavior and disagrees substantially in phase at the lower altitudes studied here. This suggests that the MSIS model is more reliable for studies involving lower thermospheric chemistry and dynamics than is the Jacchia (1971) model. In this connection it should be noted that the MSIS model exhibits the semidiurnal mode as dominant only at low latitude; at 140 km the diurnal mode is dominant above about  $30^\circ$  latitude. Our data from AE-E cannot confirm or deny this behavior, however.

The general behavior of a dominant diurnal mode at high altitudes and semidiurnal mode at low altitudes is consistent with the following picture. At high altitudes the characteristics of thermosphere are determined largely by the day-night variations in EUV heating, leading to a diurnal variation whose amplitude decreases with decreasing altitude because the heating rate per unit mass decreases with decreasing altitude. The behavior at low thermosphere altitudes arises from the propagation of tidal waves from lower regions of the atmosphere. These tides are known to be predominantly semidiurnal in character (Lindzen et al. 1976; Wand, 1976) and their amplitudes decrease with increasing altitude in the thermosphere.

Although comparison of phase in density data with that in temperature must be approached with a good deal of caution, it is worth pointing out the



qualitative similarity of the altitude dependence of the density phase measurements reported here and the temperature phase calculated for the solar semidiurnal tide by Lindzen and Hong (1974) for equinoctial conditions near the equator, including the nearly  $180^\circ$  phase change seen near 200 km and the more gradual shift toward later local time evident below 160 km. The calculations of temperature phase by Lindzen and Hong (their Figure 20) show this behavior uniformly 40-50 km below the altitudes at which it is seen in our density measurements. On the other hand, the calculations for solstitial conditions show much more extreme phase changes with altitude ( $720^\circ$  over the same altitude band, 125-175 km), and since our data base encompasses solstitial and equinoctial conditions approximately equally, the previous noted qualitative agreement may be only fortuitous. Given the general character of the altitude dependence of the semidiurnal mode, it seems to us reasonable to ascribe this behavior to the solar semidiurnal tide. With this interpretation, the transition between the part of the thermosphere dominated by tides and the EUV-dominated upper thermosphere occurs near the equator in the 200-220 km altitude range.

### Conclusion

The Atmosphere Explorer E density data suggest that the upward propagating mesospheric tidal wave dominates the local time behavior near the equator in the lower thermosphere up to about 180 km and is important to at least 200 km. Thus the common assumption of a predominantly diurnal density variation below  $\sim 200$  km may lead to serious errors in density predictions and cause errors in any calculation attempting to determine the dynamical behavior of the atmosphere below 200 km. Of those atmospheric models known to us, only the MSIS model of Hedin et al. (1977) reproduces the measured data.

PRECEDING PAGE BLANK-NOT FILMED

## References

- Broglio, L, C. Arduini, C. Buongiorno, U. Ponzi and G. Ravelli,  
San Marco 3 drag balance results below 320 km altitude in the equatorial  
atmosphere, Space Research XVI, Akademie-Verlag, Berlin, 1976.
- Hedin, A. E., J. E. Salah, J. V. Evans, C. A. Reber, G. P. Newton,  
N. W. Spencer, D. C. Kayser, D. Alcaydé, P. Bauer, L. Cogger and  
J. P. McClure, A global thermospheric model based on mass spectrom-  
eter and incoherent scatter data MSIS 1.  $N_2$  density and temperature,  
J. Geophys. Res., 82, 2139, 1977a.
- Hedin, A. E., C. A. Reber, G. P. Newton, N. W. Spencer, H. C. Brinton,  
H. G. Mayr and W. E. Potter, A global thermospheric model based on  
mass spectrometer and incoherent scatter data MSIS 2. composition,  
J. Geophys. Res., 82, 2148, 1977b.
- Jacchia, L. G., Revised static models of the thermosphere and exosphere  
with empirical temperature profiles, Smithsonian Astrophys. Obs. Spec.  
Rep. No. 332, 1971.
- Lindzen, R. S., A model decomposition of the semidiurnal tide in the lower  
thermosphere, J. Geophys. Res., 81, 2923, 1976.
- Lindzen, R. S. and S.-S. Hong, Effects of mean winds and horizontal tempera-  
ture gradients on solar and lunar semidiurnal tides in the atmosphere,  
J. Atmos. Sci., 31, 1421, 1974.

PRECEDING PAGE BLANK-NOT FILMED

- Mayr, H. G., A. E. Hedin, C. A. Reber and G. R. Carignan, Global characteristic in the diurnal variations of thermospheric temperature and composition, J. Geophys. Res., 79, 619, 1974.
- Newton, G. P., W. T. Kasprzak, S. A. Curtis and D. J. Pelz, Local time variation of equatorial thermospheric composition determined by the San Marco 3 Nace, J. Geophys. Res., 80, 2289, 1975.
- Rice, C. J., V. L. Carter, S. R. LaValle, W. T. Chater, D. A. Jones, C. G. King and D. F. Nelson, Atmosphere explorer pressure measurements: ion gauge and capacitance manometer, Radio Sci., 8, 305, 1973.
- Rice, C. J. and L. R. Sharp, Neutral atmospheric waves in the thermosphere and tropospheric weather systems, Geophys. Res. Let., 4, 315, 1977.
- Wand, R. H., Semidiurnal tide in the E region from incoherent scatter measurements at Arecibo, Radio Sci., 11, 641, 1976.

## THE IVAN A. GETTING LABORATORIES

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas lasers.

Chemistry and Physics Laboratory: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocket plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, lubrication and surface phenomena, photosensitive materials and sensors, high precision laser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semiconducting, superconducting, and crystal device physics, optical and acoustical imaging; atmospheric pollution; millimeter wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields; space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

THE AEROSPACE CORPORATION  
El Segundo, California

16