

AD-A052 590

BOJARSKI (NORBERT N) NEWPORT BEACH CA
A WAVE EQUATION FOR RADIATING SOURCE DISTRIBUTIONS. (U)
FEB 78 N N BOJARSKI

F/G 20/3

N00014-76-C-0082

NL

UNCLASSIFIED


| OF |
AD
A052 590



END
DATE
FILMED
5-78

DDC





AD A 052590

AD No. ~~AD A 052590~~
DDC FILE COPY

9

12

A WAVE EQUATION FOR RADIATING
SOURCE DISTRIBUTIONS

DR. NORBERT N. BOJARSKI

Research Contractor and Consultant
to the DEPARTMENT OF DEFENSE

Sixteen Pine Valley Lane
Newport Beach, California 92660
Telephone: (714) 640 - 7900

DDC
RECEIVED
APR 4 1978
F

February 1978

Prepared for and supported by
The Office of Naval Research
800 North Quincy Street
Arlington, Virginia 22217

see 1473

This document has been approved
for public release and sale; its
distribution is unlimited.

ABSTRACT

The Bleistein-Cohen separation of a source (for the inhomogeneous Helmholtz wave equation) distribution into radiating and non-radiating portions is reformulated into a form suitable for deriving a wave equation governing the radiating portion of the source distribution.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	B II Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
J S I 105 171	
BY	
DISTRIBUTION/AVAILABILITY GROUPS	
SP. CEN.	
A	

TABLE OF CONTENTS

<u>SECTION</u>	<u>TITLE</u>	<u>PAGE</u>
I	INTRODUCTION	2
II	AN ALTERNATIVE FORMULATION OF THE BLEISTEIN-COHEN EQUATIONS	3
III	A WAVE EQUATION FOR RADIATING SOURCE DISTRIBUTIONS	6
IV	CONCLUDING REMARKS	9
V	REFERENCES	10
	DD Form 1473: Report Documentation Page	11

SECTION I

INTRODUCTION

Bleistein and Cohen have shown [1,2] that a source distribution, giving rise to fields that satisfy the inhomogeneous Helmholtz wave equation (for acoustics, electromagnetics, etc.), can be separated into radiating and non-radiating portions; and that the radiating source distribution can be determined completely and uniquely in a well-conditioned manner from knowledge of the radiated fields on any closed surface exterior to the support of the source distribution.

An alternative formulation of this separation is developed in a form suitable for deriving a wave equation which governs the radiating portion of the source distribution.

SECTION II

AN ALTERNATIVE FORMULATION OF THE BLEISTEIN-COHEN EQUATIONS

Let $Y_{mn}(\theta, \phi)$ and $j_l(kr)$ be the normalized over $\theta \in (0, \pi)$ and $\phi \in (0, 2\pi)$ spherical harmonic and the normalized over $r \in (0, \infty)$ spherical Bessel functions respectively. It thus follows that the functions $Y_{mn}(\theta, \phi) j_l(kr)$ are orthonormal and complete in all space [3], and that any function $\rho(\theta, \phi, r)$ can be expanded as

$$\rho(\theta, \phi, r) = \sum_{n=0}^{\infty} \sum_{m=-n}^n \sum_{l=0}^{\infty} C_{mnl} Y_{mn}(\theta, \phi) j_l(kr) \quad , \quad (1)$$

where the coefficients C_{mnl} are given by

$$C_{mnl} = \int_0^{\infty} \int_0^{2\pi} \int_0^{\pi} Y_{mn}(\theta, \phi) j_l(kr) \rho(\theta, \phi, r) \sin \theta \, d\theta \, d\phi \, r^2 \, dr \quad . \quad (2)$$

If the function ρ represents a source distribution, giving rise to fields ψ , which satisfy the inhomogeneous reduced Helmholtz wave equation

$$\nabla^2 \psi + k^2 \psi = -\rho \quad , \quad (3)$$

and the support of these sources ρ is confined to the interior of a sphere of radius a , then the radiated fields ψ at the exterior of this sphere are

$$\psi(\theta, \phi, r) = ik \sum_{n=C}^{\infty} \sum_{m=-n}^n Y_{mn}^*(\theta, \phi) h_n^{(1)}(kr) \int_0^a \int_0^{2\pi} \int_0^{\pi} Y_{mn}(\theta', \phi') j_n(kr') \times \rho(\theta', \phi', r') \sin \theta' \, d\theta' \, d\phi' \, r'^2 \, dr' \quad , \quad r > a \quad , \quad (4)$$

which by (2) can be written as

$$\psi(\theta, \phi, r) = ik \sum_{n=0}^{\infty} \sum_{m=-n}^n C_{mnn} Y_{mn}^*(\theta, \phi) h_n^{(o)}(kr) ; \quad (5)$$

i.e., only the diagonal coefficients C_{mnn} (of the complete set of coefficients C_{mnl}) of the expansion (1,2) give rise to radiated fields.

If the functions G_{mnl} are defined as

$$G_{mnl} \equiv Y_{mn}(\theta, \phi) j_l(kr) , \quad (6)$$

then by the general solution in spherical coordinates of the homogeneous reduced Helmholtz wave equation,

$$\nabla^2 G_{mnn} + k^2 G_{mnn} = 0 . \quad (7)$$

Thus, by Green's theorem, (3), and (2)

$$\begin{aligned} \oint_S dS \cdot (G_{mnn} \nabla \psi - \psi \nabla G_{mnn}) &= \int_V dv (G_{mnn} \nabla^2 \psi - \psi \nabla^2 G_{mnn}) \\ &= \int_V dv G_{mnn} \rho \\ &= C_{mnn} ; \end{aligned} \quad (8)$$

i.e., the diagonal coefficients, which completely and uniquely determine the radiating portion of the source distribution, can be determined in a

well-conditioned manner from knowledge of the field ψ on any closed surface s , exterior to the support of the source distribution ρ .

SECTION III

A WAVE EQUATION FOR RADIATING SOURCE DISTRIBUTIONS

The expansion (1) of an arbitrary source distribution $\rho(\theta, \phi, r)$ can be broken into the diagonal and non-diagonal parts

$$\begin{aligned} \rho(\theta, \phi, r) = & \sum_{n=0}^{\infty} \sum_{m=-n}^n \sum_{\substack{l=0 \\ l \neq n}}^{\infty} C_{mnl} Y_{mn}(\theta, \phi) j_l(kr) \\ & + \sum_{n=0}^{\infty} \sum_{m=-n}^n C_{mnn} Y_{mn}(\theta, \phi) j_n(kr) \quad , \end{aligned} \quad (9)$$

which can be identified respectively as the non-radiating (ρ_0) and radiating (ρ_∞) portion of the source distribution; i.e.,

$$\rho(\theta, \phi, r) = \rho_0(\theta, \phi, r) + \rho_\infty(\theta, \phi, r) \quad , \quad (10)$$

where

$$\rho_0(\theta, \phi, r) = \sum_{n=0}^{\infty} \sum_{m=-n}^n \sum_{\substack{l=0 \\ l \neq n}}^{\infty} C_{mnl} Y_{mn}(\theta, \phi) j_l(kr) \quad , \quad (11)$$

$$\rho_\infty(\theta, \phi, r) = \sum_{n=0}^{\infty} \sum_{m=-n}^n C_{mnn} Y_{mn}(\theta, \phi) j_n(kr) \quad . \quad (12)$$

By the general solution in spherical coordinates of the homogeneous reduced Helmholtz wave equation, it follows from (12) that

$$\nabla^2 \rho_\omega + k^2 \rho_\omega = 0 \quad . \quad (13)$$

An alternative, and much simpler, coordinate independent derivation of (13) for the far-field radiating source distribution can be accomplished in the following fashion.

The three-dimensional (spatial) Fourier transform of any source distribution

$$\tilde{\rho}(k, \omega) = \int e^{i\mathbf{k} \cdot \mathbf{x}} \rho(\mathbf{x}, \omega) d^3x \quad , \quad (14)$$

the support of which, in general, is infinite in k-space, can always be separated into the two functions $\tilde{\rho}_o(k, \omega)$ and $\tilde{\rho}_\infty(k, \omega)$, the support of which is off and on the surface of the k-space sphere of radius $k = \frac{\omega}{c}$ respectively; i.e.,

$$\tilde{\rho}_o(k, \omega) = \begin{cases} 0 & , \quad \mathbf{V} k = \frac{\omega}{c} \\ \tilde{\rho}(k, \omega) & , \quad \mathbf{V} k \neq \frac{\omega}{c} \end{cases} \quad , \quad (15)$$

$$\tilde{\rho}_\infty(k, \omega) = \begin{cases} \tilde{\rho}(k, \omega) & , \quad \mathbf{V} k = \frac{\omega}{c} \\ 0 & , \quad \mathbf{V} k \neq \frac{\omega}{c} \end{cases} \quad , \quad (16)$$

$$\tilde{\rho}(k, \omega) = \tilde{\rho}_o(k, \omega) + \tilde{\rho}_\infty(k, \omega) \quad . \quad (17)$$

However, by (16), $\tilde{\rho}_\infty(k, \omega)$ is the complete set of the (range and phase normalized) radiated far-field of the source distribution $\rho(\mathbf{x}, \omega)$, and by (15), $\tilde{\rho}_o(k, \omega)$ contributes nothing to the radiated far-field. it thus follows that

$$\rho_o(\mathbf{x}, \omega) = \frac{1}{(2\pi)^3} \int_{\mathbf{V} \ k \neq \frac{\omega}{c}} e^{-i\mathbf{k} \cdot \mathbf{x}} \rho_o(\mathbf{k}, \omega) d^3k \quad (18)$$

is the non far-field radiating portion of the source distribution, and

$$\rho_\omega(\mathbf{x}, \omega) = \frac{1}{(2\pi)^3} \int_{\mathbf{V} \ k = \frac{\omega}{c}} e^{-i\mathbf{k} \cdot \mathbf{x}} \rho_\omega(\mathbf{k}, \omega) d^3k \quad (19)$$

is the far-field radiating portion of the source distribution; where

$$\rho(\mathbf{x}, \omega) = \rho_o(\mathbf{x}, \omega) + \rho_\omega(\mathbf{x}, \omega) \quad (20)$$

It follows directly from (19) that

$$\nabla^2 \rho_\omega + k^2 \rho_\omega = 0 \quad , \quad k = \frac{\omega}{c} \quad (21)$$

As suggested by Bleistein [4], expansion of the integrand term $e^{-i\mathbf{k} \cdot \mathbf{x}}$ in (14), (18), and (19) in spherical harmonics reveals that the radiating and non-radiating source distributions of this alternative derivation are indeed the same radiating and non-radiating source distributions given by (11) and (12) respectively.

It can similarly be shown that a vector current density distribution \mathbf{J} , which gives rise to electromagnetic fields that satisfy Maxwell's equations, can similarly be separated into radiating and non-radiating current density distributions, and that the radiating current density distribution \mathbf{J}_ω satisfies the vector wave equation

$$\nabla \times \nabla \times \mathbf{J}_\omega - k^2 \mathbf{J}_\omega = 0 \quad (22)$$

SECTION IV

CONCLUDING REMARKS

Source distributions that fail to satisfy the wave equation should be carefully excluded from solutions to the inverse source, inverse scattering, and synthesis problem. Such an exclusion might advantageously be achieved by viewing this wave equation as an additional necessary condition (e.g., facilitating regularization of ill-conditioned and/or ill-posed problems) on the solutions to the inverse source, inverse scattering, and synthesis problem.

SECTION V

REFERENCES

- [1] Bleistein, N., and Cohen, J. K., *Non-uniqueness in the inverse source problem in acoustics and electromagnetics*, J. Math. Physics, 18, 2, 194-201 (1977).
- [2] Bleistein, N., and Cohen, J. K., *A note on the inverse source problem*, J. Math. Physics, (to be published).
- [3] Watson, G. N., *A Treatise on the Theory of Bessel Functions*, Sect. 16.4., p. 533, Cambridge University Press, 1962.
- [4] Bleistein, N., Personal communication.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1 REPORT NUMBER	2 GOVT ACCESSION NO.	3 RECIPIENT'S CATALOG NUMBER
4 TITLE (and Subtitle) 6 A WAVE EQUATION FOR RADIATING SOURCE DISTRIBUTIONS.		5 TYPE OF REPORT & PERIOD COVERED 9 SCIENTIFIC rept.,
7 AUTHOR(s) 10 Norbert N. Bojarski		6 PERFORMING ORG. REPORT NUMBER
9 PERFORMING ORGANIZATION NAME AND ADDRESS Dr. Norbert N. Bojarski 16 Pine Valley Lane Newport Beach, California 92660		8 CONTRACT OR GRANT NUMBER(s) 15 N00014-76-C-0082 <i>new</i>
11 CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, Virginia 22217		10 PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14 MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12 REPORT DATE 11 Feb 1978 1978
16 DISTRIBUTION STATEMENT (of this Report) UNLIMITED		13 NUMBER OF PAGES 12 13 P. 10
17 DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15 SECURITY CLASS. (of this report) UNCLASSIFIED
18 SUPPLEMENTARY NOTES		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
19 KEY WORDS (Continue on reverse side if necessary and identify by block number) Inverse Source Inverse Scattering		
20 ABSTRACT (Continue on reverse side if necessary and identify by block number) The Bleistein-Cohen separation of a source (for the inhomogeneous Helmholtz wave equation) distribution into radiating and non-radiating portions is reformulated into a form suitable for deriving a wave equation governing the radiating portion of the source distribution.		

DISTRIBUTION STATEMENT A
 Approved for public release;
 Distribution Unlimited

389 425

ED
78