AD-AC	52 659	DEC	AL POST UCTURE 77 D	GRADUA OF SYN J BEPR	TE SCH	OOL MON	NTEREY NAVES I	CALIF N THE	TROPICA	L PACI	FIC DU	F/G 4/2 RINET	c(U)
	OF AD 52659	- 27 un parter Services							A Second	James - Andream Sterner and Minister and Angel Minister and Ang			
							And a second sec		A Description of the second se				Ensjelle Belgele Etzgele
English English English				South States						The second secon	Nortes at the second se	FILMED 5 -78	
													×
				16aC									. /



Approved for public release; distribution unlimited.

251 450

JOB

. REPORT NUMBER 2. GOVT ACCESSION NG	BEFORE COMPLETING FORM
. TITLE (and Subtitio)	3. RECIPIENT'S CATALOG NUMBER
. TITLE (and Subtitle)	
	S. TYPE OF REPORT & PERIOD COVERED
Structure of Synoptic-Scale Waves in	Master's Inesis;
the Tropical Pacific During July-	December 1977
December 19/4-19/6	. PERFORMING ORG. REPORT NUMBER
. AUTHOR(a)	S. CONTRACT OR GRANT NUMBER(+)
Donald Joseph Bepristis	
	10. PROGRAM EL EMENT PROJECT TASK
Naval Postgnaduate School	AREA & WORK UNIT NUMBERS
Monterey, California 93940	
1. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
Naval Postgraduate School Monterey, California 93940	13. NUMBER OF PAGES
Neurol Destandante Calas	Upolacoified
Monterev. California 93940	Unclassified
	ISA. DECLASSIFICATION/DOWNGRADING
SUPPLEMENTARY NOTES	
). KEY WORDS (Continue on reverse eide if necessary and identify by black number	0
. ABSTRACT (Continue on revorce olde il necessory and identify by block manber)	
. ABSTRACT (Continue on revorce olde 11 necessary and identify by block member) Composite wave structures, sea-surfa	ce temperatures (SST),
Composite wave structures, sea-surfa and mean zonal winds are studied for the	ce temperatures (SST), period July through
Composite wave structures, sea-surfa and mean zonal winds are studied for the December over the three consecutive year	ce temperatures (SST), period July through s 1974, 1975 and 1976.
Composite wave structures, sea-surfa and mean zonal winds are studied for the December over the three consecutive year These three years were found to possess	ce temperatures (SST), period July through s 1974, 1975 and 1976. an "average" SST anom-
ABITRACT (Continue on reverse olde 11 recessory and identify by block member) Composite wave structures, sea-surfa and mean zonal winds are studied for the December over the three consecutive year These three years were found to possess aly distribution, a cold anomaly pattern	ce temperatures (SST), period July through s 1974, 1975 and 1976. an "average" SST anom- and a warm anomaly
Composite wave structures, sea-surfa and mean zonal winds are studied for the December over the three consecutive year These three years were found to possess aly distribution, a cold anomaly pattern pattern respectively. Radiosonde data f for each year and each of the six locati	ce temperatures (SST), period July through s 1974, 1975 and 1976. an "average" SST anom- and a warm anomaly or the six month period ons (Koror, Yap, Truk.
ABITRACT (Continue on revorce olds if necessary and identify by block manber) Composite wave structures, sea-surfa and mean zonal winds are studied for the December over the three consecutive year These three years were found to possess aly distribution, a cold anomaly pattern pattern respectively. Radiosonde data f for each year and each of the six locati	ce temperatures (SST), period July through s 1974, 1975 and 1976. an "average" SST anom- and a warm anomaly or the six month period ons (Koror, Yap, Truk,
ABSTRACT (Continue on reverse side if receivery and identify by block member) Composite wave structures, sea-surfa and mean zonal winds are studied for the December over the three consecutive year These three years were found to possess aly distribution, a cold anomaly pattern pattern respectively. Radiosonde data f for each year and each of the six locati	ce temperatures (SST), period July through s 1974, 1975 and 1976. an "average" SST anom- and a warm anomaly or the six month period ons (Koror, Yap, Truk,



Ponape, Kwajalein and Majuro) were analyzed to derive composite wave descriptions of meridional and zonal wind components, temperature and specific humidity. The basic results such as periodicity and structural tilt were generally in agreement with previous research. Combining the SST data, mean zonal flow and wave composites leads to some new insights into the effects of the tropical SST on the semi-permanent Walker circulation and transient easterly waves. Results indicate that the east to west oscillation of the Walker circulation may be directly related to the tropical eastern Pacific SST. It is also shown that the migration of the Walker circulation and subsequent change in the upper-tropospheric flow is directly related to the structural tilt of migratory tropical waves. Consideration of the wave thermal structure and its relation to the SST indicates that the waves energetics may reflect the effects of the upstream SST. ACCESSION for White Section NTIS Buff Section DDC UNANNOUNCED JUSTIFICATION BY DISTRIBUTION/AVAILABILITY CODES Dist. AVAIL. and/or SPECIAL

DD Form 1473 1 Jan 73 5/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE(Then Date Entered)

Approved for public release; distribution unlimited.

Structure of Synoptic-Scale Waves in the Tropical Pacific During July-December 1974-1976

by

Donald Joseph Bepristis Lieutenant, United States Navy B.A., San Jose State University, 1972

Submitted in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE IN METEOROLOGY AND OCEANOGRAPHY

from the NAVAL POSTGRADUATE SCHOOL December 1977

Author LAN Approved by: Thesis Advisor Second Reader Department of Meteorology Chai an Engineering Dean ience and

ABSTRACT

Composite wave structures, sea-surface temperatures (SST), and mean zonal winds are studied for the period July through December over the three consecutive years 1974, 1975 and 1976. These three years were found to possess an "average" SST anomaly distribution, a cold anomaly pattern and a warm anomaly pattern respectively. Radiosonde data for the six month period for each year and each of the six locations (Koror, Yap, Truk, Ponape, Kwajalein and Majuro) were analyzed to derive composite wave descriptions of meridional and zonal wind components, temperature and specific humidity. The basic results such as periodicity and structural tilt were generally in agreement with previous research. Combining the SST data, mean zonal flow and wave composites leads to some new insights into the effects of the tropical SST on the semi-permanent Walker circulation and transient easterly waves. Results indicate that the east to west oscillation of the Walker circulation may be directly related to the tropical eastern Pacific SST. It is also shown that the migration of the Walker circulation and subsequent change in the upper-tropospheric flow is directly related to the structural tilt of migratory tropical waves. Consideration of the wave thermal structure and its relation to the SST indicates that the waves energetics may reflect the effects of the upstream SST.

TABLE OF CONTENTS

I.	INTR	ODU	CTIO	N 9	9
II.	DATA			12	2
	Α.	SEA-	-SUR	FACE TEMPERATURE 12	2
	в.	RAD	IOSO	NDE DATA	2
III.	RESU	ILTS	-	14	ł
	Α.	TIM	E ME.	AN FIELDS 14	ł
		1.	Sea	-Surface Temperature 14	ł
	·	2.	Mea	n Zonal Wind 15	5
	в.	TIM	E AN	D ZONAL SCALES OF THE WAVES 16	5
	с.	COM	POSI	TES OF WAVE STRUCTURE 16	5
		1.	197	4 17	7
			a.	Wind Components 17	7
			ь.	Temperature and Specific Humidity 17	,
		2.	197	5 18	3
			a.	u and v Components 18	3
			ь.	Temperature and Specific Humidity 18	3
		з.	197	6 19)
			a.	u and v Components 19)
			ь.	Temperature and Specific Humidity 20)
IV.	DISC	USS	ION	AND CONCLUSIONS 21	
LIST	OF RE	FERI	ENCE	S 36	;
TNTTT		STP	TRIT	TON LIST 3	,

LIST OF TABLES

I.	Time	and	zo	nal	. :	sca	110	es	01	f	the	е	way	ves	5 3	foi	2					
	1974-	-1978	5 -	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24

LIST OF FIGURES

1.	Sea-surface temperature anomalies for 1974, 1975 and 1976 taken from 1972- 1976 mean 2	25
2.	Sea-surface temperature analyses for 1974 through 1976 and 1972-1976 mean 2	26
3.	Radiosonde stations used in study 2	27
4.	Mean zonal winds for 1972-1976 2	28
5.	Composite vertical wave structure of u and v component winds during 1974 for all stations 2	29
6.	Composite vertical wave structure of temperature and specific humidities during 1974 for all stations 3	30
7.	Same as Figure 5 except for 1975 3	31
з.	Same as Figure 6 except for 1975 3	32
9.	Same as Figure 5 except for 1976 3	33
10.	Same as Figure 6 except for 1976 3	34
11.	 (a) Plot of 150 mb wind vs. 1000-500 mb vertical phase shift for 1972-1976. (b) Plot of average trough temperature vs. average 10° upstream SST	35

ACKNOWLEDGEMENT

The author wishes to express his appreciation and thanks for the invaluable assistance given by Prof. C.-P. Chang, without whose assistance this project could never have been completed, and to Dr. D. McCline of the Pacific Environmental group, National Marine Fisheries Services/NOAA, for his help in supplying and processing the sea-surface temperature data. Thanks are expressed also to Prof. R. T. Williams for reading the manuscript. The Radiosonde data were provided by the Director, Naval Oceanography and Meteorology, from the archives of the National Climatic Records Center, Asheville, N. C. This research was supported in part by NESS/ NOAA under contract 7-11030.

I. INTRODUCTION

Various researchers in the past have found evidence that variations in the tropical sea-surface temperature (SST) are associated with changes in the planetary-scale circulation over the tropics (Bjerknes, 1969; Winston and Kruger, 1974, and Namias, 1974). Recently several investigators have concentrated on the years 1972 and 1973 which displayed extremely anomalous and contrasting SST variations in the equatorial eastern Pacific. The 1972 anomaly was the warmest thus far recorded, with the last six month's mean temperature in excess of 2°C above the normal over broad areas. In 1973 the SST pattern was almost completely reversed with the eastern Pacific characterized by anomalies of more than 2°C below normal for the last six month's mean. It was found by Winston and Kruger (1974) that the anomalously warm eastern Pacific SST of 1972 was associated with major changes in the tropical and extratropical planetary scale circulations.

Recently Chang and Miller (1976), Maas (1977) and Delaney (1977) have found that this change in the planetary scale was accompanied by variations in the synoptic-scale tropical disturbances. Delaney's (1977) study involves a cloud cover analysis for the tropical western Pacific during summer and fall of 1972 and 1973. His results indicate that the synoptic-scale waves present during those periods were influenced both by the east-west overturning planetary scale Walker

circulation and by the local effects of SST. Other investigators have also found that variations in tropical wave behavior could be explained in part by the variations of SST and planetary scale flows. According to Chang and Miller's (1976) study of the 1972 and 1973 easterly waves in the tropical Pacific, the differences in vertical wave structure are a result of seasonal variations in the vertical shear of the mean zonal winds, which are directly linked to the east-west SST gradient and the resultant zonal Walker circulation. It was further indicated by Delaney's results that the local wave amplitudes and energetics are strongly dependent on the local variations of the SST.

The purpose of this study is to inquire further into the role of anomalous SST in affecting the synoptic-scale tropical waves. As a first step, the last six months of each year's SST were averaged and deviations from the 1972 to 1976 five-year mean were calculated (Figure 1). These charts cover the entire equatorial Pacific from $10^{\circ}-20^{\circ}N$ for the years 1972 through 1976. The 1972 and 1973 characteristics have already been described by Delaney (1977). The 1975 and 1976 anomaly charts have distinctly different patterns, while the 1974 anomaly pattern is what might be described as "average." The 1975 and 1976 anomalies, although not as intense as 1972 and 1973, are significant in that almost the entire tropical Pacific is covered by cold anomaly in 1975 and warm anomaly in 1976. By analyzing the

tropical synoptic and mean zonal features and their variations with the SST anomalies, it is hoped that more light may be shed on the inter-relationships between them.

II. DATA

A. SEA-SURFACE TEMPERATURE

The SST data used in this study and the earlier study by Delaney (1977) were produced from special monthly SST data charts compiled by the Pacific Environmental Group, National Marine Fisheries Service, for the years 1972 through 1976. These charts contained the average monthly SST for five degree blocks over the entire equatorial and northern Pacific. The data coverage was very good for all areas except the Central Pacific between 180° and $130^{\circ}W$ and south of $10^{\circ}N$. For that area data was often sparse or lacking and any analysis over that area should be considred as only tentative. The monthly SST data was averaged over the period of July to December for each year (Figure 2). These five averages were then used to produce a five-year mean chart from which the six-month mean SST anomaly charts were produced.

B. RADIOSONDE DATA

The radiosonde data were obtained from the National Climatic Center and consisted of once daily soundings (002) from July to December for 1974, 1975 and 1976, for the six stations shown in Figure 3. In total 184 soundings were used for each station (Koror, Yap, Truk, Ponape, Kwajalein and Majuro), during each year. For the 13 mandatory levels,

meridional and zonal wind components and specific humidities were calculated, and along with temperature they were smoothed over the six month periods using a band pass filter which eliminates variations with periods less than two days or greater than 10 days. The filter eliminates seven days on each end of the six month periods, thus leaving 170 days of filtered data. The filtered meridional wind components were then averaged vertically from 1000 mb to 600 mb and plotted in a time series to facilitate the selection of ridge, trough and maximum northerly and maximum southerly wind occurrences for each station and each period. If a ridge of trough passage falls between two soundings the preceding and following observations were averaged.

Although the number of significant trough passages varied between years and stations, from a low of 32 to a high of 44 (Table I), the average value for each year was 39 passages with a period of 4.4 days.

To describe the average wave structure for each station a compositing technique similar to the one used by Reed et al (1976) was used.

III. RESULTS

A. TIME MEAN FIELDS

1. Sea-Surface Temperature

Figure 1 shows the six-month averaged SST deviations for 1974, 1975 and 1976 from the five year mean (Figure 2). The year 1974 has a less distinctive anomaly pattern and may be considered as a "normal" year. The radiosonde stations studied in this work were either in or near a warm area, but in all cases there was a cold SST anomaly upstream.

The characteristics of each year's SST anomaly and the east-west SST gradient in the western Pacific may be identified by the 28° C isotherm in the actual SST distributions shown in Figure 2. In 1974 the 28° C isotherm is located just at 180° and is nearly north-south across the equatorial Pacific. In 1975 the 28° C isotherm is shifted to the west by more than 20° longitude. This indication of coldness in the eastern Pacific, and concomitant increased east-west SST gradient, are verified by the anomaly chart which shows a cold anomaly over the entire tropical Pacific east of 160° E. The stations from Truk east are all situated downstream from the broad negative anomaly area, whereas the western stations appear to be very close to the zero anomaly line.

In 1976 the 28° C isotherm is shifted eastward to 160° W near the equator, indicating a weakening of the eastwest SST gradient there.

The anomaly chart for 1976 shows that south of $10^{\circ}N$ along almost the entire equator there was a general warming over the previous two years and that the equatorial region is characterized as greater than $1^{\circ}C$ above the mean. In this year the eastern stations in our radiosonde network (Ponape, Kwajalein and Majuro) are in an opposite SST anomaly as compared to 1975, whereas the remaining stations remain near the zero anomaly line.

2. Mean Zonal Wind

The mean zonal winds for 1972 and 1973 as calculated by Chang and Miller show that in 1972, which was an extremely warm SST anomaly year in the eastern Pacific, the upper troposphere is completely dominated by easterlies and that the low levels are predominantly westerlies. In 1973, a very cold SST anomaly year in the eastern Pacific, there was almost a complete reversal of the flow at both the upper troposphere and the surface. The mean zonal winds for 1972 through 1976 are shown in Figure 4. In the central Pacific the 1975 period has a significant increase in the extent of the westerlies in the upper troposphere (near 150 mb) over 1974, while in 1976 this westerly maximum is substantially diminished. At the lower level weak westerlies are evident over the extreme western regions during 1974 and 1976, while in 1975 the surface wind was predominantly easterly.

By examining all five years together the similarities between the two cold SST anomaly years, 1973 and 1975, can be seen, likewise the two warm SST anomaly years differ

only in the eastward extent of the upper level easterlies and the lower level westerlies. It is of interest to note that 1974 falls between the two extremes of warm and cold anomalies as represented by 1972, 1976 and 1973, 1975 respectively.

During 1972, 1974 and 1976 there were also varying degrees of northward and eastward intrusions of the low-level monsoonal trough evident in the western Pacific.

B. TIME AND ZONAL SCALES OF THE WAVES

Using the subjective determinations of trough passage, taken from time series of vertically averaged 1000 mb to 600 mb meridional winds, the time scale and zonal scales of the easterly waves were estimated and shown in Table I. The average number of wave passages at each station for all three years was consistently close to 39, with an average period of 4.4 days and wavelength of 2930 km. This periodicity and wavelength corresponds very well with previous studies by Delaney (1977) and Chang and Miller (1976).

C. COMPOSITES OF WAVE STRUCTURE

Due to the proximity between the two stations and the similar SST environment and resultant wave structure, the composite data for Kwajalein and Majuro (K-M) were combined for all three years. The other stations either had distinct differences in the wave structure or were in different SST regimes, so they will be considered separately to allow for consideration of spatial and temporal variations.

1. 1974

a. Wind Components

The zonal (u) and meridional (v) wind fluctuations of the waves are shown in Figure 5. With the exception of Ponape the maximum u are located near the 200 mb level. At all stations the maximum v are located near 700 mb, and there is an indication that the intensity of the v component increases slowly from east to west.

All the stations show a northeast-southwest horizontal tilt in the lower levels which indicates that u and v are mostly in-phase. However, the vertical tilt, as indicated by the v component from 1000 mb to 300 mb, shifts from eastward at K-M and Truk to nearly vertical at Ponape, and finally westward at Yap and Koror. All the v components except Ponape show a distinct phase reversal near the 300 mb to 250 mb level. At Ponape there is a phase shift above 300 mb but due to the nearly vertical nature of the wave below 300 mb, the shift is not as significant as the other stations.

b. Temperature and Specific Humidity (Figure 6)

All stations, with the exception of Yap and Ponape, show definite warm core structure at the middle-upper troposphere. The temperature composites for both Yap and Ponape suggest very weak warm core systems, possibly reflecting less organized convective activity in the troughs.

The maximum moisture fields tend to be directly above or slightly ahead of the trough at all stations.

The vertical tilt in the moisture field does not appear to correspond to the wave tilt as indicated by the meridional winds, nor is there any consistent pattern of increase or decrease in moisture fluctuations from east to west.

2. 1975

a. u and v Components (Figure 7)

In 1975 again most stations, except Koror and K-M, possess a northeast-southwest tilt in the lower troposphere. Of the two exceptions the implied northwest-southeast tilt above 850 mb at Koror is not felt to be significant due to the very small zonal wind fluctuations.

The zonal wind fluctuations for K-M are comparable in magnitude to those of other stations, and they indicate that K-M may have a north-south to northwest-southeast tilt.

As in 1974 the vertical tilt of the waves is eastward with height at eastern stations; however, unlike 1974 the west stations do not exhibit a westward tilt, but appear nearly vertical up to 300 mb.

The maximum v fluctuation is located consistently at the 850 mb to 700 mb level, but again unlike 1974, there is no indication of amplitude change from east to west.

b. Temperature and Specific Humidity (Figure 8)

Only the westernmost station, Koror, possesses a distinct warm core. Yap indicates a weak warm core in the middle-upper troposphere and Truk, Ponape and K-M all

show weak cold core structures. Of the three eastern stations only Ponape has significant temperature values, and its temperature structure could be interpreted as either weak warm or weak cold core. Taking this into account it is best to describe the waves at stations from Truk eastward as not distinctly organized in their thermal structure.

With the exception of K-M the moisture structure for 1975 appears consistent, with an eastward tilt and the maximum values located at or behind the trough. K-M had a pronounced westward tilt with the maximum moisture found ahead of the trough at 850 mb.

3. 1976

a. u and v Components (Figure 9)

As in 1974 all stations in 1976 have a well defined northeast-southwest tilt. The intensities of the meridional winds are significantly greater at Truk and Ponape, with the maximum value located at 175-150 mb. The zonal wind maximum appears consistently located at the 700-600 mb level and all of the stations have virtually the same intensity.

Unlike the previous two years, there is no indication of change in the vertical wave tilt from east to west. The stations from Truk westward show a phase reversal in the zonal winds above 300 mb; below that they are nearly vertical. Ponape and K-M remain vertical from 1000 mb up to 175 mb and only show a slight phase shift above that.

b. Temperature and Specific Humidity (Figure 10)

K-M and Yap both possess distinct and well defined warm cores at the 400 mb to 250 mb levels. The remainder of the stations are characterized by indistinct temperature structures. Koror and Ponape appear as weak warm core systems; whereas Truk appears to be weak cold core.

Although K-M and Yap have similar thermal structures, their moisture distributions are completely dissimilar. Yap has its maximum moisture directly over the trough, as would be expected from a well developed warm core system, and K-M's maximum moisture is behind the trough and almost directly over the maximum southerly winds. The other three stations, which had indistinct thermal structures, all had their maximum moisture located upstream of the trough and tilting westward above 700 mb.

IV. DISCUSSION AND CONCLUSIONS

The main findings of this study may be summarized as follows:

1) The average wave period and wavelength for each year are relatively constant, at 4.4 days and 2930 km, respectively.

2) Except for Majuro in 1975 all stations for all three years have a definite northeast-southwest meridional tilt in the lower troposphere.

3) With the exception of 1976, the vertical tilt of the waves, as indicated by the phase shift between the 1000 mb to 500 mb levels, shifted from 1/2 cycle westward at the eastern stations to near vertical or 1/4 cycle eastward at the westernmost stations. The waves during 1976 showed little indication of tilt at all stations.

4) There are significant variations in the wave thermal structure during the entire study period.

To understand better the relationship between wave tilt and mean zonal wind, a graph was constructed with vertical tilt in cycles as the abscissa, and the station 150 mb mean zonal wind value as the ordinate (Figure 11a). The data for this plot covers 1972 to 1976 and yields a correlation coefficient of .71; thus, there appears to be a positive correlation between the vertical wave tilt and the background zonal flow. This also indicates that the waves

adjust rapidly to the mean flow as they propagate from east to west.

From the mean zonal wind data the ascending branch of the Walker circulation for the various years can be discerned from the regions of upper tropospheric divergence. Of the five years analyzed, 1973 exhibits the westernmost location of the ascending branch and 1972 the easternmost location, while 1974 through 1976 show variations between these two extremes. Because the Walker circulation seems to be related to the east-west SST gradient, it can be seen that the vertical tilt of the wave is indirectly influenced by the SST variations in the eastern Pacific via the Walker circulation.

Chang and Miller (1977) proposed that warm SST typically found in the western Pacific may result in more thermally driven "warm core" energetics, which would result in the wave vertical structure being less effected by the basic zonal flow. For 1976 this premise was partly substantiated. There was very little or not vertical tilt exhibited from east to west; however, the expected thermal structure was not well defined. To help determine what effect SST does have on the thermal structure of easterly waves, the averaged temperature difference between trough and ridge at the upper-middle tropospheric layer of 250-400 mb was plotted against the station SST. For this plot the correlation coefficient is -0.21. If the average SST at 10° longitude upstream is used instead of the local SST, the correlation coefficient becomes 0.29 (Figure 11b). If the correlation

coefficient is recalculated deleting the two easternmost stations, Kwajalein and Majuro, the correlation more than doubles to 0.60. These results indicate that the thermal structure of the waves was more influenced by the upstream SST than the local SST.

The adverse effects of Majuro and Kwajalein on the calculations suggests that the waves in the eastern-central Pacific may be instigated by mechanisms other than thermal forcing and have not undergone sufficient influence of latent heat release to become thermally driven by the time they passed Majuro and Kwajalein. As the waves continue westward the increased correlation suggests that the continued exposure to increasing SST enhances the convective activity and latent heating.

There appears to be reasonable evidence that variations in the tropical SST do effect both the large-scale and synoptic-scale features in the tropical Pacific. To determine sufficiently the extent that the thermal structure and energetics of the tropical easterly waves are influenced or modified by the local and large-scale SST variations, a more extensive data study should be undertaken. This would depend on both the expansion of data coverage from the present network and improved data quality.

Time and zonal scale of the waves. Numbers indicate number of waves passing indicated stations during 1974, 1975 and 1976. TABLE I.

KOROR YAP TRUK PONAPE KMAJURO MAJURO MAF IO 293 1974 39 36 41 40 36 41 38.6 4.4.4 293 1974 37 43 44 32 38.6 4.4.4 293 1975 40 37 4.3 4.4 32 38.6 4.4.4 293 1975 40 37 4.3 4.4 32 38.0 4.4.4 293 1976 4.1 36 4.3 38.5 4.4.4 293 1976 4.1 36 4.3 38.5 4.4.4 293								AVERAGE		WAVE-
1974 39 36 4.1 4.0 36 4.0 38.6 $\frac{4.4}{\text{days}}$ 293 1974 40 37 4.3 4.4 32 38.6 $\frac{4.4}{\text{days}}$ 293 1975 $\frac{4.0}{37}$ 37 4.3 $\frac{4.4}{32}$ 38.0 $\frac{4.4}{\text{days}}$ $\frac{293}{\text{km}}$ 1976 37 4.1 36 4.1 32 38.5 $\frac{4.4}{\text{days}}$ 293 1976 4.1 36 4.3 38 36.5 $\frac{4.4}{\text{days}}$ 293 1976 4.1 36 4.3 38 36.5 $\frac{4.4}{\text{days}}$ 293 1976 4.1 36 38 36 4.4 293		KOROR	YAP	TRUK	PONAPE	KWAJ.	MAJURO	WAVE NO.	PERIOD	LENGTH
40 37 43 44 32 38 39.0 4.4 293 1975 37 41 36 43 38 38.5 4.4 293 1076 37 41 36 43 38 36.5 4.4 293	1974	39	36	Γħ	0 4	36	0 tı	38.6	4.4 days	2930 km
37 41 36 43 38 36 4.4 293 1976 1976 38 36 43 38 36 4.4 293	1975	40	37	43	ħħ	32	38	39.0	4.4 days	2930 km
	1976	37	L H	36	H 3	38	36	38.5	4.4 days	2930 km

....







Fig. 1. Sea-surface temperature anomalies (^OC) for 1974, 1975 and 1976 taken from 5-year mean.



Fig. 2. Sea-surface temperature (°C) for 1974 through 1976 and 1972-1976 mean.























Composite vertical wave structure of temperature (^OC) and specific humidities (gm/kgx10) during 1975 for the indicated stations. 8. Fig.











shift (abscissa) for 1972-1976. Station and year indicated by letter and number, i.e. M5 is Majuro 1975. (b) Plot of average 250-400 mb trough temperature (^OC) (ordinate) vs. average 10° upstream SST (^OC) (abscissa). (a) Plot of 150 mb wind (m/sec)(ordinate) vs. 1000-500 mb vertical phase

LIST OF REFERENCES

Bjerknes, J., 1969: "Atmospheric Teleconnections from the Equatorial Pacific." Mon. Wea. Rev., 97, 163-172.

- Chang, C. P., C. R. Miller III, 1977: "Comparison of Easterly Waves in the Tropical Pacific during Two Periods of Contrasting Sea-Surface Temperature Anomalies." J. Atmos. Sci., 34, 615-628.
- Delaney, Dennis M., 1977: A Composite Satellite Study of the 1972-73 Easterly Waves in the Tropical Western Pacific, Master's Thesis, Naval Postgraduate School, Monterey.
- Krueger, A. F., and J. S. Winston, 1974: "A Comparison of Flow over the Tropics during Two Constrasting Circulation Regimes." J. Atmos. Sci., <u>31</u>, 358-370.
- _____, 1975: Large-scale Circulation Anomalies over the Tropics during 1971-72. Mon. Wea. Rev., 103, 465-473.
- Maas, Edwin, Jr., 1977: Composite Analysis of Easterly Waves in the Tropical Pacific during Two Constrasting Periods of Sea-Surface Temperature Anomalies, Master' Thesis, Naval Postgraduate School, Monterey.
- Namias, J., 1974: "Longevity of a Coupled Air-Sea-Continent System." Mon. Wea. Rev., 102, 638-648.
- Reed, R. J., and E. E. Recker, 1971: "Structure and Properties of Synoptic Scale Wave Disturbances in the Equatorial Western Pacific." <u>J. Atmos. Sci.</u>, <u>28</u>, 1117-1133.
- , D. C. Norquist, and E. E. Recker, 1976: "The Structure and Properties of African Wave Disturbances as Observed during Phase III of GATE." <u>Mon. Wea. Rev.</u>, <u>33</u>, 317-333.

INITIAL DISTRIBUTION LIST

1.

2.

3.

4.

5.

6.

7.

8.

9.

10.

11.

	No.	Copies
Defense Documentation Center Cameron Station Alexandria, Virginia 22314		2
Director, Naval Oceanography and Meteorol Naval Space Technology Laboratories Bay St. Louis, Mississippi 39529	ogy	l
Library, Code 0142 Naval Postgraduate School Monterey, California 93940		2
Professor CP. Chang, Code 63Cj Department of Meteorology Naval Postgraduate School Monterey, California 93940	:	10
Professor G. J. Haltiner, Code 63Ha Department of Meteorology Naval Postgraduate School Monterey, California 93940		1
Lieutenant D. Bepristis 394F Ricketts Drive Monterey, California 93940		2
Commanding Officer Fleet Numerical Weather Central Monterey, California 93940		l
Commanding Officer Naval Environmental Prediction Research F Monterey, California 93940	acility	,l
Dr. D. McCline Pacific Environmental Group National Marine Fisheries Service Fleet Numerical Weather Central Monterey, California 93940		1
Lieutenant D. M. Delaney USS TRIPOLI (LPH-10) FPO San Francisco 96601		1
Captain J. Shewchuk FWC/JTWC Box 2 FPO San Francisco 96630		1