

AD-A055 106

ARMY ENGINEER WATERWAYS EXPERIMENT STATION VICKSBURG MISS F/G 6/6  
PRELIMINARY GUIDE TO WETLANDS OF THE WEST COAST STATES. MAJOR A--ETC(U)  
APR 78

UNCLASSIFIED

WFC-TR-Y-78-4

1 OF 2  
AD  
A055 106



AD A 055106

DDC FILE COPY



FOR FURTHER TRAN

12



TECHNICAL REPORT Y-78-4

# PRELIMINARY GUIDE TO WETLANDS OF THE WEST COAST STATES

Major Associations and Communities Identified

Environmental Laboratory  
U. S. Army Engineer Waterways Experiment Station  
P. O. Box 631, Vicksburg, Miss. 39180

April 1978

Final Report

Approved For Public Release; Distribution Unlimited



Prepared for Office, Chief of Engineers, U. S. Army  
Washington, D. C. 20314



## PRELIMINARY GUIDE TO WETLANDS

### Major Associations and Communities Identified

<u>Technical Report No.</u>	<u>Region</u>
Y-78-2	Peninsular Florida <b>AD54191</b>
Y-78-3	Puerto Rico <b>AD55114</b>
Y-78-4	West Coast States
Y-78-5	Gulf Coastal Plain
Y-78-6	Interior-Great Lakes
Y-78-7	South Atlantic States
Y-78-8	North Atlantic States
Y-78-9	Alaska

Destroy this report when no longer needed. Do not return  
it to the originator.



DEPARTMENT OF THE ARMY  
WATERWAYS EXPERIMENT STATION, CORPS OF ENGINEERS  
P. O. BOX 631  
VICKSBURG, MISSISSIPPI 39180

IN REPLY REFER TO: WESYV

15 May 1978

SUBJECT: Transmittal of Technical Report Y-78-4

TO: All Report Recipients

1. The report transmitted herewith provides preliminary guidance on wetland determination to Corps of Engineers personnel responsible for the implementation of Section 404 of the Federal Water Pollution Control Act Amendments of 1972 (PL 92-500) in the West Coast States. This guide, sponsored by the Office, Chief of Engineers, represents one of a series of eight guides to the major wetland associations of the United States. Other guides include peninsular Florida, Puerto Rico, Alaska, South Atlantic States, Gulf Coastal Plain, North Atlantic States, and Interior-Great Lakes.
2. This guide is intended to assist in the field recognition of major wetland communities as they relate to the determination of jurisdictional boundaries in the implementation of the Section 404 permit program. It is neither a regional flora manual nor a general classification system. Several manuals that identify the flora of the West Coast States are referenced in this document and personnel requiring species identification are referred to those works. Personnel requiring a detailed wetland classification system may wish to consult "Classification of Wetland and Deep-Water Habitats of the United States (an operational draft)," prepared by the National Wetland Inventory Project of 1975-79 of the U. S. Fish and Wildlife Service.

ACCESSION for	
NTS	W & E Section <input checked="" type="checkbox"/>
DOC	B & F Section <input type="checkbox"/>
HANDLING	
J S I C A I T Y	
BY	
DISTRIBUTION/AVAILABILITY CODES	
SPECIAL	
A	

*John L. Cannon*  
JOHN L. CANNON  
Colonel, Corps of Engineers  
Commander and Director



14

WES-TR-Y-78-4

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report Y-78-4	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) PRELIMINARY GUIDE TO WETLANDS OF THE WEST COAST STATES Major Associations and Communities Identified		5. TYPE OF REPORT & PERIOD COVERED Final report
7. AUTHOR(s) Environmental Laboratory		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS U. S. Army Engineer Waterways Experiment Station Environmental Laboratory P. O. Box 631, Vicksburg, Miss. 39180		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Office, Chief of Engineers, U. S. Army Washington, D. C. 20314		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE April 1978
		13. NUMBER OF PAGES 98 p.
		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Community Plant community West Coast states Wetlands		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This guide to the major plant associations and communities found in wetlands within the West Coast States is one of a series of eight such guides, each prepared by a specialist or specialists familiar with the wetlands in the region covered by the guide. The guides are intended for distribution to the various U. S. Army Engineer Districts for use in wetland identification in the implementation of Section 404 of the Federal Water Pollution Control Act Amendments of 1972. The classification system in the guide is intended solely for use in the Section 404 permit program and is not considered a classification system for other purposes.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

438-144

hc



## SUMMARY

This report represents one of a series of eight preliminary guides to the dominant plant associations and communities found in the major wetlands of the United States. The primary purpose of the guidebook is to aid regulatory functions personnel in recognizing and delineating wetlands subject to permit regulation under Section 404 of Public Law 92-500 (Federal Water Pollution Control Act Amendments of 1972).

The guidebook is designed to be self-contained and consists of three parts. An introduction covers the purpose and use of the guidebook as well as general information about Section 404 wetlands. The second part, entitled "Wetlands of the West Coast States," consists of three major sections: Regional Environment, Regional Botanical References, and Wetland Types. The section on regional environment is brief and provides a broad context for the more detailed descriptions of the dominant plant associations and communities found in the major wetlands of the region. Because of synonymy of many scientific names, the nomenclature standard used for the guide is presented in the section on regional botanical references. Detailed description of wetland vegetation is based upon data in the literature and information from scientists having familiarity with the region. The goal of this section is to provide a description sufficiently detailed for field use but not to report minor variations of each wetland. Thus, the descriptions are a compromise between site-specific reports and extremely general discussions. The third part contains references to pertinent publications and Appendices A, B, and C and is specific to the region; a glossary that is common to all guides in the series (Appendix B) was added to aid in the user's clarity of understanding.

## PREFACE

At the request of the Office, Chief of Engineers (OCE), the Environmental Laboratory (EL) of the Waterways Experiment Station (WES) initiated production of this report, one of a series of eight preliminary guides to the dominant plant associations and communities found in the country's major wetlands. Other reports in the series apply to peninsular Florida, Alaska, Interior, Gulf Coast, North Atlantic, South Atlantic, and Puerto Rico. The reports are listed on the inside of the front cover. Funding was provided by OCE.

Dr. H. T. Harvey of Harvey, Hartesveldt, Heath, and Stanley, Inc., Santa Clara, California, provided a manuscript for initial construction of the draft guide under Purchase Order No. DACW39-76-M-2475.

Mr. Richard H. Daley, Ecologist, Missouri Botanical Gardens, St. Louis, provided major revision and rewriting of the draft copy under Purchase Order No. DACW39-76-M-5173 with the assistance of Mr. Ken Bierly of Montagne-Bierly Associates, Inc., Salem, Oregon, under Purchase Order No. DACW39-77-M-3086. Preparation of the guide was initiated by Dr. Luther F. Holloway, Research Botanist, EL. Dr. Gary E. Tucker, Research Botanist, EL, directed the production of the guide with the assistance of Dr. Robert Terry Huffman, Research Botanist, EL.

Ms. Dorothy P. Booth, EL, served as technical editor. The illustration used on the covers of this series of reports was drawn by Ms. Jane Barnes, Russellville, Arkansas.

The guide project was under the general supervision of Dr. H. K. Smith, Project Manager, Habitat Development Project; Dr. C. J. Kirby, Chief, Environmental Resources Division; Dr. Roger T. Saucier, Special Assistant, Dredged Material Research Program; and Dr. John Harrison, Chief, EL.

The Commanders and Directors of WES during the study were COL G. H. Hilt, CE, and COL J. L. Cannon, CE. Technical Director was Mr. F. R. Brown.



## CONTENTS

	<u>Page</u>
SUMMARY . . . . .	1
PREFACE . . . . .	2
PART I: INTRODUCTION . . . . .	5
Section 404 Permit Program . . . . .	6
Importance and Values of Wetlands . . . . .	7
Geographical Regions . . . . .	8
Wetland Types . . . . .	10
Organization of Guidebooks . . . . .	16
Botanical Nomenclature . . . . .	18
PART II: WETLANDS OF THE WEST COAST STATES . . . . .	21
Regional Environment . . . . .	21
Regional Botanical References . . . . .	23
Wetland Types . . . . .	23
SELECTED REFERENCES AND BIBLIOGRAPHY . . . . .	62
APPENDIX A: SCIENTIFIC AND COMMON NAMES OF PLANTS OF THE WEST COAST	
APPENDIX B: GLOSSARY	
APPENDIX C: INTERPRETATION OF WETLAND DEFINITION	

## LIST OF FIGURES

<u>No.</u>		<u>Page</u>
1	Guidebook regions . . . . .	4
2	Area of West Coast States covered by guide . . . . .	9
3	Distribution of wetland types by region . . . . .	11
4	Generalized salt marsh profile (southern) . . . . .	33
5	Generalized profile of northern low salt marsh . . . . .	34
6	Generalized profile of northern high salt marsh . . . . .	34
7	Generalized profile of California willow saltwater swamp . . . . .	39
8	Generalized profile of a freshwater aquatic wetland . . . . .	42
9	Generalized profile of a freshwater riparian or lacustrine marsh . . . . .	49
10	Generalized profile of a freshwater (wet meadow and fen) marsh . . . . .	50
11	Generalized profile of a freshwater (vernal pool) marsh . . . . .	51
12	Generalized profile of freshwater riparian swamp . . . . .	57
13	Generalized profile of freshwater montane swamp . . . . .	58






Figure 1. Eight geographical regions defined for the wetland guidebook series

## PRELIMINARY GUIDE TO THE WETLANDS OF THE WEST COAST STATES

### Major Associations and Communities Identified

#### PART I: INTRODUCTION

1. This guide to the major plant communities and associations found in wetlands within the West Coast States is one of a series of eight such regional guides, each prepared by a specialist or specialists familiar with the wetlands in the region covered by the guide. Other regional guides include Alaska, Interior, Gulf Coast, North Atlantic, South Atlantic, peninsular Florida, and Puerto Rico (Figure 1). The guides are intended for distribution to the various U. S. Army Engineer District regulatory functions personnel for use in identification of wetlands for the implementation of Section 404 of the Federal Water Pollution Control Act Amendments of 1972. The information provided is intended solely for use in the Section 404 permit program and is not considered a definitive classification system for other purposes.

2. Field personnel having need of a more detailed and definitive system of classification per se should consult one of the several wetland classification systems currently in use in the United States and Canada. The well-known Circular 39 (Shaw and Fredine, 1956) of the U. S. Fish and Wildlife Service has met with widespread use nationally despite its well-documented shortcomings. A recently published operational draft by the Fish and Wildlife Service (Cowardin et al., 1977) represents the most recent product of the National Wetland Inventory Project of 1975-79, an intensive effort that will result ultimately in the publication of a detailed and refined classification system to the wetlands of the entire nation. Numerous regional systems of classification also are available. Among the more significant regional classification systems are those of Golet and Larson (1974), Millar (1976), Odum et al. (1974), Penfound (1952), Stewart and Kantrud (1971), and Zoltai et al. (1975).

## Section 404 Permit Program

### Authority

3. Under the laws of the United States, Congress has assigned a number of nonmilitary functions to the U. S. Army Corps of Engineers. In addition to the well-known and more traditional roles in flood control, hydropower production, navigation, water supply storage, and recreation, the Corps has responsibility for some activities that are not so well known. Congress has given the Corps of Engineers regulatory responsibility to protect navigation channels and harbors against encroachments and also to preserve and restore water quality by regulating the discharge of dredged or fill material into waterways and wetlands.

4. The primary legislative basis for the Corps' regulatory authority for the disposal of dredged or fill material is the Federal Water Pollution Control Act Amendments of 1972. Section 404 of that Act gives authority to the Secretary of the Army, acting through the Chief of Engineers, to regulate the discharge of dredged or fill material in the waters of the United States.

5. Regulatory authority under Section 404 was initially considered limited to waters that are used presently, were used in the past, or could be used through reasonable improvements to transport interstate commerce. Limitation of the Corps' regulatory authority under Section 404 to navigable waters of the United States was successfully challenged in the District Court for the District of Columbia. On 27 March 1975, the Court ordered the Corps to extend its jurisdictional responsibility for the discharge of dredged or fill material under Section 404 to all waters of the United States (including the territorial seas) and adjacent wetlands and to revise its regulations accordingly.

6. In accordance with the Court's 1975 directive, the Corps of Engineers published an interim regulation in the Federal Register on 25 July 1975. The final set of permit regulations, considerably revised and reorganized, was published in the Federal Register on 19 July 1977.

### Scope

7. The Corps of Engineers permit program under Section 404 is extended to many areas that have never been regulated before. In



addition to the navigable waters of tradition, the Corps has been given jurisdictional authority over tributaries to navigable waters, including adjacent wetlands; interstate waters and their tributaries, including adjacent wetlands; and all other waters of the United States, such as lakes and rivers and streams that are not interstate waters or part of a tributary system to navigable waters of the United States; impoundments; perched wetlands; intermittent streams; and prairie potholes, the degradation or destruction of which could affect interstate commerce. In the absence of adjacent wetlands that are a part of the waters described previously, the landward limit of jurisdiction in tidal waters shall be the high tide line and the shoreward limit of jurisdiction in all other waters shall be the ordinary high water mark.

8. The term "wetlands" is a very crucial part of Section 404 and refers to those areas that are inundated or saturated by ground or surface water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions (Appendix C).

Purpose

9. The purpose of the Section 404 program, which is a part of the Corps of Engineers' overall regulatory authority, is to ensure that the chemical and biological integrity of waters of the United States is protected from unregulated discharges of dredged or fill material that could permanently alter or destroy the character of these invaluable natural resources.

Importance and Values of Wetlands

10. Wetlands are valuable and productive natural resources of national significance, and some of their major functions include the following:

- a. The provision of feeding, cover, and reproduction habitat for a great diversity of species, including endangered and threatened species.
- b. The provision of educational, study, refuge and sanctuary, and recreational areas.

- c. The maintenance of drainage, salinity, sedimentation, flushing, and current patterns.
- d. Cycling of nutrients.
- e. Reduction of contaminant loading.
- f. Protection from erosion and storm damage.

### Geographical Regions

11. Eight geographical regions have been defined for the wetlands guidebook series: Alaska, West Coast, Interior, Gulf Coast, North Atlantic, South Atlantic, peninsular Florida, and Puerto Rico. The geographical regions are based on both physiographic and pragmatic considerations; the boundaries were influenced significantly by the works of Fenneman (1931, 1938). The use of natural units rather than artificial ones, such as political boundaries, minimizes the number of wetland types described in each guidebook. Several states are covered by a combination of two guidebooks, and a very few are covered by three guidebooks. Physiographic parameters were used where possible, since both hydrologic and biotic patterns are related closely to landscape features. Each of the regions will be covered in a separate guidebook. Geographic descriptions for the guides are as follows:

- a. Alaska. The state of Alaska is the sole subject of an entire guide. Particular emphasis is placed on coastal wetlands; much of the interior region is "wet", but further study is necessary to determine the exact jurisdictional limits of Section 404.
- b. West Coast. This region includes most of California (exclusive of the southeastern part), western Oregon, and western Washington (Figure 2).
- c. Interior. The area covered by this region consists of the vast interior of the United States, including much of the Southwest, the Rockies and some of the intermontane region, the Central Plains, and the Midwest. States contained within the region are numerous.
- d. Gulf Coast. The Gulf Coast region extends from the coastal plain of Texas to western Georgia. Inland, the coastal plain extends to southern Missouri in the Mississippi embayment; other states included in the region




Figure 2. Area of West Coast States covered by guide



are all or parts of Oklahoma, Arkansas, Louisiana, Mississippi, Alabama, Florida, and Tennessee.

- e. North Atlantic. This region extends north from Sandy Hook, New Jersey, to the Canadian border and west to the Appalachian highlands. Included within the region is northern New Jersey, New York, and New England.
- f. South Atlantic. Included within this region is everything north from peninsular Florida to Sandy Hook, New Jersey, and west to the Appalachian highlands. The separation of this region from the North Atlantic region is based largely on substrate features; the exposures of bedrock throughout the North Atlantic region are strikingly different from the thick mantle of Coastal Plain sediments predominating in most of the South Atlantic region. Additionally, most of the species of the "southern" swamp forest are restricted to the South Atlantic region as defined here.
- g. Peninsular Florida. There is no clear physiographic distinction between peninsular Florida and the Gulf Coast and Atlantic Coast regions, but the vegetation of peninsular Florida has strong enough tropical affinities to warrant separate treatment. The peninsular region has been delineated by an arbitrary boundary extending from Jacksonville west to Steinhatchee, with all of Florida south of the boundary included in the region. "Subtropical Florida" as defined by Fenneman (1931) and Braun (1964) is essentially conspecific with this region.
- h. Puerto Rico. The guidebook is intended for use in Puerto Rico; however, its utility may extend to the U. S. Virgin Islands because the vegetation of the two regions has many similarities.

#### Wetland Types

##### General information

12. Nine basic wetland types are recognized in the United States (Figure 3). An interpretation of the definition of wetlands is given in Appendix C. The number of wetland types in each guidebook region, however, is either seven or eight, since no region has all possible types. The nine basic wetland types have been distinguished by a combination of differences in physiognomy (e.g., marsh versus swamp), growth form (e.g., herbaceous plants versus trees), and environmental factors (such as degree of salinity in soil and water). Terms used on a

	Alaska	West Coast	Interior	Gulf Coast	North Atlantic	South Atlantic	Peninsular Florida	Puerto Rico
Saltwater Aquatic	✓	✓	✓	✓	✓	✓	✓	✓
Saltwater Coastal Flat	✓	✓	x	✓	✓	✓	✓	✓
Saline Inland Flat	x	x	✓	x	x	x	x	x
Saltwater Marsh	✓	✓	✓	✓	✓	✓	✓	✓
Saltwater Swamp	x	✓	x	✓	x	x	✓	✓
Freshwater Aquatic	✓	✓	✓	✓	✓	✓	✓	✓
Freshwater Flat	✓	✓	✓	✓	✓	✓	✓	✓
Freshwater Marsh	✓	✓	✓	✓	✓	✓	✓	✓
Freshwater Swamp	✓	✓	✓	✓	✓	✓	✓	✓

✓ - present

x - absent

Figure 3. Distribution of wetland types by region

regional basis in the description and definition of wetlands, such as bog and pocosin, are discussed in the text at appropriate points.

#### Identification

13. The approach to the identification of wetlands in this guide-book series is to provide general classifications for each region of the country. For purposes of this series, the country has been divided into six large regions plus Alaska and Puerto Rico (as described earlier). Within each regional guide, a key (Table 1) is provided for classification of any site in question. The reader is then referred to a brief description of the type (Wetland Types, next section) for a preliminary check to see if the site was properly classified. Finally, the reader is referred to the text for a more complete description of the communities and associations in the wetland and a pictorial profile

Table 1  
Key to Wetland Types

---

A. Aquatic vegetation predominant (dominant plants free-floating or attached and having poorly developed tissues of structural support, supported and buoyed up by the water); flooded usually for long periods or permanently	
B. Coastal; below the intertidal zone; seaward to limits of vascular plant growth; permanently flooded . . . . .	SALTWATER AQUATIC
B. Inland; flooded permanently or semipermanently by fresh water . . . . .	FRESHWATER AQUATIC
A. Terrestrial vegetation predominant (dominant plants rooted and with well-developed tissues of structural support) or sometimes barren of vegetation; flooded at least occasionally, often for prolonged periods	
C. 25 percent or less vegetative cover	
D. Subject to saltwater influence	
E. Coastal, tidal . . . . .	SALTWATER COASTAL FLAT
E. Inland, nontidal . . . . .	SALINE INLAND FLAT*
D. Fresh water . . . . .	FRESHWATER FLAT
C. More than 25 percent vegetative cover	
F. Nonsaline soils	
G. 40 percent or less cover by woody plants . . . . .	FRESHWATER MARSH
G. More than 40 percent cover by woody plants . . . . .	FRESHWATER SWAMP
F. Saline (including brackish) soils	
H. 40 percent or less cover by woody plants . . . . .	SALTWATER MARSH
H. More than 40 percent cover by woody plants . . . . .	SALTWATER SWAMP

---

\* The saline inland flat does not occur in those parts of the West Coast States defined by this guide.



Table 1 (Continued)

How to use the key: A key is an artificial device constructed for the purpose of identifying an unknown object. Keys traditionally have been used in the field of biology for the identification of unidentified plant and animal species, but in this guidebook the key will be used for the identification of unidentified wetland types.

The key to wetland types consists of a series of contrasting statements or descriptions, and the user of the key is required to make decisions based on the comparison of statements in the key as related to observations on the unidentified wetland type. The user must work carefully through the key from its beginning until a wetland type has been selected for the area in question.

The key is constructed around a series of pairs of leads. The second lead of a pair usually repeats the data given in the first lead but in a negative sense. Let us assume that you, the user of the guidebook, have located a grass-dominated area that obviously is "wet" during the better part of the year and obviously under the jurisdiction of the Section 404 program. Proper use of the key should enable you to determine just what type of wetland is involved.

In order to begin use of the key, you must start with the first pair of lead sentences, in this case labelled "A." Read each lead carefully, weighing one against the other with relation to your grass-dominated area. Grasses normally do not grow as free-floating organisms nor do they depend on water to buoy them upright, since they normally have sufficient supporting tissues to grow erect; in this case, then, the second lead of the pair of choices is better descriptive of the grass-dominated area with which you are concerned. You are now ready to consider a second pair of leads. This time you will consider the leads labelled "C" (of course, if your habitat were dominated by aquatic vegetation rather than terrestrial grasses, you would be considering the choices labelled "B"). Read the two "C" leads carefully, look at your grassy area, and try to determine how much of the ground surface is covered by vegetation. If less than 25 percent of the ground surface is covered by vegetation and more than 75 percent of the area is bare ground, you will select the first "C" as indicated; if vegetative cover accounts for more than 25 percent cover, you will take the second choice labelled "C." Let us assume that your area has only 10 percent cover. You will select the first "C" and then proceed to the "D" possibilities. Is the area in question flooded by fresh water or salt water? Let us make the assumption that you are in a freshwater area; look at the key carefully and note that the second "D" lead has a series of dotted lines leading to the phrase "Freshwater Flat." After the process of first rejecting and then accepting leads, you finally have arrived at an identification of your wetland type.

Table 1 (Concluded)

After determining the wetland type of an area in question, the user should turn to the detailed description of that particular type in the guidebook. In our hypothetical case the user would turn to page 45, FRESHWATER FLAT, and carefully read the descriptive material.

The use of the key may not be as simple and easy as it may seem. After you have followed the key through until coming to an identification of the wetland type, it may appear that the wetland description does not seem to fit the site. In that case it always pays to go back to the key and make sure an error has not been made through haste or misunderstanding of terms used. Occasionally an area may be found that cannot be identified with the aid of the key; the entire guidebook is written from a regional perspective and does not cover all variations of each wetland type. If a site does not fit any of the wetland types as described but yet is suspected of being a wetland under Section 404, a professional ecologist or botanist may be required for a quantitative study of the vegetation at the site.



illustrating its dominant species. The description of each wetland association is concluded with a section entitled "Field Identification," which briefly explains how to distinguish the wetland from other wetland types and from adjacent uplands. The entire description of a wetland should be studied prior to using the field identification section, however, to familiarize the user with its major variations. Wherever feasible, characteristics of growth forms are highlighted for identification, but if classification of an area is questionable, final determination must be based upon species composition.

14. If a site "fits" the description reasonably well, then the decision is clear that the area should be classified as a wetland of that particular type. The converse is not true, however. (If the site does not closely match one of the descriptions, it cannot be concluded unequivocally that the area is not a wetland.) This text is written from a regional perspective and consequently cannot be comprehensive and describe all variations within each wetland type. If a site does not fit any of the descriptions yet is still suspected to be a wetland, a quantitative survey of the vegetation of the area will be necessary. Especially in cases where the natural vegetation cannot be ascertained, hydrologic and soil information will be required to determine whether or not a site is a wetland. The nine basic wetland types are defined as follows:

- a. Saltwater aquatic. Wetlands that are dominated by free-floating, rooted, or otherwise attached herbaceous plants (including macroscopic marine algae) and that are permanently flooded by saline or brackish water (e.g., sea grass beds).
- b. Saltwater coastal flat. Wetlands that have 25 percent or less vegetative cover and are occasionally (shallow flat) or regularly (deep flat) flooded by saline water of tidal origin (e.g., nonvegetated intertidal zone).
- c. Saline inland flat. Wetlands that have 25 percent or less vegetative cover and are occasionally or regularly flooded by saline water of nontidal origin (e.g., inland salt flat).



- d. Saltwater marsh. Wetlands that have more than 25 percent vegetative cover of herbaceous plants but 40 percent\* or less cover by woody plants and that are occasionally (high marsh) or regularly (low marsh) flooded by brackish or saline water (e.g., Smooth cordgrass marshes).
- e. Saltwater swamp. Wetlands that have more than 40 percent cover of woody plants and are occasionally or regularly flooded by brackish or saline water (e.g., mangrove swamps).
- f. Freshwater aquatic. Wetlands that are usually dominated by free-floating or rooted aquatic herbs and are semipermanently or permanently flooded by fresh water (e.g., floating duckweed mats).
- g. Freshwater flat. Wetlands that have 25 percent or less vegetative cover and are occasionally or regularly flooded by fresh water (e.g., mudflats).
- h. Freshwater marsh. Wetlands that have more than 25 percent vegetative cover of herbaceous plants but 40 percent or less cover by woody plants that are occasionally or regularly flooded by fresh water (e.g., cattail marsh).
- i. Freshwater swamp. Wetlands that have more than 40 percent cover by woody plants and are occasionally or regularly flooded by fresh water (e.g., cypress swamps).

#### Organization of Guidebooks

15. Each guidebook is designed to be self-contained. Although this necessitates repetition of general information in the introductory part, the advantages in utility outweigh the duplication. The second part of each guidebook, entitled "Wetlands by Region," is the only one of the three parts unique with each guide. The third part, containing appendixes and references to pertinent publications, is largely specific for each region, except for a glossary that is common to the entire group of regional guides.

16. Three major sections are found in Part II: Regional Environment, Regional Botanical References, and Wetland Types. The section on

---

\* The use of 40 percent as the division for woody plant cover is convenient for field work because when the tree cover is 40 percent, the distance between tree crowns equals the mean radius of a tree crown (UNESCO, 1973).

regional environment is brief and provides a broad context for the more detailed descriptions of wetland types in each region. Because of the synonymy of many scientific names, the standard used for the guide is given in the section on regional botanical references.

17. Description of each wetland type is based upon data in the literature and from discussions with scientists having familiarity with the area. The goal is to provide a sufficiently detailed description for use in the field but not to report every possible variation of each wetland type. Thus, the descriptions are a comparison between site-specific reports and extremely general discussions.

18. The description of vegetation in each wetland type is divided into the following four parts:

- a. Growth form. Growth form, such as deciduous (e.g., Alder, Bald cypress) or evergreen (e.g., pine, Sitka spruce) trees, is a concise description based upon the physiognomy of the vegetation. This should be particularly helpful to those not familiar with the species in the area.
- b. Species composition. Discussion of species composition in each case includes listings in alphabetical order (by scientific name) of the dominant plants and the most commonly associated species. Because of local variation within any wetland type, an alphabetical listing is preferred over an attempt at listing species by importance value. The choice of associated species listed sometimes is arbitrary but, in the absence of complete species lists for each type, is inescapable. Profiles are provided for most wetland types. These diagrammatic depictions of vegetation structure are meant only to reinforce the textual material. The section on transition zones outlines the plants or plant communities characteristically found between adjacent wetland types or between wetlands and uplands. Such transitions may be abrupt but more often they are gradual. The generalized structure of each wetland type and its relationship to transition zones is indicated by a pictorial vegetation profile.
- c. Physical environments. The environmental conditions, the characteristic water regimes, and soils of each wetland type, are described where available. The discussions are limited to aspects of the physical environment most often affecting the vegetation and are not intended to fully describe the environment.
- d. Field identification. The section on field identification gives the characteristics that distinguish the wetland type from other wetland types and from adjacent uplands.



19. In most cases some attempt to discuss successional relationships of wetland communities is made. In many cases, however, the successional relationships of wetlands vegetation are too poorly understood for meaningful generalizations.

20. The primary purpose of the guidebook series is to aid regulatory functions personnel in identifying wetland types. For that reason a well-organized but general approach has been attempted. The classification system in the guides is intended solely for implementation in the Section 404 permit program and is not considered a definitive classification system for other purposes.

### Botanical Nomenclature

#### Common names

21. Common names, while admittedly convenient, often vary from place to place. One species may have several names in different geographic regions, or the same name may be applied to unrelated species in different areas. Yet other species lack a common name. In the guidebook series, the common name used for a plant is the one, in the opinion of the author, most often used locally within the region. A single common name is used even though several names may be in use within the region. Those species not known to have a common name are referred to by their scientific name.

22. To assist in utility of the guides, an attempt has been made to provide a common name at each point where a scientific name appears. In a few cases, however, this has not been practical or has been considered superfluous; for that reason, in cases where assurance of communication seemed evident, a single name was employed.

#### Scientific names

23. Botanists, ecologists, and other scientists use scientific names in their technical publications and discussions. The Latin form of scientific names is definitive and uniformly adhered to by botanists around the world under the International Code of Botanical Nomenclature.



Thus, the Latin name of a plant species is understood by the scientific community throughout the world, regardless of the prevailing language in a country.

24. Scientific names used in this guidebook series consist of two words. The first word of the scientific name is that of the genus to which a plant belongs, and it is always capitalized. The second word of the scientific name is referred to as the specific epithet, and it is printed here in lower case even though it may be derived from a geographical name or the name of a person. Both words are italicized or underlined. Following the scientific name it is customary, at least in checklists, to give the name of the author or person who originally described the plant to science; the name of the author is referred to as the authority. The authority for plants in these guides is given in Appendix A and in most cases the authority is abbreviated.

25. The following example illustrates the function and meaning of a typical scientific name. The genus *Typha* was first described by the Swedish botanist Linnaeus, as was *Typha latifolia*, the Common cattail, which occurs over most of the United States. Its name, therefore, is written *Typha latifolia* L., indicating that this species was described by Linnaeus. The scientific name *latifolia* indicates that the plant has broad leaves, in this case an accurate description.

26. Occasionally, there is need to refer to an unidentified species of a particular genus; an unidentified species of *Potamogeton*, for example, would be referred to in the text as *Potamogeton* sp. Similarly, it is sometimes convenient to refer to a group of species of a particular genus without giving the complete scientific name of each. A group of species of the genus *Potamogeton* would be given as *Potamogeton* spp.

27. Within the text of a paragraph or more of material, it is considered redundant to repeat the complete scientific name repetitively after its initial use. The species *Potamogeton amplifolius* would be given in full where first mentioned but at later times might be referred to in the text as *P. amplifolius*, the *P.* being an abbreviated form of *Potamogeton*. In situations where confusion with other species might result, however, the scientific name is given in full.

#### Synonymy of scientific names

28. Many plant species have been given more than one scientific name in the course of botanical history. A species may have been described and named independently by different botanists, or two species may have been considered one and the same following a period of study. In addition, there are differences of opinion among professional botanists as to whether a variation merits recognition as a variety or as a separate species or perhaps needs no additional name.

29. Because of differences of interpretation, one will often find a particular plant referred to by different scientific names in two or more separate publications. For this reason each of the guidebooks in this series has been compiled with the use of a particular publication as a standard for botanical nomenclature. In each case the standard for botanical nomenclature is a well-known regional manual of plant identification. The standard for each guidebook is identified in the section entitled Regional Botanical References.

## PART II: WETLANDS OF THE WEST COAST STATES

### Regional Environment

30. The West Coast region extends from the Mexican to the Canadian border and includes California (except the southeastern section), western Oregon, and western Washington. This narrow strip is approximately one thousand miles long and one to two hundred miles wide. Mountain ranges form a giant H with the Oregon Coast Range and the Cascades as the upper part of the H, and the California Coast Ranges and the Sierra Nevadas as the lower part of the H. The Klamath mountains in northern California and southern Oregon form the cross bar of the H between the north-south ranging mountains. Between the Oregon Coast Range and the Cascades is the Puget Trough/Willamette Valley region, while the Central Valley of California separates the California Coast Ranges from the Sierra Nevadas. Elevations vary from slightly below sea level to over 4250 m on the highest mountain peaks, resulting in a diversity of wetland types.

31. The climate is extremely variable across the expansive latitudinal and topographic range of the region. Barbour and Major (1977) indicate that very little is known about California climatic factors, because the available climatic data is poor. Annual precipitation is less than 12.5 cm in the Central Valley of California and over 254 cm on the Olympic Peninsula in the Coast Range of Washington and also at higher elevations of the Cascades. Precipitation is highly seasonal in many parts of the region, particularly in coastal southern California, which is marked by a Mediterranean climate. Winter is the wet season throughout the region.

32. Temperature fluctuations on a seasonal basis are not pronounced. Mean July and January temperatures usually vary less than 17°C in any part of the region. July temperatures average above 27°C in the south and about 16°C in the Cascades. January temperatures average above 10°C in the south to about 4°C on the Olympic Peninsula.



33. The so-called Mediterranean climate of southern California is marked by hot, dry summers and cool but not cold, moist winters (Munz, in collaboration with Keck, 1973; Barbour and Major, 1977) and presumably has had a major influence on the evolution of the vegetation. The vegetation of coastal areas in southern California is dominated by pine forests and chaparral. Chaparral is a mixture of small trees and stout woody shrubs and typically grows on hills and lower mountain slopes. The broad-leaved evergreen species associated with the chaparral often form dense, nearly impenetrable thickets. The chaparral has a number of adaptive features that allow it to persist despite burning. Some of the shrub species have horizontal rootstocks that sprout again after the tops are removed by fire; other plants are kinds that reseed freely after fire and thus maintain themselves in the burned area. Many of the pines of the region also are favored ecologically by fire. The chaparral ascends to about 1500 m along the coast of southern California; in northern California, however, the chaparral is replaced by coastal redwood forests where rainfall and fog provide greater amounts of precipitation. The Central Valley of California is naturally a desert but has been developed agriculturally through the use of irrigation.

34. The northern portions of the West Coast region are dominated by temperate coniferous forest; Western red cedar, hemlock, Douglas fir, and spruce are dominant species. The vegetation of the Puget Trough/Willamette Valley is extremely varied; Western hemlock forest, pine forest, oak woodland, and prairie all are encountered. The lowlands and valley bottoms generally are relatively warm, dry regions from which many of the mesic species common on nearby montane slopes are absent.

35. Wetlands in the West Coast region are both widespread and diverse, but they are much less extensive today than in the past. Large acreages of both saltwater and freshwater wetlands have been destroyed through salt-production activities and draining and clearing operations associated with agriculture and expanding urbanization.

### Regional Botanical References

36. There is no single manual of plant identification applicable to the entire West Coast region. For that reason selection of a single work to serve as a standard for botanical nomenclature has not been possible.

37. Mason (1957) was used as a primary reference for the southern part of the region; that reference is well illustrated and specifically covers wetland plant species. In the northern part of the region, the work of Hitchcock et al. (1955-69) was used extensively.

38. Other taxonomic references of particular value to workers in the region include Munz (1973), Steward et al. (1963), and Hulten (1968). The ecological treatise on the vegetation of Washington and Oregon by Franklin and Dyrness (1973) also is useful, although few wetland types are treated specifically. The compendium of information on California vegetation edited by Barbour and Major (1977) similarly is of much value.

39. Resolution of the numerous discrepancies in botanical nomenclature among the several publications used as references in the construction of this guide was by necessity often arbitrary. Choice of common names is based on general usage in the guidebook region, but here also many decisions were made arbitrarily.

### Wetland Types\*

40. Of the nine possible wetland types, the West Coast region possesses eight. A brief definition of each type follows:

- a. Saltwater aquatic. Wetlands that are dominated by free-floating, rooted, or otherwise attached herbaceous plants (including macroscopic marine algae) and are permanently flooded by saline or brackish water (e.g., seagrass beds).
- b. Saltwater coastal flat. Wetlands that have 25 percent or less vegetative cover and are occasionally or regularly

---

\* See Appendix C for an interpretation of the definition of wetlands.

flooded by saline water of tidal origin (e.g., sparsely vegetated intertidal zone).

- c. Saltwater marsh. Wetlands that have more than 25 percent vegetative cover of herbaceous plants but 40 percent or less cover by woody plants and are occasionally or regularly flooded by brackish or saline water (e.g., hairgrass marsh).
- d. Saltwater swamp. Wetlands that have more than 40 percent cover of woody plants and are occasionally or regularly flooded by brackish or saline water (Goodding's and white willow swamps).
- e. Freshwater aquatic. Wetlands that are dominated usually by free-floating or rooted aquatic herbs and are semi-permanently or permanently flooded by fresh water (e.g., water lily stands).
- f. Freshwater flat. Wetlands that have 25 percent or less vegetative cover and are occasionally or regularly flooded by fresh water (e.g., mudflats in reservoir draw-down areas).
- g. Freshwater marsh. Wetlands that have more than 25 percent vegetative cover of herbaceous plants but 40 percent or less cover by woody plants and which are occasionally or regularly flooded by fresh water (e.g., cattail marsh).
- h. Freshwater swamp. Wetlands that have more than 40 percent cover by woody plants and are occasionally or regularly flooded by fresh water (e.g., Black cottonwood-willow-alder swamps).



## SALTWATER AQUATIC WETLANDS

Definition: Wetlands that are dominated by free-floating, rooted, or otherwise attached aquatic herbs or algae and are flooded permanently (or near permanently) by brackish or saline water.

41. The saltwater aquatic wetland is distributed along the entire Pacific Coast. The shoreward limit is generally the elevation of the lower mean tide, while the seaward boundary is the limit of attached macrophytic plant growth. Plants are exposed to air only during exceptionally low tides in most cases.

42. Although there is no accurate appraisal of the total area occupied by this wetland type, 2000 hectares of Eelgrass have been identified in Oregon; areas of Eelgrass adjacent to the Washington coast have been mapped, but the total area has not been calculated.

43. Seagrass beds, particularly of Eelgrass, in the saltwater subtidal zone are an extremely important contributing factor to the productivity of estuaries and coastal areas. Submerged saltwater aquatic wetlands play several other significant roles. The vascular plants of the seagrass beds serve as substrates for an epiphytic assemblage of algae and simple invertebrates. The beds also are important habitats for various fish and shellfish as well as for diving ducks and certain other waterfowl.

44. Beds of macroscopic marine algae (representing Brown algae, Red algae, and Green algae) occur in relatively shallow marine waters of the region; the algal beds attain high development in offshore coastal areas. Kelps (large Brown algae) are the most complex of all algae and have large, differentiated plant bodies that are superficially similar to those of higher plants.

45. Dense subtidal stands of kelps and other large seaweeds afford protection, breeding, spawning, and grazing areas for a variety of marine animal species. The large algal plant bodies serve as substrates to which a variety of marine life is attached. Some marine organisms feed directly on the algae, while others depend upon the detrital

material provided by these large algae. Many of the large seaweeds make important contributions to phytoplankton populations (and hence primary production) through liberation of reproductive cells (*Nereocystis* releases an estimated 3,000,000 reproductive cells per litre of water per day from June through September).

#### VEGETATION

46. Growth forms and physiognomy: submerged, narrow-leaved herbaceous plants, such as Eelgrass, and algae with leaflike blades (macroscopic forms), in sparse to dense stands.

47. Species composition of the saltwater aquatic wetland (potentially dominant species):

*Phyllospadix scouleri* (Surfgrass, Open-coast eelgrass)  
*Ruppia maritima* (Widgeon grass)  
*Zostera* spp. (Eelgrass)

*Agardhiella* spp. (Red algae)  
*Alaria* spp. (Brown algae), Winged kelp  
*Codium* spp. (Green algae), Dead man's fingers  
*Hymenena* spp. (Red algae)  
*Iridaea cordata* (Red algae)  
*Laminaria* spp. (Brown algae), Devil's apron  
*Lessonia* spp. (Brown algae)  
*Macrocystis* spp. (Brown algae), Giant or Vine kelp  
*Nereocystis* spp. (Brown algae), Bull or Bladder kelp  
*Postelsia* spp. (Brown algae), Sea palm  
*Pterygophora* spp. (Brown algae)  
*Ulva* spp. (Green algae), Sea lettuce

Dominant and associated species. Eelgrass often is the only species of higher plant present in seagrass beds. Despite the paucity of vascular plant species, these areas support an extremely diverse fauna as well as many microscopic algae (such as diatoms) that reside epiphytically on the Eelgrass. Eelgrass beds that have rocks scattered on the surface commonly have the algae *Ulva lactuca* or *Agardhiella tenera* as associates.

Algal beds (primarily of large marine Brown algae), unlike Eelgrass beds, support many important species and have the plants arranged in distinct vertical strata; species diversity in the algal beds is high and selection of dominant species in the preceding list was largely arbitrary. Large, massive kelp species (*Macrocystis*, *Nereocystis*, *Iridaea*, etc.) form a canopy that effectively reduces light penetration beneath them. Vertical layering below the canopy is evidenced by long-stalked species (*Lessonia* spp.) situated above a lower-most layer of relatively smaller algae such as *Codium* or *Hymenena*.

Long-term observations on the successional relationships of seagrass communities are few. According to den Hartog (1977), the *Phyllospadix* beds of the Pacific coast are successional to the development of the permanent submarine forests dominated by *Macrocystis* and other giant kelps. *Phyllospadix* apparently remains permanently only on those sites that are too shallow for the development of large kelps. *Zostera* beds appear to represent both pioneer and permanent stages of succession.

Transitional species. The saltwater aquatic community frequently borders other wetland types. The seaward limit of the algal community corresponds with the limit of algal attachment, while the seaward limit of the seagrass beds is the limit of rooted plant growth. The shoreward boundary is usually the lower limit of the intertidal zone. Eelgrass commonly occurs in the tidal channels of salt marshes as well as in intertidal stands; occasionally it occurs in the lower edges of the saltwater flat. Kelps extend from the lower intertidal region (where they often are dominant) to depths of 20 metres or more.

#### ENVIRONMENTAL CONDITIONS

48. Eelgrass commonly is found in estuarine areas of silty substrates that have a strong saltwater influence. The depth of the water in which it grows may be as much of 7 metres, or the water may be very shallow.

49. Kelps and other forms of macroscopic algae grow in coastal waters having salinities as high as those in the ocean proper as well as in estuarine environments having relatively low salinities. The very large kelps, such as *Nereocystis*, *Macrocystis*, etc., are found in waters from 8 to 35 metres deep. Many of the smaller forms, however, typically are found in the lower intertidal and upper subtidal areas.

50. The ocean often appears to provide a relatively uniform environment, but the underlying topography and substrate actually may be highly variable; diversity in physical and chemical factors of the marine environment is reflected in the species diversity supported by the environment. Rocky coastlines support both the greatest diversity and the largest number of organisms. Unstable substrates such as muds and sands generally support few macroscopic algal species, although microscopic and mat forming filamentous forms may be abundant. The strength of the currents likewise plays an important role in determining species distribution.



51. The macroscopic Brown algae are abundant along much of the Pacific Coast and many species are characteristic of the intertidal and shallower subtidal areas. In such areas a striking vertical zonation of species often occurs. This zonation is due to a phenomenon associated with tidal patterns in the water. The physical-chemical conditions in the shallow water along the coastline are highly variable, often changing greatly along very small vertical distances. Gradients relating to salinities, nutrient availability, light, and temperature are inter-related with the tidal phenomenon in complex patterns, contributing to distinct vertical zonation of species. Waaland (1977) discusses seaweed habitats in some detail.

## SALTWATER COASTAL FLAT

**Definition:** Wetlands that have 25 percent or less vegetative cover and are occasionally or regularly flooded by saline water of tidal origin

52. Coastal flats form a fringe along the seaward edge of salt marshes and along sandy and silty estuarine shorelines. Relatively few plants grow in these wetlands, but saltwater flats play an extremely important role in estuarine benthic production and mineral cycling. Several species of birds (including Belding's savannah sparrow, California clapper rail, and Light-footed clapper rail) frequent this wetland type.

### VEGETATION

53. Growth forms and physiognomy: nonvegetated or sparsely vegetated with succulent forbs, plants often in circular stands.

54. Species composition of the saltwater coastal flat wetland:

#### Dominant species

*Cotula coronopifolia* (Brass buttons)  
*Distichlis spicata* (Saltgrass)  
*Eleocharis parvula* (Spike rush)  
*Puccinellia pumila* (Alkali grass)  
*Ruppia maritima* (Widgeon grass)  
*Salicornia* spp. (Glasswort or Pickleweed)  
*Scirpus americanus* (Three-square bulrush)  
*Scirpus maritimus* (Saltmarsh bulrush)  
*Scirpus validus* (Soft-stemmed bulrush)  
*Spergularia canadensis* (Sand spurry)  
*Spergularia marina* (Saltmarsh sand spurry)  
*Triglochin maritima* (Sea arrowgrass)  
*Zannichellia palustris* (Horned pondweed)  
*Zostera* spp. (Eelgrass)  
  
*Cladophora* spp. (Green algae)  
*Enteromorpha* spp. (Green algae)  
*Fucus* spp. (Brown algae), Rockweed  
*Ulva* spp. (Green algae), Sea lettuce

#### Associated species

*Carex lyngbyei* (Lyngbye's sedge)  
*Spartina foliosa* (Cordgrass, California cordgrass)  
Almost any species normally found in the salt marsh also occurs here

Dominant and associated species. Coastal flats intergrade closely with low salt marshes. The species composition of the coastal flat usually is similar to that of areas that lie above it, usually the lower edges of low salt marsh. Stands composed of any one of the following species often represent the low-lying edges of more densely covered marsh communities: Sea arrowgrass, Salt-marsh bulrush, Three-square bulrush, Soft-stemmed bulrush, and Spike rush. Sea arrowgrass and the bulrushes tend to grow in circular stands, while the Spike rush has a linear growth pattern, which is due largely to its rhizomatous nature. Sea arrowgrass tends to associate with silty substrates, while Pickleweeds and Spike rush are found on sandy substrates.

Coastal flats often support assemblages of macroscopic algae of much smaller size than those found in the saltwater aquatic wetland. Algal genera found on the flats include *Cladophora*, *Enteromorpha*, *Fucus*, and *Ulva*. Eelgrass (*Zostera* spp.) often is scattered among the algal colonies.

Saltwater coastal flats often are successional in nature, ultimately developing into saltwater marshes. In Washington, for example, tideflats often are colonized by the pioneers *Distichlis spicata*, *Salicornia virginica*, and *Triglochin maritima*. In Oregon the most typical pioneer invader of flats probably is *Triglochin maritima*, but other important primary colonists are *Salicornia virginica*,\* *Scirpus* spp., and *Carex lyngbyei*. *Spartina foliosa* is a pioneer in parts of California.

Transitional species. Coastal flats usually grade into salt marsh communities. There are few differences in the species composition between saltwater coastal flats and the lower edges of saltwater marshes; the two wetland types must be distinguished quantitatively on the basis of plant cover, flats having less cover than marshes.

At their lower edges coastal flats usually support no vascular plants. These intertidal areas commonly develop a sparse cover of filamentous algae during the summer months. The lower edge of this largely barren zone is adjacent to the saltwater aquatic wetland in many areas.

#### ENVIRONMENTAL CONDITIONS

55. Coastal flats commonly are exposed only a few hours at a time during the tidal cycle. Salinities in coastal flat substrates generally range from 10 to 34 ppt. Salinities in estuarine flats may be reduced to less than 10 ppt due to freshwater influence.

---

\* See note on page 36.



#### FIELD IDENTIFICATION

56. Coastal flats characteristically are areas of exposed sediments adjacent to the ocean. Plants on the flats often grow in circular patches and are widely scattered in distribution. During the winter months, coastal flats commonly have the exhumed tubers and eroded stubble of bulrush stems.

## SALTWATER MARSH

**Definition:** Wetlands that have more than 25 percent cover of herbaceous plants but 40 percent or less cover by woody plants and that are occasionally or regularly flooded by brackish or saline water

57. Saltwater marshes are scattered along coastal shorelines protected from excessive wave action, as around edges of bays, lagoons, and estuaries or behind islands and barrier spits. The most extensive salt marshes in the West Coastal States region occur in California; the largest tracts occur around San Francisco, Los Angeles, and San Diego. Barbour and Major (eds., 1977) estimated that approximately 36,000 hectares remain in California, this figure representing less than half of the original area once covered by salt marshes. MacDonald and Barbour (1974) estimated that Washington has about 4500 hectares of salt marsh, while Oregon has about 3040 hectares.

58. A major loss of salt marshes has occurred through diking in estuarine areas, which effectively eliminates surface water inundation. Interestingly, Johannessen (1964) has shown a dramatic expansion in the salt marshes of Oregon over the last 100 years; he attributed the expansion to increased rates of soil erosion and tideflat deposition, which followed the movement of European colonists into Oregon.

59. Salt marshes are considered highly productive systems, providing large amounts of organic detrital materials that may be washed into the adjacent saltwater coastal flats and saltwater aquatic communities. This detritus is an extremely important source of food for many aquatic animal species. Salt marshes also provide habitat for fish, shellfish, and waterfowl.

### VEGETATION

60. Growth forms and physiognomy: graminoids (grasses, sedges, and rushes) and succulent forbs, varying in height from low vegetation to stands of Bull tule that are almost 2 m tall; in sparse to dense stands.

61. Species composition of the saltwater marsh:



Figure 4. Generalized salt marsh profile (southern)

The reader is cautioned that the generalized floristic profiles contained within this guide are oversimplified and are not representative of many sites that will be found in the field. Wetland systems are dynamic, and many variations will be found. Species listed as "typical" on the profiles are those that generally occur as dominants in the particular wetland types. Those listed as "transitional" are those that regularly are associated with the transition zones at the margins of the individual wetland types. Associated species are those that are of common occurrence in a particular wetland type but generally are not sufficiently abundant to be dominants.






Figure 5. Generalized profile of northern low salt marsh




Figure 6. Generalized profile of northern high salt marsh

#### Dominant species

*Carex lyngbyei* (Lyngbye's sedge)  
*Deschampsia caespitosa* (Tufted hairgrass)  
*Distichlis spicata* (Saltgrass)  
*Frankenia grandifolia* (Frankenia)  
*Jaumea carnosa* (Jaumea)  
*Juncus balticus* (Baltic rush)  
*Salicornia* spp. (Glasswort)  
*Scirpus robustus* (Bull tule)  
*Scirpus validus* (Softstem bulrush)  
*Spartina* spp. (Cordgrass)  
*Triglochin maritima* (Sea arrowgrass)

#### Associated species

*Achilla millefolium* (Yarrow)  
*Agrostis alba* (Creeping bentgrass)  
*Atriplex patula* (Fat-hen)  
*Batis maritima* (Saltwort)  
*Cordylanthus maritimum* (Bird's beak)  
*Cuscuta salina* (Dodder)  
*Eleocharis macrostachya* (Creeping spikerush)  
*Glaux maritima* (Sea milkwort)  
*Grindelia* spp. (Gum plant)  
*Hordeum jubatum* (Foxtail barley)  
*Juncus effusus* (Common rush)  
*Limonium californicum* (Sea lavender)  
*Monanthochloe littoralis* (Shore grass)  
*Myosurus minimus* (Mousetail)  
*Oenanthe sarmentosa* (Oenanthe)  
*Plantago maritima* (Goosetongue)  
*Potentilla* spp. (Silverweed)  
*Scirpus californicus* (Three-square bulrush)  
*Spergularia canadensis* (Sand spurry)  
*Trifolium wormskjoldii* (Coast clover)

Dominant and associated species. The species composition of the salt marsh is highly variable; there is a trend toward increased species diversity from north to south within the West Coast region. Species of abundance throughout most of the region are *Salicornia* spp., *Distichlis spicata*, *Jaumea carnosa*, and *Triglochin maritima*. Local dominance by other species is affected by latitude and tidal elevation. Many of the coastal marshes show a distinct zonation of plant communities as one moves up the intertidal zone to higher elevations. In the more northerly marshes, a distinction can be made between low and high marsh, but in many respects the transition becomes more gradual to the south. MacDonald and Barbour (1974) and MacDonald (1977) distinguished an intermediate marsh type (dominated by *Salicornia bigelovii*, a low form of

*S. virginica*,\* *Batis maritima*, and *Frankenia grandifolia*) in more southerly portions of the region. At the same time, when referring to California marshes, MacDonald (1977) indicated that the existence or nonexistence of vegetation zones is largely a matter of definition. The zones that often are evident generally reflect mere changes in growth habit and most often do not reflect variation of community structure.

Marshes on the Pacific coast having relatively high salinities support a limited number of dominant halophyte species. MacDonald and Barbour (1974) said the number often is less than 10 and rarely as much as 20. At lower salinities, however, the number of species increases dramatically, often with the inclusion of many weedy or upland species.

Southern low marshes are characterized by Cordgrass (*Spartina* spp.), Pickleweed (*Salicornia* spp.), Saltwort (*Batis maritima*), and Sea arrowgrass (*Triglochin maritima*). Cordgrass often occurs as the sole dominant in dense stands; however, it is absent in many places and elsewhere often shares dominance with *Salicornia* spp. Northern low marshes are characterized by spike rushes (*Eleocharis* spp.), Sand spurry (*Spergularia canadensis*), *Salicornia* spp., Three-square bulrush (*Scirpus californicus*), and Sea arrowgrass (*Triglochin maritima*). Sandy substrates often support dense mats of *Salicornia* spp. intermixed with *Jaumea*, Three-square bulrush (*Scirpus californicus*), and Sea arrowgrass (*Triglochin maritima*). Sandy substrates often support Sea arrowgrass with occasional Sand spurry. These communities of the low marsh usually have less cover than in the high marsh communities.

Southern high marshes are characterized by Tufted hairgrass (*Deschampsia caespitosa*), Saltgrass (*Distichlis spicata*), Creeping bentgrass (*Agrostis alba*), Lyngbye's sedge (*Carex lyngbyei*), *Frankenia*, Goosetongue (*Plantago maritima*), Shore grass (*Monanthochloe littoralis*), and Dodder (*Cuscuta salina*). In the extreme southern part of the region near San Diego, the following species are also frequent: *Atriplex patula*, *Cordylanthus maritimus*, and *Eleocharis macrostachya*. South of the Santa Barbara-Goleta area, *Salicornia subterminalis* and *Batis maritima* are important in this zone. Northern high marshes are characterized by Lyngbye's sedge (*Carex lyngbyei*), Tufted hairgrass (*Deschampsia caespitosa*), Creeping bentgrass (*Agrostis alba*),

---

\* Some authorities do not consider the West Coast plants conspecific with *S. virginica* of the Atlantic Coast, in which case the Pacific Coast populations are called *S. pacifica* Standley.



*Potentilla* spp., Saltgrass (*Distichlis spicata*), and Gum plants (*Grindelia* spp.). Yarrow (*Achillea millefolium*) is occasional. Areas of brackish nature are characterized by *Scirpus* spp. and Common cattail (*Typha latifolia*) along with Lyngbye's sedge (*Carex lyngbyei*).

Once formed coastal saltwater marshes appear to be long persisting and successional stable.

Transitional species. Sitka spruce (*Picea sitchensis*) often dominates the zone above the high saltwater marshes in Oregon and Washington. Sitka spruce is tolerant of salt spray and occasional inundation by extreme high tides. *Agrostis alba*, *Grindelia* spp., *Potentilla* spp., *Plantago maritima*, and *Oenothera sarmentosa* also extend into the transition zone. If the coastal marsh is backed by a freshwater marsh, Common tule (*Scirpus acutus*) and Common cattail (*Typha latifolia*) often occur in the transition zone.

Coastal saltwater marshes in California are backed by somewhat different plant communities than is the case further north. The transition zone behind marshes in the San Francisco area includes *Cordylanthus maritimus*, *Atriplex patula*, *Potentilla* spp., and *Frankenia*. These species are in the salt marsh itself, along with Sea rocket (*Cakile maritima*), Brass buttons (*Cotula coronopifolia*), and Mexican tea (*Chenopodium ambrosioides*). Sitka spruce extends south only into northern California.

The lower limit of the salt marsh normally is easily delineated; under most circumstances salt marsh on the West Coast establishes at about mean lower high water (MLHW), the average height of the lower of the two unequal high tides occurring daily under the mixed-tide regime. The upper boundary of the salt marsh often is hard to distinguish, but often it is at extreme high water (EHW). At many localities EHW corresponds to an abrupt change of slope and a line of stranded logs, algae, and other debris deposited by high tidal waters. Logs and various other forms of debris, therefore, often characterize the transition zones of many marshes along the coastline, particularly in Washington and Oregon. The logs and other materials often serve as islands on which plant species more typical of the uplands become established. When such islands are lodged out in the marsh itself, they should not be confused with the actual transition zone.

#### ENVIRONMENTAL CONDITIONS

62. Salinity of soils in coastal marshes decreases along an elevational gradient from that of sea water (34 ppt) to only slightly saline at points where freshwater flow dilutes the sea water. Salinity

levels often are significantly reduced during the rainy season. Low marshes generally are exposed for fewer than 100 diurnal periods per year, while high marshes are exposed more frequently. High marshes are exposed by 90 to 95 percent of the low tides, while low marshes are exposed to 75 to 80 percent of the low tides.

#### FIELD IDENTIFICATION

63. Coastal saltwater marshes are identified readily by their usual location adjacent to the ocean or brackish waters and by the dominance of herbaceous graminoid plants. The forbs that are present often are succulents. Saltwater marshes are found in the upper intertidal zone and usually have a dense cover of vegetation.

## SALTWATER SWAMP

**Definition:** Wetlands that have more than 40 percent cover by woody plants and that are occasionally or regularly flooded by brackish or saline water

64. There are few saltwater swamps in the West Coast region. Most of these swamps are found on islands in the delta region of California. Saltwater swamps also form a fringing edge to saltwater marshes in Oregon and Washington. The swamps provide an important nesting habitat for birds that feed in the adjacent marshes.

### VEGETATION

65. Growth forms and physiognomy: moderately dense to dense stands of deciduous trees, to 9 or 12 m tall; usually with a dense understory of shrubs and herbs.

66. Species composition of the saltwater swamp:




Figure 7. Generalized profile of California willow saltwater swamp



#### Dominant species

*Alnus oregana* (Red alder)  
*Cornus stolonifera* (Red osier dogwood)  
*Salix* spp. (Willow)

#### Associated species

*Athyrium filix-femina* (Lady fern)  
*Carex obnupta* (Slough sedge)  
*Cephalanthus occidentalis* (Buttonbush)  
*Equisetum* spp. (Horsetail)  
*Hibiscus californicus* (California hibiscus)  
*Hydrocotyle verticillata* (Marsh pennywort)  
*Limosella subulata* (Mudwort)  
*Lonicera involucrata* (Twinberry)  
*Phragmites communis* (Common reed)  
*Potentilla egedii* (Pacific silverweed)  
*Samolus parviflorus* (Water pimpernel)  
*Scirpus acutus* (Tule)  
*Scirpus californicus* (California bulrush)  
*Scirpus olneyi* (Olney's bulrush)  
*Scirpus americanus* (Three-square bulrush)  
*Tillaea aquatica* (Water stonecrop)  
*Typha angustifolia* (Narrow-leaf cattail)  
*Typha latifolia* (Common cattail)

Dominant and associated species. Saltwater swamps in the southern part of the region are dominated by Goodding's willow (*Salix gooddingii*) and White willow (*Salix lasiolepis*), usually with an occasional Sandbar willow (*Salix interior*). California hibiscus (*Hibiscus californicus*) and Twinberry (*Lonicera involucrata*) form a shrub layer with Lady fern (*Athyrium filix-femina*) often scattered in the stands. Bulrushes (*Scirpus* spp.) and other herbs are found either scattered through the stands or in dense patches.

Swamps in the northern part of the region usually are dominated by willows (*Salix scouleriana*, *S. lasiandra*, and *S. hookeriana*); Red alder (*Alnus oregana*) and Red osier dogwood (*Cornus stolonifera*) also may be important species. The understory consists of Horsetail (*Equisetum* spp.), Slough sedge (*Carex obnupta*), Lyngbye's sedge (*Carex lyngbyei*), Creeping bentgrass (*Agrostis alba*), Pacific silverweed (*Potentilla egedii*), Cattail (*Typha* spp.), and various Bulrush species (*Scirpus* spp.). These swamps are found as fringes to the lower saltwater marsh communities. Sitka spruce (*Picea sitchensis*) is found occasionally in these areas and usually has a buttressed base when growing in the swamp.

Transitional species. The transition zones between swamp and adjacent upland communities in the southern part of the

region usually include scattered willows, bulrushes, cattails, and California hibiscus. In the more northern areas, the transitions usually contain Red alder, Sitka spruce, and hemlock.

#### ENVIRONMENTAL CONDITIONS

67. West Coast saltwater swamps differ from those in many other regions, such as peninsular Florida where the plants are often subject to regular tidal flooding. On the West Coast, the saltwater swamps are flooded by tides on an irregular basis; consequently, the salinity values often are reduced considerably below those of sea water (34 ppt).

#### FIELD IDENTIFICATION

68. Saltwater swamps can be recognized by the usual dominance of willows and by their location adjacent to the ocean or to saltwater marshes.

## FRESHWATER AQUATIC WETLAND

**Definition:** Wetlands that are dominated by free-floating or rooted aquatic herbs and are semipermanently or permanently flooded by fresh water

69. Freshwater aquatic wetland communities are scattered throughout the region in ponds and along the shorelines of large lakes, rivers, and streams. Typically they are found in association with bodies of open water in which the vegetation cover is low. The plants found in the open water areas are the same as those found in the freshwater aquatic type; emphasis is placed here on the structure and composition of communities in which plant cover is relatively high. Freshwater aquatic wetlands provide food and shelter for most freshwater sport and commercial fish, for waterfowl, and for furbearers.

### VEGETATION

70. Growth forms and physiognomy: free-floating herbs, such as Water fern, and rooted aquatic herbs, such as Yellow water lily; in open to dense, though sometimes scattered, stands.

71. Species composition of the freshwater aquatic wetland:




Figure 8. Generalized profile of a freshwater aquatic wetland



#### Dominant species

*Azolla filiculoides* (Water fern)  
*Nuphar polysepalum* (Yellow water lily)  
*Potamogeton pectinatus* (Sago pondweed)  
*Potamogeton robbinsii* (Pondweed)

#### Associated species

*Callitriche* spp. (Water starwort)  
*Ceratophyllum demersum* (Hornwort, Coontail)  
*Chara* spp. (Stonewort)  
*Elodea canadensis* (Waterweed)  
*Elodea nuttallii* (Waterweed)  
*Lemna perpusilla* (Duckweed)  
*Lemna trisulca* (Duckweed)  
*Myriophyllum spicatum* (Water milfoil)  
*Najas* spp. (Water nymph)  
*Nitella* spp. (Green algae), commonly called Stonewort  
*Nymphaea odorata* (White water lily)  
*Potamogeton berchtoldii* (Pondweed)  
*Potamogeton epihydrus* (Pondweed)  
*Potamogeton richardsonii* (Pondweed)

Dominant and associated species. The deepest portions of the freshwater aquatic wetland in many cases are dominated by Sago pondweed (*Potamogeton pectinatus*) and Hornwort (*Ceratophyllum demersum*); these species are completely submerged. In more shallow areas, Yellow water lily (*Nuphar polysepalum*) and various pondweeds are most important. The leaves of *Nuphar* float on the water surface, while those of the pondweeds are mostly submerged. Water fern (*Azolla filiculoides*) and the duckweeds, all of which are free-floating, are scattered among the other plants in quiet areas. The floating *Azolla* and *Lemna* plants often are moved from one part of a water body to another by wind or currents and frequently form small dense stands to the total exclusion of attached plant species.

Freshwater aquatic communities can be successional stable where stream flow is minimal, but succession to freshwater marsh can be rapid if the lake or pond is filling with sediment. Usually the earliest invaders from the marsh are Tule (*Scirpus acutus*), Burreed (*Sparganium* spp.), Common cattail (*Typha latifolia*), Creeping spike rush (*Eleocharis macrostachya*), Reed canarygrass (*Phalaris arundinacea*), or *Spiraea* (*Spiraea douglasii*).

Transitional species. Normally the transition is to freshwater marsh in the direction of drier or more shallow areas; the species found in the transition zone usually are the same ones that tend to invade the freshwater aquatic areas (*Scirpus acutus*, *Sparganium* spp., *Typha latifolia*, etc.).

The freshwater aquatic community rarely adjoins uplands unless the intervening marsh has been destroyed. When in contact with deep open water, the transition from freshwater aquatic wetland to open water usually is a gradual diminution in plant cover with essentially no change in species composition.

#### ENVIRONMENTAL CONDITIONS

72. The freshwater aquatic community usually is flooded permanently or nearly permanently; if it becomes dry for any reason, it is invaded rapidly by freshwater marsh species. The outer limit of the freshwater aquatic wetland is the limit of rooted vascular plant growth, and the water depth at that point rarely exceeds three metres. The freshwater aquatic wetland appears to be influenced by degree of water hardness: the more solutes in the water, the denser the vegetation.

#### FIELD IDENTIFICATION

73. The freshwater aquatic community is dominated by herbaceous species, which distinguishes it from the freshwater swamp. The freshwater aquatic community is dominated by both rooted and free-floating aquatic plant species, which separates it from freshwater marshes, in which many nonaquatic species grow and in which the free-floating species (such as *Nymphaea* spp.) normally are not dominant.

## FRESHWATER FLAT

**Definition:** Wetlands that have 25 percent or less vegetative cover and are occasionally or regularly flooded by fresh water

74. Freshwater flats are most prevalent around large reservoirs where water levels fluctuate greatly during the year. They also occur on stream and river margins, gravel bars, and sand bars. Some flats are important feeding areas for shorebirds that eat invertebrates burrowing in the mudflats. Few vascular plants grow on flats, but the flats can have surprisingly high primary production rates, due to the algae that grow in the mudflats and on rocky shores.

### VEGETATION

75. Growth forms and physiognomy: barren or with scattered herbaceous plants and often with algae.

76. Species composition of the freshwater flat (potentially dominant species):

*Echinochloa crus-galli* (Barnyard grass)

*Equisetum* spp. (Horsetail)

*Juncus effusus* (Common rush)

*Nasturtium officinale* (Water cress)

*Polygonum hydropiper* (Marsh pepper)

*Rumex crispus* (Curly dock)

*Salix* spp. (Willow)

*Xanthium* spp. (Cocklebur)

*Cladophora* spp. (Green algae)

*Lyngbya* spp. (Blue-green algae)

*Mougeotiopsis* spp. (Green algae)

*Protococcus* spp. (Green algae)

*Oscillatoria* spp. (Blue-green algae)

Dominant and associated species. Many of the vascular plants found on freshwater flats occur as scattered individuals and, though of a competitive weedy nature, seldom persist through high water periods. Woody species, such as Willow (*Salix* spp.), occur as seedlings scattered in the lower edges of bars. Algae often occur in zones that probably are determined by flooding frequency.

Transitional species. Freshwater flats usually border open water at their lower edge. At their higher edge they may meet marshes, swamps, or nonwetland communities. The bound-



ary in any case can be determined by evaluation of the plant cover. If the site has 25 percent or less cover, it is an inland freshwater flat.

#### ENVIRONMENTAL CONDITONS

77. Inland freshwater flats commonly are inundated throughout high water periods and normally are exposed during periods of low water. The soils are saturated most of the year. Soil types vary from silt and clay to coarse sand and gravel. High water periods are significant to both erosion and accumulation of soils.

#### FIELD IDENTIFICATION

78. Freshwater flats may be identified by the extensive exposed substrates with a sparse vegetative cover. Frequently the exposed substrate surface is littered with remains of diatoms (a group of unicellular algae that have silica shells or frustules that are long persistent).

## FRESHWATER MARSH

**Definition:** Wetlands that have more than 25 percent vegetative cover of herbaceous plants but 40 percent or less cover by woody plants and that are occasionally or regularly flooded by fresh water

79. Freshwater marshes occur throughout the West Coast region. In coastal areas they occur as deflation plain marshes in dune landscapes, as coastal fens, or as lake margin marshes. In the Central Valley, freshwater marsh wetlands are found adjacent to lakes and rivers in the form of shallow depressions in alluvial topography. In the mountainous areas, freshwater wetlands take the form of seasonally wet meadows and fens.

80. Many common names have been applied to various types of marshes based upon whether their soils are mineral or organic (peat), whether they receive nutrients only from precipitation (ombrotrophic) or from groundwater as well (minerotrophic), and other criteria relating to drainage patterns. For this guidebook series, all of these variations are considered freshwater marshes on the basis of herbaceous plant domination. The West Coast region exhibits the four distinct subtypes given below, which are based primarily upon soil type (e.g., mineral or organic), the nature of the water table (e.g., stable or fluctuating), whether the plants are annual or perennial, and altitude.

- a. Coastal deflation plain marsh: low altitude marshes with sandy (mineral) soil and a shallow, stable water table; plants perennial
- b. Riparian and lacustrine marsh: low to medium altitude marshes with mineral soil and a fluctuating water table; plants perennial
- c. Wet meadow and fen: high altitude marshes with mineral soil (wet meadows) or peat soil (fens) and a seasonally falling water table; plants perennial
- d. Vernal pool: medium altitude marshes with mineral soil and a rapidly falling water table; plants annual

## VEGETATION

81. Growth forms and physiognomy: usually open to dense stands of

graminoids (grasses and sedges) to 1.8 m tall; occasionally dominated by forbs such as Yellow water lily (*Nuphar polysepalum*).

82. Species composition of the freshwater marsh:

a. Coastal deflation plain:

Dominant species

*Beckmannia syzigachne* (Slough grass)  
*Carex hindsii* (Hind's sedge)  
*Carex obnupta* (Slough sedge)  
*Carex senta* (Rough sedge)  
*Eleocharis macrostachya* (Creeping spikerush)  
*Hypericum anagalloides* (Bog St. John's wort)  
*Juncus acuminatus* (Small-fruited rush)  
*Lilaeopsis occidentalis* (Lilaeopsis)  
*Ludwigia palustris* (Water purslane)  
*Lycopus uniflorus* (Water horehound)  
*Paspalum distichum* (Knotgrass)  
*Potentilla anserina* (Silverweed)  
*Ranunculus flammula* (Creeping buttercup)  
*Scirpus* spp. (Bulrush)  
*Typha angustifolia* (Narrow-leaf cattail)  
*Typha domingensis* (Cattail)  
*Typha latifolia* (Common cattail)  
*Veronica scutellaria* (Marsh speedwell)

Associated species

*Alisma trivialis* (Water plantain)  
*Epilobium adenocaulon* (Willow herb)  
*Epilobium watsonii* (Watson's willow herb)  
*Galium triflorum* (Bedstraw)  
*Holcus lanatus* (Velvet grass)  
*Juncus bolanderi* (Bolander's rush)  
*Juncus leseurii* (Salt-rush)  
*Jussiaea repens* (Creeping water primrose)  
*Mimulus guttatus* (Monkey flower)  
*Nuphar polysepalum* (Yellow water lily)  
*Oenanthe sarmentosa* (Oenanthe)  
*Polygonum hydropiperoides* (Water pepper)  
*Polygonum punctatum* (Perennial smartweed)  
*Potentilla egedii* (Pacific silverweed)  
*Sagittaria* spp. (Arrowhead)  
*Sparganium eurycarpum* (Bur-reed)



b. Riparian and lacustrine marsh:




Figure 9. Generalized profile of a freshwater riparian or lacustrine marsh

Dominant species

*Deschampsia caespitosa* (Tufted hairgrass)  
*Eleocharis* spp. (Spike rush)  
*Equisetum fluviatile* (Water horsetail)  
*Juncus bufonius* (Toad rush)  
*Juncus supiniiformis* (Pointed rush)  
*Nuphar polysepalum* (Yellow water lily)  
*Nymphaea odorata* (White water lily)  
*Paspalum distichum* (Knotgrass)  
*Phalaris arundinacea* (Reed canarygrass)  
*Potentilla palustris* (Marsh cinquefoil)  
*Sagittaria latifolia* (Wapato)  
*Scirpus acutus* (Tule)  
*Scirpus californicus* (California bulrush)  
*Scirpus validus* (Softstem bulrush)  
*Typha latifolia* (Common cattail)

Associated species

*Alisma* spp. (Water plantain)  
*Bidens cernua* (Nodding beggar-tick)  
*Bergia texana* (Bergia)  
*Capsella bursa-pastoris* (Shepherd's purse)  
*Eichhornia crassipes* (Water hyacinth)  
*Epilobium watsonii* (Watson's willow herb)

*Limosella aquatica* (Mudwort)  
*Mimulus guttatus* (Monkey flower)  
*Polygonum coquimbense* (Water smartweed)

c. Wet meadow and fen:




Figure 10. Generalized profile of a freshwater (wet meadow and fen) marsh

Dominant species

*Aconitum columbianum* (Monkshood)  
*Calamagrostis nutkaensis* (Pacific needle reedgrass)  
*Caltha biflora* (Broad-leaved caltha)  
*Carex aquatilis* (Water sedge)  
*Carex exserta* (Short-hair sedge)  
*Carex nigricans* (Sedge)  
*Carex rostrata* (Beaked sedge)  
*Carex sitchensis* (Sitka sedge)  
*Carex vesicaria* (Inflated sedge)  
*Deschampsia caespitosa* (Tufted hairgrass)  
*Elymus hirsutus* (Wild rye)  
*Epilobium latifolium* (Willow herb)  
*Mimulus lewisii* (Monkey flower)

Associated species

*Carex* spp. (Sedge)  
*Eleocharis pauciflora* (Spike rush)  
*Eriophorum* spp. (Cottongrass)  
*Ledum columbianum* (Pacific Labrador tea)

*Lysichiton americanum* (Skunk cabbage)  
*Menyanthes trifoliata* (Buckbean)  
*Nuphar polysepalum* (Yellow water lily)  
*Pedicularis groenlandica* (Elephant's head)  
*Senecio triangularis* (Groundsel)

d. Vernal pool:




Figure 11. Generalized profile of a freshwater (vernal pool) marsh

Dominant species

*Callitriche* spp. (Water starwort)  
*Downingia cuspidata* (Downingia)  
*Eleocharis acicularis* (Needle spikerush)  
*Eleocharis obtusa* (Blunt spikerush)  
*Elodea* spp. (Waterweed)  
*Eryngium aristulatum* (Button snakeroot)  
*Isoetes howellii* (Quillwort)  
*Ludwigia palustris* (Water purslane)  
*Pilularia americana* (Pillwort)  
*Plagiobotrys* spp. (Popcorn flower)  
*Pogogyne abramsii* (Mesa mint)  
*Ranunculus aquatilis* (Aquatic buttercup)  
*Veronica peregrina* (Speedwell)

Associated species

*Alopecurus saccoatus* (Water foxtail)  
*Deschampsia danthonioides* (Annual hairgrass)  
*Eleocharis macrostachya* (Creeping spikerush)  
*Juncus bufonius* (Toad rush)  
*Juncus dubius* (Mariposa rush)  
*Juncus triformis* (Dwarf rush)  
*Lilaea scilloides* (Flowering quillwort)  
*Myosurus minimus* (Mousetail)  
*Phalaris lemmonii* (Canarygrass)



Dominant and associated species. Deflation plain marshes, as well as several other wetland communities in this coastal region, commonly support Slough sedge (*Carex obnupta*) and Pacific silverweed (*Potentilla egedii*); *Juncus leseurii* is a common associate. Coastal deflation plain marshes in the southern part of the region support cattails, Rough sedge (*Carex senta*), Bolander's rush (*Juncus bolanderi*), and Salt-rush (*Juncus leseurii*), as well as Velvet grass (*Holcus lanatus*) and *Potentilla* spp. Bur-reed (*Sparganium* spp.) and *Ceanothe* occur with Bulrush (*Scirpus* spp.) in deeper waters.

The most common species in the riparian (stream/river) and lacustrine (pond/lake) marshes are spikerushes (*Eleocharis* spp.), Common tule (*Scirpus acutus*), Wapato (*Sagittaria latifolia*), Softstem bulrush (*Scirpus validus*), Reed canarygrass (*Phalaris arundinacea*), California bulrush (*Scirpus californicus*), Common cattail (*Typha latifolia*), and Knotgrass (*Paspalum distichum*). These marshes occur in areas of slowly moving or slack water with silty bottoms. Although most of the species are rooted, the marshes also support floating species such as Duckweed (*Lemna* spp.) and Water fern (*Asolla* spp.).

Wet meadows and fens are cool montane wetlands typical of areas with peat soil (fens) or mineral soil (wet meadows). Wet meadows are the driest of the freshwater marshes and contain several species that cannot withstand permanently saturated soils. The species composition of wet meadows is affected strongly by local environmental conditions; because it affects snow depth and duration, slope exposure probably is the most influential factor of the microenvironment. Northern valerian, for example, dominates on steep slopes that remain wet from receding snowbanks late in the year; Sitka sedge dominates sites inundated up to three weeks by snow-melt waters. Holm's Rocky Mountain sedge (*Carex scopulorum*) is important, along with Broad-leaved caltha and *Carex lasulina*, on other snow-melt areas that are saturated but not inundated. Wet meadows in the mountainous areas have not been studied thoroughly. They often are associated with mountain streams.

Fens in the West Coast region commonly intergrade with wet meadows. The distinction between fen and wet meadow is based on soil type, whether mineral (wet meadow) or organic (fen), but many areas intergrade. Various sedges (*Carex* spp.) are often dominant; Needle spikerush (*Eleocharis acicularis*) may

share dominance with the sedges to the near exclusion of all other species. Areas with deeper water are characterized by coarse sedges (*Carex nebraskensis* and *C. rostrata*, *Eleocharis pauciflora*, and *Eriophorum* spp.) as the dominant graminoids. Elephant's head (*Pedicularis groenlandica*) and Buckbean (*Menyanthes trifoliata*) are characteristic herbs of fens.

Vernal pools are small, hardpan-floored depressions in grasslands of the Central Valley of California. They once were abundant but now are diminishing rapidly because of urbanization, grazing, and other factors of disturbance. The vegetation in vernal pools changes seasonally with pools filling up with water in the winter and drying out as the spring progresses. The sequence may progress, for example, from *Callitriche* to *Isoetes* and *Pilularia*. As the latter two mature, *Plagiobotrys* develops flowers. The following species develop leaves and flowers above the water surface: *Plagiobotrys acanthocarpus*, *P. bracteatus*, *P. leptocladus*, and *Veronica scutellaria*. Mesa mint (*Pogogyne abramsii*), which grows during the same period, does not flower until the soil dries. Button snakeroot (*Eryngium aristulatum*), also late maturing, completes its life cycle on parched cracking soil. It is difficult to specify the dominants of vernal pools because of these seasonal changes. Usually all of the plants of the pools have died by the end of summer, leaving an unvegetated area with scattered remnants of earlier growth.

Most freshwater marshes are relatively stable successional. As the landscape gradually changes, so does the vegetation change. For example, as ponds and lakes accumulate silt, the marsh communities are replaced by upland communities, but this process is usually slow unless erosion is accelerated. Wet meadows at times succeed to drier communities as the soil surface becomes raised and the soils dry. Similarly, fens may accumulate sufficient organic matter and mineral soils to become wet meadow, but this process is very slow. Fens also may be replaced by bog vegetation. Vernal pools may be invaded during dry years by more upland species, but these species cannot persist in this habitat. Consequently, this type of marsh is also stable.

Transitional species. Coastal deflation plain marshes are invaded frequently by willows (*Salix* spp.) and Lodgepole pine (*Pinus contorta*), forming a freshwater swamp. Drier areas adjoining deflation plain marshes contain thick stands of Spring bank clover

(*Trifolium willdenovii*) and Brown-headed rush (*Juncus phaeocephalus*).

Riparian and lacustrine marshes usually occupy a zone between freshwater submerged communities and freshwater swamp communities. These marshes generally form a stable community that lies in an intermediate moisture regime between freshwater aquatic and swamp communities. The communities commonly are invaded by willows (*Salix* spp.) as sedimentation raises the surface above the water table.

Wet meadow and fen communities frequently are surrounded by upland forests or by willows. Alpine wet meadows commonly grade into upland or to barren fell-field or lichen-covered rockfields. Freshwater swamps dominated by willow (*Salix* spp.), Red alder (*Alnus oregana*), or conifers commonly border these types of marshes.

The outermost zone of vernal pools is characterized by *Plagiobotrys* spp. and *Deschampsia danthonioides*. Grasses from surrounding areas mix with these species in the driest zones.

#### ENVIRONMENTAL CONDITIONS

83. The water regime in freshwater marshes varies from permanent inundation in most riparian and lacustrine marshes; to permanently saturated in most fens; to intermittently saturated in the drier wet meadows; to seasonally saturated in deflation plains and vernal pools. The water is slightly acid to alkaline.

84. The soil in deflation plain marshes is sandy and is wet most of the year. The water table fluctuates seasonally with standing water present during the winter rainy season and the water table below the surface during the summer dry season.

85. Riparian and lacustrine wetlands are primarily on wet mineral soil composed of fine sand and silt grading into highly organic muck. These freshwater marsh types are influenced directly by the water level of the adjacent water bodies. Riparian and especially lacustrine wetlands generally have a more stable water table than other freshwater marshes and commonly have standing water throughout the year.

86. Fens and wet meadows on the West Coast generally develop in basins or alluvial valleys. Wet meadows are underlain by sandy loam soil with a definite gley zone. These mineral-soil wetlands have a



fluctuating water table that usually has an oxidized surface and a reduced subsurface soil environment. Fens intergrade with wet meadows and have peat soil with a high calcium/magnesium ratio. These wetlands are mineral rich and commonly have a flow of water through the peat year round.

87. Vernal pools are seasonally flooded depressions that fill rapidly with winter and early spring rains. They usually are dry by May or June due to evaporation, not drainage. The water usually carries little sediment. Accumulation of organic matter frequently results in acidic pools, usually pH 6 to 7, but more acidic conditions are found; occasionally the pools are alkaline. The soil surface is stony with fine particles and is underlain by impermeable clay hardpans.

#### FIELD IDENTIFICATION

88. Freshwater marshes in the West Coast region are surrounded frequently by either swamps or upland forests. Consequently, the dominance of herbaceous plants is the primary key to the identification of marshes. Coastal deflation plain marshes are located on the windward sides of coastal sand dunes. Riparian and lacustrine marshes are characterized by their landscape position (being adjacent to water bodies). Fens and wet meadows characteristically are mountainous wetlands that lie in local depressions. Vernal pools are typically bare and stony in the winter and dominated by annual plants in the spring and summer months.

## FRESHWATER SWAMP

**Definition:** Wetlands that have more than 40 percent cover by woody plants and are occasionally or regularly flooded by fresh water

89. Freshwater swamps are most common along river courses in the West Coast region. Montane and coastal freshwater swamps occupy local topographic depressions and commonly are associated with freshwater marsh wetlands. These swamps generally occupy either the transition from marsh to upland or the transition from open water to upland.

90. Three major variants of swamp vegetation have been identified in the region:

- a. Coastal deflation plain swamp: swamps on wet sandy soil of dune landscapes.
- b. Riparian and lacustrine swamp: swamps on mineral soil associated with river or lake floodplains.
- c. Montane swamp: swamps occurring in local depressions of upland situations.

91. Riparian and lacustrine swamps are the most common type of freshwater swamp in the region. Freshwater swamps are important in providing wildlife habitat for a variety of birds, mammals, amphibians, and reptiles. These wetlands also contribute organic detritus to adjacent waterways.

### VEGETATION

92. Growth forms and physiognomy: open to dense stands of needle-leaf trees, such as Western red cedar, and deciduous trees, such as Willow; usually with broadleaf evergreen shrubs, particularly Huckleberry, in the understory.

93. Species composition of the freshwater swamp wetland:

- a. Coastal deflation plain swamp

#### Dominant species

*Gaultheria shallon* (Salal)

*Myrica californica* (Wax myrtle)

\**Pinus contorta* (Beach pine)

---

\* *Pinus contorta* var. *contorta* (Beach pine) is the geographical variety found primarily in coastal regions.

*Salix hookeriana* (Hooker's willow)  
*Vaccinium ovatum* (California huckleberry)

Associated species

*Alnus oregana* (Red alder)  
*Carex obnupta* (Slough sedge)  
*Fragaria chiloensis* (Coast strawberry)  
*Juncus leseurii* (Salt rush)  
*Juncus phaeocephalus* (Brown-headed rush)  
*Picea sitchensis* (Sitka spruce)  
*Spiraea douglasii* (Spiraea)

b. Riparian and lacustrine swamp




Figure 12. Generalized profile of freshwater riparian swamp

Dominant species

*Acer negundo* (Box elder)  
*Alnus oregana* (Red alder)  
*Crataegus douglasii* (Black hawthorn)  
*Fraxinus latifolia* (Oregon ash)  
*Platanus racemosa* (Sycamore)  
*Populus fremontii* (Cottonwood)  
*Populus trichocarpa* (Black cottonwood)  
*Salix* spp. (Willow)

Associated species

*Agrostis alba* (Creeping bentgrass)



*Baccharis pilularis* (Coyote bush)  
*Cephalanthus occidentalis* (Buttonbush)  
*Cornus stolonifera* (Red-osier dogwood)  
*Deschampsia caespitosa* (Tufted hairgrass)  
*Equisetum* spp. (Horsetail)  
*Glechoma hederacea* (Ground ivy)  
*Iris pseudacorus* (Yellow flag)  
*Lysichiton americanum* (Skunk cabbage)  
*Lysimachia nummularia* (Moneywort)  
*Phalaris arundinacea* (Reed canarygrass)  
*Rubus* spp. (Blackberry)  
*Sambucus callicarpa* (Red elderberry)  
*Urtica dioica* (Stinging nettle)  
*Vitis californica* (Wild grape)

c. Montane swamp




Figure 13. Generalized profile of freshwater montane swamp

Dominant species

*Alnus oregana* (Red alder)  
*Empetrum nigrum* (Crowberry)  
*Kalmia polifolia* (Mountain laurel)  
 \**Pinus contorta* (Lodgepole pine)  
*Rhododendron californicum* (California rose-bay)  
*Salix pedicellaris* (Bog willow)

\* *Pinus contorta* var. *murrayana* (Lodgepole pine) occurs primarily in interior montane situations.

*Thuja plicata* (Western red cedar)  
*Vaccinium oxycoccus* (Cranberry)

Associated species

*Alnus sinuata* (Sitka alder)  
*Athyrium filix-femina* (Lady fern)  
*Betula occidentalis* (Water birch)  
*Carex obnupta* (Slough sedge)  
*Carex* spp. (Sedge)  
*Gaultheria shallon* (Salal)  
*Lysichiton americanum* (Skunk cabbage)  
*Menyanthes trifoliata* (Buckbean)  
*Myrica californica* (Wax myrtle)  
*Oplopanax horridum* (Devil's club)  
*Polypogon monspeliensis* (Rabbit-foot grass)  
*Potentilla egedii* (Pacific silverweed)  
*Spiraea* spp. (Spiraea)  
*Vaccinium ovatum* (California huckleberry)

Dominant and associated species. Deflation plain swamps are characterized by a dense stand of Hooker's willow (*Salix hookeriana*) and Wax myrtle (*Myrica californica*). Beach pine (*Pinus contorta*) commonly dominates older deflation plain swamps. The insectivorous sundews (*Drosera* spp.) and Pitcher plants (*Darlingtonia californica*) occur locally in coastal swamps with organic soils.

Riparian and lacustrine swamps of the West Coast region are characterized by varying mixtures of a relatively few species. Willow (*Salix* spp.) swamps, cottonwood (*Populus* spp.) forests, and alder (*Alnus* spp.) and ash (*Fraxinus latifolia*) forests are all characteristic riparian swamp wetland communities in the region. Commonly there is a gradient from willows, adjacent to standing or flowing water, through ash to cottonwood, which usually is at slightly higher elevations that are inundated less frequently. These stands often have Reed canarygrass (*Phalaris arundinacea*) as the sole species in the understory. Blackberry (*Rubus* spp.), Red elderberry (*Sambucus callicarpa*), and Red-osier dogwood (*Cornus stolonifera*) are common understory shrubs.

Small streams characteristically have a riparian forest of Red alder (*Alnus oregana*) and Scouler's willow (*Salix scouleriana*). Typical understory species in these forests are Creeping bentgrass (*Agrostis alba*) and some weedy grass and forb species.

Riparian forest wetland in the southern portion of the region is dominated by cottonwood (*Populus* spp.), Sycamore (*Platanus racemosa*), willows (*Salix* spp.), and Box elder (*Acer negundo*).

Montane swamps are isolated wetlands either closely associated with streams or occurring as local depressions in the landscape, collecting and storing run-off waters. These wetlands have not been well described for the West Coast region. Commonly they are dominated by Western red cedar (*Thuja plicata*), Red alder (*Alnus oregana*), and Willow (*Salix* spp.) with Salal (*Gaultheria shallon*) and California huckleberry as dominant understory shrubs. Skunk cabbage (*Lysichiton americanum*), Slough sedge (*Carex obnupta*), and Lady fern (*Athyrium filix-femina*) are the most common herbs. Most of the trees associated with montane swamps are not obligate wetland species.

Successional trends. Coastal deflation plain swamps have been described as succeeding to forests of spruce or pine. Riparian forests are successional stable except as river geomorphology slowly changes. The montane swamps may gradually be invaded and replaced by Red alder (*Alnus oregana*), Western hemlock (*Tsuga heterophylla*), Sword fern (*Polystichum munitum*), and Oregon oxalis (*Oxalis oregana*).

Transitional species. Coastal deflation plain swamps border marshes or adjacent upland plant communities. The riparian community may border streams and rivers directly or may be separated by a narrow band of freshwater marsh or freshwater flat. The border between swamps and other wetlands usually is distinct, because the other wetland types have little, if any, cover by woody plant species. The transition to upland usually is to other forest types that lack the typically riparian species. The transition of montane swamps to upland communities is usually abrupt with sharp distinctions between the wetland and adjacent conifer forests.

#### ENVIRONMENTAL CONDITIONS

94. Deflation plain swamps occur on dune landscapes where the sand has been blown adjacent to the local water table. These wetlands commonly are flooded in the winter and dry during the summer; the water table, however, seldom is more than 0.3 m below the surface.

95. Riparian and lacustrine swamps are characterized by silty to sandy alluvial soil and are flooded seasonally. These wetlands generally occur around stream, river, pond, or lake margins; the water table seldom falls substantially below the surface.

96. Montane swamps grow in saturated soil and occasionally have standing water. The soils of the montane swamps can be either mineral or organic (peat).



#### FIELD IDENTIFICATION

97. Swamps can be separated from other wetland types by the dominance of trees. The distinction between swamps and upland forests is made on the basis of species composition. Oregon ash (*Fraxinus latifolia*), Black cottonwood (*Populus trichocarpa*), willow (*Salix* spp.), Red alder (*Alnus oregana*), or Western red cedar (*Thuja plicata*) usually identify sites as freshwater swamps. The composition of the understory also should be used to identify freshwater swamps, as several of the dominants are indicative of the wetland habitat.

#### SELECTED REFERENCES AND BIBLIOGRAPHY

1. Anderson, W., 1970. A preliminary study of the relationships of saltponds and wildlife - South San Francisco Bay. Calif. Fish & Game 56:240-252.
2. Baker, H. G., 1972. A fen on the northern California coast. Madrono 21:405-416.
3. Barbour, M. G., and Major, J., eds., 1977. Terrestrial vegetation of California. John Wiley & Sons, New York.
4. Barbour, M. G., Craig, R. B., Drysdale, F. R., and Ghiselin, M. T., 1972. Coastal ecology: Bodega Head. University of California Press, Berkeley, California.
5. Braun, E. L., 1964. Deciduous Forests of Eastern North America. Hafner Publishing Co., New York. 596 pp.
6. Carpelan, H., 1957. Hydrobiology of the Alviso saltponds. Ecology 38:375-390.
7. Chapman, V. J., 1960. Salt marshes and salt deserts of the world. Leonard Hill Limited, London.
8. Chapman, V. J., 1974. Salt marshes and salt deserts of the world. In R. J. Reimold and W. H. Queen, eds. Ecology of halophytes. Academic Press, Inc., New York.
9. Correll, D. S., and Correll, H. B., 1975. Aquatic and wetland plants of southwestern United States. Two volumes. Stanford University Press, Stanford, California.
10. Cowardin, L. M., Carter, V., Golet, F. C., and LaRoe, E. T., 1977. Classification of wetlands and deep-water habitats of the United States (an operational draft). U. S. Fish and Wildlife Service. 100 pp.
11. Daley, R. H., 1976. Guide to wetlands of the Pacific Coast States. Prepared under purchase order for the Environmental Effects Laboratory, U. S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
12. Daubenmire, R. F., 1974. Plants and environment: a textbook of plant ecology. John Wiley & Sons, New York.
13. den Hartog, C., 1977. Structure, function, and classification in seagrass communities. In C. McRoy and C. Helfferich, eds. Seagrass ecosystems. Marcel Dekker, Inc., New York.

14. Eilers, H. P., 1975. Plants, plant communities, net production, and tide levels: the ecological biogeography of the Nehalem salt marshes, Tillamook County, Oregon. PhD dissertation, Oregon State University, Corvallis.
15. Fenneman, N. M., 1931. Physiography of western United States. McGraw-Hill, New York.
16. Fenneman, N. M., 1938. Physiography of eastern United States. McGraw-Hill, New York.
17. Franklin, J. F., and Dyrness, C. T., 1973. Natural vegetation of Oregon and Washington. USDA Forest Service Ser. Gen. Tech. Rept. PNW-8.
18. Golet, F. C., and Larson, J. S., 1974. Classification of fresh-water wetlands in the glaciated northeast. U. S. Fish and Wildlife Service. Resour. Publ. 16. 56 pp.
19. Harvey, H. T., 1966. Some ecological aspects of San Francisco Bay. San Francisco Bay Conservation and Development Commission Report.
20. Harvey, H. T., Pierce, C. C., and Harvey, J. T., 1976. Guide to the wetlands of the West Coast. Prepared under purchase order for the Environmental Effects Laboratory, U. S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
21. Hinde, H. P., 1954. The vertical distribution of salt marsh phanerogams in relation to tide levels. Ecological Monographs 24:209-225.
22. Hitchcock, C. L., Cronquist, A., Ownbey, M., and Thompson, J. W., 1955-1969. Vascular plants of the Pacific Northwest. Parts 1-5. University of Washington Pub. in Biol. Vol. 17. Seattle, Washington.
23. Hood, L., ed. 1975. Inventory of California natural areas. Calif. Nat. Areas Coord. Council, Sonoma, California.
24. Hoover, R. F., 1970. Vascular plants of San Luis Obispo County. University of California Press, Berkeley, California.
25. Howell, J. T., 1970. Marin flora. 2nd ed. with supplement. University of California Press, Berkeley, California.
26. Hulten, E., 1968. Flora of Alaska and neighboring territories: a manual of the vascular plants. Stanford University Press, Stanford, California.



27. Jefferson, C., 1974. Plant communities and succession in Oregon coastal marshes. PhD dissertation, Oregon State University, Corvallis.
28. Johannessen, C. L., 1964. Marshes prograding in Oregon: aerial photographs. *Science* 146:1575-1578.
29. Johannessen, C. L., 1971. The vegetation of the Willamette Valley. *Ann. Assoc. Am. Geogr.* 61:286-302.
30. Jones, G. N., 1936. A botanical survey of the Olympic Peninsula, Washington. University of Washington Pub. in Biol. Seattle, Washington.
31. Kuramoto, R. T., and Bliss, L. C., 1970. Ecology of subalpine meadows in the Olympic Mountains, Washington. *Ecological Monographs* 40:317-347.
32. Lin, J., 1970. The floristic and plant succession in vernal pond vegetation. M.A. Thesis. San Francisco State College, San Francisco.
33. McDonald, K. B., 1977. Coastal salt marsh. In M. G. Barbour and J. Major, eds. *Terrestrial Vegetation of California*. John Wiley & Sons, New York.
34. MacDonald, K. B., and Barbour, M. G., 1974. Beach and salt marsh vegetation of the North American Pacific coast. In R. J. Reimold and W. H. Queen, eds. *Ecology of halophytes*. Academic Press, Inc., New York.
35. Mall, R. E., 1969. Soil-water-salt relationships of waterfowl food plants in the Suisun Marsh of California. *Calif. Fish and Game Dept. Bull. No. 1*. Sacramento, California.
36. Mason, H., 1957. A flora of the marshes of California. University of California Press, Berkeley, California.
37. Milar, J. B., 1976. Wetland classification in western Canada: a guide to marshes and shallow open water wetlands in the grasslands and parklands of the Prairie Provinces. *Can. Wildl. Serv. Rep. Ser. 37*. 38 pp.
38. Minore, D., and Smith, C. E., 1971. Occurrence and growth of four northwestern tree species over shallow water tables. *USDA Forest Service Res. Note PNW-160*.
39. Munz, P. A., 1973. Shore wildflowers of California, Oregon, and Washington. University of California Press, Berkeley.

40. Munz, P. A., in collaboration with Keck, D. D., 1973. A California flora. Combined edition of manual and supplement. University of California Press, Berkeley, California.
41. Newcombe, C. L., and Mason, H. L., 1972. An environmental inventory of the north San Francisco Bay-Stockton ship channel area. San Francisco Bay Marine Research Center, Inc., Lafayette, California.
42. Odum, H. T., et al., 1974. Coastal ecological systems of the United States. The Conserv. Found., Washington, D. C. 4 vols.
43. Osvald, H., 1933. Vegetation of the Pacific coast bogs of North America. Acta Phytogeographica Suecica. Almquist and Wiksells Boktryckeria, Uppsala, Sweden.
44. Peck, M. E., 1961. A manual of the higher plants of Oregon. Ed. 2. Binfords & Mort, Portland, Oregon.
45. Penfound, W. T., 1952. Southern swamps and marshes. Botan. Rev. 18:413-445.
46. Pestrong, R., 1965. The development of drainage patterns on tidal marshes. Stanford Univ. Publ. Earth Sci. 10:1-87.
47. Pestrong, R., 1972. San Francisco Bay tidelands. Calif. Geol. 25:27-40.
48. Purer, E. A., 1937. Ecological study of vernal pools, San Diego County, California. Ecology 20:217-229.
49. Purer, E. A., 1942. Plant ecology of the coastal salt marshlands of San Diego County, California. Ecological Monographs 12:81-111.
50. Riggs, G. B., 1940. Comparisons of the development of some sphagnum bogs of the Atlantic coast, the interior, and the Pacific coast. American Journal of Botany 27:1-14.
51. Shaw, S. P., and Fredine, C. G., 1956. Wetlands of the United States. U. S. Fish and Wildlife Service Circ. 39. 67 pp.
52. Steward, A. N., Dennis, L. J., and Gilkey, H., 1963. Aquatic plants of the Pacific Northwest, with vegetative keys. 2nd ed. Oregon State University Press, Corvallis.
53. Stewart, R. E., and Kantrud, H. A., 1971. Classification of natural ponds and lakes in the Glaciated Prairie Region. U. S. Bur. Sport Fish. Wildl. Resour. Publ. 92. 57 pp.

54. Thompson, K., 1961. Riparian forest of the Sacramento Valley, California. *Ann. Assoc. Am. Geogr.* 51:294-315.
55. Thorne, R. F., 1968. Flora of the Santa Rosa plateau of the Santa Ana mountains, California. *Aliso* 6:17-40.
56. Thorne, R. F., 1976. The vascular plant communities of California. In J. Latting, ed. Plant communities of southern California. Calif. Native Plant Soc. Spec. Pub. 2.
57. Thorne, R. F., and Lathrop, E. W., 1969. A vernal marsh on the Santa Rosa plateau of Riverside County, California. *Aliso* 7:85-95.
58. UNESCO. 1973. International classification and mapping of vegetation. Ecology and Conservation Series No. 6, 93 pp. and chart with map symbols.
59. Vogl, R. J., 1966. Salt marsh vegetation of upper Newport Bay, California. *Ecology* 47:80-87.
60. Waaland, J. R., 1977. Common seaweeds of the Pacific Coast. Pacific Search Press, Seattle, Washington.
61. Yoakum, J., 1971. Habitat improvement. In R. D. Teague. A manual of wildlife conservation. Wildlife Soc. Colorado State Univ. Publ.
62. Zoltai, S. C., et al., 1975. Developing a wetland classification for Canada. *Proc. N. Am. Forest Soils Conf.* 4:497-511.



APPENDIX A: SCIENTIFIC AND COMMON NAMES OF PLANTS  
OF THE WEST COAST

Scientific/Common Names

*Acer negundo* L.  
Box-elder  
*Achillea millefolium* L.  
Yarrow  
*Aconitum columbinum* Nutt.  
Monkshood  
*Agardhiella* spp.  
Red alga  
*Agrostis alba* L.  
Creeping bentgrass  
*Alaria* spp.  
Brown alga  
*Alisma* spp.  
Water plantain  
*Alisma trivialis* Pursh  
Water plantain  
*Alnus oregana* Nutt.  
Red alder  
*Alnus sinuata* (Redgel) Rydb.  
Sitka alder  
*Alopecurus saccatus* Vasey  
Water foxtail  
*Athyrium filix-femina* (L.) Roth  
Lady fern  
*Atriplex patula* L.  
Fat-hen  
*Azolla filiculoides* Lam.  
Water fern  
*Baccharis pilularia* DC.  
Coyote bush  
*Batis maritima* L.  
Saltwort  
*Beckmannia syzigachne* (Steud.) Fern.  
Slough grass  
*Bergia texana* (Hook.) Seub.  
Bergia  
*Betula occidentalis* Hook.  
Water birch  
*Bidens cernua* L.  
Nodding beggar's-tick  
*C. senta* Boott  
Rough sedge  
*Cakile maritima* Scop.  
Sea rocket

*Calamagrostis nutkaensis* (Presl) Streud.  
 Pacific needle reedgrass  
*Callitriche longipedunculata* Morong  
 Water starwort  
*Callitriche* spp.  
 Water starwort  
*Caltha biflora* DC.  
 Broad-leaved caltha  
*Capsella bursa-pastoris* (L.) Medic.  
 Shepherd's purse  
*Carex aquatilis* Wahl.  
 Water sedge  
*Carex exserta* Mackenz.  
 Short-hair sedge  
*Carex hindsii* Clarke  
 Hind's sedge  
*Carex luzulina* Olney  
 Luzula-like sedge  
*Carex lyngbyei* Hornem.  
 Lyngbye's sedge  
*Carex nebrascensis* Dewey  
 Caric-sedge  
*Carex nigricans* C. A. Mey.  
 Caric-sedge  
*Carex obnupta* L. H. Bailey  
 Slough sedge  
*Carex rostrata* Stokes ex With.  
 Beaked sedge  
*Carex scopulorum* Holm  
 Holm's Rocky Mt. sedge  
*Carex sitchensis* Prescott  
 Sitka sedge  
*Carex vesicaria* L.  
 Inflated sedge  
*Ceratophyllum demersum* L.  
 Hornwort  
*Chara* spp.  
 Stonewort  
*Chenopodium ambrosioides* L.  
 Mexican tea  
*Cladophora* spp.  
 Green alga  
*Codium* spp.  
 Green alga  
*Cordylanthus maritimus* Nutt.  
 Bird's beak  
*Cornus stolonifera* Michx.  
 Red-osier dogwood  
*Cotula coronopifolia* L.  
 Brass buttons  
*Crataegus douglasii* Lindl.  
 Black hawthorn

*Cuscuta salina* Engelm.  
     Dodder  
*Darlingtonia californica* Torr.  
     Pitcher plant  
*Deschampsia caespitosa* (L.) Beauv.  
     Tufted hairgrass  
*Deschampsia danthonioides* (Trin.) Munro ex Roth  
     Annual hairgrass  
*Distichlis spicata* (L.) Greene  
     Saltgrass  
*Downingia cuspidata* (Greene) Greene  
     Downingia  
*Drosera* spp.  
     Sundew  
*Echinochloa crus-galli* (L.) Beauv.  
     Barnyard grass  
*Eichhornia crassipes* (Mart.) Solms.  
     Water hyacinth  
*Eleocharis acicularis* (L.) R. & S.  
     Needle spikerush  
*Eleocharis macrostachya* Britt.  
     Creeping spikerush  
*Eleocharis obtusa* (Wild.) Schult.  
     Blunt spikerush  
*Eleocharis parvula* (R. & S.) Link  
     Spikerush  
*Eleocharis pauciflora* (Lightf.) Link  
     Spikerush  
*Elodea canadensis* Michx.  
     Waterweed  
*Elodea nuttallii* (Planch.) St. John  
     Waterweed  
*Elymus glaucus* Buckl.  
     Wild rye  
*Elymus hirsutus* Presl  
     Wild rye  
*Empetrum nigrum* L.  
     Crowberry  
*Enteromorpha* spp.  
     Green alga  
*Epilobium adenocaulon* Hausskn.  
     Willow herb  
*Epilobium latifolium* L.  
     Willow herb  
*Epilobium watsonii* Barb.  
     Watson's willow herb  
*Equisetum fluviatile* L.  
     Water horsetail  
*Equisetum* spp.  
     Horsetail



*Eriophorum* spp.  
     Cottongrass  
*Eryngium aristulatum* Jeps.  
     Button snakeroot  
*Fragaria chiloensis* (L.) Duchn.  
     Coast strawberry  
*Frankenia grandifolia* Cham. & Schlecht.  
     Frankenia  
*Fraxinus latifolia* Benth.  
     Oregon ash  
*Fucus* spp.  
     Brown alga  
*Galium triflorum* Michx.  
     Bedstraw  
*Gaultheria shallon* Pursh  
     Salal  
*Glaux maritima* L.  
     Sea milkwort  
*Glecoma hederacea* L.  
     Ground ivy  
*Grindelia* spp.  
     Gum plant  
*Hibiscus californicus* Kell.  
     California hibiscus  
*Holcus lanatus* L.  
     Velvet grass  
*Hordeum jubatum* L.  
     Foxtail barley  
*Hydrocotyle verticillata* Thunb.  
     Marsh pennywort  
*Hymenena* spp.  
     Red alga  
*Hypericum anagalloides* Cham. & Schlecht.  
     Bog St. John's wort  
*Iridaea* spp.  
     Brown alga  
*Iris pseudacorus* L.  
     Yellow flag  
*Isoetes howellii* Engelm.  
     Quillwort  
*Jaumea carnosa* (Less) Gray  
     Jaumea  
*Juncus acuminatus* Michx.  
     Small-fruited rush  
*Juncus balticus* Willd.  
     Baltic rush  
*Juncus bolanderi* Engelm.  
     Bolander's rush  
*Juncus bufonius* L.  
     Toad rush

*Juncus dubius* Engelm.  
 Mariposa rush  
*Juncus effusus* L.  
 Common rush  
*Juncus lesuerii* Bol.  
 Salt-rush  
*Juncus phaeocephalus* Engelm.  
 Brown-headed rush  
*Juncus supiniformis* Engelm.  
 Pointed rush  
*Juncus triformis* Engelm.  
 Dwarf rush  
*Jussiaea repens* L.  
 Creeping water primrose  
*Kalmia polifolia* Wang.  
 Mountain laurel  
*Laminaria* spp.  
 Brown alga  
*Ledum columbianum* Piper  
 Pacific Labrador-tea  
*Lemna perpusilla* Torr.  
 Duckweed  
*Lemna trisulca* L.  
 Duckweed  
*Lilaea scilloides* (Poir.) Haum.  
 Flowering quillwort  
*Lilaeopsis occidentalis* Coult. & Rose  
 Lilaeopsis  
*Limonium californicum* (Boiss.) Heller  
 Sea lavender  
*Limosella aquatica* L.  
 Mudwort  
*Limosella subulata* Ives  
 Mudwort  
*Lonicera involucrata* (Richards.) Banks  
 Twinberry  
*Ludwigia palustris* (L.) Ell.  
 Water purslane  
*Lycopus uniflorus* Michx.  
 Water horehound  
*Lyngbya* spp.  
 Bluegreen alga  
*Lysichiton americanum* Hult. & St. John  
 Skunk cabbage  
*Lysimachia nummularia* L.  
 Moneywort  
*Macrocyctis* spp.  
 Brown alga  
*Menyanthes trifoliata* L.  
 Buckbean

*Mimulus guttatus* Fisch.  
     Monkey flower  
*Mimulus lewisii* Pursh  
     Monkey flower  
*Monanthochloe littoralis* Engelm.  
     Shoregrass  
*Mougeotia* spp.  
     Green alga  
*Myosurus minimus* L.  
     Mousetail  
*Myrica californica* Cham. & Schlecht.  
     Wax myrtle  
*Myriophyllum spicatum* L.  
     Water milfoil  
*Najas* spp.  
     Water nymph  
*Nasturtium officinale* R. Br.  
     Water cress  
*Nereocystis* spp.  
     Brown alga  
*Nuphar polysepalum* Engelm.  
     Yellow water lily  
*Nymphaea odorata* Aiton  
     White water lily  
*Oenanthe sarmentosa* Presl  
     Oenanthe  
*Oplopanax horridum* (J. E. Smith) Moq.  
     Devil's club  
*Oscillatoria* spp.  
     Blue-green alga  
*Paspalum distichum* L.  
     Knotgrass  
*Pedicularis groenlandica* Retz.  
     Elephant's head  
*Phalaris arundinacea* L.  
     Reed canarygrass  
*Phalaris lemmonii* Vasey  
     Canarygrass  
*Phragmites communis* Trin.  
     Common reed  
*Phyllospadix scouleri* Hook.  
     Surf-grass  
*Picea sitchensis* (Bong.) Carr.  
     Sitka spruce  
*Pilularia americana* A. Br.  
     Pillwort  
*Pinus contorta* Dougl.  
     Lodgepole pine  
*Plagiobothrys acanthocarpus* (Piper) Jtn.  
     Popcorn flower



*Plagiobothrys bracteatus* (Howell) Jtn.  
     Bracted allocarya  
*Plagiobothrys leptocladus* (Greene) Jtn.  
     Smooth-stemmed allocarya  
*Plagiobothrys reticulatus* (Piper) Jtn.  
     Netted allocarya  
*Plantago maritima* L.  
     Goosetongue  
*Platanus racemosa* Nutt.  
     Sycamore  
*Poa* spp.  
     Bluegrass  
*Pogogyne abramsii* J. T. Howell  
     Mesa mint  
*Polygonum coccineum* Muhl.  
     Water smartweed  
*Polygonum hydropiper* L.  
     Marsh pepper  
*Polygonum hydropiperoides* Michx.  
     Water pepper  
*Polygonum punctatum* Ell.  
     Perennial smartweed  
*Polypogon monspeliensis* (L.) Desf.  
     Rabbit-foot grass  
*Populus fremontii* Wats.  
     Cottonwood  
*Populus trichocarpa* T. & G.  
     Black cottonwood  
*Potamogeton berchtoldii* Feib.  
     Pondweed  
*Potamogeton epihydrus* Raf.  
     Pondweed  
*Potamogeton pectinatus* L.  
     Sago pondweed  
*Potamogeton richardsonii* (Benn.) Rydb.  
     Pondweed  
*Potamogeton robbinsii* Oakes  
     Pondweed  
*Potentilla anserina* L.  
     Silverweed  
*Potentilla egedei* Wormsk.  
     Pacific silverweed  
*Potentilla palustris* (L.) Scop.  
     Marsh cinquefoil  
*Potamogeton* spp.  
     Pondweed  
*Protococcus* spp.  
     Green alga  
*Pterygophora* spp.  
     Brown alga

*Puccinellia pumila* (Vas.) C. Hitchc.  
 Alkali grass  
*Ranunculus aquatilis* L.  
 Asiatic buttercup  
*Ranunculus flammula* L.  
 Creeping buttercup  
*Rhododendron macrophyllum* D. Don  
 California rose-bay  
*Rubus* spp.  
 Blackberry  
*Rumex acetosella* L.  
 Sheep sorrel  
*Rumex crispus* L.  
 Curly dock  
*Ruppia maritima* L.  
 Widgeon grass  
*Sagittaria latifolia* Willd.  
 Wapato  
*Sagittaria* spp.  
 Arrowhead  
*Salicornia bigelovii* Torr.  
 Bigelow's pickleweed  
*Salicornia subterminalis* Parish.  
 Subterminal pickleweed  
*Salicornia virginica* L.  
 Glasswort  
*Salix gooddingii* Ball  
 Gooding's willow  
*Salix hindsiana* Benth.  
 Sandbar willow  
*Salix hookeriana* Barr.  
 Hooker's willow  
*Salix lasiandra* Benth.  
 Black willow  
*Salix pedicellaris* Pursh  
 Bog willow  
*Salix scouleriana* Barr.  
 Scouler's willow  
*Sambucus callicarpa* Greene  
 Red elderberry  
*Samolus parviflorus* Raf.  
 Water pimpernel  
*Scirpus acutus* Muhl.  
 Tule  
*Scirpus americanus* Pers.  
 Three-square bulrush  
*Scirpus californicus* (C. A. Mey) Steud.  
 California bulrush  
*Scirpus olneyi* Gray  
 Olney's bulrush

*Scirpus robustus* Pursh  
     Bull tule  
*Scirpus validus* Vahl  
     Softstem bulrush  
*Senecio triangularis* Hook.  
     Groundsel  
*Sparganium eurycarpum* Engelm.  
     Bur-reed  
*Sparganium* spp.  
     Bur-reed  
*Spartina foliosa* Trin.  
     Cordgrass  
*Spergularia canadensis* (Pers.) G. Don  
     Sand spurry  
*Spergularia macrotheca* (Hornem.) Heynh.  
     Sand spurry  
*Spergularia marina* (L.) Griseb.  
     Sand spurry  
*Spiraea douglasii* Hook.  
     Spiraea  
*Thuja plicata* Donn  
     Western red cedar  
*Tillaea aquatica* L.  
     Water stonecrop  
*Trifolium willdenovii* Lehm.  
     Spring bank clover  
*Trifolium wormskjoldii* Lehm.  
     Coast clover  
*Triglochin maritima* L.  
     Sea arrowgrass  
*Typha angustifolia* L.  
     Narrow-leaf cat-tail  
*Typha domingensis* Pers.  
     Cat-tail  
*Typha latifolia* L.  
     Common cat-tail  
*Ulva* spp.  
     Sea lettuce  
*Urtica* spp.  
     Stinging nettle  
*Vaccinium ovatum* Pursh  
     California huckleberry  
*Vaccinium oxycoccus* L.  
     Cranberry  
*Veronica peregrina* L.  
     Speedwell  
*Veronica scutellata* L.  
     Marsh speedwell  
*Vitis californica* Benth.  
     Wild grape



Xanthium spp.  
Cocklebur  
Zostera spp.  
Eelgrass

Common/Scientific Names

- Alkali grass  
    *Puccinellia pumila* (Vas.) C. Hitchc.  
Annual hairgrass  
    *Deschampsia danthonioides* (Trin.) Munro ex Roth  
Arrowhead  
    *Sagittaria* spp.  
Asiatic buttercup  
    *Ranunculus aquatilis* L.  
Baltic rush  
    *Juncus balticus* Willd.  
Barneyard grass  
    *Echinochloa crus-galli* (L.) Beauv.  
Beaked sedge  
    *Carex rostrata* Stokes ex With.  
Bedstraw  
    *Galium triflorum* Michx.  
Bergia  
    *Bergia texana* (Hook.) Seub.  
Bigelow's pickleweed  
    *Salicornia bigelovii* Torr.  
Bird's beak  
    *Cordylanthus maritimus* Nutt.  
Black cottonwood  
    *Populus trichocarpa* T. & G.  
Black hawthorn  
    *Crataegus douglasii* Lindl.  
Black willow  
    *Salix lasiandra* Benth.  
Blackberry  
    *Rubus* spp.  
Blue-green alga  
    *Oscillatoria* spp.  
Bluegrass  
    *Poa* spp.  
Bluegreen alga  
    *Lyngbya* spp.  
Blunt spikerush  
    *Eleocharis obtusa* (Wild.) Schult.  
Bog St. John's wort  
    *Hypericum anagalloides* Cham. & Schlecht.  
Bog willow  
    *Salix pedicellaris* Pursh  
Bolander's rush  
    *Juncus bolanderi* Engelm.  
Box-elder  
    *Acer negundo* L.  
Bracted allocarya  
    *Plagiobothrys bracteatus* (Howell) Jtn.

Brass buttons  
     *Cotula coronopifolia* L.  
 Broad-leaved caltha  
     *Caltha biflora* DC.  
 Brown alga  
     *Alaria* spp.  
 Brown alga  
     *Fucus* spp.  
 Brown alga  
     *Iridaea* spp.  
 Brown alga  
     *Laminaria* spp.  
 Brown alga  
     *Macrocystis* spp.  
 Brown alga  
     *Nereocystis* spp.  
 Brown alga  
     *Pterygophora* spp.  
 Brown-headed rush  
     *Juncus phaeocephalus* Engelm.  
 Buckbean  
     *Menyanthes trifoliata* L.  
 Bull tule  
     *Scirpus robustus* Pursh  
 Bur-reed  
     *Sparganium eurycarpum* Engelm.  
 Bur-reed  
     *Sparganium* spp.  
 Button snakeroot  
     *Eryngium aristulatum* Jeps.  
 California bulrush  
     *Scirpus californicus* (C. A. Mey) Steud.  
 California hibiscus  
     *Hibiscus californicus* Kell.  
 California huckleberry  
     *Vaccinium ovatum* Pursh  
 California rose-bay  
     *Rhododendron macrophyllum* D. Don  
 Canarygrass  
     *Phalaris lemmonii* Vasey  
 Caric-sedge  
     *Carex nebrascensis* Dewey  
 Caric-sedge  
     *Carex nigricans* C. A. Mey.  
 Cat-tail  
     *Typha domingensis* Pers.  
 Coast clover  
     *Trifolium worms kjoldii* Lehm.  
 Coast strawberry  
     *Fragaria chiloensis* (L.) Duchn.



Cocklebur  
     *Xanthium* spp.  
 Common cat-tail  
     *Typha latifolia* L.  
 Common reed  
     *Phragmites communis* Trin.  
 Common rush  
     *Juncus effusus* L.  
 Cordgrass  
     *Spartina foliosa* Trin.  
 Cottongrass  
     *Eriophorum* spp.  
 Cottonwood  
     *Populus fremontii* Wats.  
 Coyote bush  
     *Baccharis pilularia* DC.  
 Cranberry  
     *Vaccinium oxycoccus* L.  
 Creeping bentgrass  
     *Agrostis alba* L.  
 Creeping buttercup  
     *Ranunculus flammula* L.  
 Creeping spikerush  
     *Eleocharis macrostachya* Britt.  
 Creeping water primrose  
     *Jussiaea repens* L.  
 Crowberry  
     *Empetrum nigrum* L.  
 Curly dock  
     *Rumex crispus* L.  
 Devil's club  
     *Oplopanax horridum* (J. E. Smith) Moq.  
 Dodder  
     *Cuscuta salina* Engelm.  
 Downingia  
     *Downingia cuspidata* (Greene) Greene  
 Duckweed  
     *Lemna perpusilla* Torr.  
 Duckweed  
     *Lemna trisulca* L.  
 Dwarf rush  
     *Juncus triformis* Engelm.  
 Eelgrass  
     *Zostera* spp.  
 Elephant's head  
     *Pedicularis groenlandica* Retz.  
 Fat-hen  
     *Atriplex patula* L.  
 Flowering quillwort  
     *Lilaea scilloides* (Poir.) Haum.

Foxtail barley  
     *Hordeum jubatum* L.  
 Frankenia  
     *Frankenia grandifolia* Cham. & Schlecht.  
 Glasswort  
     *Salicornia virginica* L.  
 Gooding's willow  
     *Salix gooddingii* Ball  
 Goosetongue  
     *Plantago maritima* L.  
 Green alga  
     *Cladophora* spp.  
 Green alga  
     *Codium* spp.  
 Green alga  
     *Enteromorpha* spp.  
 Green alga  
     *Mougeotia* spp.  
 Green alga  
     *Protococcus* spp.  
 Ground ivy  
     *Glechoma hederacea* L.  
 Groundsel  
     *Senecio triangularis* Hook.  
 Gum plant  
     *Grindelia* spp.  
 Hind's sedge  
     *Carex hindsii* Clarke  
 Holm's Rocky Mt. sedge  
     *Carex scopulorum* Holm  
 Hooker's willow  
     *Salix hookeriana* Barr.  
 Hornwort  
     *Ceratophyllum demersum* L.  
 Horsetail  
     *Equisetum* spp.  
 Inflated sedge  
     *Carex vesicaria* L.  
 Jaumea  
     *Jaumea carnosa* (Less) Gray  
 Knotgrass  
     *Paspalum distichum* L.  
 Lady fern  
     *Athyrium filix-femina* (L.) Roth  
 Lilaeopsis  
     *Lilaeopsis occidentalis* Coult. & Rose  
 Lodgepole pine  
     *Pinus contorta* Dougl.  
 Luzula-like sedge  
     *Carex luzulina* Olney

Lyngbye's sedge  
     *Carex lyngbyei* Hornem.  
 Mariposa rush  
     *Juncus dubius* Engelm.  
 Marsh cinquefoil  
     *Potentilla palustris* (L.) Scop.  
 Marsh pennywort  
     *Hydrocotyle verticillata* Thunb.  
 Marsh pepper  
     *Polygonum hydropiper* L.  
 Marsh speedwell  
     *Veronica scutellata* L.  
 Mesa mint  
     *Pogogyne abramsii* J. T. Howell  
 Mexican tea  
     *Chenopodium ambrosioides* L.  
 Moneywort  
     *Lysimachia nummularia* L.  
 Monkey flower  
     *Mimulus guttatus* Fisch.  
 Monkey flower  
     *Mimulus lewisii* Pursh  
 Monkshood  
     *Aconitum columbinum* Nutt.  
 Mountain laurel  
     *Kalmia polifolia* Wang.  
 Mousetail  
     *Myosurus minimus* L.  
 Mudwort  
     *Limosella aquatica* L.  
 Mudwort  
     *Limosella subulata* Ives  
 Narrow-leaf cat-tail  
     *Typha angustifolia* L.  
 Needle spikerush  
     *Eleocharis acicularis* (L.) R. & S.  
 Netted allocarya  
     *Plagiobothrys reticulatus* (Piper) Jtn.  
 Nodding beggar's-tick  
     *Bidens cernua* L.  
 Oenanthe  
     *Oenanthe sarmentosa* Presl  
 Olney's bulrush  
     *Scirpus olneyi* Gray  
 Oregon ash  
     *Fraxinus latifolia* Benth.  
 Pacific Labrador-tea  
     *Ledum columbianum* Piper  
 Pacific needle reedgrass  
     *Calamagrostis nutkaensis* (Presl) Streud.



Pacific silverweed  
     *Potentilla egedei* Wormsk.  
 Perennial smartweed  
     *Polygonum punctatum* Ell.  
 Pillwort  
     *Pilularia americana* A. Br.  
 Pitcher plant  
     *Darlingtonia californica* Torr.  
 Pointed rush  
     *Juncus supiniformis* Engelm.  
 Pondweed  
     *Potamogeton berchtoldii* Feib.  
 Pondweed  
     *Potamogeton epihydrus* Raf.  
 Pondweed  
     *Potamogeton richardsonii* (Benn.) Rydb.  
 Pondweed  
     *Potamogeton robbinsii* Oakes  
 Pondweed  
     *Potamogeton* spp.  
 Popcorn flower  
     *Plagiobothrys acanthocarpus* (Piper) Jtn.  
 Quillwort  
     *Isoetes howellii* Engelm.  
 Rabbit-foot grass  
     *Polypogon monspeliensis* (L.) Desf.  
 Red alder  
     *Alnus oregana* Nutt.  
 Red alga  
     *Agardhiella* spp.  
 Red alga  
     *Hymenena* spp.  
 Red elderberry  
     *Sambucus callicarpa* Greene  
 Red-osier dogwood  
     *Cornus stolonifera* Michx.  
 Reed canarygrass  
     *Phalaris arundinacea* L.  
 Rough sedge  
     *C. senta* Boott  
 Sego pondweed  
     *Potamogeton pectinatus* L.  
 Salal  
     *Gaultheria shallon* Pursh  
 Salt-rush  
     *Juncus lesuerii* Bol.  
 Saltgrass  
     *Distichlis spicata* (L.) Greene  
 Saltwort  
     *Batis maritima* L.

Sand spurry  
     *Spergularia canadensis* (Pers.) G. Don  
 Sand spurry  
     *Spergularia macrotheca* (Hornem.) Heynh.  
 Sand spurry  
     *Spergularia marina* (L.) Griseb.  
 Sandbar willow  
     *Salix hindsiana* Benth.  
 Scouler's willow  
     *Salix scouleriana* Barr.  
 Sea arrowgrass  
     *Triglochin maritima* L.  
 Sea lavender  
     *Limonium californicum* (Boiss.) Heller  
 Sea lettuce  
     *Ulva* spp.  
 Sea milkwort  
     *Glaux maritima* L.  
 Sea rocket  
     *Cakile maritima* Scop.  
 Sheep sorrel  
     *Rumex acetosella* L.  
 Shepherd's purse  
     *Capsella bursa-pastoris* (L.) Medic.  
 Shoregrass  
     *Monanthochloe littoralis* Engelm.  
 Short-hair sedge  
     *Carex exserta* Mackenz.  
 Silverweed  
     *Potentilla anserina* L.  
 Sitka alder  
     *Alnus sinuata* (Redgel) Rydb.  
 Sitka sedge  
     *Carex sitchensis* Prescott  
 Sitka spruce  
     *Picea sitchensis* (Bong.) Carr.  
 Skunk cabbage  
     *Lysichiton americanum* Hult. & St. John  
 Slough grass  
     *Beckmannia syzigachne* (Steud.) Fern.  
 Slough sedge  
     *Carex obnupta* L. H. Bailey  
 Small-fruited rush  
     *Juncus acuminatus* Michx.  
 Smooth-stemmed allocarya  
     *Plagiobothrys leptocladus* (Greene) Jtn.  
 Softstem bulrush  
     *Scirpus validus* Vahl  
 Speedwell  
     *Veronica peregrina* L.

Spikerush  
     *Eleocharis parvula* (R. & S.) Link  
 Spikerush  
     *Eleocharis pauciflora* (Lightf.) Link  
 Spiraea  
     *Spiraea douglasii* Hook.  
 Spring bank clover  
     *Trifolium willdenovii* Lehm.  
 Stinging nettle  
     *Urtica* spp.  
 Stonewort  
     *Chara* spp.  
 Subterminal pickleweed  
     *Salicornia subterminalis* Parish.  
 Sundew  
     *Drosera* spp.  
 Surf-grass  
     *Phyllospadix scouleri* Hook.  
 Sycamore  
     *Platanus racemosa* Nutt.  
 Three-square bulrush  
     *Scirpus americanus* Pers.  
 Toad rush  
     *Juncus bufonius* L.  
 Tufted hairgrass  
     *Deschampsia caespitosa* (L.) Beauv.  
 Tule  
     *Scirpus acutus* Muhl.  
 Twinberry  
     *Lonicera involucrata* (Richards.) Banks  
 Velvet grass  
     *Holcus lanatus* L.  
 Wapato  
     *Sagittaria latifolia* Willd.  
 Water birch  
     *Betula occidentalis* Hook.  
 Water cress  
     *Nasturtium officinale* R. Br.  
 Water fern  
     *Azolla filiculoides* Lam.  
 Water foxtail  
     *Alopecurus saccatus* Vasey  
 Water horehound  
     *Lycopus uniflorus* Michx.  
 Water horsetail  
     *Equisetum fluviatile* L.  
 Water hyacinth  
     *Eichhornia crassipes* (Mart.) Solms.  
 Water milfoil  
     *Myriophyllum spicatum* L.



Water nymph  
     *Najas* spp.  
 Water pepper  
     *Polygonum hydropiperoides* Michx.  
 Water pimpernel  
     *Samolus parviflorus* Raf.  
 Water plantain  
     *Alisma* spp.  
 Water plantain  
     *Alisma trivialis* Pursh  
 Water purslane  
     *Ludwigia palustris* (L.) Ell.  
 Water sedge  
     *Carex aquatilis* Wahl.  
 Water smartweed  
     *Polygonum coccineum* Muhl.  
 Water starwort  
     *Callitriche longipedunculata* Morong  
 Water starwort  
     *Callitriche* spp.  
 Water stonecrop  
     *Tillaea aquatica* L.  
 Waterweed  
     *Elodea canadensis* Michx.  
 Waterweed  
     *Elodea nuttallii* (Planch.) St. John  
 Watson's willow herb  
     *Epilobium watsonii* Barb.  
 Wax myrtle  
     *Myrica californica* Cham. & Schlecht.  
 Western red cedar  
     *Thuja plicata* Donn  
 White water lily  
     *Nymphaea odorata* Aiton  
 Widgeon grass  
     *Ruppia maritima* L.  
 Wild grape  
     *Vitis californica* Benth.  
 Wild rye  
     *Elymus glaucus* Buckl.  
 Wild rye  
     *Elymus hirsutus* Presl  
 Willow herb  
     *Epilobium adenocaulon* Hausskn.  
 Willow herb  
     *Epilobium latifolium* L.  
 Yarrow  
     *Achillea millefolium* L.  
 Yellow flag  
     *Iris pseudacorus* L.  
 Yellow water lily  
     *Nuphar polysepalum* Engelm.

## APPENDIX B: GLOSSARY

- ABUNDANCE:** a term used in quantitative vegetation sampling, referring to density of a given species per unit area; usually expressed as the total number of individual organisms in a unit area.
- ACIDIC:** having a pH value of less than 7, nonalkaline.
- ALGAE:** a nonvascular chlorophyll-bearing organism, common to various types of wetlands and very important in productivity.
- ALKALINE:** having a pH value greater than 7, nonacidic.
- ANGIOSPERM:** a plant characterized by flowers and seeds enclosed in fruits; e.g., orchids, palms, oaks, etc.
- ANNUAL:** a plant in which the entire life cycle is completed in a single growing season.
- AQUATIC VEGETATION:** a plant characteristically growing wholly or partly submerged in water.
- AUTHORITY:** the name of the person or persons who first described a particular plant to science, appearing in conjunction with a scientific name; e.g., *Typha latifolia* L. (the L. representing the botanist Linnaeus).
- BACKWATER:** an accumulation of usually quiet water, held back by a natural dike, high tides, or unusually high water levels in creeks, rivers, or lakes.
- BANANA HOLE:** type of freshwater swamp occurring in small sinkholes in Florida.
- BARRIER ISLAND:** an offshore island, similar to a bar, except with ridges, vegetation, and swampy tracts.
- BAY:** a body of water, smaller than a gulf, located in a recess in the shoreline.
- BAYHEAD:** a regional name applied to a type of freshwater swamp in Florida, dominated by a mixture of hardwood species.
- BAYOU:** a small, sluggish secondary stream or lake, often existing as an area of backwater in an abandoned channel.
- BIENNIAL:** a plant normally requiring two growing seasons to complete its life cycle; vegetative growth appears the first year and flowering and fruiting follow in the second year.
- BOG:** a vegetation type usually denoting an area of wet, acid peat.
- BRACKISH:** referring to water or soils having salinity contents of 0.5 to 30 ppt (o/oo).
- BROADLEAF:** having broad, flat leaves; usually referring to angiosperms (flowering plants) as contrasted with the needle-leaves of many gymnosperms.



**CARR:** a poorly defined regional wetland term, used primarily in parts of the Midwest; refers to a successional community (dominated by shrubs) that appears between marsh and swamp formation.

**CLIMAX:** the terminal community of a particular plant succession sequence, maintaining itself relatively unchanged unless the environment changes.

**COASTAL FLAT:** wetland type having 25 percent or less vegetative cover and that is occasionally or regularly flooded by saline water of tidal origin.

**COLONY:** a group of organisms of the same species growing in a localized area, often used to refer to a group of plants becoming established in a new situation.

**COMMUNITY:** a distinctive combination of two or more ecologically related species, living together and interacting with each other in a characteristic natural habitat.

**CONIFER:** a common term for any gymnosperm of the order Coniferales (the group containing those gymnosperms producing definite cones, as pine, spruce, etc.).

**COVER:** a term used in quantitative vegetation sampling, referring to the amount (percent) of ground with vegetation above it; estimated by vertically projecting the outline of the crown onto the ground.

**dbh:** diameter (of a tree) at breast height.

**DECIDUOUS:** shedding of leaves at end of growing season (or sometimes, in the Southwest, under periods of environmental stress before the end of the growing season); usually referring to broad-leaved woody angiosperms (flowering plants) but sometimes referring to gymnosperms (e.g., Bald cypress).

**DEFLATION PLAIN BASIN:** a basin formed in arid areas by removal of loose material from an area by wind.

**DETRITAL:** referring to dead organic tissues, decomposed material, and organisms in an ecosystem; usually including the live microorganisms involved in the decomposition of the material.

**DISCLIMAX:** a potentially long-persisting and self-reproducing vegetation type, maintaining its composition and structure only as a consequence of continuing disturbance (as by fire, grazing, etc.).

**DOMINANT:** a prevailing species of an area; a species that to a considerable extent controls the conditions for existence of its associates within an ecosystem.

**DWARF SHRUB:** woody plants characterized by numerous stems and rarely exceeding 50 cm in height.

**ECOTONE:** the transition zone between two or more adjacent plant communities, usually containing species from each of the adjacent vegetation types.



EMERSED: standing out above the water, as the leaves of certain hydrophytes.

EMERGENT: same as EMERSED.

EPIPHYTE: a plant that grows on another plant for support but is not parasitic on it.

ESTUARY: a basin in which river water mixes with and dilutes sea water.

EVERGREEN: a perennially green plant, never losing all its leaves at one time.

FEN: a poorly defined regional term for a type of marsh; usually said to be formed on peat that is circumneutral or alkaline in pH; vegetation marked by high species diversity; equivalent to the sedge-meadow of many authors.

FLOATING-LEAVED COMMUNITY: an aquatic assemblage dominated by species having leaves that float on the water surface, often floating by virtue of long flexuous petioles (such as most water lilies).

FLORA: the vegetation of an area; also used to denote a book for identification of plant species in an area.

FORBS: associated herbaceous species other than grasses; term used in ecological description of nonwoody vegetation.

FREQUENCY: a term used in quantitative vegetation sampling, relating to the number of times a species occurs in a given number of sample plots; expressed as a fraction of the total, usually in percent.

FRESH WATER: water containing less than 0.5 ppt (o/oo) salinity.

FRESHWATER AQUATIC COMMUNITY: a wetland dominated by free-floating or rooted aquatic herbs and that is semipermanently or permanently flooded by fresh water (e.g., a patch of water lilies).

FRESHWATER INLAND FLAT: a wetland having less than 25 percent vegetative cover and that is occasionally or regularly flooded by fresh water (e.g., mudflats).

FRESHWATER MARSH: a wetland having more than 25 percent vegetative cover by terrestrial herbs but 40 percent or less cover by woody plants, occasionally or regularly flooded by fresh water (e.g., sawgrass prairie).

FRESHWATER SWAMP: a wetland having more than 40 percent cover by woody plants and that is occasionally or regularly flooded by fresh water (e.g., cypress swamp).

GENUS (plural GENERA): a taxonomic category that represents a group of closely related species (e.g., all kinds of cattail are placed in the single genus *Typha*).

GRAMINOID: a term referring to grasses or grasslike plants (including the grasses, sedges, rushes, etc.).

GRASS-SEDGE BOG: a wet peatland dominated by grasses and sedges.

GROUNDWATER: water contained in rocks below the water table.

GROWTH FORM: a descriptive concept of vegetation based on some particular characteristic, such as deciduous versus evergreen and broad-leaf versus needle-leaf.

GUT: a narrow inlet of water along a coastline.

GYMNOSPERM: any of a number of different kinds of woody seed-plants in which the seeds are not enclosed in a fruit (e.g., pine, cedar, etc.).

HALOPHYTE: any plant species capable of tolerating salinity levels of more than 0.5 ppt (o/oo).

HAMMOCK: a dense growth of broad-leaved trees on a slight elevation; not considered wet enough to be a swamp.

HARDPAN: a hard, impervious subsurface layer of clay soil, usually impervious to both water and root penetration.

HARDWOOD: a broad-leaved angiosperm (flowering plant) tree having wood characterized by the presence of specialized cells called vessels.

HERB: a nonwoody plant--annual, biennial, or perennial--whose above-ground parts are short lived (in temperate regions, only one growing season).

HERBACEOUS: the adjective used to describe plants that are herbs.

HYDRIC: aquatic.

HYDROPHYTE: a plant growing in water or in characteristically wet soil.

HYPERSALINE: soil or water with a high salt content.

IMPOUNDMENT: standing body of open water created by artificially blocking or restricting the flow of a river, stream, or tidal area.

INTERMITTENT STREAM: a stream receiving its water primarily from surface runoff.

INTERTIDAL ZONE: in coastal areas, the region between levels of high tide and low tide.

KARST TOPOGRAPHY: a topography formed over limestone, dolomite, or gypsum and characterized by sinkholes, caves, and underground drainage.

KELP: any of the various large, coarse brown seaweeds (brown algae) of marine waters.

LACUSTRINE: pertaining to a lake.

LAGOON: a shallow coastal body of water, partly separated from the sea by beaches or islands; usually a lagoon is elongate and parallel to the shoreline and characterized by higher salinity than found in an estuary.



- LAKE:** a natural depression fed by one or more streams and from which a stream may flow; occurs due to widening or natural blockage of a river or stream or occurs in an isolated natural depression that is not part of a surface river or stream; usually too deep to permit the growth of rooted plants from shore to shore.
- LIANA:** a woody or herbaceous climbing plant--a vine--with its roots in the soil.
- LITTORAL:** that portion of a body of water extending from shoreline toward the middle of the water to the limit of occupancy by rooted plants.
- MANUAL:** a handbook used in the taxonomic identification of plant species.
- MARL:** a deposit of crumbly, earthy material, usually composed of clay mixed with limestone or other carbonate.
- MARSH:** a wetland dominated by nonwoody vegetation; if woody plants are present, they account for less than 40 percent vegetative cover.
- MESIC:** pertaining to a habitat characterized by a medium amount of water, neither very wet nor very dry (much vegetation adjacent to wetlands is MESOPHYTIC in nature).
- MUCK:** a type of surface deposit in a poorly drained area, consisting of much dark, partially decomposed organic matter intermixed with mineral matter.
- MUDFLATS:** an area usually supporting only sparse vegetation or no vegetation at all, although algae may be numerous on such sites; mudflats may be intertidal in coastal areas or associated with areas of widely fluctuating water levels inland.
- MUSKEG:** a term used in several different ways but usually referring to bog (in itself a poorly defined term) habitats of the far north.
- NEEDLE-LEAF:** a descriptive term used in referring to the usually slender, often evergreen, leaves of many gymnosperms (e.g., pine).
- NONVASCULAR PLANT:** referring to the simple (and usually small and inconspicuous) plants characterized by a lack of specialized conducting and supporting tissues (e.g., algae).
- NONWOODY:** referring to a plant that does not form long-lived above ground structures; plants other than trees and shrubs.
- OPEN WATER:** areas that support very little vegetative cover (25 percent or less); such areas comprise the permanent or semi-permanent interior portions of many ponds and lakes.
- OXBOW:** a shallow, crescent-shaped lake that results when loops of a meandering stream are cut off; oxbows are very common in deltaic regions.
- PEAT:** a dark-brown or black substrate produced by the partial decomposition and disintegration of mosses, sedges, trees, and other



plants growing in areas of its deposition; peat characteristically is deposited in certain wetland types.

**PERCHED WETLANDS:** wetlands located away from significant stream influence; perched wetlands include potholes and many so-called bogs, swamps, and similar areas vegetated by marsh or swamp plants.

**PERENNIAL:** a woody or herbaceous plant living from year to year, normally not dying after once flowering.

**PERIPHYTON:** algae growing attached to rocks and vegetation.

**PERMANENT:** used in reference to bodies of water that are long persistent and not subject to the normal processes of drying out by evaporative forces.

**PHREATOPHYTE:** a plant that has roots extending into the water table, thereby attaining a permanent water supply; of major concern in arid areas.

**PHYSIOGNOMY:** a descriptive concept based on the external appearance of vegetation (e.g., forest, prairie, marsh, etc.).

**PHYTOPLANKTON:** small, free-floating or weakly swimming algae, restricted to the very upper levels of bodies of water.

**PLAYA LAKE:** a slight depression in the plains of the Interior region, containing water after heavy rains but dry at other times, often supporting distinctive vegetation.

**PNEUMATOPHORE:** slender conical roots that grow vertically out of the mud, found in certain types of mangroves; used in conduction of oxygen to underground root systems.

**POCOSIN:** a regional term applied in the Carolinas to upland bogs found in undrained, shallow depressions in pine savannahs; pocosins are dominated by evergreen shrub species.

**POND:** a small, quiet body of standing water, usually sufficiently shallow to permit the potential growth of rooted plants from shore to shore.

**POTHOLES:** wetlands occupying basins formed by melting of isolated chunks of buried ice left behind by receding glaciers.

**PRODUCTIVITY:** the rate at which energy is stored in the form of organic substances, which can be used as food materials.

**RESERVOIR:** a pond or lake build for storage of water, usually by construction of a dam across a stream or river.

**RHIZOME:** an underground stem, growing horizontally, often thickened and containing accumulations of reserve food material; important structure for vegetative reproduction in many wetland plant species.

**RIPARIAN:** pertaining to vegetation of a riverbank or streamside.

**SALINA:** the term used for coastal flat (salt flat) in Puerto Rico.

**SALINE:** referring to water having too much salinity to be considered fresh water (in common usage the term is applied to water of high salinity, i.e., in excess of 30 ppt).

**SALINE FLAT:** wetlands having 25 percent or less vegetative cover that are occasionally or regularly flooded by saline water or nontidal origin (e.g., salt flats in interior of U. S.).

**SALINE WATER:** water containing greater than 30 ppt (o/oo) salinity.

**SALINITY:** pertaining to the percentage of salt found in saline water.

**SALT FLAT:** any area having high concentrations of soil salinity and supporting little or no vegetation, may be either coastal or inland.

**SALT WATER:** water containing high concentrations of salinity; normally the term is used to refer to sea water.

**SALTWATER AQUATIC WETLAND:** a wetland that is dominated by free-floating rooted, or otherwise attached herbaceous plants (including macroscopic marine algae) and that are permanently flooded by saline or brackish water (e.g., seagrass beds).

**SALTWATER MARSH:** a wetland having saline (including brackish) soils with 40 percent or less cover by woody plants and 25 percent or more cover by terrestrial herbs that is occasionally or regularly flooded by brackish or saline water (e.g., smooth cordgrass marshes).

**SALTWATER SWAMP:** a wetland having saline (including brackish) soils with 40 percent or more cover by woody plants and occasionally or regularly flooded by brackish or saline water (e.g., mangrove swamps).

**SANDBAR:** a bar or low ridge of sand bordering the shore or near the surface of the water, built up by currents or wave action.

**SEAGRASS BEDS:** usually areas of shallow water located along the coastline that support the underwater growth of seagrasses; of great value in providing cover for spawning fish and for their great productivity.

**SEAWEED:** any of the various macroscopic forms of marine algae (either Red algae, Brown algae, or Green algae).

**SEDGE:** any member of the plant family Cyperaceae; often used to refer to the specific genus *Carex* of the Cyperaceae.

**SEMIPERMANENT:** referring to a body of water that under normal circumstances is long persisting but under certain conditions may dry up in response to the normal processes of evaporation.

**SHALLOWS:** wetlands that are not usually considered marsh; represented by shallow pools, salt pans that hold water, and shallow lakes in estuarine systems; they may be nonvegetated or vegetated with emergent or submergent vascular plants or algae.

AD-A055 106

ARMY ENGINEER WATERWAYS EXPERIMENT STATION VICKSBURG MISS F/G 6/6  
PRELIMINARY GUIDE TO WETLANDS OF THE WEST COAST STATES. MAJOR A--ETC(U)  
APR 78

UNCLASSIFIED

WFC-TR-Y-7A-4

2 OF 2  
AD  
A055 106



END  
DATE  
FILMED

7 -78

DDC



**SHRUB:** a perennial woody plant of relatively low stature (usually considered less than 20 ft) with several to many stems from at or near the ground.

**SHRUB BOG:** any permanently waterlogged peatland dominated by shrubs.

**SINKHOLE:** a characteristic feature of karst topography in limestone areas; a depression or "sink" occurs when the underlying limestone is eroded through solution processes; the sinkhole may or may not hold water.

**SLOUGH:** a channel of slow-moving water in a region having little topographic relief.

**SOUND:** a wide channel or strait connecting two large bodies of water or separating an island from the mainland.

**SPECIES:** a taxonomic category below the rank of genus representing a group of closely-related individuals that actually or potentially interbreed (e.g., the genus *Typha* contains several species of Cattail: *T. latifolia*, *T. angustifolia*, and *T. domingensis*; the species are considered to be closely related and hybridization is common in Cattails).

**SPECIFIC EPITHET:** the term referring to the scientific name applied to each species within a genus (e.g., *latifolia* is the specific epithet of the species *Typha latifolia*).

**STAND:** a group of plants on a given sample area.

**STRAND VEGETATION:** a term defined in several different ways, usually referring to the vegetation at the very edge of the shore (exclusive of adjacent areas, such as dunes).

**STREAM:** any mass of water with a unidirectional flow.

**SUBMERGED:** referring to a hydrophytic plant that grows characteristically completely under water.

**SUBMERGENT:** same as SUBMERGED.

**SUBMERSED:** same as SUBMERGED.

**SUCCESSION:** the gradual, usually orderly and sometimes predictable sequence of plant communities occupying a given area with the passage of time.

**SUCCULENT:** a plant having juicy and fleshy stems and leaves that are adapted for water storage.

**SWAMP:** a wetland in which the dominant vegetation consists of trees (greater than 40 percent cover), tidal or nontidal, saltwater or freshwater.

**TIDAL:** referring to the alternate rise and fall of waters along the coast or of those having coastal influence.

**TIDAL CREEK:** a wetland situated along channels where water flows in both directions due to tidal influence.

**TRANSITION ZONE:** also referred to as **ECOTONE**; the intermediate zone between two or more adjacent plant communities, usually containing species from each of the adjacent vegetation types.

**TREE:** a perennial woody plant usually having a single trunk or stem and usually more than 6 m in height.

**TUNDRA:** a treeless plain, either wetland or "dry," found between the northern limits of trees and the region of perpetual ice and snow in the far north, or above treeline in the high mountains.

**UPLANDS:** areas that are not flooded on a regular basis and that do not support vegetation dominated by hydrophytes.

**VASCULAR PLANT:** referring to any of the many kinds of plants having specialized conducting and supporting tissue as well as differentiation into the structures known as roots, stems, and leaves (e.g., trees and shrubs of all kinds, grasses, etc.).

**VEGETATIVE COVER:** a term used in quantitative vegetation sampling, referring to the amount (percent) of ground with vegetation above it; estimated by vertically projecting the outline of leaves onto the ground.

**VEGETATIVE REPRODUCTION:** in seed plants, referring to reproduction by any of several means other than by seeds (e.g., underground rhizome systems, formation of roots on detached stems and leaves, etc.).

**VERNAL POOL:** a regional term applied to depressions in the grassland area of California; these pools, supporting a distinctive assemblage of plant species, fill with water in winter but dry up by summer.

**WATER TABLE:** the surface of the water-saturated zone of permeable rocks.

**WETLANDS:** those areas that are inundated or saturated by ground or surface water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions.

**WET MEADOWS:** graminoid-dominated marshes, often with a wide variety of associated species, often found along floodplains where freshwater swamps have been cleared.

**WILLOW HEAD:** willow-dominated freshwater swamp occurring in southern Florida.

**XERIC:** pertaining to an area or habitat having a very low or inadequate moisture supply; plants of such habitats are **XEROPHYTIC**.



## APPENDIX C: INTERPRETATION OF WETLAND DEFINITION

1. An area of some concern with respect to policy in the interpretation of the wetland definition is inclusion of the littoral zone as a wetland. In bodies of fresh water, the littoral zone is that area extending from the shoreline into the water to the limits of occupancy by rooted plants. The littoral zone has been defined in several ways by various marine science disciplines but usually is used as more-or-less synonymous with the intertidal zone (that region between high and low tides). Most intertidal littoral habitats (such as marine seagrass beds, macrophytic algal beds, rocky shores, and flats; as well as freshwater habitats such as mud flats and submerged aquatic plant beds) were regulated prior to the Federal Water Pollution Control Act Amendments of 1972, in large part by Sections 9 and 10 of the River and Harbor Act of 1899.

2. The emphasis in this report is on plant communities and their transition zones, and, from a technical standpoint, it is unrealistic to exclude the littoral zone plant communities from technical consideration. The reason for this is that plant communities are dynamic entities that are subject to considerable variation with respect to their position along various environmental gradients, and thus cannot be delineated precisely by policy statements that fail to take field realities into account. Seagrass beds, for example, usually are considered permanently inundated habitats; den Hartog,\* however, reports that of the 12 genera of seagrasses, only three (none of which occur in American waters) occur exclusively in permanently flooded habitats.

3. For technical purposes, therefore, a broad definition of wetland has been followed in this guidebook series; although for purposes of practical delineation of wetlands from a standpoint of policy regulatory functions, personnel may find it necessary to follow a narrower definition.

---

\* C. den Hartog. 1977. Structure, function, and classification in seagrass communities. in C. McRoy and C. Helfferich, eds. Seagrass ecosystems. Marcel Dekker, Inc., New York.