

AD-A058 955

SCRIPPS INSTITUTION OF OCEANOGRAPHY LA JOLLA CALIF  
NORPAX ADS REPORT NUMBER 1.(U)

OCT 77 S PAXAN

UNCLASSIFIED

SIO-REF-77-19

F/6 4/2


N00014-75-C-0152

NL

[OF]  
AD  
A058955



END  
DATE  
FILMED  
11-78  
DDC



AD A0 58955

DDC FILE COPY


LEVEL C

# SIO REFERENCE SERIES

ADS REPORT NUMBER 1

Issued by NORPAX DATA MANAGEMENT  
S. Pazan, NORPAX Data Manager

This document has been approved  
for public release and sale; its  
distribution is unlimited.



319100

October, 1977

SIO Ref. No. 77-19

University of California      Scripps Institution of Oceanography

78 09 06 007

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE   |  | READ INSTRUCTIONS BEFORE COMPLETING FORM   |
|---|--|--|
| 1. REPORT NUMBER<br>9 Rept. for Jan 76-Feb 77   | 2. GOVT ACCESSION NO.  | 3. RECIPIENT'S CATALOG NUMBER              |
| 4. TITLE (and Subtitle)<br>ADS REPORT #1<br>ADS Report<br>Number 1.   | 5. TYPE OF REPORT & PERIOD COVERED<br>12/80 p. 1                         | 6. PERFORMING ORG. REPORT NUMBER<br>77-19  |
| 7. AUTHOR(s)<br>19 Steven Paxan   | 8. CONTRACT OR GRANT NUMBER(s)<br>15 N00014-75-C-0152                    |  |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Scripps Institution of Oceanography<br>La Jolla, CA 92093  | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS<br>11 Oct 77 |  |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>Office of Naval Research<br>Arlington, VA 22217  | 12. REPORT DATE<br>August 1978   | 13. NUMBER OF PAGES<br>79 pages            |
| 14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)<br>14 SIO-REF-77-19  | 15. SECURITY CLASS. (of this report)<br>Unclassified                     | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE |
| 16. DISTRIBUTION STATEMENT (of this Report)<br>Approved for public release: distribution unlimited  |  |  |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  |  |  |
| 18. SUPPLEMENTARY NOTES   |  |  |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  |  |  |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br>During the ADS meeting of March 16, 1977 it was decided to produce a NORPAX ADS program da-a report for trimonthly periods roughly corresponding to seasons.<br>This is the first issue of that data report, and contains contour maps of... | 319 100 <i>slu</i>   |  |

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601


SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

78 09 06 007

# NORPAX

## ADS REPORT

### NUMBER 1



This document has been approved  
for public release and sale; its  
distribution is unlimited.

SPONSORING AGENCIES  
OFFICE OF NAVAL RESEARCH  
NATIONAL SCIENCE FOUNDATION

ISSUED BY NORPAX DATA MANAGEMENT  
copyright 1977

The REGENTS of UNIVERSITY of CALIFORNIA  
SIO reference no. 77-19  
Oct. 1977

## CONTENTS

### INTRODUCTION


### PROCEDURE AND ANALYSIS

1. FNWC 63x63 Northern Hemispheric Fields
2. Objectively Analyzed TRANSPAC XBT
3. Kirwan/McNally Buoys

### FIGURES

### APPENDIX A

### REFERENCES



## INTRODUCTION

During the ADS meeting of March 16, 1977 it was decided to produce a NORPAX ADS program data report for trimonthly periods roughly corresponding to seasons.

This is the first issue of that data report, and contains contour maps of monthly meaned: FNWC air temperature, sea temperature, wind speed, wind direction, surface vapor pressure, and 700 mb heights; NORPAX calculated wind stress, wind stress curl, wind shear velocity cubed ( $U^3$ ),  <sup>$U$ -cubed</sup> sensible heat flux, latent heat flux; objectively analyzed TRANSPAC temperatures at discrete depths from White and Bernstein (SIO). This report also contains monthly drifter buoy displacement vectors from Kirwan (TAMU) and McNally (SIO).

Because of extensive back dating in this first report, the time period from June, 1976 to February, 1977 will be covered.

## PROCEDURE AND ANALYSIS

### 1. FNWC 63 x 63 Northern Hemispheric Fields

FNWC data were taken from 63 x 63 polar gridded fields of 6-hourly wind speed and direction, and 12-hourly 700 mb height, sea surface temperature, air temperature, and vapor pressure.

33 x 63 fields were extracted from the larger fields; geographic coverage of a 33 x 63 field is shown in Figure 1. 10-m level winds and  $U_{10}^3$  were calculated from FNWC winds by an iterative scheme using the neutral flux-profile relationships obtained by Businger et al. (1971). Wind stress was calculated from FNWC wind speed and direction using a bulk aerodynamic equation:

$$\tau = -\rho \overline{uw} = \rho C_d U_{10}^2 \quad (1)$$

where

$\tau$  = surface stress

$\overline{uw}$  = Reynolds stress, ( $m^2 sec^{-2}$ )

$\rho$  = air density

$U_{10}$  = wind speed at 10 m

$C_d = 1.3 \times 10^{-3}$  (drag coefficient found by Smith and Banke (1975) for  $U_{10} = 10$  m/sec).

The vertical component of wind stress curl was calculated on the FNWC northern hemispheric grid using polar projection mapping equations and finite difference approximations (Appendix A).

Sensible heat flux (cal/cm sec) was obtained using an empirical bulk formula (Friehe and Schmitt, 1976):

$$S.H.F. = \rho C_p [0.2 \times 10^{-6} + C_H \cdot U_{10} \cdot (T_s - T_A)] \quad (2)$$

where,

$$C_p = 0.24 \text{ cal/gm}^{\circ}\text{C}$$

$$C_H = 0.91 \times 10^{-7}$$

$T_s$  = sea surface temperature

$T_A$  = (10 meter) air temperature

The latent heat flux ( $\text{cal/cm}^2 \text{ sec}$ ) was also calculated from a bulk formula (Friehe and Schmitt, 1976):

$$\text{L.H.F.} = L \cdot C_E \cdot U_{10} \cdot [Q_{SEA} - Q] \quad (3)$$

$L$  = 595 cal/gm (heat of evaporation)

$$C_E = 1.32 \times 10^{-3}$$

$$Q = 0.75 \left[ \frac{\text{EAIR}}{\text{gm} \cdot \text{m}^{-3}} \right] = \text{FNWC vapor pressure}$$

$$Q_{SEA} = \text{vapor density near the sea surface} \\ = 1.667 \times 10^{-7} \left[ e^{17.191(TS + 273)/273} \right].$$

Stress, stress-curl and  $U_{10}^3$  fields were calculated at 00z, 06z, 12z, 18z GMT; heat fluxes were calculated for 00z and 12z GMT every day. These calculated fields were then averaged to obtain daily, 5-daily, and monthly means. North and east-wind components were meaned to obtain daily, 5-daily, and monthly vector means of wind speed and direction. Monthly means were contoured over the region of the North Pacific from 120E to 230E and 20N to 60N. These contour maps are grouped by month in the present report in Figures 2.1-2.11, 4.1-4.11, 7.1-7.11, 10.1-10.11, 13.1-13.11, 16.1-16.11, 19.1-19.11, 22.1-22.11, and

24.1-24.11.

## 2. Objectively Analyzed Transpac XBT

XBTs have been regularly dropped from ships of opportunity in the Pacific since 1974. Recovered temperature profile data have been analyzed at Scripps by Bernstein and White and temperature residuals from their 1968 to 1974 climatology calculated. These temperature anomalies were contoured for 0, 60, 120, 200, 300 and 400 meter depths, by month, in Figures 3.1-3.6, 5.1-5.6, 8.1-8.6, 11.1-11.6, 14.1-14.6, 17.1-17.6, 20.1-20.6, 23.1-23.6, and 25.1-25.6.

## 3. Kirwan/McNally Buoys

Several buoys drogued at 35 meters were deployed in the North Pacific ADS area from June, 1976 to the present. Because of attrition of old buoys and addition of new buoys, the number at any time varied from 12 to 26. Satellite fixes were obtained on each buoy several times a day; although the analyses of these data were not complete, monthly instantaneous buoy positions were available up to December, 1976. Monthly buoy displacement vectors have been plotted in Figures 6., 9., 12., 15., 18., and 21. Because the buoys were deployed in June, 1976, there are no buoy displacement data for that month.




FIGURE 1. 33 x 63 field geographic coverage. The area of the ADS contour maps is as indicated. This map is a polar projection of the northern hemisphere.

WIND SPEED (M/SEC) JUN 76




FIGURE 2.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

JUN 76




FIGURE 2.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

JUN 76




FIGURE 2.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

JUN 76




FIGURE 2.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

JUN 76




FIGURE 2.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.08  $\text{f (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

JUN 76




FIGURE 2.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour Intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) JUN 76




FIGURE 2.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) JUN 76




FIGURE 2.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) JUN 76




FIGURE 2.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC) JUN 76




FIGURE 2.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

JUN 76




FIGURE 2.11

Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.




FIGURE 3. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

WIND SPEED (M/SEC) JUL 76




FIGURE 4.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

JUL 76




FIGURE 4.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

JUL 76




FIGURE 4.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

JUL 76




FIGURE 4.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

JUL 76




FIGURE 4.5 Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.06 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

JUL 76




FIGURE 4.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) JUL 76




FIGURE 4.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) JUL 76




FIGURE 4.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) JUL 76




FIGURE 4.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC)

JUL 76




FIGURE 4.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

JUL 76




FIGURE 4.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of aero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.





FIGURE 5. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.



**FIGURE 6.** Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) AUG 76




FIGURE 7.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

AUG 76




FIGURE 7.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

AUG 76




FIGURE 7.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

AUG 76




FIGURE 7.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC)••3)

AUG 76




FIGURE 7.5

Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.01 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

AUG 76




FIGURE 7.6

Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) AUG 76




FIGURE 7.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) AUG 76




FIGURE 7.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) AUG 76




FIGURE 7.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC)

AUG 76




FIGURE 7.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$  SEC)

AUG 76




FIGURE 7.11

Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of aero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$  sec.




FIGURE 8. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.




FIGURE 9. Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC)

SEP 76




FIGURE 10.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

SEP 76




FIGURE 10.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

SEP 76




FIGURE 10.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

SEP 76




FIGURE 10.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>2</sup>.

U STAR CUBED ((M/SEC)••3)

SEP 76




FIGURE 10.5 Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.02 (\text{m/sec})^{••3}$ .

SEA SURFACE TEMPERATURE (DEG.C)

SEP 76




FIGURE 10.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^{\circ}\text{C}$ .

AIR TEMPERATURE (DEG. C) SEP 76




FIGURE 10.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) SEP 76




FIGURE 10.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) SEP 76




FIGURE 10.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$  SEC) SEP 76




FIGURE 10.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of sens heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

SEP 76




FIGURE 10.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of aero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.




FIGURE 11. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANS PAC XBT drops.




FIGURE 12. Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) OCT 76




FIGURE 13.1      Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION OCT 76




FIGURE 13.2      Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

OCT 76




FIGURE 13.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

OCT 76




FIGURE 13.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>2</sup>.

U STAR CUBED ((M/SEC)••3)

OCT 76




FIGURE 13.5 Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.01 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

OCT 76




FIGURE 13.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) OCT 76




FIGURE 13.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) OCT 76




FIGURE 13.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) OCT 76




FIGURE 13.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC)

OCT 76




FIGURE 13.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

OCT 76




FIGURE 13.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.




FIGURE 14. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XRT drops.

FIGURE 14.

BUOY PLOT FOR OCT. . 76




FIGURE 15. Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) NOV 76




FIGURE 16.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

NOV 76




FIGURE 16.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

NOV 76




FIGURE 16.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>3</sup>)

NOV 76




FIGURE 16.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

NOV 76




FIGURE 16.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.01 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG. C)

NOV 76




FIGURE 16.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) NOV 76




FIGURE 16.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) NOV 76




FIGURE 16.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) NOV 76




FIGURE 16.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $^2$  SEC)

NOV 76




FIGURE 16.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of sens heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $^2$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

NOV 76




FIGURE 16.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ 2 sec.




FIGURE 17. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

## BUOY PLOT FOR NOV. , 76




FIGURE 18.

Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC)      DEC 76




FIGURE 19.1      Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

DEC 76




FIGURE 19.2      Direction arrows representing directions of monthly mean wind vectors at 19.5 meters.  
Vanes on arrows indicate wind speed in knots. If speed is less than  
2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is  
plotted.

WIND STRESS (DYNES/CM<sup>-2</sup>)

DEC 76




FIGURE 19.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>-3</sup>)

DEC 76




FIGURE 19.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

DEC 76




FIGURE 19.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.04 (m/sec).

SEA SURFACE TEMPERATURE (DEG.C.)

DEC 76




FIGURE 19.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) DEC 76




FIGURE 19.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are  $2^{\circ}\text{C}$ .

700 MB HEIGHT (M) DEC 76




FIGURE 19.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) DEC 76




FIGURE 19.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$  SEC) DEC 76




FIGURE 19.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

DEC 76




FIGURE 19.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure, and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.




FIGURE 20 Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomalies are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

## BUOY PLOT FOR DEC. . 76




FIGURE 21.

Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC)

JAN 77




FIGURE 22.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

JAN 77




FIGURE 22.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters.  
Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

JAN 77




FIGURE 22.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

JAN 77




FIGURE 22.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>2</sup>.

U STAR CUBED ((M/SEC)<sup>0.3</sup>)

JAN 77




FIGURE 22.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.01 (m/sec).

SEA SURFACE TEMPERATURE (DEG.C)

JAN 77




FIGURE 22.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) JAN 77




FIGURE 22.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) JAN 77




FIGURE 22.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) JAN 77




FIGURE 22.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC)

JAN 77




FIGURE 22.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ 2 sec.





FIGURE 22.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.



**FIGURE 23.** Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomalies are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

WIND SPEED (M/SEC)

FEB 77




FIGURE 24.1     Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

FEB 77




FIGURE 24.2     Direction arrows representing directions of monthly mean wind vectors at 19.5  
meters. Vines on arrows indicate wind speed in knots. If speed is less than  
2.5 knots, there is no vine; for every 5 knots above 2.5 knots, 1/2 vine is  
plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

FEB 77




FIGURE 24.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

FEB 77




FIGURE 24.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

FEB 77




FIGURE 24.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.0 $\frac{1}{3}$  (m/sec).

SEA SURFACE TEMPERATURE (DEG.C)

FEB 77




FIGURE 24.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) FEB 77




FIGURE 24.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) FEB 77




FIGURE 24.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) FEB 77




FIGURE 24.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ SEC)

FEB 77




FIGURE 24.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.




FIGURE 24.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $^2$  sec.




FIGURE 25. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomalies are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

## APPENDIX A

The vertical component of wind stress curl,

$$\hat{r} \cdot \operatorname{curl} \vec{\tau} = \frac{1}{r} \left( \frac{\partial \tau_{NS}}{\cos \varphi \partial \theta} - \frac{\partial \tau_{EW}}{\partial \varphi} \right), \quad (1)$$

was calculated on the FNWC polar grid using the chain rule and differentiating the polar projection mapping equations,

$$\zeta = 32 + 31.205 \cdot \left( \frac{1 - \sin \varphi}{1 + \sin \varphi} \right)^{1/2} \cdot \sin(\theta + 80) \quad (2a)$$

$$\eta = 32 + 31.205 \cdot \left( \frac{1 - \sin \varphi}{1 + \sin \varphi} \right)^{1/2} \cdot \cos(\theta + 80), \quad *(2b)$$

where

$\tau_{NS}$  = north-south component of wind stress

$\tau_{EW}$  = east-west component of wind stress

$\varphi$  = latitude

$\theta$  = longitude

$r$  = earth radius

$\zeta$  = horizontal axis of rectilinear  
grid on a polar projection  
in Figure 1

$\eta$  = vertical axis of grid in Figure 1.

Finite difference expressions for equation (1) were:

\* The factors in this equation, 33 and 31.205, are specific to FNWC's 63 x 63 grid.

$$\frac{\partial \tau_{NS}}{\partial \theta} = \frac{1}{2} (\tau_{NS,i+1,j} - \tau_{NS,i-1,j}) \frac{\partial \zeta}{\partial \theta} + \frac{1}{2} (\tau_{NS,i,j+1} - \tau_{NS,i,j-1}) \frac{\partial \eta}{\partial \theta} \quad (3a)$$

$$\frac{\partial \tau_{EW}}{\partial \varphi} = \frac{1}{2} (\tau_{EW,i+1,j} - \tau_{EW,i-1,j}) \frac{\partial \zeta}{\partial \varphi} + \frac{1}{2} (\tau_{EW,i,j+1} - \tau_{EW,i,j-1}) \frac{\partial \eta}{\partial \varphi} \quad (3b)$$

where

i = 0, ..., 62; discrete values of  $\theta$

j = 0, ..., 32; discrete values of  $\varphi$ .

When equations (3a) and (3b) were substituted into equations (2a) and (2b), a finite difference expression for the vertical component of wind stress curl was obtained:

REFERENCES

- Businger, J. A., J. C. Wyngaard, Y. Izumi and E. F. Bradley  
1971: Flux-profile relationships in the atmospheric  
surface layer. *J. Atmos. Sci.*, 28, 181-189.
- Friehe, C. and K. Schmitt 1976: Parameterization of air-sea  
interface fluxes of sensible heat and moisture by the  
bulk aerodynamic formulas. *J. Phys. Ocean.*, 6: 801-809.
- Smith, S. D. and E. Banke 1975: Variation of the sea surface  
drag coefficient with wind speed. *Quart. J. Roy. Meteor.  
Soc.* 101: 665-673.

AD-A058 955

SCRIPPS INSTITUTION OF OCEANOGRAPHY LA JOLLA CALIF  
NORPAX ADS REPORT NUMBER 1.(U)

F/8 4/2

UNCLASSIFIED


OCT 77 S PAXAN  
SIO-REF-77-19

N00014-75-C-0152  
NL

[ OF ]  
AD  
A058955



END  
DATE  
FILED  
11-78  
DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A058955

DDC FILE COPY

LEVEL


C

# SIO REFERENCE SERIES

ADS REPORT NUMBER 1

Issued by NORPAX DATA MANAGEMENT  
S. Pazan, NORPAX Data Manager

This document has been approved  
for public release and sale; its  
distribution is unlimited.



319100

October, 1977

SIO Ref. No. 77-19

University of California      Scripps Institution of Oceanography

78 09 06 002

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE   |  | READ INSTRUCTIONS BEFORE COMPLETING FORM   |
|---|--|--|
| 1. REPORT NUMBER<br>9) Rept. for Jan 76-Feb 77,   | 2. GOVT ACCESSION NO.  | 3. RECIPIENT'S CATALOG NUMBER              |
| 4. TITLE (and Subtitle)<br>ADS REPORT #1<br>ADS Report<br>Number 1.   | 5. TYPE OF REPORT & PERIOD COVERED<br>12/80p/                            | 6. PERFORMING ORG. REPORT NUMBER<br>77-19  |
| 7. AUTHOR(s)<br>10 Steven Paxan   | 8. CONTRACT OR GRANT NUMBER(s)<br>15 N00014-75-C-0152                    |  |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Scripps Institution of Oceanography<br>La Jolla, CA 92093  | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS<br>11 Oct 77 |  |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>Office of Naval Research<br>Arlington, VA 22217  | 12. REPORT DATE<br>August 1978   | 13. NUMBER OF PAGES<br>79 pages            |
| 14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)<br>14 SID-REF-77-19  | 15. SECURITY CLASS. (of this report)<br>Unclassified                     | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE |
| 16. DISTRIBUTION STATEMENT (of this Report)<br>Approved for public release: distribution unlimited  |  |  |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  |  |  |
| 18. SUPPLEMENTARY NOTES   |  |  |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  |  |  |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br>During the ADS meeting of March 16, 1977 it was decided to produce a NORPAX ADS program da-a report for trimonthly periods roughly corresponding to seasons.<br>This is the first issue of that data report, and contains contour maps of... | 319 100 flue   |  |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

78 09 06 007

# NORPAX

## ADS REPORT

### NUMBER 1



This document has been approved  
for public release and sale; its  
distribution is unlimited.

SPONSORING AGENCIES  
OFFICE OF NAVAL RESEARCH  
NATIONAL SCIENCE FOUNDATION

ISSUED BY NORPAX DATA MANAGEMENT  
copyright 1977

The REGENTS of UNIVERSITY of CALIFORNIA  
SIO reference no. 77-19  
Oct. 1977

## CONTENTS

### INTRODUCTION


### PROCEDURE AND ANALYSIS

1. FNWC 63x63 Northern Hemispheric Fields
2. Objectively Analyzed TRANSPAC XBT
3. Kirwan/McNally Buoys

### FIGURES

### APPENDIX A

### REFERENCES



## INTRODUCTION

During the ADS meeting of March 16, 1977 it was decided to produce a NORPAX ADS program data report for trimonthly periods roughly corresponding to seasons.

This is the first issue of that data report, and contains contour maps of monthly meaned: FNWC air temperature, sea temperature, wind speed, wind direction, surface vapor pressure, and 700 mb heights; NORPAX calculated wind stress, wind stress curl, wind shear velocity cubed ( $U^3$ )<sup>*U-cubed*</sup>, sensible heat flux, latent heat flux; objectively analyzed TRANSPAC temperatures at discrete depths from White and Bernstein (SIO). This report also contains monthly drifter buoy displacement vectors from Kirwan (TAMU) and McNally (SIO).

Because of extensive back dating in this first report, the time period from June, 1976 to February, 1977 will be covered.

## PROCEDURE AND ANALYSIS

### 1. FNWC 63 x 63 Northern Hemispheric Fields

FNWC data were taken from 63 x 63 polar gridded fields of 6-hourly wind speed and direction, and 12-hourly 700 mb height, sea surface temperature, air temperature, and vapor pressure.

33 x 63 fields were extracted from the larger fields; geographic coverage of a 33 x 63 field is shown in Figure 1. 10-m level winds and  $U_{10}^3$  were calculated from FNWC winds by an iterative scheme using the neutral flux-profile relationships obtained by Businger et al. (1971). Wind stress was calculated from FNWC wind speed and direction using a bulk aerodynamic equation:

$$\tau = -\rho \overline{uw} = \rho C_d U_{10}^2 \quad (1)$$

where

$\tau$  = surface stress

$uw$  = Reynolds stress, ( $m^2 sec^{-2}$ )

$\rho$  = air density

$U_{10}$  = wind speed at 10 m

$C_d = 1.3 \times 10^{-3}$  (drag coefficient found by Smith and Banke (1975) for  $U_{10} = 10$  m/sec).

The vertical component of wind stress curl was calculated on the FNWC northern hemispheric grid using polar projection mapping equations and finite difference approximations (Appendix A).

Sensible heat flux (cal/cm sec) was obtained using an empirical bulk formula (Friehe and Schmitt, 1976):

$$S.H.F. = \rho C_p [0.2 \times 10^{-6} + C_H \cdot U_{10} \cdot (T_s - T_A)] \quad (2)$$

where,

$$C_p = 0.24 \text{ cal/gm}^\circ\text{C}$$

$$C_H = 0.91 \times 10^{-7}$$

$T_s$  = sea surface temperature

$T_A$  = 10 meter) air temperature

The latent heat flux (cal/cm<sup>2</sup> sec) was also calculated from a bulk formula (Friehe and Schmitt, 1976):

$$L.H.F. = L \cdot C_E \cdot U_{10} \cdot [Q_{SEA} - Q] \quad (3)$$

$L$  = 595 cal/gm (heat of evaporation)

$$C_E = 1.32 \times 10^{-3}$$

$$Q = 0.75 [EAIR (gm.m<sup>-3</sup>) = FNWC vapor pressure]$$

$$Q_{SEA} = \text{vapor density near the sea surface} \\ = 1.667 \times 10^{-7} [e^{17.19[(T_S + 273)/273]}].$$

Stress, stress-curl and  $U_{10}^3$  fields were calculated at 00z, 06z, 12z, 18z GMT; heat fluxes were calculated for 00z and 12z GMT every day. These calculated fields were then averaged to obtain daily, 5-daily, and monthly means. North and east-wind components were meaned to obtain daily, 5-daily, and monthly vector means of wind speed and direction. Monthly means were contoured over the region of the North Pacific from 120E to 230E and 20N to 60N. These contour maps are grouped by month in the present report in Figures 2.1-2.11, 4.1-4.11, 7.1-7.11, 10.1-10.11, 13.1-13.11, 16.1-16.11, 19.1-19.11, 22.1-22.11, and

24.1-24.11.

2. Objectively Analyzed Transpac XBT

XBTs have been regularly dropped from ships of opportunity in the Pacific since 1974. Recovered temperature profile data have been analyzed at Scripps by Bernstein and White and temperature residuals from their 1968 to 1974 climatology calculated. These temperature anomalies were contoured for 0, 60, 120, 200, 300 and 400 meter depths, by month, in Figures 3.1-3.6, 5.1-5.6, 8.1-8.6, 11.1-11.6, 14.1-14.6, 17.1-17.6, 20.1-20.6, 23.1-23.6, and 25.1-25.6.

3. Kirwan/McNally Buoys

Several buoys drogued at 35 meters were deployed in the North Pacific ADS area from June, 1976 to the present. Because of attrition of old buoys and addition of new buoys, the number at any time varied from 12 to 26. Satellite fixes were obtained on each buoy several times a day; although the analyses of these data were not complete, monthly instantaneous buoy positions were available up to December, 1976. Monthly buoy displacement vectors have been plotted in Figures 6., 9., 12., 15., 18., and 21. Because the buoys were deployed in June, 1976, there are no buoy displacement data for that month.




FIGURE 1. 33 x 63 field geographic coverage. The area of the ADS contour maps is as indicated. This map is a polar projection of the northern hemisphere.

WIND SPEED (M/SEC) JUN 76




FIGURE 2.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

JUN 76




FIGURE 2.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>-2</sup>)

JUN 76




FIGURE 2.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>-3</sup>)

JUN 76




FIGURE 2.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC)<sup>003</sup>)

JUN 76




FIGURE 2.5 Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.01  $(\text{m/sec})^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

JUN 76




FIGURE 2.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) JUN 76




FIGURE 2.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) JUN 76




FIGURE 2.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) JUN 76




FIGURE 2.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$  SEC)

JUN 76




FIGURE 2.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

JUN 76





FIGURE 2.11

Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isoquines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.



**FIGURE 3.** Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

WIND SPEED (M/SEC) JUL 76




FIGURE 4.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION JUL 76




FIGURE 4.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

JUL 76




FIGURE 4.3

Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

JUL 76




FIGURE 4.4

The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

JUL 76




FIGURE 4.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.08 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

JUL 76




FIGURE 4.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) JUL 76




FIGURE 4.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) JUL 76




FIGURE 4.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

## VAPOR PRESSURE (MB) JUL 76




FIGURE 4.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC)

JUL 76




FIGURE 4.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ 2 sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

JUL 76




FIGURE 4.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.





FIGURE 5. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.



**FIGURE 6.** Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) AUG 76




FIGURE 7.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

AUG 76




FIGURE 7.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

AUG 76




FIGURE 7.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

AUG 76




FIGURE 7.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>2</sup>.

U STAR CUBED ((M/SEC)••3)

AUG 76




FIGURE 7.5

Monthly mean wind shear stress velocity cubed,  $U^*{}^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.04$  (m/sec) $^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

AUG 76




FIGURE 7.6

Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^{\circ}$ C.

AIR TEMPERATURE (DEG. C) AUG 76




FIGURE 7.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) AUG 76




FIGURE 7.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) AUG 76




FIGURE 7.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$  SEC)

AUG 76




FIGURE 7.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

AUG 76




FIGURE 7.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of aero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.





FIGURE 8. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANS PAC XBT drops.



**FIGURE 9.** Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) SEP 76




FIGURE 10.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

SEP 76




FIGURE 10.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

SEP 76




FIGURE 10.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

SEP 76




FIGURE 10.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>2</sup>.

U STAR CUBED ((M/SEC)••3)

SEP 76




FIGURE 10.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.08 (m/sec)<sup>3</sup>.

SEA SURFACE TEMPERATURE (DEG.C)

SEP 76




FIGURE 10.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) SEP 76




FIGURE 10.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) SEP 76




FIGURE 10.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) SEP 76




FIGURE 10.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$  SEC)

SEP 76




FIGURE 10.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of net sensible heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

SEP 76




FIGURE 10.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.




FIGURE 11. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomalies are drawn heavily. The crosses mark the positions of the TRANSPIAC XBT drops.




FIGURE 12. Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) OCT 76




FIGURE 13.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

OCT 76




FIGURE 13.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters.  
Vaness on arrows indicate wind speed in knots. If speed is less than  
2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is  
plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

OCT 76




FIGURE 13.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>2</sup>)

OCT 76




FIGURE 13.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>5</sup>.

U STAR CUBED ((M/SEC)••3)

OCT 76




FIGURE 13.5 Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.01 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG.C)

OCT 76




FIGURE 13.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) OCT 76




FIGURE 13.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) OCT 76




FIGURE 13.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) OCT 76




FIGURE 13.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ SEC)

OCT 76




FIGURE 13.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

OCT 76




FIGURE 13.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ 2 sec.




FIGURE 14. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

# BUOY PLOT FOR OCT. . 76




FIGURE 15.

Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) NOV 76




FIGURE 16.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

NOV 76




FIGURE 16.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>-2</sup>)

NOV 76




FIGURE 16.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>-3</sup>)

NOV 76




FIGURE 16.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

NOV 76




FIGURE 16.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.01 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG. C.)

NOV 76




FIGURE 16.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) NOV 76




FIGURE 16.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) NOV 76




FIGURE 16.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) NOV 76




FIGURE 16.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$  SEC)

NOV 76




FIGURE 16.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

NOV 76




FIGURE 16.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ 2 sec.




FIGURE 17. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomaly are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

## BUOY PLOT FOR NOV. . 76




FIGURE 18.

Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC)      DEC 76




FIGURE 19.1      Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

DEC 76




FIGURE 19.2      Direction arrows representing directions of monthly mean wind vectors at 19.5 meters. Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM $\cdot$ 2)

DEC 76




FIGURE 19.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10 $\cdot$ -9 DYNES/CM $\cdot$ 3)

DEC 76




FIGURE 19.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

DEC 76




FIGURE 19.5 Monthly mean wind shear stress velocity cubed,  $U^*^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are  $0.08 \text{ (m/sec)}^3$ .

SEA SURFACE TEMPERATURE (DEG.C.)

DEC 76




FIGURE 19.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are  $1^\circ\text{C}$ .

AIR TEMPERATURE (DEG. C) DEC 76




FIGURE 19.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) DEC 76




FIGURE 19.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) DEC 76




FIGURE 19.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNUC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$  SEC) DEC 76




FIGURE 19.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNUC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$  sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

DEC 76





FIGURE 19.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure, and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$  sec.



**FIGURE 20** Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomalies are drawn heavily. The crosses mark the positions of the TRANS-PAC XBT drops.

BUOY PLOT FOR DEC. . 76




FIGURE 21.

Monthly buoy displacement vectors. Vectors are drawn from start point at beginning of month to terminal point at end of month.

WIND SPEED (M/SEC) JAN 77




FIGURE 22.1 Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

JAN 77




FIGURE 22.2 Direction arrows representing directions of monthly mean wind vectors at 19.5 meters.  
Vanee on arrows indicate wind speed in knots. If speed is less than  
2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is  
plotted.

WIND STRESS (DYNES/CM<sup>0.2</sup>)

JAN 77




FIGURE 22.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>0.2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>0.3</sup>)

JAN 77




FIGURE 22.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>0.3</sup>.

U STAR CUBED ((M/SEC)<sup>••3</sup>)

JAN 77




FIGURE 22.5 Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.01 (m/sec)<sup>3</sup>.

SEA SURFACE TEMPERATURE (DEG.C)

JAN 77




FIGURE 22.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) JAN 77




FIGURE 22.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are 2°C.

700 MB HEIGHT (M) JAN 77




FIGURE 22.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) JAN 77




FIGURE 22.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC)

JAN 77




FIGURE 22.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.

LATENT HEAT FLUX ( $10^{-3}$  CAL/CM $\cdot$ 2 SEC)

JAN 77





FIGURE 22.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $\cdot$ sec.



**FIGURE 23.** Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomalies are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

WIND SPEED (M/SEC)

FEB 77




FIGURE 24.1     Absolute value of monthly mean vector wind velocities at 19.5 meters.  
Contour intervals are 1 m/sec.

WIND DIRECTION

FEB 77




FIGURE 24.2     Direction arrows representing directions of monthly mean wind vectors at 19.5 meters.  
Vanes on arrows indicate wind speed in knots. If speed is less than 2.5 knots, there is no vane; for every 5 knots above 2.5 knots, 1/2 vane is plotted.

WIND STRESS (DYNES/CM<sup>2</sup>)

FEB 77




FIGURE 24.3 Monthly mean wind stress is the mean of 6-hourly wind stress at 10 meters calculated from FNWC wind data. Contour intervals are 0.2 dynes/cm<sup>2</sup>.

CURL OF WIND STRESS (10<sup>-9</sup> DYNES/CM<sup>3</sup>)

FEB 77




FIGURE 24.4 The vertical component of monthly mean wind stress curl is the mean of 6-hourly wind stress curl approximated by finite-differences from 6-hourly wind stresses at 10 meters. Isolines of zero curl are plotted heavily, and contour intervals are  $4.0 \times 10^{-9}$  dynes/cm<sup>3</sup>.

U STAR CUBED ((M/SEC) $\cdot\cdot\cdot$ 3)

FEB 77




FIGURE 24.5 Monthly mean wind shear stress velocity cubed,  $U^3$  is the mean of 6-hourly wind shear stress velocity cubed calculated from wind speed at 10 meters. Contour intervals are 0.01 (m/sec) $\cdot\cdot\cdot$ 3.

SEA SURFACE TEMPERATURE (DEG.C)

FEB 77




FIGURE 24.6 Monthly mean sea surface temperature is the mean of 12-hourly FNWC sea surface temperatures. Contour intervals are 1°C.

AIR TEMPERATURE (DEG. C) FEB 77




FIGURE 24.7 Monthly mean air temperature is the mean of 12-hourly FNWC air temperatures.  
Contour intervals are  $2^{\circ}\text{C}$ .

700 MB HEIGHT (M) FEB 77




FIGURE 24.8 Monthly mean 700 mb height is the mean of 12-hourly FNWC 700 mb heights.  
Contour intervals are 25 meters.

VAPOR PRESSURE (MB) FEB 77




FIGURE 24.9 Monthly mean surface vapor pressure is the mean of 12-hourly FNWC vapor pressure at 19.5 meters. Contour intervals are 1 mb.

SENSIBLE HEAT FLUX ( $10^{-4}$  CAL/CM $\cdot$ 2 SEC)

FEB 77




FIGURE 24.10 Monthly mean sensible heat flux (ocean to atmosphere) is the mean of 12-hourly sensible heat flux calculated from FNWC air and sea temperature and wind using a bulk formula. Isolines of zero heat flux are plotted heavily, and contour intervals are  $1.0 \times 10^{-4}$  cal/cm $\cdot$ sec.




FIGURE 24.11 Monthly mean latent heat flux (ocean to atmosphere) is the mean of 12-hourly latent heat flux calculated from FNWC sea temperature, vapor pressure and wind using a bulk formula. Isoines of zero heat flux are plotted heavily, and contour intervals are  $0.5 \times 10^{-3}$  cal/cm $^2$  sec.




FIGURE 25. Monthly temperature anomalies ( $^{\circ}\text{C}$ ) contoured at fixed depths. Negative anomaly areas are hatched and isolines of zero anomalies are drawn heavily. The crosses mark the positions of the TRANSPAC XBT drops.

## APPENDIX A

The vertical component of wind stress curl,

$$\hat{r} \cdot \operatorname{curl} \vec{\tau} = \frac{1}{r} \left( \frac{\partial \tau_{NS}}{\cos \varphi \partial \theta} - \frac{\partial \tau_{EW}}{\partial \varphi} \right), \quad (1)$$

was calculated on the FNWC polar grid using the chain rule and differentiating the polar projection mapping equations,

$$\zeta = 32 + 31.205 \cdot \left( \frac{1 - \sin \varphi}{1 + \sin \varphi} \right)^{1/2} \cdot \sin(\theta + 80) \quad (2a)$$

$$\eta = 32 + 31.205 \cdot \left( \frac{1 - \sin \varphi}{1 + \sin \varphi} \right)^{1/2} \cdot \cos(\theta + 80), \quad *(2b)$$

where

$\tau_{NS}$  = north-south component of wind stress

$\tau_{EW}$  = east-west component of wind stress

$\varphi$  = latitude

$\theta$  = longitude

$r$  = earth radius

$\zeta$  = horizontal axis of rectilinear  
grid on a polar projection

in Figure 1

$\eta$  = vertical axis of grid in Figure 1.

Finite difference expressions for equation (1) were:

\* The factors in this equation, 33 and 31.205, are specific to FNWC's 63 x 63 grid.

$$\frac{\partial \tau_{NS}}{\partial \theta} = \frac{1}{2} (\tau_{NS,i+1,j} - \tau_{NS,i-1,j}) \frac{\partial \zeta}{\partial \theta} + \frac{1}{2} (\tau_{NS,i,j+1} - \tau_{NS,i,j-1}) \frac{\partial \eta}{\partial \theta} \quad (3a)$$

$$\frac{\partial \tau_{EW}}{\partial \phi} = \frac{1}{2} (\tau_{EW,i+1,j} - \tau_{EW,i-1,j}) \frac{\partial \zeta}{\partial \phi} + \frac{1}{2} (\tau_{EW,i,j+1} - \tau_{EW,i,j-1}) \frac{\partial \eta}{\partial \phi} \quad (3b)$$

where

$$i = 0, \dots, 62; \text{ discrete values of } \zeta$$

$$j = 0, \dots, 32; \text{ discrete values of } \eta.$$

When equations (3a) and (3b) were substituted into equations (2a) and (2b), a finite difference expression for the vertical component of wind stress curl was obtained:

REFERENCES

- Businger, J. A., J. C. Wyngaard, Y. Izumi and E. F. Bradley  
1971: Flux-profile relationships in the atmospheric  
surface layer. *J. Atmos. Sci.*, 28, 181-189.
- Friehe, C. and K. Schmitt 1976: Parameterization of air-sea  
interface fluxes of sensible heat and moisture by the  
bulk aerodynamic formulas. *J. Phys. Ocean.*, 6: 801-809.
- Smith, S. D. and E. Banke 1975: Variation of the sea surface  
drag coefficient with wind speed. *Quart. J. Roy. Meteor.  
Soc.* 101: 665-673.