

AD-A061 007

AMERICAN CYANAMID CO BOUND BROOK N J CHEMICAL RESEAR--ETC F/G 7/4  
AN INVESTIGATION OF POTENTIAL WATER-SOLUBLE CHEMILUMINESCENT MA--ETC(U)  
OCT 78 A G MOHAN, S TSENG, L G LEWIS N00014-77-C-0634  
538-6006-78 NL

UNCLASSIFIED

1 of 1  
AD  
A061 007



END  
DATE  
FILMED  
2-79  
DDC

ADA061007

12  
SC

OFFICE OF NAVAL RESEARCH  
Contract N-00014-77-C-0634  
Annual Progress Summary Report

AN INVESTIGATION OF POTENTIAL  
WATER-SOLUBLE CHEMILUMINESCENT  
MATERIALS

BY:

A. G. Mohan, S. S. Tseng, L. G. Lewis,  
L. S. Vizcarra and M. M. Rauhut

DDC FILE COPY

DDC  
NOV 8 1978  
F

AMERICAN CYANAMID COMPANY  
CHEMICAL RESEARCH DIVISION  
BOUND BROOK, NEW JERSEY

OCTOBER, 1978

This document has been approved  
for public release and sale; its  
distribution is unlimited.

78 11 06 079

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER N0014-77-C-0634-1	2. GOVT ACCESSION NO.	3. REPORT'S CATALOG NUMBER	
6 TITLE (and Subtitle) An Investigation of Potential Water-Soluble Chemiluminescent Materials. Summary Report (9/1/77-8/31/78)		4. DATE OF REPORT, MONTH COVERED 1 Sep 77-31 Aug 78	
5. AUTHOR(s) Arthur G. Mohan, Shin-Shyong Tseng, Linda G. Lewis, Lourdes S. Vizcarra M. M. Rauhut		14. PERFORMING ORG. REPORT NUMBER 538-6006-78	15. CONTRACT OR GRANT NUMBER(s) N0014-77-C-0634
6. PERFORMING ORGANIZATION NAME AND ADDRESS American Cyanamid Company Chemical Research Division Bound Brook, New Jersey 08805		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-668	
11. CONTROLLING OFFICE NAME AND ADDRESS Sponsoring Military Activity Office of Naval Research Arlington, VA 22217		12. REPORT DATE October 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 1239p.		13. NUMBER OF PAGES 33	
		18. SECURITY CLASS. (of this report) Unclassified	
		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release: Distribution Unlimited			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Chemiluminescence Oxamide Hydrogen Peroxide			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A new highly efficient class of chemiluminescent oxalic acid derivatives, N-trifluoromethyl sulfonyl oxamides, has been discovered. One member of this class, N,N'-bis(2,4,5-trichlorophenyl)-N,N'-bis(trifluoromethylsulfonyl)oxamide, has given the highest chemiluminescence quantum yield (34%) reported for a chemiluminescent reaction in an organic solvent system. Modification of the structure of these oxamides has produced water-soluble derivatives but none were chemiluminescent in aqueous media.			

408544 ↑ 1/B  
78 11 06 079

TABLE OF CONTENTS

	<u>Page</u>
1. INTRODUCTION	2-4
2. CHEMILUMINESCENCE IN AQUEOUS MEDIA	5-7
A. Mechanism of Peroxyoxalate Chemiluminescence	
B. Aqueous Reactions	
3. SYNTHESIS OF NEW CHEMILUMINESCENT MATERIALS	8-14
A. N-Triflyl Oxamides	
B. Other Oxalic Acid Derivatives	
4. EVALUATION OF NEW CHEMILUMINESCENT MATERIALS	15-16
5. EXPERIMENTAL	17-30
6. REFERENCES	31-33

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION _____	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	SPECIAL
A	

TABLES

	<u>Page</u>
TABLE 1: N-Substituted Triflamides	11a
TABLE 2: N-Substituted Trifyl Oxamides	11b
TABLE 3: Chemiluminescent Performances of N-Trifyl Oxamides	16a

1. INTRODUCTION

The American Cyanamid Company has conducted research on chemiluminescence since 1961<sup>1</sup>, with the objective of exploiting the inherent capabilities of chemiluminescence through the discovery of practical marking and illumination systems. The practical potential of chemiluminescence is based on:

- (1) A high theoretical light capacity of 173,000  $\text{lm hr l}^{-1}$  equivalent to the light output of a 40-watt incandescent bulb burning continuously for 2 weeks. This high light capacity makes chemiluminescence especially suited for portable lighting applications and for other applications where the use of distributed power is inconvenient or impossible.
- (2) Chemiluminescence is cold light. Since heat and flame are absent, chemiluminescent systems can be used where conventional hot lights would cause fire or explosion. Automobile and aircraft accidents, coal mines, repair of gas transmission lines are examples.
- (3) Reliability through long shelf life. Conventional battery systems gradually lose energy even when in storage. Chemiluminescent systems, in principle, can have indefinite storage lifetimes.

Chemiluminescent reactions with the efficiency required for practical use were unknown at the beginning of Cyanamid's research program. Even the fundamental chemistry required for the discovery of new chemiluminescent reactions was unavailable.

Following initial feasibility studies<sup>2</sup>, a detailed mechanism study was begun in 1963 under contract with the Office of Naval Research to obtain the basic knowledge required to design efficient chemiluminescent reactions. This effort was successful<sup>3</sup> and led to the discovery of practical chemical light formulations under subsequent contracts with the U.S. Naval Ordnance Laboratory<sup>4,5</sup>. As a consequence of these programs, the light output efficiency of chemical light was increased more than 1600 times<sup>6</sup>. Under a more recent contract with the U.S. Naval Weapons Center, the light output was increased an additional threefold<sup>7</sup>. Concurrent with the later contracts, Cyanamid conducted an independent applied research and development program which developed one of the efficient formulations, oxalic ester chemiluminescence, into a practical chemical light system which is in current commercial and governmental use for a wide variety of applications.

Subsequently, the feasibility of a water-based oxalate ester chemiluminescent system was established<sup>8</sup> under contract (No. NAS 5-22303) with NASA's Goddard Space Flight Center. In a recent contract with the Office of Naval Research, the mechanism of the chemiluminescent system was investigated and further insight into the effects of key reaction variables on the excitation processes was obtained<sup>9,10</sup>.

The current oxalic ester based system utilizes an organic solvent which constitutes most of its weight and volume. A substantial gain in light output per unit weight and volume could be achieved through development of a solid-state formulation comprised primarily of active ingredients. This formulation would be activated by contact with water at the point of use<sup>11</sup>.




The objective of the present research effort is to produce a water-soluble oxalic ester derivative which when combined with the appropriate fluorescer and hydrogen peroxide will produce chemiluminescent emission efficiently. The initial phase of the program has concentrated on design and synthesis of efficient oxalic acid derivatives since these materials can be evaluated with currently available fluorescers. The optimum water-activated system will undoubtedly require new water-soluble fluorescers.



## 2. Chemiluminescence in Aqueous Media

### A. Mechanism of Peroxyoxalate Chemiluminescence

Previous research<sup>9,10</sup> on the mechanism of peroxyoxalate chemiluminescence indicates the following mechanism is in agreement with the available experimental data:



Steps (1) and (2) involve nucleophilic attack by hydrogen peroxide on the oxalic ester followed by unimolecular reaction of the peracid intermediate to generate the key chemiluminescent intermediate (K.I.), tentatively proposed to be 1,2-dioxetandione. The key intermediate then reacts with a fluorescent molecule to generate the first excited singlet state of the fluorescer (Flr\*) while liberating two molecules of carbon dioxide. The excited fluorescer then emits light characteristic of a typical fluorescence process.

Steps (1) and (2) are base-catalyzed, but recent evidence<sup>9</sup> suggests these processes may involve general base catalysis rather than specific catalysis by the hydroperoxide anion. Further mechanistic studies are needed to clarify the details of the mechanism and to unequivocally establish the structure of the key intermediate.

#### B. Aqueous Reactions

Mechanistic studies carried out to date have concentrated primarily on peroxyoxalate chemiluminescent reactions in organic solvent systems. However, calculations based on the method of Ritchie<sup>12</sup>, detailed in a recent proposal to the Office of Naval Research<sup>11</sup>, indicate that efficient chemiluminescence in aqueous media should be feasible because of the higher nucleophilic reactivity of hydrogen peroxide compared to either water or hydroxide ion<sup>13</sup>.


Since low concentrations of oxalic acid (ca.  $10^{-4}$  M) have a pronounced quenching effect on the peroxyoxalate chemiluminescent reaction, design of optimum oxalic acid derivatives must maximize the selectivity of the hydrogen peroxide nucleophilic displacement over the hydrolysis reaction which not only consumes starting material but also produces oxalic acid or an oxalic acid derivative which would quench the chemiluminescent reaction.

The synthetic program was therefore directed towards the design and synthesis of new classes of oxalic acid derivatives in which the selectivity of the hydrogen peroxide reaction over the hydrolysis reaction would be maximized for optimum efficiency in an aqueous system.

3. Synthesis of New Chemiluminescent Materials

A. N-Triflyl Oxamides

One of the major obstacles in designing chemiluminescent oxalic acid derivatives for aqueous reactions is the competitive hydrolysis to oxalic acid, a known quencher of peroxyoxalate chemiluminescent reaction. While oxamides are known in general to be hydrolytically more stable than oxalic esters, the inherent organic solvent chemiluminescence efficiencies of several substituted oxamides prepared in an earlier program were poor<sup>2</sup>. Since the efficiency of diaryl oxalic esters generally increases with increasing substitution of electron attracting groups on the aromatic nucleus, activation of the oxamide nitrogen by substitution of strong electron-attracting group, X, should enhance the effectiveness of




the leaving group and thereby improve chemiluminescence efficiency. One of the strongest neutral electron withdrawing groups known<sup>14</sup> is the trifluoromethyl ( $\text{CF}_3\text{SO}_2-$ ) or triflyl group. A series of N-triflyl oxamides was therefore prepared to determine whether sufficient activation could be achieved from this group to permit triflyl oxamides to serve as a basis for design of water-soluble analogs.

The general method for preparation of these oxamides is summarized in Scheme I shown below.

Scheme I

Synthesis of N-Triflyl Oxamides





The appropriate primary amine is treated with triflic anhydride at low temperature following procedures similar to those described by Hendrickson and Bergenson.<sup>15</sup> The acidic triflamides are then acylated with oxalyl chloride in the presence of an amine acid acceptor yielding the desired triflyloxamide. Analytical data for the N-substituted triflamides and the related triflyl oxamides is summarized in Tables 1 and 2 respectively. Chemiluminescent performance of the oxamides is discussed in section 3 of this report.

The presence of a labile chlorine atom in the 2-chloroethyl trifluoromethylsulfonamide (7a) necessitated a modification of the method described in Scheme I. The oxamide was prepared by the route summarized in Scheme 2.

Scheme 2

Synthesis of N,N'-Bis(2-chloroethyl)-N,N'-Bis(trifluoromethylsulfonyl)oxamide (7b)




\*"Proton Sponge"<sup>R</sup> is 1,8-bis-(dimethylamino)naphthalene from Aldrich Chemical Company.

Use of powdered 3A Molecular Sieve<sup>16</sup> effectively catalyzed the formation of the triflamide 7a and the use of "Proton Sponge", a strong proton abstracting agent which is a poor nucleophile<sup>17</sup> avoided displacement of the chloride and afforded the oxamide in 80% crude yield.






The presence of the pyridyl nitrogen in 8a necessitated a similar approach. The triflamide was prepared by Molecular Sieve catalysis of the reaction of 2-chloro-3-amino pyridine with triflic anhydride. The triflyl oxamide 8b was prepared by Molecular Sieve catalyzed reaction of oxalyl chloride with the triflamide 8a.

Several attempts to quaternize the pyridyl nitrogens



in 8b with methyl triflate and thereby introduce water solubility were unsuccessful. Treatment of 8b with methyl iodide gave only recovered starting material.


TABLE 1  
N-SUBSTITUTED TRIFLAMIDES:  $\text{RNSO}_2\text{CF}_3$







COMPOUND R =	Yield	mp	Elemental Analysis						
			C	H	N	S	F	Cl	
<u>1a.</u> 	93%	45-7	Calcd Found	32.43 32.49	1.93 1.80	5.41 5.55	12.36 12.21	22.01 21.79	13.51 13.30
<u>2a</u> 	97%	84-6	Calcd Found	28.67 28.55	1.37 1.44	4.78 4.85	10.88 11.15	19.39 18.95	23.89 24.00
<u>3a</u> 	82%*	104-6	Calcd Found	25.69 25.59	0.92 1.00	4.28 4.35	9.79 9.95	17.43 17.00	32.11 31.98
<u>4a</u> 	91%*	99-101	Calcd Found	25.69 25.52	0.92 0.98	4.28 4.40	9.79 10.01	17.43 17.31	32.11 32.15
<u>6a</u> -CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	99%	bp 50-1 <sup>o</sup> at .5mm	Calcd Found	23.19 22.94	3.86 3.73	6.76 6.49	15.46 15.15	27.54 26.95	
<u>7a</u> -CH <sub>2</sub> CH <sub>2</sub> Cl	31%*	bp 53-5 <sup>o</sup> at .5mm	Calcd Found	17.06 17.42	2.37 2.51	6.64 6.54	15.17 15.38	27.01 26.79	16.59 16.21
<u>8a</u> 	48%*	120-2	Calcd Found	27.69 27.85	1.54 1.41	10.77 11.00	12.31 11.95	21.92 21.50	13.46 13.66

\*CRUDE YIELD




TABLE 2  
N-SUBSTITUTED TRIFYL OXAMIDES:



COMPOUND R =	mp	ELEMENTAL ANALYSIS						
		C	H	N	S	F	Cl	
<u>1b</u> 	173-4	Calcd Found	33.50 33.51	1.40 1.38	4.88 4.74	11.17 11.52	19.90 19.61	12.40 12.10
<u>2b</u> 	148-50	Calcd Found	29.92 30.24	0.90 1.02	4.36 4.24	10.00 10.45	17.75 17.19	22.12 22.27
<u>3b</u> 	190-2	Calcd Found	27.02 27.02	0.57 0.63	3.94 3.84	9.02 8.90	16.04 16.64	29.91 30.06
<u>4b</u> 	170-2	Calcd Found	27.02 26.91	0.57 0.60	3.94 3.80	9.02 8.85	16.04 16.40	29.91 30.02
<u>5b</u> 	172-5	Calcd Found	34.16 34.46	1.42 1.25	9.96 9.80	11.39 11.78	20.28 19.82	
<u>6b</u> -CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	bp 74-6 .5 mm	Calcd Found	25.64 25.60	2.99 3.11	5.98 5.77	13.68 13.92	24.36 23.95	
<u>7b</u> -CH <sub>2</sub> CH <sub>2</sub> Cl	71-3	Calcd Found	20.17 20.05	1.68 1.49	5.88 5.92	13.45 13.20	23.95 23.60	14.71 14.98
<u>8b</u> 	104-6	Calcd Found	29.27 29.10	1.05 1.14	9.76 9.51	11.15 10.89	19.86 19.40	12.20 11.95


### B. Other Oxalic Acid Derivatives

Earlier reports<sup>2,18</sup> have indicated that 2-pyridyl oxalic acid derivatives are moderately efficient chemiluminescent compounds in non-aqueous media. Mechanistic studies indicated that an acid-catalyzed rearrangement actually occurs prior to the chemiluminescent reaction, converting the initially formed amide to the pyridyl oxalate which then undergoes the usual hydrogen peroxide light-producing pathway.



The rearrangement to the active oxalic ester would not be required in the ester of 3-hydroxypyridine and quaternization of the pyridyl nitrogens should introduce water solubility and reactivity comparable to the 2-pyridone series. Thus, bis(3-pyridyl) oxalate 9a was prepared in the usual manner from 3-hydroxypyridene and oxalyl chloride. This compound, bis(3-pyridyl)oxalate 9a, gave a bright but short-lived (ca. 5 seconds) chemiluminescent reaction in a non-aqueous (phthalate ester) solvent system. The presence of the basic pyridyl nitrogen could produce this type of effect. Quaternization of the pyridyl nitrogen with methyl triflate gave what appeared to be the desired oxalate ester, but the material was highly hygroscopic and attempts to obtain an analytically pure sample have thus far been unsuccessful.

Two derivatives of benzimidazole were converted to oxamides in the expectation that the imidazole nitrogen could be converted into a water soluble quaternary ammonium derivative.



10 x = Cl

11a x = H

Oxamide 10 was found to be unstable on standing at room temperature. The analogous methyl derivative was prepared and found to be unstable on standing at room temperature as was the corresponding quaternary ammonium compound 11b prepared by treatment of 11a with methyl triflate.

4. Evaluation of New Chemiluminescent Materials

N-triflyl oxamides substituted with electron withdrawing groups have been found to produce the highest chemiluminescence quantum yields ever reported for a non-enzymatic reaction. Table 3 summarizes the performance data for eight N-triflyl oxamides. Increasing substitution of chlorine onto N-triflyl oxanilide causes a concurrent increase in efficiency from 11% for the 4-chloro derivative (1b) to 26% (mean) for the 2,4-dichloro compound (2b) to a high of 34% measured for the 2,4,5-trichloro analog (3b). This efficiency is substantially higher than the previous best efficiency of 23% reported<sup>2b</sup> for bis-(2,4-dinitrophenyl)oxalate. The lower efficiency of the 2,4,6-trichloro isomer (12.5%) was somewhat unexpected in view of the higher performance of 2b and 3b. The 4-nitro derivative was considerably more reactive than any of the chlorinated oxanilides with a  $3/4$  lifetime of only 9 minutes.







The 2-methoxyethyl compound (6b), a liquid, gave a quantum yield of about 3% and while somewhat miscible with water, did not produce light emission on contact with aqueous hydrogen peroxide. Infrared analysis of the aqueous system (in an AgCl cell) revealed this oxamide was hydrolytically stable under these conditions, suggesting that insufficient activation by the methoxyethyl group was responsible for the low efficiency in aqueous solution.

The high efficiency of the chlorinated triflyl oxamides is very likely related to the ease of displacement of the triflyl amide by hydrogen peroxide in the chemiluminescent reaction the triflyl amides are acidic (phenyl triflamide is reported to have a  $pK_a = 4.4$ )<sup>19</sup> and the conjugate bases would be expected to function as good leaving groups in the nucleophilic displacement by hydrogen peroxide. A steric effect might play a significant role in lowering the efficiency of the 2,4,6-trichloro compound (4b) compared to the 2,4,5-isomer (3b). This steric effect would be in contrast to the results found in the diphenyl oxalates where bulky ortho substituents generally increase chemiluminescence efficiency.

The very high efficiency found in the triflyl oxamide series suggests that efficient aqueous chemiluminescence should be possible once the proper balance between hydrolytic stability and chemiluminescent reactivity is obtained. The triflyl oxamide series will be examined further so a better understanding of structure-activity can be obtained.

TABLE 3

CHEMILUMINESCENCE PERFORMANCE OF N-TRIFYL OXAMIDES,<sup>1</sup>  
Tf N(R)COCO N(R)Tf

COMPOUND R =	CONC M	Q.Y. <sup>2</sup> x 10 <sup>-2</sup>	Mean Q.Y.	T.75 <sup>3</sup> min.	Light Capacity <sup>4</sup>
<u>1b</u> 	.01	11.4		44.9	37.5
<u>2b</u> 	.10	21.4		38.8	69.9
	.01	26.2		139	85.0
	.01	25.6	25.9	116	85.4
<u>3b</u> 	.01	32.6		50.4	108
	.01	35.4	34.0	42.3	117
<u>4b</u> 	.01	11.4		100	37.8
	.01	13.6	12.5	82.9	45.0
<u>5b</u> 	.01	11.0		8.88	35.7
<u>6b</u> -CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>	.10	2.85		70.2	9.20
<u>7b</u> -CH <sub>2</sub> CH <sub>2</sub> Cl	.01	3.68		125	12.2
<u>8b</u> 	.008	15.5		129	41.3

<sup>1</sup>Chemiluminescent reactions contained the indicated concentration of the oxamide, 6.75 x 10<sup>-3</sup>M 1-chloro-9,10-bis(phenylethynyl)anthracene, .375M hydrogen peroxide and 3 x 10<sup>-4</sup>M sodium salicylate in a solvent mixture of 75% (by volume) dibutylphthalate, 20% dimethyl phthalate, 5% t-butanol.

<sup>2</sup>Chemiluminescent quantum yield in Einsteins per mole of oxamide.

<sup>3</sup>Time required for 75% of the total light to be emitted.

<sup>4</sup>Integrated visible light output in lumen-hours per liter.

## 5. Experimental

Melting points were taken on a "Mel-Temp" block and are uncorrected. IR spectra were recorded on a Perkin-Elmer 297 spectrophotometer; samples were run in Nujol or in methylene chloride solutions. NMR spectra were recorded on a Varian Associates Model EM 360A spectrometer using tetramethylsilane as an internal standard. Mass spectra were recorded on a VG Micromass Model 7070 F high resolution mass spectrometer attached to a VG Model 2035 data system, using electron impact (EI) mode or chemical ionization (CI) mode. In the CI mode, methane was used as the reagent gas. Microanalyses were performed by the Micro Analytical Laboratories of American Cyanamid Company, Bound Brook, N.J. Microanalytical data is summarized in Tables 1 and 2.

Spectra and quantum yields for both fluorescence and chemiluminescence were obtained according to procedures previously described using the spectro-radiometer-luminometer developed by Roberts and Hirt<sup>20</sup>. The instrument was modified by using a RCA C31034 photomultiplier which gave satisfactory response over 350-700 nm range. The photomultiplier and microchromater were calibrated against an NBS standard tungsten lamp using the reported fluorescence quantum yield of quinine sulfate<sup>21</sup>. The spectra thus obtained are corrected.



4-Chlorophenyl-trifluorophenylsulfonyl-amide (1a)

To a solution of 4-chloroaniline (5.12g, 0.04 mole) and triethylamine (4.04 g, 0.04 mole) in 60 mL of methylene chloride was added in portions 6.73 mL (0.04 mole) of trifluoromethanesulfonic anhydride at 0°C under a nitrogen atmosphere. The mixture was stirred at room temperature for 4 hours. Evaporation of the solvent gave 10.2g of light brown oil, which solidified upon standing at room temperature. Recrystallization of the solid from cyclohexane gave 9.6g (93%) of pure 1a: IR (CH<sub>2</sub>Cl<sub>2</sub>) 3300, 1360, 1200 and 1140 cm<sup>-1</sup>; NMR (COCl<sub>2</sub>) δ 7.2 (s, 1 NH) and 7.3 to 7.4 ppm (25, aromatic protons); mass spectrum (EI), m/e 259 (M<sup>+</sup>).

N,N-bis(4-chlorophenyl)-N,N'-bis(trifluoromethylsulfonyl) oxamide (1b)

Oxalyl chloride (0.5 me; 0.056 mole) was added dropwise to a stirred solution of 1a (2.6g; 0.01 mole) and triethylamine (1.0g; 0.01 mole) in 20 mL of 1,2-dimethoxyethane at 0°C under a nitrogen atmosphere. The mixture was stirred at 0°C for 2 hours, heated to 60°C, held at 60°C for one hour, and evaporated to obtain a yellow solid which was treated with 20 mL of water. The solid remained after the water extraction was collected and recrystallized from anhydrous ether to give 2.46g (85%) of white crystalline 1b: IR (Nujol) 1750, 1730, 1210 and 1140 cm<sup>-1</sup>; NMR (COCl<sub>2</sub>) δ 7.30 to 7.65 ppm (m, aromatic protons); mass spectrum (EI), m/e 572 (M<sup>+</sup>).

2,4-Dichlorophenyl-trifluoromethylsulfonyl-amide (2a)

To a solution of 2,4-dichloroaniline (5.0g, 0.03 mole) and triethylamine (3.1 g, 0.03 mole) in methylene chloride (50 mL) was added dropwise 5.2 ml (0.03 mole) of trifluoromethanesulfonic anhydride at 0°C under a nitrogen atmosphere. After two hours of stirring of the reaction mixture, the solvent was evaporated to give a dark brown semisolid which was extracted into about 70 mL of anhydrous ether. Ether solution was dried over sodium sulfate. Evaporation of ether then gave 8.8 g (97%) of crude product which, after recrystallization from methylcyclohexane, gave pure 2a: IR (Nujol) 3250, 1360, 1200 and 1140  $\text{cm}^{-1}$ ; NMR ( $\text{CDCl}_3$ )  $\delta$  7.20 (s, NH) and 7.30 to 7.70 ppm (m, aromatic protons); mass spectrum (EI), m/e 293 ( $\text{M}^+$ ).

N,N'-Bis(2,4-dichlorophenyl)-N,N'-Bis(trifluoromethylsulfonyl) oxamide (2b)

To a solution of 2a (5.94 g; 0.02 mole) and triethylamine (2.0 g, 0.02 mole) in 50 ml of 1,2-dimethoxyethane was added dropwise oxalyl chloride (1.5 mL, 0.015 mole) at 0°C under a nitrogen atmosphere. After the addition was completed, the mixture was heated at 70°C for 5 hours. Triethylamine hydrochloride precipitated from the solution and was filtered. Evaporation of the filtrate gave 6.05 g (94%) of crude product.

It was recrystallized from cyclohexane: IR(CH<sub>2</sub>Cl<sub>2</sub>) 1750, 1730, 1210 and 1130 cm<sup>-1</sup>; NMR (acetone-d<sub>6</sub>) δ 7.7 and 7.9 ppm (2 s. aromatic protons); mass spectrum (EI), m/e 642 (M<sup>+</sup>).

2,4,5-Trichlorophenyl-trifluoromethylsulfonyl-amide (3a).

To a solution of 2,4,5-trichloroaniline (9.8 g; 0.05 mole) in methylene chloride (50 mL) was added in portions trifluoromethanesulfonic anhydride (8.4 ml, 0.05 mole) at 0°C under a nitrogen atmosphere. The mixture was stirred at 0°C for one hour, then was heated to 50°C and stirred for 5 hours. The white solid precipitated was separated by filtration and the filtrate was evaporated to obtain a dark oil. Water was added to the oil and the resulting mixture was extracted three times with ether (150 mL). The combined ethereal extracts were then dried over sodium sulfate. Evaporation of the ethereal solution gave 13.4 g (82%) of crude product which, after recrystallization from cyclohexane, gave pure 3a: IR (CH<sub>2</sub>Cl<sub>2</sub>) 3300, 1260, 1210, and 1140 cm<sup>-1</sup>, NMR (CDCl<sub>3</sub>) δ 6.4 (s, 1NH), 7.6 and 7.8 ppm (2 s aromatic protons); mass spectrum (EI), m/e 327 (M<sup>+</sup>).

N,N'-Bis(2,4,5-trichlorophenyl)-N,N'-Bis(trifluoromethylsulfonyl) oxamide (3b)

Oxalyl chloride (1 mL, 0.01 mole) was added dropwise into a stirred solution of 3a (6.2 g; 0.02 mole) and triethyl amine (2.0 g, 0.002 mole) in methylene chloride (50 mL) at 0°C under a nitrogen atmosphere. After the addition was completed, the reaction mixture was stirred at room temperature for 4 hours and then filtered to obtain 2.1 g of crude product. Recrystallization of the crude product from methyl cyclohexane gave the desired product 3b: IR (CH<sub>2</sub>Cl<sub>2</sub>) 1750, 1730, 1340, 1210 and 1120 cm<sup>-1</sup>; NMR (acetone-d<sub>6</sub>) δ 7.7 and 7.9 ppm (2s aromatic protons); mass spectrum (EI), m/e 708 (M<sup>+</sup>).

Evaporation of the filtrate, obtained by isolating the crude product and triturating the residue with water, gave an additional 3.5 g of crude product after filtering and drying. The total yield in two crops was 78.7%.

2,4,6-Trichlorophenyl-trifluoromethylsulfonyl-amide (4a)

To a solution of 2,4,6-trichloroaniline (4.91 g; 0.025 mole) and triethylamine (2.5 g; 0.025 mole) in methylene chloride (50 mL) was added dropwise 4.2 mL (0.025 mole) of trifluoromethanesulfonic anhydride at 0°C under a nitrogen atmosphere. After the addition was completed, the mixture was stirred at room temperature for 3 hours. Evaporation of the reaction mixture obtained light brown semisolid; which was treated with water (40 mL) and then extracted three times with ether (150 mL). Ether layer then separated and was dried over sodium sulfate.

Evaporation of ether gave 7.43 g (91%) of crude product which was sublimed to obtain pure 4a; IR(CH<sub>2</sub>Cl<sub>2</sub>), 3300, 1260, 1210 and 1160 cm<sup>-1</sup>; NMR (CDCl<sub>3</sub>) δ 6.5 (s, 1NH) and 7.8 ppm (s, 2H, aromatic proton); mass spectrum (EI), m/e 327 (M<sup>+</sup>).

N,N-Bis(2,4,6-trichlorophenyl)-N,N'-Bis(trifluoromethylsulfonyl) oxamide (4b)

Oxalyl chloride (2.0 mL, 0.02 mole) was added dropwise into a solution of 4a (8.0 g, 0.025 mole) and triethylamine (2.5g, 0.025 mole) in 1,2-dimethoxyethane (100 mL) at 0°C under a nitrogen atmosphere. After the addition was completed, the mixture was heated to 70°C for 15 hours. Triethylamine hydrochloride precipitated in the reaction mixture was filtered, and the filtrate was evaporated to give 6.5 g (73%) of crude product. It was recrystallized from cyclohexane: IR(CH<sub>2</sub>Cl<sub>2</sub>) 1740, 1380, 1260, 1210 and 1140 cm<sup>-1</sup>; NMR (CDCl<sub>3</sub>) δ 7.8 ppm (s 4H aromatic protons); mass spectrum (EI), m/e 708 (M<sup>+</sup>).

4-Nitrophenyl-trifluoromethylsulfonyl-amide (5a)

To a solution of 4-nitroaniline (3.45 g; 0.025 mole) and triethylamine (2.5g; 0.025 mole) in methylene chloride (100 mL) was added in portions 4.2 mL (0.025 mole) of trifluoromethanesulfonic anhydride at 0°C under a nitrogen atmosphere. After the addition was completed, the mixture was allowed to warm to room temperature and stirred for another 2 hours. Evaporation of the solvent in reaction mixture gave 7.61 g of residual solid. To this solid was added 20 mL of water and then extracted with ether.

Ether layer that separated was dried over sodium sulfate. Evaporation of ether gave 4.83 g (72%) of crude product; IR ( $\text{CH}_2\text{Cl}_2$ ) 3250, 1350, 1220 and 1140  $\text{cm}^{-1}$ ; NMR ( $\text{CDCl}_3$ )  $\delta$  7.3 (s 1 NH) and 7.4 to 7.5 ppm (2s, aromatic protons); mass spectrum (EI), m/e 270 ( $\text{M}^+$ ).

N,N'-Bis(4-nitrophenyl)-N,N'-Bis(trifluoromethylsulfonyl) oxamide (5b)

Oxalyl chloride (0.65 ml, 0.0065 mole) was added dropwise to a stirred solution of 5a (3.61 g; 0.013 mole) and triethylamine (1.3 g 0.0013 mole) in dry tetrahydrofuran (40 mL) at  $0^\circ\text{C}$  under a nitrogen atmosphere. After the addition was completed, the reaction mixture was warmed to room temperature and stirred for 18 hours. The reaction mixture was then filtered, the filtrate was evaporated, and water (20 mL) was added to the residue. The resulting precipitate was then filtered to obtain 3.29 g (92%) of crude product. It was recrystallized from diethyl ether: IR ( $\text{CH}_2\text{Cl}_2$ ) 1750, 1730, 1350, 1210 and 1130  $\text{cm}^{-1}$ ; NMR ( $\text{CDCl}_3$ )  $\delta$  7.7 to 7.9 ppm (4s, aromatic protons) mass spectrum (EI), m/e 562 ( $\text{M}^+$ ).

2-Methoxyethyl-trifluoromethylsulfonyl-amide (6a)

To a solution of 2-methoxyethyl amine (3.75 g; 0.05 mole) and triethylamine (5.0 g, 0.05 mole) in methylene chloride (40 mL) was added in portions 8.4 mL (0.05 mole) of trifluoromethanesulfonic anhydride at  $-78^{\circ}\text{C}$  under a nitrogen atmosphere. After the addition was completed, the mixture was stirred at  $-78^{\circ}\text{C}$  for 2 hours, and then evaporated. The residue thus obtained was extracted with diethyl ether and the ethereal solution dried over sodium sulfate. Evaporation gave 9.92 g (96%) of crude liquid product. Vacuum distillation gave 7.85 g of pure 6a: IR (Neat) 3300, 3150, 1370, 1250, 1170 and  $1120\text{ cm}^{-1}$ ; NMR ( $\text{CDCl}_3$ )  $\delta$  3.5<sup>o</sup> (s 3H), 3.70 (s 4H) and 6.1 ppm (s 1NH); mass spectrum (CI), m/e 208 ( $\text{M}^{\dagger}+\text{H}$ ).

N,N'-Bis(2-Methoxyethyl)-N'N'-Bis(trifluoromethylsulfonyl) oxamides (6b)

Oxalyl chloride (1 mL, 0.01 mole) was added portionwise into a stirred solution of 6a (4.0 g; 0.02 mole) and triethyl amine (2.0 g; 0.02 mole) in 1,2-dimethoxyethane (40 mL) at  $0^{\circ}\text{C}$  under a nitrogen atmosphere. After the addition was completed, the reaction mixture was stirred at  $0^{\circ}\text{C}$  for one hour, heated to  $50^{\circ}\text{C}$  and stirred thereat for 2 hours, and concentrated to remove the solvent. The residue was then extracted with ether and then dried over sodium sulfate. Evaporation of ethereal solution obtained 4.65 g (99%) of crude product. Vacuum distillation of the crude product gave pure 6b: IR (Neat) 1740, 1720, 1420, 1320, 1200, 1160

and  $1120\text{ cm}^{-1}$ ; NMR ( $\text{CDCl}_3$ )  $\delta$  3.40 (s, 3H), 3.70 (t, 2H,  $J=6\text{ Hz}$ ) and 4.00 ppm (t, 2H,  $J=6\text{ Hz}$ ); mass spectrum (CI),  $m/e$  461 ( $M^+ + H$ ).

2-Chloroethyl-trifluoromethylsulfonyl-amide (7a)

To a suspension of 2-chloroethyl amine hydrochloride (5.80 g; 0.05 mole) and powdered 3A molecular sieves (15g, from Linde Division, Union Carbide Corporation) in dichloroethane (100 mL) was added dropwise trifluoromethanesulfonyl chloride (5.3 mL; 0.025 mole) at room temperature under a nitrogen atmosphere. After the addition was completed, the mixture was stirred at  $80^\circ\text{C}$  for 20 hours. Filtration of the solid from reaction mixture, followed by evaporation of the filtrate gave 3.26g (31%) of crude liquid product. Vacuum distillation of the liquid gave pure 7a: IR (Neat) 3310, 3150, 1420, 1370, 1220, 1200 and  $1150\text{ cm}^{-1}$ ; mass spectrum (CI),  $m/e$  212 ( $M^+ + H$ ).

N,N'-Bis(2-chloroethyl)-N,N'-Bis(trifluoromethylsulfonyl) oxamide (7b)

Oxalyl chloride (0.58 ml; 0.0067 mole) was added dropwise into a solution of 7a (2.80 g; 0.013 mole) and "Proton Sponge"<sup>R</sup> (1.44 g; 0.0067 mole, from Aldrich Chemical Company, Inc) in methylene chloride (50 mL) at  $0^\circ\text{C}$  under a nitrogen atmosphere. After the addition, the mixture was stirred at room temperature for 24 hours. Solvent was evaporated, and the residue was treated with diethyl ether. The ethereal solution



was dried over sodium sulfate. Evaporation of ether followed by the treatment of residue with petroleum ether gave 2.55 g (80%) of crude product. Recrystallization of the crude product from petroleum ether afforded pure 7b: IR ( $\text{CH}_2\text{Cl}_2$ ) 1720, 1400, 1260, 1230, 1160 and 1120  $\text{cm}^{-1}$ ; mass spectrum (CI), m/e 477 ( $\text{M}^+ + \text{H}$ ).

2-Chloro-3 pyridyl-trifluoromethylsulfonyl-amide (8a)

To a suspension of 2-chloro-3-aminopyridine (5.14 g; 0.04 mole) and powdered 3A molecular sieves (10 g) in methylene chloride (60 mL) was added dropwise trifluoromethanesulfonic anhydride (3.4 mL) at  $0^\circ\text{C}$  under a nitrogen atmosphere. After the addition was completed, the mixture was stirred at room temperature for 5 hours and then the solid was filtered. Filtrate was evaporated and treated with water to give 5.0 g (48%) of crude solid product. It was collected and recrystallized from cyclohexane to give pure 8a: IR ( $\text{CH}_2\text{Cl}_2$ ) 3300, 1370, 1230, 1210 and 1140  $\text{cm}^{-1}$ ; NMR ( $\text{CDCl}_3$ )  $\delta$  7.35 (m, 1H), 8.0 (2d, 1H,  $J=4\text{Hz}$ ), 8.35 (2d, 1H,  $J=4\text{Hz}$ ) and 8.30 ppm (s 1NH); mass spectrum (EI), m/e 260 ( $\text{M}^+$ ).

N,N'-Bis(2-chloro-3-pyridyl)-N,N'-Bis(trifluoromethylsulfonyl) oxamide (8b)

Oxalyl chloride (0.53 mL; 0.006 mole) was added dropwise into a stirring suspension of 8a (2.61 g; 0.01 mole) and powdered 3A molecular sieves (5.0g) in methylene chloride (75 mL) at  $0^\circ\text{C}$  under a nitrogen atmosphere. The mixture was then heated to  $60^\circ\text{C}$ , held thereat for 3 hours, and then at room temperature for 60 hours. The reaction mixture was

filtered and the filtrate was evaporated to dryness. The resulting residue was extracted with diethyl ether, and the combined ethereal extracts were dried over sodium sulfate. Evaporation of the dried ethereal solution obtained 2.33 g (81%) of crude product. Recrystallization of the crude product from cyclohexane gave the desired product 8b: IR (CH<sub>2</sub>Cl<sub>2</sub>) 1750, 1730, 1420, 1260, 1220 and 1130 cm<sup>-1</sup>; NMR (CDCl<sub>3</sub>) δ 7.50 (m 1H), 8.10 (2a, 1H, J=4H), and 8.5 ppm (m 1H); mass spectrum (EI), m/e 574 (M<sup>+</sup>).

Attempted Synthesis of 3,3'- [oxalyl bis ((trifluoromethyl) sulfonyl) imino] bis (2-chloro-1-methylpyridinium trifluoromethanesulfonate) (8c)

To a solution of 8b (0.5 g; 0.001 mole) in methylene chloride (20 mL) was added methyl trifluoromethanesulfonate (0.25 g; 0.0015 mole) at room temperature under a nitrogen atmosphere. The mixture was stirred at room temperature for 20 hours. Evaporation of the reaction mixture gave a yellow gummy material (0.9g). Its IR (neat) spectrum indicated the oxalate C=O band at 1740 cm<sup>-1</sup>. Attempts to obtain a pure product have not been successful. Therefore it is not certain whether 8c has actually been made. In another experiment a solution of 8b (1 g) and methyl iodide (0.5 mL) in toluene (20 mL) was heated at 80°C for 3 hours followed by stirring the mixture at room temperature for another 60 hours. Evaporation of the reaction mixture gave a residual solid, which was indicated by its IR and NMR spectra to be unchanged 8b.

Bis(3-pyridyl)oxalate (9a)

To a solution of 3-hydroxy pyridine (4.76 g; 0.05 mole) and triethylamine (5.00 g, 0.05 mole) in 1,2-dimethoxyethane (100 mL) was added dropwise oxalyl chloride (2.2 mL; 0.025 mole) at room temperature under a nitrogen atmosphere. After the addition was completed, the mixture was stirred at room temperature for one hour, then heated to reflux for two hours. The reaction mixture was evaporated to dryness, and the residue was washed three times with chloroform (90 mL). The white solid that remained after chloroform treatment was separated by filtration (weight, 5.0g). IR spectrum (Nujal) indicated the expected carbonyl band of 9a at  $1760\text{ cm}^{-1}$ .

3-Hydroxy-1-methyl pyridinium-trifluoromethanesulfonate oxalate Ester (9b)

To a solution of 9a (1g, 0.0004 mole) in methylene chloride (50 mL) was added dropwise methyl trifluoromethane sulfonate (1 mL) at  $0^{\circ}\text{C}$ . After the addition was completed, the mixture was stirred at  $0^{\circ}\text{C}$  for 1 hour, then at room temperature for another 4 hours. The white precipitate was separated on a filter and then dried under vacuum to give the desired product 9b (1.92 g, 100%): IR (Najol) 3100, 1780, 1500, 1460, 1270, 1210, 1160 and  $1100\text{ cm}^{-1}$ ; NMR (DMSO- $d_6$ )  $\delta$  4.3 (s, 6H), 7.95 (t, 2H,  $J=2\text{ Hz}$ ) and 8.50 ppm (t, 2H,  $J=2\text{ Hz}$ ). This material was highly hygroscopic and attempts to obtain an analytically pure sample have thus far been unsuccessful.

1,1'-Oxalyl bis [(2-chloromethyl)benzimidazole] (10)

Oxalyl chloride (3.5 mL; 0.04 mole) was added dropwise into a stirring solution of 2-chloro-methylbenzimidazole (10 g; 0.06 mole) in 1,2-di-methoxyethane (100 mL) at 0°C under a nitrogen atmosphere. After the addition was completed, the mixture was stirred at 0°C for an additional hour. Solid was filtered, and the filtrate was evaporated to dryness. The gummy brown material obtained from evaporation was digested with anhydrous ether. Solid thus formed was filtered immediately. Its IR spectrum (Nujol) showed the characteristic C=O band at 1700 cm<sup>-1</sup>. However, the product appeared to be unstable, it decomposed slowly upon standing at room temperature. Attempted recrystallization from organic solvents failed.

1,1'-oxalylbis [2-Methylbenzimidazole] (11a)

Oxalyl chloride (2.5 mL. 0.025 mole) was added dropwise into a solution of 2-methylbenzimidazole (5.28 g; 0.04 mole) and triethylamine (4g; 0.04 mole) in dry tetrahydrofuron (120 mL) at 0°C under a nitrogen atmosphere. After the addition was completed, the mixture was stirred at 0°C for one hour, and then at room temperature for another hour. The solid precipitate was filtered. Evaporation of the filtrate gave 5.82 g (91.5%) of crude product. Its IR spectrum (NUjol) showed a strong C=O band at 1700 cm<sup>-1</sup>.

1,1'-oxalyl bis(2,3-dimethylbenzimidazolium trifluoromethane sulfonate (11b))

To a solution of crude 11a (2.4g; 0.075 mole) in methylene chloride (80 mL) was added in portions methyl trifluoromethane sulfonate (2 mL) at 0°C. The mixture was then stirred at room temperature for 48 hours. A light green solid precipitate was filtered and dried (3.27g). Its IR spectrum (Nujol) showed bands due to C=O ( $1700\text{ cm}^{-1}$ ),  $\text{SO}_2$  ( $1160$  and  $1180\text{ cm}^{-1}$ ) and  $\text{CF}_3$  ( $1400$  and  $1380\text{ cm}^{-1}$ ). The quaternary ammonium compound (11b) was found to be unstable on standing at room temperature although it was chemiluminescent in the organic solvent system, it was not chemiluminescent in aqueous media containing hydrogen peroxide.

6. References

1. A brief review of the applied aspects of Cyanamid's chemical light research has been published: M. M. Rauhut in "Chemiluminescence and Bioluminescence", M. J. Cormier, D. M. Hercules and J. Lee, Eds., Plenum Press, New York, 1973, p. 451.
2. (a) M. M. Rauhut and R. A. Clarke, "Chemiluminescent Materials", Summary Technical Report to the Office of Naval Research and the Advanced Research Projects Agency, Contract 4200(00) 1966 (AD 653-00).  
  
(b) M. M. Rauhut, L. J. Bollyky, B. G. Roberts, M. Loy, R. H. Whitman, A. V. Iannotta, A. M. Semsel and R. A. Clarke, *J. Amer. Chem. Soc.*, 89, 6515 (1967).
3. See for example: (a) M. M. Rauhut, A. M. Semsel and B. G. Roberts, *J. Org. Chem.*, 31, 2431 (1966);  
  
(b) J. M. W. Scott and R. F. Phillips, U. S. Patent 3,366,572
4. M. M. Rauhut, L. J. Bollyky, D. R. Maulding, R. H. Whitman, A. M. Semsel, D. H. Lantz, B. G. Roberts and C. E. Eagle, "Chemiluminescent Materials", Summary Technical Report to the U.S. Naval Ordinance Laboratory, Contract N60921-67-C-0214 (1970) (AD 886-5792).
5. A. G. Mohan, L. J. Bollyky, D.E. DeWalt, E. Klingsberg, D. R. Maulding and R. L. Narburgh, "Chemiluminescent Materials", Summary Technical Report to the U.S. Naval Ordinance Laboratory, Contract N60921-70C-0198 (1971).

6. M. M. Rauhut, U. S. Patent, 3,749,679 (1973).
7. A. G. Mohan and R. L. Marburgh, "Development of High Light Capacity Chemiluminescent Formulations, Summary Technical Report to the Naval Weapons Center, Contract N00123-71-C-1069(1972).
8. A. G. Mohan, R. G. Dulina and A. A. Doering, "Investigation of Water-Soluble Chemiluminescent Materials", Final Report to the National Aeronautics and Space Administration, Contract NSA5-22303 (1976).
9. M. M. Rauhut and A. M. Semsel, "A Study of Oxalic Ester Chemiluminescent Reactions", Final Report to the Office of Naval Research, Contract N0014-73-C-0343.
10. For review see M. M. Rauhut, *Accts. Chem. Res.*, 2, 85 (1969) and references cited therein.
11. Technical Proposal to the Office of Naval Research for Continuation of Investigation of Water-Soluble Chemiluminescent Materials, April, 1978.
12. C. D. Ritchie, *Accts of Chem. Res.*, 5 348 (1972) and *J. Amer. Chem. Soc.*, 97, 1170 (1975).
13. A. G. Mohan, American Cyanamid Co., Technical Proposal to the Office of Naval Research for an Investigation of Water-Soluble Chemiluminescent Materials, January, 1977.
14. J. B. Hendrickson, D. B. Sternbach and K. W. Blair, *Accts. Chem. Res.*, 10, 306 (1977).

15. J. B. Hendrickson and R. Bergenson, *Tetrahedron Letters*, 4607 (1973).
16. A. R. Banks, R. F. Fibiger and T. Jones, *J. Org. Chem.*, 42, 3965 (1977).
17. R. W. Alder, P. S. Bowman, W.R.P. Steele and D. R. Winterman, *Chem. Commun.*, 723 (1968); M. Fieser and L. F. Fieser, "Reagents For Organic Synthesis, 3, p 22 (1972).
18. L. J. Bollyky, B. G. Roberts, R. H. Whitman, and J. E. Lancaster, *J. Org. Chem.*, 34, 836 (1969).
19. R. D. Trepka, J. K. Harrington and J. W. Belisle, *J. Org. Chem.*, 39, 1094 (1974).
20. B.G. Roberts and R. C. Hirt, *Appl. Spectrosc.*, 21, 250 (1967).
21. W. H. Melhuish, *N. Z. Sci. Tech.*, B37, 142 (1955).