



DCN

NOV 9

Im

DDC FILE COPY AD AO 61 0 47

NPS55-78-016 NAVAL POSTGRADUATE SCHOOL Monterey, California





by

D. P. Gaver and P. A. W. Lewis

August 1978

Approved for public release; distribution unlimited.

78 10 30 044

NAVAL POSTGRADUATE SCHOOL MONTEREY, CALIFORNIA

Rear Admiral T. F. Dedman Superintendent J. R. Borsting Provost

This research was supported by the National Science Foundation and the Office of Naval Research.

Reproduction of all or part of this report is authorized.

This report was prepared by:

MA

D. P. Gaver, Professor Department of Operations Research

P. A. W. Lewis, Professor

P. A. W. Lewis, Professor Department of Operations Research

Reviewed by:

Michael G. Sovereign, Chairman

Department of Operations Research

Released by:

M. Jolles

William M. Tolles Dean of Research

UNCLASSIFIED	
DEDODT DOCUMENTATION DAGE	READ INSTRUCTIONS
CEPORT NUMBER 2. GOVT ACCESSION NO.	BEFORE COMPLETING FORM 3. RECIPIENT'S CATALOG NUMBER
NPS55-78-016	
TITLE (MA SUBILIO)	5. TYPE OF REPORT & PERIOD COVERED
First Order Autoregressive Gamma Sequences	9) Technical reption
and Point Processes,	6. PERFORMING ORG. REPORT NUMBER
	B. CONTRACT OF GRANT NUMBER(4)
D. P. Gaver and P. A. W. Lewis	MCS-77-07587
PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Naval Postgraduate School	
Monterey, Ca. 93940	
1. CONTROLLING OFFICE NAME AND ADDRESS	12 REPORT DATE
Naval Postgraduate School	August 1978
Monterey, Ca. 93940	55
4. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office)	15. SECURITY CLASS. (of this report)
Washington, D.C.	Unclassified
	154. DECLASSIFICATION/DOWNGRADING
DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different tra	n Report)
. KEY WORDS (Continue on reverse elde if necessary and identify by block number) Point processes, time series, Poisson process, Ma	arkov process, simulation
ABSTRACT (Continue on reverse olde if necessary and identify by block number) An autoregressive model that generates Markov con- described. The time series have exponential or of distributions. Various properties of these time	rrelated time series is gamma distributed marginal series are investigated.
D 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE S/N 0102-014-6601 SECURITY CLA	SSIFICATION OF THIS PAGE (When Data Entered

TABLE OF CONTENTS

1.	INTRODUCTION	1
2.	EXPONENTIAL SEQUENCES	5
	2.1. The exponential first-order autoregressive sequence (EAR1)	5
	2.2. Serial correlations, stationarity and mixing	9
з.	SUMS OF STATIONARY INTERVALS	10
4.	JOINT DISTRIBUTIONS FOR THE STATIONARY SEQUENCE	12
5.	ESTIMATION AND DEGENERACY	14
6.	OTHER SOLUTIONS TO THE AUTOREGRESSIVE EQUATION	17
	6.1. The Gamma Autoregressive Process GARI.6.2. The Mixed Exponential Process MEARI6.3. Discussion	18 20 24
7.	MULTIVARIATE SEQUENCES AND NEGATIVE CORRELATION	25
8.	SUMMARY AND CONCLUSIONS	30
	REFERENCES	32

NTIS	White Section
DDC	Buff Section
UNANINGIN	CTD CTD
NISTIFICAT	100
Dista 1	
1	
	1

78 10 30 044

PAGE

FIRST ORDER AUTOREGRESSIVE GAMMA SEQUENCES AND POINT PROCESSES

D. P. Gaver *

and

P. A. W. Lewis

Naval Postgraduate School Monterey, CA 93940

1. INTRODUCTION

The Poisson process is a basic model for point processes (series of events) and can be characterized as a process in which the intervals between events are independent and exponentially distributed. In one of the earliest papers on point processes (Wold, 1948) an attempt was made to generalize the Poisson process by obtaining dependent but marginally exponentially distributed intervals between events. Similarly Cox (1955) attempted to obtain a sequence of random variables with conditionally exponential distributions. Neither of these attempts to generalize the Poisson process led to analytically tractable results. In principle, of course, it is simple to generate a sequence of marginally exponentially distributed random variables with Markov dependence if a bivariate exponential random variable is available. However, despite the recent discovery of many bivariate exponential random

Research of both authors was supported in part by NSF grant MCS-77-07587 at the Naval Postgraduate School and by the Office of Naval Research.

variable generation schemes [Marshall and Olkin (1967), Hawkes (1972); Gaver (1972); Downton (1970)], none seem to have simple enough properties, conditional and otherwise, to lead to analytically and computationally tractable models for a non-independent (Markovian) and easily simulated sequence of marginally exponentially distributed random variables. Interest in the aforementioned processes also may have been damped by the development of physically motivated point process models such as the Poisson cluster process and the self-exciting process, for example see Neyman and Scott (1972). Unfortunately, many of these point process models are somewhat awkward to handle analytically or computationally.

In this paper we show that if one starts with the usual linear, additive autoregressive equation (first-order stochastic difference equation)

 $X_n = \rho X_{n-1} + \varepsilon_n$, $n = 0, \pm 1, \pm 2, ...$ (1.1)

where the innovation sequence $\{\varepsilon_n\}$ is one of i.i.d. random variables, then there is, for $0 \leq \rho < 1$, a distribution for the ε_n 's such that the X_n 's have, marginally, an exponential distribution. The resulting exponential autoregressive process (EAR1) has several attractive features. First, the sequence is obtained as an additive random linear combination of random variables, and is easy to simulate, i.e. to realize on a computer. In simulating a queue, for instance, one can easily obtain sequences of correlated service times or interarrival times, with respectively an i.i.d. exponential sequence or Poisson process as a special case ($\rho = 0$). These

correlated sequences are useful for checking for the sensitivity of standard queueing results to departures from independence, and for other robustness studies.

A second feature is that the EARl process (1.1) is analytically tractable; one can, for instance, obtain the Laplace transform of the distribution of sums of the random variables. This is essential if the process is to be used as a model for point processes and one is to obtain the point spectrum.

A third feature of the EARI process is that its structure leads to an extended model for a sequence of marginally exponentially distributed random variables with the correlation structure of a mixed autoregressive moving average process (ARMA p,q), the orders of the autoregression and the moving average being p and q respectively. Various properties of this exponential process, called the EARMA(p,q) process, are detailed in Lawrance and Lewis (1977), Jacobs and Lewis (1977) and Lawrance and Lewis (1978).

In Section 2 of this paper we discuss the exponential solution of equation (1.1), as well as questions of stationarity and mixing. The distribution of sums of random variables from this process is discussed next; this relates in particular to the use of the process to model intervals between events in point processes. Joint distributions and conditional correlations of two and three random variables are derived in Section 4. The estimation of the exponential parameter λ , the reciprocal of $E(X_n)$, and the correlation parameter ρ are briefly considered in Section 5 for a fixed or random number of observed random variables. It is shown that, because of a certain degeneracy in the process, ρ can be estimated exactly in a long enough sequence.

correlated sequences are useful for checking for the securitie

Other solutions to equation (1.1) are considered in Section 6. In fact random variables for which solutions to (1.1) (in transformed version) exist are known as <u>self-decomposable</u> random variables or random variables of <u>type</u> L; see Feller (1971, p. 588-90). Two cases of interest in modelling point processes are discussed. These require consideration of Gamma-distributed intervals, and intervals with a mixed exponential distribution; questions connected with the latter remain unresolved.

Finally in Section 7 we consider the question of obtaining autoregressive and Markovian exponential sequences with negative correlation. It may be seen that (1.1) has no solution if ρ is negative. However, as soon as multivariate exponential sequences are considered--in particular, multivariate sequences which are antithetic realizations of each other --we discover a method for obtaining processes with exponential or other specified marginals and that exhibit negative correlation.

This process is discussed next, this relates in particular to the one of the process to model infervals between events in point processes. Joint distributions and conditional correlations of two and three tanders veriables are derived in Saction 4. The setimation of the exponential parameter λ , the radiurocal of $E(\lambda_{a})$, and the correlation parameter ρ are briefly considered in Section 5 for a fixed or random number of observed tandom variables. It is shown that, because of a cartain degeneracy in the exponential parameter is a long considered

2. EXPONENTIAL SEQUENCES

In what follows $\{E_n\}$ will always be a sequence of i.i.d. exponentially distributed random variables with parameter λ ;

$$P\{E_{n} \leq x\} = F_{E_{n}}(x) = 1 - e^{-\lambda x} \quad (x \geq 0, \ \lambda > 0)$$

= 0 (x < 0). (2.1)

Also an exponential (λ) random variable will mean a random variable with distribution (2.1).

2.1. The exponential first-order autoregressive sequence (EAR1). The starting point of the work in this paper is the question as to whether the autoregressive equation (1.7)

$$\mathbf{x}_{n} = \rho \mathbf{x}_{n-1} + \varepsilon_{n} = \sum_{j=0}^{\infty} \rho^{j} \varepsilon_{n-j} \qquad |\rho| < 1 \qquad (2.2)$$

has a solution for a given distribution of X_i (note that the expansion in (2.2) as an infinite moving average is valid because $|\rho| < 1$). Now X_{n-1} is a function only of ε_{n-1} , ε_{n-2} , ... and is therefore independent of ε_n . Therefore the Laplace-Stieltjes transform ϕ_{X_n} (s) of the distribution of X_n is, at least for s > 0,

$$\phi_{X_n}(s) = E[\exp(-sX_n)] = E[\exp(-s\rho X_{n-1} + \varepsilon_n)]$$
$$= \phi_{X_{n-1}}(\rho s) \phi_{\varepsilon_n}(s) .$$

Thus

 $\phi_{\varepsilon_{n}}(s) = \frac{\phi_{X_{n}}(s)}{\phi_{X_{n}}(\rho s)}$

Assuming that the X_n sequence is marginally stationary, we get the basic equation

$$\phi_{\varepsilon}(s) = \frac{\phi_{X}(s)}{\phi_{X}(\rho s)}$$

Now it is clear that if we require the X_n to be positive random variables, then if ρ is negative so is ρX_n and we need the error term ε_n , which is independent of X_{n-1} , to make X_n positive. Thus for positive random variables there will clearly be no solution to (2.3) for $\rho < 0$. However if we let $0 \leq \rho < 1$ and, sidestepping the general question of existence and uniqueness of solutions to (2.3), require the X_n 's to be exponential with

$$\phi_{\mathbf{X}}(\mathbf{s}) = \frac{\lambda}{\lambda + \mathbf{s}} \, ,$$

then equation (2.3) forces

$$\phi_{\varepsilon}(\mathbf{s}) = \frac{\lambda}{\lambda + \mathbf{s}} \cdot \frac{\lambda + \rho \mathbf{s}}{\lambda} = \frac{\lambda + \rho \mathbf{s}}{\lambda + \mathbf{s}} \quad (0 \leq \rho < 1, \ \lambda \geq 0)$$
$$= \rho + (1-\rho) \frac{\lambda}{\lambda + \mathbf{s}} \quad (2.4)$$

The latter expression is the Laplace-Stieltjes transform of a random variable, in fact a non-negative random variable which has an atom of mass ρ at zero and which is exponential(λ) if positive. Thus we can write the difference equation generating the series $\{x_n\}$ as

(2.3)

$$X_{n} = \rho X_{n-1} + \epsilon_{n}$$

$$= \begin{cases} \rho X_{n-1} & \text{w.p. } \rho \\ \rho X_{n-1} + E_{n} & \text{w.p. } (1-\rho) \end{cases} \quad 0 \le \rho \le 1$$
(2.5)

$$= \rho X_{n-1} + I_n E_n,$$
 (2.6)

where $\{I_n\}$ is an i.i.d. sequence in which $I_n = 1$ with probability $(1-\rho)$, $I_n = 0$ with probability ρ and $\{E_n\}$ is an i.i.d. sequence of (λ) exponentials. It is easily verified, directly from (2.2) or from the definition of ε_n , that

$$E(\varepsilon_n) = (1-\rho) \frac{1}{\lambda} ; \quad Var(\varepsilon_n) = \frac{1}{\lambda^2} (1-\rho^2) . \quad (2.7)$$

There are several points to be made about the sequence X_n:

- (i) When $\rho = 0$, then $\{X_n\}$ is an i.i.d. sequence of exponential(λ) r.v.s.
- (ii) The representations (2.5) and (2.6) of the process as an (additive) random linear combination of X_{n-1} and E_n , which has a distribution independent of ρ , are sometimes more convenient than is (2.2).
- (iii) Although X_n is strictly a linear process (AR1), it differs from the normal or Gaussian process ((2.2) with ε_n normally distributed of Gaussian) in that the mean, <u>variance and higher moments of the</u> ε_n 's are functions

of the parameter p. Thus one cannot apply general theorems about linear processes to the exponential sequence because such theorems usually assume that the ε_n 's have a distribution which is free of the parameter ρ . As an example, X_n is exponentially distributed even when ρ is close to one, despite theorems for linear processes which assert that in this case the X_n are approximately normally distributed. For instance, if the ε_n 's are independent of ρ then the skewness of the stationary distribution of X can be shown to approach zero as $\rho \uparrow 1$; in the special case of (2.4), this quantity is independent of ρ and equal to 2--the value associated with the exponential d.f.

There are many other sequences of random variables $\{X_n\}$ (iv) with exponentially distributed marginals and the first-order Markov property. In fact, given any bivariate exponential distribution $F_{E_1,E_2}(x_1,x_2)$, assumed for convenience to be absolutely continuous, with conditional density $f_{E_2|E_1}(x_2;x_1)$, then we construct a Markovian sequence with joint density for, say, X_n , X_{n-1} , ..., X_0 as

$$f_{X_{n},...,X_{0}}(x_{n},...,x_{0})$$

$$= f_{E_{n}|E_{n-1}}(x_{n}; x_{n-1}) f_{E_{n-1}|E_{n-2}}(x_{n-1}; x_{n-2})$$

$$\cdots f_{E_{1}|E_{0}}(x_{1};x_{0}) f_{E_{0}}(x_{0}) .$$

An interesting and useful feature of the EARl sequence given by the solution (2.6) of the autoregressive equation is that it is a (random) <u>linear</u> combination of i.i.d. exponential sequences, and is thus easy to simulate on a computer, gives reasonably tractable analytic results and in turn suggests the definition of processes with more complicated correlation structure and exponential marginals (Lawrance and Lewis, 1977, Jacobs and Lewis, 1977, Lawrance and Lewis, 1978). The correlation structures are essentially the same as those for linear processes (Lawrance and Lewis, 1978).

2.2. Serial correlations, stationarity and mixing

Simple computations show that, as is true of any regular Markov process, the serial correlations for the EARL process are

$$c_j = corr(x_n x_{n+j}) = \rho^j$$
, $0 \le \rho \le 1$.

Note that the correlations are always positive. The spectrum of the sequence is

$$E_{+}(\omega) = \frac{1}{\pi} \{1 + 2 \sum_{j=1}^{\infty} c_{j} \cos(j\omega)\} \quad (0 \leq \omega \leq \pi; 0 \leq \rho < 1)$$

$$= \frac{1}{\pi} = \frac{1 - \rho^2}{1 + \rho^2 - 2\rho \cos \omega} .$$

For $\rho = 0$ we have $f_{+}(\omega) = 1/\pi$ since the sequence X_{n} is i.i.d.

Moreover it is not difficult to show that the process is stationary if one takes

$$x_0 = E_0$$
 (2.8)

$$X_{n} = \rho X_{n-1} + \varepsilon_{n} = \rho X_{n-1} + I_{n} E_{n}$$
(2.9)

with ε_n defined by (2.4). This is no surprise, for the process is Markovian and is constructed to have an exponential distribution in the stationary case.

With $E_0 = X_0$ it is shown in Jacobs and Lewis (1977) that the sequence $\{X_n\}$ is strong mixing in the sense of Rosenblatt (1971).

3. SUMS OF STATIONARY INTERVALS

An important aspect of the stationary sequence $\{X_n\}$, especially when it is used to model the intervals in a point process, is the moments and distributions of sums $T_r = X_n + X_{n+1} + \cdots + X_{n+r-1}$. The Laplace-Stieltjes transform of T_r is found directly by noting that, in terms of X_n , (2.2) may be modified to give this representation:

$$x_{n+j} = \rho^{j} x_{n} + \rho^{j-1} \epsilon_{n+1} + \rho^{j-2} \epsilon_{n+2} + \dots + \epsilon_{n+j}, \quad (j = 0, 1, 2, \dots) \quad (3.1)$$

and so

$$\mathbf{T}_{\mathbf{r}} = \sum_{j=0}^{r-1} \mathbf{X}_{n+j} = \mathbf{X}_{n} \left(\frac{1-\rho^{r}}{1-\rho} \right) + \sum_{j=1}^{r-1} \epsilon_{n+j} \left(\frac{1-\rho^{r-j}}{1-\rho} \right).$$

It follows that

$$\phi_{\mathbf{T}_{\mathbf{r}}}(\mathbf{s}) = \mathbf{E}\left[\exp\left(-\mathbf{s}\mathbf{T}_{\mathbf{t}}\right)\right] = \phi_{\mathbf{X}_{\mathbf{n}}}\left[\mathbf{s}\left(\frac{1-\rho^{\mathbf{r}}}{1-\rho}\right)\right] \prod_{j=1}^{\mathbf{r}-1} \phi_{\varepsilon}\left[\mathbf{s}\left(\frac{1-\rho^{\mathbf{r}-j}}{1-\rho}\right)\right] \quad (3.2)$$

If the process starts with the stationary distribution, i.e., if X_n is exponential(λ), then

$$\phi_{\mathbf{T}_{\mathbf{r}}}(\mathbf{s}) = \frac{\lambda}{\lambda + \mathbf{s}\left(\frac{1-\rho^{\mathbf{r}}}{1-\rho}\right)} \prod_{j=1}^{\mathbf{r}-1} \left(\frac{\lambda + \mathbf{s}\rho\left(\frac{1-\rho^{\mathbf{r}-j}}{1-\rho}\right)}{\lambda + \mathbf{s}\left(\frac{1-\rho^{\mathbf{r}-j}}{1-\rho}\right)} \right)$$
(3.3)

Clearly the distribution of T_r can be found explicitly by inverting (3.3), which can be accomplished by expanding in partial fractions. So also can an expression for the interval spectrum. The result is analytically awkward, and will not be quoted. In order to obtain (i) the intensity function, $m_f(t)$, of a point process with EARl intervals, (ii) the point spectrum $g_+(\omega)$, and (iii) the distribution of the counting process N(t) (for definitions see Cox and Lewis, 1966, Sec. 4.5) one must be able to sum $\phi_{T_r}(s)$ over r; there appears to be no neat explicit formula for any of these functions. The variance of T_r may be calculated directly from (3.1) since the innovations $\{\varepsilon_n\}$ are independent:

$$\operatorname{Var}[\mathbf{T}_{\mathbf{r}}] = \frac{1}{\lambda^{2}} \left[\left(\frac{1-\rho^{\mathbf{r}}}{1-\rho} \right)^{2} + \left(\frac{1+\rho}{1-\rho} \right) \left\{ r - 1 - 2\rho \left(\frac{1-\rho^{\mathbf{r}-1}}{1-\rho} \right) + \rho^{2} \left(\frac{1-\rho^{2}(\mathbf{r}-1)}{1-\rho^{2}} \right) \right\} \right] \quad (3.4)$$

Hence for $\rho < 1$ the index of dispersion (Cox and Lewis (1966), p. 71) is

$$J = \lim_{r \to \infty} J_{r} = \lim_{r \to \infty} \frac{\operatorname{Var}[T_{r}]}{r\{E[X_{1}]\}^{2}} = \frac{1+\rho}{1-\rho} = 1 + \frac{2\rho}{1-\rho}$$
(3.5)

As a byproduct of (3.5) we get, for $\rho < 1$, values of the slope of the variance time curve, $\lim_{t \to \infty} V'(t) = V'(\infty)$, the initial points of the spectrum of counts $g_+(\omega)$ and the spectrum of intervals as (Cox and Lewis, 1966, Sect. 4.6)

$$f_{+}(0+) = \frac{J}{\pi}$$

$$V'(\infty) = \lambda J$$

$$g_{+}(0+) = \lambda J/\pi$$

Limit theorems for T_r and the counting process N(t) are given in Jacobs and Lewis (1977).

4. JOINT DISTRIBUTIONS FOR THE STATIONARY SEQUENCE

Any pair X_n , X_{n+r} in the stationary EARl sequence has a bivariate exponential distribution. Consider X_n and X_{n+1} . Directly from the definition (2.5) and (2.7) it can be shown that the conditional random variable X_{n+1} , given $X_n = x$, has an atom of mass ρ at ρx ; otherwise it is, with probability $(1-\rho)$,

an exponential(λ) random variable shifted ρx from the origin. The regression of X_{n+1} on $X_n = x$ is

$$E(X_{n+1}|X_n = x) = x + (1-\rho) \frac{1}{\lambda}$$
, (4.1)

which is linear. Moreover $var(X_{n+1}|X_n = x) = (1-\rho^2)/\lambda^2$, a constant independent of x, and X_{n+1} is never smaller than ρX_n . The Laplace-Stieltjes transform of the bivariate exponential distribution of X_{n+1} and X_n is

The bivariate distribution of $\{X_{n+1}, X_n\}$ has a singular component along the line $X_{n+1} = X_n$, with probability ρ (the first term in (4.2), and a continuous component in the space $X_{n+1} > \rho X_n$ (the second term in (4.2)) which is the joint distribution of X_n and $\rho X_n + E_n$. Note that the distribution is not symmetric in X_{n+1} and X_n , since the transform $\phi_{X_{n+1},X_n}(s_1,s_2)$ is not symmetric in s_1 and s_2 . This simply means that the process is not time-reversible, as is the normal or Gaussian ARl process.

This asymmetry appears in the conditional mean of X_n , given $X_{n+1} = x$, which is obtained from (4.2), as

This asymmetry also becomes evident in the non-linear form of the conditional mean of X_n , given $X_{n+1} = x$, which is obtained from (4.2), as

$$E(X_{n} | X_{n+1} = x) = \frac{1}{\lambda(1-\rho)} \{1 - \exp[-(\frac{1}{\rho} - 1)\lambda x]\} \neq E[X_{n+1} | X_{n} = x]$$

The conditional variance of X_n , given $X_{n+1} = x$, can also be obtained; unlike $var(X_{n+1}|X_n = x)$ it is not a constant.

Higher order directional moments of X_{n+1} , X_n are given in Jacobs and Lewis (1977), and again evidence the directionality of the process:

$$C_{2,1}(1) = E(X_n^2 X_{n+1}) - E(X_n^2) E(X_{n+1}) = \frac{4\rho}{\lambda^3};$$

$$C_{1,2}(1) = E(X_n X_{n+1}^2) - E(X_n) E(X_{n+1}^2) = \frac{2\rho(1+\rho)}{\lambda^3}.$$

Since the process is, from its definition, Markovian, the conditional correlation of X_{n+1} and X_{n-1} , given $X_n = x$, is zero. We do not discuss distributions of triples any further. In principle it is simple to write down transforms of the joint distribution of any set of k X_n 's; one obvious use of this result would be in obtaining the distribution of the sum of X_n , \dots , X_{n+k-1} . However, this was obtained by a direct argument in the previous section.

5. ESTIMATION AND DEGENERACY

Although it is not possible to write down the likelihood equation for the EARI process in a tractable form where, say, we observe X_1, \ldots, X_N , it seems at first sight that good estimates of $1/\lambda$ and ρ can be obtained from the sample mean, $\bar{X} = T_n/n$, whose variance is given at (3.4), and a first-order serial correlation coefficient estimate $\hat{\rho}_1$ with modifications to take account of the fact that the variance equals the mean squared, i.e. $Var(X) = 1/\lambda^2$.

However, there is a degeneracy in the process which makes it possible to find ρ exactly in a long enough run of X_n 's, and to estimate λ more precisely than is possible using straightforward moment methods.

To identify the degeneracy, which we call the <u>zero-defect</u>, we note that in the process there are runs of X_n 's which are equal to the previous value, X_{n-1} , times ρ . Moreover, given that there is a run of length R of this type, R has a "geometric plus one" distribution with parameter ρ ;

$$P{R = i} = (1 - \rho)\rho^{i-1}$$
, $i = 1, 2, 3, ...$ (5.1)

flever and all appreve

$$E(R) = \frac{\rho}{1 - \rho}$$
; $var(R) = \frac{\rho}{(1 - \rho)^2}$

This behavior, or a tendency towards it in data would be very evident in plots of a sample sequence of X_n values.

Thus to estimate ρ in an observed series, let $Z_n = X_{n+1}/X_n$, for $n = 2, 3, \ldots$. Then

$$Z_{n} = \frac{X_{n+1}}{X_{n}} = \rho \qquad \text{if } X_{n+1} = \rho X_{n} \quad \text{w.p. } \rho$$
$$= \rho + \frac{E_{n+1}}{X_{n}} \qquad \text{if } X_{n+1} \neq \rho X_{n} \quad \text{w.p. } (1-\rho)$$

The probability that $Z_n = \rho$, or $Z_n \neq \rho$, is independent of previous Z_n 's and $P\{Z_n = \rho\} = \rho$. Hence if we observe the minimum value of Z_n , $n = 2, 3, \ldots$ until we get a tie, that tie value is ρ . The time to obtain this tie is the sum of two "geometric plus one" random variables, with total mean $2/\rho$ and total variance $2(1-\rho)/\rho^2$. The implication is that we can find ρ exactly after a random number of observations for the present model. Moreover since one then knows those X_n 's which are made up of ρX_{n-1} plus an E_n , it is possible to compute the E_n 's exactly (except for two) and estimate $E(X) = 1/\lambda$ directly from the observed E_n 's.

For a fixed length observation of the process, say X_1, \ldots, X_N , the probability that ρ can be obtained exactly is the probability that the sum of two "geometric plus one" random variables is less than or equal to N. If two minimum values are not observed by time N, a good estimate of ρ will be the minimum of the Z_i 's. This will either be ρ , or ρ plus a small bias term. The mean can be estimated as the sample average in the usual way.

This degeneracy in sequences of positive random variables generated from the stochastic difference equation seems to be inherent in the present procedure. Slightly more realistic but complicated first-order autoregressive exponential processes without this zero defect will be discussed elsewhere.

6. OTHER SOLUTIONS TO THE AUTOREGRESSIVE EQUATION

Random variables X for which the transformed first-order autoregressive equation

$$\frac{\phi_{\mathbf{X}}(\mathbf{s})}{\phi_{\mathbf{X}}(\mathbf{\rho}\mathbf{s})} = \phi_{\varepsilon}(\mathbf{s})$$
(6.1)

has a solution $\phi_{\varepsilon}(s)$ for each $0 < \rho < 1$ which is the transform of a distribution function are called <u>random variables of class L</u> (Feller, 1971, p. 588), <u>or self-decomposable random variables</u>. Although there has been much recent interest in finding class L random variables, the connection with the first-order autoregressive process does not seem to have been made. On the other hand there have been some attempts to find non-normal solutions to the autoregressive equation without connecting them to class L theory (see, e.g. Bernier, 1970).

The limitation of the theory of class L random variables as it relates to the present work is that it requires a solution for each $0 < \rho < 1$, which is the case for exponential(λ) random variables X. This full range of ρ is desirable, but may not occur. We do not explore the full connection here, but consider only two types of random variables, Gamma distributed random variables and mixed exponential random variables. These are frequently used as alternatives to the exponential in modelling stochastic phenomena such as response times in queues.

6.1. The Gamma Autoregressive Process GARL.

A necessary condition for X to be in the class L is that it be infinitely divisible, and therefore we consider Gamma (λ, k) variables with density

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{\lambda^{\mathbf{k}} \mathbf{x}^{\mathbf{k}-1} \mathbf{e}^{-}}{\mathbf{x}^{\mathbf{k}} \Gamma(\mathbf{k})} \qquad (\lambda \ge 0; \mathbf{k} > 0; \mathbf{x} \ge 0)$$
(6.2)

and Laplace-Stieltjes transform (of the stationary distribution)

$$\phi_{\mathbf{X}}(\mathbf{s}) = \left(\frac{\lambda}{\lambda + \mathbf{s}}\right)^{\mathbf{k}} . \tag{6.3}$$

Then the solution to (6.1) is

$$\phi_{\varepsilon}(\mathbf{s}) = \left(\frac{\lambda + \rho \mathbf{s}}{\lambda + \mathbf{s}}\right)^{\mathbf{k}} = \left(\rho + (1-\rho) \frac{\lambda}{\lambda + \mathbf{s}}\right)^{\mathbf{k}}$$
(6.4)

It follows from a result reported by Feller ((1971), p. 452) or directly from Theorem 1, p. 450) that $[(\lambda + \rho s)/(\lambda + s)]$ is the transform of an infinitely divisible distribution, and hence (6.3) is actually the transform of (an infinitely divisible) distribution function for every real k > 0. Thus we can in principle generate an autoregressive process with gamma marginals (6.2) by utilizing the $\{\varepsilon_n\}$ process characterized by (6.3). Here are a few simple special cases.

$$k = 2: \quad \phi_{\varepsilon}(s) = \rho^{2} + 2\rho(1-\rho) \frac{\lambda}{s+\lambda} + (1-\rho)^{2} \left(\frac{\lambda}{s+\lambda}\right)^{2}; \quad (6.5)$$

$$k = 3: \quad \phi_{\varepsilon}(s) = \rho^{3} + 2\rho^{2}(1-\rho) \frac{\lambda}{s+\lambda} + 2\rho(1-\rho)^{2} \left(\frac{\lambda}{s+\lambda}\right)^{2} + (1-\rho)^{3} \left(\frac{\lambda}{s+\lambda}\right)^{3}. \quad (6.6)$$

Thus for k = 2 the random variable ε may be thought of and realized as a convex mixture of a degenerate random variable with mass at zero and an exponential(λ) and a Gamma (λ ,2) distribution. Alternatively if E_1 and E_2 are independent exponential(λ) random variables, the sum is chosen with probability $(1-\rho)^2$, E_1 is chosen with probability $2\rho(1-\rho)^2$, and neither of E_1 or E_2 , but rather zero, is chosen with probability ρ^2 .

Unfortunately, no really simple way of generating random variables with Laplace-Stieltjes transform (6.4) for noninteger k is known, though representation as an infinite sum of weighted Gamma variables is possible (Bernier (1970)). Moreover for a case of much interest k < 1, so that X is overdispersed relative to an exponential random variable, and the degeneracy or zero-defect of weight ρ^k becomes very prominent.

As was true for the exponential case, we have for the correlations, which are all positive,

 $c_j = \rho^j \qquad 0 \le \rho \le 1;$

also

 $\mathbf{E}(\boldsymbol{\varepsilon}) = \lambda(\mathbf{1} - \boldsymbol{\rho})$

$$Var(\varepsilon) = \lambda^2 (1-\rho^2)/k$$

As $k \neq \infty$, the degeneracy disappears and X tends to become normally distributed.

6.2. The Mixed Exponential Process MEARL

Consider now X to be a mixture of two exponential densities

$$f_{X}(x) = \pi_{1}^{\lambda_{1}}e^{-\lambda_{1}x} + \pi_{2}^{\lambda_{2}}e^{-\lambda_{2}x}$$
,

$$x \ge 0, \ \pi_1 = 1 - \pi_2 \ge 0, \ \lambda_1 < \lambda_2$$
 (6.7)

For $\pi_1 = 0$, $\pi_1 = 1$, and/or $\lambda_1 = \lambda_2$ this of course reduces to an exponential density. There are two main cases to consider, besides these special cases.

If $0 \leq \pi_1 \leq 1$, then we always have a convex mixture of exponential densities: $f_X(x)$ is a proper density function and the coefficient of variation of X lies between 1 and infinity (Cox, 1964), so that X is overdispersed relative to an exponential random variable; X may be generated as a mixture of two exponential random variables with parameters λ_1 and λ_2 , and X is infinitely divisible (Feller (1971), p. 452).

If π_1 is greater than 1, so that π_2 is negative, a necessary and sufficient condition for $f_X(x)$ to be a p.d.f. is that $\pi_1\lambda_1 + \pi_2\lambda_2 \ge 0$ or equivalently that $\pi_1 \le [1 - (\lambda_1/\lambda_2)]^{-1}$.

Here is alternative way of writing the transform of X. First, and directly,

$$\phi_{\mathbf{X}}(\mathbf{s}) = \pi_1 \frac{\lambda_1}{\lambda_1 + \mathbf{s}} + \pi_2 \frac{\lambda_2}{\lambda_2 + \mathbf{s}}$$
(6.8)

Now since $\lambda_2 > \lambda_1$ factor to obtain

$$\phi_{\mathbf{X}}(\mathbf{s}) = \frac{\lambda_2}{\lambda_2 + \mathbf{s}} \left\{ \pi_2 + \pi_1 \frac{\lambda_1}{\lambda_2} \left(\frac{\lambda_2 + \mathbf{s}}{\lambda_1 + \mathbf{s}} \right) \right\}$$
$$= \frac{\lambda_2}{\lambda_2 + \mathbf{s}} \left\{ 1 - \pi_1 \left(1 - \frac{\lambda_1}{\lambda_2} \right) + \pi_1 \left(1 - \frac{\lambda_1}{\lambda_2} \right) \frac{\lambda_1}{\lambda_1 + \mathbf{s}} \right\}. \quad (6.9)$$

Now even if $\pi_1 > 1$ ($\pi_2 < 0$) but $\gamma = \pi_1 \left(1 - \frac{\lambda_1}{\lambda_2}\right) < 1$, (6.10)

or equivalently

$$\pi_2 \lambda_2 + \pi_1 \lambda_1 > 0 \tag{6.11}$$

then $\phi_X(s)$ is the transform of the sum of an exponential (λ_2) random variable and a random variable having the distribution of the ε -innovation of an EARl process, see (2.4). It follows that X is infinitely divisible.

It is thus worth investigating whether one can generate an autoregressive process $\{x_n\}$ with mixed exponential marginals for some or all ρ , the latter being the question as to whether the mixed exponential random variable is in the class L. We have, directly

$$\phi_{\varepsilon}(\mathbf{s}) = \frac{\phi_{\mathbf{X}}(\mathbf{s})}{\phi_{\mathbf{X}}(\rho \mathbf{s})} = \frac{\frac{\pi \mathbf{1}^{\lambda} \mathbf{1}}{\mathbf{s} + \lambda_{\mathbf{1}}} + \frac{\pi \mathbf{2}^{\lambda} \mathbf{2}}{\mathbf{s} + \lambda_{\mathbf{2}}}}{\frac{\pi \lambda_{\mathbf{1}}}{\rho \mathbf{s} + \lambda_{\mathbf{1}}} + \frac{\pi \mathbf{2}^{\lambda} \mathbf{2}}{\rho \mathbf{s} + \lambda_{\mathbf{2}}}}$$

$$= \frac{(\lambda_1 + \rho \mathbf{s})(\lambda_2 + \rho \mathbf{s})(\lambda_1 \lambda_2 + \pi_1 \lambda_1 \mathbf{s} + \pi_2 \lambda_2 \mathbf{s})}{(\lambda_1 + \mathbf{s})(\lambda_2 + \mathbf{s})(\lambda_1 \lambda_2 + \pi_1 \lambda_1 \rho \mathbf{s} + \pi_2 \lambda_2 \rho \mathbf{s})}$$
(6.12)

After considerable simplification this Laplace-Stieltjes transform can be written as a mixture of a degenerate random variable with mass ρ at zero, and with probability $(1-\rho)$ (possibly) a random variable Y which has (possibly) density function

$$f_{\mathbf{Y}}(\mathbf{x}) = \gamma_1 \lambda_1 e^{-\lambda_1 \mathbf{x}} + \gamma_2 \lambda_2 e^{-\lambda_2 \mathbf{x}} + \gamma_3 \frac{C}{\rho} e^{-\mathbf{x}C/\rho}$$
(6.13)

where $\gamma_1 + \gamma_2 + \gamma_3 = 1$, $\lambda_1 < \lambda_2$, $\pi_1 > 0$, $\pi_1 \neq 1$ and

$$C = \frac{\lambda_{1}\lambda_{2}}{\pi_{1}\lambda_{1} + (1-\pi_{1})\lambda_{2}};$$

$$\gamma_{1} = \frac{(\lambda_{2} - \rho\lambda_{1})(C - \lambda_{1})}{(\lambda_{2} - \lambda_{1})(C - \rho\lambda_{1})} = p_{1} ; \qquad (6.14)$$

$$\gamma_{2} = \frac{(\lambda_{2} - C)}{(\lambda_{2} - \lambda_{1})} \times \frac{(\rho\lambda_{2} - \lambda_{1})}{(\rho\lambda_{2} - C)} = p_{2} \frac{(\lambda_{1} - \rho\lambda_{2})}{(C - \rho\lambda_{2})}; \quad (C \neq \rho\lambda_{2}) \quad (6.15)$$

$$\gamma_{3} = \frac{(C - \lambda_{1})(\lambda_{2} - C)}{(C - \rho\lambda_{1})} \times \frac{1}{(C - \rho\lambda_{2})} = p_{3} \times \frac{1}{(C - \rho\lambda_{2})}; \quad (6.16)$$

Now if π_1 is restricted to be $0 < \pi_1 < 1$, then $\lambda_1 < C < \lambda_2$ and $p_1 > 0$, $p_2 > 0$, $p_3 > 0$. Then it can be shown that the signs of γ_1 , γ_2 , γ_3 depend on whether $0 < \rho \le \lambda_1/\lambda_2$, $\lambda_1/\lambda_2 < \rho < C/\lambda_2$ or $C/\lambda_2 < \rho < 1$:

- (i) γ_1 is always positive;
- (ii) γ_2 and γ_3 are positive if $0 < \rho \le \lambda_1/\lambda_2$;
- (iii) γ_2 is negative, γ_3 is positive if $\lambda_1/\lambda_2 < \rho < C/\lambda_1$;
 - (iv) γ_2 is positive, γ_3 is negative if $C/\lambda_2 < \rho < 1$.

While this result is relatively simple, the only case in which it has been possible to establish that $f_{Y}(x)$ is a p.d.f. is that in (ii), i.e. $0 < \rho \leq \lambda_1/\lambda_2$. The ratio λ_1/λ_2 is roughly related to the overdispersion of X relative to the exponential distribution; the smaller the value λ_1/λ_2 , the greater the dispersion and the smaller the admissible range of ρ . It seems probable that $f_{Y}(x)$ is not a density function for all ρ ; at all events efforts to prove that it is in the class L using (6.10) and characterization theorems (Feller, 1971) have failed at the time of this writing. A complete understanding is not now available.

6.3. Discussion

Note the differences between the mixed exponential autoregressive sequence and the Gamma (k < 1) autoregressive sequence; the former appears to require a restricted range for ρ , ε can be simulated, and the zero defect has probability ρ ; the latter sequence is valid for $0 \leq \rho < 1$, ε cannot at present be simulated, and the zero-defect has probability $\rho^{k} > \rho$. The magnitude of this zero-defect alone probably makes the Gamma process less useful than the mixed exponential process. The tail behavior of the marginal distributions is also different.

Laplace-Stieltjes transforms of distribution of sums, T_n of X_n 's are obtained from (3.2) for both the Gamma and mixed exponential autoregressive sequences. These cases are still under study.

While this result is relatively simple, the only case in which is has have been possible to establish that $f_{\mu}(x)$ is a p.d.f. is that is (1), i.e. $0 . The ratio <math>h_1/h_2$ is roughly called to the exponential distribution; the evaluation of X relative to the exponential distribution; the evaluation of X relative to the exponential distribution; the evaluation of X relative to the evaluation of the second of the se

7. MULTIVARIATE SEQUENCES AND NEGATIVE CORRELATION

Even though the autoregressive equation (1.1) has no solution representing a positive random sequence when $\rho < 0$ it is possible to modify it so as to produce negatively correlated exponential sequences. These will not be individually Markovian, but will have Markovian properties in a bivariate sense. A suggestion as to how to proceed comes from writing X_n in terms of the innovations $\{\varepsilon_n\}$ as in (2.2):

$$x_{n} = \sum_{j=0}^{n} \rho^{j} \varepsilon_{n-j} = \varepsilon_{n} + \rho \varepsilon_{n-1} + \rho^{2} \varepsilon_{n-2} + \dots + \rho_{n} \varepsilon_{0} . \qquad (7.1)$$

Now consider generating two (or more) series in parallel, using independent pairs of innovations, $(\varepsilon_n, \varepsilon_n')$, for which if desired $Cor(\varepsilon_n, \varepsilon_n') < 0$; e.g. for $0 \le \rho < 1$ and starting with the first innovation.

Model 1:
$$X_n = \rho X'_{n-1} + \varepsilon_n = \varepsilon_n + \rho \varepsilon'_{n-1} + \rho^2 \varepsilon_{n-2} + \rho^3 \varepsilon'_{n-3} + \cdots$$

 $X'_n = \rho X_{n-1} + \varepsilon'_n = \varepsilon'_n + \rho \varepsilon_{n-1} + \rho^2 \varepsilon'_{n-2} + \rho^3 \varepsilon_{n-3} + \cdots$
(7.2)

Model 2: $x_n = \rho x_{n-1} + \varepsilon_n = \varepsilon_n + \rho \varepsilon_{n-1} + \rho^2 \varepsilon_{n-2} + \rho^3 \varepsilon_{n-3} + \cdots$ $x'_n = \rho x'_{n-1} + \varepsilon'_n = \varepsilon'_n + \rho \varepsilon'_{n-1} + \rho^2 \varepsilon'_{n-2} + \rho^3 \varepsilon'_{n-3} + \cdots$ (7.30 In order for $\{x_n\}$ and $\{x_n'\}$ to exhibit asymptotically exponential marginal distribution, it is clear that the marginal innovation sequences $\{\varepsilon_n\}$ and $\{\varepsilon_n'\}$ must themselves be iid with an atom ρ at zero, and otherwise an exponential distribution. This can be accomplished in several ways; here are two:

(i) Let

where $\{U_n\}$ is a sequence of uniform (0,1) "random numbers"; $\{E_n\}$ and $\{E'_n\}$ are then called <u>antithetic</u>, see Hammersley and Handscomb (1964), and are maximally negatively correlated, having correlation $1 - (\pi^2/6) = -0.6449$; see Moran (1967). Each series in Models 1, 2 receive independent innovations of exactly the type described in Section 2 and that were there shown to lead to exponential marginals as $n \neq \infty$. In this case

$$E[\varepsilon,\varepsilon'] = (1-\rho) \frac{1}{\lambda^2} \int_0^1 \ln u \ln (1-u) du = (1-\rho) \frac{1}{\lambda^2} \left[2 - \frac{\pi^2}{6}\right] \quad (7.5)$$

and

$$\operatorname{Cov}(\varepsilon,\varepsilon') = (1-\rho) \frac{1}{\lambda^2} \left[2 - \frac{\pi^2}{6} \right] - \frac{(1-\rho)^2}{\lambda^2}$$
(7.6)
Also
$$\operatorname{Cov}(\varepsilon_n,\varepsilon_n') > 0 \quad \text{if} \quad \rho > (\pi^2/6 - 1).$$

(ii) If the distribution function of ϵ_n is F_{ϵ_n} , then in true antithetic fashion determine

$$\varepsilon_n = F_{\varepsilon}^{-1}(U_n); \qquad \varepsilon'_n = F_{\varepsilon}^{-1}(1-U_n) \qquad (7.7)$$

In particular, if

$$F_{\varepsilon_n}(x) = \begin{cases} 0 & \text{if } x < 0 \\ \rho & \text{if } x = 0 \\ \rho + (1-\rho)(1-e^{-\lambda x}) & \text{if } x > 0 \end{cases}$$
(7.8)

Then innovation leading to marginal exponentials(λ) are obtained; see Section 2. In this case

$$\varepsilon_{n} = 0 \qquad \text{if } U_{n} \leq \rho; \quad \varepsilon_{n}' = 0 \qquad \text{if } U_{n} \geq 1-\rho;$$

$$= -\frac{1}{\lambda} \ln\left(\frac{1-U_{n}}{1-\rho}\right) \quad \text{if } U_{n} > \rho; \qquad = -\frac{1}{\lambda} \ln\left(\frac{U_{n}}{1-\rho}\right) \quad \text{if } U_{n} < 1-\rho \qquad (7.9)$$

Consequently

Free and the 27 = (strates theo

and hence

$$\operatorname{Cov}(\varepsilon_{n},\varepsilon_{n}') = \frac{1}{\lambda^{2}} \int_{\rho}^{1-\rho} \ln \frac{u}{1-\rho} \ln \frac{1-u}{1-\rho} \, du - \frac{1-\rho}{\lambda}^{2}, \qquad 0 \le \rho \le \frac{1}{2}$$

$$= -\frac{1-\rho}{\lambda}^{2}$$
, $\frac{1}{2} < \rho \le 1$. (7.11)

Scheme (ii) is capable of generating negative correlations of greater magnitude than is Scheme (i), as is clear from examining the situation $1/2 < \rho$.

The expansions (7.2) and (7.3) allow the computation of the lagged cross covariances between X_n and X'_n . For $n \rightarrow \infty$ we find

Model 1:
$$\operatorname{Cov}(X_{n+j}, X_n') \sim \frac{\rho^j}{1-\rho^2} \operatorname{Cov}(\varepsilon, \varepsilon')$$
, $j = 0, 1, 2, \dots$ (7.12)
Model 2: $\operatorname{Cov}(X_{n+j}, X_n') \sim \frac{\rho^j}{1-\rho^2} \times \begin{cases} \operatorname{Cov}(\varepsilon, \varepsilon') & \text{for } j = 0, 2, 4, \dots \\ \operatorname{Var}(\varepsilon) & \text{for } j = 1, 3, 5, \dots \end{cases}$ (7.13)

Of interest is the fact that the series of Model 2 may in fact exhibit negative correlations; for, directly from (7.2), and as $n + \infty$,

$$\operatorname{Cov}(X_{n+1}, X_n) = \operatorname{Cov}(X_{n+1}, X_n) \sim \frac{\rho}{1-\rho^2} \operatorname{Cov}(\varepsilon, \varepsilon')$$

 $\operatorname{Cov}(X_{n+2}, X_n) = \operatorname{Cov}(X_{n+2}, X_n) \sim \frac{\rho^2}{1-\rho^2} \operatorname{Var}(\varepsilon)$

...

$$\operatorname{Cov}(X_{n+j},X_n) = \operatorname{Cov}(X_{n+j}',X_n') \sim \frac{\rho^j}{1-\rho^2} \times \begin{cases} \operatorname{Cov}(\varepsilon,\varepsilon'), & j = 1,3,5,\dots \\ \operatorname{Var}(\varepsilon), & j = 2,4,6,\dots \end{cases}$$

Consequently if scheme (i) or (ii) above, or a multitude of other possibilities, generate the innovation pairs, then the odd-numbered covariances will in fact be negative.

Clearly the above generation scheme can be generalized, e.g. by including more equations, $\{X_n\}$, $\{X_n'\}$, $\{X_n''\}$, etc., and by allowing the innovations to have different marginal properties. Further investigations are planned.

Two other marginal distributions for the X₀'s have been studied, the Gamma distribution and the mixed exponential. The former is known to be a type [1,1,2, sufficient [6,1] for all b, while the mixed exponential appears to satisfy (6,1) only for a limited range of a - Other type [distributions will be investigated in the context of the modelling of independent stoches the sequences stawhere

Extensions of the Eirsi-order catorogressive structure for exponential marginals to bigoor-order autorogressions, woving averages and mixed surprogressive-moving average structure has been given by Lewrance and Lewis (1977), Jacobs and Levis (1977) and Lawrance and Lewis (1978), The possibility of extends

8. SUMMARY AND CONCLUSIONS

We have presented a simple, autoregressive Markovian sequence $\{X_n\}$ of exponential variates EARl which is an additive random linear combination of the previous value, X_{n-1} , and an independent exponential random variable. The simplicity of this structure allows one to model in an intuitive way dependencies in stochastic systems. Jacobs (1978) has considered cyclic queues with EARl service times and found that the correlation may produce a significant effect; more general queueing schemes which generate multivariate exponential sequences are given by Lewis and Shedler (1978).

Maxima of the X_n in the EARl process have been studied by Chernick (1977).

Two other marginal distributions for the X_n 's have been studied, the Gamma distribution and the mixed exponential. The former is known to be a type L, i.e. satisfies (6.1) for all ρ , while the mixed exponential appears to satisfy (6.1) only for a limited range of ρ . Other type L distributions will be investigated in the context of the modelling of independent stochastic sequences elsewhere.

Extensions of the first-order autoregressive structure for exponential marginals to higher-order autoregressions, moving averages and mixed autoregressive-moving average structures has been given by Lawrance and Lewis (1977), Jacobs and Lewis (1977) and Lawrance and Lewis (1978). The possibility of extending

the EARl structure depends on the fact that the $\{\varepsilon_n\}$ error sequence is a mixture of a random variable with mass at zero and an exponential distribution.

Three other possibilities will be detailed elsewhere-a slight complication which brings in another parameter but gets rid of the "zero-deficiency" in the EARL process; extension of mixed correlation structures to give negative correlations, and further multivariate extensions.

Finally we note that it is possible to introduce nonstationarity and dependence on concomitant variables into the sequence by multiplying X_n by, say,

$$\lambda(n) = \exp\{\sum_{j=0}^{r} \alpha_{j} z_{j}(n)\},\$$

as was done in Cox and Lewis (1966, Ch. 3, ii). It would be of interest to see how the methods given in Cox and Lewis extend to the case where there is correlation present between the X_n 's.

Lawrance A.J. and Lewis, D.A.W. (1977). A moving average exponential

REFERENCES

Bernier, J. (1970). Inventaire des modeles et processus stochastique applicables de la description des deluts journaliers des riviers. Revue de l'Institut International de Statistique, <u>38</u>, 50-71.

Chernick, M. R. (1977). A limit theorem for the maximum of an exponential autoregressive process. Tech. Report No. 14, SIMS, Dept. of Statistics, Stanford University.

Cox, D.R. (1962). Renewal Theory. Methuen, London.

Cox, D.R. (1955). Some statistical methods connected with series of events. J. Roy. Statist. Soc. B, <u>17</u>, 129-164.

Cox, D.R. and Lewis, P.A.W. (1966). The Statistical Analysis of Series of Events, Methuen, London; Wiley, New York.

Downton, F. (1970). Bivariate exponential distributions of reliability theory. J. R. Statist. Soc. B, 32, 408-417.

Feller, W. (1971). "An Introduction to Probability Theory and Its Applications." Wiley; New York.

Gaver, D.P. (1972). Point process problems in reliability. In Stochastic Point Processes, P.A.W. Lewis (ed.), Wiley, New York, 775-800,

Hammersley, J. and Handscomb, D.C. (1964). Monte Carlo Methods, Wiley, New York.

Hawkes, A.G. (1972). Mutually exciting point processes. In Stochastic Point Processes, P.A.W. Lewis (ed.), Wiley, New York, 261-271.

Jacobs, P.A. (1978). A closed cyclic queueing network with dependent exponential service times. J. Appl. Prob. (to appear).

Jacobs, P.A. and Lewis, P.A.W. (1977). A mixed autoregressivemoving average exponential sequence and point process (EARMA 1,1). Adv. Appl. Prob. 9, 87-104.

Lawrance A.J. and Lewis, P.A.W. (1977). A moving average exponential point process (EMA1). J. Appl. Prob. 14, 98-113.

Lawrance, A.J. and Lewis, P.A.W. (1978). An exponential autoregressive-moving average process EARMA(p,q): Definition and correlational properties. Naval Postgraduate School Report NPS55-78-1. Lewis, P.A.W. and Shedler, G.S. (1978). Analysis and modelling of point processes in computer systems. Bull. ISI (to appear).

Neyman, J. and Scott, E. (1972). Processes of clustering and applications. In <u>Stochastic Point Processes</u>, P.A.W. Lewis (ed.), Wiley, New York, 646-681.

Thorin, O. (1977), On the infinite divisibility of the Pareto distribution. Scand. Actuarial J., 31-40.

Wold, H. (1948). Sur les processus stationnaires ponctuels. Colloques Int. Cent. Natn. Rech. Scient. <u>13</u>, 75-86.

DISTRIBUTION LIST

No. of Copies

1

1

1

1

1

,1

1

1

1

STATISTICS AND PREBABILITY FROGRAM CFFICE OF NAVIL RESEARCH CODE 436 AFLINGTON VA 22217

CFFICE OF NAVAL RESEARCH NEW YORK AREA CFFICE 715 BRDAEWAY - STH FLOOR ATTN: ER. ROBER GRAFTEN NEW YORK, NY 10003

DIRECTOR CFFICE OF NAVAL RESEARCH ERANCH OFF 536 SCUTH CLAFK STREET ATTN: DEPUTY AND CHIEF SCIENTIST CHICAGO, IL 60605

LIERARY NAVAL OCEAN SYSTEMS CENTER SAN DIEGO CA

92152

NAVY LIBRAFY NATIONAL SPACE TECHNOLOGY LAB ATTN: NAVY LIERARIAN BAY ST. LOUIS MS 39522

NAVAL ELECTRONIC SYSTEMS COMMAND NAVELEX 32C NATIONAL CENTER NO. 1 ARLINGTON VA 20360

DIFECTOR NAVAL REAEARCH LABORATORY ATTN: LIERARY (ONRL) CODE 2025 WASHINGTON, C.C.

20375

DEFENSE COCUMENTATION CENTER CAPERON STATICN ALEXANDRIA VIFGINIA 22314

TECHNICAL INFORMATION CIVISION NAVAL RESEARCH LABORATORY WASHINGTON, D. C.

.1

1

1

1

1

1

1

OFFICE OF NAVAL RESEARCH SAN FRANCISCO AREA OFFICE 760 MARKET STREET SAN FRANCISCO CALIFORNIA 94102 TECHNICAL LIBRARY NAVAL CRENANCE STATION INCIAN HEAD MARYLAND 2064C NAVAL SHIP ENGINEERING CENTER PHILADELPHIA DIVISION TECHNICAL LIBRARY PHILADELPHIA PENNSYLVANIA 19 19112 BUREAU OF NAVAL PRESONNEL DEFARTMENT OF THE NAVY TECHNICAL LIBRARY WASHINGTON C. C. 20370 LIERARY CCCE 0212 NAVAL POSTGRACLATE SCHCCL MENTEREY CALIFERNIA 93940 PRCF. M. AECEL-HAMEED DEPARTMENT OF MATHEMATICS UNIVERSITY OF NORTH CAROLINA CHARLOTTE NC 28223 PROF. T. W. ANDERSON DEFARTMENT OF STATISTICS STANFORD UNIVERSITY STANFERE , CALIFERNIA 54365 PRCF. F. J. ANSCOMBE DEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN CONNECTICLT C6520 PROF. L. A. ARCIAN INSITIUTE OF INCUSTRIAL ACMINISTRATICN UNION COLLEGE SCIENFCIADY, NEW YORK 12308

1

1

1

1

1

1

1

1

1

1

PRCF. C. R. BAKER DEPARTMENT OF STATISTICS UNIVERSITY OF NOTRH CAFCLINA CHAPEL HILL, NOFTH CAROLINA 27514 PRCF. R. E. DECHOFER DEFARTMENT OF OPERATIONS RESEARCH CORNELL UNIVERSITY ITHACA NEW YORK 14850

FRCF. N. J. BERSHAC SCHOOL OF ENGINEERING UNIVERSITY OF CALIFORNIA IRVINE CALIFORNIA 92664

P. J. BICKEL CEFARTMENT OF STATISTICS UNIVERSITY OF CALIFORNIA BERKELEY , CALIFCRNIA

\$4720 . •

FROF. F. W. BLOCK DEPARTMENT OF MATHEMATICS UNIVERSITY OF PITTSBURGH FITTSBURGH FA

15260

PROF. JCSEPH BLUM DEPT. OF MATHEMATICS, STATISTICS AND COMPLTER SCIENCE THE AMERICAN UNIVERSITY WASHINGTON CC 20016

PROF. R. A. BRADLEY DEFARTMENT OF STATISTICS FLCRIDA STATE UNIVERSITY TALLAHASSEE , FLORIDA 323C6

FROF. R. E. BARLOW OPERATIONS RESEARCH CENTER COLLEGE OF FNGINGERING UNIVERSITY OF CALIFORNIA BERKLEY CALIFORNIA 84730 CALIFCENIA 94720

MR. C. N EFENETT NAVAL COASTAL SYSTEMS LAECRATORY CCDE P761 PANAMA CITY, FLLRIDA 32401

1_

1

1

1

1

1

1

1

1

PRCF. L. N. BHAT COMPUTER SCIENCE / OPERATIONS RESEARCH CENTER SOUTHERN METHODIST UNIVERSITY DALLAS TEXAS 75275

FRCF. W. R. ELISCHKE DEPT. OF QUANTITATIVE BUSINESS ANALYSIS UNIVERSITY CF SCUTHERN CALIFORNIA LOS ANGELES, CALIFORNIA 90007

CR. DERRILL J. BCRDELON NAVAL UNDEFWATER SYSTEMS CENTER CODE 21 NEWPORT PI

02840

J. E. COYER JR DEPT. OF STATISTICS SOUTHERN METHODIST UNIVERSITY DALLAS TX

75275

DR. J. CHANDRA U. S. ARMY RESEARCH F. G. EDX 12211 RESEARCH TRIANGLE PARK, NOFTH CARCLINA 27706

FROF. F. CHERNOFF DEPT. CF MATHEMATICS MASS INSTITUTE CF TECHNOLOGY CAMBRIDGE MASSACHUSETTS 02139

FFOF. C. EERMAN DEFARTMENT OF CIVIL ENGINEERING AND ENGINEERING MECHANICS COLUMEIA UNIVERSITY NEW YORK 10027

PRCF. R. L. DISNEY VIRGINIA PCLYTECHNIC INSTITUTE AND STATE UNIVERSITY DEFT. CF INCUSTRIAL ENGINEERING AND OPERATIONS RESEARCH ELFCKSEURG, VA 24061

MR. J. DEWLING DETENSE LOGISTICS STUDIES INFORMATICN EXCHANGE ARMY LOGISTICS MANAGEMENT CENTER FORT LEE VIFGINIA 20350

1

1

1

1

1

1

1

1

FRCF J. C. ESARY CEPT. OF OPERATIONS RESEARCH AND ADMINISTRATIVE SCIENCES NAVAL POSTGRADLATE SCHOOL MONTEREY CALIFCENIA 93940

DF. P. J. FISCHER DEFENSE COMMUNICATIONS AGENCY 1860 WIEHLE AVENUE RESTON VIRGINIA 22070

PRCF. D. P. GAVER CEPT. CF OPERATIONS RESEARCH NAVAL POSTGRADUATE SCHOOL MONTEREY CA 93940

MR. GENE F. GLEISSNER AFFLIED MATHEMATICS LABCRATCRY CAVID TAYLOR NAVAL SHIP RESEARCH AND DEVELOFMENT GENTER BETHESDA MD

FROF. S. S. GLPTA DEPARTMENT OF STATISTICS PURCUE UNIVERSITY LAFAYETTE INDIANA 47907

FFOF. C. L. HANSCH DEPT OF MATH. SCIENCES STATE UNIVERSITY OF NEW YCRK, BINGHAMTON BINGHAMTEN 13901

PROF. F. J. HARRIS CEPT. CF ELECTRICAL ENGINEERING SAN DIEGC STATE UNIVERSITY SAN DIEGC CA

92182

PROF. L. F. HERBACH DEFT. CF OPERATIONS RESEARCH AND SYSTEMS ANALYSIS FOLYTECHNIC INSTITUTE OF NEW YORK ERCOKLYN 11201 NY

FRCF. M. J. FINICH DEPARTMENT OF ECONOMICS VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACKSHURG VIRGINIA 24061

PROF. W. M. HIRSCH INSTITUTE OF MATHEMATICAL SCIENCES NEW YORK UNIVERSITY NEW YORK NEW YORK 1C453 1 FRCF. D. L. IGLEHART LEPARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSITY STANFORD, CALIFORNIA 54350 1 FRCF. J. E. KACANE DEFARTMENT OF STATISTICS CAFNEGIE-MELLON FITTSEURGF. PENNSYLVANIA 15213 1 DR. RICHARD LAU LIFECTOR CFFICE CF NAVAL RESEARCH ERANCH OFF 1C30 EAST GREEN STREET PASADENA CA S1101 1 DF. A. R. LAUFER DIRECTCR CFFICE OF NAVAL RESEARCH BRANCH OFF 1030 EAST GREEN STREET PASACENA CA 91101 1 1 PROF. N. LEADBETTER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL NOFTH CAROLINA 27514 CR. J. S. LEE J. S. LEE ASSOCIATES, INC. 2001 JEFFERSCN DAVIS HIGHWAY SUITE 802 ARLINGTON VA 1 22202 FRCF. L. C. LEE DEPARTMENT CF STATISTICS VIRGINIA FCLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACKSBURG VA 24061 CONSTRUCT AND STATE UNIVERSITY •-

1

1

1

1

1

1

1

1

FRCF. P. A. W. LEW IS CEPT. CF CFERATIONS RESEARCH AND ADMINISTRATIVE SCIENCES NAVAL FCST GRACLATE SCHCOL MONTEREY , CALIFORNIA 93540

FRCF G. LIEBERMAN STANFORD INIVERSITY DEFARIMENT OF GPERATIONS RESEARCH STANFORD CALIFORNIA \$4305

DR. JAMES R. MAAR NATIONAL SECURITY AGENCY FORT MEADE , MARYLAND 20755

FRCF. R. W. MACSEN DEPARTMENT CF STATISTICS UNIVERSITY OF MISSCURI COLUMBIA MO

65201

YI ANALA TR

92093

96503

DR. N. R. MANN SCIENCE CENTER ROCKWELL INTERNATIONAL CORPORATION P.C. BOX 1085 THOUSAND C4KS CALIFORNIA S136C

CR. W. H. MARLCW PROGRAM IN LOCISTICS THE GEORGE WASHINGTON UNIVERSITY 707 22NC STREET , N. W. WASHINGTON , D. C. 20037

PROF. E. MASRY DEFT. APPLIEC PHYSICS AND INFORMATICA SERVICE UNIVERSITY OF CALIFORN IA LA JULLA CALIFORNIA

CF. BFLCE J. MCCONALD SCIENTIFIC DIRECTCR SCIENTIFIC LIAISON GROUP OFFICE CF NAVAL RESEARCH AMERICAN EMBASSY - TOKYC AFC SAN FFANCISCO

503 1 PROF. J. A. MUCK STADT DEFT. CF CPERATIONS RESEARCH GCFNELL UNIVERSITY ITFACA, New YCFK 19850

1

1

1

1

1

1

1

1

CR. JANET M. MYHRE THE INSTITUTE OF DECISION SCIENCE FOR BUSINESS AND PUBLIC POLICY CLAREMONT MEN'S COLLEGE CLAREMONT \$1711 MR. F. NISSELSCN BLREAU OF THE CENSUS ROCM 2025 FREERAL FUILCING 3 WASHINGTCN , D. C. 2033 A RUSSENALATI MISS B. S. CRLEANS NAVAL SEA SYSTEMS COMMAND (SEA 03F) FM 10SC3 ARLINGTON VIRGINIA 20360 FRCF. C. E OWEN DEPARTMENT CF STATISTICS SOUTHERN METHODIST UNIVERSITY CALLAS TEXAS 75222 PROF. E. PARZEN STATISTICAL SCIENCE DIVISION STATE UNIVERSITY OF NEW YORK AT BUFFALC AMFERST NEW YORK 14 14226 DR. A. PETRASOVITS RCCM 2078 , FCCC AND DRLG PLDG. TUNNEY'S PASTLRE CTTOWA , ENTARIE KIA-CL2 , CANADA FRCF. S. L. PHCENIX SIBLEY SCHOOL OF MECHAN ICAL AND AEROSPACE ENGINEERING CORNELL UNIVERSITY ITHACA NY DP. A. L. POWELL CIRECTCR CFFICE OF NAVAL RESEARCH BRANCH OFF 495 SUMMER STREET BESTEN 02210 MA MR. F. R. FRICFI CODE 224 CPERATIONSL TEST AND ONRS EVALUATION FORCE (OPTEVFOR) NORFOLK . VIRGINIA

20:00

1

1

1

1

1

1

1

1

1

PROF. M. L. PURI DEFT. CF MATHEMATICS P.G. BOX F INCIANA UNIVERSITY FOUNDATION PLCOMINGTON IN 474 47401

FROF, H RCEBINS DEFARTMENT OF MATHEMATICS COLUPEIA UNIVERSITY NEW YORK, NEW YORK, 10327

PFOF. M ROSENBLATT DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA SAN DIEGO LA JOLLA CALIFORNIA

92093

PROF. S. M. RCSS College of Engineering UNIVERSITY OF CALIFORNIA BERKELEY CA

94723

PROF. I RUBIN SCHOUL OF ENGINEERING AND AFPLIED SCIENCE UNIVERSITY OF CALIFORNIA LOS ANGELES CALIFORNIA 90024

FRCF. I. R. SAVAGE CEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN. CCNNECTICUT C6520

:

FRCF. L. L., SCHARF JR DEPARTMENT CF ELECRICAL ENGINEER ING COLORACO STATE UNIVERSITY FT. CCLLINS, CCLORACO E0521

PRCF. R. SERFLING CEPARTMENT CF STATISTICS FLCRIDA STATE UNIVERSITY

32306 TALLAHASSEE FLOR IDA

PROF. W. R. SCHLCANY DEFARTMENT OF STATISTICS SULTHERN METHICDIST UNIVERSITY CALLAS . TEXAS 75222

1

1

1

1

1

1

1

LOITATIAN TO AND A

. .

PROF. C. C. SIEGMUND CEPT. OF STATISTICS STANFORD LAIVEFSITY STANFORD CA

\$4305

SUDIO 2122 CHUR OF

FRCF. M. L. SHGOMAN DEPT. CF ELECTRICAL ENGINEERING POLYTECHNIC INSTITUTE CF NEW YORK BRCCKLYN, NEW YORK 11201

PROF. N. SINGPURWALLA LEPT. CF OPERATIONS RESEARCH THE GEORGE WASHINGTON UNIVERSITY 7C7 22ND ST . N. W. WASHINGTON, C. C. 20052

DR. A. L. SLAFKOSKY SCIENTIFIC ADVISOR COMMANCANT OF THE MARINE CORPS D. C. 20380

WR. CHARLES S. SMITH CASD (IGL), PENTAGEN WASHINGTON

20301

CR. C. E. SMITH DE SMATICS INC. P.C. BCX 618 STATE COLLEGE PENNSYLVANIA 16801

PROF. W. L. SMITH DEFARTMENT OF STATISTICS UNIVERSITY OF NORTH CARCLINA CHAPEL HILL NOFTH CARCLINA 27514

PRCF. H SCLOMCN CEPARTMENT CF STATISTICS STANFCRC UNIVERSITY STANFCRD ; CALIFORNIA \$4305

MF. GLENN F. STAFLY NATIONAL SECURITY AGENCY FORT MEACE MARYLAND 20755

•

1

1

1

1

1

1

1

1

1

04101114 27514

ADITATISTICS ADITATISTICS

A PARTY A PARTY A PERKEY

MR. EAVIE A. SWICK ADVANCED PROJECTS GROUP CODE 8123 NAVAL RESEARCH LAB. MASHINGTEN CC

20375

MR. WENDELL G. SYKES ARTHUR C. LITTLE, INC. ACERN PARK CAMBRIDGE MA

02140

PROF. J. R. THEMPSON DEPARTMENT OF MATHEMATICAL SCIENCE RICE UNIVERSITY HOUSTON, TEXAS 77001

PROF. W. A. THCMPSCN DEFARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMEIA, MISSOURI 65201

FREF. F. A. TILLMAN DEPT. CF INDUSTRIAL ENGINEERING KANSAS STATE UNIVERSITY MANHATTAN KS

66506

PRCF J. W. TUKEY CEFARIMENT OF STATISTICS FRINCETON UNIVERSITY

FRINCETON , N. J. 08540

PRCF. A . F . VEINOTT DEFARTMENT CF GPERATIONS RESEARCH STANFORD UNIVERSTITY STANFORD CALIFORNIA 94305

CANIEL H. WAGNER STATION SQLARE ONE FACLI , FENNSYLVANIA 19301

PRCF. GRACE WAHBA CEFT. CF STATISTICS UNIVERSITY CF WISCONSIN MADISON WI

FRCF. K. T. WALLENIUS DEFARTMENT OF MATHEMATICAL SCIENCES. CLEMSON UNIVERSITY CLEMSON, SOLTH CARCLINA 29631 1 PRCF. G. S. WATSCN DEFARTMENT OF STATISTICS FRINCETON , N. J. C8540 1 PRCF, BERNARD WIDRCH STANFORD ELECTRONICS LAB STANFORD UNIVERSITY STANFORD CA 1 94305 FRCF. G. E. WEITEFOUSE DEPT. OF INCLSTRIAL ENGINEERING LEFIGE UNIVERSITY BETHLEHEM 1 PA . 18015 FIDF. S. ZACKS DEPT. OF MATHEMATICS AND STATISTICS CASE WESTERN RESERVE UNIVERSITY CLEVELANC., CHIO 44106 1 PROF. M. ZIA-HASSAN CEPARTMENT OF INDUSTRIAL AND SYSTEMS ENGINEERING ILLINDIS INSTITUTE OF TECHNOLOGY CHICAGO IL 1 HEAD, MATH. SCI SECTION NAT. SCIENCE FOUNDATION 18CO G STREET, N.W. WASHINGTON, C.C. 1 20550 PROF. A. F. ANDRUS DEFT. CF CF NAVAL POSTGRADUATE SCHCCL MONTEREY CALIFORNIA 1 \$3940 FFOF. C. R. BARR DEFT. CF CP NAVAL POSTGRACLATE SCHOCL MONTEREY CALIFCENIA 1 93940 2104133

PRCF. PETER ELCCMFIELC STATISTICAL DEPT. PRINCETON UNIVERSITY PRINCETUN, N. J.

C8540

\$3940

15025 AATTONA

RAJ 20101. DELYARD

ANTH SCIENCE SCIECK

PRCF. G. G. BROWN EEFT. CF DR NAVAL POSTGRACUATE SCHCCL MONTEREY CALIFERNIA

R. W. EUTTERWORTH Systems Exploration WEESTER ST. MONTEREY CALIFORNIA

93940

9430%

DR. JAMES CAPRA 7218 CELFIELC STREET CHEVY CHASE MARYLANC

20015

ATTES CR. C.R. CCX DEPT. CF MATHEMATICS IMPERIAL COLLEGE LCNDCN SW7

E NG LA N

DEFENSE DOCUMENTATION CTR. CAMERON STATION ALEXANDRIA VIFGINIA

22314

ECCN. AND MAN. SCI. CTR. NORTHWESTERN UNIV. EVANSTON ILLINCIS

602 C1

L VAL CETTRADEATE SCHOOL MAN. SCE. RES. CTR. FACULTY OF COM. AND BUS. ADMIN. UNIV. CF ERITISH COLUMEIA VANCOUVER BRITISH COLUMEIA VOT 165 CAN CANADA LETT: CE C BARR LETT: CE C BARR LETT: CE CE BARR

CR. M. CWASS MAIN. DEFT. NORTHWESTERN UNIV. EVANSTEN ILLINDIS

602 01

.

93960

No. of Copies

2 72 12 14 ATA 10 14 ATA 11 A

1

1

1

. 1

1

1

1

1

1

CR. R. ELASHCFF BIOMATHEMATICS UNIV. CF CALIF. LCS ANGELES CALIFORNIA

90024

PROF. GECRGE S. FISHMAN UNIV. CF NORTH CAROLINA CUR. IN DR AND SYS. ANALYSIS PHILLIPS ANNEX CHAPEL HILL, NORTH CAROLINA 20742

DR. R. GNANACESIKAN EELL TELEPHENE LAB HOLMDEL, N. J.

07733

20234

DR. A. J. GCLEMAN CHIEF, OR DIV. 205.02, ADMIN. A428 U.S. DEPT. CF COMMERCE WASHINGTON, E.C.

DR. H. FIGGINS 53 BONN 1, POSTFACH 585 NASSESTRASSE 2

.

..

WEST GERMANY

DR. P. T. HOLMES DEPT. OF MATH. CLEMSCN LNIV. CLEMSON SCUTH CAROLINA

•

29631

CR. J. A. HOCKI BELL TELEPHONE LABS HOLMDEL NEW JERSEY

07733

DR. ROBERT HOCKE MATH. DEPT. WEST INCHOUSE RES. LABS CHURCHILL BEFE FITTSPURGE, PENNSYLVANIA

LR. D. L. ICLEFART CEPT. CF C.F. STANFCRD LNIV. STANFORD CALIFLENIA

94305

1 52 35

1

1

1

SELECT F. RCCIMMIS

es

. `

No. of Cap			No.	of Copic
CR. PATRICIA JACCBS CR DEPI. NAVAL POSTGRADLATE SCH MCNTEREY CALIFORNIA	ICCL 93		THEFASHU THEFASTIC NEEFSSIES NEEFFSSIES ORNIA	C
DR. H. KOEAYASHI IBM Yorktown feigfts New York	.YSIS 0174	S. FISHMAN F CARGLINA D SYS. ANAL X NCRTH C 898	GECRGE CF NCRI IR DR AN IFS ANNEL	PROF. LLIV. CUR. PHIL. CHARS
ER. JOHN LEHOCZKY STATISTICS DEPARTMENT CARNEGIE-MELLON UNIVER FITTSBURGH PENNSYLVANIA	20702 SITY	451123 5 L 48 *	GNANAC TELEPHCN EL, N. J	8 .30 1353 01164
L JERARY CCCE 55 NAVAL POSTGRADUATE SCH MONTEREY	ICGL	EMAN ADMIN, A428 ÇEMMERCE	. J. 60L 205.02.	DI CHIEF DI V:
CALIFERNIA DR. A. LEMOINE 1020 GUINEA ST. FALO ALTC. CALIFORNIA	2023 2023	340 340 Sirach 585	NULDA L	HADHI DA 53 80 NA 556
DR. J. MACCUSEN	уидчязэ ⁹⁴	301 23%	unit T	a h a
UNIV. CF CALIF. LOS ANGELES CALIFORNIA	. 90	024	OF MATH	CLENT, SCLEMSC SCUTH
FFCF. K. T. MARSHALL DETT. CF CF Naval Postgracuate SCH Monterey California	CCL 53	284J	ELERFON	е .
DR. M. MAZUMCAR MATH. DEPT. ESTINCHOLSE RZS. LABS CHURCHILL BCFC FITTSBLRGH PENNSYLVANJA	15	IKE ES. LABS MNSYLVAI CES	лерт, нас изетт. сноиза к тс всес укор, ре	CHURCH WATH WESTIN CHURCH FITTS
DR. LEON F. MCCINNIS SCHOLL CF INC. ANC SYS GEORGIA INST. OF TECH. ATLANTA CECRGIA	• EN G •	тяан 332	L. ICLE CF C.F. En CNIV.	1 1940 1947 1947 1947 1947 1947 1947 1947 1947

1

1

TENER OF HIGH TOPS

ARTAGATON UNIV. MDINGATON YASANG MAN

ASPTAR ... APTSON

1/15523 144.000 1

CR. D. R. MCNEIL DEPT. CF STATISTICS PRINCETON UNIV. FRINCETON NEW JERSEY

08540

PRCF. P. R. MILCH CEFT. OF OR NAVAL POSTGRAEUATE SCHECL MONTEREY CALIFORNIA 93940

CR. F. MOSTELLER STAT. CEPT. HARVARC UNIV. CAMBRIEGE MASSACHUSETTS

02139

FRCF. R. R. READ DEFT. CF CR NAVAL POSTGRAEUATE SCHOOL MONTEREY CALLFORNIA

93940

DR. M. REISER IEM THOMAS J. WATSEN RES. CTR. YERKTOWN HEIGHTS NEW YERK 10598

10598

DEAN CF RESEAFCH CODE 013 NAVAL FESTGRAELATE SCHOOL MENTEREY CALIFERNIA 93940 93940

FRCF. F. F. FICHARDS DEFT. GF OR NAVAL FCSTGFACLATE SCHOOL MONTERY CALIFORNIA

93940

DR. J. RICRCAN CEPT. OF MATHEMATICS PCCKEFELLER UNIV. NEW YORK NEW YORK

100 21

DR. LINUS SCHRAGE UNIV. CF CFICAGE CFAD. SCHOOL OF BLS. 5836 GREENWECE AVE. CHICAGE, ILLINDIS

1

1

1

1.140

1

N82 133 - 1 - 861 584

DR. PAUL SCHWEITZER THOMAS J. WATSON RESEARCH CTR9 POST OFFICE BOX 218 YORKTOWN FEIGHTS NEW YOFK

10598

CR. RICHARE SCRENSEN CODE 303 NPRDC 271 CATALINA BLVD. SAN DIEGE CALIFORNIA

No. 'bi Copicy

92152

PFCF. M. G. SOVEREIGN DEFT. CF OR NAVAL POSTGRACUATE SCHCCL MONTEREY CALIFERNIA

53 94 0

DR. V. SFINIVASAN GRADUATE SCHCCL CF BUSINESS STANFORD UNIVERSITY STANFORD CALIFORNIA

94305

DR. R. M. STARK STATISTICS AND COMPUTER SCI. UNIV. OF DELAWARE NEWARK NEWARK CELAWARE 19711

:

FFOF. RICHARE VANSLYKE RES. ANALYSIS CCRP. BEECHWCOD CLD TAFPEN FOAC GLEN CCVE, NEW YORK

11 542

CF_RESEAFCH ...

HRCAL F. F. FICHAROS DEF. GF GR LIVAL FOSTERACLETE SCHOOL

CRAPTIC SCHRAGE CREWE TF CLECKCO ERROW TF CLECKCO SCHRAGE OF ALS SCHRAGE TRANSIT SCHRAGE TRANSIT SCHRAGE TRANSIT

PRCF. JDFN N. TUKEY FINE FALL FRINCETON UNIV. PRINCETON NEW JERSEY

08540

CR. THEMAS C. VARLEY CFFICE OF NAVAL RESEARCH CODE 434 ARLINGTEN VA

22217

PRCF. G. WATSCN FINE HALL FRINCFTON UNIV. PRINCFTON NEW JERSEY

C8540

1

Dr. Roy Welsch M.I.T., Sloan School Cambridge, MA 02139

Dean of Research 012 Naval Postgraduate School Monterey, Ca. 93940

Professor D. P. Gaver Code 55Gv Naval Postgraduate School Monterey, Ca. 93940

Professor Peter Lewis Code 55Lw Naval Postgraduate School Monterey, Ca. 93940

•.

R. J. Stampfel Code 55 Naval Postgraduate School Monterey, Ca. 93940 1

50

50