

AD-A061 972

METEOROLOGY INTERNATIONAL INC MONTEREY CALIF

F/6 8/10

SOUND PROPAGATION IN THE SEA--RAY TRACING, CURVATURE CORRECTION--ETC(U)

N62271-67-M-1597

JUN 67 M M HOLL


NL

UNCLASSIFIED

|OF|  
AD  
A061972



END  
DATE  
FILMED  
2-79  
DDC



MICROCOPY RESOLUTION TEST CHART

METEOROLOGY INTERNATIONAL INCORPORATED

P. O. BOX 349 • MONTEREY, CALIFORNIA 93940



**LEVEL III**

①

AD A061972

⑥ Sound Propagation in the Sea -- Ray Tracing.  
CURVATURE CORRECTIONS FOR THE EARTH (SEA LEVEL)  
SHAPED AS AN ELLIPSOID OF REVOLUTION.

⑨ Project M-138: Technical Summary Report,  
June 1967

⑫ 22 p.

⑩ Prepared by: M. M. Holl  
Under Contract No.: N62271-67-M-1597

⑮

DDC FILE COPY

For the Officer-in-Charge  
Fleet Numerical Weather Facility  
U. S. Naval Postgraduate School  
Monterey, California

ACCESSION FOR	
DTIC	White Section <input checked="" type="checkbox"/>
DDC	Diff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
Per Basic rpt.	
BY AD-731734	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

DDC  
RECEIVED  
DEC 11 1978

⑮

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

227 450

78 12 11 058 LB

Sound Propagation in the Sea -- Ray Tracing  
CURVATURE CORRECTIONS FOR THE EARTH (SEA LEVEL) SHAPED AS AN  
ELLIPSOID OF REVOLUTION

by: M. M. Holl


ABSTRACT

1. The Problem

The three-dimensional tracing of a sound ray in the sea may be referred to the surface coordinate system of latitude and longitude and to the depth below the sea surface. This coordinate system involves curvature terms.

~~For~~ the earth shaped as an ellipsoid of revolution ~~we~~ may refer to two definitions of latitude: the geocentric latitude  $\phi$  and the astronomical latitude  $\phi'$ . These are illustrated in Fig. 2. There is a one-to-one relationship between these definitions and the transformation is simple as will be shown. ~~We choose to use~~ the geocentric latitude  $\phi$  as principal latitude coordinate.

Let the direction of propagation of a sound ray at an arbitrary point in its progress -- latitude  $\phi$ , longitude  $\theta$  and depth  $z$  -- be defined by the unit vector  $\mathbf{T}$ . It makes the angle  $\gamma$  with the horizontal as shown in Fig. 1. The horizontal component of the unit vector  $\mathbf{T}$  defines the direction of the unit vector  $\mathbf{t}$ .



RECEIVED  
DEC 11 1958

APPROVED FOR PUBLICATION  
BY THE DIRECTOR OF THE NATIONAL BUREAU OF STANDARDS




Fig. 1 Definition of Parameters in the Vertical Plane of the Ray

In Fig. 2 we define and illustrate geometric parameters which are relevant to the problem. These include the angle  $\beta$  which  $\mathbf{t}$  makes with the unit vector  $\mathbf{i}$ :  $\mathbf{t}$  is directed  $\beta$  radians north of due east.


The coordinate-curvature terms arise in prescribing the ray direction  $\mathbf{T}$  by the angles  $\gamma$  and  $\beta$ . The angle  $\gamma$  is referred to the horizontal which itself is turning in space from location to location. The angle  $\beta$  is referred to the east which itself is turning from location to location.

We may formulate the curvature terms directly. The reference directions turn only by horizontal progress and the angles are defined in specific planes. The terms are

$$(\partial\gamma/\partial s)_c = + \mathbf{t} \cdot \nabla \mathbf{t} \cdot \mathbf{k} \quad (1)$$

$$(\partial\beta/\partial s)_c = - \mathbf{t} \cdot \nabla \mathbf{i} \cdot \mathbf{j} \quad (2)$$

Fig. 2 Definition of Parameters for the Surface



$\phi$  : geocentric latitude

$\Phi$  : astronomical latitude

$$\alpha = \Phi - \phi$$

$k$  : unit vector, surface normal, aligned with local gravity, subtends angle  $\beta$  with equatorial plane as shown.

$i$  : unit vector, surface tangent, directed eastward.

$j$  : unit vector, surface tangent, directed northward.

$$i \times j = k$$

$A$  : equatorial radius of earth  $A > B$   
 $B$  : polar radius of earth

$E$  : radius at geocentric latitude  $\phi$  (astronomical latitude  $\Phi$ )

$m, h$  : Cartesian coordinates in arbitrary meridional plane of longitude  $\theta$ .  
 Solid lines lie in the meridional plane; dashed lines do not.

$t$  : unit vector, tangent to surface trace.

$m$  : unit vector, surface tangent, normal to  $t$ .

$$k \times t = m$$

$$t = \cos \beta i + \sin \beta j$$

$$m = -\sin \beta i + \cos \beta j$$

## 2. Geometry of the Surface

The intersect of the arbitrary meridional plane and the surface is an ellipse:

$$\frac{h^2}{B^2} + \frac{m^2}{A^2} = 1 \quad (3)$$

We substitute

$$\begin{aligned} h &= E \sin \phi \\ m &= E \cos \phi \end{aligned} \quad (4)$$

and obtain

$$E^2 = \frac{A^2 B^2}{A^2 \sin^2 \phi + B^2 \cos^2 \phi} \quad (5)$$

We also note that

$$\tan \phi = \frac{h}{m} \quad (6)$$

An incremental northward displacement on the ellipse may be expressed by

$$h \frac{\delta h}{B^2} + m \frac{\delta m}{A^2} = 0 \quad (7)$$

where  $\delta h$  is positive and  $\delta m$  is negative. This displacement is directed along  $j$ . Hence

$$\begin{aligned} \tan \bar{\phi} &= - \frac{\delta m}{\delta h} = \frac{A^2}{B^2} \frac{h}{m} \\ &= \frac{A^2}{B^2} \tan \phi \end{aligned} \quad (8)$$

This is the relationship between the astronomical latitude  $\bar{\phi}$  and the geocentric latitude  $\phi$ . Their difference angle

$$\alpha = \bar{\phi} - \phi \quad (9)$$

is given by

$$\begin{aligned} \tan \alpha &= \tan (\bar{\phi} - \phi) = \frac{\tan \bar{\phi} - \tan \phi}{1 + \tan \bar{\phi} \tan \phi} \\ &= \frac{(A^2 - B^2) \tan \phi}{B^2 + A^2 \tan^2 \phi} \end{aligned} \quad (10)$$

We also require the ratio of latitude increments: Differentiation of Eq. (8) yields

$$\sec^2 \Phi \delta \Phi = \frac{A^2}{B^2} \sec^2 \phi \delta \phi$$

$$\frac{\delta \Phi}{\delta \phi} = \frac{A^2}{B^2} \frac{\cos^2 \Phi}{\cos^2 \phi}$$

$$= \frac{A^2 B^2}{\cos^2 \phi (B^4 + A^4 \tan^2 \phi)} \quad (11)$$

### 3. Curvatures of the Coordinate Unit Vectors

We are now in position to express and develop the curvatures of the coordinate unit vectors. These are expressed by an incremental directed turning of the unit vector divided by the space increment, in the pertinent direction, over which this turning takes place:

$$i \cdot \nabla i = \frac{\delta \theta}{E \cos \phi \delta \theta} (j \sin \Phi - k \cos \Phi) \quad (12)$$

$$i \cdot \nabla j = - \frac{i \sin \Phi \delta \theta}{E \cos \phi \delta \theta} \quad (13)$$

$$i \cdot \nabla k = \frac{i \cos \Phi \delta \theta}{E \cos \phi \delta \theta} \quad (14)$$

$$j \cdot \nabla i = 0 \quad (15)$$

$$j \cdot \nabla j = - \frac{k \delta \Phi}{E \sec \alpha \delta \phi} \quad (16)$$

$$\mathbf{j} \cdot \nabla \mathbf{k} = \frac{\mathbf{j} \delta \Phi}{E \sec \alpha \delta \phi} \quad (17)$$

$$\mathbf{k} \cdot \nabla \mathbf{i} = \mathbf{k} \cdot \nabla \mathbf{j} = \mathbf{k} \cdot \nabla \mathbf{k} = 0 \quad (18)$$

In Eqs. (16) and (17) we substitute from Eq. (11):

$$\mathbf{j} \cdot \nabla \mathbf{j} = - \frac{\mathbf{k} \cos \alpha A^2 B^2}{E \cos^2 \phi (B^4 + A^4 \tan^2 \phi)} \quad (19)$$

$$\mathbf{j} \cdot \nabla \mathbf{k} = \frac{\mathbf{j} \cos \alpha A^2 B^2}{E \cos^2 \phi (B^4 + A^4 \tan^2 \phi)} \quad (20)$$

#### 4. Development of the Curvature Terms

Equation (1) is developed as follows

$$\begin{aligned} (\partial \delta / \partial s)_c &= \mathbf{t} \cdot \nabla \mathbf{t} \cdot \mathbf{k} \\ &= (\cos \beta \mathbf{i} + \sin \beta \mathbf{j}) \cdot \nabla (\cos \beta \mathbf{i} + \sin \beta \mathbf{j}) \cdot \mathbf{k} \\ &= \cos^2 \beta \mathbf{i} \cdot \nabla \mathbf{i} \cdot \mathbf{k} + \cos \beta \sin \beta \mathbf{i} \cdot \nabla \mathbf{j} \cdot \mathbf{k} \\ &\quad + \sin \beta \cos \beta \mathbf{j} \cdot \nabla \mathbf{i} \cdot \mathbf{k} + \sin^2 \beta \mathbf{j} \cdot \nabla \mathbf{j} \cdot \mathbf{k} \\ &= - \frac{\cos^2 \beta \cos \Phi}{E \cos \phi} - \frac{\sin^2 \beta \cos \alpha A^2 B^2}{E \cos^2 \phi (B^4 + A^4 \tan^2 \phi)} \quad (21) \end{aligned}$$

Equation (2) is developed as follows

$$\begin{aligned}
 (\partial\beta / \partial s)_c &= - \mathbf{t} \cdot \nabla \mathbf{i} \cdot \mathbf{j} \\
 &= - (\cos \beta \mathbf{i} + \sin \beta \mathbf{j}) \cdot \nabla \mathbf{i} \cdot \mathbf{j} \\
 &= - \cos \beta \mathbf{i} \cdot \nabla \mathbf{i} \cdot \mathbf{j} - \sin \beta \mathbf{j} \cdot \nabla \mathbf{i} \cdot \mathbf{j} \\
 &= - \frac{\cos \beta \sin \Phi}{E \cos \phi} \tag{22}
 \end{aligned}$$

We now wish to eliminate  $E$ ,  $\Phi$  and  $\alpha$  by substitution in terms of  $\phi$  according to Eqs. (5), (8) and (10). For abbreviation, where convenient, we adapt

$$K \equiv \tan \phi ; \quad F \equiv \frac{A^2}{B^2} \tag{23}$$

We may rewrite Eqs. (5), (8) and (10) as follows:

$$E^2 = \frac{A^2 (1 + K^2)}{(1 + FK^2)} \tag{24}$$

$$\tan \Phi = FK \tag{25}$$

$$\tan \alpha = \frac{(F - 1) K}{(1 + FK^2)} \tag{26}$$

It follows that

$$\cos \bar{\Phi} = \frac{1}{(1 + F^2 K^2)^{1/2}} \quad (27)$$

$$\sin \bar{\Phi} = \frac{FK}{(1 + F^2 K^2)^{1/2}} \quad (28)$$

$$\cos \alpha = \frac{(1 + FK^2) \cos \phi}{(1 + F^2 K^2)^{1/2}} \quad (29)$$

With the appropriate substitutions, Eqs. (21) and (22) become

$$(\partial \delta / \partial s)_C = - \frac{1}{A} \left\{ \frac{1 + FK^2}{1 + F^2 K^2} \right\}^{1/2} \frac{\cos^2 \beta + F \sin^2 \beta + F^2 K^2}{1 + F^2 K^2} \quad (30)$$

$$(\partial \beta / \partial s)_C = - \frac{\cos \beta}{A} FK \left\{ \frac{1 + FK^2}{1 + F^2 K^2} \right\}^{1/2} \quad (31)$$

These are the coordinate-curvature terms, on  $\delta$  and  $\beta$ , as functions of the geocentric latitude  $\phi$  and the ray-direction angle  $\beta$ .

In the special limit of a sphere:

$$A = B \equiv R; \quad \phi = \bar{\Phi}; \quad F = 1$$

the curvature terms simplify to

$$(\partial \gamma / \partial s)_{C,R} = - \frac{1}{R} \quad (32)$$

$$(\partial \beta / \partial s)_{C,R} = - \frac{\cos \beta \tan \phi}{R} \quad (33)$$

### 5. The Ray Tracing Equations

The ray tracing equation, in vector form, may be expressed by

$$\mathbf{T} \cdot \nabla \mathbf{T} = \mathbf{T} \times (\mathbf{T} \times \nabla \ln C) \quad (34)$$

where  $\mathbf{T}$  is the unit-vector ray direction and  $C$  is the sound speed. We take the components along  $\mathbf{N}$  and  $\mathbf{n}$  -- unit vectors defined in Figs. (1) and (2) respectively -- and obtain

$$\mathbf{T} \cdot \nabla \mathbf{T} \cdot \mathbf{N} = - \mathbf{N} \cdot \nabla \ln C \quad (35)$$

$$\mathbf{T} \cdot \nabla \mathbf{T} \cdot \mathbf{n} = - \mathbf{n} \cdot \nabla \ln C \quad (36)$$

The left-hand sides may be transformed by geometrical consideration:

$$\mathbf{T} \cdot \nabla \mathbf{T} \cdot \mathbf{N} = \partial \gamma / \partial S - \cos \gamma \mathbf{t} \cdot \nabla \mathbf{t} \cdot \mathbf{k} \quad (37)$$

$$\mathbf{T} \cdot \nabla \mathbf{T} \cdot \mathbf{n} = \cos \gamma \left\{ \partial \beta / \partial S + \cos \gamma \mathbf{t} \cdot \nabla \mathbf{i} \cdot \mathbf{j} \right\} \quad (38)$$

where  $S$  is a linear measure along the ray trace:

$$\delta s = \delta S \cos \gamma \quad (39)$$

The coordinate-curvature terms are expressed by Eqs. (1) and (2) and the ray-tracing equations may be written

$$\frac{\partial \gamma}{\partial s} = \left( \frac{\partial \gamma}{\partial s} \right)_c - \sec \gamma \mathbf{N} \cdot \nabla \ln C \quad (40)$$

$$\frac{\partial \beta}{\partial s} = \left( \frac{\partial \beta}{\partial s} \right)_c - \sec^2 \gamma \mathbf{n} \cdot \nabla \ln C \quad (41)$$

The curvature terms are given by Eqs. (30) and (31).

#### 6. Linearization as to Oblateness of Earth

If we define  $\epsilon$  by

$$A \equiv (1 + \epsilon) B \quad (42)$$

where  $A$  is the equatorial radius and  $B$  is the polar radius of the earth, then

$$0 < \epsilon \ll 1, \quad (43)$$

and we may linearize Eqs. (30) and (31) by dismissing squares and higher powers of  $\epsilon$  as negligible. In this linearization

$$A^2 = (1 + 2\epsilon) B^2 \quad (44)$$

$$F = (1 + 2\epsilon) \quad (45)$$

$$F^2 = (1 + 4\epsilon) \quad (46)$$

We also define as the earth's mean radius

$$R \equiv \frac{1}{2} (A + B) \quad (47)$$

It follows that

$$A = (1 + \frac{1}{2}\epsilon) R \quad (48)$$

With these substitutions we linearize Eqs. (30) and (31) and obtain

$$\left(\frac{\partial \gamma}{\partial s}\right)_C = -\frac{1}{R} \left\{ 1 + \epsilon (2 \sin^2 \beta \cos^2 \phi - \sin^2 \phi - \frac{1}{2}) \right\} \quad (49)$$

$$\left(\frac{\partial \beta}{\partial s}\right)_C = -\frac{\cos \beta \tan \phi}{R} \left\{ 1 + \epsilon \left(\frac{1}{2} + \cos^2 \phi\right) \right\} \quad (50)$$

We note that as  $\epsilon \rightarrow 0$  we arrive at Eqs. (32) and (33) respectively, for the sphere; this, however, is not a check on the linear terms.

It is convenient, at this point, to note that the effect of the depth of the ray on the curvature terms can be taken into account by replacing  $R$  by  $R - z$ . This is equivalent to replacing  $A$  and  $B$  by  $A - z$  and  $B - z$  in all earlier expressions of the curvature terms; the effect on  $\epsilon$  is negligible. In general the depth correction is smaller than the oblateness correction.

For convenience we also exhibit Eqs. (49) and (50) in terms of the astronomical latitude,  $\bar{\Phi}$  :

$$\left(\frac{\partial \gamma}{\partial s}\right)_c = -\frac{1}{R} \left\{ 1 + \epsilon (2 \sin^2 \beta \cos^2 \bar{\Phi} - \sin^2 \bar{\Phi} - \frac{1}{2}) \right\} \quad (51)$$

$$\left(\frac{\partial \beta}{\partial s}\right)_c = -\frac{\cos \beta \tan \bar{\Phi}}{R} \left\{ 1 - \epsilon \left(\frac{1}{2} + \sin^2 \bar{\Phi}\right) \right\} \quad (52)$$

By linearization of Eq. (26) we find that the difference between the astronomical latitude  $\bar{\Phi}$  and the geocentric latitude  $\phi$  is

$$\alpha = \bar{\Phi} - \phi = \epsilon \sin 2\phi$$

We note that

$$\alpha = \sin \alpha = \tan \alpha = \epsilon \sin 2\phi = \epsilon \sin 2\bar{\Phi} \quad (53)$$

$$\cos \alpha = \sec \alpha = 1 \quad (54)$$

The angle  $\alpha$  is a maximum at about  $\phi = \pm 45^\circ$ .

#### 7. Magnitude of Corrections for Oblateness

The oblateness factor does not enter in sea-surface or sea-bottom reflections. These reflections are treated in the same way as in the spherical case; no oblateness correction enters; this is because the angle  $\gamma$  is defined relative to the local vertical (see Fig. 1).

The values

$$A = 6.378 \times 10^6 \text{ meters}$$

$$B = 6.357 \times 10^6 \text{ meters} \quad (55)$$

have been quoted; perhaps better values are now available. These values give

$$\epsilon = \frac{A}{B} - 1 \approx 0.0033 \approx \frac{1}{300} \quad (56)$$

The oblateness corrections have the following ranges in the curvature terms:

$$\left(\frac{\partial \gamma}{\partial s}\right)_c = -\frac{1}{R} \left\{ 1 + (-0.005 \text{ to } +0.005) \right\} \quad (57)$$

for both, Eqs. (49) and (51).

$$\left(\frac{\partial \beta}{\partial s}\right)_c = -\frac{\cos \beta \tan \phi}{R} \left\{ 1 + (0.0017 \text{ to } 0.005) \right\} \quad (58)$$

$$\left(\frac{\partial \beta}{\partial s}\right)_c = -\frac{\cos \beta \tan \phi}{R} \left\{ 1 + (-0.005 \text{ to } -0.0017) \right\} \quad (59)$$

for Eqs. (50) and (52), respectively.

In moving northward at the equator, we have

$$\left(\partial \gamma / \partial s\right)_c = -\frac{1}{R} (1 + 0.005) \quad (60)$$

In a northward displacement of

$$\delta s = 0.1 R \approx 637,000 \text{ meters} \quad (61)$$

the oblateness correction amounts to 0.0005 radians. Spreading this correction linearly over the displacement results in a depth correction of

$$637,000 \times 0.00025 \approx 160 \text{ meters} \quad (62)$$

This is an upper-limit approximation to the displacement errors made in neglecting the earth's oblateness.

We also note from Eq. (53) that maximum  $\alpha$  (at  $\phi = \pm 45^\circ$ ) is

$$0.0033 \text{ radians} \approx 11 \text{ minutes} \quad (63)$$

#### 8. The Geodesic: Surface Ray with no Refraction

Consider a ray which is trapped in the sea surface by continuous glancing reflection, but undergoing no refraction. Its trace describes a geodesic:

$$\mathbf{t} \cdot \nabla \mathbf{t} = \mathbf{t} \cdot \nabla \mathbf{t} \cdot \mathbf{kk} \quad (64)$$

Its geodesic curvature (i.e. curvature in the surface) is zero:

$$\mathbf{t} \cdot \nabla \mathbf{t} \cdot \mathbf{n} = 0 \quad (65)$$

We have seen that Eq. (65) transforms into

$$\partial \beta / \partial s = (\partial \beta / \partial s)_c \quad (66)$$

We choose Eq. (22) to write

$$\frac{\delta \beta}{\delta s} = - \frac{\cos \beta \sin \Phi}{E \cos \phi} \quad (67)$$

for the governing equation for a geodesic.

The governing equation can be integrated once and obtains that, for a geodesic,  $\beta$  is a function of latitude only. We achieve this by substituting

$$\sin \beta \delta s = E \sec \alpha \delta \phi \quad (68)$$

as seen from Fig. 2, into Eq. (67) to eliminate  $\delta s$ . We obtain

$$\tan \beta \delta \beta = - \frac{\sin \Phi}{\cos \phi \cos \alpha} \delta \phi \quad (69)$$

We transform Eq. (28) to obtain

$$\sin \Phi = \sin \phi \frac{F}{\left\{ 1 + (F^2 - 1) \sin^2 \phi \right\}^{1/2}} \quad (70)$$

We transform Eq. (29) to obtain

$$\cos \alpha = \frac{1 + (F - 1) \sin^2 \phi}{\left\{ 1 + (F^2 - 1) \sin^2 \phi \right\}^{1/2}} \quad (71)$$

Introduction of Eqs. (70) and (71) in (69) yields

$$\tan \beta \delta \beta = - \tan \phi \frac{F}{1 + (F - 1) \sin^2 \phi} \delta \phi \quad (72)$$

We integrate Eq. (72) from the equator,  $\phi = 0$ , where  $\beta = \beta_0$ , to an arbitrary latitude  $\phi$ . The result is

$$\cos \beta = \cos \beta_0 (1 + F \tan^2 \phi)^{1/2} \quad (73)$$

The maximum latitude which this ray can attain is reached when  $\beta = 0$ :

$$\tan^2 \phi_m = \frac{\tan^2 \beta_0}{F} \quad (74)$$

The analytical equation for the geodesic requires one more integration. We first transform Eq. (73) to obtain

$$\tan \beta = \left\{ \frac{\sec^2 \beta_0}{1 + F \tan^2 \phi} - 1 \right\}^{1/2} \quad (75)$$

From the geometry of Fig. 2 we obtain

$$\tan \beta = \frac{R \sec \alpha \delta \phi}{R \cos \phi \delta \theta} \quad (76)$$

It follows that, for the geodesic,

$$\frac{d\phi}{d\theta} = \cos \phi \cos \alpha \left\{ \frac{\sec^2 \beta_0}{1 + F \tan^2 \phi} - 1 \right\}^{1/2} \quad (77)$$

Equation (77) may also be expressed by

$$\int_{\theta_0}^{\theta} d\theta = \int_0^{\phi} \frac{\sec \phi \sec \alpha (1 + F \tan^2 \phi)^{1/2}}{(\tan^2 \beta_0 - F \tan^2 \phi)^{1/2}} d\phi \quad (78)$$

The geodesic trace is obtained by  $\phi$  cycling back and forth between  $\phi_m$  and  $-\phi_m$ , the range given by Eq. (74).

In the linearization of Eq. (78), as to oblateness, we must concern ourselves with the range of  $\phi$ . We may immediately set  $\sec \alpha$  equal to one; its departure is second order in  $\epsilon$ . Again using  $K \equiv \tan \phi$  we transform Eq. (78) into

$$\theta - \theta_0 = \int_0^K \frac{(1 + FK^2)^{1/2} dK}{(1 + K^2)^{1/2} (\tan^2 \beta_0 - FK^2)^{1/2}} \quad (79)$$

We change the variable of integration once more to

$$P \equiv F^{1/2} K \quad (80)$$

and obtain

$$\theta - \theta_0 = \int_0^P \frac{F^{-1/2} (1 + P^2)^{1/2} dP}{(1 + F^{-1} P^2)^{1/2} (\tan^2 \beta_0 - P^2)^{1/2}} \quad (81)$$

Linearization, following  $F = 1 + 2\epsilon$ , yields

$$\theta - \theta_0 = \int_0^P \frac{dP}{(\tan^2 \beta_0 - P^2)^{1/2}} - \epsilon \int_0^P \frac{dP}{(1 + P^2)(\tan^2 \beta_0 - P^2)^{1/2}} \quad (82)$$

where the range of  $P$  is  $\pm \tan \beta_0$ .

We introduce  $\omega$  by

$$\sin \omega = P \cot \beta_0$$

$$\tan \omega = P (\tan^2 \beta_0 - P^2)^{-1/2} \quad (83)$$

Equation (82) transforms into

$$\theta - \theta_0 = \int_0^{\arcsin(P \cot \beta_0)} d\omega - \epsilon \int_0^{\arcsin(P \cot \beta_0)} \frac{d\omega}{1 + \tan^2 \beta_0 \sin^2 \omega} \quad (84)$$

The first integral is solved; the second is transformed once more:

$$\theta - \theta_0 = \arcsin(P \cot \beta_0) - \epsilon \int_0^{\left\{ \tan^2 \beta_0 - P^2 \right\}^{-1/2}} \frac{dy}{1 + \sec^2 \beta_0 y^2} \quad (85)$$

where  $y$  replaces the dummy variable,  $\tan \omega$ .

The linearized solution for the geodesic is

$$\theta - \theta_0 = \arcsin (P \cot \beta_0) - \epsilon \cos \beta_0 \arctan \left\{ \sec \beta_0 P (\tan^2 \beta_0 - P^2)^{-1/2} \right\} \quad (86)$$

where

$$P = (1 + \epsilon) \tan \phi \quad (87)$$

has the range  $\pm \tan \beta_0$ . The longitude at the maximum latitude is

$$\theta_m - \theta_0 = \pi/2 (1 - \epsilon \cos \beta_0) \quad (88)$$

For  $\epsilon = 0$  the surface becomes a sphere, and the geodesic is a great circle: Equation (86) reduces to

$$\sin (\theta - \theta_0) = \tan \phi \cot \beta_0 \quad (89)$$

This result could have been obtained directly from the spherical trigonometry of a right spherical triangle.

In Fig. 3 we show the extent to which orbits of the geodesic precess. For

$$\epsilon = \frac{1}{300} \quad \text{and} \quad \beta_0 = 45 \text{ degrees} \quad (90)$$




Fig. 3 Schematic of the Geodesic

the latitude maximum attained in Fig. 3 is located at

$$\theta = 89.7_9 \text{ degrees}$$

$$\phi = 44.9_0 \text{ degrees}$$

$$\bar{\phi} = 45.1_0 \text{ degrees}$$

(91)

The oblateness correction appears to be negligible within the practical limitations in sound-ray tracing in the sea.