

DDC FILE COPY. AD A063219

**LEVEL**

2

NPS55-78-029

**NAVAL POSTGRADUATE SCHOOL**  
Monterey, California



DDC  
JAN 11 1979  
F

SEARCH FOR A MOVING TARGET:  
UPPER BOUND ON DETECTION PROBABILITY

by

Alan R. Washburn

October 1978

Approved for public release; distribution unlimited.

79 01 09 030


NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CALIFORNIA

Rear Admiral T. F. Dedman  
Superintendent

J. R. Borsting  
Provost


Reproduction of all or part of this report is authorized.


This report was prepared by:

  
Alan R. Washburn, Associate Professor  
Department of Operations Research

Reviewed by:

Released by:

  
Michael G. Sovereign, Chairman  
Department of Operations Research

  
William M. Tolles  
Dean of Research

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
14. REPORT NUMBER NPS55-78-029	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
6. TITLE (and Subtitle) Search for a Moving Target: Upper Bound on Detection Probability.	9. TYPE OF REPORT & PERIOD COVERED Technical rept.	5. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Alan R. Washburn	8. CONTRACT OR GRANT NUMBER(s)	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, Ca. 93940	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 12/17a	
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, Ca. 93940	13. REPORT DATE 11/ Oct 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) SVP	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. PJR		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Search Moving Target		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Existing algorithms for computing the optimal distribution of effort in search for a moving target operate by producing a sequence of progressively better distributions. This report shows how to compute an upper bound on the detection probability for each of those effort distributions in the case where the detection function is concave. A		

DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE  
1 JAN 73 5/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

70 01 09 030 251 450 Liu

SEARCH FOR A MOVING TARGET:  
UPPER BOUND ON DETECTION PROBABILITY

by

Alan R. Washburn  
Naval Postgraduate School  
Monterey, CA 93940

ABSTRACT

Existing algorithms for computing the optimal distribution of effort in search for a moving target operate by producing a sequence of progressively better distributions. This report shows how to compute an upper bound on the detection probability for each of those effort distributions in the case where the detection function is concave.

SEARCHED	INDEXED
SERIALIZED	FILED
APR 1980	
FBI - MONTEREY	
BY	AG
DATE	APR 1980
RECEIVED	AG
SPECIAL	

## TABLE OF CONTENTS

	PAGE
1. INTRODUCTION .....	3
2. THE GENERAL CASE .....	3
3. THE CASE OF MARKOV MOTION AND EXPONENTIAL DETECTION FUNCTION .....	6
4. AN EXAMPLE .....	9
REFERENCES .....	11

## 1. INTRODUCTION

The object is to detect a randomly moving target at one of the discrete times  $0, 1, \dots, \tau$ . The searcher determines a non-negative effort distribution  $\psi(x, t)$  such that the total effort applied at time  $t$  does not exceed  $m(t)$ . Our purpose here is to establish an upper bound on the detection probability for every effort distribution. This is intended to supplement existing iterative procedures that develop sequences of effort distributions that improve monotonically.

## 2. THE GENERAL CASE

Let  $w(x, t)$  be an "effectiveness coefficient" for search effort applied at position  $x$  at time  $t$ . If  $X_t$  is the position of the target at time  $t$ , then the probability of detection depends, we assume, only on the total effective search effort

$$Z = \sum_{t=0}^{\tau} w(X_t, t) \psi(X_t, t) .$$

Specifically the probability of detection is  $P(\psi) = E(b(Z))$ , where the expectation operator is needed because  $X_t$  is a stochastic process. We assume that  $b(z') - b(z) \leq s(z)(z' - z)$  for some function  $s(z)$  and for all  $z'$ ; if the detection

function  $b$  is concave and differentiable, then  $s(z)$  is just  $\frac{d}{dz} b(z)$ . In most applications,  $b(z) = 1 - \exp(-z)$  and  $s(z) = \exp(-z)$ .

Let  $\psi$  and  $\psi'$  be two effort distributions. Then

$$(1) \quad P(\psi') - P(\psi) = E(b(Z') - b(Z)) \leq E(s(Z) (Z' - Z))$$

We also have

$$(2) \quad E(s(Z) (Z' - Z)) = \sum_{t=0}^T E(s(Z) w(X_t, t) [\psi'(X_t, t) - \psi(X_t, t)])$$

We now consider two cases, depending on whether  $X_t$  is discrete or continuous. If  $X_t$  is discrete, we let  $p_t(x)$  be the probability mass function of  $X_t$ , and require

$$\sum_x \psi'(x, t) = \sum_x \psi(x, t) = m(t) .$$

We define

$$(3) \quad D_\tau(\psi, x, t) = w(x, t) p_t(x) E(s(Z) | X_t = x)$$

Then (2) can be written

$$(4) \quad E(s(Z) (Z' - Z)) = \sum_{t=0}^T \sum_x D_\tau(\psi, x, t) [\psi'(x, t) - \psi(x, t)]$$

If  $X_t$  is continuous, let  $p_t(x)$  be the probability density function of  $X_t$ , and require  $\int \psi'(x,t) dx = \int \psi(x,t) dx = m(t)$ . Then (2) can be written

$$(4') \quad E(s(Z)(Z'-Z)) = \sum_{t=0}^{\tau} \int D_{\tau}(\psi, x, t) [\psi'(x, t) - \psi(x, t)] dx .$$

In either case, suppose that  $D_{\tau}(\psi, x, t) \leq \bar{\lambda}(t)$  for all  $x$ , and  $D_{\tau}(\psi, x, t) \geq \underline{\lambda}(t)$  when  $\psi(x, t) > 0$ . In the discrete case, from (4),

$$(5) \quad E(s(Z)(Z'-Z)) \leq \sum_{t=0}^{\tau} \left[ \sum_x \bar{\lambda}(t) \psi'(x, t) - \sum_x \underline{\lambda}(t) \psi(x, t) \right] \\ = \sum_{t=0}^{\tau} (\bar{\lambda}(t) - \underline{\lambda}(t)) m(t)$$

Combining (5) and (1),

$$(6) \quad P(\psi') - P(\psi) \leq \sum_{t=0}^{\tau} (\bar{\lambda}(t) - \underline{\lambda}(t)) m(t) \equiv \Delta(\psi)$$

A formula similar to (5) shows that (6) must also hold in the continuous case.

Now  $\psi'$  is not needed to compute any of the quantities on the right-hand side of (6), so every effort density  $\psi'$  has associated with it an upper bound on the detection probability  $P(\psi) + \Delta(\psi)$ . In the event  $\bar{\lambda}(t) = \underline{\lambda}(t)$  for all  $t$ ,  $\psi$  must actually be optimal--this has been observed by Stone [1], and in fact our whole development is a modification of his



sufficiency proof. The main issue is now computational: is determination of  $\Delta(\psi)$  worth the effort?

### 3. THE CASE OF MARKOV MOTION AND EXPONENTIAL DETECTION FUNCTION.

Let

$$z = z_t^- + w(x_t, t) \psi(x_t, t) + z_t^+ \quad \text{for } 0 \leq t \leq \tau,$$

where

$$z_t^- = \sum_{u=0}^{t-1} w(x_u, u) \psi(x_u, u),$$

$$z_t^+ = \sum_{u=t+1}^{\tau} w(x_u, u) \psi(x_u, u),$$

and

$$z_0^- = z_\tau^+ = 0.$$

Then

$$D_\tau(\psi, x, t) = w(x, t) p_t(x) E(s(z_t^- + w(x, t) \psi(x, t) + z_t^+) | X_t = x).$$

If the motion is Markov, then  $z_t^+$  is independent of  $z_t^-$  when  $X_t = x$  is given. If  $b(z) = 1 - \exp(-z)$ , then  $s(z) = \exp(-z)$ . If both conditions hold, then

$$D_\tau(\psi, x, t) = w(x, t) P(\psi, x, t) \exp(-w(x, t) \psi(x, t)) Q(\psi, x, t),$$

where

$$P(\psi, x, t) \equiv p_t(x) E(s(z_t^-) | X_t = x)$$

and

$$Q(\psi, x, t) \equiv E(s(z_t^+) | X_t = x).$$

$P(\psi, x, t)$  is the joint probability that  $X_t = x$  and that the target is not detected by any of the searches at  $0, 1, \dots, t-1$ ; note that  $P(\psi, x, t)$  does not depend on  $\psi(y, u)$  for  $u \geq t$ .  $Q(\psi, x, t)$  is the conditional probability that the target is not detected by any of the searches at  $t+1, \dots, \tau$  given that  $X_t = x$ ; note that  $Q(\psi, x, t)$  does not depend on  $\psi(y, u)$  for  $u \leq t$ . Given  $\psi$ ,  $P(\psi, x, t)$  and  $Q(\psi, x, t)$  can be easily computed recursively.  $P(\psi, x, 0)$  is a given initial distribution, and  $P(\psi, x, t+1)$  can be obtained from  $P(\psi, x, t)$  using the Markov transition rule and  $\psi(\cdot, t)$ . Similarly,  $Q(\psi, x, \tau) \equiv 1$ , and  $Q(\psi, x, t-1)$  can be obtained from  $Q(\psi, x, t)$  using the Markov transition rule and  $\psi(\cdot, t)$ . After obtaining  $P(\psi, x, t)$  and  $Q(\psi, x, t)$ , it is a simple matter to compute  $D_\tau(\psi, x, t)$  and then  $\Delta(\psi)$ .

Algorithms for finding the optimal effort distribution  $\psi^*$  typically operate by generating a sequence  $\psi_1, \psi_2, \dots$  that approaches  $\psi^*$ . The method of computation is such that  $\Delta(\psi_1)$  can be computed with only slightly more effort. Consider the discrete case. For any  $t$ , the probability of detection is

$$(7) \quad P(\psi) = 1 - \sum_x P(\psi, x, t) \exp(-w(x, t) \psi(x, t)) Q(\psi, x, t)$$

In order to find  $\psi^*$ , we first make an initial guess  $Q^0(x, t)$  and then calculate  $\psi^r$  the function that minimizes

$$\sum_x P(\psi, x, t) \exp(-w(x, t) \psi(x, t)) Q^{n-1}(x, t) \quad \text{for } n \geq 1,$$

with  $Q^n(x, t) = Q(\psi^n, x, t)$  for  $n \geq 1$ . Each of the minimization problems is relatively simple, and it can be shown that  $P(\psi^n)$  increases with  $n$  [1,2,3].

For each of the minimization problems, there must exist a function  $\lambda_n(t)$  such that  $D_n(x, t) = \lambda_n(t)$  when  $\psi^n(x, t) > 0$  and  $D_n(x, t) \leq \lambda_n(t)$  when  $\psi^n(x, t) = 0$ , where

$$D_n(x, t) = w(x, t) P(\psi^n, x, t) \exp(-w(x, t) \psi^n(x, t)) Q^{n-1}(x, t).$$

These are simply the Kuhn-Tucker conditions; in practice, some sort of a search is made until  $\lambda_n(t)$  is found such that  $\sum_x \psi(x, t) = m(t)$ . But

$$(8) \quad D_T(\psi^n, x, t) \begin{cases} = \lambda_n(t) Q^n(x, t)/Q^{n-1}(x, t) & \text{when } \psi^n(x, t) > 0 \\ \leq \lambda_n(t) Q^n(x, t)/Q^{n-1}(x, t) & \text{when } \psi^n(x, t) = 0 \end{cases}$$

so computation of  $D_T(\psi^n, x, t)$  is a small burden given that  $\lambda_n(t)$ ,  $Q^n(x, t)$ , and  $Q^{n-1}(x, t)$  have to be computed in any case.

In the continuous case,  $\sum_x$  is replaced by  $\int dx$ ; otherwise, the development remains the same.

#### 4. AN EXAMPLE

We set  $\tau = 79$ , so there are 80 looks. The target does a random walk over the cells  $1, \dots, 67$ , starting at cell 34. At each opportunity, it moves left with probability .3, right with probability .3, or else does not move. The boundary is reflecting. We set  $w(x,t) = .001625$  and  $m(t) = 100$ ; if all 100 units of search effort are used in a single cell, the probability of detection would be  $1 - \exp(-.1625) = .15$  at each of the 80 opportunities. We also set  $Q^0(x,t) \equiv 1$ .

The first five detection probabilities (and values for  $\Delta(\psi)$ ) are: .7054 (.0393), .7075 (.0246), .7086 (.0120), .7090 (.0067), and .7092 (.0036). The amounts of CPU time required on the NPS IBM 360/67 for computation of each of these five pairs were 2.4, 2.8, 2.4, 2.0, and 2.0 seconds, respectively. The second pass always takes a relatively long time because  $Q^1(x,t)$  is radically different from  $Q^0(x,t)$ . The function  $\exp(-w(x,t) \psi^5(x,t))$  is shown to three decimal places in Figure 1, with \*\*\*\* indicating no search. Time reads down the page (there are 80 rows). Only the central cells are shown; the rest are not searched.

It is interesting to compare these results with those that result when all effort must be placed in a single cell at each time. The best probability of detection in that case is apparently .64--sufficient conditions for optimality are not known in that case, but the author has done sufficient

experimentation to conjecture that .64 is the answer. The increase from .64 to .7092 can, of course, be attributed to the relaxation of a constraint.

## REFERENCES

- (1) "Numerical Optimization of Search for a Moving Target," Daniel H. Wagner, Associates Report to Office of Naval Research, by L. D. Stone et al., 23 June 1978.
- (2) "Optimal Search for a Moving Target in Discrete Time and Space," by S. S. Brown (submitted to Opns. Res.).
- (3) "On Search for a Moving Target," by A. R. Washburn (submitted to NRLQ).

FIGURE 1. Survival Probabilities for Near Optimal Search.

Table with multiple columns of survival probability data, organized in a grid-like structure. The data points are numerical values representing survival probabilities, ranging from approximately 0.55 to 0.99.

0.550	0.650	0.750	0.850	0.950
0.557	0.657	0.757	0.857	0.957
0.564	0.664	0.764	0.864	0.964
0.571	0.671	0.771	0.871	0.971
0.578	0.678	0.778	0.878	0.978
0.585	0.685	0.785	0.885	0.985
0.592	0.692	0.792	0.892	0.992
0.599	0.699	0.799	0.899	0.999
0.606	0.706	0.806	0.906	1.006
0.613	0.713	0.813	0.913	1.013
0.620	0.720	0.820	0.920	1.020
0.627	0.727	0.827	0.927	1.027
0.634	0.734	0.834	0.934	1.034
0.641	0.741	0.841	0.941	1.041
0.648	0.748	0.848	0.948	1.048
0.655	0.755	0.855	0.955	1.055
0.662	0.762	0.862	0.962	1.062
0.669	0.769	0.869	0.969	1.069
0.676	0.776	0.876	0.976	1.076
0.683	0.783	0.883	0.983	1.083
0.690	0.790	0.890	0.990	1.090
0.697	0.797	0.897	0.997	1.097
0.704	0.804	0.904	1.004	1.104
0.711	0.811	0.911	1.011	1.111
0.718	0.818	0.918	1.018	1.118
0.725	0.825	0.925	1.025	1.125
0.732	0.832	0.932	1.032	1.132
0.739	0.839	0.939	1.039	1.139
0.746	0.846	0.946	1.046	1.146
0.753	0.853	0.953	1.053	1.153
0.760	0.860	0.960	1.060	1.160
0.767	0.867	0.967	1.067	1.167
0.774	0.874	0.974	1.074	1.174
0.781	0.881	0.981	1.081	1.181
0.788	0.888	0.988	1.088	1.188
0.795	0.895	0.995	1.095	1.195
0.802	0.902	1.002	1.102	1.202
0.809	0.909	1.009	1.109	1.209
0.816	0.916	1.016	1.116	1.216
0.823	0.923	1.023	1.123	1.223
0.830	0.930	1.030	1.130	1.230
0.837	0.937	1.037	1.137	1.237
0.844	0.944	1.044	1.144	1.244
0.851	0.951	1.051	1.151	1.251
0.858	0.958	1.058	1.158	1.258
0.865	0.965	1.065	1.165	1.265
0.872	0.972	1.072	1.172	1.272
0.879	0.979	1.079	1.179	1.279
0.886	0.986	1.086	1.186	1.286
0.893	0.993	1.093	1.193	1.293
0.900	1.000	1.100	1.200	1.300
0.907	1.007	1.107	1.207	1.307
0.914	1.014	1.114	1.214	1.314
0.921	1.021	1.121	1.221	1.321
0.928	1.028	1.128	1.228	1.328
0.935	1.035	1.135	1.235	1.335
0.942	1.042	1.142	1.242	1.342
0.949	1.049	1.149	1.249	1.349
0.956	1.056	1.156	1.256	1.356
0.963	1.063	1.163	1.263	1.363
0.970	1.070	1.170	1.270	1.370
0.977	1.077	1.177	1.277	1.377
0.984	1.084	1.184	1.284	1.384
0.991	1.091	1.191	1.291	1.391
0.998	1.098	1.198	1.298	1.398
1.005	1.105	1.205	1.305	1.405
1.012	1.112	1.212	1.312	1.412
1.019	1.119	1.219	1.319	1.419
1.026	1.126	1.226	1.326	1.426
1.033	1.133	1.233	1.333	1.433
1.040	1.140	1.240	1.340	1.440
1.047	1.147	1.247	1.347	1.447
1.054	1.154	1.254	1.354	1.454
1.061	1.161	1.261	1.361	1.461
1.068	1.168	1.268	1.368	1.468
1.075	1.175	1.275	1.375	1.475
1.082	1.182	1.282	1.382	1.482
1.089	1.189	1.289	1.389	1.489
1.096	1.196	1.296	1.396	1.496
1.103	1.203	1.303	1.403	1.503
1.110	1.210	1.310	1.410	1.510
1.117	1.217	1.317	1.417	1.517
1.124	1.224	1.324	1.424	1.524
1.131	1.231	1.331	1.431	1.531
1.138	1.238	1.338	1.438	1.538
1.145	1.245	1.345	1.445	1.545
1.152	1.252	1.352	1.452	1.552
1.159	1.259	1.359	1.459	1.559
1.166	1.266	1.366	1.466	1.566
1.173	1.273	1.373	1.473	1.573
1.180	1.280	1.380	1.480	1.580
1.187	1.287	1.387	1.487	1.587
1.194	1.294	1.394	1.494	1.594
1.201	1.301	1.401	1.501	1.601
1.208	1.308	1.408	1.508	1.608
1.215	1.315	1.415	1.515	1.615
1.222	1.322	1.422	1.522	1.622
1.229	1.329	1.429	1.529	1.629
1.236	1.336	1.436	1.536	1.636
1.243	1.343	1.443	1.543	1.643
1.250	1.350	1.450	1.550	1.650
1.257	1.357	1.457	1.557	1.657
1.264	1.364	1.464	1.564	1.664
1.271	1.371	1.471	1.571	1.671
1.278	1.378	1.478	1.578	1.678
1.285	1.385	1.485	1.585	1.685
1.292	1.392	1.492	1.592	1.692
1.299	1.399	1.499	1.599	1.699
1.306	1.406	1.506	1.606	1.706
1.313	1.413	1.513	1.613	1.713
1.320	1.420	1.520	1.620	1.720
1.327	1.427	1.527	1.627	1.727
1.334	1.434	1.534	1.634	1.734
1.341	1.441	1.541	1.641	1.741
1.348	1.448	1.548	1.648	1.748
1.355	1.455	1.555	1.655	1.755
1.362	1.462	1.562	1.662	1.762
1.369	1.469	1.569	1.669	1.769
1.376	1.476	1.576	1.676	1.776
1.383	1.483	1.583	1.683	1.783
1.390	1.490	1.590	1.690	1.790
1.397	1.497	1.597	1.697	1.797
1.404	1.504	1.604	1.704	1.804
1.411	1.511	1.611	1.711	1.811
1.418	1.518	1.618	1.718	1.818
1.425	1.525	1.625	1.725	1.825
1.432	1.532	1.632	1.732	1.832
1.439	1.539	1.639	1.739	1.839
1.446	1.546	1.646	1.746	1.846
1.453	1.553	1.653	1.753	1.853
1.460	1.560	1.660	1.760	1.860
1.467	1.567	1.667	1.767	1.867
1.474	1.574	1.674	1.774	1.874
1.481	1.581	1.681	1.781	1.881
1.488	1.588	1.688	1.788	1.888
1.495	1.595	1.695	1.795	1.895
1.502	1.602	1.702	1.802	1.902
1.509	1.609	1.709	1.809	1.909
1.516	1.616	1.716	1.816	1.916
1.523	1.623	1.723	1.823	1.923
1.530	1.630	1.730	1.830	1.930
1.537	1.637	1.737	1.837	1.937
1.544	1.644	1.744	1.844	1.944
1.551	1.651	1.751	1.851	1.951
1.558	1.658	1.758	1.858	1.958
1.565	1.665	1.765	1.865	1.965
1.572	1.672	1.772	1.872	1.972
1.579	1.679	1.779	1.879	1.979
1.586	1.686	1.786	1.886	1.986
1.593	1.693	1.793	1.893	1.993
1.600	1.700	1.800	1.900	2.000

DISTRIBUTION LIST

	No. of Copies
Defense Documentation Center Cameron Station Alexandria, VA 22314	2
Library, Code 0212 Naval Postgraduate School Monterey, CA 93940	2
Library, Code 55 Department of Operations Research Naval Postgraduate School Monterey, Ca. 93940	2
Dean of Research, Code 012 Naval Postgraduate School Monterey, CA 93940	1
COMPATWINGSPAC Naval Air Station, Code 532 Moffett Field, CA 94035	1
COMSUBPAC, ATTN: Cdr. Hayne FPO San Francisco, CA 96610	1
Office of Naval Research Code 431 Arlington, VA 22217	1
Center for Naval Analyses, 1401 Wilson Blvd., Arlington, VA 22209 Attn: Marc Mangel	1
Operations Research Center Room 24-215 Massachusetts Institute of Technology Cambridge, MA 02139	1
ORI, Inc. 1400 Spring Street Silver Spring, MD 20910	1



DISTRIBUTION LIST CONT.

	No. of Copies
Director Strategic Systems Project Office 1931 Jefferson Davis Highway Arlington, VA 20376 ATTN: Code 2024	1
Professor Takasi Kisi Department of Applied Physics Defense Academy Yokosuka, Japan	1
Professor A. R. Washburn Code 55Ws Naval Postgraduate School Monterey, CA 93940	10
R. J. Stampfel Code 55 Naval Postgraduate School Monterey, CA 93940	1
R. N. Forrest Code 71 Naval Postgraduate School Monterey, Ca. 93940	1