

UNITED STATES ARMY ENVIRONMENTAL HYGIENE AGENCY

ABERDEEN PROVING GROUND, MD 21010

*

SENTINEL

ADA064243

FILE COPY

JOC

DETERMINATION OF URINE METABOLITE LEVELS FOLLOWING INHALATION OF THE INSECTICIDE PERMETHRIN IN RATS STUDY NO. 75-53-0053-79 MAY - AUGUST 1978

FEB 6 1979

Approved for public release; distribution unlimited.

79 02 02 029

REFORT DOCOMENTATION FAC	E READ INSTRUCTIONS BEFORE COMPLETING FORM
. REPORT NUMBER 2. GO	TACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER
75-53-0053-79	
TITLE (and Sublice) Determination of Urine Metabolite Leve	1s / AJR & DT. FAW
Following Inhalation of the Insecticid	May - August 1978
Permethrin in Rats.	6. PERFORMING UNG. REPORT HUMBER
· AUTHOR(a)	S. CONTRACT OR GRANT NUMBER(+)
Jeffrey A. Geres Ph.D.	
Roger E. Boldt. / Ph.D.	
- PERFORMING ORDERIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TAS
Aberdeen Proving Ground, MD 21010	
Commander	May - August 1078
US Army Health Services Command	13. NUMBER GE PAGES
FORT Sam HOUSTON, TX 78234 4. MONITORING AGENCY NAME & ADDRESS(II different from	Controlling Office) 15. SECURITY CLASS. (of this report)
	linelandet
-	
	SCHEDULE
6. DISTRIBUTION STATEMENT (of this Report)	
Approved for public release: distribut	ion unlimited.
(I) A	
$(12) 9 p_1$	
7. DISTRIBUTION STATEMENT (of the obstract entered in Big	ock 20, 11 different from Report)
AND A -WA MN NA	NANA HAT
(1) 45A EHA-15-53	-0053-11/
. SUPPLEMENTARY NOTES	-
	tily by block number)
9. KEY WORDS (Continue on revoree elde if necessary and iden	
9. KEY WORDS (Continue on reverse elde if necessary and iden 3-phenoxybenzyl-(+) cis, trans-3-(2,2-	dichloroviny1)-2,2-dimethylcyclopropane
 KEY WORDS (Continue on reverse elde if necessary and iden 3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity 	dichlorovinyl)-2,2-dimethylcyclopropane
KEY WORDS (Continue on reverse elde if necessary and iden 3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine	Rat (+ or -)
 KEY WORDS (Continue on reverse olde if necessary and iden 3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine ADDITACT (Continue on prove olde if measures and iden 	Rat (+ or -)
9. KEY WORDS (Continue on reverse elde II necessary and iden 3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine ABSTRACT (Continue on reverse of M messessiry and ident the Pyrethroid insecticide Permethrin	Rat (+ or -) ity by block mumber) 3-phenoxybenzyl-bd-cis, trans-3-(2,2)
3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine ABSTRACT (Continue or covers at 11 mesocory of 14mm The Pyrethroid insecticide Permethrin, dichlorovinyl)-2,2-dimethylcyclopropan	Rat (+ or -) HT by block number) 3-phenoxybenzy1-pht-cis, trans-3-(2,2) lecarboxylate, was administered to rats
 KEY WORDS (Continue on reverse olds if necessary and iden 3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine ABSTRACT (Continue on reverse of M messary and iden The Pyrethroid insecticide Permethrin, dichlorovinyl)-2,2-dimethylcyclopropan as an aerosol. Metabolite studies of 	Rat (+ or -) 3-phenoxybenzyl-pd-cis, trans-3-(2,2) lite tr block mumber) 3-phenoxybenzyl-pd-cis, trans-3-(2,2) lecarboxylate, was administered to rats the urine showed the presence of
3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine ABSTRACT (Continue on reverse of M measuremy of Idea The Pyrethroid insecticide Permethrin, dichlorovinyl)-2,2-dimethylcyclopropan as an aerosol. Metabolite studies of significant quantities of Permethrin m exposure. The pattern of metabolite in	Rat (+ or -) A s-phenoxybenzyl-bd-cis, trans-3-(2,2) recarboxylate, was administered to rats the urine showed the presence of retabolites for several days following excretion indicates an extremely
3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine ABSTRACT (Contract of Marcasson of Marc	Rat (+ or -) A-phenoxybenzyl-bd-cis, trans-3-(2,2) recarboxylate, was administered to rats the urine showed the presence of retabolites for several days following excretion indicates an extremely and excretion of this compound.
3-phenoxybenzyl-(+) cis, trans-3-(2,2- carboxylate toxicity Permethrin Metabolic SBP-1513 Urine ASSTRACT (Continue on reverse of N measurement of the intermethrin Metabolic SBP-1513 Urine ASSTRACT (Continue on reverse of N measurement of the intermethrin intermethrin intermethr	Rat (+ or-) A 3-phenoxybenzyl-bd-cis, trans-3-(2,2) recarboxylate, was administered to rats the urine showed the presence of retabolites for several days following excretion indicates an extremely and excretion of this compound.
ABSTRACT (Continue on reverse olde if necessary and iden ABSTRACT (Continue on reverse olde if necessary and iden ABSTRACT (Continue on reverse olde if necessary and iden The Pyrethroid insecticide Permethrin, dichlorovinyl)-2,2-dimethylcyclopropan as an aerosol. Metabolite studies of significant quantities of Permethrin m exposure. The pattern of metabolite e efficient mechanism for the metabolism D 1000 1473 EDTTION OF 1 NOV 68 IS OBSOLETE	Rat (+ or -) ity by block number) 3-phenoxybenzyl-pki-cis, trans-3-(2,2) lecarboxylate, was administered to rats the urine showed the presence of wetabolites for several days following excretion indicates an extremely and excretion of this compound.



HSE-LR-B/WP

DEPARTMENT OF THE ARMY U.S. ARMY ENVIRONMENTAL HYGIENE AGENCY ABERDEEN PROVING GROUND, MARYLAND 21010

CPT Gere/1m/584-2810

24 JAN 1979

SUBJECT: Determination of Urine Metabolite Levels Following Inhalation of the Insecticide Permethrin in Rats, Study No. 75-53-0053-79, May-August 1978

Executive Secretary Armed Forces Pest Control Board Forest Glen Section, WRAMC Washington, DC 20012

A summary of the pertinent findings of the inclosed report follows:

The pattern of primary excretion in rats of Permethrin during and following inhalation exposure was measured. The urine was found to contain metabolite that quickly reaches a dose related concentration permitting measurement of adsorption and excretion rates.

FOR THE COMMANDER:

1 Incl as (5 cy)

BRENDAN E. JOYCE, Ah.D. LTC, MSC Director, Laboratory Services

CF: HQDA (DASG-PSP) Cdr, HSC (HSPA-P) Supt, AHS (HSA-IHE) Dir, Advisory Ctr on Tox, NRC USDA-ARS, Southern Region/Dr. Weidhaas USDA-ARS, Southern Region/Cdr Grotehaus

NTIS	White Section 🖠
DDC	S C Section [
UNANNOLIN	icia (1
10341-103	1949

••••••	*****
5 Y	
BY DISTINCTION	MANAL ANT ITY DROES
BY DISTINCT Dist. A.	MI/AYAC ABILITY CODES
BY Distinctin	MAYAF ABILITY OBDES
DIST. A.	MU/AVALIABILITY OBDES
BY DISTINGUT	MI/AVAL'ABILITY OBDES

79 02 02 029



DEPARTMENT OF THE ARMY U.S. ARMY ENVIRONMENTAL HYGIENE AGENCY ABERDEEN PROVING GROUND, MARYLAND 21010

HSE-LR-B/WP

DETERMINATION OF URINE METABOLITE LEVELS FOLLOWING INHALATION OF THE INSECTICIDE PERMETHRIN IN RATS STUDY NO. 75-53-0053-79 MAY - AUGUST 1978

1. AUTHORITY.

a. Memorandum of Understanding between the Department of the Army, Office of The Surgeon General, the US Army Health Services Command, the US Army Environmental Hygiene Agency, the Armed Forces Pest Control Board (AFPCB), and the US Department of Agriculture, effective December 1970 with Amendment No. 1, effective August 1974.

b. Letter, AFPCB, Armed Forces Pest Control Board, 21 October 1975, subject: Request for Toxicological Evaluation.

c. Letter, AFPCB, Armed Forces Pest Control Board, 5 April 1977, subject: Request for Toxicological Evaluation.

2. REFERENCES.

a. Interim Report, HSE-LT, this Agency, Preliminary Assessment of Relative Toxicity of Candidate Insect Repellent AI3-29158, 3-Phenoxybenzyl cis/trans 2,2-dimethyl-3-(2,2-dichlorovinyl) cyclopropanecarboxylate Study No. 51-031-76, December 1975 - April 1976.

b. Report, HSE-LT, this Agency, Toxicological Evaluation of 3-(phenoxyphenyl) methyl (+)-cis, trans-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate (Permethrin), Study No. 51-0831-78, December 1975 - April 1977.

c. Toxicology Division Procedural Guide, USAEHA, 1972, revised 1976.

3. PURPOSE. The purpose of this investigation was to determine the pattern of Permethrin metabolite excretion in the urine of rats which were being used in an inhalation toxicology study. The results of the toxicology study will be detailed in a separate USAEHA report. Results of the present investigation are for advising the AFPCB on the possible hazard in use of this compound with human subjects.

> The use of company names or their product does not imply indorsements by the US Army, but is intended only to assist in identification of a specific compound.

Approved for public release; distribution unlimited.

4. GENERAL.

a. Permethrin [3-phenoxybenzyl- (\pm) -cis, trans-3-(2,2-dichlorovinyl)-2,2dimethylcyclopropanecarboxylate], a synthetic pyrethroid, is an amber colored liquid having a specific gravity of 1.2, melting point of 35°F and boiling point of 220°F. It is soluble in organic solvents (acetone, hexane). The sample studied was SBP-1513, lot No. RAX-6*.

b. Detailed descriptions of the inhalation exposures will be reported elsewhere¹. Four male Sprague-Dawley rats per group were exposed to Permethrin aerosols at concentations of 0, 125, 250, and 500 mg/M³. The animals were exposed for 6 hours per day, 5 days per week for 13 weeks. No exposures were performed on Saturdays or Sundays. The metabolite excretion determinations were begun on Monday of the thirteenth week of exposure.

c. The animals were removed from the inhalation chambers or holding cages (Saturday and Sunday), where they were individually caged, and placed into individual metabolism cages. Urine was collected in plastic cups until 8:00 AM the following morning, when the animals were again placed in inhalation chambers or holding cages.

d. The volume of the individual urine samples was measured each morning and the concentration of 3-(2,2-dichlorovinyl)-2,2dimethylyclopropanecarboxylic acid (C1VA) was determined by the as yet unpublished procedure of Cridland and Weatherley². Excreted almost exclusively as the glucuronide and sulfate esters, this metabolite is known to appear rapidly in the urine of animals exposed to Permethrin,^{3 4} and to represent over 50% of an administrered oral dose⁴. The method of analysis includes the enzymatic hydrolysis of the metabolite, esterification and determination by gas chromatography. Authentic C1VAt was used for standarization.

e. Literature cited in this report is shown in the Appendix.

5. FINDINGS.

a. Extraction of C1VA, after enzymatic hydrolsis of the metabolite, was accomplished with hexane followed by centrifugation to separate the phases. Phase separation was difficult because of an apparent increase in lipid content of the rat urine, becoming impossible in those urine samples collected within 24 hours after exposure of the animals to the highest dose of Permethrin. Phases were well resolved in urine from rats receving lower doses, or on Saturday or Sunday when the animals were not exposed to Permethrin.

2

^{*} SBP-1513 is a company designation of CPC International, Inc., S.B. Penick and Co., 160 Church Street, NY, NY 10007. † Obtained from Dr. James Hubbell, Burroughs-Wellcome Co., Research Triangle Park, NC 27709.

b. Permethrin metabolite concentration analyses were performed on the urine collected as described. The results of analyses are shown in the Table and Figure. The concentration of Permethrin metabolite increased during the week, reaching a maximum on Thursday and Friday. A dramatic decrease took place on Saturday, when no exposure occurred; a further decrease was apparent on Sunday. This pattern was true for all dose levels, animals receiving the highest dose excreted approximately three times more of the metabolite than those exposed to the lower dose levels. Control animals uniformly excreted a very low (below 20 g/day) quantity of the metabolite.

TABLE.	Average C1VA Conter 500 mg/M ³	t of Rat Urine 250 mg/M ³	(g/Sample/Rat) 125 mg/M ³	13th Week. 0 mg/M ³
Monday	2140 + 172	401 + 67	365 + 16	< 20
Tuesday	2360 7 401	540 + 100	550 + 4	< 20
Wednesday	y 2843 + 343	415 + 30	788 + 76	< 20
Thursday	2879 + 190	663 + 63	1058 + 124	< 20
Friday	2777 7 60	827 7 24	959 + 35	< 20
Saturday	884 + 88	446 + 37	195 + 19	< 20
Sunday	598 + 36	329 7 26	229 + 15	< 20

6. DISCUSSION. Administration of the pyrethroid insecticide, Permethrin, by inhalation caused a gradual increase in the rate of urinary Permethrin metabolite excretion, reaching a maximum in 3-4 days. It is especially interesting that the urine concentration rapidly fell after inhalation exposures were discontinued. Animals receving the highest dose of Permethrin excreted approximately three times as much of the metabolite as those receiving the lower doses. Only 2 days after the cessation of exposures, the excretion rate of all animals had dropped to roughly the same level. These data indicate that the animals possess a very efficient mechanism for the detoxification and excretion of Permethrin. This permits a drop in the excretion level to only 20-45 percent of the test levels by the second day after the end of the exposure period. Support for this view comes from a recent report⁵ on the purification and characterization, from rat liver microsomes, of the enzyme pyrethroid carboxyesterase, which hydrolyzes esters of chrysanthemumic acid such as Permethrin.

3





FIGURE I: PERMETHRIN CONTENT OF RAT URINE, µg/RAT/DAY

4

7. CONCLUSION. The urine of rats exposed to Permethrin by inhalation has been found to contain a pyrethroid metabolite which very quickly reaches a dose-related concentration showing that an inhaled aerosol of the pesticide is readily absorbed, metabolized and excreted in the urine.

CPT, MSC Biochemist Radiological & Biological Chemistry Division

E. Break

ROGER E. BOLDT, Ph.D. Chief, Biochemistry Branch Radiological & Biological Chemistry Division

APPROVED:

ALLHUS LA JONES Chief, Radiological & Biological Chemistry Division

APPENDIX

LITERATURE CITED

1. Metker, L. and M. Weeks, USAEHA Internal Report, in preparation.

2. Cridland, J.S. and B. C. Weatherley, Wellcome Internal Report 77-2, 1977, Wellcome Research Laboratories, Beckenham, England.

3. Unai, T. and J.E. Casida, J. Agric. Food Chem., 25, 979 (1977).

4. Gaughan, L.C., T. Unai and J.E. Casida, J. Agric. Food Chem., <u>25</u>, 9 (1977).

5. Suzuki, T. and J. Miyamoto, Pestic. Biochem. Physiol., 8, 186 (1978).