



CAN ANY STATIONARY ITERATION USING LINEAR INFORMATION BE GLOBALLY CONVERGENT?

G. W. Wasilkowski Department of Computer Science Carnegie-Mellon University Pittsburgh, Pennsylvania 15213 (On leave from the University of Warsaw)

December 1978

C MANNE 6 1979 FFR U

This research was supported in part by the National Science Foundation under Grant MCS 75-222-55 and the Office of Naval Research under Contract N00014-76-C-0370, NR 044-422.

ABSTRACT

All known globally convergent iterations for the solution of a nonlinear operator equation f(x) = 0 are either non-stationary or use nonlinear information. We ask whether there exists a globally convergent stationary iteration which uses linear information. We prove that even if global convergence is defined in a weak sense, there exists no such iteration for as simple a class of problems as the set of all analytic complex functions having only simple zeros. We conjecture that even for the class of all real polynomials which have real simple zeros there does not exist a globally convergent stationary iteration using linear information.

ACCESSION for White Section NTIS certion. 000 Sin CIAL

1. INTRODUCTION

Suppose we solve a nonlinear operator equation f(x) = 0 by an iteration which constructs a sequence of approximations $\{x_i\}$. For most convergence theorems it is assumed that f is sufficiently smooth and starting points are "sufficiently close" to a solution α . In practice it is very hard to verify the second assumption. One therefore wants to use iterations which are globally convergent.

All known globally convergent iterations are either non-stationary or use nonlinear information. For instance, Laguerre's iteration (see for instance Ralston and Rabinowitz [78]) is globally convergent for the class of all real polynomials having only real zeros. However this iteration uses the degree of the polynomial whose zero is approximated. This means that Laguerre's iteration uses <u>nonlinear information</u> (see Section 6). An example of a <u>non-stationary</u> iteration using linear information which is globally convergent for analytic operator equations may be found in Traub and Woźniakowski [76].

However, most commonly used iterations are stationary and use linear information. Therefore it is important to know whether there exist globally convergent stationary iterations which use linear information. In this paper we prove that for as simple a class of problems as the set of all analytic complex functions with simple zeros there exists no such iteration. We conjecture that the same negative result holds even for the class of all real polynomials having real simple zeros.

We summarize the contents of the paper. In Section 2 we remind the reader of the definitions of <u>information</u> and <u>stationary iteration without</u>

<u>memory</u>. In Section 3 we discuss the concept of global convergence. In Section 4 we show that no stationary iteration without memory which uses linear information can be globally convergent for the class of all analytic complex functions. In Section 5 we extend this result to all stationary iterations with or without memory using linear information. In Section 6 we pose a conjecture that for the class of all real polynomials with real simple zeros there does not exist a globally convergent stationary iteration using linear information.

2. STATIONARY ITERATIONS WITHOUT MEMORY

We recall the definition of <u>information</u> and <u>stationary iteration</u>. See Traub and Woźniakowski [78]. For the reader's convenience, in Sections 2 to 4 we deal only with one-point iterations without memory. The extension to the general case is given in Section 5.

Consider the solution of a nonlinear scalar equation

(2.1) f(x) = 0

for $f \in \mathfrak{J}$ where \mathfrak{J} is a subset of a space \mathfrak{H} of functions $f: \mathbb{D}_{f} \subset \mathbb{C} \to \mathbb{C}$. To solve (2.1) iteratively we need to know something about f. Let $L_{i}: \mathbb{D}_{L_{i}} \subset \mathfrak{H} \times \mathbb{C} \to \mathbb{C}$ be a functional which is linear with respect to the first argument, i.e.,

 $L_{i}(c_{1}f_{1}+c_{2}f_{2},x) = c_{1}L_{i}(f_{1},x) + c_{2}L_{i}(f_{1},x) \text{ whenever } x \in D_{f_{1}} \cap D_{f_{2}}, \quad i=1,2,\ldots,n.$ Consider the <u>linear information operator</u> $\mathfrak{N}, \mathfrak{N}: D_{\mathfrak{N}} \subset \mathbb{H} \times \mathbb{C} \to \mathbb{C}^{n}$, defined as $(2.2) \quad \mathfrak{N}(f,x) = [L_{1}(f,x), L_{2}(f,x), \ldots, L_{n}(f,x)], \quad \forall f \in \mathbb{H}, \quad \forall fx \in D_{f}.$

Let x_0 be an approximation of a solution of (2.1). Let $\varphi: D_{\varphi} \subset \mathbb{C}^{n+1} \to \mathbb{C}$ be a functional. We construct the sequence of approximations x_i by the formula

(2.3) $x_{i+1} = \varphi(x_i; \Re(f, x_i)).$

The functional φ is called a <u>one-point stationary iterative operator without</u> <u>memory using a linear information operator \Re </u>. For brevity φ is called an iteration. Let $\Phi(\Re)$ be the class of all such iterations.

Note that most iterations use values of f and its derivatives. A linear information operator is a generalization of this. For example, the information operator \Re used by Newton iteration

$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$

is $\Re(f,x) = [f(x),f'(x)]$. This operator is linear and $\varphi \in \Phi(\Re)$.

3. WHAT DO WE MEAN BY GLOBAL CONVERGENCE?

Let \Im be the class of all functions f,f: $D_f \subset \mathbb{C} \to \mathbb{C}$, analytic in D_f and having only simple zeros. Let S(f) be the set of all zeros of f.

Consider any iteration $\varphi \in \Phi$, where Φ is the class of all stationary iterations without memory which use a linear information operator \Re per iterative step, i.e., $\Phi = \bigcup_{\mathfrak{M}} \Phi(\mathfrak{R})$.

Which properties should ϕ have to be called a globally convergent iteration? To motivate our definition consider first the problem f(x) = 0 where f is defined in $D_f = \{x: |x-\alpha| \le R(f)\}$ and α is its unique simple zero. Suppose we apply φ to this problem. Let $r(f, \varphi)$ be the maximal number such that for any starting point x_0 satisfying $|x_0 - \alpha| < r(f, \varphi)$ the sequence $x_{i+1} = \varphi(x_i, \Re(f, x_i))$ is convergent to α . Then the ball $\{x: |x-\alpha| < r(f, \varphi)\}$ is called the <u>ball of convergence of φ for f</u>. Of course, $r(f, \varphi)$ depends on R(f) and in particular $r(f, \varphi) \leq R(f)$. Suppose there exists a positive constant $c = c(\phi)$ such that $r(f,\phi) \ge c R(f)$ for any f. Then ϕ enjoys a type of global convergence since the ball of convergence has radius proportional to the radius R(f) of the domain D_f . However for problems with R(f) = + ∞ we get $r(f, \varphi) = +\infty$ which means that any choice of a starting point x_0 yields convergence. This seems to be too strong. For $R(f) = +\infty$ we would like to have $r(f, \phi)$ large but not necessarily equal to infinity. This is the motivation of the following. Let K,L be two given constants such that $K \ge 0$ and $0 < L \le +\infty$. Define $R_f = \min\{R(f), K | \alpha| + L\}$. Now the existence of a positive constant $c = c(\phi)$ such that $r(f,\phi) \ge c R_f$ implies a type of global convergence φ . This discussion shows that we should compare $r(f,\varphi)$ with R_f.

If f has more than one simple zero we proceed as follows. Let dist(α , ∂D_f) be the distance of α , $\alpha \in S(f)$, to the boundary ∂D_f of the domain D_f . Define

$$R_{f}(\alpha) = \min\{dist(\alpha,\partial D_{f}), K|\alpha| + L\}, (K \ge 0, 0 < L \le + \infty).$$

$$B(b,f) = \bigcup_{\alpha \in S(f)} \{x: |x-\alpha| < b \cdot R_f(\alpha)\}$$

where $b \ge 0$. Note that if $S(f) = \{\alpha\}$ then $R_f(\alpha) = R_f$. For any iteration φ , $\varphi \in \Phi(\mathbb{R})$, we define a number $c(\varphi)$ such that

- (i) for any f ∈ ℑ and for any starting point x₀ ∈ B(c(φ), f) the sequence {x_i}, x_{i+1} = φ(x_i; ℜ(f,x_i)), is well-defined and lim x_i ∈ S(f).
- (ii) for any ε > 0 there exist a problem f ∈ 3 and a starting point x₀ ∈ B(c(φ) + ε, f) such that either the sequence {x_i} is not well-defined or lim x_i ∉ S(f).

Note that for any iteration $c(\phi) \in [0,1]$. The set $B(c(\phi),f)$ is a convergence domain of ϕ for the function f since taking any starting point $x_0 \in B(c(\phi), f)$ we get convergence of $\{x_i\}$. Note, however, that we do not specify which element from S(f) is the limit of $\{x_i\}$.

Definition 3.1

Let

We shall say that an iteration $\phi, \ \phi \in \Phi,$ is globally convergent for the class 3 iff

 $c(\phi) > 0.$

Definition 3.1 imposes only a weak condition on φ . However, we shall show that for any iteration from Φ , $c(\varphi) = 0$. This means that even in the sense of Definition 3.1 there exists no globally convergent stationary iteration using linear information for the class 3.

4. MAIN RESULT

Theorem 4.1

No iteration ϕ from ϕ is globally convergent for the class 3.

Proof

Suppose there exist $\mathfrak{N}: D_{\mathfrak{N}} \to \mathbb{C}^n$ and $\varphi \in \Phi(\mathfrak{N})$ with $c = c(\varphi) > 0$. Let

$$\mathbf{a} = \begin{cases} 1 & \text{if } cK \ge 1, \\ \frac{cL}{2(1-cK)} & \text{if } cK < 1. \end{cases}$$

Define f(x) = x-a. Then $f \in \mathcal{J}$, $S(f) = \{a\}$ and $R_f(a) = K|a| + L$. Let $x_0 = 0$. Since $x_0 \in B(c, f) = \{x; |x-a| < cR_f(a)\}$, the sequence $\{x_i\}$, $x_{i+1} = \varphi(x_i; \Re(f, x_i))$, tends to a, $a \neq 0$. Thus, there exists a unique integer k, $k \geq 1$, such that

(4.1)
$$|x_0|, |x_1|, \dots, |x_{k-1}| < \frac{a}{2}$$
 and $|x_k| \ge \frac{a}{2}$.

Consider a polynomial w of the form

(4.2) w(x) =
$$\sum_{i=0}^{n \cdot k} a_i x^{i+1}$$

which satisfies

(4.3)
$$\Re(f-w,x_j) = \Re(f,x_j)$$
 for $j = 0,1,...,k-1$.

This is equivalent to the following homogeneous system of $n \cdot k$ linear equations

(4.4)
$$\sum_{i=0}^{n \cdot k} a_{i}L_{s}(x^{i+1}, x_{j}) = 0$$
 for $s = 1, 2, ..., n$ and $j = 0, 1, ..., k-1$.

Since we have more unknowns than equations, it is obvious that there exists a non-zero polynomial w of the form (4.2). Then there exist positive integers r, $1 \le r \le nk+1$, and p, $1 \le p \le r$, and non-zero z_{p+1}, \ldots, z_r such that

$$w(x) = x^{p}(x-z_{p+1}) \cdot \ldots \cdot (x-z_{r}).$$

For $\varepsilon > 0$, define

$$f_{e}(x) = \begin{cases} f(x) + \frac{1}{e}w(x) & \text{if } |x| < \frac{a}{2}, \\ \\ undefined & \text{if } |x| \ge \frac{a}{2}. \end{cases}$$

From the general theory of algebraic functions (see for instance Wilkinson [63]) it is known that for sufficiently small ϵ there exist p simple zeros $\lambda_1(\epsilon), \ldots, \lambda_p(\epsilon)$ such that

(4.5)
$$\lambda_{i}(\epsilon) = \left| \frac{p!f(0)}{w^{(p)}(0)} \right|^{\frac{1}{p}} \epsilon_{i} + O(\epsilon_{i}^{2})$$

where ϵ_i is the i-th complex root of the equation $x^p = \epsilon$. Note that $\lim_{\epsilon \to 0} \lambda_i(\epsilon) = 0$, $\forall i$. Therefore for sufficiently small ϵ , we get $f_{\epsilon} \in \Im$, $\lambda_1(\epsilon), \lambda_2(\epsilon), \dots, \lambda_p(\epsilon) \in S(f_{\epsilon})$ and $\lim_{\epsilon \to 0} R_f(\lambda_i(\epsilon)) = \min\{\frac{a}{2}, L\}$. Then the starting point $y_0 = x_0 = 0$ belongs to $B(c, f_{\epsilon})$. This means that the sequence $\{y_i\}, y_{i+1} = \varphi(y_i; \Re(f_{\epsilon}, y_i))$, is well-defined. Observe that $\Re(f_{\epsilon}, x_j) = \Re(f_{\epsilon}, y_j)$ for $j = 0, 1, \dots, k-1$ and $y_0 = x_0$. Therefore $y_{i+1} = x_{i+1} = \varphi(x_i; \Re(f, x_i))$ for $i = 0, 1, \dots, k-1$. From (4.1) we know that $|y_k| = |x_k| \ge \frac{a}{2}$ which means that y_k does not belong to the domain $D_{f_{\epsilon}}$. Thus $\Re(f_{\epsilon}, y_k)$ is not well-defined, see (2.2), which contradicts $c(\phi) > 0$. Theorem 4.1 says that there exists no globally convergent iteration without memory using a linear information operator for the class \Im of analytic complex functions having only simple zeros. This negative result also holds for real problems. Let \Im_1 be the class of real functions f, f: $D_f \subseteq R \rightarrow R$, analytic in D_f and having an unique simple zero. Consider real information operators, i.e., $\Re(f,x) \in \mathbb{R}^n$, $\forall f \in \Im_1$, $\forall x \in D_f$.

Theorem 4.2

For any real linear information operator \mathfrak{N} there exists no iteration \mathfrak{D} from $\Phi(\mathfrak{N})$ which is globally convergent for the class \mathfrak{Z}_1 .

Proof

Suppose that for a real linear information operator \mathfrak{N} , \mathfrak{N} : $D_{\mathfrak{N}} \rightarrow \mathbb{R}^n$ and for $\varphi \in \mathfrak{H}(\mathfrak{N})$, $c = c(\varphi) > 0$. Let $f(\mathbf{x}) = \mathbf{x}$ -a be defined as in the proof of Theorem 4.1. Since $\mathbf{x}_0 = 0 \in B(c,f)$, the sequence $\{\mathbf{x}_i\}, \mathbf{x}_{i+1} = \varphi(\mathbf{x}_i; \mathfrak{N}(f,\mathbf{x}_i))$ tends to $a \neq 0$. Therefore

(4.6)
$$x_1 = \varphi(0; \Re(f, 0)) \neq 0.$$

Consider now a real polynomial w,

(4.7) w(x) =
$$\sum_{i=0}^{n} a_i x^{2i+1}$$
,

satisfying

(4.8) $\Re(f-w,0) = \Re(f,0)$.

The equation (4.8) is equivalent to the following homogeneous system of n real linear equations.

(4.9)
$$\sum_{i=0}^{n} a_{i}L_{s}(x^{i+1},x_{0})$$
 for $s = 1,2,...,n$.

It is obvious that there exists a non-zero real polynomial w of the form (4.7) and satisfying (4.8). Let $w(x) = x^p (x-z_{p+1}) \cdots (x-z_r)$ where $1 \le r \le 2n+1$, $1 \le p \le r$ and $z_i \ne 0$ for $i = p+1, \ldots, r$. Due to (4.7) p is odd.

Define

$$\mathbf{r} = \begin{cases} \min\{\frac{|\mathbf{x}_1|}{2}, \frac{|\mathbf{z}_{p+1}|}{2}, \dots, \frac{|\mathbf{z}_r|}{2}\} & \text{if } p < r, \\ \frac{|\mathbf{x}_1|}{2} & \text{otherwise,} \end{cases}$$

and

$$f_{\varepsilon}(\mathbf{x}) = \begin{cases} f(\mathbf{x}) + \frac{1}{\varepsilon} \mathbf{w}(\mathbf{x}) & \text{if } |\mathbf{x}| < \Gamma, \\\\ \text{undefined} & \text{if } |\mathbf{x}| \geq \Gamma \end{cases}$$

for $\epsilon > 0$. For sufficiently small ϵ , the solutions of $f_{\epsilon}(x) = 0$ in the complex plane are given by (4.5). Since p is odd, only one of $\lambda_{i}(\epsilon)$ in (4.5) is real. Thus, f_{ϵ} has a unique simple real zero, say $\lambda(\epsilon)$. Since $\lambda(\epsilon)$ tends to zero with ϵ , $\lim_{\epsilon \to 0} R_{f_{\epsilon}}(\lambda(\epsilon)) = \min\{\Gamma, L\}$ and $x_{0} = 0 \in B(c, f_{\epsilon})$ for sufficiently small ϵ . Therefore the sequence $\{x_{i}\}, x_{i+1} = \phi(x_{i}; \Re(f_{\epsilon}, x_{i}))$ is well-defined. Observe that

$$x_1 = \varphi(0; \Re(f_0, 0)) = \varphi(0; \Re(f, 0)).$$

Due to (4.6) $x_1 \notin D_{f_e}$ which means that x_2 is not well-defined. This contradiction ends the proof.

5. THE GENERAL CASE

In Section 4 we showed that there exists no globally convergent onepoint stationary iteration without memory which uses linear information. In this section we prove the same result for multipoint iterations with or without memory.

Let L_1, L_2, \ldots, L_n be functionals defined as in Section 2. Then a <u>multi-</u> point linear information operator $\mathfrak{A}: D_{\mathfrak{A}} \subset \mathfrak{H} \times \mathbb{C} \to \mathbb{C}^n$, is defined as

(5.1) $\mathfrak{N}(\mathbf{f},\mathbf{x}) = [L_1(\mathbf{f},\mathbf{z}_1), L_2(\mathbf{f},\mathbf{z}_2), \dots, L_n(\mathbf{f},\mathbf{z}_n)], \quad \forall \mathbf{f} \in \mathbb{H}, \forall \mathbf{x} \in D_{\mathbf{f}}$ where

(5.2) $z_1 = x$ and $z_{k+1} = \xi_{k+1}(z_1; L_1(f, z_1), L_2(f, z_2), \dots, L_k(f, z_k))$, $k = 1, 2, \dots, n-1$, for certain functions $\xi_2, \xi_3, \dots, \xi_n$. Note that if $\xi_k = x$, $k = 2, \dots, n$, then \Re is a one-point linear information operator defined in Section 2.

For given integer m, let $x_0, x_{-1}, \dots, x_{-m}$ be distinct approximations of a solution of f(x) = 0, $f \in \mathcal{J}$. Suppose we construct a sequence of approximations by the formula

(5.3)
$$x_{i+1} = \varphi(x_i, x_{i-1}, \dots, x_{i-m}; \Re(f, x_i), \Re(f, x_{i-1}), \dots, \Re(f, x_{i-m}))$$

where $\varphi: D_{\varphi} \subset \mathbb{C}^{(m+1)(n+1)} \to \mathbb{C}$ is a functional. Then φ is called a <u>multipoint stationary iteration operator with memory</u> if $m \ge 1$ and <u>without memory</u> if m = 0 using a linear information operator \mathfrak{N} . For brevity φ is also called an <u>iteration</u>. Let $\tilde{\Phi}_m(\mathfrak{N})$ be the class of all such iterations.

For particular \Re and m we get commonly used iterations. For instance, m = 1, $\Re(f,x) = [f(x)]$ and

$$\varphi(\mathbf{x},\mathbf{y}; \ \Re(\mathbf{f},\mathbf{x}), \Re(\mathbf{f},\mathbf{y})) = \mathbf{x} - \frac{\mathbf{x}-\mathbf{y}}{\mathbf{f}(\mathbf{x})-\mathbf{f}(\mathbf{y})}\mathbf{f}(\mathbf{x})$$

is the secant iteration. An example of two-point stationary iteration without memory is provided by Steffensen iteration (see Steffensen [33]) which is defined as follows

$$\Re(f,x) = [f(z_1), f(z_2)]$$
 where $z_1 = x$ and $z_2 = z_1 - f(z_1)$

and

$$\varphi(\mathbf{x}; \ \Re(\mathbf{f}, \mathbf{x})) = \mathbf{z}_1 - \frac{\mathbf{z}_1 - \mathbf{z}_2}{\mathbf{f}(\mathbf{z}_1) - \mathbf{f}(\mathbf{z}_2)} \ \mathbf{f}(\mathbf{z}_1).$$

We extend the definition of global convergence as follows. Let \mathfrak{R} be a one- or multipoint linear information operator and let m be an integer. Recall that B(b,f) is defined in Section 3. For any iteration φ , $\varphi \in \Phi_{\mathfrak{m}}(\mathfrak{R})$, define $c = c(\varphi)$ as a number that

- (i) for any f ∈ ℑ and any choice of distinct starting points
 x₀,x₋₁,...,x_{-m} from B(c,f) the sequence {x_i},
 x_{i+1} = φ(x_i,x_{i-1},...,x_{i-m}; ℜ(f,x_i),ℜ(f,x_{i-1}),...,ℜ(f,x_{i-m}))
 is well-defined and lim x_i ∈ S(f).
- (ii) for any € > 0 there exist f ∈ ℑ and distinct points
 x₀,x₋₁,...,x_{-m} ∈ B(c+e,f) such that either the sequence {x_i} is not well-defined or lim x_i ∉ S(f).

Definition 5.1

We shall say that an iteration ϕ is globally convergent for the class 3 iff

c(m) > 0

Now let Φ be the class of all stationary iterations with or without memory which use a linear information operator \mathfrak{A} , i.e., $\Phi = \bigcup_{\mathfrak{A}} \bigcup_{\mathfrak{m}} \Phi_{\mathfrak{m}}(\mathfrak{A})$.

Theorem 5.1

No iteration φ from $\Phi(\mathfrak{N})$ is globally convergent for the class 3.

Proof

Apply the proof of Theorem 4.1 with starting points $x-j = \frac{j}{4(m+1)} \min\{a,L\}$, $j = 0, 1, \dots, m$, and with n k in (4.2) replaced by n(m+k).

Theorem 5.1 says that knowing only the value of a finite number of linear functionals on f it is impossible to find a globally convergent stationary iteration for the class 3. Therefore if we want to solve f(x) = 0by a stationary iteration we <u>have</u> to assume that the starting points are sufficiently close to a solution. By contrast it is known that for some <u>non-linear</u> information operators there exist globally convergent stationary iterations. An example is provided by Laguerre iteration. Also for linear information operators there exist globally convergent <u>non-stationary</u> iterations for the class 3. An example may be found in Traub and Woźniakowski [76] where global convergence of the sequence of interpolatory iterations I_k is proved. For this case, the k-th iteration requires the knowledge of k linear functionals of f. Furthermore there exist globally convergent non-stationary iterations which are based on the use of increasing size of memory. This will be reported in Wasilkowski [79].

6. FINAL COMMENTS

In Section 5 we showed that no stationary iteration using linear information is globally convergent for the class of analytic problems having simple zeros. The existence of globally convergent iterations depends on the class \Im of functions whose zeros we want to approximate. For some simple classes there exist well known globally convergent stationary iterations which use linear information. For example if \Im is the class of real functions f: $R \rightarrow R$ whose first derivative is monotonic then Newton iteration is globally convergent.

For many interesting classes the existence of global convergent iterations is unknown. Even for the class Π of all real polynomials with simple real zeros this problem is open. All known globally convergent iterations for the class Π are either non-stationary or use nonlinear information. For example Newton iteration with a suitably chosen starting point, Bernoulli's method and Laguerre iteration are globally convergent for Π . See for instance Ralston and Rabinowitz [78]. Either implicitly or explicitly these iterations use the degree k of the polynomial whose zero is desired.

Note that the degree k is nonlinear information. Indeed, suppose there exists a function g (in general nonlinear) such that $k = g(x; \Re(f,x))$ where \Re is a multipoint <u>linear</u> information operator. For any x_0 there exists a polynomial w of degree greater than one such that $\Re(w,x_0) = 0$. Taking $f_{\varepsilon}(x) = x + \frac{1}{\varepsilon} w(x)$ we get $\Re(f_{\varepsilon},x_0) = \Re(x,x_0)$ and $f_{\varepsilon} \in \Pi$ for sufficiently small ε . Therefore $g(x_0,\Re(x,x_0)) = g(x_0,\Re(f_{\varepsilon},x_0))$, but the degrees of the polynomials x and $f_c(x)$ are different. This contradicts the assumption $k = g(x; \Re(f,x))$, with \Re linear.

We believe that any globally convergent iteration for Π has to be nonstationary or use nonlinear information. Therefore we propose the following conjecture.

Conjecture 6.1

where the statement of the

There exists no globally convergent stationary iteration using linear information for the class of all real polynomials with simple real zeros.

ACKNOWLEDGMENT

I thank J. F. Traub and H. Woźniakowski for their help. Without it this paper could not have been completed.

A.1

REFERENCES

Ralston and Rabinowitz [78] Ralston, A. and P. Rabinowitz, A First Course in Numerical Analysis, Second Edition, McGraw-Hill Book Co., New York, 1978. Steffensen [33] Steffensen, J. F., "Remarks on Iteration," Skand. Aktuar Tidskr, 16, 1933, 64-72. Traub and Woźniakowski [76] Traub, J. F. and H. Wozniakowski, "Optimal Radius of Convergence of Interpolatory Iterations for Operator Equations," Department of Computer Science Report, Carnegie-Mellon University, 1976. To appear in Aequationes Mathematicae. Traub and Woźniakowski [78] Traub, J. F. and H. Woźniakowski, 'General Theory of Optimal Error Algorithms and Analytic Complexity, Part B: Iterative Information Model," Department of Computer Science Report, Carnegie-Mellon University, 1978. To appear in <u>A General Theory of</u> Optimal Error Algorithms, Academic Press,

New York, 1979.

Wasilkowski [79]

Wasilkowski, G. W., "Global Convergence of a Sequence of Stationary Iterations Using Linear Information," in progress.

Wilkinson [63]

Wilkinson, J. H., <u>Rounding Errors in Alge-</u> braic Processes, Prentice-Hall, Inc., 1963.

UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered) READ INSTRUCTIONS BEFORE COMPLETING FORM REPORT DOCUMENTATION PAGE PORT NUMBER 2. GOVT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER CMU-CS-78-155 TITLE (and Subilitie) S. TYPE OF REPORT & PERIOD COVERED Interim Phi CAN ANY STATIONARY ITERATION USING LINEAR INFORMATION BE GLOBALLY CONVERGENT? 6. PERFORMING AUTHOR() S. CONTRACT OR GRANT NUMBER(... 10 G.W. Wasilkowski N00014-76-C-0370 PROGRAM ELEMENT. PROJECT, TASK . PERFORMING ORGANIZATION NAME AND ADDRESS Carnegie-Mellon University Computer Science Dept. Pittsburgh, PA 15213 11. CONTROLLING OFFICE NAME AND ADDRESS RT DATI December 1978 Office of Naval Research Arlington, VA 22217 14. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office) 15. SECURITY CLASS. (of this report) Same as above UNCLASSIFIED 184. DECLASSIFICATION/ DOWNGRADING 16. DISTRIBUTION STATEMENT (of this Report) DISTRIBUTION STATEMENT ? Approved for public selectory · Distribution Unlimited 17. DISTRIBUTION STATEMENT (of the obstract entered in Black 20, it dillerent from Report) NOOD14-76-2-03705NSF-MES-75-223 55 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) 20. ABSTRACT (Continue on reverse side if necessary and identity by block number) DD 1 JAN 73 1473 EDITION OF I NOV 65 IS OBSOLETE UNCLASSIFIED S/N 0102-014-6601 | SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

