

AD-A066 268

FOREIGN TECHNOLOGY DIV WRIGHT-PATTERSON AFB OHIO
POWERFUL LASERS ON SOLUTIONS OF DYES, (U)
MAR 78 M I DZYUBENKO, A M KOROBV

F/G 20/5

UNCLASSIFIED

FTD-ID(RS)T-0297-78

NL

| OF |
ADA
066268



END
DATE
FILMED

5-79
DDC



AD-A066268


FOREIGN TECHNOLOGY DIVISION



POWERFUL LASERS ON SOLUTIONS OF DYES

by

M. I. Dzyubenko, A. M. Korobov, I. G. Naumenko



Approved for public release;
distribution unlimited.

78 12 22 394

ACCESSION NR	
RT13	White Section <input checked="" type="checkbox"/>
DD?	Ref Section <input type="checkbox"/>
ANNOUNCED	<input type="checkbox"/>
INSTRUCTIONS	
BY	
DISTRIBUTION/AVAILABILITY CODES	
SPECIAL	
A	

EDITED TRANSLATION

FTD-ID(RS)T-0297-78

9 March 1978

MICROFICHE NR: *FTD-78-C-000318*

POWERFUL LASERS ON SOLUTIONS OF DYES

By: M. I. Dzyubenko, A. M. Korobov, I. G. Naumenko

English pages: 2

Source: Ukrainskiy Fizicheskiy Zhurnal, Vol. 14,
No. 11, Nov 1969, pp. 1929.

Country of origin: USSR

Translated by: Charles T. Ostertag, Jr.

Requester: FTD/TQTD

Approved for public release; distribution unlimited.

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-AFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
А а	А а	A, a	Р р	Р р	R, r
Б б	Б б	B, b	С с	С с	S, s
В в	В в	V, v	Т т	Т т	T, t
Г г	Г г	G, g	У у	У у	U, u
Д д	Д д	D, d	Ф ф	Ф ф	F, f
Е е	Е е	Ye, ye; E, e*	Х х	Х х	Kh, kh
Ж ж	Ж ж	Zh, zh	Ц ц	Ц ц	Ts, ts
З э	З э	Z, z	Ч ч	Ч ч	Ch, ch
И и	И и	I, i	Ш ш	Ш ш	Sh, sh
Й й	Й й	Y, y	Щ щ	Щ щ	Shch, shch
К к	К к	K, k	Ъ ъ	Ъ ъ	"
Л л	Л л	L, l	Ы ы	Ы ы	Y, y
М м	М м	M, m	Ь ь	Ь ь	'
Н н	Н н	N, n	Э э	Э э	E, e
О о	О о	O, o	Ю ю	Ю ю	Yu, yu
П п	П п	P, p	Я я	Я я	Ya, ya

*ye initially, after vowels, and after ъ, ь; e elsewhere.
When written as ë in Russian, transliterate as yë or ë.

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	sinh ⁻¹
cos	cos	ch	cosh	arc ch	cosh ⁻¹
tg	tan	th	tanh	arc th	tanh ⁻¹
ctg	cot	cth	coth	arc cth	coth ⁻¹
sec	sec	sch	sech	arc sch	sech ⁻¹
cosec	csc	csch	csch	arc csch	csch ⁻¹

Russian English

rot curl
lg log

FIRST LINE OF TEXT

POWERFUL LASERS ON SOLUTIONS OF DYES

Dzyubenko, M. I., Korobov, A. M., Naumenko, I. G.

The authors of a number of works [1, 2], in noting the advantages of lasers on solutions of organic dyes, indicated the possibility of obtaining output energies which are comparable with the energy of radiation of solid-state lasers. However, the energies and powers which have been attained experimentally up to the present time are considerably less than in solid-state lasers. Thus the output energy of generation of an alcohol solution of rhodamine 6G in work [3] comprised 0.05-0.07 J (power 7-10 kW), and in [4] - 0.07-0.1 J (power 250-300 kW). Using flash lamps of a special design for the excitation of dyes, the authors of [5] were able to obtain an output energy of ~0.2 J (power ~1 MW) on rhodamine 6G, and ~0.001 J on sodium fluorescein.

In this work we are reporting on some of the energy characteristics of lasers on aqueous and alcohol solutions of the following organic compounds: rhodamine S [C], rhodamine 6G [Ж], sodium fluorescein, and 4-methyl-7-hydroxycoumarin. These compounds yielded the required radiation in the red, yellow, green and blue areas of the spectrum. The solutions were poured into quartz cuvettes with an inner diameter of 5 and 10.5 mm and length of 160 and 280 mm respectively.

Direct xenon lamps were used for excitation. The least threshold energy was observed in the case of rhodamine 6G, dissolved in alcohol, and comprised 6-7 J at a concentration of 5.0×10^{-5} mole/l in a cuvette 160 mm long. Let us also note that in the case of an excitation energy of an order of 100 J generation was observed in

the resonator, formed by the ends of the cuvette (reflection coefficient of an end ~4%).

After selection of the optimal feedback the output energy of the laser on a solution of rhodamine 6G was ~0.7 J (concentration 2.0×10^{-4} mole/L, excitation energy 540 J, length of cuvette 160 mm). In a cuvette 280 mm in length at the same excitation energy the energy of radiation was ~1.2 J, which corresponds to ~2.5 MW and a conversion efficiency of ~0.22%. The maximum output energy of the laser on a solution of sodium fluorescein was ~0.03 J, and on a solution of 4-methyl-7-hydroxycoumarin - ~0.1 J.

The output energy of a laser on aqueous solutions of xanthene dyes was considerably lower, and the threshold of generation was higher than for a laser on alcohol solutions.

Thus our experimental results show that lasers on solutions of organic dyes can be serious competitors of solid-state lasers.

Institute of Radiophysics
and Electronics
AS USSR
Kharkov

Submitted 30 Jul 1969

BIBLIOGRAPHY

1. D. L. Stockman, W. R. Mallory, Proc. IEEE, 52, № 3, 341, 1964.
2. Л. Н. Темная, Обзоры научно-технической литературы по электронной технике, сер. «Электровакuumные приборы», № 25, 1967.
3. А. Н. Рубинов, В. А. Мостовников, ДАН БССР, 12, № 7, 602, 1968.
4. М. И. Дзюбенко, И. Г. Науменко, А. М. Коробов, УФЖ, 14, № 4, 1969.
5. P. P. Sorokin, J. R. Lankard, V. L. Moruzzi, E. C. Hammond, J. Chem. Phys., 48, № 10, 4726, 1968.

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

ORGANIZATION	MICROFICHE	ORGANIZATION	MICROFICHE
A205 DMATC	1	E053 AF/INAKA	1
A210 DMAAC	2	E017 AF/RDXTR-W	1
B344 DIA/RDS-3C	8	E404 AEDC	1
C043 USAMIIA	1	E408 AFWL	1
C509 BALLISTIC RES LABS	1	E410 ADTC	1
C510 AIR MOBILITY R&D LAB/FIO	1	E413 ESD	2
C513 PICATINNY ARSENAL	1	FTD	
C535 AVIATION SYS COMD	1	CCN	1
C591 FSTC	5	TQIS	3
C619 MIA REDSTONE	1	NIA/PHS	1
D008 NISC	1	NICD	2
H300 USAICE (USAREUR)	1		
P005 ERDA	1		
P055 CIA/CRS/ADD/SD	1		
NAVORDSTA (50L)	1		
NASA/KSI	1		
AFIT/LD	1		

FTD-ID (RS) T-0297-78