

AD-A066 588

NAVAL SURFACE WEAPONS CENTER DAHLGREN LAB VA
THE INFLUENCE OF DISTANT ZONES ON STOKES' EQUATION CONSIDERING --ETC(U)
NOV 78 P J FELL

F/G 8/5

UNCLASSIFIED


NSWC/DL-TR-3911

NL

| OF |
AD
A066588



END
DATE
FILMED
'5--79
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A0 66588

DDC FILE COPY

12
LEVEL II

DDC
RECORDED
MAR 30 1979
RECEIVED
X E

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

14) REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NSWC/DL-TR-3911	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) THE INFLUENCE OF DISTANT ZONES ON STOKES' EQUATION CONSIDERING THE REMOVAL OF LOWER-DEGREE HARMONICS FROM S(ψ) OR Δg .	5. TYPE OF REPORT & PERIOD COVERED Final rept.	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Patrick J./Fell	8. CONTRACT OR GRANT NUMBER(s)	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Surface Weapons Center (K12) Dahlgren Laboratory Dahlgren, VA 22448	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 11221N:J0094:J0094000	
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Strategic Systems Project Office Washington, DC 20376	12. REPORT DATE November 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12) 24p.	13. NUMBER OF PAGES 22	15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Lower-Degree Harmonics Stokes' Equation Geoid Undulation Cap Radii		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The expected error in the computation of geoid undulations using Stokes' equation due to neglected distant zones is analyzed considering the removal of lower-degree harmonics from limited gravity anomaly data and from Stokes' kernel. Evaluation of the derived error equations assuming Kaula's rule for anomaly degree variances indicates that if the lower-degree components of geoid undulation are assumed known, removal of an equivalent number of such harmonics from the anomaly data alone produces the least-expected error for cap radii of less than 60 deg.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

391 598

19 03

25 036

LB

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

[A large rectangular box with a black border, containing faint, illegible text and a small white checkmark in the upper left quadrant.]


SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

FOREWORD

The work reported herein was conducted by the Ocean Geodesy Branch of the Naval Surface Weapons Center under the sponsorship of the Strategic Systems Project Office (Project Number J0094).

This report has been reviewed and approved by P. Ugincius, Head, Ocean Geodesy Branch and R. J. Anderle, Head, Astronautics and Geodesy Division.

Released by



R. A. NIEMANN, Head
Strategic Systems Department

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION _____	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	SECRET
A	

CONTENTS

	Page
INTRODUCTION	1
INFLUENCE OF DISTANT ZONES ON STOKES' EQUATION	1
CASE I: STANDARD CASE	1
CASE II: REMOVAL OF M LOWER-DEGREE HARMONICS FROM $S(\psi)$	3
CASE III: REMOVAL OF M LOWER-DEGREE HARMONICS FROM Δg	5
CASE IV: REMOVAL OF M LOWER-DEGREE HARMONICS FROM BOTH $S(\psi)$ AND Δg	7
NUMERICAL EVALUATION OF ERROR FORMULAE	10
CONCLUSIONS	12
REFERENCES	14
DISTRIBUTION	

INTRODUCTION

Stokes' equation gives the mathematical relationship between the geometric shape of the geoid and the magnitude of gravity on the geoid. The equation is

$$N = \frac{R}{4\pi G} \iint S(\psi) \Delta g \, d\sigma \quad (1)$$

where

R = mean radius of the earth

G = mean value of gravity over the earth

$S(\psi)$ = Stokes' kernel

The integration, extended over the unit sphere using a global set of gravity anomalies Δg , produces the geoid-ellipsoid separation at a given point.

In practice, the integration is not extended over the entire globe, but only up to a spherical distance ψ_0 from the computation point due to a lack of sufficient surface gravity data. The error δN that results from neglecting data beyond ψ_0 has been analyzed by Molodenskii (Reference 1) and is described in Moritz (Reference 2). The magnitude of this error may be reduced if the gravitational potential is known in terms of a finite spherical harmonic expansion derived from satellite observations by applying, for instance, the computational technique of Rummel and Rapp (Reference 3). Equivalently, the error may be reduced by removing known lower-degree harmonics from the gravity anomaly data used when Equation (1) is applied to a spherical cap assuming the lower-order harmonic components of N are known. In addition, one might also consider the possibility of modifying or replacing Stokes' kernel $S(\psi)$ with a function which produces a better estimate of N using a limited data set. The purpose of this report is to determine what decrease in δN may be expected if harmonics are removed from $S(\psi)$ or Δg individually, or from both simultaneously.

INFLUENCE OF DISTANT ZONES ON STOKES' EQUATION

CASE I: STANDARD CASE

The error in the geoid undulation due to neglecting gravity anomaly data outside a spherical cap σ of radius ψ_0 is given by Moritz (Reference 2) as

$$\delta N_I = \frac{R}{4\pi G} \int_{\psi=\psi_0}^{\pi} \int_{\alpha=0}^{2\pi} S(\psi) \Delta g \sin \psi \, d\psi \, d\alpha \quad (2)$$

where the subscript I refers to the removal of harmonics from the data only (Table 1). Defining a new function $\bar{S}(\psi)$ as

$$\bar{S}(\psi) = \begin{cases} 0 & 0 \leq \psi < \psi_0 \\ S(\psi) & \psi_0 \leq \psi \leq \pi \end{cases} \quad (3)$$

Equation (2) becomes

$$\delta N_1 = \frac{R}{4\pi G} \int_{\psi=0}^{\pi} \int_{\alpha=0}^{2\pi} \bar{S}(\psi) \Delta g \sin \psi \, d\psi \, d\alpha \quad (4)$$

The function $\bar{S}(\psi)$ has the following spherical harmonic expansion:

$$\bar{S}(\psi) = \sum_{n=0}^{\infty} \frac{2n+1}{2} Q_n P_n(\cos \psi) \quad (5)$$

where

$$Q_n = \int_0^{\pi} \bar{S}(\psi) P_n(\cos \psi) \sin \psi \, d\psi$$

(6)

or

$$= \int_{\psi_0}^{\pi} S(\psi) P_n(\cos \psi) \sin \psi \, d\psi$$

Following Moritz (Reference 2), the RMS error due to neglecting distant zones is the square root of

$$\overline{\delta N^2} = \frac{R^2}{4G^2} \sum_{n=2}^{\infty} Q_n^2 c_n \quad (7)$$

where it is assumed the zero and first-order harmonics of the gravity anomaly field are zero.

$$\Delta g(\theta, \lambda) = \sum_{n=2}^{\infty} \Delta g_n(\theta, \lambda) \quad (8)$$

The quantity c_n is the anomaly degree variance for degree n defined by

$$c_n = E \left\{ \Delta g_n^2 \right\} \quad (9)$$

where E is the expected value operator. The anomaly degree variance is a function of the spherical harmonic coefficients of degree n

$$c_n = \sum_{m=0}^n (\bar{a}_{nm}^2 + \bar{b}_{nm}^2) \quad (10)$$

where the bar indicates normalized coefficients. The coefficients Q_n are known as the Molodenskii coefficients, and much is known of their properties.

CASE II: REMOVAL OF M LOWER-DEGREE HARMONICS FROM S(ψ)

Equation (1) may be written in the following form:

$$N = \frac{R}{4\pi G} \int_{\psi=0}^{\psi_0} \int_{\alpha=0}^{2\pi} S(\psi) \Delta g \sin \psi \, d\psi \, d\alpha$$

$$+ \frac{R}{4\pi G} \int_{\psi=\psi_0}^{\pi} \int_{\alpha=0}^{2\pi} S(\psi) \Delta g \sin \psi \, d\psi \, d\alpha$$
(11)

The kernel $S(\psi)$ has the spherical harmonic expansion

$$S(\psi) = \sum_{n=2}^{\infty} \frac{2n+1}{n-1} P_n(\cos \psi)$$
(12)

Now define

$$\tilde{S}(\psi) = S(\psi) - \sum_{n=2}^m \frac{2n+1}{n-1} P_n(\cos \psi)$$
(13a)

or

$$S(\psi) = \tilde{S}(\psi) + S_m(\psi)$$
(13b)

Substituting Equation (13b) into Equation (11) gives

$$N = I_1 + I_2 + I_3$$
(14a)

where

$$I_1 = \frac{R}{4\pi G} \int_{\psi=0}^{\psi_0} \int_{\alpha=0}^{2\pi} \tilde{S}(\psi) \Delta g \, d\sigma$$
(14b)

$$I_2 = \frac{R}{4\pi G} \int_{\psi=\psi_0}^{\pi} \int_{\alpha=0}^{2\pi} \tilde{S}(\psi) \Delta g \, d\sigma$$
(14c)

$$I_3 = \frac{R}{4\pi G} \int_{\psi=0}^{\pi} \int_{\alpha=0}^{2\pi} S_m(\psi) \Delta g \, d\sigma$$
(14d)

Considering the definition of $S_m(\psi)$ and Equation (8), the integral I_3 becomes

$$I_3 = N_2 + N_3 + \dots + N_m \quad (15)$$

which are the first m harmonic components of N at the computation point. Assuming I_3 to be known, the error due to integration over a limited cap becomes

$$\delta N_{II} = \frac{R}{4\pi G} \int_{\psi=\psi_0}^{\pi} \int_{\alpha=0}^{2\pi} \tilde{S}(\psi) \Delta g \, d\sigma \quad (16)$$

Define the function \tilde{S} as

$$\tilde{S}(\psi) = \begin{cases} 0 & 0 \leq \psi < \psi_0 \\ \tilde{S}(\psi) & \psi_0 \leq \psi \leq \pi \end{cases} = \begin{cases} 0 & 0 \leq \psi < \psi_0 \\ S(\psi) - S_m(\psi) & \psi_0 \leq \psi \leq \pi \end{cases} \quad (17)$$

The function \tilde{S} has the Legendre series expansion

$$\tilde{S}(\psi) = \sum_{n=0}^{\infty} \frac{2n+1}{2} \tilde{Q}_n P_n(\cos \psi) \quad (18)$$

where

$$\begin{aligned} \tilde{Q}_n &= \int_0^{\pi} \tilde{S}(\psi) P_n(\cos \psi) \sin \psi \, d\psi \\ &= \int_{\psi_0}^{\pi} S(\psi) P_n(\cos \psi) \sin \psi \, d\psi - \int_{\psi_0}^{\pi} S_m(\psi) P_n(\cos \psi) \sin \psi \, d\psi \\ &= Q_n - \int_{\psi_0}^{\pi} S_m(\psi) P_n(\cos \psi) \sin \psi \, d\psi \end{aligned} \quad (19)$$

Therefore,

$$\begin{aligned} \delta N_{II} &= \frac{R}{4\pi G} \int_{\psi_0}^{\pi} \int_0^{2\pi} \tilde{S}(\psi) \Delta g \, d\sigma \\ &= \frac{R}{4\pi G} \int_0^{\pi} \int_0^{2\pi} \tilde{S}(\psi) \Delta g \, d\sigma \end{aligned}$$

$$\begin{aligned}
&= \frac{R}{8\pi G} \sum_{n=0}^{\infty} (2n+1) \tilde{Q}_n \int_0^{\pi} \int_0^{2\pi} \Delta g P_n(\cos \psi) \sin \psi \, d\psi \, d\alpha \\
&= \frac{R}{8\pi G} \sum_{n=0}^{\infty} (2n+1) \tilde{Q}_n \frac{4\pi \Delta g_n}{(2n+1)} \\
&= \frac{R}{2G} \sum_{n=0}^{\infty} \tilde{Q}_n \Delta g_n \tag{20}
\end{aligned}$$

The RMS error is, following Moritz (Reference 2),

$$\overline{\delta N_{II}^2} = \frac{R^2}{4G^2} \sum_{n=2}^{\infty} \tilde{Q}_n^2 c_n \tag{21}$$

CASE III: REMOVAL OF M LOWER-DEGREE HARMONICS FROM Δg

From Equation (8),

$$\Delta g = \sum_{n=2}^{\infty} \Delta g_n(\theta, \lambda)$$

where the n'th degree harmonic Δg_n is given by

$$\Delta g_n = \sum_{m=0}^n (\bar{a}_{nm} \bar{R}_{nm}(\theta, \lambda) + \bar{b}_{nm} \bar{S}_{nm}(\theta, \lambda)) \tag{22}$$

Removing the first m harmonics from the data set gives

$$\tilde{\Delta g} = \Delta g - \sum_{n=2}^m \Delta g_n \tag{23}$$

Equation (1) becomes

$$\begin{aligned}
N &= \frac{R}{4\pi G} \int_0^{\pi} \int_0^{2\pi} S(\psi) [\tilde{\Delta g} + \sum_{n=2}^m \Delta g_n] \sin \psi \, d\psi \, d\alpha \\
&= I_1 + I_2 \tag{24a}
\end{aligned}$$

where

$$I_1 = \frac{R}{4\pi G} \int_0^\pi \int_0^{2\pi} S(\psi) \tilde{\Delta g} \sin \psi \, d\psi \, d\alpha \quad (24b)$$

$$I_2 = \frac{R}{4\pi G} \sum_{n=2}^m \int_0^\pi \int_0^{2\pi} S(\psi) \Delta g_n \sin \psi \, d\psi \, d\alpha \quad (24c)$$

The second integral reduces to

$$I_2 = N_2 + \dots + N_m \quad (25)$$

which is assumed known since $\Delta g_2, \dots, \Delta g_m$ are assumed given. Therefore,

$$\begin{aligned} N = N_2 + \dots + N_m + \frac{R}{4\pi G} \int_0^{\psi_0} \int_0^{2\pi} S(\psi) \tilde{\Delta g} \, d\sigma \\ + \frac{R}{4\pi G} \int_0^\pi \int_0^{2\pi} S(\psi) \tilde{\Delta g} \, d\sigma \end{aligned} \quad (26)$$

The error due to distant zones is the second integral in Equation (26).

$$\begin{aligned} \delta N_{III} &= \frac{R}{4\pi G} \int_{\psi_0}^\pi \int_0^{2\pi} S(\psi) \tilde{\Delta g} \, d\sigma \\ &= \frac{R}{4\pi G} \int_0^\pi \int_0^{2\pi} \bar{S}(\psi) \tilde{\Delta g} \, d\sigma \\ &= \frac{R}{8\pi G} \sum_{n=0}^{\infty} (2n+1) Q_n \int_0^\pi \int_0^{2\pi} \tilde{\Delta g} P_n(\cos \psi) \sin \psi \, d\psi \, d\alpha \end{aligned} \quad (27)$$

For $n = 0, 1, \dots, m$

$$\int_0^\pi \int_0^{2\pi} \tilde{\Delta g} P_n(\cos \psi) \sin \psi \, d\psi \, d\alpha = 0 \quad (28)$$

since $\tilde{\Delta g}$ contains no harmonics of these degrees. Therefore,

$$\delta N_{III} = \frac{R}{2G} \sum_{n=m+1}^{\infty} Q_n \Delta g_n \quad (29)$$

and

$$\overline{\delta N_{III}^2} = \frac{R^2}{4G^2} \sum_{n=m+1}^{\infty} Q_n^2 c_n \quad (30)$$

It should be noted that this error is the same as that obtained using the Rummel and Rapp (Reference 3) procedure with the first m harmonics applied (neglecting here corrections for atmospheric effects and zero-order undulation correction).

CASE IV: REMOVAL OF M LOWER-DEGREE HARMONICS FROM BOTH $S(\psi)$ AND Δg

Using equations (13b) and (23),

$$S = \tilde{S} + S_m \quad (13b)$$

$$\Delta g = \tilde{\Delta g} + \sum_{n=2}^m \Delta g_n \quad (23)$$

Equation (1) becomes

$$N = I_1 + I_2 + I_3 + I_4 \quad (31a)$$

where

$$I_1 = \frac{R}{4\pi G} \int_0^{\pi} \int_0^{2\pi} \tilde{S}(\psi) \tilde{\Delta g} \, d\sigma \quad (31b)$$

$$I_2 = \frac{R}{4\pi G} \sum_{n=2}^m \int_0^{\pi} \int_0^{2\pi} \tilde{S}(\psi) \Delta g_n \, d\sigma \quad (31c)$$

$$I_3 = \frac{R}{4\pi G} \int_0^{\pi} \int_0^{2\pi} S_m(\psi) \tilde{\Delta g} \, d\sigma \quad (31d)$$

$$I_4 = \frac{R}{4\pi G} \sum_{n=2}^m \int_0^\pi \int_0^{2\pi} S_m(\psi) \Delta g_n d\sigma \quad (31e)$$

By orthogonality, I_2 and I_3 vanish. Also, by definition of S_m (Equation 13a),

$$I_4 = \frac{R}{4\pi G} \sum_{n=2}^m \sum_{j=2}^m \frac{2j+1}{j-1} \int_0^\pi \int_0^{2\pi} P_j(\cos \psi) \Delta g_n d\sigma$$

If $n \neq j$, the partial sum is zero by orthogonality. When $n = j$, the nonvanishing terms sum to

$$I_4 = \frac{R}{4\pi G} \sum_{n=2}^m \frac{2n+1}{n-1} \Delta g_n \frac{4\pi}{2n+1} \quad (32)$$

since

$$\int_0^\pi \int_0^{2\pi} P_n(\cos \psi) \Delta g_n d\sigma = \frac{4\pi \Delta g_n}{2n+1}$$

Therefore,

$$I_4 = N_2 + \dots + N_m \quad (33)$$

Thus, Equation (31a) becomes

$$\begin{aligned} N = N_2 + \dots + N_m + \frac{R}{4\pi G} \int_{\psi=0}^{\psi_0} \int_{\alpha=0}^{2\pi} \tilde{S}(\psi) \tilde{\Delta g} d\sigma \\ + \frac{R}{4\pi G} \int_{\psi=\psi_0}^{\pi} \int_{\alpha=0}^{2\pi} \tilde{S}(\psi) \tilde{\Delta g} d\sigma \end{aligned} \quad (34)$$

Assuming the first m harmonics of N are known, the error in N due to neglecting distant zones is

$$\delta N_{IV} = \frac{R}{4\pi G} \int_{\psi_0}^{\pi} \int_0^{2\pi} \tilde{S}(\psi) \tilde{\Delta g} d\sigma$$

$$\begin{aligned}
&= \frac{R}{4\pi G} \int_0^\pi \int_0^{2\pi} \tilde{S}(\psi) \tilde{\Delta g} \, d\sigma \\
&= \frac{R}{8\pi G} \sum_{n=0}^{\infty} (2n+1) \tilde{Q}_n \int_{\psi=0}^{\pi} \int_{\alpha=0}^{2\pi} \tilde{\Delta g} P_n(\cos \psi) \, d\sigma
\end{aligned} \tag{35}$$

For $n = 0, 1, \dots, m$ the integral in Equation (35) vanishes giving

$$\delta N_{IV} = \frac{R}{2G} \sum_{n=m+1}^{\infty} \tilde{Q}_n \Delta g_n \tag{36}$$

The RMS error is the square root of

$$\overline{\delta N_{IV}^2} = \frac{R^2}{4G^2} \sum_{n=m+1}^{\infty} \tilde{Q}_n^2 c_n \tag{37}$$

Tables 1 and 2 present summaries of Cases I through IV and of RMS error formulae, respectively.

Table 1. Summary of Cases I through IV

Case	Equation for Undulation
I	$N = \frac{R}{4\pi G} \iint_{\sigma} S(\psi) \Delta g \, d\sigma + \delta N_I$
II	$N = N_2 + \dots + N_m + \frac{R}{4\pi G} \iint_{\sigma} \tilde{S}(\psi) \Delta g \, d\sigma + \delta N_{II}$
III	$N = N_2 + \dots + N_m + \frac{R}{4\pi G} \iint_{\sigma} S(\psi) \tilde{\Delta g} \, d\sigma + \delta N_{III}$
IV	$N = N_2 + \dots + N_m + \frac{R}{4\pi G} \iint_{\sigma} \tilde{S}(\psi) \tilde{\Delta g} \, d\sigma + \delta N_{IV}$

Table 2. Summary of RMS Error Formulae

<u>Case</u>	<u>M Harmonics Removed</u>	<u>Square of RMS Error</u>
I	-	$\frac{R^2}{4G^2} \sum_{n=2}^{\infty} Q_n^2 c_n$
II	$S(\psi)$	$\frac{R^2}{4G^2} \sum_{n=2}^{\infty} \tilde{Q}_n^2 c_n$
III	Δg	$\frac{R^2}{4G^2} \sum_{n=m+1}^{\infty} Q_n^2 c_n$
IV	$S(\psi), \Delta g$	$\frac{R^2}{4G^2} \sum_{n=m+1}^{\infty} \tilde{Q}_n^2 c_n$

NUMERICAL EVALUATION OF ERROR FORMULAE

The RMS error for each case was computed for m equal to 6, 12, and 18. In these computations the anomaly degree variances were computed using Kaula's rule

$$c_n = \frac{192}{n + 1.5} \text{ mgals}^2 \quad n = 3, 4, \dots \quad (38)$$

and c_2 was taken as 10 mgal^2 . The constant R^2/G^2 was taken as $42.3 \text{ m}^2/\text{mgal}^2$.

The Molodenskii (Reference 1) coefficients Q_n were computed using two algorithms. Initially, the Q_n were computed using an algorithm of Hagiwara (Reference 4); however, due to numerical difficulties, only the first 50 terms in the error expressions could be summed. The computations were repeated using the algorithm of Paul (Reference 5). An increased number of terms (250) were summed in this case.

The coefficients \tilde{Q}_n were computed using the following procedure:

$$\tilde{Q}_n = Q_n - \int_{\psi_0}^{\pi} S_m P_n(\cos \psi) \sin \psi \, d\psi \quad (39)$$

$$S_m = \sum_{j=2}^m \frac{2j + 1}{j - 1} P_j(\cos \psi) \quad (40)$$

$$\tilde{Q}_n = Q_n - \sum_{j=2}^m \frac{2j+1}{j-1} \int_{\psi_0}^{\pi} P_j(\cos \psi) P_n(\cos \psi) \sin \psi d\psi \quad (41)$$

where $Q_n = Q_n(\psi_0)$
 $\tilde{Q}_n = \tilde{Q}_n(\psi_0)$

Taking $x = \cos \psi$

$$\begin{aligned} & \int_{\psi_0}^{\pi} P_j(\cos \psi) P_n(\cos \psi) \sin \psi d\psi \\ &= \int_{-1}^{\cos \psi_0} P_j(x) P_n(x) dx, \end{aligned} \quad (42)$$

and using the identities

$$\begin{aligned} \int_a^b P_n^2(x) dx &= \frac{1}{2n+1} \left\{ (2n-1) \int_a^b P_{n-1}^2(x) dx \right. \\ & \left. + x [P_n^2(x) + P_{n-1}^2(x)] - 2P_{n-1} P_n \right\} \Bigg|_a^b \end{aligned} \quad (43)$$

$$\int_a^b P_n(x) P_k(x) dx = \frac{[x(n-k) P_n(x) P_k(x) + k P_n(x) P_{k-1}(x) - n P_{n-1}(x) P_k(x)]}{(n-k)(n+k+1)} \Bigg|_a^b$$

$n \neq k$ (44)

Equation (41) may be evaluated.

CONCLUSIONS

Figures 1 through 4 give the root mean square error for Cases I through IV where 250 terms have been summed in the error equations. In Cases II through IV, the first m harmonic components of the geoid undulation N were assumed known. An examination of these figures shows that for spherical caps of radii $\psi_0 < 60^\circ$, removal of known lower-degree harmonics from the cap data alone produces the least-expected error due to neglected zones. Removal of lower-degree harmonic components from the kernel function $S(\psi)$ does not minimize the expected error for cap sizes normally considered. Thus, the former procedure (Case III) is recommended if the assumptions, upon which these error formulae are based, are satisfied.

Finally, a comparison of the results based on 50 and 250 terms in the summation demonstrated that satisfactory convergence of the series had been obtained.




Figure 1. Root Mean Square Error, Case I




Figure 2. Root Mean Square Error,
Cases II Through IV, $m = 6$




Figure 3. Root Mean Square Error,
Cases II Through IV, $m = 12$




Figure 4. Root Mean Square Error,
Cases II through IV, $m = 18$

REFERENCES

1. M. S. Molodenskii, V. F. Eremeev, and M. I. Yurkina: *Methods for Study of the External Gravitational Field and Figure of the Earth*, TT61-31207 National Technical Information Service, Springfield, VA, 1962.
2. W. Heiskanen and H. Moritz, *Physical Geodesy*, W. H. Freeman, San Francisco, CA, 1967.
3. R. Rummel and R. H. Rapp, *The Influence of the Atmosphere on Geoid and Potential Coefficient Determinations from Gravity Data*, JGR Vol. 81, No. 32, 1976.
4. Y. Hagiwara, "Truncation Error Formulas for Geoidal Height and the Deflection of the Vertical," *Bulliten Geodesique*, 106, 453, 1973.
5. M. K. Paul, "A Method of Evaluating the Truncation Error Coefficients for Geoidal Height," *Bulliten Geodesique*, 110, 413, 1973.

DISTRIBUTION

Defense Documentation Center
Cameron Station
Alexandria, VA 22314 (12)

Library of Congress
Washington, DC 20540
Attn: Gift and Exchange Division (4)

Director Defense Mapping Agency
U.S. Naval Observatory
Bldg. 56
Washington, DC 20360
Attn: O. W. Williams
C. Martin
P. M. Schwimmer
D. A. Rekenhaller

Commander
Naval Oceanographic Office
NSTL Station
Bay St. Louis, MS 39522
Attn: T. Davis
J. Hankins

Office of the Oceanographer
Hoffman 2
200 Stovall Street
Alexandria, VA 22332
Attn: CAPT D. Brown

Oceanographer of the Navy
Hoffman 2
200 Stovall St.
Alexandria, VA 22332

Defense Mapping Agency
Aerospace Center
St. Louis, MO 63118
Attn: R. Ballew
J. Finkland (Code GDG) (2)
M. Shultz (Code GDGA) (4)

Defense Mapping Agency
Hydrographic/Topographic Center
Washington, DC 20315
Attn: R. Smith (3)

Air Force Geophysical Laboratory
L. G. Hanscom Field
Bedford, MA 01730
Attn: G. Hadgigeorge

U.S. Army Topographic Laboratory
Ft. Belvoir, VA 22060

NOAA-National Ocean Survey
6001 Executive Blvd.
Rockville, MD 20852
Attn: B. H. Chovitz
T. Soler
E. Herbrechtsmier
R. Snay

Battelle Columbus Laboratories
505 King Avenue
Columbus, OH 43201
Attn: A. G. Mourad
S. Gopalapillai

Smithsonian Astrophysical Observatory
60 Garden Street
Cambridge, MA 02138
Attn: E. M. Gaposchkin

U.S. Naval Postgraduate School
Monterey, CA 93940
Attn: Library

U.S. Naval Academy
Annapolis, MD 21402
Attn: Technical Library

The Ohio State University
Department of Geodetic Science
164 W. Nineteenth Avenue
Columbus, OH 43210
Attn: Dr. I. I. Mueller
Dr. U. A. Uotila
Dr. R. Rapp

Aerospace Corporation
2350 East El Segundo Boulevard
El Segundo, CA
Attn: R. Farrar
L. Wong

Department of Civil Engineering
Purdue University
Lafayette, IN
Attn: Prof. L. Kivioja

Division of National Mapping
P.O. Box 667
Canberra City, A.C.T. 2601
Australia
Attn: J. Luck

Technische Hochschule Graz
Technikerstr. 4, A-8010 Graz
Austria
Attn: Prof. P. Meissl
Prof. H. Moritz
Prof. K. Rinner

Universidade Federal Do Parana
Curso De Pos
Graduacao Em Ciencias Geodesicas
CX. P. 1611
60.000, Curitiba, Parana, Brazil
Attn: Prof. C. Gemael
Prof. J. Bittencourt de Andrade

Geodetic Survey of Canada
615 Booth St., Ottawa, Canada
Attn: J. Kolba
A. Peterson
A. Porter

Department of Survey Engineering
University of New Brunswick
Fredericton, N.B., Canada
Attn: Dr. P. Vanicek

Geodetic Institute of the Delft
Technological University
Delft, Netherlands
Attn: Dr. J. E. Alberda
Dr. L. Aardoy
Dr. B. Van Gelder

Local:
E41
K10 (50)
K50 (10)
K01
K05
X210 (2)
X2101 (GIDEP)