

AD-A066 800

COLORADO STATE UNIV FORT COLLINS DEPT OF CHEMISTRY  
CHEMICAL AND ELECTROCHEMICAL STUDIES IN ROOM TEMPERATURE ALUMIN--ETC(U)  
MAR 79 H L CHUM, R A OSTERYOUNG  
N00014-77-C-0004  
NL

UNCLASSIFIED

TR-9

| OF |

AD  
A066800



END

DATE

FILMED

5-79

DDC

**LEVEL**

Q

OFFICE OF NAVAL RESEARCH

Contract N00014-77-C-0004

TECHNICAL REPORT NO. 9

CHEMICAL AND ELECTROCHEMICAL STUDIES IN ROOM TEMPERATURE

ALUMINUM HALIDE-CONTAINING MELTS

by

Helena Li Chum and R. A. Osteryoung

Prepared for Publication in

"Ionic Liquids"

Department of Chemistry  
Colorado State University  
Fort Collins, Colorado 80523

March 1979

Reproduction in whole or in part is permitted for any purpose of the United States Government.

Approved for Public Release; Distribution Unlimited.



79 04 03 021

AD A0 66800

DDC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report 19	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CHEMICAL AND ELECTROCHEMICAL STUDIES IN ROOM TEMPERATURE ALUMINUM HALIDE-CONTAINING MELTS.		5. TYPE OF REPORT & PERIOD COVERED 9 Technical rept.
7. AUTHOR(s) 10 Helena Li Chum and R. A. Osteryoung		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Colorado State University Fort Collins, Colorado 80523		8. CONTRACT OR GRANT NUMBER(s) 15 N00014-77-C-0004
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Program Office of Naval Research Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 1241p.		12. REPORT DATE 11 March 1979 13. NUMBER OF PAGES 36
16. DISTRIBUTION STATEMENT (of this Report)  Approved for Public Release; Distribution Unlimited.		15. SECURITY CLASS. (of this report) Unclassified
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication in "Ionic Liquids," edited by David Lovering and Douglas Inman, Plenum Publishing Company.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Chloroaluminate melts; n-butylpyridinium chloride; Raman spectroscopy; acid-base chemistry; electrochemistry		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This review of recent work in the area covers a brief overview of work in sodium tetrachloroaluminate melts, with emphasis on acid-base chemistry, species characterization, chemistry of chalcogenides and chalcogens and the stability of radical ions. Work in the aluminum chloride-n-butylpyridinium room temperature melt is covered in more detail with regard to species characterization by Raman spectroscopy, acid-base chemistry, the electrochemical and spectroscopic studies of aromatic hydrocarbons, and an indication of other studies of inorganic species in these melts.		

DD FORM 1473  
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

79 04 03 021

404 992

JP

# TABLE OF CONTENTS

1.	INTRODUCTION. . . . .	1
2.	ALUMINUM HALIDE-ALKALI METAL HALIDE MOLTEN SALT SYSTEMS . . . . .	3
2a.	Acid-Base Chemistry. . . . .	3
2b.	Species Characterization . . . . .	4
2c.	Electrochemistry of Chalcogenide Ions and Chalcogens . . . . .	5
2d.	Stability of Radical Ions . . . . .	7
3.	ROOM TEMPERATURE MOLTEN SALT SYSTEMS - INTRODUCTORY REMARKS . . . . .	9
4.	ALUMINUM CHLORIDE - n-Butylpyridinium Chloride (BuPyCl) MOLTEN SALT SYSTEMS . . . . .	11
4a.	Species Characterization . . . . .	11
4b.	Acid-Base Chemistry. . . . .	13
4c.	Electrochemical and Spectroscopic Studies of Aromatic Hydrocarbons. . . . .	16
4d.	Electrochemical Study of the Fe(III)/Fe(II) System . . . . .	18
5.	CONCLUSIONS . . . . .	20
6.	ACKNOWLEDGEMENTS. . . . .	21
7.	REFERENCES. . . . .	22

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
PLANNING	<input type="checkbox"/>
JSTICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	SPECIAL
A	



## 1. INTRODUCTION

Molten salts have been used in industrial processes, such as for the production of aluminum, for many years. Applications in areas of molten salt battery systems, molten salt nuclear reactors, extraction from metal ores, electrorefining, etc., have prompted researchers to study the fundamental properties of molten salt systems for the past 30 years.<sup>1</sup> Recently, due partly to the relatively low working temperatures (150-200°C), the molten aluminum halides have been widely studied, e.g.,  $AlX_3/MX$ , where  $M^+$  is an alkali metal cation.<sup>1</sup> Very recently, work on "room temperature"  $AlX_3$  based systems has been initiated.

The primary scope of this review is the room temperature molten salt systems based on aluminum halides; however, it is pertinent to review some of the properties of the higher temperature aluminum halide-alkali metal halide melts as background for the discussion of the low temperature systems.

Several aspects of the chemistry of chloroaluminate melts have been reviewed. Boston<sup>2</sup> reviewed the molten salt chemistry of haloaluminates up to 1971 covering physical properties, solvent properties, some aspects of electrochemical and spectroscopic studies. Fung and Mamantov<sup>3</sup> detailed the electroanalytical chemistry of molten salts, particularly of molten haloaluminates, emphasizing the electrochemical behavior of the inorganic solutes that had been studied with emphasis on transition metal ions. Jones and Osteryoung<sup>4</sup>, in 1975, reviewed organic reactions in molten tetrachloroaluminate solvents of the type: condensation-addition, dehydrogenation-addition (Scholl reactions) and rearrangement-isomerizations. Plambeck<sup>5</sup> summar-




ized the various inorganic systems studied in these melts, hydrogen, elements of the groups IIIA, IVA, VA, VIA, VIIA, as well as transition metal elements, including actinides. A summary of electromotive force series in the 50:50 mole percent  $\text{AlCl}_3$ :NaCl at  $175^\circ\text{C}$  is presented. Mamantov and Osteryoung<sup>1</sup> described the acid-base dependent redox chemistry in molten chloroaluminates, basically from the point of view of the authors' own research.

With this brief introduction it is now appropriate to consider aspects of the chemistry which makes the chloroaluminate systems unique.

## 2. ALUMINUM HALIDE-ALKALI METAL HALIDE MOLTEN SALT SYSTEMS

### 2a. Acid-Base Chemistry

The aluminum halide-alkali metal halide melts display wide acidity changes with varying melt compositions. The acid-base properties can determine the redox and coordination chemistry in these media and several research groups have investigated this particular aspect over the past ten years.<sup>6-10</sup> The acid-base chemistry of the  $\text{AlCl}_3/\text{NaCl}$  molten salt system, in the range 49-70 mole percent of  $\text{AlCl}_3$ , can be represented<sup>9,10</sup> by the following three equilibria (175°C):



where the K values are on the mole fraction scale. The dominant acid-base equilibrium is described by Eq. 3 where  $\text{Al}_2\text{Cl}_7^-$  is the Lewis acid and the  $\text{Cl}^-$  is the base. This equilibrium is an autosolvolysis reaction, analogous to water autoprotolysis. The  $\text{pCl}^-$  ( $-\log[\text{Cl}^-]$ ), on the mole fraction scale, of a neutral melt (1:1  $\text{AlCl}_3:\text{NaCl}$ ) is 3.5 at 175°C. At this composition, the system can be considered essentially as  $\text{Na}^+\text{AlCl}_4^-$ . This melt can be made more acid (excess  $\text{AlCl}_3$ ) by anodizing an Al wire<sup>10</sup> or by addition of  $\text{AlCl}_3$ . The melt can be made more basic (excess  $\text{NaCl}$ ) by cathodizing an Al wire or by addition of  $\text{NaCl}$ . An Al electrode is employed to monitor the melt's acidity.<sup>9</sup> At 175°C, the  $\text{pCl}$  of the most basic melt saturated in  $\text{NaCl}$  is 1.1. The major acidity ( $\text{pCl}$ ) changes occur in the vicinity of the 1:1  $\text{AlCl}_3:\text{NaCl}$  mole ratio.

## 2b. Species Characterization.

The nature of the species present in these equilibria has been studied by Raman spectroscopy.<sup>11-15</sup> Mamantov *et al.*<sup>11</sup> have reported Raman spectra of sodium chloroaluminate melts and have reported evidence for species  $\text{AlCl}_4^-$ ,  $\text{Al}_2\text{Cl}_7^-$ ,  $\text{Al}_2\text{Cl}_6$ , and possibly  $\text{Al}_3\text{Cl}_{10}^-$ .<sup>11</sup> For the  $\text{AlCl}_4^-$  species, the major Raman vibrational frequencies and their respective polarizations were: 351 (P); 121 (D); 490 (D); and 186 (D)  $\text{cm}^{-1}$ , in order of decreasing intensity. For pure molten  $\text{Al}_2\text{Cl}_6$ , the major Raman-active frequencies were 341 (P); 218 (P); 119 (D); 104 (P)  $\text{cm}^{-1}$ . By assuming that in the 50/50 to 58/42 mole percent  $\text{AlCl}_3/\text{NaCl}$  range, the only Raman-active species are  $\text{AlCl}_4^-$  and  $\text{Al}_2\text{Cl}_7^-$ , the following vibrational frequencies were assigned to  $\text{Al}_2\text{Cl}_7^-$ : 313 (P); 432 (P); ~100 (D); and ~165 (D)  $\text{cm}^{-1}$ . Due to overlap of broad bands of  $\text{AlCl}_4^-$  at 121 and 186  $\text{cm}^{-1}$ , the frequencies at 100 and 165  $\text{cm}^{-1}$  were only estimated. Since the spectra in the high acidity range (>66 mole percent  $\text{AlCl}_3$ ) displayed other bands which could not be explained solely on the basis of  $\text{AlCl}_4^-$ ,  $\text{Al}_2\text{Cl}_7^-$ , and  $\text{Al}_2\text{Cl}_6$ , species such as  $\text{Al}_3\text{Cl}_{10}^-$  were postulated.<sup>11</sup> Raman spectra of the corresponding bromide and iodide melts have also been reported.<sup>13</sup>

Raman spectroscopic data for the potassium chloroaluminate melts have been reported by Oeye and coworkers.<sup>12,14</sup> Spectral data for  $\text{AlCl}_4^-$  agree well with those found in the sodium chloroaluminate melt.<sup>11</sup> The four absorption bands of  $\text{AlCl}_4^-$  verify  $T_d$  symmetry. Better low frequency spectroscopic data for  $\text{Al}_2\text{Cl}_7^-$  were also obtained<sup>12,14</sup>: 99 (D); 164 (D); 312 (P); and 435 (D)  $\text{cm}^{-1}$ . An earlier report<sup>15</sup> of Raman spectra of the 50/50 mole percent  $\text{AlCl}_3/\text{KCl}$  indicated a much larger number of bands (9) than observed for  $\text{AlCl}_3/\text{NaCl}$  (4). This melt was related to a distorted  $\text{AlCl}_4^-$  ( $C_{2v}$ ).<sup>15</sup>




According to Maroni and Cairns,<sup>16</sup> a number of the reported bands are weak and poorly defined and may be spectral artifacts.

A normal coordinate analysis for  $\text{Al}_2\text{Cl}_7^-$  assuming a  $D_{3d}$  model with a linear Al-Cl-Al bridge does not reproduce the experimental frequencies.<sup>12</sup> Structural determinations<sup>17</sup> by X-ray crystallography on the analogous  $\text{Al}_2\text{Br}_7^-$  ion either as a potassium or ammonium salt show that the anion consists of two  $\text{AlBr}_4^-$  tetrahedra sharing one corner in a staggered arrangement with a bent Al-Br-Al bridge ( $109.3^\circ$  and  $107.7^\circ$  respectively for the  $\text{K}^+$  and  $\text{NH}_4^+$  salts). The symmetry of the  $\text{Al}_2\text{Br}_7^-$  anion is close to  $C_{2v}$ . Structural determinations by the NQR double-resonance technique confirmed the structure of  $\text{Al}_2\text{Br}_7^-$  and led to a reasonable qualitative bond model for this ion.<sup>18</sup> The structure of the compound  $\text{Te}_4(\text{Al}_2\text{Cl}_7)_2$ <sup>19</sup> was determined and the  $\text{Al}_2\text{Cl}_7^-$  anion also consists of two tetrahedra sharing one corner in staggered arrangement with a bent Al-Cl-Al bridge ( $110.8^\circ$ ). Normal coordinate calculations assuming similar symmetry for the  $\text{Al}_2\text{Cl}_7^-$  anion might yield better agreement between calculated and experimental frequencies, but is clearly more difficult to do.


## 2c. Electrochemistry of the Chalcogenide Ions and Chalcogens

Mamantov et al.,<sup>20,21</sup> and Paulsen and Osteryoung<sup>22</sup> have studied the electrochemistry of sulfur and sulfide. Sulfur has been quantitatively reduced in both acid and basic melts at  $175^\circ\text{C}$ . Nernst plots for the sulfur/sulfide redox couple have a slope corresponding to a four-electron process, described as




The electrochemistry of oxide and chalcogenide ions in these molten salt systems was more quantitatively investigated by Osteryoung and coworkers<sup>23-26</sup>.

By titrating  $O^{2-}$ ,  $S^{2-}$ ,  $Se^{2-}$ , or  $Te^{2-}$  species in the melt with electrogenerated  $Al_2Cl_7^-$  ions, it was shown that in the basic to neutral melts these ions behave as dibases as described by the equilibrium:



where  $X^-$  may be  $O^-$ ,  $S^-$ ,  $Se^-$ , or  $Te^-$ , whereas in neutral to acidic melts they became tribases, the equilibrium then becoming:



The base strength of the ions in the acid melt was found to be in the order  $O^{2-} > S^{2-} > Se^{2-} \approx Te^{2-}$ .<sup>24</sup>

Oxide does not appear to be electroactive<sup>23</sup> in this molten salt system (cf. ref. 1) but the electrochemistry of sulfur<sup>22</sup>, selenium<sup>26</sup>, tellurium<sup>25</sup>, and their compounds has been thoroughly investigated using a variety of electrochemical methods. Sulfur was shown to oxidize very close to the anodic limit of the melt and it is therefore uncertain what the nature of the oxidation product is though in basic melts it appears to be  $S_2^{2+}$  whereas in acidic melts some higher oxidation state species appears likely. Spectrophotometric, potentiometric, and Raman data<sup>27</sup> indicate that the oxidation product is a S(IV) species. Addition of  $SCl_3^+ AlCl_4^-$ <sup>28</sup> to the melts reproduces the electrochemical and spectral behavior<sup>1,27</sup> observed in the electrochemical oxidation of sulfur.

The electrochemical behavior of selenium<sup>26</sup> is similar; it is reduced to selenide, more reversibly than sulfur is to sulfide, and can be oxidized to Se(IV). In basic melts this oxidation goes via a stable Se(II) intermediate whereas in acidic melts the oxidation is directly to Se(IV). The behavior of tellurium<sup>25</sup> is rather different as the metal is insoluble in the melt. In the most basic melts at temperatures above 250°C, Te(IV) is initially reduced

to a soluble Te(II) species, and then to the metal. In more acidic melts and at lower temperatures the Te(IV) is reduced directly to Te. Te metal rapidly reacts with Te(IV) to form  $\text{Te}_4^{2+}$ .

The nature of the chalcogen species involved in chloroaluminate melts, as determined by an interesting combination of potentiometric and spectrophotometric studies by Bjerrum and coworkers has been reviewed recently<sup>27</sup>. For instance, entities such as  $\text{Te}_2^{2+}$ ,  $\text{Te}_4^{2+}$ ,  $\text{Te}_6^{2+}$ , and  $\text{Te}_8^{2+}$ , stabilized by the large anions,  $\text{AlCl}_4^-$  and  $\text{Al}_2\text{Cl}_7^-$ , are present in the melt as well as  $\text{Te}^{2+}$ , which is stabilized by  $\text{Cl}^-$ .<sup>27,29,30</sup> Bjerrum and coworkers<sup>31,32</sup> have also studied the chloro-complexes of the  $\text{Te}^{4+}$  cation in  $\text{AlCl}_3/\text{KCl}$  melts and found the complexes  $\text{TeCl}_6^{2-}$ ,  $\text{TeCl}_5^{2-}$ ,  $\text{TeCl}_4$ , and  $\text{TeCl}_3^+$ , which are linked by melt acidity dependent equilibria.  $\text{Te}_4(\text{AlCl}_4)_2$  and  $\text{Te}_4(\text{Al}_2\text{Cl}_7)_2$ , where formally the tellurium is in a  $\frac{1}{2}$  oxidation state, have been isolated and characterized.<sup>18,33</sup> This agrees with earlier data by Bjerrum and Smith.<sup>34</sup>

Similarly, Bjerrum and coworkers<sup>27</sup> have presented evidence for ions:  $\text{Se}_4^{2+}$ ,  $\text{Se}_8^{2+}$ ,  $\text{Se}_2^{2+}$ ,  $\text{Se}_{16}^{2+}$  as the most likely other low oxidation states formed by reduction of Se(IV) with elemental selenium. The Se(IV) species is  $\text{SeCl}_3^+$  in most of the basic and most of the acidic range as deduced from spectroscopic and potentiometric data. For sulfur<sup>35</sup> the most consistent interpretation of potentiometric and spectrophotometric data was  $\text{S}_2^{2+}$ ,  $\text{S}_4^{2+}$ ,  $\text{S}_8^{2+}$ ,  $\text{S}_{16}^{2+}$ , together with S(II) and S(IV) most likely as  $\text{SCl}_3^+$ .

## 2d. Stability of Radical Ions

From a chemical point of view, another interesting finding is the stability of radical ions in these melts. Several aromatic amine radical cations were shown to be remarkably stable in these melts<sup>36</sup> at 175°C, in comparison to the poor stability found in aprotic organic solvents.<sup>4</sup> It has been suggested that this stability is a result of the "totally" anhydrous

nature of these melts, whereas most organic solvents contain varying residual amounts of water depending on the purification procedure. Triphenylamine,<sup>36</sup> N,N-dimethylaniline,<sup>36,37</sup> N,N,N',N'-tetramethylbenzidine,<sup>38</sup> quinone-hydroquinone systems,<sup>38</sup> sulfur heterocycles,<sup>40</sup> were studied and showed the interesting interplay of acid-base chemistry and redox processes with radical cation stabilization.



### 3. ROOM TEMPERATURE MOLTEN SALT SYSTEMS - INTRODUCTORY REMARKS

All the electrochemical and chemical reactions mentioned so far involve temperatures greater than  $150^{\circ}\text{C}$ . It is very clear 1) that the operating temperatures were above the boiling point of many organic solutes whose electrochemistry could be of interest and 2) that rapid homogeneous reactions could take place between melts and organic compounds.<sup>4,41</sup> For most other applications of the haloaluminates, a melt system with similar solvent properties operating at lower temperatures, e.g., room temperature, would be advantageous.

In 1951 a melt composed of 67 mole percent aluminum chloride and 33 mole percent of the quaternary organic salt ethylpyridinium bromide (a 2:1 molar ratio), which is molten at room temperature, was reported by Hurley and Wier.<sup>42</sup> Osteryoung and coworkers utilized this solvent for electrochemical and photochemical investigations.<sup>43-46</sup> The electrochemical oxidation of iron diimine complexes was shown to be a reversible process. Upon irradiation with low energy photons, the iron(II) diimine complexes were shown to be converted to the corresponding stable iron (III) compounds, with electron transfer to the acceptor ethylpyridinium cations.<sup>46</sup>

An interesting feature of the 2:1  $\text{AlCl}_3$ :ethylpyridinium bromide melt is its miscibility with co-solvents, e.g., benzene<sup>42</sup>, toluene, etc. Hurley and Wier<sup>42</sup> had observed that this mixed solvent system was better than the pure melt for electroplating of aluminum. Upon dilution with benzene, the conductivity of the solutions increases compared to the pure melt, while the viscosity decreases. This solvent system, melt plus benzene, proved very suitable to the study of organic<sup>43,49</sup> and organometallic<sup>45</sup> compounds. An advantage of the mixed solvent is that a number of organic and organometallic compounds, insoluble in the pure melt, are soluble in the melt plus benzene.<sup>45</sup> In the 50%(v/v) melt and benzene solution hexamethyl-

benzene (HMB) can be anodically oxidized to pentamethylbenzene and diphenylmethane.<sup>44</sup> This was the first reasonable synthesis in a molten salt system, where HMB can be selectively and quantitatively demethylated to mesitylene and diphenylmethane in a Friedel-Crafts type reaction.<sup>44</sup> The same solvent system has been employed to study the electrochemical oxidation of six metal carbonyls.<sup>45</sup> Chromium hexacarbonyl and iron pentacarbonyl were found to be reversibly oxidized to the corresponding seventeen electron cations.<sup>45</sup> In acetonitrile, only the former compound was found to undergo reversible one-electron oxidation.<sup>48</sup> However, this melt is not very suitable for investigating acid-base dependent reactions, since decreasing the acidity by increasing the ethylpyridinium bromide content raises the melting point sharply at the 50:50 mole percent composition.

Similar systems based on toluene plus  $\text{AlBr}_3$  and  $\text{KBr}$  in a 2:1 mole ratio have been employed by Gileadi and coworkers<sup>49-53</sup> primarily with interest in electroplating of aluminum.

#### 4. ALUMINUM CHLORIDE - N-(n-butylpyridinium) CHLORIDE (BuPyCl) MOLTEN SALT SYSTEMS

Recently, the  $\text{AlCl}_3$ -N-(n-butylpyridinium) chloride system (1:1 mole ratio) was found to melt at  $27^\circ\text{C}$ , and the 2:1 mole ratio exists as a liquid at temperatures well below ambient. In fact, a mole ratio range of 0.75:1 to 2:1 can be employed at temperatures only slightly greater than  $27^\circ\text{C}$ , very close to room temperature.<sup>47,54</sup>


##### 4a. Species Characterization<sup>54</sup>

Raman spectroscopy has been used to study the species composition of the molten salt system  $\text{AlCl}_3$ :BuPyCl over the 0.75:1 to 2.1:1 mole ratio range<sup>54</sup> at  $30^\circ\text{C}$ . Table I summarizes the experimental frequency shifts ( $\text{cm}^{-1}$ ) for the vibrational peaks. To characterize these spectra, the peak positions were compared to the assignments for  $\text{AlCl}_4^-$  and  $\text{Al}_2\text{Cl}_7^-$  (see above) in the  $\text{AlCl}_3/\text{MCl}$  ( $\text{M} = \text{Na}^+, \text{K}^+$ )<sup>11,12,14</sup> melts, and the spectra of  $\text{Al}_2\text{Cl}_6$ <sup>11</sup> (see above) and n-butylpyridinium cation (major frequencies in order of decreasing intensity in  $\text{cm}^{-1}$ : 1024; 125; 88-103; 644; 774). There is a straightforward correlation between the frequencies: 351 (P); 126 (D); 484 (D); and 184 (D) found in the low  $\text{AlCl}_3$ -content melts (0.75:1 - 1.0:1.0) and the frequencies assigned to  $\text{AlCl}_4^-$  anion<sup>11,12,14</sup>. As the  $\text{AlCl}_3$ -content increases, new vibrational peaks at 102 (shoulder) (D), 163 (D), 315 (P), and 434 (P) become dominant, and the bands assigned to  $\text{AlCl}_4^-$  do not contribute to the observed spectra within the experimental error. These frequencies compare very well with the previously assigned frequencies for the  $\text{Al}_2\text{Cl}_7^-$  anion, observed by Mamantov *et al.*<sup>11</sup> and Oeye *et al.*<sup>12,14</sup> (*vide supra*). Due to the overlap of the strong peak at  $315 \text{ cm}^{-1}$  in the high aluminum chloride content region or the strong peak at  $351 \text{ cm}^{-1}$  in the basic side, and the major  $\text{Al}_2\text{Cl}_6$  Raman active frequency at  $341 \text{ cm}^{-1}$ ,  $\text{Al}_2\text{Cl}_6$  cannot be

detected if present in small concentrations.


The molten salt system  $\text{AlCl}_3:\text{BuPyCl}$  can be described as follows as a function of composition.

2:1 mole ratio of  $\text{AlCl}_3:\text{BuPyCl}$  - At this mole ratio, the molten salt is best described as n-butylpyridinium dialuminum heptachloride,  $\text{BuPy}^+\text{Al}_2\text{Cl}_7^-$ . There is no evidence from the Raman spectra for the presence of  $\text{AlCl}_4^-$  anions, contrary to what was observed in the  $\text{AlCl}_3$ -alkali metal halides.<sup>11,12,14</sup> These facts suggest that at this composition the association:



must be virtually complete.

1:1 mole ratio of  $\text{AlCl}_3:\text{BuPyCl}$  - The major aluminum containing component anion in this composition is the  $\text{AlCl}_4^-$  anion. A small amount of the  $\text{Al}_2\text{Cl}_7^-$  anion may be present at this composition (cf. band intensities at  $433 \text{ cm}^{-1}$ ). Therefore, at this composition the melt contains  $\text{AlCl}_4^-$ ,  $\text{Al}_2\text{Cl}_7^-$ ,  $\text{Cl}^-$ , and possibly traces of  $\text{Al}_2\text{Cl}_6$ . Thus, the equilibrium:



identical to the dominant equilibrium in the  $\text{AlCl}_3/\text{NaCl}$  systems, (Eq. 3), may be considered to exist in the room temperature molten salt.

At intermediate compositions it is possible to estimate the ratio of the mole fractions of  $\text{Al}_2\text{Cl}_7^-$  and  $\text{AlCl}_4^-$  species by assuming that these mole fractions are proportional to the integrated intensity of the strongest bands for each species.<sup>54,12</sup> The ratios of 1.2:1 and 3.3:1 were obtained



respectively for the 1.5:1.0 and 1.75:1.0 molar ratios of  $\text{AlCl}_3:\text{BuPyCl}$ . Therefore, as the melt becomes  $\text{AlCl}_3$ -rich, the molar excess of  $\text{AlCl}_3$  is converted into  $\text{Al}_2\text{Cl}_7^-$ . At the limit, the 2:1 mole ratio  $\text{AlCl}_3:\text{BuPyCl}$ , this species is the major and possibly the only aluminum-containing species in the melt.

The large organic pyridinium cation does not influence the absorption frequencies of the anionic species,  $\text{AlCl}_4^-$  or  $\text{Al}_2\text{Cl}_7^-$ . This finding agrees well with previous data by Rytter and Oeye<sup>58</sup> on the influence of the alkali metal cation size on the absorption frequencies of  $\text{AlCl}_4^-$  anion. The smaller cation,  $\text{Li}^+$ , affects strongly asymmetric vibrations, whereas practically identical frequencies are obtained for  $\text{Na}^+$  and  $\text{K}^+$  within experimental error.

The major effect of the organic cation is to promote more extensive  $\text{Al}_2\text{Cl}_7^-$  formation at the 2:1 mole ratio than is found in the higher temperature haloaluminate-alkali metal halide molten salt systems.

#### 4b. Acid-Base Chemistry<sup>55</sup>

In order to quantify<sup>55</sup> the equilibrium described by Eq. 7 in the basic region, potentiometric titrations were carried out, by using an aluminum reference electrode, which behaves as a reversible chloride activity indicator electrode. From the Nernst equation:

$$E = E_o + (RT/3F) \ln(a_{\text{Al}^{3+}}/a_{\text{Al(O)}}) \quad (8)$$

incorporating the association constants for reactions (7) and formation of  $\text{AlCl}_4^-$ :



one obtains the following relation:

$$E = (RT/3F) \ln (a_{\text{AlCl}_4}^0 / a_{\text{AlCl}_4}^1) + (4RT/3F) \ln (a_{\text{Cl}^-}^1 / a_{\text{Cl}^-}^0) \quad (10)$$

where superscripts 0 and 1 refer to reference and variable states, respectively.

The description of the electrochemical cell can be given as follows:



where BuPyCl = n-butylpyridinium chloride. Figure 1 shows some results of potentiometric titrations in the 2.0:1.0 to 0.6:1.0 molar ratio of  $\text{AlCl}_3$ :BuPyCl at 30, 60, and 120°C. The sigmoidal curves in Figure 1 differ considerably from those obtained for the  $\text{AlCl}_3/\text{NaCl}$  system.<sup>9,10</sup> First, the potential difference between the two plateau regions is 2-3 times larger than that observed in the high temperature melts indicating a greater pCl change in the low temperature melt. Second, a change in the slope of the curve in the acid region is observed in the low temperature but not in the  $\text{AlCl}_3/\text{NaCl}$  system.<sup>9</sup> Third, in the basic portion of the curve, potentiometric data are less reproducible than those in the acid region, due to the appearance of a blue colored species at the Al electrode, indicating that secondary reactions are taking place.

From the potentiometric data it is possible to assess an approximate conditional equilibrium constant for Eq. 7 by using Eq. 10

$$K = \frac{[\text{Al}_2\text{Cl}_7^-][\text{Cl}^-]}{[\text{AlCl}_4^-]^2} = 3.8 \times 10^{-13} \quad (30^\circ\text{C})$$

The solid line in Figure 1 is a potential function represented by Eq. 10 with this equilibrium constant. The model is inappropriate in the acid region above 2:1 ratio.

In the basic region mixed potentials are measured due to the

spontaneous reduction of  $\text{BuPy}^+$  cations by elemental aluminum. Therefore, the constant  $K$ , determined by calculations over the entire acid and basic range, is a lower limit because of the positive shift in the measured potential due to the  $\text{BuPy}^+$  reduction.

A plot of the species distribution in this system from the 0.6:1.0 to the 1.8:1.0 mole ratio of  $\text{AlCl}_3$ : $\text{BuPyCl}$  is shown in Figure 2. It is interesting to note that the chloride ion mole fraction almost spans the range found in the proton/hydroxyl equilibrium in water.

The potentiometric curves at higher temperatures up to  $175^\circ\text{C}$  can also be characterized solely on the basis of Eq. 7. At  $175^\circ\text{C}$ ,  $K < 1.19 \times 10^{-8}$ , similar, but less than  $1.06 \times 10^{-7}$ ,<sup>9</sup> the constant for this equilibrium in the  $\text{AlCl}_3$ / $\text{NaCl}$  system. These findings are consistent with the data of Torsi and Mamantov<sup>59</sup> for the  $\text{AlCl}_3$ /alkali metal halide systems which indicate that the greater the polarizing power of the cation, the larger the autosolvolytic constant at a given temperature.

When comparing spectroscopic and potentiometric data for the  $\text{AlCl}_3$ /alkali metal halide and  $\text{AlCl}_3$ / $\text{BuPyCl}$  systems, one can observe that whereas in the former  $\text{Al}_2\text{Cl}_6$  is a species present throughout the studied composition and temperature ranges, being particularly important on the acid side, this Lewis acid does not seem to be a component of the room temperature molten salt. Even at temperatures as high as  $175^\circ\text{C}$ , little aluminum chloride vapor is detected over the 2:1  $\text{AlCl}_3$ : $\text{BuPyCl}$  melt. For the  $\text{AlCl}_3$ : $\text{BuPyCl}$  system in the molar ratios of 0.75:1 to 2:1, the acid-base chemistry is described by Eq. 7, whereas for the  $\text{AlCl}_3$ /alkali metal halides two additional equilibria involving  $\text{AlCl}_3$  and  $\text{Al}_2\text{Cl}_6$  (Eqs. 1 and 2) must be invoked.

#### 4c. Electrochemical and Spectroscopic Studies of Aromatic Hydrocarbons<sup>56</sup>

The electrochemical oxidation of twelve aromatic hydrocarbons: mesitylene, biphenyl, naphthalene, durene, phenanthrene, pentamethylbenzene, hexamethylbenzene, pyrene, anthracene, 9,10-diphenylanthracene, benzo(a)pyrene, and benzo(e)pyrene, was studied in the  $\text{AlCl}_3\text{:BuPyCl}$  systems in the 0.75:1 to 2:1 molar ratio, at 40°C.<sup>56</sup> All the hydrocarbons studied were shown to undergo a one-electron oxidation to the radical-cation at a potential that is independent of the melt composition. The radical-cations in these molten salt systems were found to be more stable than in acetonitrile.<sup>60-63</sup>

Figure 3 illustrates a linear correlation between the half-peak potential (proportional to the half-wave potential) and the first ionization potential (IP) of the parent hydrocarbon determined from photoelectron spectroscopy.<sup>64-65</sup> The equation:

$$E_{p/2} = 0.61 \text{ IP} - 3.34 \quad (11)$$

was found to fit the experimental data with a correlation coefficient  $\text{CC}=0.97$ . An analogous correlation with similar slope and intercept was obtained using the data from acetonitrile experiments<sup>60-63</sup> ( $\text{CC}=0.96$ ). These findings suggest that there is no strong interaction between the acid species in the melt and the hydrocarbons.


After referring all  $E_{p/2}$  potentials to that of the ferrocene/ferricinium ion couple in the melt and in acetonitrile, it has been observed that the oxidation potentials in the melt are about 0.1 V positive to those in acetonitrile, implying that it is slightly more difficult to oxidize these compounds in the melt than in acetonitrile.

Though the electrochemical data and correlations with IP seem to



suggest that the hydrocarbons do not interact with acid species in the melt, it was observed that some of the hydrocarbon solutions change color in acid melts suggesting that the hydrocarbons interacts with the solvent. When these solutions are made basic by addition of BuPyCl, they can be reversibly decolorized.

There are several reports in the literature of the existence of solid state  $\sigma$ -complexes between  $\text{AlCl}_3$  and hydrocarbons.<sup>68-71</sup> Table II compares the UV-visible spectra of solutions of some of the hydrocarbons studied in the 1.2:1.0  $\text{AlCl}_3$ :BuPyCl melt and the solid state  $\sigma$ -complexes.<sup>66-69</sup> It has been found that in melts more basic than 1.0:1.0, the spectra obtained correspond to those of the parent hydrocarbons. Though extinction coefficients were not reported, it has been observed that as the acidity increases, the intensity of the colored species increased. This behavior indicates that an acid-base type equilibrium between acid species in the melt,  $\text{Al}_2\text{Cl}_7^-$ , and the hydrocarbons is occurring. It has been suggested than an equilibrium of the type:



similar to that involved in the formation of carbonium ions<sup>70</sup> may exist, as in the example shown with anthracene. From the absence of any acid-base effects in the electrochemical oxidations of these hydrocarbons in the  $\text{AlCl}_3$ /BuPyCl melts it was concluded that either the equilibrium of Eq. 12 is very fast and the hydrocarbon is more readily oxidized than the  $\sigma$ -complex or alternatively the  $\sigma$ -complex and the hydrocarbon oxidize at the same potential.

An interesting but so far puzzling phenomenon was observed in acid melts for hydrocarbons with a  $E_{p/2}$  of less than 1.4 V at a glassy carbon electrode vs. Al reference electrode. In the more acid melts, these hydrocarbons (hexamethylbenzene, pyrene, anthracene, 9,10-diphenylanthracene, benzo(a)pyrene and benzo(e)pyrene) are slowly but spontaneously oxidized to the cation radicals by the molten salt itself. The nature of the oxidizing agent or its reduction product is unknown. The spontaneous oxidation ceases when the acidity of the melt is lowered.

#### 4d. Electrochemical Study of the Fe(III)/Fe(II) System

Hussey and King<sup>57</sup> have studied the Fe(III)/Fe(II) electrode reaction in the 2:1  $AlCl_3$ :BuPyCl melt at 30°C and in the 1:1  $AlCl_3$ :BuPyCl melt at 35°C, employing a rotating disc electrode, cyclic voltammetry and potentiometry.

At a glassy carbon or tungsten working electrode in the acid melt the reduction  $Fe(III) + e^- \rightarrow Fe(II)$  was found to be moderately rapid, with a standard heterogeneous electron transfer rate constant of  $1.3 \times 10^{-4} \text{ cm s}^{-1}$  and  $\alpha = 0.4$ . The heat of activation for this process is  $\sim 60 \text{ kJ mol}^{-1}$ . From Nernst plots the standard electrode potential ( $E^0$ ) for the Fe(III)/Fe(II) couple at 30°C is 2.01 V, with a  $dE^0/dt = 9.5 \times 10^{-4} \text{ mV/}^\circ\text{C}$ . From rotating disc data, the diffusion coefficient of the electroactive species was found to be  $4.4 \times 10^{-7} \text{ cm}^2 \text{ s}^{-1}$ , with an activation energy for diffusion of  $21 \text{ kJ mol}^{-1}$ .

The electrochemical reduction of Fe(III) in the basic melt is a one-electron reversible process. The standard electrode potential is 0.125 V (35°C) with  $dE^0/dt = 4.7 \times 10^{-4} \text{ mV/}^\circ\text{C}$ . The diffusion coefficient of the electroactive species in this medium is  $4.6 \times 10^{-7} \text{ cm}^2 \text{ s}^{-1}$  (35°C).

The marked increase in the rate of the Fe(III)/Fe(II) electrode reaction in going from the 2:1 to the 1:1  $\text{AlCl}_3\text{:BuPyCl}$  melts, and also the almost 2 V difference in standard potentials coincides with the marked difference in anionic species observed in these melts, reported above. That the Fe(III) species present in these two media are different solvates and/or complexes is to be expected, and is further supported by the different colors exhibited by the ion in these media: orange (acid) to canary yellow (basic). A more complete conclusion about the nature of the species present in these media must await further investigations. The electrochemical behavior of Fe(III) ions illustrates the influence of the melt composition, therefore acidity, on the redox and coordination chemistry of this transition metal ion in these media.

## 5. CONCLUSIONS

The newly developed molten salt systems based on aluminum halides and alkylpyridinium halides, principally the  $\text{AlCl}_3$ :n-butylpyridinium chloride system, appear to be very suitable media for the study of chemical and electrochemical acid dependent reactions at a very convenient temperature range close to room temperature. It is worth emphasizing that these new ionic liquids offer new possibilities for structural studies over a wide range of temperatures. Miscibility of these molten salt systems with aromatic co-solvents such as benzene, toluene, etc., suggests the possibility of expanding the number of organic and organometallic compounds which may be investigated, in addition to improving the physical properties of the solvent system by decreasing the viscosity and increasing the electrical conductivity.

From a more practical point of view, it appears that these systems may also be useful battery electrolytes. Research of possible electrode systems is therefore another branch for future research.



6. ACKNOWLEDGEMENTS

Support from the Air Force Office of Scientific Research and the Office of Naval Research is acknowledged. Helpful discussions with Drs. J. H. Christie and R. I. Gale are appreciated.

7. REFERENCES

- (1) G. Mamantov and R. A. Osteryoung in "Characterization of Solutes in Nonaqueous Solvents," G. Mamantov, Ed., Plenum Press, 1976, p. 223-249.
- (2) C. R. Boston in "Advances in Molten Salt Chemistry," Vol. 1, J. Braunschtein, G. Mamantov and G. P. Smith, Eds., Plenum Press, NY, 1971, p. 129-163.
- (3) K. W. Fung and G. Mamantov, *ibid.*, Vol. 2, 1973, p. 199-254.
- (4) H. L. Jones and R. A. Osteryoung, *ibid.*, Vol. 3, 1975, p. 121-176.
- (5) J. A. Plambeck, "Fused Salt Systems," Vol. X of "Encyclopedia of Electrochemistry of the Elements," A. J. Bard, Ed., Marcel Dekker, NY, 1976, p. 233-254.
- (6) B. Tremillon and G. Letisse, J. Electroanal. Chem. 17, 37 (1968).
- (7) G. Torsi and G. Mamantov, Inorg. Chem. 10, 1900 (1971).
- (8) A. A. Fannin, L. A. King and D. W. Seegmiller, J. Electrochem. Soc. 119, 801 (1972).
- (9) L. G. Boxall, H. L. Jones and R. A. Osteryoung, J. Electrochem. Soc. 120, 223 (1973).
- (10) L. G. Boxall, H. L. Jones and R. A. Osteryoung, J. Electrochem. Soc. 121, 212 (1974).
- (11) G. Torsi, G. Mamantov and G. M. Begun, Inorg. Nucl. Chem. Letters 6, 553 (1970).
- (12) S. J. Cyvin, P. Klaeboe, E. Rytter and H. A. Oeye, J. Chem. Phys. 52, 2776 (1970); H. A. Oeye, E. Rytter, P. Klaeboe and C. J. Cyvin, Acta Chem. Scand. 25, 559 (1971).
- (13) G. M. Begun, C. R. Boston, G. Torsi and G. Mamantov, Inorg. Chem. 10, 886 (1971).
- (14) E. Rytter, H. A. Oeye, S. J. Cyvin, B. N. Cyvin and P. Klaeboe, J. Inorg. Nucl. Chem. 35, 1185 (1973).
- (15) K. Balasubrahmanyam and L. Nanis, J. Chem. Phys. 42, 676 (1965).
- (16) V. A. Maroni and E. J. Cairns in "Molten Salts - Characterization and Analysis," G. Mamantov, Ed., Marcel Dekker, NY, London, 1969, p. 245.
- (17) E. Rytter, B.E.D. Rytter, H. A. Oeye and J. Krogh-Moe, Acta Crystallogr. 29B, 1541 (1973); 31B, 2177 (1975).

- (18) N. Weiden and A. Weiss, *J. Magn. Resonance* 30, 403 (1978).
- (19) T. W. Couch, D. A. Lokken and J. D. Corbett, *Inorg. Chem.* 11, 357 (1972).
- (20) R. Marassi, G. Mamantov and J. Q. Chambers, *Inorg. Nucl. Chem. Letters* 11, 245 (1975).
- (21) R. Marassi, G. Mamantov and J. Q. Chambers, *J. Electrochem. Soc.* 123, 1128 (1976).
- (22) K. A. Paulsen and R. A. Osteryoung, *J. Amer. Chem. Soc.* 98, 6866 (1976).
- (23) B. Gilbert and R. A. Osteryoung, *J. Amer. Chem. Soc.* 100, 2725 (1978).
- (24) J. Robinson, B. Gilbert and R. A. Osteryoung, *Inorg. Chem.* 16, 3040 (1977).
- (25) J. Robinson and R. A. Osteryoung, *J. Electrochem. Soc.* 125, 1784 (1978).
- (26) J. Robinson and R. A. Osteryoung, *J. Electrochem. Soc.* 125, 1454 (1978).
- (27) N. Bjerrum in "Characterization of Solutes in Nonaqueous Solvents," G. Mamantov, Ed., Plenum Press, NY, 1976, p. 251-271.
- (28) H. E. Doorenbos, J. C. Evans and R. U. Kagel, *J. Phys. Chem.* 74, 3385 (1970).
- (29) N. J. Bjerrum, *Inorg. Chem.* 9, 1965 (1970); 10, 2578 (1971); 11, 2648 (1972).
- (30) R. Fehrmann, N. J. Bjerrum and M. A. Andreassen, *Inorg. Chem.* 15, 2187 (1976).
- (31) J. M. vonBarnes, N. J. Bjerrum and D. R. Nielsen, *Inorg. Chem.* 13, 1708 (1974).
- (32) F. W. Paulsen, N. J. Bjerrum and D. R. Nielsen, *Inorg. Chem.* 13, 2693 (1976).
- (33) D. J. Prince, J. D. Corbett and B. Garbisch, *Inorg. Chem.* 9, 2731 (1970).
- (34) N. J. Bjerrum and G. P. Smith, *J. Amer. Chem. Soc.* 90, 4472 (1968).
- (35) R. Fehrmann, N. J. Bjerrum and F. W. Paulsen, *Inorg. Chem.* 17, 1195 (1978).
- (36) H. L. Jones, L. G. Boxall and R. A. Osteryoung, *J. Electroanal. Chem.* 38, 476 (1972).

- (37) H. L. Jones and R. A. Osteryoung, J. Electroanal. Chem. 49, 281 (1974).
- (38) D. E. Bartak and R. A. Osteryoung, J. Electrochem. Soc. 122, 600 (1975).
- (39) D. E. Bartak and R. A. Osteryoung, J. Electroanal. Chem. 74, 68 (1976).
- (40) K. W. Fung, J. Q. Chambers and G. Mamantov, J. Electroanal. Chem. 47, 81 (1973).
- (41) V. R. Koch, L. L. Miller and R. A. Osteryoung, J. Org. Chem. 39, 2416 (1974).
- (42) F. H. Hurley and J. P. Wier, J. Electrochem. Soc. 98, 203 (1951).
- (43) H. L. Chum, V. R. Koch, L. L. Miller and R. A. Osteryoung, J. Amer. Chem. Soc. 97, 3264 (1975).
- (44) V. R. Koch, L. L. Miller and R. A. Osteryoung, J. Amer. Chem. Soc. 98, 5277 (1976).
- (45) H. L. Chum, D. Koran and R. A. Osteryoung, J. Organometallic Chem. 140, 349 (1977).
- (46) H. L. Chum, D. Koran and R. A. Osteryoung, J. Amer. Chem. Soc. 100, 310 (1978).
- (47) J. Robinson, R. C. Bugle, H. L. Chum, D. Koran and R. A. Osteryoung, J. Amer. Chem. Soc., in press (1979).
- (48) C. J. Pickett and D. Pletcher, J. Chem. Soc. Chem. Comm., 660 (1974).
- (49) E. Peled and E. Gileadi, Plating 62, 342 (1975); J. Electrochem. Soc. 123, 15 (1976).
- (50) A. Reger, E. Peled and E. Gileadi, J. Electrochem. Soc. 123, 639 (1976).
- (51) E. Peled, A. Mitavsky, A. Reger and E. Gileadi, J. Electroanal. Chem. Int. Electrochem. 75, 677 (1977).
- (52) E. Peled, A. Mitavsky and E. Gileadi, Z. Phys. Chem. 96, 111 (1976).
- (53) S. Ziegel, E. Peled and E. Gileadi, Electrochim. Acta 23, 363 (1978).
- (54) R. J. Gale, B. Gilbert and R. A. Osteryoung, Inorg. Chem. 17, 2728 (1978).
- (55) R. J. Gale and R. A. Osteryoung, Inorg. Chem., in press,
- (56) J. Robinson and R. A. Osteryoung, J. Amer. Chem. Soc., 101, 323 (1979).



- (57) C. L. Hussey and L. A. King, J. Electroanal. Chem., submitted for publication, 1978.
- (58) E. Rytter and H. A. Oeye, J. Inorg. Nucl. Chem. 35, 4311 (1973).
- (59) G. Torsi and G. Mamantov, Inorg. Chem. 11, 1439 (1972).
- (60) H. Lund, Acta Chem. Scand. 11, 1232 (1957).
- (61) W. C. Niekam and M. M. Desmond, J. Amer. Chem. Soc. 86, 4811 (1964).
- (62) E. S. Pysch and M. C. Yang, J. Amer. Chem. Soc. 85, 2124 (1963).
- (63) R. E. Sioda, J. Phys. Chem. 72, 2322 (1968).
- (64) W. C. Herndon, J. Phys. Chem. 98, 887 (1976).
- (65) F. I. Vilesov, Zn. Fiz. Khim. 35, 2010 (1961).
- (66) M. Sato and Y. Aoyama, Bull. Chem. Soc. Japan 46, 631 (1973).
- (67) S. J. Costanzo and W. B. Jurinski, Tetrahedron 23, 2571 (1967).
- (68) M. M. Perkampus and Th. Kranz, Z. Physik Chem. Neue Folge 34, 213 (1962).
- (69) M. M. Perkampus and Th. Kranz, Z. Physik Chem. Neue Folge 38, 295 (1963).

TABLE I

Summary of experimental Raman frequency shifts ( $\text{cm}^{-1}$ ) for aluminum chloride:  
1-butylpyridinium chloride mixtures at room temperatures  
from reference 54.

Molar Ratio $[\text{AlCl}_3]:[\text{RCl}]$				
0.75:1.0	1.0:1.0	1.5:1.0	1.75:1.0	2.0:1.0
			-95 sh	-102 sh
-126 sh (0.9)	-127 sh (1.3)	-126 sh		
		-159 (1.2)	-163 (1.5)	-163 (2.3)
-186 (2.1)	-184 (2.0)	-183 (2.8)	-183 (1.6)	-182 (0.9)
-298 (0.9)	-295 (0.7)	-296 sh (1.4)	-295 sh (1.6)	-295 sh (1.1)
		-314 (8.8)	-316 (10.0)	-315 (10.0)
-351 (10.0)	-352 (10.0)	-351 (10.0)	-352 (5.0)	
-433 (0.1)	-433 (0.1)	-434 (2.7)	-433 (3.3)	-434 (2.2)
-485 (0.5)	-483 (0.4)			
-653 (2.8)	-650 (2.4)	-651 (3.2)	-647 (4.7)	-651 (2.7)
-775 (0.7)	-770 (0.5)	-774 (0.2)	-772 (0.2)	-769 (1.0)

TABLE II

Spectroscopic data for aromatic hydrocarbons in a 1.2:1 aluminum chloride:n-butylpyridinium chloride molten salt<sup>56</sup> and for solid state complexes with  $\text{AlCl}_3$ .<sup>68,69,71</sup>

Aromatic Hydrocarbon	$\lambda_{\text{max}}$ /nm Melt <sup>56</sup>	$\lambda_{\text{max}}$ /nm Solid State Complex
Mesitylene	367	365 <sup>69</sup>
Durene	382	380 <sup>69</sup>
Pentamethylbenzene	380	385 <sup>69</sup>
Hexamethylbenzene	398	400 <sup>69</sup>
Naphthalene	393	410, <sup>69</sup> 377 <sup>71</sup>
Phenanthrene	417	425, <sup>69</sup> 653 <sup>71</sup>
Pyrene	451, 465, 475, 492	520, <sup>69</sup> 480 <sup>71</sup>
Anthracene	417	418, <sup>71</sup> 560, <sup>69</sup> 420 <sup>68</sup>

TABLE III

Physical properties of some alkyl pyridinium halide - aluminum halide melts from reference 47.

Melt	Temp/ $^{\circ}\text{C}$	Density/ $\text{g}\cdot\text{cm}^{-3}$	Viscosity c.p.	Electrochemical Window/ $\text{V}^{\text{a}}$
2:1 $\text{AlCl}_3$ :BuPyCl	25	1.33	27.0	0.0 to 2.1
1:1 $\text{AlCl}_3$ :BuPyCl	40	1.24	20.9	-0.4 to 2.0
2:1 $\text{AlCl}_3$ :EtByPr	25	1.52	25.0	-0.2 to 1.8
2:1 $\text{AlBr}_3$ :EtByPr	25	2.20	~50	0.2 to 1.6

a) At a glassy carbon electrode versus an Al wire reference electrode in the same melt.



Figure 1

Potentiometric data for  $\text{AlCl}_3$ :1-butylpyridinium chloride versus  $\text{Al(O)}/2:1$  mole ratio mixture reference; x  $30^\circ\text{C}$ , o  $60^\circ\text{C}$ , ●  $120^\circ\text{C}$ ; — theory ( $30^\circ\text{C}$ ).  
After Gale and Osteryoung.<sup>55</sup>




Figure 2

Mole fraction,  $N$ , of the major species in the melt at  $30^{\circ}\text{C}$  as a function of the net  $\text{AlCl}_3:\text{RCI}$  mole ratio. After Gale and Osteryoung.<sup>55</sup>







Figure 3

The correlation between  $E_{p/2}$  values for the oxidation of aromatic hydrocarbons in the  $AlCl_3$ :n-butylpyridinium chloride molten salt and their photoionization potentials. (1) Mesitylene; (2) Biphenyl; (3) Napthalene; (4) Durene; (5) Phenanthrene; (6) Pentamethylbenzene; (7) Hexamethylbenzene; (8) Pyrene; (9) Anthracene; (10) 9,10-Diphenylanthracene; (11) Benzo(a)pyrene; and (12) Benzo(e)pyrene. After Robinson and Osteryoung.<sup>56</sup>



TECHNICAL REPORT DISTRIBUTION LIST, 051C

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. M. B. Denton University of Arizona Department of Chemistry Tucson, Arizona 85721	1	Dr. K. Wilson University of California, San Diego Department of Chemistry La Jolla, California	1
Dr. R. A. Osteryoung Colorado State University Department of Chemistry Fort Collins, Colorado 80521	1	Dr. A. Zirino Naval Undersea Center San Diego, California 92132	1
Dr. B. R. Kowalski University of Washington Department of Chemistry Seattle, Washington 98105	1	Dr. John Duffin United States Naval Postgraduate School Monterey, California 93940	1
Dr. S. P. Perone Purdue University Department of Chemistry Lafayette, Indiana 47907	1	Dr. G. M. Hieftje Department of Chemistry Indiana University Bloomington, Indiana 47401	1
		Dr. Victor L. Rehn Naval Weapons Center Code 3813 China Lake, California 93555	1
Dr. D. L. Venezky Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. Christie G. Enke Michigan State University Department of Chemistry East Lansing, Michigan 48824	1
Dr. H. Freiser University of Arizona Department of Chemistry Tucson, Arizona 85721		Dr. Kent Eisentraut, MBT Air Force Materials Laboratory Wright-Patterson AFB, Ohio 45433	1
Dr. Fred Saalfeld Naval Research Laboratory Code 6110 Washington, D.C. 20375	1	Walter G. Cox, Code 3632 Naval Underwater Systems Center Building 148 Newport, Rhode Island 02840	1
Dr. H. Chernoff Massachusetts Institute of Technology Department of Mathematics Cambridge, Massachusetts 02139	1		

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. George Sandoz	1	U.S. Army Research Office P.O. Box 1211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Weapons Center China Lake, California 93555 Attn: Dr. A. B. Amster Chemistry Division	1
ONR Area Office One Hallidie Plaza, Suite 601 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Naval Civil Engineering Laboratory Port Hueneme, California 93401 Attn: Dr. R. W. Drisko	1
ONR Branch Office Building 114, Section D 666 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Professor K. E. Woehler Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Naval Ship Research and Development Center Annapolis, Maryland 21401 Attn: Dr. G. Bosmajian Applied Chemistry Division	1
		Naval Ocean Systems Center San Diego, California 91232 Attn: Dr. S. Yamamoto, Marine Sciences Division	1

Encl 1



TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Paul Delahay New York University Department of Chemistry New York, New York 10003	1	Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1
Dr. R. A. Osteryoung Colorado State University Department of Chemistry Fort Collins, Colorado 80521	1	Dr. P. J. Hendra University of Southampton Department of Chemistry Southampton SO9 5NH United Kingdom	1
Dr. E. Yeager Case Western Reserve University Department of Chemistry Cleveland, Ohio 44106	1	Dr. Sam Perone Purdue University Department of Chemistry West Lafayette, Indiana 47907	1
Dr. D. N. Bennion University of California Chemical Engineering Department Los Angeles, California 90024	1	Dr. Royce W. Murray University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus California Institute of Technology Department of Chemistry Pasadena, California 91125	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Technical Library	1
Dr. J. J. Auborn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. J. H. Ambrus The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Telephone Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co, Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Oakland University Department of Chemistry Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated Five Lee Street Cambridge, Massachusetts 02139	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No.</u> <u>Copies</u>
Dr. H. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1
Larry E. Plew Naval Weapons Support Center Code 3073, Building 2906 Crane, Indiana 47522	1
S. Ruby DOE (STOR) 600 E Street Washington, D.C. 20545	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California 92803	1
Dr. M. G. Sceats University of Rochester Department of Chemistry Rochester, New York 14627	1