

REPORT SAMSO-TR-79-26

FVF

Analysis and Computation of the Derivative Function for an Isolated Lorentz Line

> STEPHEN J. YOUNG Chemistry and Physics Laboratory Laboratory Operations The Aerospace Corporation El Segundo, Calif. 90245

D APR 12 1979 LIT

20 March 1979

Interim Report

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

Prepared for

SPACE AND MISSILE SYSTEMS ORGANIZATION AIR FORCE SYSTEMS COMMAND Los Angeles Air Force Station P.O. Box 92960, Worldway Postal Center Los Angeles, Calif. 90009

04 09 119

79



DOC FILE COPY

This interim report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract No. F04701-78-C-0079 with the Space and Missile Systems Organization, Contracts Management Office, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by S. Siegel, Director, Chemistry and Physics Laboratory. Gerhard E. Aichinger was the project officer for Mission-Oriented Investigation and Experimentation (MOIE) Programs.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

Gerhard E. Aichinger Project Officer

FOR THE COMMANDER

Frank J. Bane, Chief Contracts Management Office

UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered) READ INSTRUCTIONS BEFORE COMPLETING FORM REPORT DOCUMENTATION PAGE REPORT NUMBER 2. GOVT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER SAMSOHTR-79-26 TYPE OF REPORT & PERIOD COVERED TITLE (and Subtitle) ANALYSIS AND COMPUTATION OF THE Interim rept DERIVATIVE FUNCTION FOR AN ISOLATED LORENTZ LINE . 6. PERFORMING ORS. REPORT NUMBER TR-0079(4970-40)-2 CONTRACT OR GRANT NUMBER(.) AUTHOR(.) Stephen J. Young FØ47Ø1-78-C-ØØ79 PERFORMING ORGANIZATION NAME AND ADDRESS PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS The Aerospace Corporation El Segundo, Calif. 90245 11. CONTROLLING OFFICE NAME AND ADDRESS 12. REPORT DATE Space and Missile Systems Organization 20 March 1979 Air Force Systems Command NUMBER OF PAG 22 Los Angeles, Calif. 90009 15. SECURITY CLASS. 14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office) Unclassified 154. DECLASSIFICATION/DOWNGRADING SCHEDULE 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different from Report) 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Line Absorptance (2/pi) integral from & to infinity of (exp <-2x/(1+(+to-square)(2-square))) **Radiative Transfer** 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) d = / (1+ =-squared) Analysis and methods of computation of the function $y(x, \rho) = \int_{T} \int \left(exp \left(- \frac{1}{2} \right) \right) dx$ $\frac{-2x}{+\rho^2 z^2}$ $\frac{dz}{1+z}$ 1 + zare presented. Power series, asymptotic series, and other expansions are developed for six regions of the positive quadrant of the xp plane. These expansions allow efficient calculation of $y(x, \rho)$ to a relative accuracy of 0.01% or better. FORM 1473 UNCLASSIFIED CSIMILE SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

CONTENTS

INTRODUC	CTION 3
METHODS	OF COMPUTATION
А.	Expansion for Small x (Region 1) 7
в.	Expansion for Small ρ (Region 2)
с.	Expansion for Large ρ (Region 3) 12
D.	Expansion for Large x (Region 4) 14
E.	Expansion for Large x and ρ (Region 5)
F.	Intermediate x and ρ (Region 6)
REFEREN	ICES 21



INTRODUCTION

Treatment of radiative transfer for an isolated Lorentz line in the Lindquist-Simmons approximation^(1, 2) requires the evaluation of the derivative function $y(x, \rho)$ defined by

$$y(x, \rho) = \frac{2}{\pi} \int_0^\infty \exp\left\{-\frac{2x}{1+\rho^2 z^2}\right\} \frac{dz}{1+z^2}$$
(1a)

or, with the change of variable $tan(\theta/2) = \rho z$, by

$$y(x, \rho) = \frac{2\rho}{\pi} \int_0^{\pi} \frac{e^{-x(1+\cos\theta)}}{(\rho^2+1) + (\rho^2-1)\cos\theta} d\theta.$$
 (1b)

This same function is used in the resonance absorption band model formulation of Cobb. ⁽³⁾ In both applications, x is a measure of optical depth, and ρ is a ratio of line widths. The range of both x and ρ is zero to infinity.

Prior work on the evaluation of $y(x, \rho)$ includes the rational approximation of Lindquist and Simmons⁽¹⁾ (accuracy not stated), the tabulation of Young⁽²⁾ (accuracy of one part in 10⁵), and the series expansions of $Cobb^{(3)}$ (accuracy of < 1% for $0.01 \le x \le 10^5$, $10^{-3} \le \rho \le 10^3$, and $10^{-3} \le y \le 1$). The present work considers the efficient calculation of $y(x, \rho)$ to a relative accuracy of $\le 0.01\%$ for the entire positive quandrant of the xp plane.

The general features of the function are shown in Fig. 1. The function limits are



-3-



Fig. 1. The Function $y(x, \rho)$. From Ref. 2.

$$y(\mathbf{x}, \rho) \rightarrow \begin{cases} 1 & \mathbf{x} \rightarrow 0 \\ \rho/\sqrt{2\pi\mathbf{x}} & \mathbf{x} \rightarrow \infty \\ e^{-2\mathbf{x}} & \rho \rightarrow 0 \\ e^{-\mathbf{x}}I_0(\mathbf{x}) & \rho \rightarrow 1 \\ 1 & \rho \rightarrow \infty \end{cases}$$
(2)

where $I_0(x)$ is the modified Bessel function of order zero.

The dramatic difference in basic mathematical form that the function assumes in different regions of the $x\rho$ plane makes it impractical to use only one method of calculation. Six approximation expansions are used in the present method. These approximations are considered in the following section, and the regions in which they apply are indicated in Fig. 2.



Fig. 2. Regions for which the Various Approximations to $y(x, \rho)$ Apply

METHODS OF COMPUTATION

A. EXPANSION FOR SMALL x (REGION 1)

For $x \le 1$, an expansion of the exponential factor of equation (1a) is made, and $y(x, \rho)$ is written as

$$y(x, \rho) = \sum_{n=0}^{\infty} (-1)^n \frac{(2x)^n}{n!} E_n(\rho)$$
 (3)

where

$$E_{n}(\rho) = \frac{2}{\pi} \int_{0}^{\infty} \frac{dz}{(1+z^{2})(1+\rho^{2}z^{2})^{n}}.$$
 (4)

Explicit evaluation of $E_n(\rho)$ by the method of residues for the first few values of n leads by induction to the convenient recurrence relation

$$E_{n}(\rho) = \int \frac{2n-1}{2n} E_{n-1}(\rho) \qquad \rho = 1$$
(5a)

$$\left(\frac{1}{\rho^{2}-1}[\rho C_{n} - E_{n-1}(\rho)] \quad \rho \neq 1$$
 (5b)

with

$$E_{0}(\rho) = 1 \text{ (for all } \rho),$$

$$C_{n} = \frac{2n-3}{2n-2} C_{n-1} = \frac{(2n-3)!}{2^{2n-3}(n-1)!(n-2)!},$$
(6)

and

$$C_1 = 1.$$

The explicit solutions for the first five $E_n(\rho)$ functions are

$$E_{0}(\rho) = 1,$$

$$E_{1}(\rho) = 1/(\rho + 1),$$

$$E_{2}(\rho) = (\rho + 2)/2(\rho + 1)^{2},$$

$$E_{3}(\rho) = (3\rho^{2} + 9\rho + 8)/8(1 + \rho)^{3},$$

$$E_{4}(\rho) = (5\rho^{3} + 20\rho^{2} + 29\rho + 16)/16(1 + \rho)^{4}.$$
(7)

Additional properties of $E_n(\rho)$ that can be derived are

$$E_{n}(\rho) \rightarrow \begin{cases} 1 & \rho \rightarrow 0 \\ C_{n}/\rho & \rho \rightarrow \infty \\ C_{n+1} & \rho \rightarrow 1 \\ 1/\rho \sqrt{\pi n} & \rho \rightarrow \infty \end{cases}$$
(8)

In practice, $y(x, \rho)$ is computed from equation (3) and the recurrence relations equations (5) and (6). Since equation (3) is an alternating series of decreasing terms, the accuracy of the approximation can be checked as each term is added to the series by testing the absolute value of the term added. The slowest convergence of equation (3) occurs for $\rho = 0$, in which case terms through n = 11 must be retained for a relative accuracy of 0.01%. The forward recurrence relation equation (5b) is unstable for ρ near unity. For $|1 - \rho| \le 0.044$, this stability sets in below n = 11. However, for this condition of ρ and for $x \leq 1$, $y(x, \rho)$ may be computed to the desired accuracy with only the first two terms (i.e., n = 0 and 1) of the Bessel function expansion considered in the following section.

B. EXPANSION FOR SMALL ρ (REGION 2)

For this and the following three approximations, the starting point is Cobb's expansion⁽³⁾ of $y(x, \rho)$ as a series of modified Bessel functions. The expansion is

$$y(x, \rho) = e^{-x} \left[I_0(x) + 2 \sum_{n=1}^{\infty} \alpha^n I_n(x) \right]$$
 (9)

where $\alpha = (\rho - 1)/(\rho + 1)$. For small ρ , $\alpha \rightarrow -1$, and we expand $y(x, \rho)$ in a Taylor series about $\alpha = -1$. At $\alpha = -1$, the series converges⁽⁴⁾ to give $y(x, 0) = e^{-2x}$. The first derivative of equation (9) evaluated at $\alpha = -1$ is

$$\frac{\mathrm{d}\mathbf{y}}{\mathrm{d}\alpha}\Big|_{\alpha = -1} = -2e^{-\mathbf{x}}\sum_{n=1}^{\infty}n(-1)^{n}\mathbf{I}_{n}(\mathbf{x}).$$
(10)

Use of the recurrence relation (4)

$$I_{n}(x) = \frac{x}{2n} [I_{n-1}(x) - I_{n+1}(x)]$$
(11)

in equation (10) and shifts of summation indices such that the sums over I_{n-1} and I_{n+1} are transformed into sums over I_n yield

$$\frac{dy}{d\alpha}\Big|_{\alpha = -1} = xe^{-x}\left[\sum_{n=0}^{\infty} -\sum_{n=2}^{\infty}\right](-1)^{n}I_{n}(x) = xe^{-x}[I_{0}(x) - I_{1}(x)].$$
(12)

Higher-order derivatives can be obtained by the same procedure. The resulting expansion to third order in ρ for $y(x, \rho)$ is

$$\mathbf{y}(\mathbf{x},\boldsymbol{\rho}) = \mathbf{e}^{-2\mathbf{x}} + \left(\frac{2\boldsymbol{\rho}}{1+\boldsymbol{\rho}}\right)\mathbf{A}_{1} + \left(\frac{2\boldsymbol{\rho}}{1+\boldsymbol{\rho}}\right)^{2}\mathbf{A}_{2} + \left(\frac{2\boldsymbol{\rho}}{1+\boldsymbol{\rho}}\right)^{3}\mathbf{A}_{3}$$
(13)

where

$$A_{1} = xe^{-x} [I_{0}(x) - I_{1}(x)],$$
$$A_{2} = \frac{A_{1}}{2} - \frac{x}{2}e^{-2x},$$

and

$$A_3 = -\frac{x}{6}e^{-x}[(2x - 3)I_0(x) - (2x - 2)I_1(x)] - \frac{x}{2}e^{-2x}.$$

The slowest rate of convergence of equation (13) occurs for large x. In this case, the coefficients A_n approach

$$A_{n} \sim \frac{1}{2^{n}\sqrt{2\pi x}}$$
(14)

and successive coefficients decrease by the factor two. For smaller x, the decrease is faster. Thus, a worst case estimate of the accuracy of equation (13) can be obtained by substituting equation (14) into (13) to obtain

$$y = e^{-2x} + \frac{1}{\sqrt{2\pi x}} [\alpha + \alpha^2 + \alpha^3 + \alpha^4 + \cdots],$$

-10-

where $\alpha = \rho/(1 + \rho)$. If we retain only terms through α^3 in the sum, the absolute error of truncation is

$$\epsilon = \alpha^4 + \alpha^5 + \cdots = \frac{\alpha^4}{1 - \alpha}.$$

The largest α that can be tolerated for a maximum relative error of 10⁻⁴ (in the sum) is given by

$$(\alpha + \alpha^2 + \alpha^3) \ge 10^4 \frac{\alpha^4}{1 - \alpha}$$

for which $\alpha \le 0.047$ ($\rho \le 0.045$) is the solution. Similar analyses show that the α^3 term may be neglected for $\rho \le 0.01$, and the α^2 term neglected when $\rho \le 0.0001$.

Computation of the I_0 and I_1 Bessel functions required to evaluate A_1 , A_2 , and A_3 of equation (13) are handled with rational approximations⁽⁴⁾ that provide sufficient accuracy for $1 \le x \le 20$. Beyond $x \simeq 20$, these rational approximations are not accurate enough to yield the difference $I_0 - I_1$ to sufficient accuracy. For $x \ge 20$, use is made of the asymptotic expansion

$$y = e^{-2x} + \frac{1}{\sqrt{2\pi x}} \left[1 + \frac{3}{8x} + \frac{45}{128x^2} \right] \frac{\alpha}{2} \left(1 + \frac{\alpha}{2} + 0.266\alpha^2 \right).$$
(15)

Up to the a^2 term, this expansion is obtained from equation (13) and the asymptotic expansions⁽⁴⁾ for I₀ and I₁. The coefficient 0.266 on the a^3 term was determined empirically so as to yield 0.01% accuracy for all $x \ge 20$ and $\rho \le 0.05$.

C. EXPANSION FOR LARGE ρ (REGION 3)

The expansion for this case is obtained exactly as for the small ρ expansion, except that the Taylor expansion is made about $\alpha = 1$. The result to third order in ρ^{-1} is ⁽³⁾

$$y(x, \rho) = 1 - \left(\frac{2}{1+\rho}\right)B_1 + \left(\frac{2}{1+\rho}\right)^2 B_2 - \left(\frac{2}{1+\rho}\right)^3 B_3$$
 (16)

where

and

$$B_{1} = xe^{-x}[I_{0}(x) + I_{1}(x)],$$
$$B_{2} = \frac{x}{2} - \frac{B_{1}}{2},$$

$$B_{3} = \frac{x}{6}e^{-x}[(2x + 3)I_{0}(x) + (2x + 2)I_{1}(x)] - \frac{x}{2}.$$

As for the small ρ expansion, the worst case of convergence of equation (16) occurs for large x. In this case, the first four coefficients approach

$$B_{1} \sim \frac{1}{\sqrt{\pi}} (2x)^{1/2},$$

$$B_{2} = \frac{1}{4} (2x),$$

$$B_{3} \sim \frac{1}{6\sqrt{\pi}} (2x)^{3/2},$$

and

$$B_4 = \frac{1}{32} (2x)^2$$
.

Substitution of these coefficients into equation (16) yields

$$y \sim 1 - \frac{2\alpha^{1/2}}{\sqrt{\pi}} + \alpha - \frac{4}{3\sqrt{\pi}} \alpha^{3/2} + \frac{\alpha^2}{2} - \ldots$$

where $\alpha = 2x/(\rho + 1)^2$. Since the series alternates in sign, we can be assured that the maximum absolute error incurred by truncating equation (16) at the cubed term is less than or equal to $\alpha^2/2$. The condition for relative error less than 0.01% is

$$\frac{\alpha^2}{2} \leq 10^4 \left[1 - \frac{2\alpha^{1/2}}{\sqrt{\pi}} + \alpha - \frac{8\alpha^{3/2}}{\sqrt{\pi}} \right]$$

from which we obtain $\alpha \leq 0.01328$ or

$$\rho \ge 12.3 x^{1/2} - 1.$$

In addition, the third; second; and first-order terms of equation (16) may be neglected, respectively, when

$$\rho \ge 28.2x^{1/2} - 1,$$

 $\rho \ge 142x^{1/2} - 1,$

and

$$\rho \ge 16000 x^{1/2} - 1$$
.

1 13

As in the small ρ approximation, the functions I_0 and I_1 required to compute the B coefficients are obtained from accurate rational approximations.⁽⁴⁾

D. EXPANSION FOR LARGE x (REGION 4)

Substitution of the asymptotic expansions⁽⁴⁾ for $I_n(x)$ into equation (9) and a collection of terms on like powers of x yields

$$y(x, \rho) \sim \frac{1}{\sqrt{2\pi x}} \left[\rho + \frac{F(\rho)}{8x} + \frac{G(\rho)}{2!(8x)^2} + \cdots \right]$$
 (17)

where

$$\begin{split} \mathbf{F}(\rho) &= \left(\frac{\rho+1}{2}\right)^3 [1 - 9\alpha - 9\alpha^2 + \alpha^3],\\ \mathbf{G}(\rho) &= \left(\frac{\rho+1}{2}\right)^5 3 [3 - 25\alpha + 150\alpha^2 + 150\alpha^3 - 25\alpha^4 + 3\alpha^5], \end{split}$$

and $\alpha = (\rho - 1)/(\rho + 1)$. For $\rho \ge 1.225$, $F(\rho)$ is negative and decreases monatonically to $F = -2\rho^3$ as $\rho \rightarrow \infty$. Similarly, $G(\rho)$ is positive and increases monatonically to $G = 24\rho^5$ as $\rho \rightarrow \infty$. The next higher coefficient approaches $H = -720\rho^7$ for large ρ . Thus, the worst case of convergence for $\rho \ge 1.225$ appears to the alternating series

$$\mathbf{y} \longrightarrow \frac{\rho}{\sqrt{2\pi\mathbf{x}}} \left[1 - \frac{1}{2} \left(\frac{\rho^2}{2\mathbf{x}} \right) + \frac{3}{4} \left(\frac{\rho^2}{2\mathbf{x}} \right)^2 - \frac{15}{8} \left(\frac{\rho^2}{2\mathbf{x}} \right)^3 + \cdots \right].$$

The absolute error of truncation at the x^{-2} term is less than $15(\rho^2/2x)^3/8$, and the condition for a relative accuracy of less than 0.01% is

-14-

$$\frac{15}{8} \left(\frac{p^2}{2x} \right)^3 \leq 10^{-4} \left[1 - \frac{1}{2} \left(\frac{p^2}{2x} \right) + \frac{3}{4} \left(\frac{p^2}{2x} \right)^2 \right],$$

which yields the condition $\rho^2/2x \le 0.0375$ or

$$x \ge 13.3\rho^2$$
. (18)

Comparisons of the results of equation (17) with exact calculations⁽²⁾ confirm this bound by predicting the less stringent bound $x \ge 7.25\rho^{2.22}$ -for $20 \le x \le 10^4$. In addition, these comparisons show that x must be greater than ~20 in order to obtain the desired accuracy of 0.01%. This result is consistent with equation (18) and the lower bound $\rho = 1.225$ set on ρ for this analysis.

For $\rho < 1.225$, the preceding error analysis does not apply because the series is not an alternating series. Here, comparisons with exact calculations confirm that equation (17) is accurate to the desired accuracy for $x \ge 20$ and ρ down to at least the upper bound ($\rho = 0.045$) of Region 2.

E. EXPANSION FOR LARGE x AND ρ (REGION 5)

The large x expansion equation (17) is restricted by the condition $\rho \le 0.274x^{1/2}$, whereas the large ρ expansion equation (16) is restricted by $\rho \ge 12.3x^{1/2}-1$. The following method provides a means of computation in this excluded region. In equation (1a), we make the transformation

$$u = \frac{1}{\sqrt{1 + \rho^2 z^2}}$$

to obtain

$$y(x, \rho) = \frac{2\rho}{\pi} \int_0^1 \frac{f(u)}{\sqrt{1 - u^2}} du$$
 (19)

where

$$f(u) = \frac{e^{-2xu^2}}{1 + u^2(\rho^2 - 1)}.$$

When x is large and $\rho > 1$, f(u) peaks sharply at u = 0, and most of the contribution to equation (19) will come from this region. This phenomenon suggests that not much error will result if we set the upper limit of integration to infinity and expand the square root term of equation (19) in a power series. This latter expansion is

$$\frac{1}{\sqrt{1-u^2}} = \sum_{n=0}^{\infty} B_n u^{2n}.$$
 (20)

The expansion coefficient B_n is

$$B_n = \frac{(2n)!}{2^{2n}(n!)^2}$$

which, for computational purposes, can be written in the recurrence form

$$B_n = \left(\frac{2n-1}{2n}\right) B_{n-1} \tag{21}$$

with

$$B_0 = 1.$$

Substitution of equation (20) into (19), an interchange of the orders of integration and summation, and setting the upper limit of integration to infinity yields

-16-

$$y(x, \rho) = \frac{2\rho}{\pi} \sum_{n=0}^{\infty} B_n \int_0^{\infty} \frac{u^{2n}e^{-2xu^2}}{1+u^2(\rho^2-1)} du.$$

With the transformation $z = u\sqrt{\rho^2 - 1}$, this becomes

$$y(x, \rho) = \rho \sum_{n=0}^{\infty} \frac{B_n F_n(\alpha)}{(\rho^2 - 1)^{n+1/2}},$$
 (22)

where

$$F_{n}(\alpha) = \frac{2}{\pi} \int_{0}^{\infty} \frac{z^{2n} e^{-\alpha^{2} z^{2}}}{1 + z^{2}} dz$$
(23)

and

$$\alpha = \sqrt{\frac{2x}{\rho^2 - 1}}.$$

A recurrence relation can be obtained for $F_n(\alpha)$ as follows. Differentiate equation (23) with respect to α to obtain

$$F_{n}(\alpha) = -\frac{1}{2\alpha} \frac{dF_{n-1}(\alpha)}{d\alpha}.$$
 (24)

From the solution (5)

$$F_0(\alpha) = e^{\alpha^2} [1 - erf(\alpha)], \qquad (25)$$

-17-

equation (24) can be used to evaluate the first few $F_n(\alpha)$ functions, from which, by induction, we can obtain either the explicit solution

$$F_{n}(\alpha) = (-1)^{n} \left[F_{0}(\alpha) \times \frac{2}{\sqrt{\pi}} \sum_{i=1}^{n} (-1)^{i} \frac{(2i-1)(2i-3) \cdot \cdot \cdot 5 \cdot 3 \cdot 1}{2^{i} \alpha^{(2i-1)}(2i-1)} \right]$$

or the recurrence relations

$$F_{n}(\alpha) = C_{n}(\alpha) - F_{n-1}(\alpha)$$
(26)

and

 $C_{n}(\alpha) = \frac{2n-3}{2\alpha^{2}}C_{n-1}(\alpha)$

with

$$C_1(\alpha) = \frac{1}{\sqrt{\pi}\alpha}.$$

In application, $y(x, \rho)$ is computed from equation (22) with B_n evaluated by equation (21) and $F_n(\alpha)$ by equations (25) and (26).

The accuracy of this expansion was determined by comparison with exact calculations. In the region $x \ge 5$ and $\rho \ge \sqrt{2}$, the approximation yields the desired accuracy with only a few terms in the expansion. Near x = 5, $\rho = \sqrt{2}$, about ten terms are required, whereas for $x \ge 10^4$ or $\rho \ge 2 \times 10^3$ (and $x \ge 1$), only the first term of equation (22) is required, i.e.,

$$y \sim e^{\alpha^2} [1 - erf\alpha].$$

-18-

In practice, the sum is terminated when the relative term contribution to the accumulated sum is less than 10^{-6} .

Computation of the function $F_0(\alpha)$ defined by equation (25) is made to a relative accuracy of 5×10^{-6} by using the approximation of Matta and Reichel.⁽⁶⁾

F. INTERMEDIATE x AND ρ (REGION 6)

For intermediate values of x and ρ , none of the methods considered to now can yield the accuracy desired in $y(x, \rho)$. For this excluded region (Region 6), we resort to the Bessel function expansion equation (9)

$$y(x, \rho) = e^{-x} \left[I_0(x) + 2 \sum_{n=1}^{N} \left(\frac{\rho - 1}{\rho + 1} \right)^n I_n(x) \right].$$
 (27)

Although this expression (with $N \rightarrow \infty$) is an exact solution of equation (1), its use for efficient computation is limited, especially for large x, unless ρ is very close to unity. For large x, the functions $I_n(x)$ fall off very slowly with n, and, consequently, a very large number of terms must be included in the summation. The number of terms (N) that must be included within Region 6 for a relative accuracy of 0.01% are tabulated in Table 1. Even for x only as large as 20, as many as 27 terms must be included for the desired accuracy.

In practice, the number of terms used in the expansion is determined by the worst case of $\rho \sim 0.05$, for which

$$N \simeq 8x^{0.393}.$$

The Bessel functions $I_n(x)$ are computed with a backward recurrence algorithm⁽⁷⁾ initiated at order

$$M = 14x^{0.307}$$

to ensure 10^{-4} relative accuracy in I_n(x) through n = N.

-19-

	x					
ρ	1	2	5	10	20	
20		9	13			
10	7	9	12			
5	7	8	11			
2	6	7	8	10	11	
1	1	1	1	1	1	
0.5	6	7	9	10	11	
0.2	7	9	12	15	19	
0.1	8	10	14	18	23	
0.05	8	10	15	19	26	

Table 1. Number of terms required in Bessel function expansion of $y(x, \rho)$ for 0.01% accuracy.

REFERENCES

- G. H. Lindquist and F. S. Simmons, J. Quant. Spectros. Radiat. Transfer <u>12</u>, 807 (1972).
- 2. S. J. Young, J. Quant. Spectros. Radiat. Transfer 15, 483 (1975).
- 3. D. D. Cobb, J. Quant. Spectros. Radiat. Transfer 16, 919 (1976).
- 4. M. Abramowitz and I. A. Stegun, <u>Handbook of Mathematical</u> Functions, National Bureau of Standards, Washington, D.C. (1972).
- 5. I. S. Gradshteyn and I. M. Ryzhik, <u>Tables of Integrals, Series</u>, and Products, Academic Press, Now York (1965).
- 6. F. Matta and A. Reichel, Math. Comp. 25, 339 (1971).
- 7. Y. I. Luke, <u>Mathematical Functions and their Approximations</u>, Academic Press, New York (1975).

LABORATORY OPERATIONS

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

<u>Aerophysics Laboratory</u>: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas lasers.

<u>Chemistry and Physics Laboratory</u>: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocket plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, lubrication and surface phenomena, photosensitive materials and sensors, high precision laser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semiconducting, superconducting, and crystal device physics, optical and acoustical imaging: atmospheric pollution; millimeter wave and far-infrared technology.

<u>Materials Sciences Laboratory</u>: Development of new materials: metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields: space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere: the effects of optical, electromagnetic, and particulate radiations in space on space systems.

> THE AEROSPACE CORPORATION El Segundo, California

• • •