

AD-A071 126

NAVAL HEALTH RESEARCH CENTER SAN DIEGO CA

F/6 6/5

A MODEL FOR MEASURING THE CONSISTENCY OF DIAGNOSTIC CLASSIFICAT--ETC(U)

UNCLASSIFIED

JUL 76 S FICHMAN, D EDWARDS

NAVHLTHRSCHC-76-47

ML

| OF |
AD
A071126



END
DATE
FILMED
8 -79
DDC

AD A 071126

LEVEL ^H



A MODEL FOR MEASURING THE CONSISTENCY OF DIAGNOSTIC CLASSIFICATION


S. FICHMAN

D. EDWARDS

See 1473 in facts

REPORT NO. 76-47

DDC FILE COPY



NAVAL HEALTH RESEARCH CENTER

Naval Health Research Center
SAN DIEGO, CALIFORNIA 92152
San Diego, CA 92138

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

79 07 11 033

A Model for Measuring the Consistency of Diagnostic Classification¹

Susan Fichman and Darrel Edwards

Naval Health Research Center

San Diego, California

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced Justification	<input type="checkbox"/>
By _____	
Distribution/ _____	
Availability Codes	
Dist.	Avail and/or special
A	

¹Report No. 76-47. The views presented in this paper are those of the authors. No endorsement by the Department of the Navy has been given or should be inferred.

A Model for Measuring the Consistency of Diagnostic Classification¹

Susan Fichman and Darrel Edwards

One goal of clinical practice is to develop a rational system for making and using a diagnosis in treatment. During clinical practice, clinicians develop decision making rules for evaluating their clients. These rules will guide future interactions. If the rules are sound, the clinician will be able to successfully deal with his clients. The measurement of the actual accessibility of these cognitive rules to the trainee in managing clinical cases is difficult. When experimenters rely on observable behaviors to evaluate clinical functioning, they ignore the fact that different clinical styles may be equally effective in dealing with a patient. Observable clinical behaviors may vary widely. There are alternative decision making processes which lead to successful treatment outcomes (Edwards, Gunderson, Brown, and Taylor, 1973).

This study examined the conceptual process of diagnosis as used by psychiatrists in their clinical practice. Psychiatrists evaluated their patients using dimensions unique to their clinical philosophy. These evaluations were analyzed to produce each psychiatrist's conceptual scheme as it relates to the diagnosis of his patients. The purpose of this study was to demonstrate that the diagnostic structure of an individual clinician can be objectively measured, producing a method for evaluating consistency, clarity, or other dimensions of the clinical decision.

Method

Procedure

A grid method (Bannister and Mair, 1968) was used to collect data from 13 psychiatrists at two Navy inpatient psychiatric services. The doctors had a mean time in practice post-residency of 2.0 years with a range of 1 to 12 years.

Grids were administered in small groups of no more than five clinicians at one time. The clinicians were asked to compile a list of 16 patients from their caseloads: four psychotics, three neurotics, three personality disorders, three situational maladjustments, and three alcoholics. Adjustments in the distribution were made to accommodate differences in individual caseloads where a clinician was not able to fill all classes. Adjustments were made by having the therapists substitute patients from other diagnostic categories when they had exhausted their cases from the requested category. At least two patients from each diagnostic group were required.

Each clinician was instructed to evaluate his patients on 16 clinical dimensions which the therapist chose as important in his practice. Patients were scored with an "X" to indicate whether that patient possessed a specific characteristic. This technique produced a 16 dimensional profile for each patient. The result was a grid of 16 patients by 16 dimensions (Figure 1).

Insert Figure 1 about here

Data Analysis

The intention of the analysis was to determine if a clinician exhibited a conceptual scheme which indicated that similarly diagnosed cases were treated with some consistency. Each grid was analyzed individually with the following

Figure 1
CONCEPTUAL GRID

		Name: _____ Date: _____	
		CONSTRUCT	CONTRAST
PORT NUMBER			
1	Manic depressive psychosis		
2	Depressive neurosis		
3	Schizoid personality		
4	Adult situational stress		
5	Acute Alcoholism		
6	Acute paranoid reaction		
7	Hysterical neurosis		
8	Antisocial personality		
9	Adult situational stress		
10	Chronic alcoholism		
11	Latent schizophrenia		
12	Obsessive compulsive neurosis		
13	Explosive personality		
14	Adult situational stress		
15	Acute alcoholism		
16	Schizophrenia paranoid type		
1		Honest	Dishonest
2		Mature	Immature
3		Insight	No insight
4		Acceptance of illness	Denial of illness
5		Considerate	Egocentric
6		Independent	Dependence
7		Self-conscious	Self-assured
8		Hyper-active	Psycho-motor retarded
9		Many friends	Improvised interpersonal relationships
10		Oral Character	Not oral character
11		Obsessive Compulsive	Not obsessive compulsive
12		Manipulative	Not manipulative
13		Poor impulse control	Good impulse control
14		Poor ego image	Good ego image
15		Covert hostility	No covert hostility
16		Lack of energy	Adequate energy

procedure. Every patient's 16-dimensional profile was compared with every other patient's profile for each clinician's grid. Paired profiles were examined to determine the number of matches in dimensions ascribed to the pairs of patients. A match occurred when two patients were rated as having the same attribute in common or when both patients did not possess a given attribute. Profiles did not match when one patient was rated as having an attribute and the other patient was not. The number of matches was a measure of similarity between patients. The higher the total, the greater the perceived similarity.

The similarity totals produced a new patient profile listing the similarity between that patient and every other patient. This information was recorded in a 16 by 16 matrix. Intercolumnar correlations were computed, i.e., each patient's similarity profile was correlated with every other patient's similarity profile. A cluster analysis (Johnson, 1967) was performed on these correlations. Patients whose profiles were highly correlated were combined into clusters. The program scanned the matrix of correlations and selected the highest correlation. The similarity score profiles for the two most highly correlated patients were collapsed to form a single profile. Correlations were recomputed to form a new matrix containing one fewer pair of correlations and the new correlation matrix was scanned to again select the highest correlation. This process was stopped after the tenth reiteration to maximize the possibility of grouping patients into five clusters representing the five diagnostic categories.

A cluster analysis was performed on each clinician's data. Four representative cluster patterns are presented in Figures 2-5. The patient's diagnosis is represented by an abbreviation (psychosis (P), neurosis (N), situational

maladjustment (SM), personality disorder (PD), and alcoholism (A), the patient's number is a subscript. The relative lengths of the "branches" represent the correlation for the cluster; longer distances represent lower correlations.

Results and Discussion

Clinician I. The first cluster ($r=1.0$) was a pair of situational maladjustments. At later stages, clusters contained combinations of unlike diagnoses. One alcoholic defined a single member case. Final clusters showed some appropriate pairings of diagnoses but little overall structure for diagnostic clarity.




Figure 2. Diagnostic Structure for Clinician I.

Clinician 2. The initial clusters ($r > .90$) showed diagnostic clarity. The first contained two alcoholics, the second contained two psychotics, the third contained two personality disorders, the fourth contained two psychotics, and the fifth contained one psychotic and one neurotic. Later clusterings produced groups of mixed diagnoses. Two personality disorders were single member cases that did not combine with any other cases. This diagnostic structure began with pairings of similar diagnoses but these pairings did not generalize to include more members of the same diagnostic class.




Figure 3. Diagnostic Structure for Clinician 2.

Clinician 3. At early stages ($\bar{r} > .80$), there is evidence for diagnostic clarity with one cluster containing two psychotics, a second cluster containing two neurotics, a third containing a psychotic and an alcoholic, and a fourth containing a psychotic and a personality disorder. Later stages ($\bar{r} > .60$) showed increased diagnostic clarity. One cluster contained two of four psychotics, a second cluster contained all neurotics, a third cluster contained all situational maladjustments and one personality disorder, a fourth contained a psychotic and an alcoholic and a fifth contained a psychotic and a personality disorder. One alcoholic and one personality disorder defined single member classes.




Figure 4. Diagnostic Structures for Clinician 3.

Clinician 4. Initial clusters ($r > .90$) showed a clear diagnostic strategy. Of the five clusters produced, only one cluster contained unmatched diagnoses. Later clusters ($r > .970$) showed continued diagnostic clarity. One cluster contained all alcoholics, a second cluster contained one psychotic and one situational maladjustment, a third cluster contained all neurotics, a fourth cluster contained two of three situational maladjustments, and a fifth cluster contained all personality disorders. One psychotic defined a single member case. This diagnostic structure used distinct categories for every diagnosis but psychosis.




Figure 5. Diagnostic Structure for Clinician 4.

Summary of all Clinicians

Individual cluster patterns were unique in composition, but there were some similarities in structure across clinicians. The four examples described above represent the four types of structure in diagnosis: (1) General inconsistency; (2) Immediate consistency followed by inconsistency; (3) Developing consistency; and (4) General consistency. Cluster analysis of each profile showed the diagnostic structure of each clinician. A summary of the profiles had the following characteristics: (a) four clinicians exhibited a consistency in conceptually sorting all patients; (b) six clinicians had a consistent structure for one or two of the diagnostic classes with each diagnosis showing some consistency for at least one doctor in the sample; and (c) three clinicians appeared to have no conceptually clear way to manage their patient sample.

Clarity of diagnostic structure appeared often at the concrete level of association and continued to build as associations became more abstract. Most doctors appeared to have some systematic structure associated with diagnosis. The complexity of these structures ranged from single concrete pairings to more complex structures handling one diagnostic category to those structures that dealt efficiently with a wide range of diagnostic classes. Individual diagnostic systems differed in the level of abstraction associated with diagnostic clarity and confusion. These differences were objectified by the grid method used in this research. Overall, the results defined the level of functioning for each clinician in diagnosing patients.

Comment

Although variations in clinical cases and individual structures are expected, some consistency in diagnostic practices is also anticipated. Diagnostic similarity is expected to be greater within diagnostic categories than across

diagnostic categories, even if the diagnostic processes were different for each clinician. Most doctors in this study demonstrated clear, consistent conceptual structure for the diagnostic categories.

This study clearly demonstrates that clinical decision making processes can be documented. Documentation of a subjective process would permit comparisons between doctors, between doctors and an idealized standard, and between teachers and students in clinical training settings. Objectifying diagnostic decisions allows a clearer communication between supervisor or teacher and trainee about clinical rules and subjective processes (Ivery, et al., 1968). Previous use of introspective processes and behavioral descriptions of diagnosis have failed to delimit the dynamics of the process (Hansen and Barker, 1964).

Information on decision making would be useful in a training situation. In an ongoing training program, both the instructor and the student could complete grids on patients familiar to both. The instructor could compare the student's structure and his own and evaluate differences to determine where the student might need additional feedback. Thus, the instructor could spot student weaknesses and direct his attention toward these areas. Repeated examination, comparing grids produced by students and teachers at additional checkpoints, would show the students' progress.

Program evaluation may also be measured with these procedures. Selected observation points might be chosen and students evaluated on the critical dimensions to be taught in a training program. The results could be compared to benchmarks for the program. Effectiveness of training methods, trainers, or program material may be reflected against criteria developed by the documentation methods described in this report (Carkhuff & Truax, 1965). This technique provides a method for measurement which is relatively easy to use and shows possibilities for criteria development. Incorporating cognitive information with current behavioral observations will give a more complete picture of clinical functioning.

References

- Carkhuff, R. & Truax, C. Training in counseling and psychotherapy: An evaluation of an integrated didactic and experiential approach. Journal of Consulting Psychology, 1965, 29, 333-336.
- Edwards, D., Gunderson, E. K. E., Brown, D. C., & Taylor, W. R. Clinical Decisions and outcomes in a Navy psychiatric service. Military Medicine, 1973, 138, 430-432.
- Hansen, J. C. & Barker, E. N. Experiencing and the supervisory relationship. Journal of Counseling Psychology, 1964, 11, 107-111.
- Ivery, A. E., Normington, C. J., Miller, D. C., Merrill, W. H., & Haase, R. F. Microcounseling and attending behavior: An approach to prepracticum counselor training. Journal of Counseling Psychology, 1968, 15 (Monogr. Suppl. 5)
- Johnson, S. C. Hierarchical clustering schemes. Psychometrika, 1967, 32, 241-254.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 76-47	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Model for Measuring the Consistency of Diagnostic Classification.		5. TYPE OF REPORT & PERIOD COVERED final
7. AUTHOR(s) Susan Fichman and Darrel Edwards		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Health Research Center P.O. Box 85122 San Diego, CA 92138		8. CONTRACT OR GRANT NUMBER(s) M4305.07-3013DXB5
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command Bethesda, Maryland 20014		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS M4305.07-3013DXB5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Bureau of Medicine and Surgery Department of Navy Washington, D. C. 20375		12. REPORT DATE July 1976
		13. NUMBER OF PAGES 10
		15. SECURITY CLASS. (of this report) unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited. 9 Final rept. (12) 15 p.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Approved for public release; distribution unlimited. 14		
18. SUPPLEMENTARY NOTES NAVHLTHRSCHC-76-47		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Diagnosis, clinical training, cluster analysis, grid method, cognitive process		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Appearance of this material in the professional literature is in support of the Chief of Naval Material's program for Military-Civilian Technology Transfer and Cooperative Development. Psychiatric screening, evaluations, and treatment strategies represent 300,000 manhours lost to the Navy each year. With this heavy responsibility, there is a great need for good methods to promote uniform quality practices in diagnosis and treatment. The model presented can be used to assure high quality screening, evaluation, and treatment in the naval		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

391 642

elt

unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

psychiatric environment. Evaluation of clinical training is a complex task. Traditional methods have relied on directly observable criteria and often produced disappointing results. The real object of clinical training is to develop good decision making procedures. This paper presents a methodology for examining the conceptual process of diagnosis. Thirteen psychiatrists evaluated their own patients using a grid method. Each doctor's grid was examined individually using cluster analysis. The Doctor's cluster structures were evaluated to determine whether the psychiatrists were systematically using diagnosis to classify the patients. Application of this technique to the training setting was presented.

unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)