



REPORT DOCUMENTATION PAGE	4 READ INSTRUCTIONS
1. REPORT HUMBER	ION NO. RECIPIENT'S CATALOG NUMBER
AFUSR-TR- 80-0929 AD 40	89650
4. TITLE (and Sublilio) AUS 5022	5. TYPE OF REPORT & PERIOD COVERS
SCHEME-INDEPENDENT STABILITY CRITERIA FOR	Interim
DIFFERNCE APPROXIMATIONS OF HYPERBOLIC INITIA	AL - 6. PERFORMING ORG. REPORT NUMBER
-BOUNDARY VALUE PROBLEMS. II 7. AUTHOR(2)	THE SONTRACT OR GRANT NUMBER(S)
Moshe Goldberg and Eitan Tadmor	AFOSR 76-3046
9. PERFORMING ORGANIZATION NAME AND ADDRES	AREA & WORK UNIT NUMBERS
University of California	2304/A3
Los Angeles, CA 90024	61102F
1. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE August 1980
Bolling AFB, Wasington, D.C. 20332	13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling)	49 Office) 15. SECURITY CLASS. (of this report)
	UNCLASSIFIED
	15. DECLASSIFICATION/DOWNGRADING
	SCHEDULE
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, if diff	Inlimited.
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff	terent from Report) DTIC SEP 2 9 1980
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, 18 diff	rent from Report) SEP 2 9 1980
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, if diff	Indimited. Derent from Report) SEP 2 9 1980 A
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff 18. SUPPLEMENTARY NOTES	Areant from Report)
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block Hunarholic Pontial Differential Equations : Fi	rement from Report) Terent from Report) SEP 2 9 1980 A
Approved for public release; distribution u Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse eide if necessary and identify by block Hyperbolic Partial Differential Equations; Fir Boundary Conditions; Stability of Difference S	rement from Report) Terent from Report SEP 2 9 1980 A Inte Difference Schemes; Schemes.
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessary and identify by block Hyperbolic Partial Differential Equations; Fin Boundary Conditions; Stability of Difference S	Indimited. Terent from Report) DIC SEP 2 9 1980 A A Inte Difference Schemes; Schemes.
Approved for public release; distribution u Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 diff 18. SUPPLEMENTARY NOTES 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block Hyperbolic Partial Differential Equations; Fin Boundary Conditions; Stability of Difference S 10. ABTRACT (Continue on reverse side if necessary and identify by block Convenient stability criteria are obtained	Intervent from Report) Formula (from Report) Difference Schemes; Schemes.
Approved for public release; distribution u Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse eide if necessary and identify by block Hyperbolic Partial Differential Equations; Fin Boundary Conditions; Stability of Difference S 10. ADSTRACT (Continue on reverse eide if necessary and identify by block Convenient stability criteria are obtained hyperbolic initial-boundary value problems. The	rumber) number) number) number) nite Difference Schemes; Schemes.
Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse eide if necessary and identify by block Hyperbolic Partial Differential Equations; Fin Boundary Conditions; Stability of Difference S 10. AMSTRACT (Continue on reverse eide if necessary and identify by block Convenient stability criteria are obtained hyperbolic initial-boundary value problems. The arbitrary basic schemes and a wide class of boundary conditions of the outflow mart of the basis	Interview Interview
Approved for public release; distribution u Approved for public release; distribution u T. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if diff supplementary notes KEY WORDS (Continue on reverse side if necessary and identify by block Hyperbolic Partial Differential Equations; Fin Boundary Conditions; Stability of Difference S Convenient stability criteria are obtained hyperbolic initial-boundary value problems. The arbitrary basic schemes and a wide class of boundary conditions of the basic scheme. The results of the provent of the basic scheme. The results of the basic scheme. The results of the basic scheme.	International consist of undary conditions and are easily imply that a number of well
Approved for public release; distribution u Approved for public release; distribution u 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 diff 18. SUPPLEMENTARY NOTES 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block Hyperbolic Partial Differential Equations; Fin Boundary Conditions; Stability of Difference S 10. ABSTRACT (Continue on reverse side if necessary and identify by block Convenient stability criteria are obtained hyperbolic initial-boundary value problems. The arbitrary basic schemes and a wide class of bou are given in terms of the outflow part of the binder known boundary treatments, when used in combined schemes, always maintain stability. Consequent	Internations consist of undary conditions and are easily imply that a number of well ation with arbitrary stable basic
Approved for public release; distribution u Approved for public release; distribution u TO DISTRIBUTION STATEMENT (of the ebetrect entered in Block 20, 11 diff SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES Supplementary notes Approved for public reverse elde 11 necessary and identify by block Hyperbolic Partial Differential Equations; Fin Boundary Conditions; Stability of Difference S Convenient stability criteria are obtained hyperbolic initial-boundary value problems. The arbitrary basic schemes and a wide class of bou are given in terms of the outflow part of the h independent of the basic scheme. The results of known boundary treatments, when used in combined schemes, always maintain stability. Consequent in recent literature are generalized.	International consist of undary conditions and are easily imply that a number of well ation with arbitrary stable basic thy, many special cases studied
Approved for public release; distribution u Approved for public release; distribution u T. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 diff SUPPLEMENTARY NOTES Supplementary NOTES Supplementary notes Astronomical Equations; Fin Boundary Conditions; Stability of Difference S Convenient stability criteria are obtained hyperbolic initial-boundary value problems. The arbitrary basic schemes and a wide class of boundary diversion of the outflow part of the findependent of the basic scheme. The results of known boundary treatments, when used in combined schemes, always maintain stability. Consequent in recent literature are generalized.	number) in for difference approximations to he approximations consist of undary conditions. The new criter boundary conditions and are easily imply that a number of well ation with arbitrary stable basic tly, many special cases studied

AFOSR-TR- 80-0929 SCHEME-INDEPENDENT STARILITY CRITERIA FOR DIFFERENCE APPROXIMATIONS OF HYPERBOLIC INITIAL-BOUNDARY VALUE PROBLEMS. II by AInterim rept. (16) 23741 Moshe Goldberg **Eitan Tadmor** Department of Mathematical Department of Mathematics Technion--Israel Institute of Sciences Tel Aviv University Technology Tel Aviv, Israel Haifa, Israel and and Department of Applied Mathematics Institute for the Interdisciplinary Applications of Algebra and California Institute of Technology Combinatorics Pasadena, California 91125 University of California Santa Barbara, California 93106 (1)/11-- Ir/ Mr. 10 / Geeld K. m. / Eiten / Thimer / 1. MICSR-76- Eq. 16, AICSR-77 9127 AMS (MOS) Subject Classification (1970). Primary 65M10; Secondary 65N10. "The research of the first author was sponsored in part by the Air Force Office of Scientific Research, Air Force System Command, United States Air Force Grants. No. 4505-76-3046 and AFOSR-79-0127.0 The research of the second author is part of his Ph.D. dissertation and has been sponsored in part by the Air Force Office of Scientific

072266

Research (NAM) through the European Office of Aerospace Research, Air Force System Command, United States Air Force, under Grant 76-3035.

203 pergad

ABSTRACT

Convenient stability criteria are obtained for difference approximations to initial-boundary value problems associated with the hyperbolic system $u_t = Au_x + Bu + f$ in the quarter plane $x \ge 0$, $t \ge 0$. The approximations consist of arbitrary basic schemes and a wide class of boundary conditions. The new criteria are given in terms of the outflow part of the boundary conditions and are independent of the basic scheme. The results easily imply that a number of well known boundary treatments, when used in combination with arbitrary stable basic schemes, always maintain stability. Consequently, many special cases studied in recent literature are generalized.

Accession For	this technical report has been reviewed and is
NTIS U.A. AL	a proved for public release IAW AFR 190-18 (7b).
Unampounted	A. D. BLOSE
Justification	
Бу	
Fistribution/	
the Stability Codes	
< dil and/or	
H	

and a second second

0. Introduction.

In this paper we extend the results of [2] to obtain easily checkable stability criteria for difference approximations of initial-boundary value problems associated with the linear hyperbolic differential system $u_t = Au_x + Bu + f$ in the quarter plane . $x \ge 0$, $t \ge 0$. The difference approximations, introduced in Section 1, consist of arbitrary basic schemes -- explicit or implicit, dissipative or unitary, two-level or multi-level -and boundary conditions of a rather general type.

The first step in our stability analysis is made in Section 2 where we prove that the approximation is stable if and only if the scalar outflow components of its principal part are stable. This reduces the global stability question to that of a scalar, homogeneous, outflow problem which thereafter becomes the main object of the paper.

Investigating the stability of the reduced problem our main results are restricted to the case where the boundary conditions are translatory, i.e., determined at all boundary points by the same coefficients. Such boundary conditions are commonly used in practice; and in particular, when the numerical boundary consists of a single point the boundary conditions are translatory by definition.

The main stability criteria for the translatory case, stated without proof in Section 3, are given essentially in terms of the boundary conditions. Such scheme-independent criteria elliminate the need to analyze the intricate and often

complicated interaction between the basic scheme and the boundary conditions; hence providing convenient alternatives to the well known stability criterion of Gustafsson, Kreiss and Sundström [3], which is the basis for our work.

As in [3], we assume that the basic scheme is stable for the pure Cauchy problem and that the approximation is solvable. Under these basic assumptions -- which are obviously necessary for stability -- we obtain, for example, in Theroems 3.3 and 3.4, that the reduced problem is stable if the (translatory) boundary conditions are solvable and satisfy the von Neumann condition as well as an additional simple inequality. If the basic scheme is unitary it is also required that the boundary conditions be dissipative.

Having the new stability criteria, we continue in Section 3 to study several examples. First, we reestablish the known fact that if the basic scheme is two-level and dissipative, then outflow boundary conditions determined by horizontal extrapolation always maintain stability. Surprisingly, we show that this result is false if the basic scheme is of more than two levels. Next, for arbitrary multi-level dissipative basic schemes we find that if the outlfow boundary conditions are generated, for example, by oblique extrapolation, by the Box-Scheme, or by the right-sided Euler scheme, then overall stability is assured. Finally, for basic schemes (dissipative or unitary) we show that overall stability holds if the outflow boundary conditions are determined by the right-sided explicit or implicit Euler schemes. These examples incorporate many special cases discussed in recent literature [1,2,3,4,5,6,9,10].

In Sections 4 and 5 we prove the results stated in Section 3.

It should be pointed out that there is no difficulty in extending our stability criteria to cases with two boundaries. In fact, if the corresponding left and right quarter-plane problems are stable, then by Theorem 5.4 of [3], the original two-boundary problem is stable as well.

Thanks are due to Bjorn Engquist and Stanley Osher for most helpful discussions.

1. The Difference Approximation

Consider the first order hyperbolic system of partial differential equations

(1.1a)
$$\partial u(x,t)/\partial t = A \partial u(x,t)/\partial x + B u(x,t) + f(x,t), \quad x \ge 0, t \ge 0,$$

where $u(x,t) = (u^{(1)}(x,t),...,u^{(n)}(x,t))$ is the vector of unknowns (prime denoting the transpose), $f(x,t) = (f^{(1)}(x,t),...,f^{(n)}(x,t))$ is a given vector, and A and B are fixed $n \times n$ matrices so that A is Hermitian and nonsingular. Without restriction we may assume that A is diagonal of the form

(1.2)
$$A = \begin{pmatrix} A^{II} & 0 \\ 0 & A^{II} \end{pmatrix}, \quad A^{II} < 0, \quad A^{III} > 0,$$

where A^{II} and A^{III} are of orders $l \times l$ and $(n-l) \times (n-l)$, respectively

The solution of (1.1) is uniquely determined if we prescribe initial values

(1.1b)
$$u(x,0) = \dot{v}(x), \quad x \ge 0,$$

and boundary conditions

فأفتعهم والمنازع والارتزار فلأحمص والانتفاق الالمولار المحارج فترعما

(1.1c)
$$y^{I}(0,t) = Sy^{II}(0,t) + g(t), \quad t \ge 0,$$

where S is a fixed $\ell \times (n-\ell)$ matrix, g(t) is a given ℓ -vector, and

(1.3)
$$u^{I} = (u^{(1)}, \dots, u^{(\ell)})', \quad u^{II} = (u^{(\ell+1)}, \dots, u^{(n)})'$$

is a partition of <u>u</u> into inflow and outflow unknowns, respectively, corresponding to the partition of A. In order to solve the initial-boundary value problem (1.1) by difference approximations we introduce a mesh size $h \equiv \Delta x > 0$, $k \equiv \Delta t > 0$, such that $\lambda \equiv h/k = \text{constant}$. Using the notation $y_{\nu}(t) = y(\nu h, t)$ we approximate (1.1a) by a consistent, two-sided, general multi-step basic scheme of the form

$$Q_{1,v_{v}}(t+k) = \sum_{\sigma=0}^{s} Q_{\sigma,v_{v}}(t-\sigma k) + kb_{v}(t), \qquad v = r,r+1,\ldots,$$

(1.4a)

$$Q_{\sigma} = \sum_{j=-r}^{\nu} A_{j\sigma} E^{j}, \quad E_{\nu} = \nu_{\nu+1}, \quad \sigma = -1, \dots, s ,$$

where the $n \times n$ matrices $A_{j\sigma}$ are polynomials in A and kB, and the n-vectors $b_{\nu}(t)$ depend smoothly on f(x,t) and its derivatives.

To solve (1.4a) uniquely, we provide initial values

(1.4b)
$$\underline{v}_{\nu}(\sigma \mathbf{k}) = \overset{\circ}{\underline{v}}_{\nu}(\sigma \mathbf{k}), \quad \sigma = 0, \dots, s, \quad \nu = 0, 1, \dots,$$

where in addition we must specify, at each time step $t = \sigma k \ge sk$, boundary values $v_{\mu}(t+k)$, $\mu = 0, \dots, r-1$. The required boundary values will be determined by two sets of boundary conditions, the first of which is obtained by taking the last $n-\ell$ components of general boundary conditions of the form

$$\mathbf{T}_{-1}^{(\mu)} \underbrace{\mathbf{y}}_{\mu}(\mathbf{t}+\mathbf{k}) = \sum_{\sigma=0}^{\mathbf{q}} \mathbf{T}_{\sigma}^{(\mu)} \underbrace{\mathbf{y}}_{\mu}(\mathbf{t}-\sigma\mathbf{k}) + \underbrace{\mathbf{k}}_{\sigma}_{\mu}(\mathbf{t}),$$

$$\mathbf{T}_{\sigma}^{(\mu)} = \sum_{j=0}^{\mathbf{m}} \mathbf{C}_{j\sigma}^{(\mu)} \mathbf{E}^{j}, \quad \mu = 0, \dots, \mathbf{r}-1, \quad \sigma = -1, \dots, q$$

1 and the second second second

where the matrixes $C_{j\sigma}^{(\mu)}$ are polynomials in A and kB, the $C_{j(-1)}^{(\mu)}$ are nonsingular, and the n-vectors $d_{\mu}(t)$ are functions of f(x,t), g(t) and their derivatives. If we put

$$\mathbf{c}_{\mathbf{j}\sigma}^{(\mu)} = \begin{pmatrix} \mathbf{c}_{\mathbf{j}\sigma}^{\mathbf{I}\,\mathbf{I}(\mu)} & \mathbf{c}_{\mathbf{j}\sigma}^{\mathbf{I}\,\mathbf{I}(\mu)} \\ \mathbf{c}_{\mathbf{j}\sigma}^{\mathbf{I}\,\mathbf{I}(\mu)} & \mathbf{c}_{\mathbf{j}\sigma}^{\mathbf{I}\,\mathbf{I}(\mu)} \end{pmatrix}, \quad \mathbf{x}_{\mu} = \begin{pmatrix} \mathbf{x}_{\mu}^{\mathbf{I}} \\ \mathbf{x}_{\mu} \\ \mathbf{x}_{\mu}^{\mathbf{I}} \end{pmatrix}, \quad \mathbf{d}_{\mu} = \begin{pmatrix} \mathbf{d}_{\mu}^{\mathbf{I}} \\ \mathbf{d}_{\mu} \\ \mathbf{d}_{\mu} \\ \mathbf{d}_{\mu} \end{pmatrix}$$

in accordance with the partitions of A and U in (1.2), (1.3), this set of conditions takes the form

(1.4c)
$$T_{-1}^{\Pi I(\mu)} \chi_{\mu}^{I}(t+k) + T_{-1}^{\Pi \Pi(\mu)} \chi_{\mu}^{\Pi}(t+k)$$

$$= \sum_{\sigma=0}^{q} [T_{\sigma}^{\Pi I(\mu)} \chi_{\mu}^{I}(t-\sigma k) + T_{\sigma}^{\Pi \Pi(\mu)} \chi_{\mu}^{\Pi}(t-\sigma k)] + k \sum_{\mu}^{\Pi} (t),$$

$$T_{\sigma}^{\Pi \alpha(\mu)} \equiv \sum_{j=0}^{m} C_{j\sigma}^{\Pi \alpha(\mu)} E^{j}, \quad \alpha = I, \Pi, \quad \mu = 0, \dots, r-1.$$

For the second set of boundary conditions we use the analytic condition

(1.4d)
$$y_0^{I}(t+k) = Sy_0^{II}(t+k) + g(t+k)$$

together with r-l additional conditions of the form

(1.4e)
$$y_{\mu}^{I}(t+k) = \sum_{j=1}^{\rho} [D_{j}^{II(\mu)} y_{j}^{I}(t+k) + D_{j}^{III(\mu)} y_{j}^{II}(t+k)] + k g_{\mu}^{I}(t),$$

 $\mu = 1, \dots, r-1,$

where the matrices $D_{j}^{II(\mu)}$ and $D_{j}^{II(\mu)}$ -- of orders $\ell \times \ell$ and $\ell \times (n-\ell)$, respectively -- are polynomials in the blocks $(A^{\alpha\alpha})^{-1}$ and $kB^{\alpha\beta}$, $\alpha,\beta = I, II$, of the matching partitions

$$A^{-1} = \begin{pmatrix} (A^{II})^{-1} & 0 \\ & & \\ 0 & (A^{IIII})^{-1} \end{pmatrix}, \quad kB = k \begin{pmatrix} B^{II} & B^{III} \\ & & \\ B^{III} & B^{IIII} \end{pmatrix},$$

so that $D_j^{II(\mu)}$ are homogeneous in B^{II} and B^{II} , and the *k*-vectors $g_{\mu}^{I}(t)$ are again functions of f(x,t), g(t) and their derivatives.

We remark that (1.4c,d,e) can be solved uniquely for the required boundary values $v_{\mu}(t + k)$, $\mu = 0, \ldots, r - 1$, in terms of neighboring values of y, at least for sufficiently small k. Indeed, since B introduces an O(k) perturbation of the matrix coefficients in (1.4c,e), it suffices to prove this statement for B = 0. But then, using the properties of $C_{j\sigma}^{(\mu)}$ and $D_{j}^{II(\mu)}$, it is not hard to see that $C_{j}^{III(\mu)} = D_{j}^{II(\mu)} = 0$ and that the $C_{j(-1)}^{III(\mu)}$ are nonsingular; hence (1.4c)uniquely determines the vectors $v_{\mu}^{II}(t + k)$, $\mu = r - 1, \ldots, 0$ (in that order) and substituting in (1.4d,e) we explicitly obtain $v_{\mu}^{I}(t + k)$, $\mu = 0, \ldots, r - 1$.

We also remark that while it is a standard matter to construct boundary conditions of the form (1.4c) to any degree of accuracy, the construction of (1.4e) is less obvious: For example, using (1.1) we find by induction on $j \ge 1$ that

(1.5)
$$\frac{\partial^{\mathbf{j}}}{\partial \mathbf{x}^{\mathbf{j}}} \underline{u}(\mathbf{x},t) = (\mathbf{L}_{t})^{\mathbf{j}} \underline{u}(\mathbf{x},t) - \chi_{\mathbf{j}}(\mathbf{x},t)$$

and

(1.6)
$$\frac{\partial^{\mathbf{j}}}{\partial t^{\mathbf{j}}} \mathfrak{u}(\mathbf{x},t) = (\mathbf{L}_{\mathbf{x}})^{\mathbf{j}} \mathfrak{u}(\mathbf{x},t) + \mathfrak{z}_{\mathbf{j}}(\mathbf{x},t),$$

where the operators L_{t}^{L} , and the vectors $\chi_{j}(x,t)$, $z_{j}(x,t)$ are given by

$$L_{t} = A^{-1} \left(\frac{\partial}{\partial t} - B \right), \quad \chi_{j}(x,t) = \sum_{i=0}^{j-1} \left(L_{t} \right)^{i} A^{-1} \frac{\partial^{j-i-1}}{\partial x^{j-i-1}} f(x,t)$$
$$L_{x} = A \frac{\partial}{\partial x} + B, \quad z_{j}(x,t) = \sum_{i=0}^{j-1} \left(L_{x} \right)^{i} \frac{\partial^{j-i-1}}{\partial t^{j-i-1}} f(x,t).$$

Now, if conditions of the form (1.4e) are required to ρ order of accuracy, we take a Taylor expansion of $u_{\mu}^{I}(t)$ and use (1.5) and (1.1c) to obtain

$$(1.7) \qquad \mathfrak{U}_{\mu}^{I}(t+k) = \sum_{j=0}^{\rho} \frac{(\mu h)^{j}}{j!} \frac{\partial^{j}}{\partial x^{j}} \mathfrak{U}^{I}(0,t+k) + O(h^{\rho+1}) \\ = \sum_{j=0}^{\rho} \frac{(\mu h)^{j}}{j!} [(L_{t})^{j} \mathfrak{U}(0,t+k) - \chi_{j}(0,t+k)]^{I} + O(h^{\rho+1}) \\ = \sum_{j=0}^{\rho} \frac{(\mu h)^{j}}{j!} \left[(L_{t})^{j} \left(\sum_{u=1}^{N^{II}} (0,t+k) + \mathfrak{g}(t+k) \right) - \chi_{j}(0,t+k) \right]^{I} \\ + O(h^{\rho+1}),$$

where $[\cdot]^{I}$ denotes the first ℓ components of the enclosed vectors. We see that $u_{\mu}^{I}(t+k)$ depends on time derivatives of $u^{I}(0,t+k)$ which, using (1.6), may be replaced by space derivatives of $u^{I}(0,t+k)$ and $u^{II}(0,t+k)$. Approximating these space derivatives by ρ -order accurate linear combinations of $u_{0}^{I}(t+k), \ldots, u_{\rho}^{I}(t+k)$ and $u_{0}^{II}(t+k), \ldots, u_{\rho}^{II}(t+k)$, respectively, we finally obtain (1.4e) if u is replaced by v and terms of order $O(h^{\rho+1})$ are dropped.

A concrete example of a second order accurate boundary condition of form (1.4e) for the special case B = f = 0, is given in [2].

2. The Reduced Problem

The difference approximation is completely defined now by (1.4) and we wish to apply to it the stability theory of Gustafsson, Kreiss and Sundström [3]. Trying to fit our approximation into the form discussed in [3] we realize, however, that while in the present paper the vector \underline{b} of the basic scheme (1.4a) is a general combination of \underline{f} and its derivatives, in [3] we have $\underline{b} = \underline{f}$. Indeed, the general \underline{b} admitted by us here is necessary if arbitrary high order approximations to (1.1a) are desired.^{*} Yet, it is not hard to see that this generalization does not affect the results of [3]. We conclude, therefore, that <u>making the same</u> <u>assumptions about our difference approximation as were made in [3], the</u> <u>theory of Gustafsson et al. holds for our case</u> and we raise the question of stability in the sense of Definition 3.3 of [3].

In Theorem 2.1 below we shall reduce the above stability question to that of a scalar outflow approximation with homogeneous boundary conditions. To obtain this theorem we begin by recalling Lemma 10.3 of [3] which provides a necessary and sufficient determinental stability criterion given entirely in terms of the <u>principal part</u> of the approximation, i.e., the part obtained by neglecting B and eliminating all inhomogeneity vectors. The mere existence of such a criterion implies that for stability furposes we may study (1.4) with $b_{\nu}(t) = d_{\mu}(t) = e_{\mu}(t) = g(t) = C_{j\sigma}^{\Pi I(\mu)} = D_{j}^{I I(\mu)} = 0$; hence, instead of (1.4) we may consider a basic scheme of the form

For example, the Lax-Wendroff scheme [7] for (1.1a) is

$$Q_{\underline{k}} v_{\nu}(t + k) = \sum_{\sigma=0}^{s} Q_{\sigma} v_{\nu}(t - \sigma k) ,$$

(2.1a)

$$Q_{\sigma} = \sum_{j=-r}^{p} A_{j\sigma} E^{j}, E_{\nu} = V_{\nu+1}, \qquad \nu = -1, \dots, s,$$

with initial values

(2.1b)
$$y_{\nu}(\sigma k) = y_{\nu}(\sigma k)$$
, $\sigma = 0,...,s$, $\nu = 0,1,...,$

and boundary conditions

(2.1c)

$$T_{-1}^{\Pi\Pi(\mu)} \underbrace{v_{\mu}^{\Pi}(t+k)}_{\sigma=0} = \sum_{\sigma=0}^{\underline{q}} T_{\sigma}^{\Pi\Pi(\mu)} \underbrace{v_{\mu}^{\Pi}(t-\sigma k)}_{\sigma=0}$$

$$T_{\sigma}^{\Pi\Pi(\mu)} = \sum_{j=0}^{\underline{m}} C_{j\sigma}^{\Pi\Pi(\mu)} E^{j}, \quad \mu = 0, \dots, r-1,$$

(2.1d)
$$y_0^{I}(t + k) = S y_0^{II}(t + k),$$

(2.1e)
$$y_{\mu}^{I}(t+k) = \sum_{j=1}^{\rho} D^{II(\mu)} y_{j}^{I}(t+k), \qquad \mu = 1,...,r-l,$$

where (2.1a) is now consistent with

(2.2)
$$\frac{\partial u}{\partial t} = A \frac{\partial u}{\partial x}$$

and the $A_{j\sigma}$ and $C_{j\sigma}^{III(\mu)}$ are polynomials in A and in A^{III} , respectively.

We thus obtain,

LEMMA 2.1. Approximation (1.4) is stable if and only if its principal part (2.1) is stable.

Setting

$$\mathbf{A}_{j\sigma} = \begin{pmatrix} \mathbf{A}_{j\sigma}^{\mathbf{I},\mathbf{I}} & \mathbf{0} \\ \mathbf{J}_{\sigma} & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_{j\sigma}^{\mathbf{I},\mathbf{I}} \end{pmatrix}, \quad \mathbf{j} = -\mathbf{r}, \dots, \mathbf{p}, \quad \sigma = -1, \dots, s,$$

2.2

 $v = r, r+1, \ldots,$

according to the partition of A in (1.2), our next step is to split the basic scheme (2.1a) and the initial values (2.1b) into

(2.3a)

$$\begin{aligned}
Q_{-1}^{II} & x_{\nu}^{I}(t+k) = \sum_{\sigma=0}^{8} Q_{\sigma}^{II} \frac{1}{v_{\nu}}(t-\sigma k) , & \nu = r, r+1, \dots, \\
Q_{\sigma}^{II} &= \sum_{j=-r}^{p} A_{j\sigma}^{II} E^{j},
\end{aligned}$$

(2.3b)
$$\vec{v}_{\nu}^{I}(\sigma k) = \vec{v}_{\nu}^{I}(\sigma k), \quad \sigma = 0,...,s, \quad \nu = 0,1,...,$$

and

$$Q_{-1}^{\Pi \Pi} \underbrace{y_{\nu}^{\Pi}}_{\nu} (t+k) = \sum_{\sigma=0}^{s} Q_{\sigma}^{\Pi \Pi} \underbrace{y_{\nu}^{\Pi}}_{\nu} (t-\sigma k), \qquad \nu = r, r+1, \dots,$$

(2.4a)

$$Q_{\sigma}^{\Pi\Pi} = \sum_{j=-r}^{p} A_{j\sigma}^{\Pi\Pi} E^{j},$$

(2.4b)
$$y_{\nu}^{II}(\sigma k) = y_{\nu}^{OII}(\sigma k), \quad \sigma = 0,...,s; \quad \nu = 0,1,...;$$

thus viewing approximation (2.1) as consisting of inflow and outflow parts given by (2.3) (2.1d,e) and (2.4) (2.1c), respectively. Obviously, (2.1) is stable if and only if both parts are.

We observe that the outflow part (2.4) (2.1c) is self-contained and provides, via (2.1d,e), the boundary values $\chi^{I}_{\mu}(t+k)$, $\mu = 0, \dots, r-1$. We may therefore consider (2.1d,e) as arbitrary inhomogeneous boundary values for the inflow part. So by the argument involving Lemma 10.3 of [3] preceding Lemma 2.1, we may replace (2.1d,e) -- without affecting stability -- by homogeneous boundary values

(2.5)
$$y^{I}_{\mu}(t+k) = 0, \qquad \mu = 0, \dots, r-1.$$

This gives us a new self-contained inflow part, (2.3) (2.5), whose

stability together with that of (2.4) (2.1c) is equivalent to the overall stability of (2.1).

Since the $A_{j\sigma}$ and $C_{j\sigma}^{III(\mu)}$ are diagonal, we write

$$A_{j\sigma} = diag(a_{j\sigma}), \quad C_{j\sigma}^{III(\mu)} = diag(c_{j\sigma}^{(\mu)}),$$

and split (2.3) (2.5) and (2.4) (2.1c) into n scalar components, each of the form

$$Q_{1}v_{\nu}(t+k) = \sum_{\sigma=0}^{s} Q_{\sigma}v_{\nu}(t-\sigma k), \qquad \nu = r, r+1, \dots,$$

(2.6a)

$$\mathbf{Q}_{\sigma} = \sum_{\mathbf{j}=-\mathbf{r}}^{\mathbf{p}} \mathbf{a}_{\mathbf{j}\sigma} \mathbf{E}^{\mathbf{j}},$$

(2.6b)
$$v_{\nu}(\sigma k) = v_{\nu}(\sigma k), \quad \sigma = 0,...,s, \quad \nu = 0,1,...,$$

(2.6c)
$$T_{-1}^{(\mu)}v_{\mu}(t+k) = \sum_{\sigma=0}^{q} T_{\sigma}^{(\mu)}v_{\mu}(t-\sigma k), \qquad \mu = 0,...,r-l,$$

where (2.6a) is consistent with a corresponding component of (2.2),

(2.7)
$$\partial u/\partial t = a \partial u/\partial x$$
, $a \neq 0$;

and the boundary conditions (2.6c) are either homogeneous, i.e.,

(2.8)
$$T_{-1}^{(\mu)} = 1, T_{\sigma}^{(\mu)} = 0, \mu = 0, \dots, r-1, \sigma = 0, \dots, s,$$

or are given by

(2.9)
$$T_{\sigma}^{(\mu)} = \sum_{j=0}^{m} c_{j\sigma}^{(\mu)} E^{j}, c_{0(-1)}^{(\mu)} \neq 0, \mu = 0, \dots, r-1, \sigma = -1, \dots, q$$

depending on whether a < 0 or a > 0, respectively.

Since (2.1) is stable if and only if (2.3) (2.5) and (2.4)(2.1c) are stable, and since the latter are stable if and only if their scalar components are, we obtain immediately,

LEMMA 2.2. <u>Approximation</u> (2.1) is stable if and only if the scalar <u>components of</u> (2.3) (2.5) and (2.4) (2.1c), given by (2.6) (2.8) and (2.6) (2.9), are stable.

In Section 4 we shall prove:

IEMMA 2.3. The inflow approximation (2.6) (2.8) is unconditionally stable.

This lemma -- due to Kreiss [4] in the special case when the basic scheme is dissipative, explicit and two-level -- combined with the previous two, finally yields the main result of this section:

THEOREM 2.1. <u>Approximation</u> (1.4) is stable if and only if the scalar outflow components of its principal part are stable.

The above discussion implies that from now on we may reduce our stability study to scalar approximations of form (2.6) with either (2.8) or (2.9). We thus conclude this section by stating the basic assumptions of [3] relating to these approximations which will hereafter hold throughout the paper.

ASSUMPTION 2.1 (Assumption 3.1, [3]). Approximation (2.6) is solvable, i.e., there exists a constant K > 0 such that for each $w \in \ell_2(x)$ there is a unique solution $y \in \ell_2(x)$ to

> $Q_{-1}y_{\nu} = W_{\nu}, \quad \nu = r, r+1, ...,$ $T_{-1}^{(\mu)}y_{\mu} = W_{\mu}, \quad \mu = 0, ..., r-1,$

with

 $\|\mathbf{y}\| \leq \mathbf{K} \|\mathbf{w}\|.$

Here, $l_2(\mathbf{x})$ is the space of all grid functions $\mathbf{w} = \{\mathbf{w}_v\}_{v=0}^{\infty}$ with $\|\mathbf{w}\|^2 = h \sum_{\nu=0}^{\infty} |\mathbf{w}_{\nu}|^2 < \infty$.

ASSUMPTION 2.2 (Assumption 5.1 [3]). The basic scheme (2.6a) is stable for the pure Cauchy problem, $-\infty < v < \infty$. That is, putting

(2.10)
$$a_j(z) \equiv a_{j(-1)} - \sum_{\sigma=0}^{s} z^{-\sigma-1} a_{j\sigma}, \qquad j = -r, ..., p,$$

we have:

(1) The von Neumann condition; i.e., the solutions $z(\xi)$ of the equation

(2.11)
$$\sum_{j=-r}^{p} a_{j}(z)e^{ij\xi} = 0,$$

satisfy $|z(\xi)| \leq 1$ for all $|\xi| \leq \pi$.

(i1) Those $z(\xi)$ which lie on the unit circle, are simple roots of (2.11).

ASSUMPTION 2.3 (Assumption 5.4 [3]). The basic scheme (2.6a) is either <u>dissipative</u>, i.e., the roots of (2.11) satisfy

(2.12)
$$|z(\xi)| < 1, \quad 0 < |\xi| \le \pi;$$

or it is <u>unitary</u>, namely

(2.13)
$$|z(\xi)| = 1, \quad |\xi| \leq \pi.$$

Finally, for convenience only we make,

ASSUMPTION 2.4 (Assumption 5.5 [3]).

(2.14) $a_{r}(z), a_{p}(z) \neq 0 \text{ for } |z| \geq 1$.

3. Statement of Main Results and Examples.

The purpose of this section is to provide easily checkable stability criteria for outflow approximations of form (2.6) (2.9). In view of Theorem 2.1 this is the key to the overall stability question of approximation (1.4).

Our results -- stated below and proved in Section 5 -- are essentially independent of the basic scheme (2.6a) and are given solely in terms of the boundary conditions. These results, however, do not apply to general boundary conditions of form (2.6c) (2.9); instead we are concerned in this section with the <u>translatory</u> case where (2.6c) (2.9) are of the form

3.1)

$$T_{-1}v_{\mu}(t + k) = \sum_{\sigma=0}^{q} T_{\sigma}v_{\mu}(t - \sigma k),$$

$$T_{\sigma} = \sum_{j=0}^{m} c_{j\sigma} E^{j}, c_{0(-1)} \neq 0, \mu = 0, \dots, r-1.$$

As mentioned in the introduction, such boundary conditions are widely used in practice since the coefficients $c_{j\sigma}$ are independent of μ and all boundary values are conveniently determined by the same procedure. Especially, when the numerical boundary consists of a single grid point (r = 1), the computation at the boundary is translatory by definition.

We associate now with the boundary conditions (3.1) the <u>boundary</u> characteristic function

(3.2)
$$R(z,\kappa) \equiv \sum_{j=0}^{m} c_j(z)\kappa^{j},$$

where

(3.3)
$$c_j(z) = c_{j(-1)} - \sum_{\sigma=0}^{q} z^{-\sigma-1} c_{j\sigma}, \quad j = 0,...,n.$$

This allows us to state

THEOREM 3.1. Let the basic scheme (2,6a) be two-level and dissipative. Then the outflow approximation (2.6a,b) (3.1) is stable if

(3.4)
$$R(z,\kappa) \neq 0 \quad \forall |z| \ge 1, \quad 0 < |\kappa| < 1.$$

EXAMPLE 3.1 (Kreiss [4, Theorem 6]; see also [1] and [2, Example (4.5)].) Let the basic scheme (2.6a) be two-level and dissipative, and let the boundary conditions be determined by horizontal extrapolation of order $\omega - 1$, i.e.,

(3.5a)
$$\mathbf{v}_{\mu}(\mathbf{t}+\mathbf{k}) = \sum_{j=1}^{\omega} {\omega \choose j} (-1)^{j+1} \mathbf{v}_{\mu+j}(\mathbf{t}+\mathbf{k}), \quad \mu = 0, \dots, r-1.$$

The boundary characteristic function -- which for one-level boundary conditions is always z-independent -- satisfies

(3.5b)
$$R(\kappa) = 1 - \sum_{j=1}^{\omega} {\binom{\omega}{j}} (-1)^{j+1} \kappa^{j} = (1-\kappa)^{\omega} \neq 0, \quad 0 < |\kappa| < 1.$$

Hence, (3.4) holds and by Theorem 3.1, (2.6a,b) (3.5) is stable.

It should be pointed out that Theorem 3.1 is generally false if the basic scheme is of more than two levels. Surprisingly, even the well known result in Example 3.1 may fail to hold; namely, <u>outflow dissipative</u> <u>multi-level basic schemes</u> $(s \ge 3)$ with boundary values determined by <u>extrapolation of type</u> (3.5a), <u>are not always stable</u>. For example, consider the 3-level, 5-point basic scheme

(3.6)
$$v_{\nu}(t+k) = [I - \frac{\varepsilon}{16} (E-I)^2 (I - E^{-1})^2] v_{\nu}(t-k)$$

+ $\lambda a(E - E^{-1}) v_{\nu}(t), \quad 0 < \varepsilon < 1, \quad \lambda a \le 1 - \varepsilon, \quad \nu = 2, 3, \dots,$

with boundary values $v_{\mu}(t+k)$, $\mu = 0,1$, determined by (3.5a). As shown in Section 9 of [6] the basic scheme is dissipative, and it is not hard to verify that the rest of our basic assumptions are fulfilled as well.

Yet, although condition (3.4) of Theorem 3.1 is satisfied as exhibited by (3.5b), we prove in Example 4.1 below that approximation (3.6) (3.5a) is unstable.

Despite the above observation we can strengthen Theorem 3.1 for multi-level dissipative basic shcemes as follows.

THEOREM 3.2. Let the basic scheme (2.6a) be dissipative. Then the outflow approximation (2.6a,b) (3.1) is stable if (3.4) holds and if

(3.7)
$$R(z, \kappa = 1) \neq 0 \quad \forall \quad |z| = 1, \quad z \neq 1.$$

Evidently, Example (3.6) (3.5a) implies that the additional condition (3.7) is essential for Theorem 3.2.

Having stated Theorems 3.1, 3.2, we see that when the boundary conditions (3.1) are not single-level (as in Examples 3.1), condition (3.4) may become a cumbersome inequality in two variables, z and κ . Seeking a convenient alternative to these theorems, we extend the range of μ in (3.1) to obtain the <u>boundary scheme</u>,

(3.8)

5

$$\sigma = \sum_{j=0}^{m} c_j \sigma^{j}, \qquad \sigma = -1, \dots, r,$$

μ = 0,1,2,...,

and in analogy to the definitions in Assumptions 2.1, 2.2, we introduce:

 $\mathbf{T}_{-1}\mathbf{v}_{\mu}(\mathbf{t}+\mathbf{k}) = \sum_{\sigma=0}^{q} \mathbf{T}_{\sigma}\mathbf{v}_{\mu}(\mathbf{t}-\sigma\mathbf{k}),$

DEFINITION 3.1. The boundary scheme (3.8) is said to be <u>solvable</u> if there exists a constant K > 0 so that for each $y \in l_2(x)$ there is a unique solution $w \in l_2(x)$ to

(3.9)
$$T_{-1} W_{\mu} = Y_{\mu}, \qquad \mu = 0, 1, 2, ...,$$

$$\|\mathbf{w}\| \leq \mathbf{K} \|\mathbf{y}\|.$$

DEFINITION 3.2. The boundary scheme is said to fulfill the von Neumann condition if the roots $z(\xi)$ of

(3.10)
$$R(z, \kappa = e^{i\xi}) = \sum_{j=0}^{m} c_j(z)e^{ij\xi} = 0$$

satisfy $|z(\xi)| \leq 1$ for all $|\xi| \leq \pi$.

We can now state,

THEOREM 3.3 (1st Main Theorem). Let the basic scheme (2.6a) be dissipative. If (3.7) holds, and if the boundary scheme (3.8) is solvable and satisfies the von Neumann condition, then the outflow approximation (2.6a,b) (3.1) is stable.

This result is an extended analogue of the main theorem (Theorem 2.2) of [2].

Useful sufficient conditions for (3.7) as well as for the solvability of the boundary scheme (3.8), are given in the next two lemmas.

IEMA 3.1. Condition (3.7) holds if any of the following is satisfied:

- (i) <u>The boundary conditions</u> (3.1) are two-level (i.e., q = 0) and accurate of order zero at least.
- (ii) The boundary conditions are three-level, accurate of order zero at least, and in addition $R(z = -1, \kappa = 1) \neq 0$.

LENNA 3.2. (i) The boundary scheme (3.8) is solvable if

$$\mathbf{T}_{1}(\kappa) \equiv \sum_{j=0}^{\mathbf{q}} c_{j(-1)} \kappa^{j} \neq 0, \qquad 0 < |\kappa| \leq 1.$$

(ii) In particular, explicit boundary schemes are always solvable.

EXAMPLE 3.2. (Compare the special cases [3,6.11] and [2, Example 1].) Let the basic scheme (2.6a) be dissipative and determine the boundary conditions by oblique extrapolation of order $\omega - 1$:

(3.11a)
$$v_{\mu}(t+k) = \sum_{j=1}^{\omega} {\binom{\omega}{j}(-1)^{j+1}} v_{\mu+j}[t-(j-1)k], \quad \mu = 0, \dots, r-1.$$

The boundary characteristic function associated with (3.11a) is given by

(3.11b)
$$R(z,\kappa) = 1 - \sum_{j=1}^{\omega} {\binom{\omega}{j}} {(-1)^{j+1}} z^{-j} \kappa^{j} = (1 - z^{-1} \kappa)^{\omega},$$

so obviously (3.7) holds. Further, the roots of (3.11b) satisfy

$$|z(\kappa = e^{i\xi})| = |e^{i\xi}| = 1;$$

thus, the associated boundary scheme -- which by Lemma 3.2(11) is solvable -fulfills the von Neumann condition, and by Theorem 3.3, (2.6a,b) (3.11a) is stable.

EXAMPLE 3.3. (Compare the special cases [3,(6.3c)],[9,(3.4)] and [4, Example 4].) Let the basic scheme (2.6a) be dissipative and let the boundary conditions be generated by the second order accurate Box-Scheme,

$$\mathbf{v}_{\mu}(\mathbf{t}+\mathbf{k}) + \mathbf{v}_{\mu+1}(\mathbf{t}+\mathbf{k}) - \lambda \mathbf{a}[\mathbf{v}_{\mu+1}(\mathbf{t}+\mathbf{k}) - \mathbf{v}_{\mu}(\mathbf{t}+\mathbf{k})]$$

(3.12)

$$= \mathbf{v}_{\mu}(t) + \mathbf{v}_{\mu+1}(t) + \lambda \mathbf{a}[\mathbf{v}_{\mu+1}(t) - \mathbf{v}_{\mu}(t)], \quad \mu = 0, \dots, r-1.$$

By Lemma 3.1(i), (3.7) is fulfilled; and since

$$\operatorname{Re}[T_{-1}(\kappa)] = 1 + \operatorname{Re}(\kappa) + \lambda a[1 - \operatorname{Re}(\kappa)] \neq 0, \qquad |\kappa| \leq 1,$$

then by Lemma 3.2(1) the boundary scheme is solvable. The boundary characteristic function is

$$R(z,\kappa) = 1 + \kappa - \lambda a(\kappa - 1) - z^{-1}[1 + \kappa + \lambda a(\kappa - 1)],$$

so its root satisfies

$$|\mathbf{z}(\kappa = e^{i\xi})| = \left| \frac{1 + e^{i\xi} + \lambda a(e^{i\xi} - 1)}{1 + e^{i\xi} - \lambda a(e^{i\xi} - 1)} \right| = 1, \qquad |\xi| \le \pi;$$

hence, the von Neumann condition holds as well, and Theorem 3.3 implies stability.

EXAMPLE 3.4. (Compare the special case [10, (6.26)].) Let the basic scheme (2.6a) be dissipative and define the boundary conditions by the right-sided weighted Euler scheme

(3.13)
$$\mathbf{v}_{\mu}(\mathbf{t}+\mathbf{k}) = \mathbf{v}_{\mu}(\mathbf{t}-\mathbf{k}) + \lambda \mathbf{a}[2\mathbf{v}_{\mu+1}(\mathbf{t}) - \mathbf{v}_{\mu}(\mathbf{t}+\mathbf{k}) - \mathbf{v}_{\mu}(\mathbf{t}-\mathbf{k})],$$

 $0 < \lambda \mathbf{a} \leq 1, \quad \mu = 0, \dots, r-1.$

The characteristic function for (3.13) is

$$R(z,\kappa) = 1 - z^{-2} - \lambda a(2\kappa z^{-1} - 1 - z^{-2})$$

and by equating to zero we find its roots,

$$\mathbf{z}(\kappa = e^{\mathbf{i}\xi}) = e^{\mathbf{i}\xi} \frac{\lambda \mathbf{a} \pm \mathbf{b}(\xi)}{\lambda \mathbf{a} + \mathbf{1}}, \quad \mathbf{b}(\xi) = \sqrt{(\lambda \mathbf{a})^2 + e^{-2\mathbf{i}\xi}[1 - (\lambda \mathbf{a})^2]};$$

hence

$$\mathbf{z}(\kappa=1)=\frac{\lambda\mathbf{a}\pm1}{\lambda\mathbf{a}+1}\neq e^{\mathbf{i}\boldsymbol{\varphi}}, \qquad 0<|\boldsymbol{\varphi}|\leq\pi,$$

and (3.7) holds. In addition, since $0 < \lambda a \le 1$ then $|b(\xi)| \le 1$, $|\xi| \le \pi$; so

$$|\mathbf{z}(\kappa = \mathbf{e}^{\mathbf{i}\xi})| \leq \frac{\lambda \mathbf{a} + |\mathbf{b}(\xi)|}{\lambda \mathbf{a} + \mathbf{1}} \leq \mathbf{1}, \qquad |\xi| \leq \pi,$$

and the boundary scheme satisfies the von Neumann condition. Finally, since the boundary scheme is explicit, Lemma 3.2(11) implies solvability, and by Theorem 3.3 stability follows.

A second second

We remark that solvability of the boundary scheme is necessary for Theorem 3.3. To see this, consider any dissipative basic scheme with zero-order accurate boundary conditions of the form

(3.14a)
$$v_{\mu}(t+k) - \Theta v_{\mu+1}(t+k) = v_{\mu}(t) - \Theta v_{\mu+1}(t)$$
, $\Theta > 1, \mu = 0, \dots, r-1$.

By Lemma 3.1(i), (3.7) is fulfilled. Also the boundary characteristic function is

(3.14b)
$$R(z,\kappa) = (1-z^{-1})(1-\Theta\kappa);$$

hence its single root, z = 1, satisfies the von Neumann condition. As shown in Example 4.2, however, the approximation is unstable, which is explained by the failure of the associated boundary scheme to be solvable. Indeed, taking y = 0 in (3.9) we find that the grid function $w = \{\theta^{-\mu} w_0\}_{\mu=0}^{\infty}$ with arbitrary w_0 , belongs to $\ell_2(x)$ and satisfies (3.9); thus we have neither the uniqueness nor the boundedness of w required by Definition 3.1.

Condition (3.7) is also necessary for Theorem 3.3 as can be shown by taking (3.6) with $0 < \lambda a < \frac{1}{2}$ and consistent boundary conditions of the form

(3.15a)
$$\mathbf{v}_{\mu}(t+k) = \mathbf{v}_{\mu}(t-k) + 2\lambda \mathbf{a}[\mathbf{v}_{\mu+1}(t-k) - \mathbf{v}_{\mu}(t-k)], \ \mu = 0, 1.$$

As mentioned before, the basic scheme is dissipative, and by Lemma 3.2(ii) the boundary scheme is solvable. The boundary characteristic function is

(3.15b) $R(z,k) = 1 - z^{-2} [1 + 2\lambda a(k - 1)]$,

so it is not hard to verify that the boundary scheme satisfies the von Neumann condition (and in fact, even dissipative.) Yet, as demonstrated by Example 4.3 below, (3.6) (3.15a) is unstable. The reason Theorem 3.3 does not apply in this case is that $R(z = -1, \kappa = 1) = 0$, i.e., (3.7) is violated.

So far we have treated in this section the case where the basic scheme is dissipative. For the general case where the basic scheme might also be unitary we need,

DEFINITION 3.3. The boundary scheme (3.8) is said to be dissipative if the roots of equation (3.10) satisfy $|z(\xi)| < 1$ for $0 < |\xi| \le \pi$.

This enables us to state,

THEOREM 3.4 (2nd Main Theorem). Let the basic scheme (2.6a) be dissipative or unitary, let (3.7) hold, and let the boundary scheme (3.8) be solvable and dissipative. Then the outflow approximation (2.6a,b) (3.1) is stable.

EXAMPLE 3.5. (Compare the special cases [3, (6.3a)], [8, (3.2)] and [2, Example 2].) Let the basic scheme (2.6a) be dissipative or unitary and let the boundary conditions be generated by the right-sided explicit Euler Scheme,

(3.16) $v_{\mu}(t+k) = v_{\mu}(t) + \lambda a[v_{\mu+1}(t) - v_{\mu}(t)], \quad 0 < \lambda a < 1, \mu = 0, \dots, r-1.$

10. 10 St.

The boundary characteristic function is now

$$R(z,\kappa) = 1 - z^{-1}[1 + \lambda a(\kappa - 1)],$$

and since $0 < \lambda a < 1$, its root satisfies

$$|\mathbf{z}(\mathbf{x} = e^{\mathbf{i}\xi})|^{2} = (\lambda \mathbf{a})^{2} + (1 - \lambda \mathbf{a})^{2} + 2\lambda \mathbf{a}(1 - \lambda \mathbf{a})\cos \xi$$

$$\leq (\lambda \mathbf{a})^{2} + (1 - \lambda \mathbf{a})^{2} + 2\lambda \mathbf{a}(1 - \lambda \mathbf{a}) = 1, \ 0 < |\xi| \le \pi;$$

hence the corresponding boundary scheme is dissipative. Moreover, since (3.16) is two-level, first order accurate and explicit, Lemmas 3.1(i) and 3.2(ii) imply that (3.7) holds and that the boundary scheme is solvable. The hypotheses of Theorem 3.4 are fulfilled therefore, and approximation (2.6a,b) (3.16) is stable.

EXAMPLE 3.6. (Compare the special cases [8, (3.3)] and [2, Example 3].) Let the basic scheme (2.6a) be dissipative or unitary, and define the boundary conditions by the right-sided, first order accurate, implicit Euler scheme:

(3.17)
$$v_{\mu}(t+k) - \lambda a[v_{\mu+1}(t+k) - v_{\mu}(t+k)] = v_{\mu}(t), \quad \mu = 0, \dots, r-1.$$

The characteristic function associated with (3.17) is given by

$$R(z,\kappa) = 1 - \lambda a(\kappa - 1) - z^{-1}$$

so its root satisfies

$$|z(\kappa = e^{i\xi})|^{2} = [(\lambda a)^{2} + (1 + \lambda a)^{2} - 2\lambda a(1 + \lambda a)\cos \xi]^{-1}$$

$$\leq [(\lambda a)^{2} + (1 + \lambda a)^{2} - 2\lambda a(1 + \lambda a)]^{-1} = 1, \ 0 < |\xi| \le \pi ,$$

and the boundary scheme is dissipative. Also, Lemma 3.1(i) implies (3.7); and since

$$\operatorname{Re}[T_{1}(\kappa)] = 1 + \lambda a[1 - \operatorname{Re}(\kappa)] \neq 0$$
, $|\kappa| \leq 1$,

then by Lemma 3.2(i) the boundary scheme is solvable. Thus, Theorem 3.4 applies and stability is assured.

In concluding this section we claim that condition (3.7) is necessary for Theorem 3.4 (as well as for Theorem 3.3.) For example, consider the nondissipative, 3-level Leap-Frog scheme

(3.18)
$$v_{v}(t+k) = v_{v}(t-k) + \lambda a[v_{v+1}(t) - v_{v-1}(t)], 0 < \lambda a < \frac{1}{2}, v=1,2,...$$

where for the boundary condition take (3.15a) with $\mu = 0$. As mentioned earlier, the corresponding boundary scheme is solvable and dissipative whereas (3.7) is violated. In Example 4.4 we show that approximation (3.18) (3.15a) is unstable, thus proving our claim.

We conjecture that the solvability and dissipativity of the boundary scheme are essential for Theorem 3.4.

4. A priliminary Stability Criterion.

In this section we use the theory of Gustafsson et al.[3] to obtain, in Theorem 4.2 below, a preliminary stability criterion for approximation (2.6). This criterion will be a major tool in proving Lemma 2.3 and Theorems 3.1-3.4.

Following [3] we associate with approximation (2.6) the <u>resolvent</u> <u>equation</u> (4.1a) $\left(Q_{-1} - \sum_{\sigma=0}^{s} z^{-\sigma-1} Q_{\sigma}\right) w_{\nu} = 0$, $\nu = r, r + 1, ...,$

(4.1b)
$$\left(T_{-1}^{(\mu)} - \sum_{\sigma=0}^{q} z^{-\sigma-1} T_{\sigma}^{(\mu)}\right) w_{\mu} = 0$$
, $\mu = 0, \dots, r-1$,

where $z \neq 0$ is a fixed arbitrary complex number. This can be written as

(4.2)
$$G(z)w = 0$$
,

where G(z) is a linear bounded operator on $\ell_2(x)$ defined by the left hand sides of (4.1). We say that z is an eigenvalue of approximation (2.6) if (4.2) has a nontrivial solution $w = w(z) \in \ell_2(x)$. If z is not an eigenvalue but there exists a sequence $\{w(z)^{(j)}\}_{j=1}^{\infty} \subseteq \ell_2(x)$ with $\|w(z)^{(j)}\| = 1$ such that

$$G(z) w(z)^{(j)} \frac{1}{j \to \infty} 0$$
,

we call z a generalized eigenvalue of approximation (2.6).

Having these definitions, we restate the main result of [3] in the language of [5]:

THEOREM 4.1 (Gustafsson et al. [3].) Approximation (2.6) is stable if and only if it has no eigenvalues nor generalized eigenvalues z with $|z| \ge 1$.

Seeking a practical version of Theorem 4.1, we first need the following characterization of solvability.

LEMMA 4.1 (essentially Osher, [8].) Approximation (2.6) is solvable if and only if

(i) The difference equations

 $Q_{-1} w_{\nu} = 0$, $\nu = r, r + 1, ...$

(4.2)

$$T_{-1}^{(\mu)}w_{\mu} = 0$$
, $\mu = 0, ..., r-1$,

<u>have no nontrivial solution</u> we $\ell_2(x)$.

(ii) The equation

(4.3a)
$$Q_{-1}(\kappa) \equiv \sum_{j=-r}^{p} a_{j(-1)} \kappa^{j} = 0$$

(4.3b) $r_0 = \max\{-j: a_{j(-1)} \neq 0\}$

(iii) $Q_{-1}(\kappa)$ does not vanish on the unit circle, i.e., $Q_{-1}(\kappa) \neq 0$, $|\kappa| = 1$.

<u>Proof</u>. Conditions (i) and (iii) coincide with Osher's conditions (d) and (g) in [8]. Regarding (ii), we note that $r_0 \ge 0$ (or else the basic scheme is unnaturally shifted to the right.) Hence,

$$Q_{-1}(\kappa) = \kappa \frac{r_0}{\sum_{j=-r_0}} a_{j(-1)}^{\kappa} a_{j(-1)}^{\kappa}$$

has a single pole of order r_0 at the origin. Since by (iii), $Q_{-1}(\kappa)$ does not vanish on the unit circle, we use the Argument Principle to find that there is no change in $\arg[Q_{-1}(\kappa = e^{i\xi})]$ as ξ varies from $-\pi$ to π if and only if (ii) holds. Thus, conditions (i) - (iii) are equivalent to (d) (e) (g) in [8], and Theorem I of [8] completes the proof.

Recalling the functions $a_j(z)$ in (2.10), we introduce now the <u>characteristic</u> equation of the basic scheme (2.6a),

$$(4.4) \quad P(z,\kappa) \equiv \sum_{j=-r}^{p} a_{j}(z)\kappa^{j} = 0 ,$$

whose r + p roots $\kappa_j(z)$ play a central role in determining the eigenvalues of approximation (2.6).

LEMMA 4.2 (compare Lemmas 5.1 and 5.2, [3].) For |z| > 1, the characteristic equation (4.4) has precisely r roots with $0 < |\kappa_j(z)| < 1$, p roots with $|\kappa_j(z)| > 1$, and no roots with $|\kappa_j(z)| = 1$.

<u>Proof</u>. By Assumption 2.4, the leading coefficients of $P(z,\kappa)$ do not vanish for |z| > 1; hence (4.4) has r + p roots, all satisfying $|\kappa_j(z)| > 0$. Since these roots are the solutions of the the polynomial equation

(4.5) $P_{r}(z,\kappa) \equiv \kappa^{r}P(z,\kappa) = 0$,

we may study (4.5) rather than (4.4).

By Assumption 2.2 (i),

$$P_{r}(z,\kappa = e^{i\xi}) \neq 0$$
, $|z| > 1$, $|\xi| \leq \pi$,

i.e., for |z| > 1, $P_r(z,\kappa)$ does not vanish on the unit circle $|\kappa| = 1$. Since the roots of $P_r(z,\kappa)$ are continuous functions of z, it follows that for |z| > 1 the number of roots satisfying $0 < |\kappa_j(z)| < 1$ is independent of z. In particular, consider the limit case

$$P_{r}(z \to \infty, \kappa) = \sum_{j=-r}^{p} a_{j(-1)} \kappa^{j+r}$$

By Lemma 4.1 (iii),

$$P_{\mathbf{r}}(\mathbf{z} \to \infty, \, \mathbf{\kappa} = \mathbf{e}^{\mathbf{i}\,\boldsymbol{\xi}}) \neq 0 \,, \, \, \left|\boldsymbol{\xi}\right| \leq \pi \,,$$

so by continuity again, the number of roots of $P_r(z,\kappa)$ satisfying $0 < |\kappa_j(z)| < 1$ may be determining by counting the roots $\kappa_j, |\kappa_j| < 1$, of $P_r(z \to \infty, \kappa)$. We have

$$P_{\mathbf{r}}(\mathbf{z} \to \infty, \kappa) = \kappa \sum_{\mathbf{j}=-\mathbf{r}_{0}}^{\mathbf{r}-\mathbf{r}_{0}} a_{\mathbf{j}(-1)}^{\mathbf{j}+\mathbf{r}_{0}}$$

where r_0 is defined in (4.3b). Moreover, by Lemma 4.1(ii), $\sum_{j=-r_0}^{p} a_{j(-1)}^{\kappa j} \text{ hence } \sum_{j=-r_0}^{p} a_{j(-1)}^{\kappa j+r_0} \text{ have precisely } r_0 \text{ roots}$ with $|\kappa| < 1$. Thus, with its additional $r - r_0$ zeroes, $P_r(z \to \infty, \kappa)$ has r roots with $|\kappa| < 1$ and the lemma follows.

According to the above lemma the roots of the characteristic equation (4.4) split for |z| > 1 into two groups: r <u>inner roots</u> satisfying $0 < |\kappa_j(z)| < 1$ and p <u>outer roots</u> with $|\kappa_j(z)| > 1$. By continuity, therefore, these groups of inner and outer roots remain well defined for $|z| \ge 1$ as well, where milder inequalities,

and a start

 $|\kappa_j(z)| \le 1$ and $|\kappa_j(z)| \ge 1$, hold, respectively. Since by Assumption 2.4, $\kappa = 0$ is never a root of $P(z,\kappa)$ for $|z| \ge 1$, we finally obtain:

LEMMA 4.3. For $|z| \ge 1$, the r + p roots $\kappa_j(z)$ of the characteristic equation (4.4) split into r inner roots with $0 < |\kappa_j(z)| \le 1$ and p outer roots with $|\kappa_j(z)| \ge 1$.

Now, let z be given. It is well known (e.g. [5]) that z is an eigenvalue or a generlized eigenvalue of approximation (2.6) if and only if equations (4.1) have a nontrivial solution of the form

(4.6)
$$W_{\nu} = \sum_{\alpha=1}^{N} \sum_{\beta=0}^{M_{\alpha}-1} \tau_{\alpha\beta} \Phi_{\alpha\beta}(\nu) \kappa_{\alpha}(z)^{\nu}, \quad \nu = 0, 1, 2, \dots,$$

where $\kappa_{\alpha}(z)$, $l \leq \alpha \leq N$, are the distinct inner roots of the characteristic equation (4.4) each with multiplicity $M_{\alpha} = M_{\alpha}(z)$. Here, $\Phi_{\alpha\beta}(v)$ are arbitrary polynomials in v with $deg[\Phi_{\alpha\beta}(v)] = \beta$, and $\tau_{\alpha\beta}$ are coefficients whose number, by Lemma 4.3, is precisely

$$\sum_{\alpha=1}^{N} M_{\alpha} = r$$

To find the $\tau_{\alpha\beta}$ we substitute (4.6) in (4.1b) and obtain a linear homogeneous system of r equations with r unknowns,

(4.7)
$$J(z)\tau' = 0$$

where J(z) is an $r \times r$ coefficient matrix and $\underline{\tau} = (\underline{\tau}_{Q\beta})$ is the unknown vector. Obviously, w = w(z) in (4.6) does not vanish if and only if (4.7) has a nontrivial solution $\underline{\tau}$; hence, z is an eigenvalue or a generalized eigenvalue of approximation (3.6) if and only if det $J(z) \neq 0$.

This observation combined with Theorem 3.1 gives the following equivalent of Lemma 10.3 of [3]:

LEMMA 4.4 (Gustafsson et al. [3].) Approximation (2.6) is stable if and only if

$$\det J(z) \neq 0 , |z| \geq 1 .$$

Now for z, $|z| \ge 1$, with corresponding distinct inner roots $\kappa_{\alpha} = \kappa_{\alpha}(z)$, $1 \le \alpha \le N$, each with multiplicity M_{α} , we make a specific choice for the polynomials $\Phi_{\alpha\beta}(v)$ in (4.6):

$$\phi_{\alpha\beta}(\nu) = \kappa_{\alpha}(z)^{-\beta} \beta! \begin{pmatrix} \nu \\ \beta \end{pmatrix}.$$

Thus, (4.6) becomes

$$\mathbf{w}_{\nu} = \sum_{\alpha=1}^{N} \sum_{\beta=0}^{M_{\alpha}-1} \tau_{\alpha\beta}\beta! \binom{\nu}{\beta} \kappa_{\alpha}^{\nu-\beta}, \qquad \nu = 0, 1, 2, \dots,$$

and substituting in (4.1b) -- with $T_{\sigma}^{(\mu)}$ given by either (2.8) or (2.9) -- we obtain explicit expressions for the system J(z)T'=0, namely

(4.8)
$$\sum_{\alpha=1}^{N} \sum_{\beta=0}^{M_{\alpha}-1} \beta! {\mu \choose \beta} \kappa_{\alpha}^{\mu-\beta} \tau_{\alpha\beta}, \quad \mu=0,\ldots,r-1,$$

or

$$(4.9) \sum_{\alpha=1}^{N} \sum_{\beta=0}^{M_{\alpha}-1} \sum_{j=0}^{m} \left(c_{j(-1)}^{(\mu)} - \sum_{\sigma=0}^{q} z^{-\sigma-1} c_{j\sigma}^{(\mu)} \right)_{\beta} \left(\left(\begin{array}{c} \mu+j \\ \beta \end{array}\right)_{\kappa} \left(z \right)^{\mu+j-\beta} \overline{c}_{\alpha\beta} = 0 ,$$

according to the inflow and outflow cases (a < 0 or a > 0), respectively. To further simplify these expressions we use the boundary coefficients to introduce r <u>boundary functions</u>

$$R_{\mu}(z,\kappa) \equiv \begin{cases} \kappa^{\mu} & , \quad a < 0 \\ \sum \\ j=0 \end{cases} \begin{pmatrix} c_{j(-1)}^{(\mu)} - \sum \\ \sigma=0 \end{cases} z^{-\sigma-1} c_{j\sigma}^{(\mu)} \kappa^{\mu+j} , \quad a > 0 \end{cases}$$

$$(4.10)$$

 $\mu = 0, \dots, r-1$.

Since

$$\frac{\partial^{\beta}R_{\mu}(z,\kappa)}{\partial \kappa^{\beta}} = \begin{cases} \beta : \begin{pmatrix} \mu \\ \beta \end{pmatrix} \kappa^{\mu-\beta} & a < 0 , \\ \sum_{j=0}^{m} \left(c_{j(-1)}^{(\mu)} - \sum_{\sigma=0}^{q} z^{-\sigma-1} c_{j\sigma}^{(\mu)} \right) \beta : \begin{pmatrix} \mu+j \\ \beta \end{pmatrix} \kappa^{\mu+j-\beta}, \quad a > 0 , \end{cases}$$

systems (4.8) and (4.9) both take the unified compact form

$$\sum_{\alpha=1}^{N} \sum_{\beta=0}^{M_{\alpha}-1} \left[\frac{\partial^{\beta} R_{\mu}(z,\kappa)}{\partial \kappa^{\beta}} \right]_{\kappa = \kappa_{\alpha}(z)} \cdot \tau_{\alpha\beta} = 0 , \quad \mu = 0, \dots, r-1;$$

so the coefficient matrix J(z) can be conveniently written as

(4.11a)
$$J(z) = [H(z, \kappa_1, M_1), \dots, H(z, \kappa_N, M_N)]$$

where $H(z,\kappa_{\alpha},M_{\alpha})$, $l \leq \alpha \leq N$, are the $r \times M_{\alpha}$ blocks

$$(4.11b) \quad H(z_{j}\kappa_{\alpha},M_{\alpha})$$

$$= \begin{bmatrix} \begin{bmatrix} R_{0}(z,\kappa) \\ R_{1}(z,\kappa) \\ \vdots \\ R_{r-1}(z,\kappa) \end{bmatrix}, \frac{\partial}{\partial \kappa} \begin{bmatrix} R_{0}(z,\kappa) \\ R_{1}(z,\kappa) \\ \vdots \\ R_{r-1}(z,\kappa) \end{bmatrix}, \dots, \frac{M_{\alpha-1}}{\partial \kappa} \begin{bmatrix} R_{0}(z,\kappa) \\ R_{1}(z,\kappa) \\ \vdots \\ R_{r-1}(z,\kappa) \end{bmatrix} \end{bmatrix}_{\kappa = \kappa_{\alpha}(z)}$$

This expression for J(z) gives us a concrete analogue of Lemma 4.4:

LEMMA 4.5. <u>Approximation</u> (2.6) is stable if and only if for <u>every</u> z, $|z| \ge 1$, (4.12) det $J(z) \equiv det[H(z,\kappa_1,M_1),\ldots,H(z,\kappa_N,M_N)] \neq 0$ <u>where</u> $\kappa_{\alpha} = \kappa_{\alpha}(z), 1 \le \alpha \le N$, <u>are the distinct inner roots of</u> (4.4), <u>each with multiplicity</u> $M_{\alpha} = M_{\alpha}(z)$.

A milder version of this theorem is given in Theorem 3.2, [2]. We return now to the case where the inflow boundary conditions are, as before, homogeneous and given by (2.6c) (2.8), whereas the outflow conditions (2.6c) (2.9) are translatory as described in (3.1). In this case -- where both the inflow and outflow conditions are of translatory type -- the boundary functions in (4.10) become

4.13)

$$R_{\mu}(z,\kappa) = \begin{cases} \kappa^{\mu} , a < 0 \\ \prod_{j=0}^{m} \left(c_{j(-1)} - \sum_{\sigma=0}^{q} z^{-\sigma-1} c_{j\sigma} \right) \kappa^{\mu+j} , a > 0 , \\ \mu = 0, \dots, r = 1 . \end{cases}$$

Hence, denoting

$$\mathbf{R}(\mathbf{z},\boldsymbol{\kappa}) \equiv \mathbf{R}_{\mathbf{n}}(\mathbf{z},\boldsymbol{\kappa})$$

(note: for a > 0 this definition coincides with the one in (3.2)), we find that

$$R_{\mu}(z,\kappa) = \kappa^{\mu}R(z,\kappa), \quad \mu = 0,...,r-1;$$

so the blocks of J(z) in (4.11b) become

$$= \begin{bmatrix} R(z,\kappa) \\ R(z,\kappa) \\ \vdots \\ \kappa^{r-1}R(z,\kappa) \end{bmatrix}, \frac{\partial}{\partial \kappa} \begin{bmatrix} R(z,\kappa) \\ \kappa R(z,\kappa) \\ \vdots \\ \kappa^{r-1}R(z,\kappa) \end{bmatrix}, \dots, \frac{\partial}{\partial \kappa} \begin{bmatrix} R(z,\kappa) \\ \kappa R(z,\kappa) \\ \vdots \\ \kappa^{r-1}R(z,\kappa) \end{bmatrix} \end{bmatrix}_{\kappa = R_{\alpha}} K^{\alpha-1} \begin{bmatrix} R(z,\kappa) \\ R(z,\kappa) \\ \vdots \\ R(z,\kappa) \\ \vdots \\ R(z,\kappa) \end{bmatrix}_{\kappa = R_{\alpha}} K^{\alpha-1} K^{\alpha$$

This allows us to simplify Lemma 4.5 as follows.

THEOREM 4.2 (<u>compare Theorem</u> 4.1, [2].) <u>Approximation</u> (2.6) <u>with</u> <u>translatory boundary conditions given by either</u> (3.1) for a > 0 or (2.8) for a < 0, <u>is stable if and only if for every</u> z, $|z| \ge 1$, <u>with corresponding inner roots</u> $\kappa_{\alpha}(z)$, $1 \le \alpha \le N$, <u>we have</u>

(4.15)
$$R(z,\kappa_{\alpha}) \neq 0$$
, $\alpha = 1,...,N$.

<u>Proof</u>. Suppose $R(z,\kappa_{\alpha}) = 0$ for some z, $|z| \ge 1$, with a corresponding inner root κ_{α} . Then clearly, the first column of $H(z,\kappa_{\alpha},M_{\alpha})$ in (4.14) vanishes; thus det J(z) = 0, and by Lemma 4.5 we have instability.

Conversely, let (4.15) hold and take an arbitrary z, $|z| \ge 1$, with distinct inner roots $\kappa_{\alpha}(z)$, $1 \le \alpha \le N$. To prove stability it suffices, by Lemma 4.5, to verify that the rows of

(4.16)
$$J(z) = [H(z, \kappa_1, M_1), \dots, H(z, \kappa_N, M_N)]$$

are linearly independent, where the $H(z,\kappa_{\alpha},M_{\alpha})$ are given by (4.14). For that purpose, let

(4.17)
$$\sum_{\mu=0}^{r-1} \gamma_{\mu} \begin{bmatrix} \kappa_{1}^{\mu}R(z,\kappa_{1}) \\ \vdots \\ \partial^{N} \left[\kappa_{N}^{\mu}R(z,\kappa_{N})\right] / \partial\kappa_{N} \end{bmatrix} = 0$$

be a vanishing linear combination of the rows of (4.16), and let us rewrite (4.17) as r scalar equations

(4.18)
$$\frac{\partial \mathbf{j}}{\partial \mathbf{k}^{\mathbf{j}}} \left\{ \mathbb{R}(\mathbf{z},\mathbf{\kappa}) \left[\sum_{\mu=0}^{\mathbf{r}-1} \gamma_{\mu} \mathbf{\kappa}^{\mu} \right] \right\}_{\mathbf{k} = \mathbf{k}_{\alpha}} = 0 , 1 \le \alpha \le \mathbb{N} , 0 \le \mathbf{j} \le M_{\alpha} - 1 .$$

Since by hypothesis

$$\mathbb{R}(z,\kappa)|_{\kappa = \kappa_{\alpha}} \neq 0$$
, $1 \leq \alpha \leq N$,

we expend the partial derivatives in (4.18) by Leibnitz' rule and use induction on $j \ge 0$ to find that the sum in (4.18) must have vanishing derivatives, i.e.,

$$\frac{d \mathbf{j}}{d \mathbf{k}^{\mathbf{j}}} \begin{bmatrix} \mathbf{r} - \mathbf{l} \\ \sum \gamma_{\mu} \mathbf{k}^{\mu} \\ \mu = 0 \end{bmatrix}_{\mathbf{k}} = \mathbf{k}_{\alpha} = 0 , \quad \mathbf{l} \leq \alpha \leq \mathbf{N} , \quad \mathbf{0} \leq \mathbf{j} \leq \mathbf{M}_{\alpha - \mathbf{l}} .$$

Consequently, the polynomial

1

$$\Psi(\kappa) \equiv \sum_{\mu=0}^{r-1} \gamma_{\mu} \kappa^{\mu}$$

which is of degree r-l at most, has r roots $(\kappa_{\alpha}, l \leq \alpha \leq N)$, each with multiplicity M_{α}) so $\Psi(\kappa) \equiv 0$ and the coefficients γ_{μ} must vanish. By (4.17), therefore, the rows of (4.16) are linearly independent and stability follows.

The proof of Lemma 2.3 and the counterexamples of Section 3 are almost at hand now.

<u>Proof of Lemma 2.3</u>. By (4.10), the boundary function associated with the homogeneous boundary conditions (2.6c) (2.8) is

$$R(z,\kappa) \equiv R_{0}(z,\kappa) = 1$$

Thus, (4.15) holds trivially, and by the last theorem approximation (2.6) (2.8) is stable.

EXAMPLE 4.1. Consider the dissipative basic scheme (3.6)with the boundary conditions in (3.5a). The boundary function is given by (3.5b) and for z = -1 it can be shown (as in Lemma 6.2, [3]) that the characteristic equation has exactly one inner root satisfying $\kappa(z = -1) = 1$. Hence, $R(z = -1, \kappa = 1) = 0$, and by Theorem 4.2 we have instability.

EXAMPLE 4.2. Take the zero-order accurate boundary conditions (3.14a) in combination with any basic scheme (dissipative or even unitary.) By (3.14b), $R(z = 1,\kappa) = 0$ for all κ ; so at z = 1 the characteristic boundary function vanishes for all inner roots, and Theorem 4.2 implies instability.

EXAMPLE 4.3. Take the same basic scheme as in Example 4.1 with the boundary conditions in (3.15a). As in Example 4.1 we have an inner root $\kappa(z \approx -1) = 1$ for which, by (3.15b), $R(z,\kappa) = 0$. Hence, (4.15) is violated and approximation (3.6) (3.15) is unstable.

EXAMPLE 4.4. Consider the Leap-Frog scheme (3.18) with a boundary condition as in Example 4.3. In Lemma 6.2 [3] it is shown that the characteristic equation of (3.18) has a single inner root $\kappa(z = -1) = 1$. So as in the previous example, $R(z = -1, \kappa = 1) = 0$, and by Theorem 4.2 instability follows.

5. Proof of Main Results.

We turn now to prove the results stated in Section 3, beginning with the following lemma. 5.1

LEMMA 5.1. For z = 1, the characteristic equation (4.4) has exactly one root satisfying $\kappa(z = 1) = 1$. In the outflow case (a > 0) this is always an outer root.

Proof. Since the basic scheme (2.6a) is consistent with

$$0 \neq a$$
, $x_6/u_6a = t_6/u_6$

the coefficients a must satisfy the ordinary consistency conditions $J\sigma$

$$\sum_{j=-r}^{p} a_{j(-1)} = \sum_{\sigma=0}^{s} \sum_{j=-r}^{p} a_{j\sigma}$$

and

$$\sum_{j=-r}^{p} ja_{j(-1)} = \sum_{\sigma=0}^{s} \sum_{j=-r}^{p} ja_{j\sigma} - \lambda a \sum_{\sigma=0}^{s} (\sigma+1) \sum_{j=-r}^{p} a_{j\sigma},$$

which can be written as

(5.1a)
$$\sum_{j=-r}^{p} a_{j}(z) = 0$$
, $z=1$

(5.1b)
$$\sum_{j=-r}^{p} ja_{j}(z) \Big|_{z=1} = -\lambda a \sum_{j=-r}^{p} \frac{d}{dz} a_{j}(z) \Big|_{z=1}$$

or equivalently as

(5.2a)
$$P(z,\kappa) = 0$$
,
 $z=\kappa = 1$

(5.2b)
$$\frac{\partial}{\partial \kappa} P(z,\kappa) \Big|_{z=\kappa=1} = -\lambda a \frac{\partial}{\partial z} P(z,\kappa) \Big|_{z=\kappa=1}$$

Here, $a_j(z)$ and $P(z,\kappa)$ are defined in (2.1) and (4.4), respectively. By (5.1a), z = 1 is a solution of

$$\sum_{j=-r}^{p} a_j(z) = 0$$

and by Assumption 2.2(ii) this solution is simple. Hence,

$$\frac{\mathrm{d}}{\mathrm{d}z} \sum_{j=-r}^{p} a_{j}(z) \Big|_{z=1} \neq 0$$

so by (5.2b)

(5.3)
$$\frac{\partial}{\partial \kappa} P(z,\kappa) \Big|_{z=\kappa=1} = -\lambda a \frac{\partial}{\partial z} P(z,\kappa) \Big|_{z=\kappa=1} = -\lambda a \frac{d}{dz} \sum_{j=-r}^{r} a_j(z) \Big|_{z=1} \neq 0$$
.

Having (5.2a) and (5.3), we employ the implicit function theorem to find that in the neighborhood of z = 1 the characteristic equation (4.4) can be uniquely solved for κ as a differentiable function of z such that

(5.4)
$$\kappa(z = 1) = 1$$

This is the first part of the lemma.

To complete the proof consider the outflow case a > 0. Then (5.2b) yields

(5.5)
$$\frac{d\kappa(z)}{dz}\Big|_{z=1} = \left[-\frac{\partial P}{\partial z}\Big/\frac{\partial P}{\partial \kappa}\right]_{z=\kappa=1} = \frac{1}{\lambda a} > 0 ;$$

so by (5.4) (5.5), for $z = 1 + \varepsilon$ with sufficiently small $\varepsilon > 0$,

$$\kappa(z) = 1 + (\lambda a)^{-1} \epsilon + 0(\epsilon^2) > 1$$
.

That is, for z in the right real neighborhood of z = 1,

 $|\kappa(z)| > 1$,

and by Lemma 4.2 this inequality is valid for all z, |z| > 1. By definition, therefore, x(z) of (5.4) is an outer root of (4.4) and the lemma follows.

<u>Proof of Theorem 3.1.</u> Take an arbitrary z, $|z| \ge 1$, and let $\kappa_{\alpha} = \kappa_{\alpha}(z)$ be a corresponding inner root. In order to prove stability it suffices, by Theorem 4.2, to show that

$$(5.4) \qquad R(z_{,\kappa}\alpha) \neq 0 .$$

By Lemma 4.3, we have $0 < |\kappa_{\alpha}(z)| \leq 1$, where for $0 < |\kappa_{\alpha}(z)| < 1$, (5.4) is implied by (3.4). Hence, we may restrict attention to inner roots on the unit circle, i.e., $\kappa_{\alpha}(z) = e^{i\xi}$, $|\xi| \leq \pi$. Since the basic scheme is dissipative then by (2.12), the solutions $z = z(\kappa)$ of (4.4) satisfy

(5.5)
$$|\mathbf{z}(\kappa = \mathbf{e}^{\mathbf{i}\xi})| < 1, \quad 0 < |\xi| \le \pi.$$

Thus, for $|z| \ge 1$, (4.4) has no roots $\kappa = e^{i\xi}$, $0 < |\xi| \le \pi$, and our discussion is further reduced to $|z| \ge 1$, $\kappa_{\alpha}(z) = 1$. Next, by continuity, (5.5) yields

$$|z(\kappa = 1)| \leq 1;$$

so $\kappa = 1$ is ruled out as an inner root for |z| > 1 and it remains to consider |z| = 1, $\kappa_{\alpha}(z) = 1$. Finally, by Lemma 5.1, $\kappa = 1$ is is excluded as an inner root for z = 1 and we are left with

(5.6)
$$|z| = 1$$
, $z \neq 1$, $\kappa_{0}(z) = 1$.

Since the basic scheme is consistent then by (5.2a),

(5.7)
$$P(z = 1, \kappa = 1) = 0$$

Moreover, since the basic scheme is two-level, $P(z,\kappa)$ is a polynomial of first degree in z^{-1} where, by (5.7), its only root is $z^{-1} = 1$. Thus,

$$P(z_{,\kappa} = 1) \neq 0$$
, $|z| = 1$, $z \neq 1$,

and the proof is complete.

For Theorem 3.2 we repeat the previous proof to the point where the remaining values of z and $\kappa_{\alpha}(z)$ to be studied are given in (5.6). At this point, (3.7) implies (5.4) and stability is assured.

To prove Theorems 3.3 and 3.4 we need yet another result.

IEMMA 5.2. Let the boundary scheme (3.8) be solvable and satisfy the von Neumann condition. Then,

(5.8)
$$R(z,\kappa) \neq 0$$
, $|\kappa| \leq 1$, $|z| > 1$,
and
(5.9) $R(z,\kappa) \neq 0$, $|\kappa| < 1$, $|z| \geq 1$.

<u>Proof</u>. Apply Lemma 4.1 to the solvable boundary scheme, rather than to the basic scheme, with equation (4.3a) replaced by its boundary counterpart

$$\mathbf{T}_{-1}(\kappa) \equiv \sum_{\mathbf{j}=0}^{m} c_{\mathbf{j}(-1)} \kappa^{\mathbf{j}} = 0 \quad .$$

Since the boundary scheme is one sided where by (3.1) $c_{j(-1)} \neq 0$, then parts (ii) - (iii) of the lemma imply that

$$T_{1}(\kappa) \neq 0$$
 , $0 < |\kappa| \leq 1$.

We also have,

$$r_{-1}(0) = c_{0(-1)} \neq 0;$$

so all told

$$(5.10) T_{-1}(\kappa) \neq 0 , |\kappa| \leq 1 .$$

Since the boundary scheme satisfies von Neumann condition then by (3.10), for |z| > 1, $R(z,\kappa)$ does not vanish on the unit circle $|\kappa| = 1$. Hence -- as in the proof of Lemma 4.2 -- the number of roots of $R(z,\kappa)$ satisfying $|\kappa(z)| \le 1$ is independent of z, |z| > 1, and, by continuity, equals that of the roots κ , $|\kappa| \le 1$, of

$$R(z \to \infty, \kappa) \equiv T_{-1}(\kappa) = 0$$

By (5.10), therefore, $R(z,\kappa)$ has no roots $|\kappa(z)| \le 1$ for |z| > 1, and (5.8) holds.

To obtain (5.9), we merely note that by (5.8), the roots $\kappa(z)$ of $R(z,\kappa)$ satisfy $|\kappa(z)| > 1$ if |z| > 1. Thus, for $|z| \ge 1$, those continuous roots satisfy $|\kappa(z)| \ge 1$ and the lemma follows.

<u>Proof of Theorem 3.3</u>. Since the boundary scheme is solvable and satisfies the von Neumann condition, Lemma 5.1 implies (3.4), and by Theorem 3.2, approximation (2.6a,b) (3.1) is stable.

<u>Proof of Theorem 3.4.</u> As in the proof of Theorem 3.1, let z satisfy $|z| \ge 1$, let $\kappa_{\alpha}(z)$, $|\kappa_{\alpha}(t)| \le 1$, be a corresponding inner root, and let us prove that

(5.11)
$$R(z, \kappa_{\alpha}) \neq 0$$
.

By Theorem 4.2, this will implies stability.

Comparing Definitions 3.2 and 3.3, we immediately see that since the boundary scheme is dissipative, it satisfies the von Neumann condition; so Lemma 5.2 applies, and it remains to verify (5.11) for z and $\kappa_{\alpha}(z)$ with

$$|z| = 1$$
, $|\kappa_{\alpha}(z)| = 1$.

Indeed, for

$$|z| = 1$$
, $|\kappa_{\alpha}(z)| = 1$, $\kappa_{\alpha}(z) \neq 1$,

(5.11) follows from the dissipativity of the boundary scheme as described in Definition 3.3; for

$$|z| = 1$$
, $z \neq 1$, $\kappa_{\alpha}(z) = 1$,

(5.11) is implied by (3.7); and finally, by Lemma 5.1, $\kappa = 1$ is never an inner root for z = 1. Thus, (5.11) is verified and the theorem is proven.

We conclude the paper by proving Lemmas 3.1 and 3.2.

<u>Proof of Lemma 3.1.</u> (i) Let the boundary scheme (3.1) be two-level and accurate of order zero at least. By zero-order accuracy, the boundary coefficients satisfy

$$\sum_{j=0}^{m} c_{j(-1)} = \sum_{\sigma=0}^{q} \sum_{j=0}^{m} c_{j\sigma};$$

hence

(5.12) $R(z,\kappa)|_{z=\kappa=1} = 0$.

Since the boundary scheme is two-level, $R(z, \kappa = 1)$ is a first degree polynomial in z^{-1} whose single root, by (5.12), is $z^{-1} = 1$. Consequently,

 $R(z_{j}\kappa = 1) \neq 0$, |z| = 1, $|z| \neq 1$,

and (3.7) holds.

(ii) In the three-level case, $R(z, \kappa = 1)$ is a 2nd degree polynomial in z^{-1} with real coefficients. By (5.12) again, $z^{-1} = 1$ is one of the roots, so the other is real as well, namely

(5.13)
$$R(z, \kappa = 1) \neq 0$$
, $|z| = 1$, $z \neq \pm 1$.

Combining (5.13) with our hypothesis

$$R(z = -1, \kappa = 1) \neq 0$$

(3.7) follows and the proof is complete.

<u>Proof of Lemma 3.2.</u> As in the proof of Lemma 5.2, apply Lemma 4.1 to the boundary scheme (3.8). Since the boundary scheme is right sided with $c_{0(-1)} \neq 0$, we find that it is solvable if

(a) the difference equations

(5.14)
$$T_{-1} w_{\nu} \equiv \sum_{j=0}^{m} c_{j(-1)} w_{\nu+j} = 0$$
, $\nu = 0, 1, 2, ...,$

have no nontrivial solution $w \in \ell_2(x)$, and

(b) the hypothesis of the present lemma is fulfilled, i.e.,

(5.15)
$$T_{-1}(\kappa) \equiv \sum_{j=0}^{m} c_{j(-1)} \kappa^{j} \neq 0$$
, $0 < |\kappa| \le 1$

Now, it is well known that the most general solution of (5.14) in $l_2(\mathbf{x})$ is a combination of powers of the roots of $T_{-1}(\kappa)$ which lie inside the unit disc. Thus, if (5.15) holds then the only solution of (5.14) is the trivial one, namely (b) implies (a) and the lemma follows.

REFERENCES

- M. Goldberg, On a boundary extropolation theorem by Kreiss, <u>Math. Comp.</u>, v. <u>31</u>, 1977, pp. 469-477.
- M. Goldberg and E. Tadmor, Scheme-independent stability criteria for difference approximations of hyperbolic initial-boundary value problems. I, <u>Math. Comp.</u>, v. <u>32</u>, 1978, pp. 1097 - 1107.
- B. Gustafsson, H.-O. Kreiss and A. Sundström, Stability theory of difference approximations for mixed initial boundary value problems. II, <u>Math. Comp.</u>, v. <u>26</u>, 1972, pp. 649-686.
- H.-O. Kreiss, Difference approximations for hyperbolic differential equations, "<u>Numerical Solution of Partial Differential</u>
 <u>Equations</u>" (Proc. Sympos. Univ. of Maryland, 1965), Academic
 Press, New York, 1966, pp. 51-58.
- H.-O. Kreiss, Stability theory for difference approximations of mixed initial boundary value problems. I, <u>Math. Comp.</u>, v. <u>22</u>, 1968, pp. 703 - 714.
- H.-O. Kreiss and J. Oliger, "<u>Methods for the Approximate Solu-</u> <u>tion of Time Dependent Problems</u>", GARP Publication Series No. 10, 1973.
- 7. P.D. Lax and B. Wendroff, Systems of conservation laws, <u>Comm</u>. Pure Appl. Math., v. <u>13</u>, 1960, pp. 217-237.
- 8. S. Osher, Stability of parabolic difference approximations to certain mixed initial boundary value problems, <u>Math. Comp.</u>,
 v. <u>26</u>, 1972, pp. 13-39.

R.1

- G. Skölermo, <u>How the boundary conditions affect the stability and</u> <u>accuracy of some implicit methods for hyperbolic equations</u>, Report No. 62, 1975, Dept. of Comp. Sci., Uppsala Univ., Uppsala, Sweden.
- G. Sköllermo, <u>Error analysis for the mixed initial boundary value</u> problem for hyperbolic equations, Report No. 63, 1975, Dept. of Comp. Sci, Uppsala Univ., Uppsala, Sweden.

R.2