

AD-A093 183

SOUTH CAROLINA UNIV COLUMBIA DEPT OF MATHEMATICS AND--ETC F/6 12/1
ON THE SOLUTION OF A CLASS OF STOCHASTIC INTEGRAL SYSTEMS.(U)

UNCLASSIFIED

NOV 80 A N RAO, W J PADGETT


F49620-79-C-0140

AFOSR-TR-80-1298

NL



END
DATE
FILED
L 8h
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AFOSR-TR- 80 - 1298



Department of Mathematics
and Statistics
The University of South Carolina
Columbia, South Carolina 29208

5

LEVEL

AD A093183

DTIC
ELECTED
DEC 22 1980



80 12 22 105

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER (18) AFOSR/TR-80-1298	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER (9)	
4. TITLE (and Subtitle) (6) ON THE SOLUTION OF A CLASS OF STOCHASTIC INTEGRAL SYSTEMS,		5. TYPE OF REPORT & PERIOD COVERED Interim rept.	
7. AUTHOR(s) (10) A.N.V. Rao and W. J. Padgett		8. CONTRACT OR GRANT NUMBER(s) (15) F49620-79-C-0140	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of South Carolina Dept. of Mathematics & Statistics Columbia, South Carolina 29208		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F (16) 2304/AS (17)	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM. Bolling AFB, Washington, D. C. 20332		12. REPORT DATE (11) November 1980 (12) 10/26	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (14) TR-62		13. NUMBER OF PAGES 25	
		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Stochastic systems; McShane integrals; Stochastic stability.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) General stochastic integral systems involving McShane integrals are studied. These systems include classes of stochastic integro-differential systems. Existence, uniqueness, and stability of solutions to such systems are investigated using contractor theory. In particular, the results yield frequency-type conditions for the stability of a class of stochastic systems.			

410-12

UNCLASSIFIED

5

ON THE SOLUTION OF A CLASS OF STOCHASTIC
INTEGRAL SYSTEMS

by

A.N.V. Rao^{*} and W. J. Padgett^{**}

University of South Carolina
Statistics Technical Report No. 68
60H20-2

DTIC
COLLECTED
DEC 22 1980
C

November, 1980

Department of Mathematics and Statistics
University of South Carolina
Columbia, S. C. 29208

* On leave from the University of South Florida, Tampa, Fla.

** This author's research supported by the United States Air Force Office
of Scientific Research under Contract No. F49620-79-C-0140.

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

Abstract

General stochastic integral systems involving McShane integrals are studied. These systems include classes of stochastic integro-differential systems. Existence, uniqueness, and stability of solutions to such systems are investigated using contractor theory. In particular, the results yield frequency-type conditions for the stability of a class of stochastic systems.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or Special
A	

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is approved for public release IAW AFR 190-12 (7b). Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

1. Introduction

Stochastic equations are important in stochastic systems theory as well as in other areas of science and engineering [9]. Existence, uniqueness, stability, and approximation of solutions to such equations have been studied by several authors, for example, Bharucha-Reid [2], Blankenship [3], Lee and Padgett [4], Padgett and Rao [6], Padgett and Tsokos [7], to name a few. In this paper, a general stochastic integral system which includes classes of stochastic integro-differential systems is studied. In particular, this paper generalizes the work of Padgett and Tsokos [7] and of Rao and Tsokos [8] on integro-differential systems and yields frequency-type conditions for the stability of a class of stochastic systems, extending the work of Blankenship [3].

Consider the stochastic integral system

$$\begin{aligned}
 x_1(t; \omega) = & x_{1,0}(\omega) + \int_0^t h_1(s, x(s; \omega); \omega) ds \\
 & + \sum_{j=1}^p \int_0^t r_{11j}(t, \tau) \int_0^\tau K_{11j}(\tau, s; \omega) f_{11j}(s, x(s; \omega)) dz_j(s; \omega) d\tau \\
 & + \sum_{j, l=1}^p \int_0^t r_{21jl}(t, \tau) \int_0^\tau K_{21jl}(\tau, s; \omega) f_{21jl}(s, x(s; \omega)) dz_j(s; \omega) dz_l(s; \omega) d\tau, \\
 & i=1, 2, \dots, n, \qquad (1.1)
 \end{aligned}$$

where

- (i) $t \in R^+ \equiv [0, \infty)$, $\omega \in \Omega$, the supporting set of a complete probability measure space (Ω, \mathcal{A}, P) ;
- (ii) $h_1: R^+ \times R^n \times \Omega \rightarrow R \equiv (-\infty, \infty)$;
- (iii) $r_{11j}(t, \tau)$, $r_{21jl}(t, \tau)$ are continuous real-valued functions defined for $0 \leq \tau \leq t < \infty$;

- (iv) $K_{1ij}(t,s;\omega)$ and $K_{2ijl}(t,s;\omega)$ are real-valued functions defined for $0 \leq s \leq t < \infty$ and $\omega \in \Omega$;
- (v) $f_{1ij}, f_{2ijl}: \mathbb{R}^+ \times \mathbb{R}^n \rightarrow \mathbb{R}$;
- (vi) $z(t;\omega)$ with subscript is a real-valued stochastic process satisfying certain conditions; and
- (vii) $x(t;\omega)$ is a vector stochastic process with components $x_i(t;\omega)$, $i = 1, 2, \dots, n$.

The integrals involving the process $z(s;\omega)$ in the equation (1.1) are to be understood as McShane integrals [5]. Throughout the paper we consider sums involving subscripts i, j and l . For notational simplicity, we let

$$\sum_i \equiv \sum_{i=1}^n \quad \text{and} \quad \sum_{j,l} \equiv \sum_{j,l=1}^p \quad \text{unless otherwise stated.}$$

2. Preliminaries

Let (Ω, \mathcal{A}, P) be a complete probability measure space. We shall assume that there is a family of sub- σ -algebras $\mathcal{A}_t, t \in \mathbb{R}^+$, such that for $s < t$, $\mathcal{A}_s \subset \mathcal{A}_t$. We shall further assume with respect to equation (1.1) that

(H₁) every process denoted by $z(t;\omega)$ with subscript is a real-valued stochastic process adapted to \mathcal{A}_t , with almost surely (a.s.) continuous sample functions, and satisfies the condition

$$E[(z(t;\omega) - z(s;\omega))^T | \mathcal{A}_s] \leq K(t-s), \text{ where } 0 \leq s \leq t < \infty, r = 1, 2, 4$$

and K is some constant;

(H₂) $x_0(\omega)$ is measurable with respect to \mathcal{A} and is mean-square continuous;

(H₃) $K_{1ij}(t,s;\omega)$ and $K_{2ijl}(t,s;\omega)$ are adapted to \mathcal{A}_t for all $t \geq s$ and are continuous as maps from $\Delta = \{(t,s): 0 \leq s \leq t < \infty\}$ into L_∞ .

the set of all P-essentially bounded random variables (define the norm $|||K(t,s;\omega)||| = P - \text{ess sup}_{\omega \in \Omega} |K(t,s;\omega)|$); and

(H₄) for any n-dimensional mean-square continuous process $x(t;\omega)$ adapted to A_t , $h(t,x(t;\omega);\omega)$, $f_{1ij}(t,x(t;\omega))$ and $f_{2ijl}(t,x(t;\omega))$ are adapted to A_t and are mean-square continuous.

Under the above assumptions, it is known [5, pp. 61-70, 138-139] that for an n-dimensional mean-square continuous process $x(s;\omega)$ adapted to A_s , the McShane integrals in equation (1.1) exist, are adapted to A_t , and satisfy the following inequalities:

$$\begin{aligned} & [E\{\int_0^t K_{1ij}(t,s;\omega)f_{1ij}(s,x(s;\omega))dz_j(s;\omega)\}^2]^{1/2} \\ & \leq 2K\int_0^t [E\{K_{1ij}(t,s;\omega)f_{1ij}(s,x(s;\omega))\}^2]^{1/2} ds \\ & \quad + E\{\int_0^t K_{1ij}(t,s;\omega)f_{1ij}(s,x(s;\omega))dz_j(s;\omega)\}^2 \\ & \leq c^2\int_0^t E\{K_{1ij}(t,s;\omega)f_{1ij}(s,x(s;\omega))\}^2 ds \end{aligned} \quad (2.1)$$

and

$$\begin{aligned} & E\{\int_0^t K_{2ijl}(t,s;\omega)f_{2ijl}(s,x(s;\omega))dz_j(s;\omega)dz_l(s;\omega)\}^2 \\ & \leq c^2\int_0^t E\{K_{2ijl}(t,s;\omega)f_{2ijl}(s,x(s;\omega))\}^2 ds \end{aligned} \quad (2.2)$$

where $c = 2K\sqrt{t} + \sqrt{K}$, K is the constant defined in (H₁), and E denotes expectation.

We shall now define some specific function spaces that will be used in this study. Let $y(t;\omega)$ be a second-order scalar process adapted to A_t .

Denote

$$|||y(t;\omega)|||_2 = [E\{y(t;\omega)\}^2]^{1/2}. \quad (2.3)$$

Definition 2.1. $C \equiv C(\mathbb{R}^+, L_2(\Omega, \mathcal{A}, P))$ will denote the space of scalar mean-square continuous functions $y(t; \omega)$ adapted to \mathcal{A}_t . We shall induce a topology on C by the family of semi-norms

$$\|y(t; \omega)\|_{C_n} = \sup_{t \in [0, n]} \|y(t; \omega)\|_2, n = 1, 2, \dots \quad (2.4)$$

It is known [9] that this topology is metrizable and the resulting metric space is a Fréchet space.

Definition 2.2. C_1 will denote the set of functions $y(t; \omega)$ in C such that

$$\sup_{t \geq 0} \|y(t; \omega)\|_2 < \infty.$$

Then, C_1 is a Banach space with norm $\|\cdot\|_{C_1}$ defined by

$$\|y(t; \omega)\|_{C_1} = \sup_{t \geq 0} \|y(t; \omega)\|_2.$$

Definition 2.3. Let B and D be Banach spaces in C and let T be an operator on C . Then the pair (B, D) is said to be admissible with respect to T if $TB \subset D$.

Definition 2.4. A Banach space B in C is said to be stronger than C if every sequence which converges in the norm of B also converges in the topology of C .

Definition 2.5. C_∞ will denote the space of continuous real functions $u(t) (t \geq 0)$ with norm $\|u\|_\infty$ defined by

$$\|u\|_\infty = \sup_{t \geq 0} |u(t)|.$$

Definition 2.6. Let $x(t; \omega)$ be a vector process with components $x_1(t; \omega) (i = 1, 2, \dots, n)$. Then $x(t; \omega) \in C^n (B^n \text{ or } D^n)$ if and only if $x_1(t; \omega) \in C(B \text{ or } D)$.

The following lemma is well-known [9].

Lemma 2.1. Let T be a continuous linear operator on C . Let B and D be Banach spaces stronger than C such that (B,D) is admissible with respect to T . Then T is continuous from B to D .

We shall now introduce the concept of a bounded integral vector contractor for a set of vector valued functions. Let B and D be Banach spaces stronger than C . Define the linear operators T , T_{ij} , and T_{ijl} on C by

$$(Ty)(t;\omega) = \int_0^t y(s;\omega) ds, \quad (2.5)$$

$$(T_{ij}y)(t;\omega) = \int_0^t r_{1ij}(t,\tau) \int_0^\tau K_{1ij}(\tau,s;\omega) y(s;\omega) dz_j(s;\omega) d\tau \quad (2.6)$$

$$(T_{ijl}y)(t;\omega) = \int_0^t r_{2ijl}(t,\tau) \int_0^\tau K_{2ijl}(\tau,s;\omega) y(s;\omega) dz_j(s;\omega) dz_l(s;\omega) d\tau. \quad (2.7)$$

Assume that the pair (B,D) is admissible with respect to each of the operators T, T_{ij} , and T_{ijl} . Let $h(t,x;\omega)$, $f_{1ij}(t,x;\omega)$, and $f_{2ijl}(t,x;\omega)$ be real-valued functions such that $h(t,x(t;\omega);\omega)$, $f_{1ij}(t,x(t;\omega);\omega)$, $f_{2ijl}(t,x(t;\omega);\omega)$ are in B whenever $x(t;\omega) \in D^n$. Let f_1 denote the $n \times p$ matrix with elements f_{1ij} and let f_2 denote the $n \times p \times p$ three-dimensional matrix with elements f_{2ijl} .

Definition 2.7. The set of functions (h, f_1, f_2) is said to have a bounded integral vector contractor $(\Gamma, \Gamma_1, \Gamma_2)$ with respect to (B^n, D^n) if

- (1) $\Gamma \equiv \Gamma(t,x)$ is a bounded linear operator from D to B for each $t \in R^+$ and $x \in R^n$. The function $||\Gamma(t,x)||$ is continuous in (t,x) and $||\Gamma(t,x)|| \leq Q(t)$ where $Q(t)$ is a bounded continuous function;
- (2) Γ_2 is an $(n \times p)$ matrix of operators $\Gamma_{1ij}(t,x)$ such that for each $t \in R^+$ and $x \in R^n$, $\Gamma_{1ij}(t,x)$ is a bounded linear operator from D to B . The function $||\Gamma_{1ij}(t,x)||$ is continuous in (t,x) and

- $\sum_{i,j} \|\Gamma_{1ij}(t,x)\| \leq Q_1(t)$, where $Q_1(t)$ is bounded and continuous;
- (3) Γ_2 is a three-dimensional matrix of operators $\Gamma_{2ijl}(t,x)$ such that for each $t \in \mathbb{R}^+$, $x \in \mathbb{R}^n$, $\Gamma_{2ijl}(t,x)$ is a bounded linear operator from D to B . The function $\|\Gamma_{2ijl}(t,x)\|$ is continuous in (t,x) and $\sum_{i,j,l} \|\Gamma_{2ijl}(t,x)\| \leq Q_2(t)$ where $Q_2(t)$ is bounded and continuous; and
- (4) for $x(t;\omega), y(t;\omega) \in D^n$, the following inequalities hold:

$$\begin{aligned}
 & \|h(t,x(t;\omega) + y(t;\omega) + (T\Gamma y)(t;\omega) \\
 & \quad + ((\sum_j T_{1ij} \Gamma_{1ij})y)(t;\omega) \\
 & \quad + ((\sum_{j,l} T_{2ijl} \Gamma_{2ijl})y)(t;\omega); \omega) \\
 & \quad - h(t,x(t;\omega); \omega) - (\Gamma y_1)(t;\omega)\|_B \leq \alpha_1 \|y\|_{D^n};
 \end{aligned}$$

$$\begin{aligned}
 & \|f_{1ij}(t,x(t;\omega) + y(t;\omega) + (T\Gamma y)(t;\omega) \\
 & \quad + ((\sum_j T_{1ij} \Gamma_{1ij})y)(t;\omega) \\
 & \quad + ((\sum_{j,l} T_{2ijl} \Gamma_{2ijl})y)(t;\omega)) \\
 & \quad - f_{1ij}(t,x(t;\omega)) - (\Gamma_{1ij} y_1)(t;\omega)\|_B \leq \alpha_{1ij} \|y\|_{D^n};
 \end{aligned}$$

$$\begin{aligned}
 & \|f_{2ijl}(t,x(t;\omega) + y(t;\omega) + (T\Gamma y)(t;\omega) \\
 & \quad + (\sum_j T_{1ij} \Gamma_{1ij} y)(t;\omega) \\
 & \quad + (\sum_{j,l} T_{2ijl} \Gamma_{2ijl} y)(t;\omega)) \\
 & \quad - f_{2ijl}(t,x(t;\omega)) - (\Gamma_{2ijl} y_1)(t;\omega)\|_B \leq \alpha_{2ijl} \|y\|_{D^n}.
 \end{aligned}$$

where Ty , $(\sum_j T_{11j} \Gamma_{11j})y$, and $(\sum_{j,l} T_{21jl} \Gamma_{21jl})y$ are n -vectors with components Ty_1 , $(\sum_j T_{11j} \Gamma_{11j})y_1$, and $(\sum_{j,l} T_{21jl} \Gamma_{21jl})y_1$, respectively. Let α denote n -vector with components α_1 , α_1 the $n \times p$ matrix of constants α_{11j} and α_2 , the three-dimensional $n \times p \times p$ matrix of constants α_{21jl} . Then the triplet $(\alpha, \alpha_1, \alpha_2)$ will be called the vector of contractor constants.

We remark that from equation (2.5) and the assumption (H_3) on K_1 and K_2 and from the inequalities (2.1) and (2.2), it is readily demonstrated that the operator T , T_{11j} , and T_{21jl} are continuous on C . Hence, if a pair of spaces (B, D) are both stronger than C and admissible with respect to T , T_{11j} and T_{21jl} , it follows from Lemma 2.1 that these are bounded linear operators from B to D . Therefore, there are constants k , k_{11j} and k_{21jl} such that

$$\|T\| \leq k, \quad (2.8)$$

$$\|T_{11j}\| \leq k_{11j}, \quad \text{and} \quad (2.9)$$

$$\|T_{21jl}\| \leq k_{21jl} \quad (2.10)$$

Definition 2.8. The random function $x(t; \omega)$ will be called a solution of the equation (1.1) if $x(t; \omega) \in C^n$ and satisfies (1.1) a.s.

3. Existence of Solutions

In this section the concept of a bounded integral vector contractor, as defined in the previous section, will be used to obtain the existence and uniqueness of solutions of the general stochastic integral system (1.1). The conditions under which the existence and uniqueness will be proven are

very general, and as a specific application, Theorem 3.3 will be used in Section 4 to study stability of stochastic systems under frequency-type conditions.

In the following theorems, B and D will denote Banach spaces stronger than C and B^n and D^n will be the corresponding product spaces.

Theorem 3.1 Let the system (1.1) satisfy the following conditions:

- (1) $x_0(\omega) \in D^n$;
- (2) the pair (B, D) is admissible with respect to each of the operators T , T_{1ij} and T_{2ijl} ;
- (3) the set of functions (h, f_1, f_2) has a bounded integral vector contractor $(\Gamma, \Gamma_1, \Gamma_2)$ with the vector of contractor constants $(\alpha, \alpha_1, \alpha_2)$.

Then, if $k\alpha_i + \sum_j k_{1ij}\alpha_{1ij} + \sum_{j,l} k_{2ijl}\alpha_{2ijl} < 1$, there exists a solution to equation (1.1) in D^n .

Proof. Consider the sequence $\{x^{(m)}\}$ defined by

$$\begin{aligned} x_i^{(m+1)}(t; \omega) &= x_i^{(m)}(t; \omega) - [y_i^{(m)}(t; \omega) + (Ty_i^{(m)})(t; \omega) \\ &\quad + \sum_j (T_{1ij}\Gamma_{1ij} y_i^{(m)})(t; \omega) \\ &\quad + \sum_{j,l} (T_{2ijl}\Gamma_{2ijl} y_i^{(m)})(t; \omega), \\ m &= 0, 1, 2, \dots; \quad i = 1, 2, \dots, n, \end{aligned} \quad (3.1)$$

where

$$\begin{aligned} y_i^{(m)}(t; \omega) &= x_i^{(m)}(t; \omega) - x_{i0}(\omega) \\ &\quad - (Th(s, x^{(m)}(s; \omega); \omega))(t; \omega) \end{aligned}$$

$$\begin{aligned}
& - \left(\sum_j T_{11j} f_{11j}(s, x^{(m)}(s; \omega)) \right) (t; \omega) \\
& - \left(\sum_{j,l} T_{21jl} f_{21jl}(s, x^{(m)}(s; \omega)) \right) (t; \omega), \tag{3.2}
\end{aligned}$$

and

$$x^{(m)} = (x_1^{(m)}, x_2^{(m)}, \dots, x_n^{(m)}), x_0(\omega) \in D^n.$$

It is clear from the definition of the operators Γ , Γ_{11j} , and Γ_{21jl} and from the admissibility of (B, D) with respect to T , T_{11j} , and T_{21jl} that $x^{(m)}, y^{(m)} \in D^n$. We shall now show that $\|y_i^{(m)}\|_D \rightarrow 0$ as $m \rightarrow \infty$. From equations (3.1) and (3.2), we have

$$\begin{aligned}
y_i^{(m+1)}(t; \omega) &= (Th(s, x^{(m)}(s; \omega); \omega))(t; \omega) \\
&+ \sum_j (T_{11j} f_{11j}(s, x^{(m)}(s; \omega)))(t; \omega) \\
&+ \sum_{j,l} (T_{21jl} f_{21jl}(s, x^{(m)}(s; \omega)))(t; \omega) \\
&- (Ty_i^{(m)})(t; \omega) \\
&- \sum_j (T_{11j} \Gamma_{11j} y_i^{(m)})(t; \omega) \\
&- \sum_{j,l} (T_{21jl} \Gamma_{21jl} y_i^{(m)})(t; \omega) \\
&- Th(s, x^{(m)}(s; \omega) - y^{(m)}(s; \omega)) \\
&- (Ty^{(m)})(s; \omega) - \left[\sum_j (T_{11j} \Gamma_{11j} y^{(m)}) \right] (s; \omega) \\
&- \left[\sum_{j,l} T_{21jl} \Gamma_{21jl} y^{(m)} \right] (s; \omega; \omega) \\
&- \sum_j T_{11j} f_{11j}(s, x^{(m)}(s; \omega) - y^{(m)}(s; \omega))
\end{aligned}$$

$$\begin{aligned}
& - (\Gamma y^{(m)})(s; \omega) - [\sum_j \Gamma_{11j} \Gamma_{11j} y^{(m)}] \\
& - [\sum_{j,l} \Gamma_{21jl} \Gamma_{21jl} y^{(m)}](s; \omega) (s; \omega) \\
& - \sum_{j,l} \Gamma_{21jl} f_{21jl}(s, x^{(m)}(s; \omega) - y^{(m)}(s; \omega) \\
& - (\Gamma y^{(m)})(s; \omega) - [\sum_j \Gamma_{11j} \Gamma_{11j} y^{(m)}](s; \omega) \\
& - (\sum_{j,l} \Gamma_{21jl} \Gamma_{21jl} y^{(m)})(s; \omega). \tag{3.3}
\end{aligned}$$

Equation (3.3) together with equations (2.8) and (2.9) enable us to conclude

$$\begin{aligned}
\|y_i^{(m+1)}\|_D & \leq [k\alpha_i + \sum_j k_{11j} \alpha_{11j} \\
& + \sum_j k_{21jl} \alpha_{21jl}] \|y_i\|_D \leq \alpha_i^* \|y_i^{(m)}\|_D, \tag{3.4}
\end{aligned}$$

where $\alpha_i^* < 1$. It therefore follows that $\|y_i^{(m)}\|_D \rightarrow 0$ as $m \rightarrow \infty$. Also from equation (2.1) we have

$$\begin{aligned}
\|x_i^{(m+1)} - x_i^{(m)}\|_D & \leq \|y_i^{(m)}\|_D + k \|\Gamma y_i^{(m)}\|_B \\
& + \sum_j k_{11j} \|\Gamma_{11j} y_i^{(m)}\|_B \\
& + \sum_{j,l} k_{21jl} \|\Gamma_{21jl} y_i^{(m)}\|_B \\
& \leq [1 + Q^* k + Q_1^* \sum_j k_{11j} \\
& + Q_2^* \sum_{j,l} k_{21jl}] \|y_i^{(m)}\|_D \tag{3.5}
\end{aligned}$$

where Q^*, Q_i^* , $i = 1, 2$ are upper bounds on $Q(t)$, $Q_i(t)$, $i = 1, 2$.

It follows from (3.4) and (3.5) that the sequence $\{x_1^{(m)}\}$ is a Cauchy sequence in D . Hence, there exists an $x_1 \in D$ such that $\lim_{m \rightarrow \infty} x_1^{(m)} = x_1$.

In view of the continuity of operators and the functions involved in equation (3.2), it follows that x_1 is a solution of equation (1.1) and this completes the proof. ///

We next prove a theorem concerning the uniqueness of solutions of equation (1.1).

Theorem 3.2. Let the hypotheses of Theorem 3.2 be satisfied. Assume in addition that the linear integral system

$$\begin{aligned} \eta_1(t; \omega) = & y_1(t; \omega) \\ & + \sum_j \int_0^t r_{11j}(t, \tau) \int_0^\tau K_{11j}(t, s; \omega) (\Gamma_{11j}(s, x(s; \omega)) y_1(s; \omega) dz_j(s; \omega) d\tau \\ & + \sum_{j, l} \int_0^t r_{21jl}(t, \tau) \int_0^\tau K_{21jl}(t, s; \omega) \Gamma_{21jl}(s, x(s; \omega)) y_1(s; \omega) dz_j(s; \omega) dz_l(s; \omega) d\tau \end{aligned} \quad (3.6)$$

has a solution $y_1(t; \omega)$ in D for every $x(t; \omega) \in D^n$ and $\eta_1(t; \omega) \in D$. Then equation (1.1) has a unique solution $x(t; \omega) \in D^n$.

Proof. Let $x_1^{(1)}(t; \omega)$ and $x_1^{(2)}(t; \omega)$ be two solutions in D of (1.1) corresponding to two initial random variables $x_{1,0}^{(1)}(\omega)$ and $x_{1,0}^{(2)}(\omega)$, $i = 1, 2, \dots, n$. Then

$$\begin{aligned} x_1^{(1)}(t; \omega) - x_1^{(2)}(t; \omega) = & x_{1,0}^{(1)}(\omega) - x_{1,0}^{(2)}(\omega) + \left[\int_0^t h_1(s, x^{(1)}(s; \omega); \omega) \right. \\ & \left. - h_1(s, x^{(2)}(s; \omega); \omega) \right] ds \\ & + \sum_j \int_0^t r_{11j}(t, \tau) \int_0^\tau K_{11j}(\tau, s; \omega) [f_{11j}(s, x^{(1)}(s; \omega)) \\ & - f_{11j}(s, x^{(2)}(s; \omega))] dz_j(s; \omega) d\tau \end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j} \int_0^t r_{ijl}(t,\tau) \int_0^\tau k_{2ijl}(\tau,s;\omega) [f_{2ijl}(s,x^{(1)}(s;\omega)) \\
& \quad - f_{2ijl}(s,x^{(2)}(s;\omega))] dz_j(s;\omega) dz_l(s;\omega) d\tau \\
& \quad i = 1, 2, \dots, n. \tag{3.7}
\end{aligned}$$

Letting $\eta_i(s;\omega) = x_i^{(1)}(t;\omega) - x_i^{(2)}(t;\omega)$ we can conclude from the fact that (3.6) has a solution for every z and x , that there is a $y_i(t;\omega) \in D$ such that

$$\begin{aligned}
x_i^{(1)}(t;\omega) &= x_i^{(2)}(t;\omega) + y_i(t;\omega) \\
& + \sum_j r_{1ij}(t,\tau) \int_0^\tau k_{1ij}(\tau,s;\omega) (\Gamma_{1ij}(s,x^{(2)}(s;\omega)) y_j)(s;\omega) dz_j(s;\omega) d\tau \\
& + \sum_{j,l} r_{2ijl}(t,\tau) \int_0^\tau k_{2ijl}(\tau,s;\omega) (\Gamma_{2ijl}(s,x^{(2)}(s;\omega)) y_j)(s;\omega) dz_j(s;\omega) dz_l(s;\omega) d\tau. \tag{3.8}
\end{aligned}$$

From (3.7) and (3.8), and upon using the contractor condition and simplifying, we obtain

$$\begin{aligned}
\|y_i(t;\omega)\|_D &\leq \|x_{i,0}^{(1)}(\omega) - x_{i,0}^{(2)}(\omega)\|_D \\
& + (ak + \sum_j a_{1ij} k_{1ij} + \sum_{j,l} a_{2ijl} k_{2ijl}) \|y_i(t;\omega)\|_D.
\end{aligned}$$

from which we get

$$\|y_i\|_D \leq (1 - ak - \sum_j a_{1ij} k_{1ij} - \sum_{j,l} a_{2ijl} k_{2ijl})^{-1} \times \|x_{i,0}^{(1)} - x_{i,0}^{(2)}\|_D. \tag{3.9}$$

If $x_{i,0}^{(1)} = x_{i,0}^{(2)}$, it follows that $y_i \equiv 0$, which in view of (3.8) implies that $x_i^{(1)} = x_i^{(2)}$. ///

We now consider the following special case of the equation (1.1) because of its usefulness in applications:

$$\begin{aligned}
x(t; \omega) &= h(t; \omega) + \int_0^t r_0(t, \tau) \int_0^\tau K_0(\tau, u; \omega) f_0(u, x(u; \omega)) du d\tau \\
&+ \sum_j \int_0^t r_{1j}(t, \tau) \int_0^\tau K_1(\tau, u; \omega) f_{1j}(u, x(u; \omega)) dz_j(u; \omega) d\tau \\
&+ \sum_{j, l} \int_0^t r_{2jl}(t, \tau) \int_0^\tau K_2(\tau, u; \omega) f_{2jl}(u, x(u; \omega)) dz_j(u; \omega) dz_l(u; \omega) d\tau.
\end{aligned}
\tag{3.10}$$

Let T_0 be the operator

$$(T_0 x)(t; \omega) = \int_0^t r_0(t, \tau) \int_0^\tau K_0(\tau, u; \omega) x_0(u; \omega) du d\tau,$$

and let T_{1j} and T_{2jl} be as defined in equations (2.6) and (2.7), respectively, with $1 = 1$, $r_{11j} = r_{1j}$, $r_{21jl} = r_{2jl}$, $K_{11j} = K_{1j}$ and $K_{21jl} = K_{2jl}$. Also, let C_1^* be a subspace of the Banach space C_1 .

Theorem 3.3. Assume the following:

- (i) $h(t; \omega) \in C_1^*$.
- (ii) The pair (C_1, C_1^*) is admissible with respect to each of the operators T_0, T_{1j} , and T_{2jl} .
- (iii) There are positive constants c_0, c_{1j} and c_{2jl} such that the operator $(I - c_0 T_0)$ is a.s. invertible on C_∞ and the operator $(I - (I - c_0 T_0)^{-1} (\sum_j c_{1j} T_{1j} + \sum_{j, l} c_{2jl} T_{2jl}))$ is invertible as a map from C_1 to C_1 .
- (iv) $|f_0(t, utv) - f_0(t, u) - c_0 v| \leq \gamma_0 |v|$,
 $|f_{1j}(t, utv) - f_{1j}(t, u) - c_{1j} v| \leq \gamma_{1j} |v|$, and
 $|f_{2jl}(t, utv) - f_{2jl}(t, u) - c_{2jl} v| \leq \gamma_{2jl} |v|$
for $u, v \in \mathbb{R}$ where γ_0, γ_{1j} , and γ_{2jl} are positive constants.

Then, if

$$\begin{aligned} & \| (I - (I - c_0 T_0)^{-1} (\sum_j c_{1j} T_{1j} + \sum_{j,l} c_{2jl} T_{2jl}))^{-1} \| \\ & [\gamma_0 \| (I - c_0 T_0)^{-1} T_0 \| + \sum_j \gamma_{1j} \| (I - c_0 T_0)^{-1} T_{1j} \| \\ & + \sum_{j,l} \gamma_{2jl} \| (I - c_0 T_0)^{-1} T_{2jl} \|] < 1, \end{aligned}$$

the equation (3.9) has a unique solution in C_1^* such that

$$\sup_{t \geq 0} \| x(t; \omega) \|_2 \leq M \sup_{t \geq 0} \| h(t; \omega) \|_2.$$

Proof. Using the invertibility of the operator $(I - c_0 T_0)$, equation (3.10) can be written as

$$\begin{aligned} x(t; \omega) &= \bar{h}(t; \omega) + \bar{T}_0 \bar{f}_0(t, x(t; \omega)) \\ &+ \sum_j \bar{T}_{1j} f_{1j}(t, x(t; \omega)) \\ &+ \sum_{j,l} \bar{T}_{2jl} f_{2jl}(t, x(t; \omega)), \end{aligned} \tag{3.11}$$

where

$$\bar{h}(t, \omega) = (I - c_0 T_0)^{-1} h(t; \omega),$$

$$\bar{f}_0(t, x) = f_0(t, x) - c_0 x,$$

$$\bar{T}_{1j} = (I - c_0 T_0)^{-1} T_{1j},$$

and

$$\bar{T}_{2jl} = (I - c_0 T_0)^{-1} T_{2jl}.$$

From condition (iv) of this theorem, it is easily verified that with the choice

$$\begin{aligned}
 \Gamma_0 &= 0 \\
 \Gamma_{1j} &= c_{1j} [I - \sum_j c_{1j} \bar{T}_{1j} - \sum c_{2jl} \bar{T}_{2jl}]^{-1} \\
 \Gamma_{2jl} &= c_{2jl} [I - \sum_j c_{1j} \bar{T}_{1j} - \sum c_{2jl} \bar{T}_{2jl}]^{-1}, \quad (3.12)
 \end{aligned}$$

the set of functions (f_0, f_{1j}, f_{2jl}) has a vector of contractor constants $\alpha_0 = \gamma \gamma_0$, $\alpha_{1j} = \gamma \gamma_{1j}$, and $\alpha_{2jl} = \gamma \gamma_{2jl}$, where

$$\gamma = \|(I - \sum_j c_{1j} \bar{T}_{1j} - \sum_{j,l} c_{2jl} \bar{T}_{2jl})^{-1}\|.$$

The existence of a solution now follows from Theorem 3.1.

To prove the uniqueness, we note that for the choice of contractors given by equations (3.12), the equality

$$I - \sum_j \bar{T}_{1j} \Gamma_{1j} - \sum_{j,l} \bar{T}_{2jl} \Gamma_{2jl} = I - \sum_j c_{1j} \bar{T}_{1j} - \sum_{j,l} c_{2jl} \bar{T}_{2jl}, \quad (3.13)$$

holds. Thus, the invertibility of the operator $(I - \sum_j c_{1j} \bar{T}_{1j} - \sum_{j,l} c_{2jl} \bar{T}_{2jl})$ implies that the equation

$$\eta(t; \omega) = (I - \sum_j c_{1j} \bar{T}_{1j} - \sum_{j,l} c_{2jl} \bar{T}_{2jl}) \eta(t; \omega)$$

has a unique solution for each $\eta \in C_1^*$. Therefore, by Theorem 3.2 and from equation (3.9) the solution is unique and satisfies the condition

$$\sup_{t \geq 0} \|x(t; \omega)\|_2 \leq M \sup_{t \geq 0} \|h(t; \omega)\|_2.$$

///

4. Application to the Stability of Stochastic Systems

In this section we shall discuss the stability of solutions of a special

class of feedback systems with convolution kernels, sector-bounded nonlinearities, and disturbances comprised of a Brownian motion and a Poisson point process. Using the contractor theory, we shall obtain sufficient conditions under which the feedback system is mean-square stable [3]. The conditions involve the sector parameters of the nonlinearities, the parameters of the stochastic disturbances and the Laplace transforms of the kernels.

Consider the feedback system

$$\begin{aligned} x(t;\omega) = & h(t;\omega) - \int_0^t r_0(t-\tau) \int_0^\tau K_0(\tau-u) f_0(u, x(u;\omega)) du d\tau \\ & - \int_0^t r_1(t-\tau) \int_0^\tau K_1(\tau-u) f_1(u, x(u;\omega)) dz_1(u;\omega) d\tau \\ & - \int_0^t r_2(t-\tau) \int_0^\tau K_2(\tau-u) f_2(u, s(u;\omega)) dz_2(u;\omega) d\tau. \end{aligned} \quad (4.1)$$

The process $z_1(t;\omega)$ and $z_2(t;\omega)$ in (4.1) satisfy:

- (i) $z_1(t;\omega)$ is a random process such that $z_1(0;\omega) = 0$ and

$$z(t;\omega) - z_1(s;\omega) = \sum_{s < \tau \leq t} v(\tau;\omega)(N(\tau;\omega) - N(\tau^-;\omega)).$$

The random function $v(\tau;\omega)$ has mean μ and variance σ^2 and, corresponding to any finite set τ_1, \dots, τ_k , the random variables $v(\tau_i;\omega)$ are independent, and $N(t;\omega)$ is a Poisson process with parameter λ (see [5], page 88). Furthermore, $v(t;\omega)$ is independent of all increments $N(t;\omega) - N(s;\omega)$.

- (ii) $z_2(t;\omega)$ is a standard Brownian motion process.

Under the above assumptions the following results are known [5]:

$$\mathbb{E} \int_0^t f(t,\tau) dz_2(\tau;\omega) = 0; \quad \mathbb{E} \left\{ \int_0^t f(t,\tau) dz_2(\tau;\omega) \right\}^2 = \int_0^t \mathbb{E} f^2(t,\tau) d\tau; \quad (4.2)$$

$$E \int_0^t f(t, \tau) dz_1(\tau; \omega) = \lambda \mu \int_0^t f(t, \tau) d\tau; \quad (4.3)$$

$$E \left\{ \int_0^t f(t, \tau) d[z_1(\tau; \omega) - \mu \lambda \tau] \right\}^2 = \int_0^t E f^2(t, \tau) \lambda (\mu^2 + \sigma^2) d\tau. \quad (4.4)$$

We shall further assume that z_1 and z_2 are independent and h is admissible and non-anticipating [3]. The functions r_1 , K_1 , and f_1 ($i = 0, 1, 2$) are all continuous. The nonlinearities f_1 are sector-bounded and satisfy the condition

$$|f_1(t, u + v) - f_1(t, u) - c v| \leq \frac{b-a}{2} |v|$$

where $c = \frac{a+b}{2}$ ($i = 0, 1, 2$), $u, v \in \mathbb{R}$.

In what follows, we shall denote the Laplace transform of a function $r(t)$ by $\hat{r}(s)$ and the convolution of two functions f and g by $f * g$. We will now present a stability theorem for the system (4.1).

Theorem 4.1 Assume

$$(i) \quad \left(-\frac{1}{c}, 10\right) \notin \bigcup_{\text{Re } s \geq -\beta_0} \hat{R}_0(s)$$

$$\text{where } \hat{R}_0(s) = [\hat{r}_0(s) \hat{K}_0(s) + \mu \lambda \hat{r}_1(s) \hat{K}_1(s)]$$

and $\beta_0 > 0$ is such that

$$\int_0^\infty e^{-\beta_0 t} (|r_0(t)K_0(t)| + |r_1(t)K_1(t)|) dt < \infty.$$

$$(ii) \quad \text{Let } \hat{R}_1(s) = (1 - \hat{R}_0(s))^{-1} \hat{r}_1(s), \quad i = 1, 2 \quad \text{and}$$

$$\hat{W}(s) = \lambda (\mu^2 + \sigma^2) [(\hat{R}_1(s) \hat{K}_1(s)) * (\hat{R}_1(s) \hat{K}_1(s)) + (\hat{R}_1(s) \hat{K}_2(s)) * (\hat{R}_1(s) \hat{K}_2(s))].$$

$$\text{so that } \left(-\frac{2}{a+b}, 10\right) \notin \bigcup_{\text{Re } s \geq -\beta_0} \hat{W}(s).$$

$$(iii) \sup_{s \geq -\beta_0} \left| \left(1 - \frac{a^2 + b^2}{2} \hat{W}(s) \right)^{-1} \right|$$

$$\times \left\{ \sup_{f \in R} \left| \int_{-\infty}^{+\infty} \frac{\hat{r}^*(\mu + j(\xi - \xi_0)) \hat{r}^*(\mu + j\xi_0)}{(\mu + j(\xi - \xi_0))(\mu + j\xi_0)} d\xi_0 \right| \right.$$

$$\left. + \sup_{s \geq -\beta_0} W(s) \right\} < \frac{2}{b-a}$$

$$\text{where } \hat{r}^*(s) = (1 - \hat{R}_0(s))^{-1} [\hat{r}_0(s) \hat{K}_0(s) + \hat{r}_1(s) \hat{K}_1(s)]$$

$$\text{and } \sup_{s \geq -\beta_0} \hat{W}(s) > \frac{2}{a^2 + b^2}.$$

(iv) $h \in C_1$.

Then there exists a unique solution $x(t; \omega)$ of (4.1) such that

$$\sup_{t \geq 0} \|x(t; \omega)\|_2 \leq M \sup_{t \geq 0} \|h(t; \omega)\|_2$$

for some $M > 0$.

Before we proceed with the proof of the theorem, we shall first prove the following lemma.

Lemma 4.1. Let T_1^* and T_2^* be operators on C_∞ defined by

$$T_1^* v(t) = \int_0^t (r_1 * K_1)^2(t-\tau) v(\tau) d\tau, \quad (4.5)$$

$$T_2^* v(t) = \int_0^t (r_2 * K_2)^2(t-\tau) v(\tau) d\tau. \quad (4.6)$$

Let T_1 and T_2 be operators on C_1 defined by

$$T_1 x(t; \omega) = \int_0^t r_1(t-\tau) \int_0^\tau K_1(\tau-u) x(u; \omega) d(z_1(u; \omega) - \mu \lambda) du, \quad (4.7)$$

$$T_2 x(t; \omega) = \int_0^t r_2(t-\tau) \int_0^\tau K_2(\tau-u) x(u; \omega) dz_2(u; \omega) d\tau, \quad (4.8)$$

where r_1, r_2, K_1, K_2, z_1 and z_2 are as in equation (4.1). Assume that there exist constants $c_1, c_2 > 0$ such that $\|c_1 T_1 + c_2 T_2\| > 1$ and $(I - c_1^2 T_1^* - c_2^2 T_2^*)$ is invertible. Then the operator $(I - c_1 T_1 - c_2 T_2)$ is invertible as a map from C_1 to C_1^* ($C_1^* = \text{Range of } T_1 + T_2$) and

$$\|(I - c_1 T_1 - c_2 T_2)^{-1}\|^2 \leq \|(I - c_1^2 T_1^* - c_2^2 T_2^*)^{-1}\|.$$

Proof. The assumption that $\|c_1 T_1 + c_2 T_2\| > 1$ implies that for some $\lambda(t) > 1$, there is a $y^* \in C_1$ such that

$$\|(c_1 T_1 + c_2 T_2) y^*\|_2 = \lambda(t) \|y^*\|_2. \quad (4.9)$$

$$\text{Let } A \equiv \{y^* \mid \|(c_1 T_1 + c_2 T_2) y^*\|_2 = \lambda(t) \|y^*\|_2\}. \quad (4.10)$$

$$\text{Let } x, y \in C_1 \text{ such that } y = x + c_1 T_1 y + c_2 T_2 y. \quad (4.11)$$

Then

$$\|y\|_2^2 \leq \|x\|_2^2 - \|c_1 T_1 y\|_2^2 - \|c_2 T_2 y\|_2^2 + 2\|y\|_2 [\|c_1 T_1 y + c_2 T_2 y\|_2]. \quad (4.12)$$

Choose $y^* \in A$ such that $\|y^*\|_2 = \|y\|_2$. Then

$$\begin{aligned} \|y\|_2^2 &= \|y^*\|_2^2 \leq \|x\|_2^2 - \|c_1 T_1 y^*\|_2^2 - \|c_2 T_2 y^*\|_2^2 \\ &\quad + 2\|c_1 T_1 y^* + c_2 T_2 y^*\|_2^2 [\lambda(t)]^{-1} \\ &\leq \|x\|_2^2 + (c_1^2 T_1^* + c_2^2 T_2^*) \|y\|_2^2. \end{aligned} \quad (4.13)$$

Hence,

$$\|y\|_2^2 \leq \|(I - c_1^2 T_1^* - c_2^2 T_2^*)^{-1}\| \|x\|_2^2. \quad (4.14)$$

Equations (4.11) and (4.14) allow us to conclude that

$$\|(I - c_1 T_1 - c_2 T_2)^{-1}\|^2 \leq \|(I - c_1^2 T_1^* - c_2^2 T_2^*)^{-1}\|.$$

To show that $(I - c_1 T_1 - c_2 T_2)$ has an inverse, we note that the operator is an onto map from C_1 to C_1^* . Therefore, we have only shown that it is one-to-one. Let $y \in C_1^*$ such that $y = (I - c_1 T_1 - c_2 T_2)x_1 = (I - c_1 T_1 - c_2 T_2)x_2$ for some $x_1 \neq x_2$. Then

$$E(x_1 - x_2)^2 - [c_1^2 T_1^* + c_2^2 T_2^*] E(x_1 - x_2)^2 = 0. \quad (4.15)$$

From the invertibility of the operator $(I - c_1^2 T_1^* - c_2^2 T_2^*)$ it follows that $x_1 = x_2$ almost surely, which shows that the map $(I - c_1 T_1 - c_2 T_2)$ has an inverse. ///

Proof of Theorem 4.1. Rewrite equation (4.1) as

$$x(t; \omega) = \bar{h}(t; \omega) + \sum_{i=0}^1 \bar{T}_i \bar{f}_i(t, x(t; \omega)) + \sum_{i=2}^3 \bar{T}_i f_i(t, x(t; \omega)), \quad (4.16)$$

where

$$\begin{aligned} \bar{h} &= (I - c T_0 - c T_1)^{-1} h, \\ \bar{T}_i &= (I - c T_0 - c T_1)^{-1} T_i, \quad i = 0, 1, 2, 3 \\ \bar{f}_i(t, x) &= f_i(t, x) - c x, \quad i = 0, 1, \text{ and } c = \frac{a+b}{2} \end{aligned} \quad (4.17)$$

The operators T_0, T_2 and T_3 are defined as in Theorem 3.3 (with kernels being convolution) and T_1 is defined by

$$T_1 x(t; \omega) = \int_0^t r_1(t-\tau) \int_0^\tau K_1(\tau-u) x(u; \omega) \mu \lambda \, du d\tau. \quad (4.18)$$

The existence of $(I - cT_0 - cT_1)^{-1}$ follows from assumption (i) of the theorem [3].

We shall now show that the assumptions of Theorem 4.1 imply those of Theorem 3.3. Condition (ii) of the theorem implies that the inverse of $(I - c^2 \bar{T}_1^* - c^2 \bar{T}_2^*)$ exists [3]. Therefore, from Lemma 4.1 it follows that the operator $(I - c(I - c(T_0+T_1))^{-1}(T_1+T_2))$ is invertible. Further, condition (iii) [3] and Lemma 4.1 together imply that

$$\begin{aligned} & \frac{(b-a)}{2} \left[\left\| (I - c(I - c(T_0+T_1))^{-1}(T_1+T_2))^{-1} \right\| \right. \\ & \quad \times \left[\left\| (I - c(T_0+T_1))^{-1}(T_0+T_1) \right\| \right. \\ & \quad \left. \left. + \left\| (I - c(T_0+T_1))^{-1}(T_1+T_2) \right\| \right] \right] < 1 \end{aligned}$$

The other conditions are easily verified and the conclusion follows from Theorem 3.3. ///

References

- [1] M. Altman, Inverse differentiability contractors and equations in Banach space, Studia Mathematica 46 (1973), 1-15.
- [2] A. T. Bharucha-Reid, Random Integral Equations, Academic Press, New York, 1972.
- [3] G. L. Blankenship, Frequency domain stability criteria for stochastic nonlinear feedback systems, SIAM J. Control and Optimization 14 (1976), 1107-1123.
- [4] A. C. H. Lee and W. J. Padgett, Random contractors and the solution of random nonlinear equations, J. Nonlinear Analysis, Theory Meth. and Appl. 1 (1977), 175-185.
- [5] E. J. McShane, Stochastic Calculus and Stochastic Models, Academic Press, New York, 1974.
- [6] W. J. Padgett and A.N.V. Rao, Existence and stability of solutions for generalized McShane systems, J. Integral Equations 1 (1979), 47-59.
- [7] W. J. Padgett and C. P. Tsokos, On a stochastic integro-differential equation of Volterra type, SIAM J. Appl. Math. 23 (1972), 499-512.
- [8] A.N.V. Rao and C. P. Tsokos, On the existence, uniqueness and stability behavior of a random solution to a nonlinear perturbed stochastic integro-differential equation, Information and Control 27 (1975), 61-74.
- [9] C. P. Tsokos and W. J. Padgett, Random Integral Equations with Applications to Life Sciences and Engineering, Academic Press, New York, 1974.

END

DATE
FILMED

1. 8/4

DTIC