

AD-A093 609

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER  
SOME APPROXIMATION PROPERTIES IN ORLICZ-SOBOLEV SPACES. (U)

F/6 12/1

UNCLASSIFIED

OCT 80 J GOSSEZ  
MRC-TSR-2126

DAAG29-80-C-0041  
NL

1 of 1  
ADA  
09 1673



END  
DATE  
FILMED  
2-81  
DTIC

AD A 093609

MRC Technical Summary Report # 2126

*P<sub>g</sub>*  
*P<sub>g</sub>*

SOME APPROXIMATION PROPERTIES  
IN ORLICZ-SOBOLEV SPACES

Jean-Pierre Gossez

LEVEL 11

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

SPIC  
JAN 9 1981  
C

October 1980

(Received June 13, 1980)

Approved for public release  
Distribution unlimited

DDC FILE COPY

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

80 12 22 039

(14) MFZ-T-R-2224 (11) + 8A

(15) 271

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

(1)

SOME APPROXIMATION PROPERTIES IN

ORLICZ-SOBOLEV SPACES.

10) Jean-Pierre/Gossez <sup>97</sup>

9) Technical Summary Report # 2126

October 1980

(15) D. 29-89-2-0041

ABSTRACT

We prove that weak derivatives in general Orlicz spaces are globally strong derivatives with respect to the modular convergence. Other approximation theorems involving the modular convergence are presented, which improve known density results of interest in the existence theory for strongly nonlinear boundary value problems.

AMS (MOS) Subject Classifications: 46E35, 46E30.

Key Words: weak and strong derivatives, Orlicz spaces, Orlicz-Sobolev spaces, modular convergence, approximation theorem, strongly nonlinear boundary value problem.

Work Unit Number 1 - Applied Analysis

---

(1) Département de Mathématique, C.P. 214, Université Libre de Bruxelles, 1050 Bruxelles, Belgium.

---

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

### SIGNIFICANCE AND EXPLANATION

One classical result in partial differential equations is the equality of weak derivatives (i.e. derivatives in the distribution sense) and strong derivatives (i.e. derivatives obtained from smooth functions by a limiting process). Such an equality holds in the setting of  $L^p$  spaces, as was proved locally by Friedrichs in 1944 and later globally by Meyers-Serrin in 1964.

Orlicz spaces are generalizations of  $L^p$  spaces where, roughly speaking, the defining function  $t \rightarrow |t|^p$  is replaced by a function whose growth at infinity is not necessarily of polynomial type. They have been successfully used in recent years in the study of several questions from partial differential equations, as the limiting case of the Sobolev imbedding theorem or the existence theory for strongly nonlinear boundary value problems. In the latter theory, problems related to the equality of weak and strong derivatives in the setting of Orlicz spaces play an important role. Such an equality does not hold for general Orlicz spaces. The main purpose of this paper is to provide an adequate substitute by slightly modifying the limiting process involved in the definition of strong derivatives.

|               |                                     |
|---------------|-------------------------------------|
| Accession For |                                     |
| RIIS GRAB     | <input checked="" type="checkbox"/> |
| ERIC TR       | <input type="checkbox"/>            |
| Unannounced   | <input type="checkbox"/>            |
| Justification | <input type="checkbox"/>            |
| By _____      |                                     |
| Distribution/ |                                     |
| A             |                                     |

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

SOME APPROXIMATION PROPERTIES IN  
ORLICZ-SOBOLEV SPACES

Jean-Pierre Gossez<sup>(1)</sup>

STATEMENT OF RESULTS

Let  $L_M(\Omega)$  be the Orlicz space on an open subset  $\Omega$  of  $\mathbb{R}^N$  corresponding to a  $N$ -function  $M$  and let  $E_M(\Omega)$  be the norm closure in  $L_M(\Omega)$  of the  $L^\infty(\Omega)$  functions with compact support in  $\bar{\Omega}$ . The Sobolev space of functions  $u$  such that  $u$  and its distributional derivatives up to order  $m$  lie in  $L_M(\Omega)$  (resp.  $E_M(\Omega)$ ) is denoted by  $W^m L_M(\Omega)$  (resp.  $W^m E_M(\Omega)$ ). Standard references about these spaces include [11,1,12].

A well-known theorem of Meyers-Serrin [13] states that for  $1 < p < \infty$ ,  $C^\infty(\Omega) \cap W^{m,p}(\Omega)$  is norm dense in  $W^{m,p}(\Omega)$ , i.e. weak derivatives in  $L^p(\Omega)$  are globally strong derivatives (the local version goes back to Friedrichs and his mollifiers [7]). This was extended to the Orlicz spaces setting by Donaldson-Trudinger [4] who proved that  $C^\infty(\Omega) \cap W^m E_M(\Omega)$  is norm dense in  $W^m E_M(\Omega)$ . The corresponding statement with  $E_M$  replaced by  $L_M$  is not true, even locally, simply because a  $L_M(\Omega)$  function may not belong to  $E_M(\Omega')$  for  $\Omega' \subset\subset \Omega$  (take  $\Omega = ]-1,+1[$ ,  $M(t) = e^{t^2} - 1$  and  $u(x) = (\log |x|^{-1})^{1/2}$ ). Our first result concerns the density of  $C^\infty(\Omega) \cap W^m L_M(\Omega)$  in  $W^m L_M(\Omega)$  with respect to a weaker convergence, the so-called modular convergence [14].

**THEOREM 1.** Let  $u \in W^m L_M(\Omega)$ . Then there exist  $\lambda > 0$  and a sequence  $u_k \in C^\infty(\Omega) \cap W^m L_M(\Omega)$  such that for  $|\alpha| \leq m$ ,  $\int_\Omega M((D^\alpha u_k - D^\alpha u)/\lambda) \rightarrow 0$  as  $k \rightarrow \infty$ .

---

(1) Département de Mathématique, C.P. 214, Université Libre de Bruxelles, 1050 Bruxelles, Belgium.

As will be seen in the proof, it suffices to choose  $\lambda$  such that  $16/\lambda D^\alpha u \in L_M(\Omega)$  for  $|\alpha| = m$ , where  $L_M(\Omega)$  denotes the Orlicz class. Consequently, when  $u \in W^m E_M(\Omega)$ ,  $\lambda$  can be taken arbitrary small, and we recover from theorem 1 the result of [4] mentioned above.

The space  $W^m L_M(\Omega)$  will, as usual, be identified to a subspace of the product  $\prod_{|\alpha| \leq m} L_M(\Omega) \equiv \Pi L_M$ . Denoting by  $\bar{M}$  the N-function conjugate to  $M$ , we consider the weak topologies  $\sigma(\Pi L_M, \Pi E_{\bar{M}})$  and  $\sigma(\Pi L_M, \Pi L_{\bar{M}})$ . Density results involving the latter play an important role in the existence theory for strongly nonlinear boundary value problems (cf. [8,3]). Comparing the modular convergence with  $\sigma(\Pi L_M, \Pi L_{\bar{M}})$  (lemma 6 below), we obtain the following

COROLLARY 2.  $C^\infty(\Omega) \cap W^m L_M(\Omega)$  is  $\sigma(\Pi L_M, \Pi L_{\bar{M}})$  dense in  $W^m L_M(\Omega)$ .

We now turn to the approximation by functions which are smooth up to the boundary, assuming some regularity on  $\Omega$ . Recall that  $\Omega$  is said to have the segment property when there exist an open covering  $\{U_i\}$  of  $\bar{\Omega}$  and corresponding vectors  $\{y_i \in \mathbb{R}^N\}$  such that for  $x \in \bar{\Omega} \cap U_i$  and  $0 < t < 1$ ,  $x + ty_i \in \Omega$ ;  $\Omega$  has the cone property when there exists a finite cone  $C$  such that each  $x \in \Omega$  is the vertex of a cone  $C_x \subset \Omega$  congruent to  $C$ . It was proved in [8] that if  $\Omega$  has the segment property, then  $\mathcal{D}(\bar{\Omega})$  is  $\sigma(\Pi L_M, \Pi L_{\bar{M}})$  dense in  $W^m L_M(\Omega)$ , where  $\mathcal{D}(\bar{\Omega})$  denotes the restrictions to  $\Omega$  of the functions in  $\mathcal{D}(\mathbb{R}^N)$ . This is improved in our next result under the assumption that  $\Omega$  also has the cone property.

**THEOREM 3.** Suppose that  $\Omega$  satisfies both the segment and the cone property. Let  $u \in W^m L_M(\Omega)$ . Then there exist  $\lambda > 0$  and a sequence  $u_k \in \mathcal{D}(\bar{\Omega})$  such that for  $|\alpha| \leq m$ ,  $\int_{\Omega} M((D^\alpha u_k - D^\alpha u)/\lambda) \rightarrow 0$  as  $k \rightarrow \infty$ .

The proof will show that it suffices to choose  $\lambda$  such that  $16(N+1)/\lambda D^\alpha u \in L_M(\Omega)$  for  $|\alpha| \leq m$ . Moreover the cone property is only used to guarantee that an element  $v \in W^1 L_M(\Omega)$  with compact support in  $\bar{\Omega}$  lies in  $E_M(\Omega)$  (the imbedding theorem of [4] implies  $v \in C(\Omega) \cap L^\infty(\Omega)$  or  $v \in L_{M^*}(\Omega)$  with  $M^*$  a  $N$ -function which, by lemma 4.14 of [8], increases essentially more rapidly than  $M$ , and so, in any case,  $v \in E_M(\Omega)$ ). This fact probably holds under a weaker assumption on  $\Omega$ . Anyway, taking again  $\lambda$  arbitrary small, we recover from theorem 3 the result of [4] that if  $\Omega$  has the segment property, then  $\mathcal{D}(\bar{\Omega})$  is norm dense in  $W^m E_M(\Omega)$ .

Finally we consider the analogue of the  $W_0^{m,p}$  spaces.  $W_0^m L_M(\Omega)$  is defined as the  $\sigma(\Pi L_M, \Pi E_M)$  closure of  $\mathcal{D}(\Omega)$  in  $W^m L_M(\Omega)$  and  $W_0^m E_M(\Omega)$  as the norm closure of  $\mathcal{D}(\Omega)$  in  $W^m L_M(\Omega)$  (or equivalently in  $W^m E_M(\Omega)$ ). When  $\partial\Omega$  is sufficiently regular, one can define the trace on  $\partial\Omega$  of  $D^\alpha u$  for  $u \in W_0^m L_M(\Omega)$  and  $|\alpha| \leq m-1$ , and prove that the functions in  $W_0^m L_M(\Omega)$  (resp.  $W_0^m E_M(\Omega)$ ) are precisely those in  $W^m L_M(\Omega)$  (resp.  $W^m E_M(\Omega)$ ) whose trace and normal derivative up to order  $m-1$  on  $\partial\Omega$  vanish (cf. [6,9]).

**THEOREM 4.** Suppose that  $\Omega$  satisfies the segment property. Let  $u \in W_0^m L_M(\Omega)$ . Then there exist  $\lambda > 0$  and a sequence  $u_k \in \mathcal{D}(\Omega)$  such that for  $|\alpha| \leq m$ ,  $\int_{\Omega} M((D^\alpha u_k - D^\alpha u)/\lambda) \rightarrow 0$  as  $k \rightarrow \infty$ .

The same estimate on  $\lambda$  as in theorem 3 holds here. Theorem 4 improves our result of [8] that if  $\Omega$  has the segment property, then  $\mathcal{D}(\Omega)$  is  $\sigma(\Pi L_M, \Pi L_{\bar{M}})$  dense in  $W_0^m L_M(\Omega)$ .

The proofs of the  $W^m E_M(\Omega)$  results of [4] referred to above as well as those of the  $\sigma(\Pi L_M, \Pi L_{\bar{M}})$  versions of theorems 3 and 4 given in [8] are rather simple modifications of the standard  $L^p$  proofs. To get corollary 2 or theorem 1, 3 and 4 requires more involved calculations, although the construction of the approximants is basically the same. An inequality which will be used repeatedly is

$$M\left(\sum_i r_i\right) \leq r^{-1} \sum_i M(r r_i)$$

where  $r_i \in \mathbb{R}$  and  $r$  is the maximum number of nonzero  $r_i$ 's. When the  $r_i$ 's originate from a partition of unity, a control on  $r$  can be obtained from simple topological considerations (lemma 7 below).

PROOFS.

Given a function  $u(x)$ , we denote by  $u_y$  its translate  $u_y(x) = u(x-y)$ , by  $u_\delta$  its regularization  $u_\delta = u * \rho_\delta$  where  $\rho_\delta \in \mathcal{D}(\mathbb{R}^N)$ ,  $\rho_\delta(x) = 0$  for  $|x| > \delta$ ,  $\rho_\delta > 0$  and  $\int_{\mathbb{R}^N} \rho_\delta = 1$ , and by  $u_r$  the function  $u_r = \varphi_r u$  where  $\varphi_r(x) = \varphi(x/r)$  with  $\varphi \in C_c^\infty(\mathbb{R}^N)$ ,  $0 \leq \varphi \leq 1$ ,  $\varphi(x) = 1$  for  $|x| \leq 1$  and  $\varphi(x) = 0$  for  $|x| > 2$ . The following lemma will be needed.

LEMMA 5. If  $u \in L_M(\mathbb{R}^N)$  with  $2u \in L_M(\mathbb{R}^N)$ , then  $\int_{\mathbb{R}^N} M(u_y - u) \rightarrow 0$  as  $|y| \rightarrow 0$  and  $\int_{\mathbb{R}^N} M(u_\delta - u) \rightarrow 0$  as  $\delta \rightarrow 0$ . If  $u \in L_M(\mathbb{R}^N)$ , then  $\int_{\mathbb{R}^N} M(u_r - u) \rightarrow 0$  as  $r \rightarrow \infty$ .



PROOF. We consider  $u_\delta$ ; the case of  $u_\gamma$  is treated similarly and that of  $u_r$  is immediate. Since  $u_\delta \rightarrow u$  in  $L^1_{loc}(\mathbb{R}^N)$ , one has, for a subsequence,  $u_\delta \rightarrow u$  a.e. and so  $M(u_\delta - u) \rightarrow 0$  a.e. Moreover

$$M(u_\delta - u) < \frac{1}{2} M(2u_\delta) + \frac{1}{2} M(2u) < \frac{1}{2} (M(2u))_\delta + \frac{1}{2} M(2u)$$

where we have used Jensen's integral inequality. Since  $(M(2u))_\delta \rightarrow M(2u)$  in  $L^1(\mathbb{R}^N)$ , we conclude by Vitali's theorem that  $M(u_\delta - u) \rightarrow 0$  in  $L^1(\mathbb{R}^N)$ .

Q.E.D.

Thus we see that for any  $u \in L_M(\mathbb{R}^N)$ ,  $u_\delta \rightarrow u$  with respect to the modular convergence (i.e. there exists  $\lambda > 0$  such that  $\int_{\mathbb{R}^N} M((u_\delta - u)/\lambda) \rightarrow 0$ ). In general  $u_\delta$  does not converge to  $u$  in the mean (i.e. the above with  $\lambda = 1$ ) and a fortiori in norm. When  $u \in E_M(\mathbb{R}^N)$ , then  $2u/\lambda \in L_M(\mathbb{R}^N)$  for all  $\lambda > 0$  and by taking  $\lambda$  arbitrary small, we eventually derive from lemma 5 that  $u_\delta$  converges to  $u$  in norm, a result originally given in [2]. Similar remarks apply to  $u_\gamma$ .

PROOF OF THEOREM 1. Let  $u \in W^m L_M(\Omega)$  and choose  $\lambda > 0$  such that  $16/\lambda D^\alpha u \in L_M(\Omega)$  for  $|\alpha| = m$ . Let  $\varepsilon > 0$ ,  $\varepsilon < 1$ . We will prove that there exists  $v \in C^\infty(\Omega) \cap W^m L_M(\Omega)$  such that

$$(1) \quad \int_{\Omega} M((D^\alpha v - D^\alpha u)/\lambda) < \varepsilon$$

for  $|\alpha| < m$ .

Define for  $i = 1, 2, \dots$ ,

$$\Omega_i = \{x \in \Omega; |x| < i \text{ and } \text{dist}(x, \partial\Omega) > 1/i\} ,$$

and also, for convenience,  $\Omega_{-2} = \Omega_{-1} = \Omega_0 = \emptyset$ . Let  $\{\psi_i; i = 1, 2, \dots\}$  be a  $C^\infty$  partition of unity on  $\Omega$  such that  $\text{supp } \psi_i \subset \Omega_{i+1} \setminus \Omega_{i-1}$ . For each  $i = 1, 2, \dots$ , let  $\rho_{\delta_i} \equiv \rho_i$  be a mollifier satisfying

$$(2) \quad \delta_i < 1/(i+1)(i+2) ,$$

$$(3) \quad \|(D^\beta \psi_i D^\gamma u) * \rho_i - D^\beta \psi_i D^\gamma u\|_{M, \Omega} < \epsilon \lambda / a 2^{i+1}$$

for all  $|\beta + \gamma| \leq m$  with  $|\gamma| < m$ , and

$$(4) \quad \int_{\Omega} M(\beta((\psi_i D^\alpha u) * \rho_i - \psi_i D^\alpha u) / \lambda) < \epsilon / 2^i$$

for  $|\alpha| = m$ . Here  $\|\cdot\|_{M, \Omega}$  denotes the Luxemburg norm in  $L_M(\Omega)$ :

$$\|w\|_{M, \Omega} = \inf\{h > 0; \int_{\Omega} M(w/h) < 1\}$$

and the number  $a$  is defined in terms of the coefficients which appear in Leibnitz formula:

$$(5) \quad a = \max\left\{ \sum_{\beta \leq \alpha} \binom{\alpha}{\beta}; |\alpha| \leq m \right\} .$$

Condition (3) can be fulfilled because on a smooth bounded domain,  $W^m L_M$  is (compactly) imbedded into  $W^{m-1} E_M$  (see the introduction) and so  $D^\gamma u$ ,  $|\gamma| < m$ , lies in  $E_M^{loc}(\Omega)$ . Condition (4) can be fulfilled by applying lemma 5. It follows from (2) that  $(\psi_i u) * \rho_i$  has support in  $\Omega_{i+2} \setminus \Omega_{i-2}$ . Thus the series

$$v = \sum_{i=1}^{\infty} (\psi_i u) * \rho_i$$

is trivially convergent and  $v \in C^\infty(\Omega)$ . The fact that  $v \in W^m L_M(\Omega)$  will follow once (1) is proved.

To verify (1), take  $j = 1, 2, \dots$  and write

$$\begin{aligned} \int_{\Omega_j} M((D^\alpha v - D^\alpha u)/\lambda) &= \int_{\Omega_j} M\left(\sum_{i=1}^{j+1} (D^\alpha(\psi_i u) * \rho_i - D^\alpha(\psi_i u))/\lambda\right) \\ &< 2^{-1} \int_{\Omega_j} M\left(2 \sum_{i=1}^{j+1} \sum_{\beta+\gamma=\alpha, |\gamma|<m} \binom{\alpha}{\beta} ((D^\beta \psi_i D^\gamma u) * \rho_i - D^\beta \psi_i D^\gamma u)/\lambda\right) \\ &\quad + 2^{-1} \int_{\Omega_j} M\left(2 \sum_{i=1}^{j+1} ((\psi_i D^\alpha u) * \rho_i - \psi_i D^\alpha u)/\lambda\right) = I_1 + I_2, \end{aligned}$$

where the term  $I_2$  does not appear when  $|\alpha| < m$ . We have

$$\begin{aligned} I_1 &< \varepsilon 2^{-1} a^{-1} \sum_{i=1}^{j+1} 2^{-i} \sum_{\beta+\gamma=\alpha, |\gamma|<m} \binom{\alpha}{\beta} M(2^{i+1} a ((D^\beta \psi_i D^\gamma u) * \rho_i - D^\beta \psi_i D^\gamma u)/\varepsilon \lambda) \\ &< \varepsilon/2 \end{aligned}$$

by (3) and the definition of the Luxemburg norm. To study  $I_2$ , we observe that for a.e.  $x \in \Omega_j$ ,

$$M(2 \sum_{i=1}^{j+1} ((\psi_i D^\alpha u) * \rho_i - \psi_i D^\alpha u) / \lambda) < \frac{1}{4} \sum_{i=1}^{j+1} M(8((\psi_i D^\alpha u) * \rho_i - \psi_i D^\alpha u) / \lambda)$$

since at most 4 terms of the sum are nonzero at  $x$ . Consequently, using (4), we get  $I_2 < \epsilon/8$ . So

$$\int_{\Omega_j} M((D^\alpha v - D^\alpha u) / \lambda) < \epsilon$$

and letting  $j \rightarrow \infty$ , we obtain (1).

Q.E.D.

By a simple modification of this proof one can show that the  $u_k$ 's in the statement of theorem 1 can be taken so that  $D^\alpha u_k \rightarrow D^\alpha u$  in norm for  $|\alpha| < m$ .

Corollary 2 is a direct consequence of theorem 1 and the following

**LEMMA 6.** Let  $u_k, u \in L_M(\Omega)$ . If  $u_k \rightarrow u$  with respect to the modular convergence, then  $u_k \rightarrow u$  for  $\sigma(L_M, L_{\frac{1}{M}})$ .

**PROOF.** Let  $\lambda > 0$  be such that  $\int_{\Omega} M((u_k - u) / \lambda) \rightarrow 0$ . Thus, for a subsequence,  $u_k \rightarrow u$  a.e. in  $\Omega$ . Take  $v \in L_{\frac{1}{M}}(\Omega)$ . Multiplying  $v$  by a suitable constant, we can assume  $\lambda v \in L_{\frac{1}{M}}(\Omega)$ . By Young's inequality,

$$|(u_k - u)v| < M((u_k - u) / \lambda) + \bar{M}(\lambda v),$$

which implies, by Vitali's theorem, that  $\int_{\Omega} |(u_k - u)v| \rightarrow 0$ .

Q.E.D.

The proof of theorems 3 and 4 uses the following lemma from general topology.

LEMMA 7. Let  $A$  be a closed subset of  $R^N$  and let  $\{U_i\}$  be an open covering of  $A$ . Then  $\{U_i\}$  can be refined into a locally finite countable open covering  $\{V_j\}$  of  $A$  with the property that at most  $N + 1$  distinct  $V_j$ 's have a nonempty intersection.

Lemma 7 follows from the fact that the covering dimension of  $R^N$  is  $N$  (cf. [5]). All we need actually for our purposes is an estimate on the maximum number of intersecting  $V_j$ 's which is independent of  $A$  and of the covering  $\{U_i\}$ . Such an estimate can be obtained directly by elementary means, as we now briefly indicate.

Recall Lebesgue's lemma (cf. [10]): given an open covering of a compact set  $K$  in a metric space, there exists  $r > 0$  such that the open  $r$ -ball about each point of  $K$  is contained in some member of the covering. We apply this lemma successively to the covering  $\{U_i\}$  and to the sets

$A \cap \bar{B}(0,1)$ ,  $A \cap (\bar{B}(0,2) \setminus B(0,1))$ , ... and get a (possibly finite) sequence of radius  $r_1, r_2, r_3, \dots$  which clearly can be taken such that each  $r_p < 1/3$  and  $r_1 > r_2 > \dots$ . Now we cover  $A \cap \bar{B}(0,1)$  with a finite number of balls  $B(y_{1,\ell}, r_1)$  chosen in such a way that  $y_{1,\ell} \in A \cap \bar{B}(0,1)$  but  $y_{1,\ell} \notin B(y_{1,1}, r_1) \cup \dots \cup B(y_{1,\ell-1}, r_1)$ . Next we cover  $A \cap (\bar{B}(0,2) \setminus B(0,1))$  with the balls  $B(y_{1,\ell}, r_1)$  and a finite number of balls  $B(y_{2,k}, r_2)$  chosen in such a way that  $y_{2,k} \in A \cap (\bar{B}(0,2) \setminus B(0,1))$  but  $y_{2,k} \notin \cup_{\ell} B(y_{1,\ell}, r_1)$  and  $y_{2,k} \notin B(y_{2,1}, r_2) \cup \dots \cup B(y_{2,k-1}, r_2)$ . And so on. The family  $\mathcal{B}$  of the balls  $B(y_{p,q}, r_p)$  constructed in this way constitutes a locally finite

countable open covering of  $A$  and is a refinement of  $\{U_i\}$ . Moreover, since each  $r_p$  is  $< 1/3$ , a ball  $B(y_{p,q}, r_p)$  can only intersect either balls with radius  $r_p$  and  $r_{p-1}$  or balls with radius  $r_p$  and  $r_{p+1}$ . Also, since  $r_1 > r_2 > \dots$ , the only ball of the family  $\mathcal{B}$  to which  $y_{p,q}$  belongs is  $B(y_{p,q}, r_p)$ . Using these facts it is then easy to verify that any subfamily of  $\mathcal{B}$  with nonempty intersection contains at most  $2d_N$  members, where  $d_N$  denotes the maximum number of points  $x_s$  which we put inside the unit ball  $B(0,1)$  of  $\mathbb{R}^N$  with  $|x_s - x_t| > 1$  for  $s \neq t$ .

PROOF OF THEOREM 3. Let  $u \in W_{L_M}^m(\Omega)$  and choose  $\lambda > 0$  such that  $16(N+1)/\lambda D^\alpha u \in L_M(\Omega)$  for  $|\alpha| \leq m$ . We will show that for  $|\alpha| \leq m$ ,

$$(6) \quad \int_{\Omega} M(2(D^\alpha u - D^\alpha u_r)/\lambda) \rightarrow 0$$

as  $r \rightarrow \infty$ , where  $u_r = \varphi_r u$  is the function involved in lemma 5. We will also show that for each  $r$  and for each  $\eta > 0$ ,  $\eta \leq 1$ , there exists  $v \in \mathcal{D}(\bar{\Omega})$  such that for  $|\alpha| \leq m$ ,

$$(7) \quad \int_{\Omega} M(2(D^\alpha u_r - D^\alpha v)/\lambda) \leq \eta .$$

The conclusion of theorem 3 then follows easily.

To verify (6), we write

$$\int_{\Omega} M(2(D^{\alpha}u - D^{\alpha}u_r)/\lambda)$$

$$\leq 2^{-1} \int_{\Omega} M(4(D^{\alpha}u - \varphi_r D^{\alpha}u)/\lambda)$$

$$+ 2^{-1} a^{-1} \sum_{\beta+\gamma=\alpha, |\beta|>0} \binom{\alpha}{\beta} \int_{\Omega} M(4a r^{-|\beta|} (D^{\beta}\varphi)(x/r) D^{\gamma}u/\lambda) .$$

Here  $a$  is the number defined by (5). Each term on the right hand side goes to zero as  $r \rightarrow \infty$ , the first one by lemma 5 and the second one by direct examination.

To verify (7) we use the covering  $\{U_i\}$  of  $\bar{\Omega}$  given by the segment property. Refining  $\{U_i\}$  if necessary, we can assume that it satisfies the properties stated in lemma 7. In particular  $i = 1, 2, \dots$ . Let  $\{\psi_i\}$  be a  $C^{\infty}$  partition of unity on  $\bar{\Omega}$  subordinate to  $\{U_i\}$ . Clearly  $\text{supp } \psi_i \subset U_i'$  for some open set  $U_i'$  with compact closure  $\bar{U}_i' \subset U_i$ . Write  $\Gamma_i = \bar{U}_i' \cap \partial\Omega$  and  $\Gamma_{i,t} = \Gamma_i - ty_i$  where  $y_i$  is the vector associated to  $U_i$  by the segment property. Extend  $u_r$  outside  $\Omega$  by zero and note that  $\psi_i u_r$  vanishes identically for  $i >$  some  $i_r$ . Note also that the translate  $(\psi_i u_r)_t(x) \equiv (\psi_i u_r)(x + ty_i)$ ,  $0 < t < 1$ , belongs to  $W_{L,M}^m(\mathbb{R}^N \setminus \Gamma_{i,t})$  and that, by the segment property,  $\text{dist}(\Gamma_{i,t}, \Omega) > 0$ . For each  $i = 1, 2, \dots, i_r$ , choose  $0 < t_i < 1$  and  $\rho_{\delta_i} \equiv \rho_i$ ,  $\delta_i < \text{dist}(\Gamma_{i,t_i}, \Omega)$ , such that

$$(8) \quad \|(D^{\beta}\psi_i D^{\gamma}u_r)_{t_i} * \rho_i - D^{\beta}\psi_i D^{\gamma}u_r\|_{M,\Omega} \leq \lambda\eta/2^{i+2}a$$

for all  $|\beta + \gamma| \leq m$  with  $|\gamma| < m$ , and

$$(9) \quad \int_{\Omega} M(4(N+1)((\psi_i D^{\alpha}u_r)_{t_i} * \rho_i - \psi_i D^{\alpha}u_r)/\lambda) \leq \eta/2^i$$

for  $|\alpha| = m$ . Condition (8) can be fulfilled because  $D^Y u_r \in E_M(\Omega)$  for  $|\gamma| < m$  (the cone property is used here, see the introduction). Condition (9) can be fulfilled by two consecutive applications of lemma 5. Taking  $t_i$  and  $\delta_i$  smaller if necessary, we can assume that  $\text{supp}(\psi_i u_r)_{t_i} * \rho_i \subset U_i$ .

Define

$$(10) \quad v = \sum_{i=1}^r (\psi_i u_r)_{t_i} * \rho_i \in \mathcal{D}(\bar{\Omega})$$

and observe that by the property of lemma 7, at each  $x \in \Omega$ , the sum above contains at most  $(N + 1)$  nonzero terms. We have

$$\begin{aligned} \int_{\Omega} M(2(D^\alpha v - D^\alpha u_r)/\lambda) &= \int_{\Omega} M(2 \sum_{i=1}^r ((D^\alpha(\psi_i u_r))_{t_i} * \rho_i - D^\alpha(\psi_i u_r))/\lambda) \\ &< 2^{-1} \int_{\Omega} M(4 \sum_{i=1}^r \sum_{\beta+\gamma=\alpha, |\gamma|<m} \binom{\alpha}{\beta} ((D^\beta \psi_i D^\gamma u_r)_{t_i} * \rho_i - D^\beta \psi_i D^\gamma u_r)/\lambda) \\ &\quad + 2^{-1} \int_{\Omega} M(4 \sum_{i=1}^r ((\psi_i D^\alpha u_r)_{t_i} * \rho_i - \psi_i D^\alpha u_r)/\lambda) = I_1 + I_2 \end{aligned}$$

where the term  $I_2$  does not appear when  $|\alpha| < m$ . Now

$$\begin{aligned} I_1 &< n 2^{-1} a^{-1} \sum_{i=1}^r 2^{-i} \sum_{\beta+\gamma=\alpha, |\gamma|<m} \int_{\Omega} M(2^{i+2} a ((D^\beta \psi_i D^\gamma u_r)_{t_i} * \rho_i - D^\beta \psi_i D^\gamma u_r)/\lambda) \\ &< n/2 \end{aligned}$$

by (8) and the definition of the Luxemburg norm. Since at a.e.  $x \in \Omega$ ,

$$\begin{aligned} M(4 \sum_{i=1}^r ((\psi_i D^\alpha u_r)_{t_i} * \rho_i - \psi_i D^\alpha u_r)/\lambda) \\ &< (N+1)^{-1} \sum_{i=1}^r M(4(N+1)((\psi_i D^\alpha u_r)_{t_i} * \rho_i - \psi_i D^\alpha u_r)/\lambda) \end{aligned}$$



we obtain from (9) that  $I_2 < \eta/2$ , and so (7) is proved.

Q.E.D.

PROOF OF THEOREM 4. It is essentially the same as that of theorem 3 except that one replaces in (10)  $t_i$  by  $-t_i$  and chooses  $\delta_i$  with

$$\delta_i < \text{dist} ((\text{supp } \psi_i \cap \bar{\Omega}) + t_i y_i, \mathbb{R}^N \setminus \Omega) .$$

One also uses the fact that by extending a  $W_0^{m, L_M}(\Omega)$  function by zero outside  $\Omega$ , one gets a  $W^{m, L_M}(\mathbb{R}^N)$  function. This allows us to avoid the cone property for  $\Omega$ .

Q.E.D.

We remark that when  $\Omega$  is bounded, the above arguments can be carried through without using lemma 7. The coefficient  $\lambda$  is then chosen so that  $8b/\lambda D^\alpha u \in L_M(\Omega)$  for  $|\alpha| = m$ , where  $b$  is the number of pieces of the covering  $\{U_i\}$  needed to cover  $\bar{\Omega}$ .

As for theorem 1, one can show by a simple modification of the proofs that the  $u_k$ 's in the statement of theorems 3 and 4 can be taken so that  $D^\alpha u_k \rightarrow D^\alpha u$  in norm for  $|\alpha| < m$ .

#### REFERENCES

1. R. ADAMS, Sobolev spaces, Ac. Press, 1975.
2. G. DANKERT, Sobolev imbedding theorems in Orlicz spaces, Thesis, Cologne, 1966.
3. T. DONALDSON, Inhomogeneous Orlicz-Sobolev spaces and nonlinear parabolic initial value problems, J. Diff. Eq., 16 (1974), 201-256.
4. T. DONALDSON and N. S. TRUDINGER, Orlicz-Sobolev spaces and imbedding theorems, J. Funct. Anal., 8 (1971), 52-75.
5. R. ENGELKING, Outline of general topology, North Holland and J. Wiley, 1968.
6. A. FOUGERES, Théorèmes de trace et de prolongement dans les espaces de Sobolev et de Sobolev-Orlicz, C.R. Ac. Sc. Paris, Serie A, 274 (1972), 181-184.
7. K. FRIEDRICHS, The identity of weak and strong extensions of differential operators, Trans. Amer. Math. Soc., 55 (1944), 132-151.
8. J. P. GOSSEZ, Nonlinear elliptic boundary value problems for equations with rapidly (or slowly) increasing coefficients, Trans. Amer. Math. Soc., 190 (1974), 163-205.
9. J. P. GOSSEZ, A remark on strongly nonlinear elliptic boundary value problems, Bol. Soc. Brasileira Mat., 8 (1977), 53-63.
10. J. KELLEY, General topology, Van Nostrand, 1955.
11. M. KRASNOSEL'SKII and J. RUTICKII, Convex functions and Orlicz spaces, Noordhoff, 1961.
12. A. KUFNER, O. JOHN and S. FUCIK, Function Spaces, Academia, Praha, 1977.
13. N. MEYERS and J. SERRIN,  $H = W$ , Proc. Nat. Ac. Sc. U.S.A., 51 (1964), 1055-1056.
14. J. MUSIELAK and W. ORLICZ, On modular spaces, Studia Mat., 18 (1959), 49-65.

JPG/jvs

| REPORT DOCUMENTATION PAGE  |  | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM |
|--|--|---|
| 1. REPORT NUMBER<br>#2126  | 2. GOVT ACCESSION NO.<br>AD A093 609   | 3. RECIPIENT'S CATALOG NUMBER               |
| 4. TITLE (and Subtitle)<br><br>SOME APPROXIMATION PROPERTIES IN ORLICZ-SOBOLEV SPACES  | 5. TYPE OF REPORT & PERIOD COVERED<br>Summary Report - no specific reporting period                        |   |
|  | 6. PERFORMING ORG. REPORT NUMBER   |   |
| 7. AUTHOR(s)<br><br>Jean-Pierre Gossez   | 8. CONTRACT OR GRANT NUMBER(s)<br><br>DAAG29-80-C-0041   |   |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Mathematics Research Center, University of<br>610 Walnut Street<br>Madison, Wisconsin 53706   | 10. PROGRAM ELEMENT, PROJECT, TASK<br>AREA & WORK UNIT NUMBERS<br>Work Unit Number 1 -<br>Applied Analysis |   |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>U. S. Army Research Office<br>P. O. Box 12211<br>Research Triangle Park, North Carolina 27709   | 12. REPORT DATE<br>October 1980  |   |
|  | 13. NUMBER OF PAGES<br>14  |   |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)  | 15. SECURITY CLASS. (of this report)<br><br>UNCLASSIFIED   |   |
|  | 15a. DECLASSIFICATION/DOWNGRADING<br>SCHEDULE  |   |
| 16. DISTRIBUTION STATEMENT (of this Report)<br><br>Approved for public release; distribution unlimited.  |  |   |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)   |  |   |
| 18. SUPPLEMENTARY NOTES  |  |   |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br><br>weak and strong derivatives, Orlicz spaces, Orlicz-Sobolev spaces, modular convergence, approximation theorem, strongly nonlinear boundary value problem.  |  |   |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br><br>We prove that weak derivatives in general Orlicz spaces are globally strong derivatives with respect to the modular convergence. Other approximation theorems involving the modular convergence are presented, which improve known density results of interest in the existence theory for strongly nonlinear boundary value problems.. |  |   |