

AD-A097 696

ROYAL AIRCRAFT ESTABLISHMENT FARNBOROUGH (ENGLAND)  
THE ORBIT OF 1972-05B IN ITS FINAL PHASE, WITH GEOPHYSICAL INFE--ETC(U)  
SEP 80 H HILLER

F/6 22/3

UNCLASSIFIED

RAE-TR-80119

DRIC-BR-77047

NL

1-1  
AD  
A097696



END  
DATE  
FILMED  
5 81  
DTIC

UNLIMITED

10/10

Technical report



RAE

ROYAL AIRCRAFT ESTABLISHMENT

RAE-TR

Technical Report 5075

11 September 1950

AD A 097 696

6 THE ORBIT OF 1972-658 IN ITS FINAL PHASE, WITH GEOPHYSICAL INFERENCES

by

19/50

16 H. Miller

Procurement Executive, Ministry of Aircraft Production

UNLIMITED

81

31/10/50

FILE COPY

UDC 521.3 : 521.6 : 551.557

ROYAL AIRCRAFT ESTABLISHMENT

Technical Report 80119

Received for printing 26 September 1980

THE ORBIT OF 1972-05B IN ITS FINAL PHASE, WITH GEOPHYSICAL INFERENCES

by

H. Hiller

SUMMARY

The polar orbit of Heos 2 second-stage rocket, 1972-05B, has been determined on each of the final 16 days before its decay in September 1978, using the RAE orbit refinement program, PROP 6, with about 1360 observations. An accuracy of 30-70 m, both radial and across track, was achieved.

Eleven values of density scale height have been determined from the decrease in perigee height, with a 2 per cent error; seven of these values are within 6 per cent of the *CIRA 1972* reference-atmosphere values, the rms value being 4 per cent higher than *CIRA*.

The rotation rate of the upper atmosphere,  $\Lambda$ , was determined from the decrease in orbital inclination as  $\Lambda = 1.40 \pm 0.05$  rev/day; *ie* a strong west-to-east zonal wind of  $160 \pm 20$  m/s, at a mean height of about 240 km. The local time was 01-02 h; solar activity was high; and the latitude of perigee moved steadily from  $10^{\circ}\text{N}$  to  $67^{\circ}\text{S}$ .

Departmental Reference: Space 587

Copyright

©

Controller HMSO London  
1980

LIST OF CONTENTS

	<u>Page</u>
1 INTRODUCTION	3
2 THE ORBIT	3
3 VARIATION IN PERIGEE HEIGHT	3
4 DENSITY SCALE HEIGHT	5
5 UPPER-ATMOSPHERE ZONAL WIND	7
5.1 Results	7
5.2 Discussion	8
6 CONCLUSIONS	9
Table 1 Orbital parameters for Heos 2 second-stage rocket, with standard deviations	4
References	10
Illustrations	Figures 1-9
Report documentation page	inside back cover

Accession For		
NIS	GRA&I	X
DTIC	TAB	
Unannounced		
Justification		
By		
Distribution/		
Availability Codes		
Avail and/or		
Dist	Special	
A		

## 1 INTRODUCTION

The ESRO highly-eccentric-orbit satellite Heos 2 (1972-05A) was launched into an orbit of eccentricity 0.95 on 1972 January 31 by NASA. The second-stage Ablestar rocket, 1972-05B, entered a much lower eccentricity polar orbit ( $e \approx 0.06$ ) with a life of about 6½ years and numerous orbits are being determined at the University of Aston; here the orbit is analysed only in the final 16 days of its orbital life, 1978 September 12-27. The North American Air Defense Command (NORAD) kindly supplied about 2500 observations for this 16-day period.

The RAE orbit refinement program<sup>1</sup>, in the PROP 6 version (which has a limit of 100 observations per orbit), has been used to determine 16 daily orbits with about 1500 of the NORAD observations, and an additional 70 from the US Navy Navspasur system.

The upper-atmosphere rotation rate has been determined from the decrease in orbital inclination to give the zonal wind speed near 240 km height, and the density scale height has been evaluated over the height band 200-270 km from the decrease in perigee height.

## 2 THE ORBIT

About 1360 observations were used, with a further 200 or so rejected, in the final determination of the 16 daily orbits for 1978 September 12-27, the last orbit being at an epoch 13 hours before decay. The orbital elements derived are listed in Table 1, with standard deviations.

The sd in inclination,  $i$ , varies between  $0.0003$  and  $0.0006^\circ$ , and in eccentricity,  $e$ , from  $4 \times 10^{-6}$  to  $7 \times 10^{-6}$ . These are equivalent to between 30 and 70 m in position for both  $i$  and  $e$ . This is an excellent accuracy for an orbit so near decay. The sd in right ascension of the node,  $\Omega$ , is between  $0.0002$  and  $0.0008^\circ$ ; for argument of perigee,  $\omega$ , or mean anomaly at epoch,  $M_0$ , the sd varies from  $0.03$  to  $0.19^\circ$ . Figs 1 to 5 show the variations of  $i$ ,  $a$ ,  $e$ ,  $\Omega$  and  $\omega$ . Although the overall drop in inclination over the 16 days was only  $0.02^\circ$  (see Fig 1), it was sufficient to determine accurately the upper-atmosphere wind (section 5).

## 3 VARIATION IN PERIGEE HEIGHT

Perigee height is required for determining density scale height and the heights applicable to density scale height and atmospheric rotation. Values of semi major axis,  $a$ , and eccentricity,  $e$ , from Table 1 are used to calculate perigee height over a spherical Earth,  $h_p$ , where

$$h_p = a(1 - e) - R, \quad (1)$$

where  $R$ , the Earth's mean equatorial radius, is taken to be 6378.14 km. Next, the perturbation  $\Delta e$ , in  $e$ , due to zonal harmonic and lunisolar effects, is obtained using the PROD computer program<sup>2</sup> and cleared from  $h_p$  to give  $Q$  where

$$Q = h_p + a\Delta e, \quad (2)$$

Table 1  
Orbital parameters for Heos 2 second-stage rocket, with standard deviations

Orbit	Date 1978 Sep	MJD	a	e	i	$\Omega$	$\omega$	$M_0$	$M_1$	$M_2$	$M_3$	$M_4$	$\epsilon$	D	N
1	12	43763.0	6659.3724 11	0.007233 6	89.7371 5	18.6781 5	9.66 3	90.66 3	5753.312 1	2.058 6			0.39	0.9	95
2	13	64.0	6656.0261 10	0.006900 4	89.7366 4	18.6376 4	5.62 3	85.85 3	5757.654 1	2.228 4			0.32	0.9	87
3	14	65.0	6652.6892 9	0.006591 5	89.7360 5	18.5984 4	1.42 3	85.61 3	5761.989 1	2.091 5			0.32	0.9	81
4	15	66.0	6649.4293 10	0.006280 5	89.7348 5	18.5580 4	357.25 4	89.59 4	5766.228 1	2.070 5			0.33	0.9	75
5	16	67.0	6646.1362 8	0.005979 5	89.7341 4	18.5179 4	352.96 3	97.92 3	5770.517 1	2.224 4			0.28	0.9	72
6	17	68.0	6642.5396 37	0.005662 4	89.7337 4	18.4784 4	348.06 4	111.34 4	5775.206 5	2.408 11			0.21	0.6	76
7	18	69.0	6638.7040 13	0.005378 7	89.7338 5	18.4373 4	343.50 4	129.25 4	5780.215 2	2.556 4			0.32	1.0	91
8	19	70.0	6634.7378 8	0.005098 4	89.7325 3	18.3974 3	338.57 3	152.64 3	5785.401 1	2.643 3			0.29	0.9	92
9	20	71.0	6630.4886 11	0.004780 5	89.7332 4	18.3562 2	333.63 4	181.37 3	5790.966 1	2.876 5			0.29	0.9	80
10	21	72.0	6625.8551 23	0.004516 5	89.7319 4	18.3144 4	328.52 5	216.07 5	5797.045 3	3.285 5	0.27 2		0.29	0.9	90
11	22	73.0	6620.2125 9	0.004197 6	89.7289 5	18.2716 6	323.53 7	257.33 7	5804.462 1	3.934 5			0.36	0.9	83
12	23	74.0	6613.4266 14	0.003885 5	89.7276 5	18.2317 5	317.63 6	307.52 6	5813.402 2	5.058 8			0.30	0.8	71
13	24	75.0	6604.9725 26	0.003537 5	89.7261 4	18.1873 4	311.83 9	7.70 9	5824.573 3	6.129 5	0.20 2		0.38	0.9	96
14	25	76.0	6594.8032 20	0.003114 4	89.7245 4	18.1460 3	306.15 11	79.99 11	5838.057 3	7.763 7	0.75 2		0.35	0.9	92
15	26	77.0	6579.6208 40	0.002604 7	89.7227 6	18.1006 8	300.54 16	168.44 16	5858.286 5	12.24 2	1.73 3	2.3 1	0.43	0.9	69
16	27	78.0	6551.8756 67	0.001815 6	89.7175 5	18.0595 4	292.69 19	285.98 19	5895.556 9	30.25 3	16.8 2	16.5* 2	0.48	0.9	94

\*  $M_5 = 11.2 \pm 0.8$

Key: MJD Modified Julian Day  
a semi major axis (km)  
e eccentricity  
i inclination (deg)  
 $\Omega$  right ascension of node (deg)  
 $\omega$  argument of perigee (deg)  
 $M_0$  mean anomaly at epoch (deg)  
 $M_1$  mean motion, n (deg/day)  
 $M_2$ - $M_5$  additional coefficients in polynomial for M  
 $\epsilon$  measure of fit  
D time coverage of observations (days)  
N number of observations used

which should be free of gravitational perturbations. These values of  $Q$  are plotted in Fig 6, and its decrease  $\Delta Q$  is used in the determination of scale height (section 4).

Perigee height,  $y_p$ , over an oblate Earth is obtained by allowing for (a) the variation in local Earth radius at perigee latitude and (b) the dynamical effects of the Earth's oblateness<sup>3</sup>, to give

$$y_p = h_p + 21.38 \sin^2 \omega + (1.67 - 3.32 \sin^2 \omega) ,$$

for 1972-05B at inclination  $90^\circ$ . Hence

$$y_p = h_p + 18.06 \sin^2 \omega + 1.67 \quad \text{km} \quad (3)$$

and is also plotted in Fig 6.

#### 4 DENSITY SCALE HEIGHT

The decrease in  $Q$ , shown in Fig 6 to be very well defined, depends on the density scale height  $H$ , and analysis of the variation in  $Q$  should therefore yield reliable values of  $H$ .

Low-eccentricity theory is needed because  $z$  ( $= ae/H$ ) is less than 3, and, for an orbit in an oblate atmosphere, we have<sup>4</sup>

$$\frac{da}{dx} = y_0 + \frac{1}{2}e(4 - 3y_0^2 - y_0y_2) - \frac{1}{2}c \cos 2\omega(y_0 - 2y_2 + y_0y_3) , \quad (4)$$

where  $x = ae$ ,  $y_r = I_r/I_1$ ,  $I_r$  being the Bessel function of the first kind and imaginary argument, of order  $r$  and argument  $z$ ; and  $c = \epsilon'a(1-e) \sin^2 i/2H$ , with  $\epsilon'$ , the ellipticity of the atmosphere, taken equal to the Earth's ellipticity, 0.00335.

In previous orbital analyses using low-eccentricity theory, the perigee height has been below 200 km and equation (4) has been used as it stands. Here, however, values of  $H$  are being determined at heights up to about 260 km, where the day-to-night variation in air density is large enough to affect equation (4) appreciably. In an atmosphere with maximum daytime density  $\rho_{\max}$  (at 14 h local time) and minimum night-time density  $\rho_{\min}$  (at 02 h local time), the equation for  $da/dx$  for a low-eccentricity orbit is<sup>5</sup>

$$\frac{da}{dx} = \frac{y_0 + F \cos \phi_p}{1 + \frac{1}{2}F \cos \phi_p (y_0 + y_2)} (1 + O(e)) , \quad (5)$$

where  $\phi_p$  is the angular distance of perigee from the centre of the diurnal bulge,  $F = (f-1)/(f+1)$  and  $f = \rho_{\max}/\rho_{\min}$ . In the conditions experienced by 1972-05B (exospheric temperature at night  $\approx 900$  K), the *COSPAR International Reference Atmosphere 1972* (Ref 6) gives  $f = 1.3$  at 200 km increasing to  $f = 1.7$  at 260 km height, so that  $0.15 < F < 0.25$ . Also  $z \approx 1$ , so that  $\frac{1}{2}(y_0 + y_2) \approx 1$ . Assuming that equation (5) can be expanded in powers of  $F$ , we find

$$\frac{da}{dx} = y_0 - \frac{1}{2}F \cos \phi_p (y_0^2 + y_0 y_2 - 2) + O(F^2, e) \quad (6)$$

If we assume that the effects of  $c$  and  $F$  are simply additive, the equation for  $da/dx$  in an oblate atmosphere with day-to-night variation in density may, from (4) and (6), be written

$$\begin{aligned} \frac{da}{dx} = y_0 + \frac{1}{2}e(4 - 3y_0^2 - y_0 y_2) - \frac{1}{2}c \cos 2\omega(y_0 - 2y_2 + y_0 y_3) \\ - \frac{1}{2}F \cos \phi_p (y_0^2 + y_0 y_2 - 2) + O(c^2, F^2, e^2, \text{etc}) \quad (7) \end{aligned}$$

For 1972-05B the local time at perigee is near 02 h, diametrically opposite the centre of the diurnal bulge when  $\omega = 0$ , so that, by a convenient chance,  $\phi_p = |180^\circ - \omega|$ . Thus, for 1972-05B in its final days, equation (7) becomes

$$\begin{aligned} \frac{da}{dx} = y_0 + \frac{1}{2}e(4 - 3y_0^2 - y_0 y_2) - \frac{1}{2}c \cos 2\omega(y_0 - 2y_2 + y_0 y_3) \\ + \frac{1}{2}F \cos \omega (y_0^2 + y_0 y_2 - 2) = \beta, \text{ say} \quad (8) \end{aligned}$$

This theoretical value of  $da/dx$  must be matched with the observational value which, following Ref 7, is given by

$$\frac{da}{dx} = 1 / \left( 1 + \frac{3\Delta Q}{4a} \cdot \frac{M_{1A} + M_{1B}}{M_{1B} - M_{1A}} \right) = \alpha, \text{ say} \quad (9)$$

where  $\Delta Q$  is the change in  $Q$  over a time interval  $\Delta t$ , selected to ensure that  $|\Delta Q| \geq 3$  km, and  $M_{1A}$  and  $M_{1B}$  are the initial and final values of the mean motion  $M_1$  during the time interval  $\Delta t$ . In (8) and (9) the values of  $a$ ,  $e$ ,  $\cos 2\omega$  and  $\cos \omega$  are taken as the mean during this time interval.

The density scale height  $H$  can now be determined by guessing two values of  $H$ , say  $H_1$  and  $H_2$ , which give  $\beta_1$  and  $\beta_2$  from equation (8), and then by interpolation using

$$H = \frac{H_1(\beta_2 - \alpha) + H_2(\alpha - \beta_1)}{\beta_2 - \beta_1} \quad (10)$$

These observational values of  $H$  are plotted in Fig 7 as circles with vertical rms error bars (calculated to be near 1 km for all values of  $H$ ) and horizontal bars representing the time intervals over which the values of  $H$  are calculated.

Values of  $H$  have also been calculated using the CIRA 1972 reference atmosphere<sup>6</sup>, for the appropriate heights and exospheric temperatures,  $T_\infty$ , and are plotted in Fig 7 as crosses. The values of  $T_\infty$  are calculated using appropriate values of solar 10.7 cm



radiation energy (plotted in Fig 7) and geomagnetic index, with allowance for semi-annual effects<sup>8</sup>. Care must be taken here to ensure that local time used to calculate  $T_{\infty}$  is not that at perigee but averaged over a wide arc of the orbit, as discussed in Ref 7. Using local time at perigee would result in errors of up to 100 K in  $T_{\infty}$ . Also plotted in Fig 7 is the height  $y$  ( $= y_p + \gamma H_p$ ) at which  $H$  applies. Values of  $\gamma$ , obtained from Ref 9, vary between 0.4 and 0.8.

Of the eleven observational values of  $H$  calculated, with a 2 per cent error, seven are within 6 per cent of the *CIRA 1972* values, the rms value being 4 per cent higher than *CIRA*. This agrees with a previously<sup>10</sup> determined value of 7 per cent higher, also for high solar activity. The four remaining values are 10-15 per cent higher than *CIRA* and Fig 7 shows this occurs during a period of high solar radiation energy. A medium magnetic storm occurred during the final 3 days in orbit (1978 September 25-27), where  $A_p$  increased from 36 to 51, but this does not seem to have made any significant difference.

The observational values of density scale height,  $H$ , plotted in Fig 7 are replotted in Fig 8 against height  $y$ , as crosses against a background of density scale height curves, for various  $T_{\infty}$ , obtained from *CIRA 1972*. Adjacent to each cross is the value of  $T_{\infty}$ , in brackets, which was used to calculate the *CIRA* value of  $H$ .

Both atmospheric oblateness  $\epsilon'$  and the day-to-night variation in density, measured by  $F$ , strongly influence the values of  $H$  obtained. If  $c$  is taken as zero in equation (8), *ie* for zero  $\epsilon'$ , the value of  $H$  obtained is reduced by 20 per cent in the first few days, is unchanged on September 23-24 (when  $\cos 2\omega = 0$ ), and is increased by 20 per cent on the day of decay. If  $F$  is taken as zero in equation (8),  $H$  increases by 30 per cent except for the final 4 days (to decay) when the increase is about 20 per cent. There is no reason to suspect that the values of  $\epsilon'$  or  $F$  used are appreciably in error: but if they were in error by 10 per cent, say, errors of up to 2½ per cent in  $H$  would occur.

## 5 UPPER-ATMOSPHERE ZONAL WIND

### 5.1 Results

The 16 daily values of inclination determined are listed in Table 1 and plotted in Fig 1. The values were then cleared of lunisolar and geopotential perturbations using the PROD program<sup>2</sup>, with numerical integration at 1-day intervals. The modified values are plotted in Fig 9, with standard deviations. The overall decrease in inclination is about  $0.023^{\circ}$ , about 50 times the mean sd of the values.

Since the orbital inclination is near  $90^{\circ}$ , the meridional wind has no effect and so the variation in inclination is due to the zonal wind.

The theoretical change in inclination has been calculated for several values of atmospheric rotation rate (expressed as  $\Lambda$  times the Earth's rotation rate) using the RAE computer program ROTATM (for an oblate atmosphere), with daily integration steps. The best fit to the modified inclination values is given by  $\Lambda = 1.40 \pm 0.05$  rev/day and is shown in Fig 9, where most of the points are within 2 sd of the curve. This corresponds to a zonal west-to-east wind of  $160 \pm 20$  m/s. Since zonal wind speed depends on local

time, this is plotted along the top of Fig 9; it shows little variation, always being between 01 and 02 h. The orbits are determined for 1978 September, i.e. equinox and high solar activity conditions; and the latitude of perigee steadily changes from  $10^{\circ}\text{N}$  to  $67^{\circ}\text{S}$ . So, although the effect of aerodynamic forces is spread over a wide arc around perigee, there is a bias towards equatorial rather than polar regions.

The wind speed is also an average over a range of heights, but the mean height  $y_{\Lambda}$  is given by

$$y_{\Lambda} = y_p + \lambda H_p$$

where  $\lambda = 0.5$ , from Ref 11. The mean perigee height  $y_p$  is estimated to be near 220 km from Fig 6, while the mean perigee scale height  $H_p$  is estimated to be near 40 km. Hence  $y_{\Lambda} \approx 240$  km.

## 5.2 Discussion

The analysis of numerous orbits by King-Hele and Walker<sup>12</sup> has indicated that zonal winds at heights of 200-400 km are usually from west to east at local times from 18 to 24 h, and east to west from 04 to 12 h local time. So the west-to-east wind of 160 m/s at 240 km height at 01-02 h local time obtained here is unexpectedly strong. However, it is not so anomalous as appears at first sight, for three reasons.

(1) The zonal winds are more strongly from west to east near the equator than at higher latitudes (see Fig 14 of Ref 12), and the perigee of 1972-05B travelled from latitude  $10^{\circ}\text{N}$  to latitude near  $60^{\circ}\text{S}$ , so that the effective average latitude was probably about  $20^{\circ}$ .

(2) The variation of wind speed with local time depends on latitude, and the maximum west-to-east wind probably occurs at a later local time near the equator. For example, the airglow measurements of Sipler and Biondi<sup>13</sup>, for 20 August 1977 at latitude  $9^{\circ}\text{N}$ , indicate a west-to-east wind exceeding 100 m/s between 21 and 02 h, with a value of 125 m/s at 01 h.

(3) The variation with local time may also depend on solar activity, with the maximum west-to-east wind being somewhat later when solar activity is high. For example, in 1970, when solar activity was high, Roble, Salah and Emery<sup>14</sup> found that, at 240 km height at latitude  $42^{\circ}\text{N}$ , zonal winds were west to east with velocity exceeding 100 m/s from 18 to 01 h on 23-24 February and from 19 to 01 h on 18-19 May. But in 1974, when solar activity was low, Hernandez and Roble<sup>15</sup> found west-to-east winds exceeding 100 m/s from 19 to 23 h at 240 km on 15 August.

The high value of  $\Lambda$  obtained here,  $1.40 \pm 0.05$  rev/day, is paralleled by results from three other satellites, 1969-20B, 1969-94B and 1970-43B, which all give  $\Lambda \approx 1.3 \pm 0.1$  for heights near 240 km at times of high solar activity, for local times of 18-03 h (see Ref 12).

The zonal wind speed in the thermosphere probably varies with height, season, latitude, solar activity, local time and day of the year, as well as being correlated with geomagnetic disturbances and events in the lower atmosphere. The value from 1972-05B, being so well defined, should help in unravelling this complex pattern.

6 CONCLUSIONS

The near-polar orbit of Heos 2 second-stage rocket, 1972-05B, has been determined daily for the last 16 days in orbit, including an orbit for the day of decay 1978 September 27. About 1360 observations were used, most of which were kindly supplied by the North American Air Defense Command, NORAD.

For the 16 orbits obtained, given in Table 1, the sd in inclination varies between 0.0003 and 0.0006°, while the sd in eccentricity varies from  $4 \times 10^{-6}$  to  $7 \times 10^{-6}$ ; these variations are both equivalent to 30-70 m in position.

Values of density scale height have been determined from the decrease in perigee height and compared with the *CIRA 1972* reference atmosphere; the values obtained are probably accurate to about 2 per cent and are about 4 per cent above *CIRA*, on average.

A value for the atmospheric rotation rate  $\Lambda$  was determined from the decrease in orbital inclination and found to be  $\Lambda = 1.40 \pm 0.05$  rev/day, equivalent to a strong west-to-east zonal wind of  $160 \pm 20$  m/s at a mean height of about 240 km. The conditions prevailing were: local time 01-02 h; equinox; high solar activity; and latitude of perigee between 10°N and 67°S.

## REFERENCES

- | <u>No.</u> | <u>Author</u>                                | <u>Title, etc</u>   |
|------------|--|---|
| 1          | R.H. Gooding                                 | The evolution of the PROP 6 orbit determination program, and related topics.<br>RAE Technical Report 74164 (1974)   |
| 2          | G.E. Cook                                    | Basic theory for PROD, a program for computing the development of satellite orbits.<br><i>Celestial Mechanics</i> , <u>7</u> , 301-314 (1972)<br>RAE Technical Report 71007 (1971)                                    |
| 3          | Y. Kozai                                     | The motion of a close Earth satellite.<br><i>Astronom. J.</i> , <u>64</u> , 367-377 (1959)  |
| 4          | D.G. King-Hele<br>G.E. Cook<br>D.M.C. Walker | The contraction of satellite orbits under the influence of air drag. Part 2: with oblate atmosphere.<br><i>Proc. Roy. Soc. A</i> , <u>264</u> , 88-121 (1961)<br>RAE Technical Note GW 565 (1960)                     |
| 5          | G.E. Cook<br>D.G. King-Hele                  | The contraction of satellite orbits under the influence of air drag. Part 5: with day-to-night variation in air density.<br><i>Phil. Trans. A</i> , <u>259</u> , 33-67 (1965)<br>RAE Technical Report 64029 (1964)    |
| 6          | -  | <i>CIRA 1972</i> (Cospar International Reference Atmosphere 1972).<br>Akademie-Verlag, Berlin (1972)  |
| 7          | H. Hiller                                    | Determination and geophysical interpretation of the orbit of China 2 rocket (1971-18B).<br><i>Planet. Space Sci.</i> , <u>27</u> , 1247-1267 (1979)<br>RAE Technical Report 78107 (1978)                              |
| 8          | D.M.C. Walker                                | Variations in air density from January 1972 to April 1975 at heights near 200 km.<br><i>Planet. Space Sci.</i> , <u>26</u> , 291-309 (1978)<br>RAE Technical Report 77078 (1977)                                      |
| 9          | D.G. King-Hele<br>G.E. Cook                  | The contraction of satellite orbits under the influence of air drag. Part 4: with scale height dependent on altitude.<br><i>Proc. Roy. Soc. A</i> , <u>275</u> , 357-390 (1963)<br>RAE Technical Note Space 18 (1962) |
| 10         | H. Hiller                                    | Upper-atmosphere rotation rate determined from the orbit of China 6 rocket (1976-87B).<br><i>Planet. Space Sci.</i> , <u>28</u> , 549-558 (1980)<br>RAE Technical Report 79115 (1979)                                 |

## REFERENCES (concluded)

- | <u>No.</u> | <u>Author</u>                          | <u>Title, etc</u>  |
|------------|--|--|
| 11         | D.G. King-Hele<br>D.W. Scott           | The effect of atmospheric rotation on a satellite orbit,<br>when scale height varies with height.<br><i>Planet. Space Sci.</i> , <u>17</u> , 217-232 (1969)<br>RAE Technical Report 68066 (1968) |
| 12         | D.G. King-Hele<br>D.M.C. Walker        | Upper-atmosphere zonal winds: variation with height and<br>local time.<br><i>Planet. Space Sci.</i> , <u>25</u> , 313-336 (1977)<br>RAE Technical Report 76055 (1976)                            |
| 13         | D.P. Sipler<br>M.A. Biondi             | Equatorial F-region neutral winds from nightglow OI<br>630.0 nm doppler shifts.<br><i>Geophys. Res. Letters</i> , <u>5</u> , 373-376 (1978)  |
| 14         | R.G. Roble<br>J.E. Salah<br>B.A. Emery | The seasonal variation of the diurnal thermospheric winds<br>over Millstone Hill during solar cycle maximum.<br><i>Journ. Atmos. Terr. Phys.</i> , <u>39</u> , 503-511 (1977)                    |
| 15         | G. Hernandez<br>R.G. Roble             | Direct measurements of nighttime thermospheric winds and<br>temperatures.<br><i>Journ. Geophys. Res.</i> , <u>81</u> , 2065-2074 (1976)  |

Fig 1




Fig 1 Observational values of inclination, with sd

Fig 2




Fig 2 Observational values of semi major axis, a

Fig 3




Fig 3 Observational values of eccentricity,  $e$



Fig 4




Fig 4 Observational values of right ascension of the node,  $\Omega$

Fig 5




Fig 5 Observational values of argument of perigee,  $\omega$

Fig 6




Fig 6 Perigee heights:  $y_p$  over an oblate Earth; and  $Q$  over a spherical Earth, cleared of gravitational perturbations

TR 60119

Fig 7




Fig 7 Density scale height H , its height of application y , and solar radiation energy S<sub>10.7</sub>

Fig 8




Fig 8 Values of density scale height, with CIRA 1972 curves

Fig 9




Fig 9 Inclination values, cleared of perturbations, with fitted  $\Lambda$ -curve

REPORT DOCUMENTATION PAGE

Overall security classification of this page

UNLIMITED

As far as possible this page should contain only unclassified information. If it is necessary to enter classified information, the box above must be marked to indicate the classification, e.g. Restricted, Confidential or Secret.

1. DRIC Reference (to be added by DRIC)	2. Originator's Reference RAE TR 80119	3. Agency Reference N/A	4. Report Security Classification/Marking <del>UNCLASSIFIED</del> <b>UNLIMITED</b>
5. DRIC Code for Originator 7673000W	6. Originator (Corporate Author) Name and Location Royal Aircraft Establishment, Farnborough, Hants, UK		
5a. Sponsoring Agency's Code N/A	6a. Sponsoring Agency (Contract Authority) Name and Location N/A		
7. Title <b>The orbit of 1972-05B in its final phase, with geophysical inferences</b>			
7a. (For Translations) Title in Foreign Language			
7b. (For Conference Papers) Title, Place and Date of Conference			
8. Author 1. Surname, Initials Hiller, H.	9a. Author 2	9b. Authors 3, 4 ....	10. Date    Pages    Refs. September 1980    20    15
11. Contract Number N/A	12. Period N/A	13. Project	14. Other Reference Nos. Space 587
15. Distribution statement (a) Controlled by —    Head of Space Department, RAE (b) Special limitations (if any) —			
16. Descriptors (Keywords)                    (Descriptors marked * are selected from TEST) Upper atmosphere winds. Upper atmosphere scale height. HEOS 2. Orbit determination.			
17. Abstract    The polar orbit of Heos 2 second-stage rocket, 1972-05B, has been determined on each of the final 16 days before its decay in September 1978, using the RAE orbit refinement program, PROP 6, with about 1360 observations. An accuracy of 30-70 m, both radial and across track, was achieved.  Eleven values of density scale height have been determined from the decrease in perigee height, with a 2 per cent error; seven of these values are within 5 per cent of the CIRA 1972 reference-atmosphere values, the rms value being 4 per cent higher than CIRA.  The rotation rate of the upper atmosphere, $\dot{\alpha}$ , was determined from the decrease in orbital inclination as $\dot{\alpha} = 1.40 \pm 0.05$ rev/day; to a strong west-east zonal wind of $160 \pm 20$ m/s, at a mean height of about 240 km. The local time was 01-02 h; solar activity was high; and the latitude of perigee varied from $10^{\circ}N$ to $67^{\circ}S$ .			

DATE  
FILMED  
— 8