

AD-A099 296

MATERIALS RESEARCH LABS ASCOT VALE (AUSTRALIA)
MYTILUS EDULIS AS AN INDICATOR OF TRACE METAL POLLUTION IN NAVA--ETC(U)
JUL 80 I C DUNSTAN, A DE FOREST, R W PETTIS

F/G 13/2

UNCLASSIFIED NRL-R-781

NL

for
reference



END
DATE
FILMED
8-84
DTIC

LEVEL

12



AD A 099 296

**DEPARTMENT OF DEFENCE
DEFENCE SCIENCE AND TECHNOLOGY ORGANISATION
MATERIALS RESEARCH LABORATORIES**

MELBOURNE, VICTORIA


REPORT

MRL-R-781

MYTILUS EDULIS AS AN INDICATOR OF TRACE METAL POLLUTION
IN NAVAL DOCKYARD WATERS WITH PRELIMINARY RESULTS
FROM WILLIAMSTOWN NAVAL DOCKYARD, VICTORIA

Ian C. Dunstan, Albert de Forest and Russell W. Pettis*

Approved for Public Release



E

* Present address: Monsanto Australia Ltd.,
West Footscray, Victoria.

DTIC FILE COPY



© COMMONWEALTH OF AUSTRALIA 1980

JULY, 1980

81 5 26 011

THE UNITED STATES NATIONAL
TECHNICAL INFORMATION SERVICE
IS AUTHORIZED TO
REPRODUCE AND SELL THIS REPORT

DEPARTMENT OF DEFENCE
MATERIALS RESEARCH LABORATORIES

REPORT

(12) 2/88

(17)

MRL-R-781

1 = 01 88

MYTILUS EDULIS AS AN INDICATOR OF TRACE METAL POLLUTION
IN NAVAL DOCKYARD WATERS WITH PRELIMINARY RESULTS
FROM WILLIAMSTOWN NAVAL DOCKYARD, VICTORIA,

(11) Ian C./Dunstan, Albert/de Forest and Russell W./Pettis*

ABSTRACT

↓

Literature on the accumulation of trace metals by marine organisms suggest *Mytilus edulis* as a potential indicator of trace metal levels. A technique for trace metal determination in mussel tissue has been developed and used for the determination of Pb, Zn, Cu, Cd, Ni, Mn, Ag, Fe and Co in whole tissues and individual organs of mussels from Williamstown Naval Dockyard. From these results, the suitability of *M. edulis* as an indicator of trace metal pollution in dockyard waters has been assessed.

Approved for Public Release

Section For	
Dist	
Special	

* Present address: Monsanto Australia Ltd.,
West Footscray, Victoria.

Availability Codes	
Dist	Special
A	

© COMMONWEALTH OF AUSTRALIA 1980

POSTAL ADDRESS: Chief Superintendent, Materials Research Laboratories
P.O. Box 50, Ascot Vale, Victoria 3032, Australia

409027
alt

DOCUMENT CONTROL DATA SHEET

Security classification of this page:

UNCLASSIFIED

1. DOCUMENT NUMBERS:		2. SECURITY CLASSIFICATION:	
a. AR Number:	AR-002-166	a. Complete document:	UNCLASSIFIED
b. Series & Number:	REPORT MRL-R-781	b. Title in isolation:	UNCLASSIFIED
c. Report Number:	MRL-R-781	c. Abstract in isolation:	UNCLASSIFIED
3. TITLE: MYTILUS EDULIS AS AN INDICATOR OF TRACE METAL POLLUTION IN NAVAL DOCKYARD WATERS WITH PRELIMINARY RESULTS FROM WILLIAMSTOWN NAVAL DOCKYARD, VICTORIA			
4. PERSONAL AUTHOR(S):		5. DOCUMENT DATE:	
DUNSTAN, Ian C., de FOREST, Albert PETTIS, Russell W.		JULY, 1980	
6. TYPE OF REPORT & PERIOD COVERED:			
7. CORPORATE AUTHOR(S):		8. REFERENCE NUMBERS:	
Materials Research Laboratories		a. Task: DST 77/006	
		b. Sponsoring Agency:	
		9. COST CODE: 284410	
10. IMPRINT (Publishing establishment)		11. COMPUTER PROGRAMME(S):	
Materials Research Laboratories, P.O. Box 50, Ascot Vale, Vic.3032		(Title(s) and language(s)):	
JULY, 1980			
12. RELEASE LIMITATIONS (of the document):			
Approved for Public Release			
12-0. OVERSEAS: <input type="checkbox"/> N.O. <input type="checkbox"/> P.R. <input checked="" type="checkbox"/> 1 <input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C <input type="checkbox"/> D <input type="checkbox"/> E <input type="checkbox"/>			
13. ANNOUNCEMENT LIMITATIONS (of the information on this page):			
No limitation			
14. DESCRIPTORS:			
	630	Water pollution	Mussels
	636	Heavy metals	
15. COSATI CODES: 0603 1302			
16. ABSTRACT (if this is security classified, the announcement of this report will be similarly classified):			
Literature on the accumulation of trace metals by marine organisms suggest <i>Mytilus edulis</i> as a potential indicator of trace metal levels. A technique for trace metal determination in mussel tissue has been developed and used for the determination of Pb, Zn, Cu, Cd, Ni, Mn, Ag, Fe and Co in whole tissues and individual organs of mussels from Williamstown Naval Dockyard. From these results, the suitability of <i>M. edulis</i> as an indicator of trace metal pollution in dockyard waters has been assessed.			

C O N T E N T S

	<u>Page No.</u>
1. INTRODUCTION	1
2. METHOD	3
2.1 <i>Sample Collection</i>	5
2.2 <i>Sample Analysis</i>	5
2.3 <i>Data Analysis</i>	5
3. RESULTS AND DISCUSSION	5
3.1 <i>Trace Metal Uptake by Different Organs</i>	5
3.2 <i>Comparison with other Areas</i>	6
3.3 <i>Mytilus edulis as an Indicator Organism in Dockyard Waters</i>	6
4. CONCLUSIONS	7
5. REFERENCES	7

* * *

MYTILUS EDULIS AS AN INDICATOR OF TRACE METAL POLLUTION
IN NAVAL DOCKYARD WATERS WITH PRELIMINARY RESULTS
FROM WILLIAMSTOWN NAVAL DOCKYARD, VICTORIA

1. INTRODUCTION

Contamination of the marine environment by trace metals can produce deleterious effects on upper levels of marine food chains, and subsequently man [1,2,3]. Assessment of trace metal loads in regions of effluent discharge ensures that contamination by trace metals can be detected before dangerous levels are reached.

The possibility of heavy metal input from dockyard activities has led authorities at Williamstown Naval Dockyard (WND) Victoria to request a study of trace metal levels in waters adjacent to the dockyard in Hobson's Bay.

Trace metal concentrations in the marine environment can be measured in water [e.g. 4,5], sediments [e.g. 4,6] or organisms [e.g. 7,8,9,10]. Measurements performed on marine organisms have the following advantages over the alternative sources :

- (i) trace metal levels in organisms are higher and hence less subject to analytical error than those from sediments or water samples [5,7],
- (ii) levels in organisms are buffered against short term fluctuations in ambient levels of trace metals and therefore reflect past environmental conditions more accurately than those gained from water samples [11], and
- (iii) measurements from organisms provide data on the trace metal load in the marine food system not provided by measurements of water or sediment trace metal content [12].

Numerous marine organisms have been used to assess trace metal pollution. These include various algae [13,14], crustacea [12,15,16], ascidians [17,18], polychaetes [19], bivalves [20,21,22,23] and fish [24,25].

Barbaro [12] lists the following criteria required for an organism to be an effective pollution indicator :

- (1) sessile living habits,
- (2) availability in all seasons,
- (3) ease of sampling,
- (4) ubiquitous occurrence and abundance in the areas to be monitored,
- (5) predisposition for a consistent uptake of one or more pollutants,
- (6) high capacity for accumulation above environmental levels, and
- (7) predisposition for retaining pollutants for a sufficient period after a reduction in environmental levels.

Mussels, particularly *Mytilus edulis*, have been used as pollution indicators in local [10,26,27] and overseas studies [28,29,30,31].

Mussels are sessile in habit and occur abundantly throughout the year in easily accessible locations at WND. They are also found in waters adjacent to Naval facilities in Sydney Harbort [32] and Cockburn Sound [33].

Experimental evidence [34] has shown the rate of Pb uptake by *Mytilus edulis* to be linearly dependent upon the concentration in the surrounding medium. Uptake of Cu, Pb, Co, Cd and Ni by *Mytilus edulis* has been quantified under experimental conditions [34]. The mussel has been considered to reflect ambient levels of Zn [10,26,29,34,36], Pb [1,10,26,29,34], Ni [35], Ag [29], Cu [1,29], Cd [1,10,26], Cr [29], Mn [10,36], Fe [10,36] and Co [36], although other writers express doubts on the ability of *Mytilus edulis* to monitor environmental levels of Ni and Cu which show atypical uptake behaviour in molluscs [26,29]. The Zn, Cd and Pb content of mussels have been shown to reflect industrial effluent discharge in Port Phillip Bay [26]. Retention of metals for varying periods of time after termination of supply has been demonstrated experimentally for Cd, Co, Ag [1] and Pb [1,34].

Mytilus edulis therefore satisfies the necessary criteria for indicator organisms and can be used to assess trace metal levels in dockyard environments. Trace metal contamination can be measured for the whole organism [10,26,30,31,37] or in specific organs [27,28,29]. Analysis of the various organs affords additional data on the uptake pathways of trace metals, although *in situ* information needs to be supplemented with experimental evidence before firm conclusions can be made. Experimental studies [34] have shown the kidney to be important in the Pb exchange process between

mussels and their environment. The exchange rate of Pb was also high for the stomach, as was metal uptake by the gills. Uptake of Zn, Mn, Co and Fe followed the order : stomach > gills > foot > mantle > gonad > adductor muscle [36]. Schulz-Baldes [34] proposed a general pattern of trace metal uptake in bivalves, with the kidney, gills and stomach having the highest, and the gonads, mantle and muscle the lowest rate of uptake.

Concentrations of several trace metals in New Zealand mussels were found to be highest in visceral mass and intestine, although the kidney was not isolated [28]. Elevated concentrations of lead were found in the stomach and gills of Port Phillip Bay mussels [27]. Again, no measurements were obtained from the kidney.

Food ingestion has been cited as a greater source of ^{65}Zn , ^{54}Mn , ^{58}Co and ^{59}Fe uptake than sorption from the water for mussels [36]. However, Pb accumulated in the stomach and kidney of mussels at the same rate in Pb-labelled medium experiments as it did for Pb-labelled food experiments [34]. Metals were taken up primarily across gill and adductor muscle membranes, passed into the blood stream and then transported to the kidney and other organs. Similar processes of ion adsorption at water-membrane interfaces and absorption across semi-permeable membranes were postulated by Romeril [22] as important in the experimental accumulation of Zn by oysters. Some doubt exists therefore regarding the pathway of trace metal accumulation in mussels. Measurements of the concentrations of metals in different organs of mussels in their natural habitat are required to enable the determination of metal uptake pathways.

This report investigates the use of the mussel *Mytilus edulis* as an indicator of trace metal concentration in dockyard waters, and presents preliminary results for a range of metals in whole mussels and various organs from mussels at WND.

2. METHOD

2.1 *Sample Collection*

Mussels were collected from the intertidal zone of pilings at the end of Dockyard Pier, at Williamstown Naval Dockyard, Hobson's Bay. The dockyard waters are estuarine with a mean salinity of 28‰. The water temperature varies seasonally from 10°C to 22°C [32].

Twenty mussels of between 3.5 and 5.0 cm in length were collected and placed in a bucket of ambient seawater for forty-eight hours to allow defecation of gut contents. They were then frozen pending analysis.

2.2 *Sample Analysis*

The lengths of all mussels were measured. The foot, mantle, gills, posterior adductor muscle, kidney, stomach (including digestive gland) and remaining soft parts were dissected from ten mussels. These organs and the whole soft parts of the other ten mussels were isolated, weighed and freeze-dried to constant weight.

Several methods of digestion [38,39] were tested and the results compared to the biological reference material NBS-SRM 1571 - Orchard Leaves (Table 1). The nitric acid/hydrogen peroxide mixture yielded the most accurate results. The method of Topping [38] was unsuitable as it led to the formation of unstable perchlorate salts.

TABLE 1

COMPARATIVE DIGESTION PROCEDURES FOR BIOLOGICAL TISSUE

Digestion Procedure	Metal Concentration ($\mu\text{g/g}$) dry wt			
	Cd	Pb	Cu	Zn
NBS Orchard Leaves	0.11 \pm 0.02	45 \pm 3	12 \pm 1	25 \pm 3
Nitric Acid	0.79	44.1	15.3	36.3
Nitric/Perchloric Acid	1.11	38.8	14.6	33.1
Nitric Hydrogen Peroxide	0.81	44.1	12.6	31.9

The nitric acid/hydrogen peroxide method subsequently adopted was as follows :

- (i) tissue samples were refluxed with 10 ml concentrated nitric acid and evaporated to near dryness,
- (ii) samples and a mixture of 10 ml water, 10 ml nitric acid and 10 ml hydrogen peroxide were refluxed and the step repeated until a colourless solution indicated oxidation was complete,
- (iii) samples were then evaporated to near dryness and the residue diluted to 25 ml with distilled water.

Fe and Zn concentrations were determined by flame mode analysis in a Varian Techtron Model AA-6 Atomic Absorption Spectrophotometer (AAS). The manufacturers recommendations were used for all instrument settings [40]. Cu, Cd, Pb, Ni, Co, Mn and Ag were analysed by flameless AAS using the Varian Techtron Model 63 carbon rod furnace. Samples were injected into the carbon furnace using a 5 μl teflon-tipped microsyringe. The furnace settings used for all metals are described in Table 2.

TABLE 2

FURNACE SETTINGS FOR ALL TRACE METAL ANALYSES

	Temp (°C)	Time (sec)
Dry	150	30
Ash	400	30
Atomise	2000	4

Analyses were calibrated with standard solutions in the working range of the analytes. Standard addition methods of calibration were not used as the matrix was destroyed prior to analysis of the nitric acid/hydrogen peroxide mixture.

2.3 Data Analysis

The kidney sample from one mussel was lost during preparation. Calculations which involved the kidney were therefore made from only nine samples.

Final trace metal concentrations were calculated/in parts per million (ppm) dry weight. The relationship between wet weight (w) and dry weight (D), calculated by linear regression, was $w = 9.01D - 0.03$ (correlation coefficient = 0.97).

As the statistical distribution of trace metal concentrations was not known, the Friedman's non-parametric two-way analysis of variance by ranks was used to determine the significance of differences between several groups of data. Where differences occurred, a critical range method was used to test the significance of differences' between all possible pairs in the analysis [41].

3. RESULTS AND DISCUSSION

3.1 Trace Metal Uptake by Different Organs

Trace metal concentrations varied significantly ($P < 0.01$) between organs (Table 3). For each metal, pairs of organs which differed significantly ($P < 0.05$) are shown in Table 4. The kidney was the primary site for metal concentration. 25% of Pb, 16% of Co, 15% of Zn and 13% of Ni found in the whole soft tissue were present in the kidney whilst the dry weight of the organ constituted only 2.7% of the total body weight. Similarly, Schulz-Baldes [34] found 50% to 70% of total Pb in the kidney whilst the organ formed only 6% to 8% of the dry weight of mussels sampled in that study.

Some metals, particularly Fe, Cu, Co, Pb and Ni are also strongly concentrated in the stomach.

The results suggest that the various metals may have different accumulation pathways. Higher concentrations of Co, Fe and Cu occurred in the stomach than in the gills and muscle, and these metals may have been taken up primarily during food ingestion. The other metals do not show higher concentrations in the stomach and may be primarily accumulated by sorption processes directly from the water. The high concentrations of metals in the kidney, for all metals, support observations [34] that the metals are transferred to the circulatory systems from either the stomach or gills and transported to the kidney where they are removed from the blood.

Further experimental and *in situ* evidence is required before the uptake pathways of each metal can be determined.

3.2 Comparison with other Areas

Trace metal concentrations from the present investigation are similar in magnitude to those from other studies (Table 5). Whenever possible results from Port Phillip Bay and Westernport Bay, Victoria, have been cited. However, the absence of data for some metals necessitated the inclusion of overseas results. The only data available on Co accumulation by mussels was for *Mytilus galloprovincialis* from a study in the Gulf of Spezia, Italy [30].

Uptake of trace metals by mussels can vary considerably with salinity and temperature of the water [11,43], size and reproductive condition of the mussels [42,44] and season and depth of sampling [11]. Several of these parameters would vary between studies and interstudy comparisons should be made with some reservations.

The concentrations of Zn, Cd, Cu and Pb fall below permissible levels recommended for shell fish by the National Health and Medical Research Council [45]. However, as our results are from a single sampling experiment, an assessment of trace element pollution at WND cannot be made at this time.

3.3 *Mytilus edulis* as an Indicator Organism in Dockyard Waters

Mussels have been used to assess trace metal levels in both Australia and overseas regions. This investigation has noted the availability of mussels in WND and exemplified the ability of *Mytilus edulis* to take up and concentrate trace metals far above environmental levels. As mussels can also be obtained from Sydney Harbour and Cockburn Sound, they provide an effective monitor of trace metal levels in waters adjacent to the major Australian Naval Facilities.

Where *Mytilus* is to be used as an indicator of trace metal pollution, sampling methods need to be planned to maximise the usefulness of results. Mussel length and depth of collection should be constant, and water temperature and salinity, sampling season and reproductive condition of the mussels recorded. This data would enable valid comparisons of trace metal levels from different studies to be made.

Comparisons of concentrations in organs that respond rapidly to environmental conditions, such as the kidney, to those that react more slowly, such as the foot, can provide additional information on past environmental levels of trace metal concentrations. In addition to the determination of

metals concentrations in whole mussels, future sampling programmes should include measurement of heavy metal concentrations in several organs.

4. CONCLUSIONS

1. A nitric acid/hydrogen peroxide mixture was a suitable digestion agent for heavy metal analyses of mussel tissue.
2. Trace metal concentrations varied significantly between organs of mussels collected from Williamstown Naval Dockyard.
3. Concentrations of Pb, Zn, Cu and Cd in mussels from WND were below those recommended for molluscs by the National Health and Medical Research Council.
4. *Mytilus edulis* satisfies criteria for indicator organisms of trace metals in Australian Naval Dockyard waters.

5. REFERENCES

1. Majori, L. and Petronio, F. (1973). Marine pollution by metals and their accumulation of biological indicators (accumulation factor). *Rev. Intern. Oceanogr. Med.*, 31/32, 55-89.
2. Thrower, S.J. and Eustace, I.J. (1973). Heavy metal accumulation in oysters grown in Tasmanian waters. *Ed. Technol. Aust.*, 25, (11), 546-547.
3. Waldichuk, M. (1974). Some biological concerns in heavy metal pollution. In *Pollution and Physiology of Marine Organisms*. Academic Press.
4. French, H.I. and Thistlewaite, P.J. (1976). Trace metal levels in the waters and sediments of Corio Bay. *Proc. Roy. Aust. Chem. Inst.*, 43, (3), 73-79.
5. de Forest, A., Pettis, R.W. and Fabris, G. (1978). Analytical acceptability of trace metal levels found in oceanic waters around Australia. *Aust. J. mar. Freshwat. Res.*, 29, 193-204.
6. Smith, J.D. (1976). Occurrence of heavy metals in the sediments of Mordialloc Creek. *Proc. Roy. Aust. Chem. Inst.*, 43, 305-306.
7. Bryan, G.W. (1971). The effects of heavy metals on marine and estuarine organisms. *Proc. Roy. Soc. Lond.*, 177, 389-410.
8. Stenner, R.D. and Nickless, G. (1975). Heavy metals in organisms of the Atlantic coast of S.W. Spain and Portugal. *Mar. Poll. Bull.*, 6, 89-92.

9. Bryan, G.W. and Hummerstone, L.G. (1977). Indicators of heavy metal contamination in the Looe Estuary, Cornwall, with particular reference to silver and lead. *J. mar. biol. Assoc. U.K.*, 57, (1), 75-92.
10. Harris, J.E., Fabris, G.J., Statham, P.J. and Tawfik, F. (1979). Biogeochemistry of selected heavy metals in Westernport Bay, Victoria, and the use of invertebrates as indicators with emphasis on *Mytilus edulis planulatus*. *Aust. J. mar. Freshwat. Res.*, 30, 159-178.
11. Phillips, D.J.H. (1976). The common mussel, *Mytilus edulis* as an indicator of pollution by Zn, Cd, Pb and Cu. I. Effects of environmental variables on uptake of metals. *Mar. Biol.*, 38, 59-69.
12. Barbaro, A., Francescon, A., Polo, B. and Bilio, M. (1978). *Balanus amphitrite* as a potential indicator of fluoride, copper, lead, chromium and mercury in north Atlantic lagoons. *Mar. Biol.*, 46, (3), 247-258.
13. Bryan, A.W. and Hummerstone, L.G. (1973). Brown seaweed as an indicator of heavy metals in estuaries in S.W. England. *J. mar. biol. Assoc. U.K.*, 53, 705-720.
14. Fuge, R. and James, K.H. (1974). Trace metal concentration in *Fucus* from the Bristol Channel. *Mar. Poll. Bull.*, 5, 9-12.
15. Bryan, A.W. (1968). Concentrations of zinc and copper in the tissues of decapod crustacea. *J. mar. biol. Assoc. U.K.*, 48, 303-321.
16. Walker, G., Rainbow, P.S., Foster, P. and Crisp, D.J. (1975). Barnacles: possible indicators of zinc pollution. *Mar. Biol.*, 30, 57-65.
17. Papadopoulou, C. and Kaniias, G. (1977). Tunicate species as marine pollution indicators. *Mar. Poll. Bull.*, 8, (10), 229-231.
18. Danskin, G.P. (1978). Accumulation of heavy metals by some solitary tunicates. *Can. J. Zool.*, 56, 547-555.
19. Reish, D.J., Piltz, F., Martin, N. and Word, J. (1976). The effect of heavy metals on laboratory populations of two species of polychaetes with comparisons of water quality conditions and standards in southern Californian marine waters. *Wat. Res.*, 10, 299-302.
20. Bryan, C.W. (1973). The occurrence and seasonal variation of trace metals in the scallop *Pecten maximus* (L) and *Chlamys opercularis* (L). *J. mar. biol. Assoc. U.K.*, 53, 145-166.
21. Hugget, R.J., Bender, M.E. and Stone, H.D. (1973). Utilizing metal concentration relationship in the eastern oyster (*Crassostrea virginica*) to detect heavy metal pollution. *Wat. Res.*, 7, 451-460.

22. Romeril, M.G. (1971). The uptake and distribution of Zn in oysters. *Mar. Biol.*, 9, 347-354.
23. Bryan, G.W. and Hummerstone, L.G. (1978). Heavy metals in the burrowing bivalve *Scrobicularia plana* from contaminated and uncontaminated estuaries. *J. mar. biol. Assoc., U.K.*, 58, 401-420.
24. Badsha, K.S. and Sainsbury, M. (1977). Uptake of zinc, cadmium and lead by young whiting in the Severn Estuary. *Mar. Poll. Bull.*, 8, (7), 164-166.
25. Bebbington, G.N., Machay, N.J., Chvojka, R., Williams, R.J., Dunn, A. and Auty, E.H. (1977). Heavy metals selenium and arsenic in nine species of Australian commercial fish. *Aust. J. mar. Freshwat. Res.*, 28, 277-286.
26. Phillips, D.J.H. (1976). The common mussel, *Mytilus edulis*, as an indicator of pollution by Zn, Cd, Pb and Cu. II. Relationship of metals in the mussel to these discharged by industry. *Mar. Biol.*, 38, 71-80.
27. Talbot, V., Magee, R.J. and Hussain, M. (1976). Lead in Port Phillip Bay mussels. *Mar. Poll. Bull.*, 7, 234-237.
28. Brooks, R.R. and Rumsby, M.G. (1965). The biogeochemistry of trace element uptake by some New Zealand bivalves. *Limnol. Oceanog.*, 10, 521-528.
29. Alexander, G.V. and Young, D.R. (1976). Trace metals in Southern Californian mussels. *Mar. Poll. Bull.*, 7, (1), 7-9.
30. Capelli, R., Contardi, V., Fassone, B. and Zanicchi, G. (1978). Heavy metals in mussels from the Gulf of Spezia and from the promontory of Portofino, Italy. *Mar. Chem.*, 6, 179-186.
31. Davenport, J. and Mandy, A. (1978). The detection of heightened seawater copper concentrations by the mussel *Mytilus edulis*. *J. mar. biol. Assoc. U.K.*, 58, 845-850.
32. Russ, G. (1977). A comparison of the marine fouling occurring at the two principal Australian naval dockyards. *Report MRL-R-688*. Materials Research Laboratories, Melbourne, Victoria.
33. Dunstan, I.C. (1978). Marine fouling at HMAS Stirling, WA, October 1976 - April 1978. *Report MRL-R-731*. Materials Research Laboratories, Melbourne, Australia.
34. Schulz-Baldes, M. (1974). Lead uptake from seawater and food, and lead loss in the common mussel, *Mytilus edulis*. *Mar. Biol.*, 25, 177-193.
35. Friedrich, A.R. and Filice, F.P. (1976). Uptake and accumulation of the nickel ion by *Mytilus edulis*. *Bull. environ. Contamin. Toxicol.*, 16, (6), 750-755.

36. Pentreath, R.J. (1973). The accumulation from water of ^{65}Zn , ^{54}Mn , ^{58}Co and ^{59}Fe by *Mytilus edulis*. *J. mar. biol. Assoc. U.K.*, 53, 127-143.
37. Fowler, S.W. and Oregioni, B. (1976). Trace metals in mussels from the N.W. Mediterranean. *Mar. Poll. Bull.*, 7, 26-29.
38. Florence, T.M. (1972). Determination of trace metals in marine samples by anodic stripping voltammetry. *J. Electroanal. Chem.*, 35, 237.
39. Topping, G. (1972). Analysis of reference materials used for heavy metal analyses of biological tissue. In *Baseline Studies of Pollutants in the Marine Environment - Heavy Metals, Halogenated Hydrocarbons and Petroleum*. IDOE, Brookhaven National Laboratory, USA.
40. Anonymous. (1975). Basic atomic absorption spectroscopy. *Varian Techtron Manual*, Springvale, Australia.
41. Colquhoun, D. (1971). *Lectures on Biostatistics*. Oxford: Clarendon Press.
42. Boyden, C.R. (1974). Trace element content and body size in molluscs. *Nature*, 253, 311-314.
43. Phillips, D.J.H. (1977). Effects of salinity on net uptake of zinc by the common mussel *Mytilus edulis*. *Mar. Biol.*, 41, (1), 79-88.
44. Simpson, R.D. (1979). Uptake and loss of zinc and lead by *Mytilus edulis* and relationships with body weight and reproductive cycle. *Mar. Poll. Bull.*, 10, 74-78.
45. Thompson, J.D. (1979). Heavy metals in the native oyster and mussel from Port Davey, Tasmania. *Aust. J. mar. Freshwat. Res.*, 30, (3), 421-424.

TABLE 3

TRACE METAL CONCENTRATION (mean \pm s.d., ppm dry weight)
IN *MYTILUS EDULIS*

	Whole Soft Tissue	Gills	Mantle	Foot	Kidney	Stomach	Poserior Adductor Muscle	Soft Remains
Fe	253 \pm 99	247 \pm 105	499 \pm 400	469 \pm 388	1453 \pm 529	950 \pm 278	208 \pm 88	290 \pm 147
Zn	279 \pm 102	560 \pm 115	470 \pm 981	193 \pm 75	1477 \pm 519	187 \pm 103	145 \pm 64	195 \pm 56
Cu	10.3 \pm 3.1	15.6 \pm 5.6	37.5 \pm 68.2	27.6 \pm 14.6	135 \pm 149	41.8 \pm 15.7	8.4 \pm 7.1	21.3 \pm 7.6
Pb	11.4 \pm 3.2	31.0 \pm 14.6	67.8 \pm 52.4	77.3 \pm 28.1	506 \pm 306	95.0 \pm 3.8	27.4 \pm 15.1	7.1 \pm 3.0
Cd	6.4 \pm 2.4	17.2 \pm 5.3	22.3 \pm 21.9	28.5 \pm 14.4	40.9 \pm 21.2	12.3 \pm 13.1	23.1 \pm 31.9	1.1 \pm 0.4
Ni	14.3 \pm 3.9	74.0 \pm 70.6	171.4 \pm 196.3	134.6 \pm 79	445 \pm 259	97.3 \pm 45.1	97.4 \pm 63.9	13.6 \pm 4.3
Mn	7.1 \pm 2.0	6.3 \pm 3.4	23.3 \pm 23.6	9.5 \pm 2.2	50.5 \pm 17.4	26.8 \pm 21.8	23.4 \pm 16.5	22.9 \pm 6.5
Ag	0.66 \pm 0.30	1.14 \pm 0.56	3.95 \pm 3.02	7.77 \pm 8.34	10.6 \pm 5.21	2.11 \pm 1.53	2.23 \pm 1.59	3.5 \pm 1.8
Co	4.16 \pm 2.48	19.5 \pm 6.6	51.4 \pm 38.6	42.2 \pm 17.6	231 \pm 118	72.7 \pm 65.2	14.4 \pm 7.1	14.7 \pm 3.9

TABLE 4

ORGAN PAIRS WITH SIGNIFICANTLY DIFFERENT CONCENTRATIONS

OF THE TRACE METALS LISTED

Higher Concentration	Kidney	Stomach	Foot	Mantle	Muscle	Gills	Soft Remains
Lower Concentration							
Soft Remains	Fe, Ni, Cd Pb, Cu, Co	Ni, Pb, Co	Ni, Cd, Pb	Ni, Cd	Pb, Cd	Cd	
Gills	Fe, Zn, Ni Pb, Mn, Ag Co	Fe	Ag				Mn
Muscle	Fe, Zn, Ni Pb, Cu, Ag Co	Fe, Cu, Co					
Stomach	Zn, Cd, Ag						
Mantle	Zn, Cu						
Foot	Zn, Mn						

TABLE 5

COMPARISON OF TRACE METAL LEVELS (ppm dry wt) IN MUSSELS FROM
OTHER STUDIES TO THOSE OBTAINED FROM THIS INVESTIGATION

Metal	Permissible Limit + :	Hobson's Bay		Frankston (PPB) (26) ⁺	Crib Point WPB (10)	Gulf of Speize Italy (30) [*]	Tasman Bay, NZ (28)
		Our study	(26) ⁺				
Fe		253			250-465		1960
Zn	9000	279	719	264	212-238	203-379	91
Cu	270	10.3	5.8	3.5	5.6-6.4	6.9-33.7	9
Pb	18	11.4	16.6	5.8	0.5	13.9-44.6	12
Cd	18	6.4	13.4	3.8	0.6-0.9	2.0-6.8	10
Ni		14.3				1.3-10.9	7
Mn		7.1				11.8-37.8	27
Ag		0.7					
Co		4.2				0.8-3.2	0.1

+ Calculated from wet weight

: Recommended by National Health and Medical Research Council

* *Mytilus galloprovincialis*





FIG. 1 - Locality Diagram

DISTRIBUTION LIST

MATERIALS RESEARCH LABORATORIES

Chief Superintendent
Superintendent, Physical Chemistry Division
Mr. J.H. Bishop
Mr. I.C. Dunstan
Mr. A. de Forest
Mr. R.W. Pettis
Library
Librarian, Materials Testing Laboratories, N.S.W. Branch
(Through Officer-in-Charge)

DEPARTMENT OF DEFENCE

Chief Defence Scientist
Deputy Chief Defence Scientist
Controller, Projects and Analytical Studies
Controller, Service Laboratories and Trials
Scientific Adviser - Army
Air Force Scientific Adviser
Navy Scientific Adviser
Chief Superintendent, Aeronautical Research Laboratories
Chief Superintendent, Weapons Systems Research Laboratory,
Defence Research Centre
Chief Superintendent, Electronics Research Laboratory,
Defence Research Centre
Chief Superintendent, Advanced Engineering Laboratory,
Defence Research Centre
Superintendent, Trials Resources Laboratory, Defence Research
Centre
Senior Librarian, Defence Research Centre
Librarian, R.A.N. Research Laboratory
Officer-in-Charge, Document Exchange Centre (16 copies)
Technical Reports Centre, Defence Central Library
Central Office, Directorate of Quality Assurance - Air Force
Deputy Director Scientific and Technical Intelligence
Head, Engineering Development Establishment
Librarian, Bridges Library, Royal Military College
Superintendent, Services Research Laboratories
Superintendent, Propulsion and Marine Physics
Hydrographer, RAN
Assistant Secretary, Defence Facilities Policy and Development
Officer-in-Charge, Joint Tropical Trials and Research Establishment
Officer-in-Charge, RANTAU
Director, Naval Quality Assurance
Director, Fleet Maintenance
Director, Naval Use Requirements

DISTRIBUTION LIST

(Continued)

DEPARTMENT OF DEFENCE (Cont.)

Director, Naval Ship Design
Project Director, Base Development
Director, Trials
Controller Publications - Navy
Chief of Naval Technical Services

DEPARTMENT OF PRODUCTIVITY

NASA Canberra Office
Head of Staff, B.D.R.S.S. (Aust.)

OTHER FEDERAL AND STATE DEPARTMENTS AND INSTRUMENTALITIES

The Chief Librarian, Central Library, C.S.I.R.O.
Australian Atomic Energy Commission Research Establishment
Division of Fisheries and Oceanography, C.S.I.R.O.
Australian Institute of Marine Science
Ministry of Conservation
Marine Chemistry Section, School of Chemistry, University of
Melbourne

MISCELLANEOUS - OVERSEAS

Defence Scientific and Technical Representative, Australian
High Commission, London, England
Assistant Director/Armour and Materials, Military Vehicles and
Engineering Establishment, Surrey, England
Reports Centre, Directorate of Materials Aviation, Kent, England
Library - Exchange Desk, National Bureau of Standards,
Washington, U.S.A.
U.S. Army Standardization Representative, C/o DGAD (NSO),
Canberra, A.C.T.
The Director, Defence Scientific Information & Documentation
Centre, Delhi, India
Colonel B.C. Joshi, Military, Naval and Air Adviser, High
Commission of India, Red Hill, A.C.T.
Director, Defence Research Centre, Kuala Lumpur, Malaysia
Exchange Section, British Library, Lending Division, Yorkshire,
England
Periodicals Recording Section, Science Reference Library, British
Library, Holborn Branch, London, England
Library, Chemical Abstracts Service, Columbus, Ohio, U.S.A.
INSPEC: Acquisition Section, Institution of Electrical Engineers,
Herts, England
Overseas Reports Section, Defence Research Information Centre,
Orpington, Kent, England

(MRL-R-781)

DISTRIBUTION LIST

(Continued)

MISCELLANEOUS - OVERSEAS (Cont.)

Engineering Societies Library, New York, U.S.A.
Naval Ship R & D Centre, Annapolis, U.S.A.
(Attention: Dr. H. Preiser and Dr. E. Fischer)
Central Dockyard Laboratory, H.M. Naval Base, Portsmouth, U.K.
(Attention: Dr. D. Houghton)
Naval Research Laboratory, Washington, U.S.A.
(Attention: Dr. R.R. Rojas)
Naval Undersea Center, San Diego, U.S.A.
(Attention: Dr. D.E. Andrews, Jnr.)
Scripps Institute of Oceanography, La Jolla, U.S.A.
(Attention: Dr. E. Goldberg)
Woods Hole Oceanographic Institution, Woods Hole, Mass., U.S.A.
(Attention: Dr. P.G. Brewer)
Biologische Anstalt Helgoland, Hamburg, West Germany.
(Attention: Dr. O. Kinne)
Oceanography Division, St. Louis, U.S.A.
(Attention: Mr. A. Pressman, Code 330)
Laboratorio de Ensayo de Materiales e Investigaciones Tecnologicas,
Argentina. (Attention: Dr. V. Rascio)
Department of Zoology, National Science Museum, Tokyo, Japan.
(Attention: Dr. S. Mawatari)
Department of Oceanography, University of Southampton, U.K.
(Attention: Dr. J.D. Burton)
Department of Oceanography, University of Liverpool, U.K.
(Attention: Dr. J. Riley)
National Institute of Oceanographic Sciences, Surrey, U.K.
(Attention: Dr. Culkin)
Miami Marine Research Inc., Florida, U.S.A.
(Attention: Dr. S. Miller)
Defence Research Establishment Pacific (DREP), Canada
(Attention: Mr. J. Carson)
Marine Science Laboratories, North Wales, U.K.
(Attention: Dr. Crispe)
Scottish Marine Biological Association Laboratory, Scotland.
(Attention: Dr. H. Barnes)
Centre ORSTOM, B.P. No.A5, Noumea, New Caledonia.
(Attention: Dr. J.R. Donguy)
U.S. Environmental Protection Agency, West Kingston, U.S.A.
Fisheries Research Board of Canada, Marine Ecology Laboratory,
Bedford Institute, Dartmouth, Canada
(Attention: Dr. I. Duedall)
Graduate School of Oceanography, University of Rhode Island, U.S.A.
(Attention: Dr. D.R. Kester)
Defence Scientific Establishment, HMNZ Naval Dockyard, N.Z.
(Attention: Dr. D. Barnes)
Secretariat General, Comite International Permanent pour la, France
Department of Biological Sciences, Portsmouth Polytechnic,
Portsmouth, England (Attention: Dr A. Chamberlain)